1   **Discovering paracrine regulators of cell type composition from spatial transcriptomics using SPER**

3   Tianxiao Zhao[1,2], Adam L. Haber[2]

6   [1] Institute of Systems Genetics, New York University Grossman School of Medicine, New York, NY

7   [2] Department of Environmental Health, Harvard T.H. Chan School of Public Health, Boston, MA.

10   Correspondence to Adam L. Haber (ahaber@hsph.harvard.edu)

SUMMARY:

A defining characteristic of biological tissue is its cell type composition. Many pathologies and chronic diseases are associated with perturbations from the homeostatic composition, and these transformations can lead to aberrant or deleterious tissue function. Spatial transcriptomics enables the concurrent measurement of gene expression and cell type composition, providing an opportunity to identify transcripts that co-vary with and potentially influence nearby cell composition. However, no method yet exists to systematically identify such intercellular regulatory factors. Here, we develop Spatial Paired Expression Ratio (SPER), a computational approach to evaluate the spatial dependence between transcript abundance and cell type proportions in spatial transcriptomics. We demonstrate the ability of SPER to accurately detect paracrine drivers of cellular abundance using simulated data. Using publicly available spatial transcriptomics data from mouse brain and human lung, we show that genes identified by SPER show statistical enrichment for both extracellular secretion and participation in known receptor-ligand interactions, supporting their potential role as compositional regulators. Taken together, SPER represents a general approach to discover paracrine drivers of cellular compositional changes from spatial transcriptomics.

Keywords: Spatial transcriptomics, spatial data analysis, spatial compositional analysis, systems biology, computational biology, cell circuits, network biology, cellular composition, cell-cell communication.

INTRODUCTION:

Given the specialized character of each cell type, the distribution of types that compose each biological tissue is fundamental to mediating physiological functions. Accordingly, compositional changes, such as influx of immune cells, proliferation of a stem cell pool, or metaplasia of fibroblasts, are tightly regulated by paracrine signaling networks, and these networks are fundamental organizing principles of biological tissues [1]. For example, inflammation can lead to cytokine-mediated changes in cell type composition, such as the canonical 'weep and sweep' response to intestinal parasites, involving mucus-producing goblet cell hyperplasia, eosinophilia and basophilia [2]. Many diseases involve this type of change in corresponding tissues: in Alzheimer's disease for example, neuropathology is associated with aberrant proliferation of microglia and astrocytes [3], which may be mediated by paracrine CX3CL1/CX3CR1 signaling [4]; in asthma, airway remodeling driven by IL-13 leads to increased numbers of smooth muscle and goblet cells accompanied by the abnormal recruitment of immune cells [5]. Triggering factors, like environmental exposures [6] and microenvironmental signals [7], can induce cellular composition changes. To gain insight into the normal physiology of tissue composition and transformations associated with disease, it is essential to understand *which signals regulate the compositions of cell types* in a tissue.

Single-cell RNA-seq (scRNA-seq), combined with computational tools, now provides the transformative ability to simultaneously estimate cell-type composition and measure the expression of factors that regulate it. However, scRNA-seq is unable to determine whether the signals and the receptor cell types are sufficiently near to each other that a putative signaling process is physiologically possible. Spatial transcriptomics (ST) promises to remove this hurdle by simultaneously revealing (1) the expression of paracrine ligands and receptors, (2) the pattern of cell type compositions, and (3) the spatial relationships between the relevant pairs of transcripts and cell populations. ST is an overarching term for a range of methods designed to capture transcriptome-wide gene expression while retaining information about the spatial tissue context [8]. The most widely-used and relatively affordable technologies, including ST Visium [9] and Slide-seq [10], capture spatial information in voxels around 55 microns in diameter, larger than a single cell. Such restriction results in the generation of multi-cell resolution spatial expression data where the absolute number of cells in each voxel is difficult to determine. To address this, recovering the relative proportions of component cell types, can be straightforwardly achieved by available deconvolution algorithms. Probabilistic-based approaches, like RCTD [11], cell2location [12], and SONAR [13], model the expression data with certain distributions under a Bayesian likelihood framework. All these methods produce data which has the constraint that in each spot all proportions sum to 1, and thus their output is compositional. When combined with gene expression data, this type of spatial compositional data can potentially be used to discover paracrine drivers of tissue transformation, but no computational tools yet exist to do so.

In lieu of appropriate methods, several studies have used classical statistical correlations, such as Pearson's R and Spearman's ρ, to identify such signals in spatial transcriptomics data [14-16]. However, spatial compositional data possesses unique characteristics that requires tailored analytical approaches [17], as treating each individual cell type's composition as an independent measurement can result in spurious correlations [18]. Additionally, classical metrics face other limitations in this context. For a given cell type A, transcriptional signals with spatial dependence on its composition fall into two categories. The first includes A-type 'marker' genes—those specifically expressed by A—which exhibit a trivial spatial relationship with A's composition simply because A cells are their primary source. The second, and more

72 intriguing, category includes genes not expressed by *A* cells but whose spatial patterns correlate with *A*'s
73 composition, indicating that they are instead produced by neighboring, non-*A* cells. These genes are
74 potential paracrine regulators influencing *A*'s distribution and abundance.
75
Machine learning methods have also been applied to spatial transcriptomics to model spatial patterns and
cell-cell interactions. For example, MISTy [19] builds multiple components to model the spatial contexts
and extracts the spatial relationships between spatial marker genes. SpatialDM [20] uses a bivariate Moran's
statistic to identify the direct spatial association between ligand-receptor pairs, while SpaTalk [21] analyzes
cell-level ligand-receptor interactions by testing for enrichment in co-expressed spots. Although these
methods are powerful approaches to analyze the spatial distributions of transcriptional signals, most of them,
as well as correlation metrics and methods like scHOT [22] and SpatialCorr [23], focus on direct spatial
cell-cell interactions and therefore are unable to capture the second category of genes above, as spatial
signals in this category may exhibit no direct overlap at all. In addition, none of these methods explicitly
model cell type composition, or offer a mechanism to detect drivers of compositional changes.
86
To overcome this problem, we introduce Spatial Paired Expression Ratio (SPER), a new computational
framework to detect paracrine signals regulating the composition of target cell types, even when these
signals exhibit non-overlapping spatial dependence. SPER is inspired by the pair correlation function (radial
distribution) from physics, which quantifies spatial correlations between two particle types by estimating
the likelihood of one particle appearing at a given distance from another [24]. To denoise the spatial
compositional signal while appropriately modeling its compositionality, we apply co-kriging, a
geostatistical method [25], prior to computing pair correlations to identify the transcripts most strongly co-
varying with spatial compositions.
95
In the following sections, we describe the SPER algorithm and demonstrate, using simulated data, that it
substantially outperforms all other applicable metrics: Pearson correlation, Spearman correlation, and Earth
Mover's Distance (EMD). We apply SPER to published ST profiles from adult mouse brain and human
lung, and demonstrate that genes identified by SPER are strongly and significantly enriched for transcripts
encoding proteins known to be extracellularly secreted, and also known to participate in intercellular
receptor-ligand interactions, both consistent with potential roles in paracrine interactions. Moreover, SPER
identified novel, biologically significant interactions in the mouse brain data, such as identifying the
neuropeptide precerebellin (*Cbln1*) as a novel factor associated with abundance of GABA-ergic Meis2$^+$
neuron, which express its cognate receptor *Nrxn2*, as well as the Wnt ligand R-spondin 3 (*Rspo3*) associated
with the same Meis2$^+$ neuron subset, which also express its receptor *Lgr5*. In human lung data, SPER detects
the fibroblast-derived signal fibrillin-1 (*FBN1)* associated with alveolar type II (AT2) cells, which express
the integrin αvβ6 (*ITGB6*), one of the known receptors for FBN1. These novel putative paracrine
interactions detected by SPER were not identified using other available computational methods. Taken
together, our findings establish SPER as a general and effective approach for identifying putative regulators
of cell type composition from spatial transcriptomics data.
111
112

RESULTS:

SPER scores paracrine signals by evaluating spatial dependence between genes and cell types.

Spatial Paired Expression Ratio (SPER) is a computational framework designed to uncover paracrine signals that regulate cell type compositions within spatial transcriptomics datasets. Unlike classical correlation-based approaches that are prone to detecting trivial spatial associations, SPER explicitly models the spatial interplay between genes, potentially encoding paracrine signal molecules, and cell types within a certain distance. Studies suggest that paracrine signals typically function within a maximal distance of 25 cell diameters, approximately 250-300 μm [26]. To detect these interactions, SPER employs the pair correlation function that can describe how spatial correlation changes as a function of distance. For every gene-cell type pair, SPER evaluates how the dependence of the gene's expression level on the target cell type's abundance varies with distance to the target cell type and identifies those pairs showing strong a signal of paracrine spatial dependence.

The method integrates three key components: spatial gene expression, spatially resolved cell-type compositions (derived from deconvolution algorithms), and spot-to-spot spatial distances (**Fig. 1**). As the first step, SPER applies geostatistical techniques, specifically co-kriging, to de-noise compositional data while accounting for inherent spatial dependencies. For each gene-cell type pair, it then computes the gene's expression ratio to its global average expression level, defined as 'paired expression ratio', across different distance lags. The outcome, as a paired expression ratio spatial density distribution, will be convolved with a spatial weight into one score to prioritize biologically relevant interaction scales. After adjusted with the expression prevalence in the target cell type, a SPER score matrix can be eventually calculated based on every pair of gene and cell type. This workflow enables SPER to differentiate between direct, overlapping spatial associations and more subtle, non-overlapping dependencies indicative of paracrine regulation.

While no method yet exists to mine ST data for paracrine signals regulating cell type composition, the major analytical field to derive inferential tools for spatial compositional data has been geostatistics. This branch of statistics focuses on modeling multivariate data that show codependency related to spatial proximity, originally developed in predicting the co-distribution of metal ores for mining applications [27]. Co-kriging is a geostatistical interpolation approach that models' compositionality as well as spatial dependance between components of a mixture. It uses kernel functions to describe the pattern of variograms between each pair of components as a function of geographic lag distance [18]. SPER leverages these desirable properties of co-kriging to de-noise the spatial compositional data, modeling variograms between cell type proportions, and adjusting for spatial autocorrelation. Then, using the distance information obtained from the physical coordinates of all ST spots, calculates the paired expression ratio as a series of matrices, which includes all pairs of cell types and genes; finally, SPER convolves the ratios at each distance with a weight function to integrate over length scales of interest and obtain a scalar score representing the spatial relationship between the given cell type and the gene (**Fig. 1** and **Methods**).

153
Figure 1. Schematic of spatial paired expression ratio (SPER) algorithm. The workflow of SPER: (Left) **Pre-processing**:
SPER requires three different inputs: spatial transcriptomics data, spatial cell-type compositional data, and the spot-spot distance
matrix. Any cell-type deconvolution method can be used to generate the spatial compositional data. (Center) **Core algorithms**:
SPER models the spatial dependence between genes and their potential target cell types by measuring the expression changes over
the interaction distance of a cell type-gene pair. As the first step, we apply co-kriging algorithms to model and de-noise the spatial
compositional data. Next, we evaluate the spatial association by calculating a gene's ratios to its mean expression over the distance
towards the target cell type. These ratios are defined as the paired expression ratios, and a spatial weight function on these ratios is
applied at the last step to emphasize paracrine signal patterns. (Right) **Results**: the algorithm generates a SPER score matrix as the
final output, where higher scores indicate a stronger confidence of this cell type-gene pair as a putative paracrine signal, with a
schematic example transcript cell-type pair shown (bottom right).

SPER detects simulated paracrine drivers of cell type composition.

To evaluate how well SPER detects paracrine regulators of cell composition, we applied it to three different
simulated ST datasets (**Fig. 2**) and compared it with traditional methods of detecting spatial co-variation,
specifically: Pearson correlation, Spearman correlation, and Earth Mover's Distance (EMD) [28]. To set
up these synthetic benchmark datasets, we simulated 80-by-80 grids where each tile represents a spot in the
physical space. Next, different cell types and gene expression patterns were assigned to tiles in the following
manner: cell type specific 'marker genes' for each cell type, only expressed in the corresponding cells, were
defined across the space; the 'paracrine signal genes', which we are interested in, can be mathematically
described as genes having a similar but not directly corresponding spatial distribution as their target cell

type. Such genes can be expressed either by a sole source cell type or be jointly expressed by multiple cell
types. Therefore, we setup three different scenarios to simulate the paracrine signals: simplified single-
source, realistic single-source, and realistic multi-source paracrine signal patterns (**Fig. 2, left**), using
radially dependent association for a simplistic and Markov random fields for more realistic simulations
(**Methods**). With these simulated ST data, cell-type deconvolution methods were then applied to generate
spatial cell type compositional data. SPER, as well as the other canonical statistical metrics, were then
applied to detect potential paracrine signals affecting target cell composition.
181
In the simplified single-source simulation, paracrine signal genes are expressed by a single source cell type
located at a fixed average radial distance of 10 units from the target cell type. This setup ensures that the
spatial distributions of signal genes spatially depend on, *but do not substantially overlap with*, the
distributions of the corresponding cell types (**Fig. 2, top row**). We reasoned that this lack of overlap would
be characteristic of a paracrine signaling interaction where a neighboring cell population provides a ligand
to a target cell population. To enhance the biological fidelity of the synthetic data, we developed a more
realistic single-source simulation where the spatial distribution of cell types is modeled using a Markov
random field [29]. In this scenario, target cell abundance is spatially associated with, but again remain
distinct from, the abundance of their simulated paracrine ligand in a more biologically realistic arrangement
(**Fig. 2, middle row**). Furthermore, to account for signaling pathways involving multiple sources, we
simulated a multi-source scenario in which target cells are surrounded by two source cell types expressing
the signal genes (**Fig. 2, bottom row**).
194
195

Figure 2. Evaluation of SPER on different simulated spatial transcriptomics data. **Left column**. Schematic of the three paracrine signal patterns - simplified single-source, realistic single-source, realistic multi-source - simulated in this study. **Middle column**. Visualizations of representative simulated spatial compositional data for each pattern type (rows) in an 80x80 grid, with each tile colored by its major cell type (color legend). In the realistic scenarios (bottom two rows), Markov random fields were used to simulate the spatial dependence between the sources and target. **Right column**. Area under ROC curve (AUROC) values are calculated to evaluate how well each metric distinguishes simulated paracrine signals from other spatially unrelated genes.

To compare the paracrine pattern to other genes, we also simulated a control set of genes that are uniformly distributed in the grid and therefore have no spatial association with any cell type. We considered the marker genes expressed by target cell types as part of control group. Thus, we can evaluate a method's performance

207 on detecting paracrine gene signals by how well it can distinguish them from the control set. Four different
208 metrics, SPER, Pearson and Spearman correlation, and Earth Mover's Distance (EMD), were applied in the
209 simulations (**Fig. 2, right**). While all other metrics completed in minutes, EMD, a method with extremely
210 high computational complexity, failed to complete within two hours. Therefore, we decide to exclude this
211 metric from the list as it is therefore impractical to apply on real data.
212
We repeated each simulation 10 times to generate robust results. Given the output scores from each method,
we calculated the ROC curve of discriminating the paracrine signal genes from other genes. While all three
methods exhibited a good detection accuracy in the simplified simulation, SPER consistently showed the
best performance, accurately detecting signals in all three scenarios with AUROC scores from 0.92 to 0.99.
Pearson and Spearman correlations, which did well in the simplified scenario, failed to capture the paracrine
signals in two more realistic simulations with the AUROC scores from 0.56 to 0.76. Taken together, our
simulation studies demonstrate the ability of SPER to detect paracrine signals that might be missed by other
traditional metrics, especially in a more complex and realistic settings.
221
SPER outperforms classical metrics in detecting potential paracrine signals in real ST data
To evaluate its performance on real data we next applied SPER and the other metrics to ST profiles of
mouse brain generated using 10X Visium [9]. After preprocessing, 18270 unique genes were detected
across 2695 spots sampled from a sagittal anterior mouse brain section. Based on benchmarking studies
evaluating performance on this task [30-32], we selected RCTD [11], which consistently ranks among the
top-performing methods, in this study to estimate the cell type composition with matched scRNA-seq as
the reference [33]. (**Fig. S1A**). SPER's co-kriging models showed good fits to the observed variograms of
top ten cell types (**Fig. S2**), and visualizing the compositional data after co-kriging demonstrated its ability
to remove noise (**Fig. S1B**). For each of the ten neuronal cell types in the dataset, we used a published
scRNA-seq reference dataset to define cell-type-specific marker genes that were strongly upregulated in
one particular cell type (at least two-fold higher, FDR<0.001, calculated using MAST [34]). Given the cell
type source of these transcripts, we expect their expression distribution in the ST data to be spatially strongly
coupled to the abundance of that cell type. Using Pearson and Spearman correlations, we confirmed that
these marker genes showed significant spatial dependency to their corresponding cell types (**Fig. S3**).
236
However, our goal is to identify paracrine signals, we hope to detect those genes that are not expressed (or
only lowly expressed) in the target cell type but are instead expressed by neighboring source cells. We
defined 'putative paracrine transcripts' as those whose spatial dependency was strongly associated with a
target cell type but had only low expression in that target cell type. To maximize detection of these
transcripts associated with—but not expressed by—a cell type, the SPER score also considered the gene
expression prevalence matrix from the scRNA-seq expression data (**Methods**). To visualize the extent to
which putative paracrine transcripts were enriched in each metrics top ranked genes, we plotted the metric
scores against the proportion of cells expressing each gene in the reference scRNA-seq data [33] for each
cell type-gene pair (**Fig. 3A-B, S5**). In L4 (Layer 4) neurons for example (**Fig. 3A-B**), all genes above the
95[th] percentile of Pearson correlation also showed high expression prevalence in L4 neurons; and so, it
detected just 3 putative paracrine signals whose expression prevalence less than 0.5 at the 95[th] percentile
threshold (**Fig. 3A**), indicating that virtually all transcripts associated with L4 neurons are expressed by
them, and many of them are trivially L4 marker genes. On the other hand, in the case of SPER, there are no
marker genes within the top 95[th] percentile of association scores, and the majority meet our baseline

criterion to be putative paracrine transcripts , which we define as genes whose expression prevalence is less
than 20% in the target cell type (**Fig. 3B**). This trend was consistent across all cell types after varying the
metric percentile thresholds from 75th to 95th, where SPER detected markedly more putative paracrine genes
meeting this criterion (**Fig. 3C**), supporting SPER's capability to detect putative paracrine regulators of cell
type composition.

257
Figure 3. SPER detects non-trivial spatial relationships. A-B. Scatter plots show the distribution of Pearson correlations (A)
and SPER scores (B) with the proportion of layer 4 (L4) cortical neurons in which each gene (dot) is detected in scRNA-seq profiles.
The dark blue dots mark the marker genes for each cell type (**Methods**). Red dots stand for the genes whose metric scores are
above the 95% percentile. Purple dots represent genes that meets both conditions. (C) Proportion of top-ranked (above the given
percentile, x axis) genes that are not trivially expressed by the target cell type (defined as expression prevalence less than 20% in
the target cell type) for each metric (color legend, top).

Transcripts with high SPER scores are enriched for extracellular proteins and paracrine ligands.
If the genes spatially related to cell type composition are indeed participating in paracrine interactions, then
we reasoned that they must encode proteins capable of performing these functions in facilitating
intercellular communication. We therefore tested whether the putative paracrine transcripts identified by
SPER and other methods were statistically significantly enriched for (a) genes known to be secreted into
the extracellular space and (b) genes known to act as paracrine ligands (**Fig. 4**). As before, we set thresholds
on the metrics, requiring them to have a SPER or other metric score larger than the 95th percentile of all
genes. As expected, transcripts identified by SPER were significantly enriched ($p<0.001$) for both
extracellular genes (**Fig. 4A**) and paracrine ligands (**Fig. 4B**) for all ten cell types in the mouse brain dataset.
The other two metrics were significantly enriched ($p<0.05$) only in three and four cell types for Pearson
and Spearman correlation respectively for extracellular genes (**Fig. 4A**), and in four and three respectively
for known paracrine ligand genes (**Fig. 4B**). With few exceptions, SPER identified more of both types of
transcripts compared to Pearson and Spearman correlations (**Fig. 4**). Lastly, we performed a sensitivity
analysis to confirm the utility of co-kriging in denoising our compositional data (**Methods**). We found that
while transcripts highly ranked by SPER without co-kriging still showed enrichment, the addition of co-
kriging improved the enrichment for both extracellular location and known paracrine ligand activity for the
majority of cell types, demonstrating its utility in improving SPER's sensitivity (**Fig. S4**).

Together, these results demonstrate that candidate paracrine transcripts identified by SPER have three
important characteristics: (1) they are predominantly expressed by a neighboring cell type, not the target

cell, but they are nonetheless spatially related to the target cell's composition, and they are also statistically
enriched for (2) genes known to be secreted into the extracellular space, and (3) genes known to participate
as ligands in paracrine interactions.
288

289
Figure 4. SPER's top-ranked genes are enriched for both extracellular genes and known paracrine ligands. Dot plots
visualize hypergeometric enrichment analysis of extracellular (A) and known paracrine ligand signals (B) within candidate gene
sets. Extracellular and ligand genes annotations were collected from COMPARTMENTS, FANTOM5, and CellPhoneDB
databases respectively (see **Methods**). The candidate sets include genes whose scores for each spatial dependency metric (x-axis)
are above than the 95% percentile of all genes. The size of dots represents the number of extracellular genes or ligand genes in
the given gene set. The color represents the significance the enrichment in the set (hypergeometric test), non-significant ($p>0.05$)
overlaps are shown as light grey.
297

SPER detects single- and multi-source putative paracrine signals in mouse brain.
Using SPER, we have detected a set of putative ligand genes that have high spatial association with the
composition of a putative target cell type. Across all cell types there are 5569 pairs of cell types and putative
paracrine regulators above the 95th percentile of SPER scores. Of these, 250 also showed substantial
expression of the cognate receptor in the target cell types: a total of 506 ligand-receptor pairs, each
potentially active in regulating the composition of the brain tissue in the sample. For example, *Cbln1*
(precerebellin) and Meis2+ neurons (a cell type defined by expression of the homeobox transcription factor
Meis2 [33, 35]) showed a strong spatial association, but also non-overlapping spatial distribution, indicating
possible paracrine compositional regulation (**Fig. 5B**). Notably, it was not identified by Pearson correlation,
ranking 349th ($R=0.32$). On the other hand, using SPER we found that the *Cbln1*-Meis2+ neurons pair had
the 4th highest score among all Meis2+ neuron-related signals where the appropriate cognate receptor was
also expressed. Consistently, the scRNA-seq data showed that *Cbln1* is only lowly expressed in Meis2+
neurons, but is indeed highly expressed by the neighboring L6 IT neurons (**Fig. 5A**). In this case, reference

scRNA-seq data showed that the paracrine signal is expressed only by this single source cell type. We
examined *Nrxn2*, a receptor for *Cbln1* (**Fig. 5C**), and found as expected that it was expressed on Meis2[+]
neurons, supporting the possibility of Cbln1 acting on them. In the cerebellum, *Cbln1* is known to play an
important role in the Purkinje cell synapse formation [36] as well as the growth and guidance of axon in
multiple neural regions within mouse embryos [37], but to our knowledge, its role as a paracrine signal in
the motor cortex has not yet been described.
317

318
Figure 5. SPER detects single-source (*Cbln1*) and multi-source (*Rspo3*) paracrine signals in mouse brain. (A, D) Violin plots
show expression levels (y-axis) of *Cbln1* and *Rspo3* in relevant mouse brain cell types (x-axis) from reference scRNA-seq data. L6
IT, L5 PT, L5 IT, L4, and Pvalb neurons are the neighboring cell types of Meis2[+] neurons. (**B, E**) ST plots visualize the spatial
dependence between transcript-target cell type pairs *Cbln1*/Meis2[+] neurons (B) and *Rspo3*/Meis2[+] neurons (E). The color of spots
(color legend) shows whether *Cbln1* (B) or *Rspo3* (E) is detected, a proportion of Meis2 cell larger than 35%, both (overlap), or
otherwise. (**C, F**) The expression levels of the relevant cognate receptor *Nrxn2* (C) and *Lgr5* (F) in the reference scRNA-seq data.
325

Paracrine signals can of course also be derived from multiple cell types. For example, SPER identified
Rspo3, which encodes the Wnt ligand R-Spondin 3. *Rspo3* is produced by several source cells: Pvalb[+], L6
IT, and L4 neurons (**Fig. 5D**), and shows a non-overlapping spatial association with the identified target
cells, again Meis2[+] neurons (**Fig. 5E**). While *Rspo3* is identified as a significant putative signal by SPER,

it cannot be found as a paracrine signal using any correlation metrics (**Supplementary Table 1**). We tested
for expression of the cognate receptor for *Rspo3* in Meis2 cells and found that *Lgr5* (Leucine-rich repeat-
containing G-protein coupled receptor 5) was strongly and selectively expressed in the target cell type,
consistent with an active paracrine interaction (**Fig. 5F**). This suggests that *Rspo3*, released by Pvalb, L6
IT and L4 cells, may serve as a paracrine regulator to Meis2 cells, activating the *Wnt* signaling pathways,
which could potentially shape the spatial compositions of Meis2$^+$ neurons. While this interaction is well-
studied in the intestine [38], to our knowledge its role in the mouse motor cortex is not yet known. The link
between *Rspo3* and Meis2 cells, therefore, provides a representative example of multi-source paracrine
compositional regulators that can be discovered using SPER.
339
SPER enables discovery of paracrine signals from established human lung spatial dataset.
To validate SPER's sensitivity to detect such paracrine signals, we also applied it to a second published ST
dataset of the human lung [39]. The processed Visium spatial transcriptomics data was deconvoluted by
cell2location [12], based on the paired reference scRNA-seq data with a total of 1147 genes in
approximately 200 thousand cells. The deconvolved ST data (**Fig. 6**) contains 3234 spots and the relative
compositions of 80 annotated cell types. We applied SPER (Methods) to identify potential paracrine
regulators of human lung cell type abundances (**Supplementary Table 2**). In addition to identifying many
known paracrine interactions—such as club cell-derived *SCGB3A2* and alveolar macrophage abundance,
via the *MARCO* receptor [40] or basal epithelial cell-derived *CXCL14* and B cell abundance, via the receptor
CXCR4 [41]—SPER identified a novel fibroblast-derived signal: fibrillin-1 (*FBN1*), an extracellular matrix
(ECM) protein [42], was spatially associated with the composition of alveolar type II (AT2) cells (**Fig. 6A-**
B). The association for FBN1 and AT2 cells was very strong, ranking in the 98th percentile of SPER scores
(20th of 1147 tested genes), but both Pearson and Spearman correlation scores failed to capture this signal
with R=0.056 and ρ=0.061, outside the top 200 ranked transcripts for both metrics. Consistent with a
plausible paracrine signaling role of fibroblast-derived FBN1, integrin αvβ6 (*ITGB6*), an epithelial-
exclusive integrin that acts as a receptor for FBN1, is strongly expressed on AT2 cells (**Fig. 6C**). In lung
organoid models, diverse ECM proteins including FBN1 are known to be critical for differentiation and
proliferation of AT2 cells [43], but to our knowledge, the direct link between fibroblast derived *FBN1* and
ITGB6 on AT2 cells has not been previously identified as active *in vivo*. Mutations in *FBN1* result in
increased availability of TGFβ and are linked to the congenital disease Marfan's syndrome, the effects of
which are predominantly cardiovascular, skeletal, and ocular, but also induce a respiratory phenotype [44]
including lung lesions and spontaneous pneumothorax [42].
362

363
364 **Figure 6. SPER detects non-overlapping paracrine signals (*FBN1*) in human lung.** (**A**) Violin plots show expression levels (y-
365 axis) of *FBN1* in relevant human lung cell types (x-axis) from reference scRNA-seq data. (**B**) ST plots visualize the spatial
366 dependence between transcript-target cell type pairs *FBN1*/AT2 cells. The color of spots (color legend) shows whether *FBN1* is
367 detected, a proportion of AT2 cell larger than 15%, both (overlap), or otherwise. (**C**) The expression levels of the relevant cognate
368 receptor *ITGB6* in the reference scRNA-seq data.
369
370

DISCUSSION:

Sequencing technologies and computational methods for spatial transcriptomics are both developing rapidly, promising an exciting new wave of biological discoveries informed by spatial insight. However, despite the importance of cell type compositional changes, which are frequently the first analytical question asked in spatial and single-cell studies, and the potential of ST data to map them, and the well-established role of paracrine signals in dictating tissue composition [1], to our knowledge there has been no method to detect putative compositional regulators; SPER provides the first such method.

Although we performed evaluations using ST data generated with 10X Genomics Visium, SPER is not specific to this modality, and naturally generalizes to other spot-level technologies, such as Slide-seq [10] and XYZeq [45]. In these cases, the SPER workflow remains unchanged, requiring input ST expression data in the form of a genes-by-location matrix, spot locations, and estimated cell-type spatial compositional data from a deconvolution algorithm. For newer spatial technologies, such as CosMx (NanoString), MERSCOPE (Vizgen), and Xenium In Situ (10X Genomics), the spot-level and deconvolved cell-type compositional data is replaced by the cell-level spatially resolved data. This shift requires additional preprocessing for the data, treating every single cell as a spot with only one cell type. While the workflow remains functional, the computational burden for the single-cell data, often involving millions of cells, could be significantly higher than for Visium data, which typically includes two to four thousand spots. Since workhorse ST methods such as 10X Visium do not have single-cell resolution, analysis of their data using SPER requires estimating cell type compositions from ST data using any deconvolution method: SPER then models the compositionality of the spatial data using co-kriging, which can reduce the impact of noisy estimates while preserving core spatial patterns [18, 46]. Methods with higher deconvolution accuracy will increase the performance of SPER to capture biologically meaningful signals. Hence, we expect the SPER algorithm to be more powerful in the future as improved ST technologies and superior deconvolution approaches emerge.

Algorithms for paracrine signal inference such as SPER face two related challenges given the complexity of receptor ligand interactions. First, if a receptor-ligand interaction has been previously described, and SPER identifies it as highly significant in a particular dataset, since it is based on transcriptional measurements, this is only an inference, rather than definitive proof that the interaction is physiologically relevant. In this context, SPER operates as a discovery tool, generating hypotheses regarding the paracrine interactions that could be active in the tissue, and follow-up experiments are then necessary to translate these potential interactions into definitive new biological mechanisms. As we've shown, even a cursory inspection of the results of analysis by SPER in a well-known tissue (mouse brain), identifies both known and novel potential interactions. We thus expect that SPER can provide the first step in an analytical pipeline to discover active intercellular circuits that underly tissue structure, and if applied to perturbations or disease, that underly tissue transformation. A second challenge is that not all paracrine interactions are currently annotated—for this reason, SPER does not ignore putative regulatory transcripts that are unannotated as ligands or receptors, but rather enables a user to examine *all* highly ranked transcript-cell type pairs. In this way SPER can potentially uncover previously unknown intercellular signals that are associated with changes in tissue composition, providing the ability to identify regulators even, for example, if their receptor is unknown or is not detected due to lack of sensitivity.

Taken together, our results demonstrate that the application of SPER to ST data has the potential to identify
which paracrine signals are involved in the intercellular circuits controlling homeostatic tissue composition,
stress responses, tissue inflammation, and perhaps most importantly, pathological tissue transformations
associated with disease. Identification of these signals is a critical step towards deepening systemic
understanding of these complex processes and pinpointing novel biomarkers and potential therapeutic
targets.

METHODS:

SPER: Spatial Paired Expression Ratio. In general, SPER algorithm requires the normalized spatial
transcriptomics data $E_{n \times m}$ (the expression level of genes divided by their global average across all spots),
the spots distance matrix $M_{n \times n}$, and the cell-type compositional data $C_{k \times n}$ as the input, where n is the
number of spots, m as the number of genes and k as the number of cell types. As the first step, the distance
matrix of spots M is separated into l adjacency matrices A_i, each recording whether two cells lie within
certain distance range $d_{i-1} \sim d_i$ for $i \in \{0, 1, \dots, l\}$ with $d_0 = 0$. As a result, the distance matrix M can be
estimated as the summarization of $d_i A_i$, where the original distances are binned into different groups. The
default bin of the distance discretization is the distance between two spots, and all distances which are larger
than the given largest distance limits are trimmed to that value. These adjacency matrices $A_i \in \text{Boolean}^{n \times n}$
are normalized by multiplying a diagonal matrix $F_{i,n \times n}$, where each element is the reciprocal of each row's
sum of A_i. Then, they are multiplied with the spatial transcriptomics data $E_{n \times m}$ and the cell-type
compositional data $C_{k \times n}$ as:

439
$$M \approx \sum_{i=1}^{l} d_i A_i$$

440
$$F_i = \text{diag}(\{\frac{1}{\sum_{k=1}^{n} A_{i,j,k}}\}_{j=1}^{n})$$

441
$$R_i = C \times A_i \times F_i \times E$$

442 where R_i represents the $k \times m$ paired expression ratio matrix at the distance range $D_{i-1} \sim D_i$. Finally, a
443 weight function $\mathbf{\Phi} = (\varphi_1, \dots, \varphi_l)$ is applied on each gene/cell type pair across $R_1, \dots, R_l$. The default weight
444 function follows a Poisson distribution $\text{Pois}(\bar{\varphi})$ which has an extra penalty on overlapping and distal
445 regions, where the parameter $\bar{\varphi}$ is chosen based on the potential functional distance of paracrine signal.
446 Eventually, we can get the SPER score matrix S as:

447
$$S = \sum_{i=1}^{l} \varphi_i R_i = \sum_{i=1}^{l} \varphi_i C A_i F_i E$$

Spatial ST data simulation. The simulations are created by the following steps: 1) A 80x80 grids is created
and its distance matrix is calculated; 2) Cell type labels are assigned to each tile of the grid, which follows
different mechanisms in three scenarios; 3) For each cell type, we create the corresponding marker genes
and simulated their expression levels across all cells. Corresponding cell types will have higher counts of
their markers with their counts following Poisson distributions $\text{Pois}(\lambda)$ where $\lambda \sim \text{Unif}(15, 30)$. For other
genes, including the non-marker and the control genes, their counts follow Poisson distributions $\text{Pois}(\lambda')$
where $\lambda' \sim \text{Unif}(0, 3)$. 4) The compositional data for each cell type is computed from marker genes'
expression level. The count of each cell type within a grid is sampled from another Poisson distribution.
The final cell-type proportions are the fractions of each cell types within the tiles. Mathematically it can be
shown as:

459
$$\tilde{y}_{g,k} \sim \text{Pois}(\lambda_{g,k})$$

460
$$c_{i,k} \sim \text{Pois}(\sum_{g \in \text{marker of } i} \alpha \lambda_{g,k} \tilde{y}_{g,k})$$

461
$$\tilde{c}_{i,k} = \frac{c_{i,k}}{\sum_i c_{i,k}}$$

$$\lambda_{g,k} \sim \begin{cases} \text{Unif}(15, 30), & \text{if } g \text{ is a marker} \\ \\ \text{Unif}(0, 3), & \text{otherwise} \end{cases}$$

where $\tilde{y}_{g,k}$ is the expression level of gene g at tile k, $\tilde{c}_{i,k}$ is the cell-type proportion of cell type i at tile k,
and α is the scale factor.

Simplified single-source scenario simulates the source and target cells in a spatial relationship as ring (**Fig.**
2). Within the 80x80 grid, five cell types, two pairs of source and target cells and a control, are simulated.
For each simulated cell type, we created 20 marker genes from two-dimensional normal distributions and
calculated the cell-type spatial composition as described above. While the target cells have their marker
genes simulated with given means and variances, the markers for source genes have their maximal values
10-unit away from the centers of target cell types. This spatial dependence characterizes the paracrine
patterns between the source genes and target cells.

Realistic scenarios, both single-source and multi-source, we use Markov random field algorithm from the
R package 'mrf2d' to simulate the spatial relationships between cells. Neighbors with L1 distance less than
3 are considered when calculating the 2D Markov random field. Source and target cells are set to spatially
dependent on each other, while the target cells have a greater spatial autocorrelation. The single-source
scenario simulated one pair of source and target cells and two control cell types, while the multi-source
having two sources, one target, and two control cell types. The Pearson and Spearman correlation, and
SPER scores are respectively calculated on three simulations.

We repeated each scenario 10 times given the random seed from 1 to 10. The resulting matrices from all
repeats are used to predict the spatial dependence between genes and cell types, particularly the target cells
and source genes. For each type, the ROC curve is plotted in distinguishing its designed paracrine signals
from the other genes.

Collection and pre-processing of scRNA-seq reference data. The scRNA-seq data were collected and
pre-processed by Tasic, B., et al. in the Allen Institute for Brain Science, generated by the SMART-seq2
protocol [33]. The dataset contains 14249 adult mouse brain cortical cells and 34617 genes. The raw data
was normalized by Seurat using SCTransform function [47], where 3000 subsampling cells were used to
build the NB regression model and other parameters set as the default. The label 'subclass' was used as the
annotation of cell type, which included 23 different subclass of mouse brain cells. Cajal–Retzius (CR) cell
type, the count of which was only 7 in the scRNA-seq data, was removed from the data as the low cell
counts weakened its statistical power. After this removal, 14242 cells with 22 distinct cell types were kept
as the reference in the downstream analysis.

Identifying cell-type-specific marker genes in scRNA-seq data. The cell-type-specific marker genes
were found in the normalized scRNA-seq data by Seurat. For each cell type, their marker genes were then
identified by Seurat using MAST one-versus-all test [34]. In these tests, each gene was tested whether it
was differentially expressed in each cell type compared to rest of cells. Genes are selected as marker genes
by the following criteria: fold change > 2, adjusted p-value < 10^{-30}, fraction of expressing cells > 70%. All
the other parameters were set as the default in the 'FindMarkers' function in the Seurat package [47].

Collection and pre-processing of Visium spatial transcriptomics data in mouse brain. The mouse brain spatial transcriptomics data is collected from the 10X Genomics Visium public database [9]. It contains 2696 spots with 124367 mean reads per spot. The raw count matrix is normalized by Seurat using SCTransform function with default settings. The physical distances of cells are calculated from the provided spatial coordinates and the scale factor in the metadata. In the downstream analysis where ST and scRNA-seq data are needed, we only kept the genes detected in both datasets and having the total UMI counts greater than 40 in the normalized ST data.

Cell-type deconvolution and co-kriging modeling in Visium mouse brain data. In the Visium mouse brain example, we used the normalized ST data to perform the cell-type deconvolution. The cell-type spatial compositional data C is estimated using RCTD full mode [11] with a reference using the scRNA-seq reference data. We select top 10 cell types which have the highest prevalence among all spots in the compositional data generated by RCTD. They are L2/3 IT, L5 IT, Meis2, L4, L6 IT, Oligo, Pvalb, L6 CT, Sst, and L6b. The other cell types are summed together as 'Other' aside from those 10 types. The cell-type spatial compositional matrix is further denoised using the co-kriging algorithms. 15 distance bins, 120 μm each, are used to compute the log-ratio variograms. To model the variograms between cell types, we use a compositional linear model of spatial coregionalization that includes a nugget effect term and 3 Gaussian variogram terms whose respective effective as 600, 1300, and 3000 micrometers. These 3 terms respectively represent the short-range, middle-range, and long-range interactions between cell types.

SPER calculation. In the Visium mouse brain and human lung example, we used the normalized ST data and cell-type spatial compositional data as the SPER inputs. Calculation was performed using SPER R package. The weight function is derived from a Poisson distribution (λ=2.5) to give penalty on overlapping and distal regions while keeping its sum as 1, where the Poisson parameter 2.5 represents 180 nm in reality as the assumed average functional distance of paracrine signals. The choice of weight function is not unique, as users can manually fine-tune it to match the desired functional distance of the signals.

The final SPER score $\tilde{s}_{i,j}$ was calculated as:

$$\tilde{s}_{i,j} = \frac{s_{i,j}}{\exp{(p_{i,j} - 1)}}$$

Where $s_{i,j}$ stands for the raw SPER score and $p_{i,j}$ for the expression prevalence of gene i and cell type j.

Collection of extracellular gene set and ligand-receptor pair data. To identify the subcellular locations, we used the all channels integrated version of mouse extracellular gene set that includes knowledge, experiments, text mining, and predictions channel from the COMPARTMENTS database [48]. To filter out the intracellular genes, we select the genes whose locations are in the extracellular region (including extracellular space, matrix, exosome, and vesicle). Also, we only kept the genes whose confident level greater than 3. As a result, 4797 piece of genes' information and 2622 different genes are left as the set of extracellular genes. After doing the intersection with the ST data, 1784 genes are included in the final extracellular gene set. The ligand-receptor interaction dataset is collected from FANTOM5 ligand-receptor datasets [49] and CellPhoneDB database [50]. Since all these databases only have the human genome data, we transformed the gene ID into the homologous gene in mouse genome to compare with the ST data we have got. There are totally 1888 ligand-receptor pairs that have the homologous genes in mouse, where 1636 of them are from FANTOM5 and 252 of them from CellPhoneDB.

Calculating correlations and EMD of gene-cell type pairs. The Pearson and Spearman correlations are calculated between the spatial compositional data after co-kriging modeled and the spatial transcriptomics data. The same matrices, but size reduced, are used to calculate the EMD in comparing the computing complexity. We use the physical distance between spots as the ground distances, and the proportions of each cell-type composition are inputted as the weights when comparing to a gene's spatial distribution. R package 'emdist' is used to calculate EMD in the simulations.

Spatial-dependent signal detection. The SPER score matrix, Pearson correlation matrix and Spearman correlation matrix are used to filter the candidate genes and spatial-dependent signals with the help of scRNA-seq data as the reference of expression levels. When studying the enrichment of extracellular and ligand genes, we used the hypergeometric test to see if they are enriched within the gene set given thresholds. In both cases, the population size is 13436; the number of successes in the population is 1784 for the extracellular and 1888 for ligands. The sample size and the number of successes in the sample are dependent on the expression-level and measurement-level thresholds. If the p-values calculated from the tests are less than 0.05, we will take it as significantly enriched.

Ablation study of co-kriging. We conducted an ablation sensitivity analysis to explore the contribution of co-kriging in modeling the cell type compositional data and SPER's sensitivity. Following the same procedure as previously (**Fig. 4**), we applied SPER directly on the raw compositional data estimates generated from RCTD, with and without co-kriging to stabilize and model the compositional estimates. We used both statistical enrichment of extracellular cell location and participation as a known paracrine ligand as metrics to benchmark the performance with and without co-kriging (**Fig. S4**). We reasoned that after co-kriging modeling, SPER would be more likely to detect transcripts meeting these criteria, because of the ability of co-kriging to remove noise in compositional estimates. Consistent with this hypothesis, we observed that co-kriging led to an increase in the number of detected transcripts across most cell types, with 9 out of 10 cell types exhibiting a higher number of transcripts with extracellular localization and 8 out of 10 showing an increase in putative paracrine ligands.

Acknowledgements
The authors thank Leslie Gaffney for preparing and generation of figures for this article, Evan Lemire for programming assistance, and Yu Sun for helping improve the data visualization.

Author contributions
T.Z. designed and implemented the SPER algorithm, performed data analysis and visualization and wrote the manuscript. A.H. conceived the study and wrote the manuscript.

Declaration of interests
The authors declare no competing interests.

Data and Code availability
An R package to implement the SPER algorithm is available for download at github.com/TianxiaoNYU/SPER. Extracellular gene set in mouse is available from the

COMPARTMENTS database: https://compartments.jensenlab.org/Downloads; scRNA-seq reference data is obtained from Allen Institute for Brain Science: https://portal.brain-map.org/atlases-and-data/rnaseq; Visium spatial transcriptomics data from mouse brain is available from 10X Genomics dataset: https://www.10xgenomics.com/resources/datasets/mouse-brain-serial-section-1-sagittal-anterior-1-standard-1-0-0. An R markdown tutorial to help generate the main results of this article using SPER package, as well as all other necessary data files in this study can be found at: https://www.dropbox.com/scl/fo/pep07jjid71rr72voeeik/h?rlkey=2agyojo1ugf5l61bqtal6hjy0&dl=0.

REFERENCES:
1.   Meizlish ML, Franklin RA, Zhou X, Medzhitov R. Tissue Homeostasis and Inflammation. Annu Rev Immunol. 2021;39:557-81.
2.   Anthony RM, Rutitzky LI, Urban JF, Stadecker MJ, Gause WC. Protective immune mechanisms in helminth infection. Nat Rev Immunol. 2007;7(12):975-87.
3.   Johnson TS, Xiang S, Dong T, Huang Z, Cheng M, Wang T, et al. Combinatorial analyses reveal cellular composition changes have different impacts on transcriptomic changes of cell type specific genes in Alzheimer's Disease. Sci Rep. 2021;11(1):353.
4.   Pawelec P, Ziemka-Nalecz M, Sypecka J, Zalewska T. The Impact of the CX3CL1/CX3CR1 Axis in Neurological Disorders. Cells. 2020;9(10).
5.   Knoop KA, Newberry RD. Goblet cells: multifaceted players in immunity at mucosal surfaces. Mucosal Immunol. 2018;11(6):1551-7.
6.   Carr EJ, Dooley J, Garcia-Perez JE, Lagou V, Lee JC, Wouters C, et al. The cellular composition of the human immune system is shaped by age and cohabitation. Nat Immunol. 2016;17(4):461-8.
7.   Ansell SM, Vonderheide RH. Cellular composition of the tumor microenvironment. Am Soc Clin Oncol Educ Book. 2013.
8.   Palla G, Fischer DS, Regev A, Theis FJ. Spatial components of molecular tissue biology. Nat Biotechnol. 2022;40(3):308-18.
9.   Genomics X. Mouse Brain Serial Section 1 (Sagittal-Anterior) 2019 [Available from: https://www.10xgenomics.com/resources/datasets/mouse-brain-serial-section-1-sagittal-anterior-1-standard-1-0-0.
10.   Rodriques SG, Stickels RR, Goeva A, Martin CA, Murray E, Vanderburg CR, et al. Slide-seq: A scalable technology for measuring genome-wide expression at high spatial resolution. Science. 2019;363(6434):1463-7.
11.   Cable DM, Murray E, Zou LS, Goeva A, Macosko EZ, Chen F, et al. Robust decomposition of cell type mixtures in spatial transcriptomics. Nat Biotechnol. 2022;40(4):517-26.
12.   Kleshchevnikov V, Shmatko A, Dann E, Aivazidis A, King HW, Li T, et al. Cell2location maps fine-grained cell types in spatial transcriptomics. Nat Biotechnol. 2022;40(5):661-71.
13.   Liu Z, Wu D, Zhai W, Ma L. SONAR enables cell type deconvolution with spatially weighted Poisson-Gamma model for spatial transcriptomics. Nat Commun. 2023;14(1):4727.
14.   Tuong ZK, Loudon KW, Berry B, Richoz N, Jones J, Tan X, et al. Resolving the immune landscape of human prostate at a single-cell level in health and cancer. Cell Rep. 2021;37(12):110132.
15.   Park J, Choi W, Tiesmeyer S, Long B, Borm LE, Garren E, et al. Cell segmentation-free inference of cell types from in situ transcriptomics data. Nat Commun. 2021;12(1):3545.
16.   Grisanti Canozo FJ, Zuo Z, Martin JF, Samee MAH. Cell-type modeling in spatial transcriptomics data elucidates spatially variable colocalization and communication between cell-types in mouse brain. Cell Syst. 2022;13(1):58-70 e5.
17.   Baker EAG, Schapiro D, Dumitrascu B, Vickovic S, Regev A. In silico tissue generation and power analysis for spatial omics. Nat Methods. 2023;20(3):424-31.
18.   Tolosana-Delgado R, Mueller U, van den Boogaart KG. Geostatistics for Compositional Data: An Overview. Math Geosci. 2019;51(4):485-526.

19.   Tanevski J, Flores ROR, Gabor A, Schapiro D, Saez-Rodriguez J. Explainable multiview
framework for dissecting spatial relationships from highly multiplexed data. Genome Biol. 2022;23(1):97.
20.   Li Z, Wang T, Liu P, Huang Y. SpatialDM for rapid identification of spatially co-expressed ligand-
receptor and revealing cell-cell communication patterns. Nat Commun. 2023;14(1):3995.
21.   Shao X, Li C, Yang H, Lu X, Liao J, Qian J, et al. Knowledge-graph-based cell-cell communication
inference for spatially resolved transcriptomic data with SpaTalk. Nat Commun. 2022;13(1):4429.
22.   Ghazanfar S, Lin Y, Su X, Lin DM, Patrick E, Han ZG, et al. Investigating higher-order interactions
in single-cell data with scHOT. Nat Methods. 2020;17(8):799-806.
23.   Bernstein MN, Ni Z, Prasad A, Brown J, Mohanty C, Stewart R, et al. SpatialCorr identifies gene
sets with spatially varying correlation structure. Cell Rep Methods. 2022;2(12):100369.
24.   Satoh A. Chapter 10 - Typical Properties of Colloidal Dispersions Calculable by Molecular-
Microsimulations. In: Satoh A, editor. Studies in Interface Science. 17: Elsevier; 2003. p. 153-9.
25.   Myers DE. Matrix Formulation of Co-Kriging. J Int Ass Math Geol. 1982;14(3):249-57.
26.   Francis K, Palsson BO. Effective intercellular communication distances are determined by the
relative time constants for cyto/chemokine secretion and diffusion. Proc Natl Acad Sci U S A.
1997;94(23):12258-62.
27.   Krige DG. A Statistical Approach to Some Basic Mine Valuation Problems on the Witwatersrand.
J S Afr I Min Metall. 1994;94(3):95-111.
28.   Rubner Y, Tomasi C, Guibas LJ. A metric for distributions with applications to image databases.
Sixth International Conference on Computer Vision. 1998:59-66.
29.   Freguglia V, Garcia NL. Inference Tools for Markov Random Fields on Lattices: The R Package
mrf2d. Journal of Statistical Software. 2022;101(8):1 - 36.
30.   Sang-Aram C, Browaeys R, Seurinck R, Saeys Y. Spotless, a reproducible pipeline for
benchmarking cell type deconvolution in spatial transcriptomics. Elife. 2024;12.
31.   Yan L, Sun X. Benchmarking and integration of methods for deconvoluting spatial transcriptomic
data. Bioinformatics. 2023;39(1).
32.   Li H, Zhou J, Li Z, Chen S, Liao X, Zhang B, et al. A comprehensive benchmarking with practical
guidelines for cellular deconvolution of spatial transcriptomics. Nat Commun. 2023;14(1):1548.
33.   Tasic B, Menon V, Nguyen TN, Kim TK, Jarsky T, Yao ZZ, et al. Adult mouse cortical cell
taxonomy revealed by single cell transcriptomics. Nature Neuroscience. 2016;19(2):335-+.
34.   Finak G, McDavid A, Yajima M, Deng JY, Gersuk V, Shalek AK, et al. MAST: a flexible statistical
framework for assessing transcriptional changes and characterizing heterogeneity in single-cell RNA
sequencing data. Genome Biology. 2015;16.
35.   Frazer S, Prados J, Niquille M, Cadilhac C, Markopoulos F, Gomez L, et al. Transcriptomic and
anatomic parcellation of 5-HT3AR expressing cortical interneuron subtypes revealed by single-cell RNA
sequencing. Nature Communications. 2017;8(1):14219.
36.   Ito-Ishida A, Okabe S, Yuzaki M. The role of Cbln1 on Purkinje cell synapse formation. Neurosci
Res. 2014;83:64-8.
37.   Han P, She Y, Yang Z, Zhuang M, Wang Q, Luo X, et al. Cbln1 regulates axon growth and guidance
in multiple neural regions. PLoS Biol. 2022;20(11):e3001853.
38.   Barker N, van Es JH, Kuipers J, Kujala P, van den Born M, Cozijnsen M, et al. Identification of
stem cells in small intestine and colon by marker gene Lgr5. Nature. 2007;449(7165):1003-7.
39.   Madissoon E, Oliver AJ, Kleshchevnikov V, Wilbrey-Clark A, Polanski K, Richoz N, et al. A
spatially resolved atlas of the human lung characterizes a gland-associated immune niche. Nat Genet.
2023;55(1):66-77.
40.   Blackburn JB, Li NF, Bartlett NW, Richmond BW. An update in club cell biology and its potential
relevance to chronic obstructive pulmonary disease. Am J Physiol Lung Cell Mol Physiol.
2023;324(5):L652-L65.
41.   Rodriguez LR, Emblom-Callahan M, Chhina M, Bui S, Aljeburry B, Tran LH, et al. Global Gene
Expression Analysis in an in vitro Fibroblast Model of Idiopathic Pulmonary Fibrosis Reveals Potential
Role for CXCL14/CXCR4. Sci Rep. 2018;8(1):3983.

42. Li L, Huang J, Liu Y. The extracellular matrix glycoprotein fibrillin-1 in health and disease. Front
Cell Dev Biol. 2023;11:1302285.
43. Hoffman ET, Uriarte JJ, Uhl FE, Eckstrom K, Tanneberger AE, Becker C, et al. Human alveolar
hydrogels promote morphological and transcriptional differentiation in iPSC-derived alveolar type 2
epithelial cells. Sci Rep. 2023;13(1):12057.
44. Tun MH, Borg B, Godfrey M, Hadley-Miller N, Chan ED. Respiratory manifestations of Marfan
syndrome: a narrative review. J Thorac Dis. 2021;13(10):6012-25.
45. Lee Y, Bogdanoff D, Wang Y, Hartoularos GC, Woo JM, Mowery CT, et al. XYZeq: Spatially
resolved single-cell RNA sequencing reveals expression heterogeneity in the tumor microenvironment. Sci
Adv. 2021;7(17).
46. Payares-Garcia D, Osei F, Mateu J, Stein A. Multivariate Poisson cokriging: A geostatistical model
for health count data. Stat Methods Med Res. 2024;33(9):1637-59.
47. Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, 3rd, et al. Comprehensive
Integration of Single-Cell Data. Cell. 2019;177(7):1888-902 e21.
48. Binder JX, Pletscher-Frankild S, Tsafou K, Stolte C, O'Donoghue SI, Schneider R, et al.
COMPARTMENTS: unification and visualization of protein subcellular localization evidence. Database-
Oxford. 2014.
49. Ramilowski JA, Goldberg T, Harshbarger J, Kloppman E, Lizio M, Satagopam VP, et al. A draft
network of ligand-receptor-mediated multicellular signalling in human. Nature Communications. 2015;6.
50. Efremova M, Vento-Tormo M, Teichmann SA, Vento-Tormo R. CellPhoneDB: inferring cell-cell
communication from combined expression of multi-subunit ligand-receptor complexes. Nat Protoc.
2020;15(4):1484-506.
713
714
715
716

717 SUPPLEMENTARY FIGURES:

718
719 **Supplementary Figure 1. Co-kriging enables SPER to effectively denoise spatial compositional data**. Spatial distribution of
720 composition of primary cell types decomposed by RCTD in full mode before (A) and after co-kriging modeling (B). The top ten
721 most abundant cell types are modeled individually, and the other 11 cell types are summed as the 'Other' category. In each
722 subplot, X and Y-axis represent the physical coordinates of spots, and the color represents the normalized proportion of the
723 relevant cell type.

724
725

726
Supplementary Figure 2. Co-kriging fits visualized as variograms. Co-kriging method fits on the ten neuronal cell types from
an example dataset from mouse brain. A compositional linear model of co-regionalization (red line) with a 'nugget' effect term
for short scale randomness and three Gaussian variogram terms to model effects at different ranges (600, 1300, and 3000 μm)
showed excellent fit to the original covariance data (black line) collected and calculated at 15 uniformly distributed lags. The
labels at the diagonal identify the cell type for each row and column, while 'Other' represents the sum of all remaining cell types.
The x-axis in each subpanel represents the physical distance in μm, and y-axis represents the relative value of the covariance
between each pair of cell types.
734

735
736 **Supplementary Figure 3. Pearson and Spearman correlations detect marker genes with overlapping spatial dependency.**
737 Boxplots showing Pearson and Spearman correlations between cell type spatial compositions and gene's spatial distribution.
738 Marker genes for each cell types are selected from the scRNA-seq reference data by MAST. The four-star labels represent the
739 difference (adjusted p-value $< 10^{-4}$) in the measurement scores between the marker and non-marker groups.
740

Supplementary Figure 4. Ablation study demonstrates SPER detect more paracrine ligand genes with co-kriging. Dot plots visualize hypergeometric enrichment analysis of known paracrine ligand signals with and without co-kriging analysis. Ligand genes annotations were collected from COMPARTMENTS, FANTOM5, and CellPhoneDB databases respectively (see **Methods**). The candidate sets include genes whose scores for each spatial dependency metric (x-axis) are above than the 95% percentile of all genes. The size of dots represents the number of extracellular or ligand genes in the given gene set. The color represents the significance the enrichment in the set (hypergeometric test), non-significant ($p>0.05$) overlaps are shown as light grey. With co-kriging, the number of transcripts in both categories increased for most cell types, with 9 out of 10 showing an increase for extracellular location and 8 out of 10 for paracrine ligands.

752
753 **Supplementary Figure 5. Comparison of trivial spatial associations and genuine putative regulators between SPER and**
754 **Pearson correlation.** Scatter plots show the distribution of SPER (A) and Pearson correlation (B) scores (x axis) with the
755 proportion of cells of each type (plots) in which each gene (points) is detected (y axis) in scRNA-seq data. The dark blue dots mark
756 the marker genes for each cell type (found by MAST, details in Methods). Red dots stand for the genes whose measurement scores
757 larger than 95% percentile. Purple dots represent genes that meets both conditions.
758