mTORC1/S6K1 signaling promotes sustained oncogenic translation through
modulating CRL3[IBTK]-mediated non-degradative ubiquitination of eIF4A1

Running Title: CRL3[IBTK] promotes eIF4A1-related oncogenic translation

Huiru Sun[1,4], Dongyue Jiao[1,4], Xiaying Zhao[1], Yingji Chen[1], Qing Shi[1], Zeheng Lv[2], Yao
Li[1], Kun Gao[2,3,*], and Chenji Wang[1,*]

[1]Shanghai Stomatological Hospital & School of Stomatology, State Key Laboratory of
Genetic Engineering, MOE Engineering Research Center of Gene Technology,
Shanghai Engineering Research Center of Industrial Microorganisms, School of Life
Sciences, Fudan University, Shanghai 200438, China

[2]Department of Clinical Laboratory, Shanghai First Maternity and Infant Hospital,
School of Medicine, Tongji University, Shanghai 200092, China

[3]Shanghai Key Laboratory of Maternal and Fetal Medicine, Shanghai First Maternity
and Infant Hospital, Shanghai 200092, China

[4]These authors contributed equally

*Correspondence: chenjiwang@fudan.edu.cn (C.W); kungao@tongji.edu.cn. (K.G)

Abstract

Enhanced protein synthesis is a crucial molecular mechanism that allows cancer cells to survive, proliferate, metastasize, and develop resistance to anti-cancer treatments, and often arises as a consequence of increased signaling flux channeled to mRNA-bearing eukaryotic initiation factor 4F (eIF4F). However, the post-translational regulation of eIF4A1, an ATP-dependent RNA helicase and subunit of the eIF4F complex, is still poorly understood. Here, we demonstrate that IBTK, a substrate-binding adaptor of Culllin 3-RING ubiquitin ligase complex (CRL3), interacts with eIF4A1. The non-degradative ubiquitination of eIF4A1 by catalyzed CRL3[IBTK] complex promotes cap-dependent translational initiation, nascent protein synthesis, oncogene expression, and tumor cell growth both *in vivo* and *in vitro*. Moreover, our results show that mTORC1 and S6K1, two key regulators of protein synthesis, directly phosphorylate IBTK to augment eIF4A1 ubiquitination and sustained oncogenic translation. This link between the CRL3[IBTK] complex and the mTOR signaling pathway, frequently dysregulated in cancer, represents a promising target for anticancer therapies.

Statement of Significance: IBTK overexpression contributes to cervical cancer tumorigenesis by translation regulation and represents a promising target for anticancer therapies.

Keywords: Tumorigenesis, Ubiquitination, Phosphorylation, mTOR signaling

1. Introduction

Enhanced protein synthesis is a critical process that allows cancer cells to survive, multiply, metastasize, and resist anti-cancer treatments(1, 2). This process typically results from increased signaling through the eukaryotic initiation factor 4F (eIF4F), which binds to mRNAs. eIF4F complex is composed of four proteins: eIF4E, eIF4G, eIF4B, and eIF4A. Of these, eIF4A is responsible for unwinding the secondary structure in the 5'-UTR region of the mRNA, allowing the ribosomal subunit to bind and begin translation(3). The eIF4A protein family consists of three paralogs, eIF4A1, eIF4A2, and eIF4A3. Although eIF4A1 and eIF4A2 have similar roles in translation initiation, eIF4A3 is not involved in translation control but an important component of the exon junction complex (EJC) and plays a critical role in nonsense-mediated mRNA decay(4). Increased eIF4F activity promotes the translation of mRNAs involved in cell proliferation and survival, and tumor immune evasion, which are hallmarks of cancer(5-8). Several eIF4A inhibitors, such as eFT226, silvestrol, hippuristanol, and pateamine A, have demonstrated promising anti-tumor activity in various studies conducted *in vitro* and *in vivo*(5-7). In particular, eFT226 is currently being evaluated in multiple clinical trials(9).

The mTORC1 signaling pathway plays a crucial role in promoting cell growth and anabolism while inhibiting catabolism. It integrates various signals from nutrients, cellular energy levels, growth factors, and environmental stimuli to regulate this process. Translation is a highly energy-intensive process and tightly regulated by the mTORC1 signaling. Ribosomal protein S6 kinases (S6Ks) and 4E-binding proteins (4E-BPs) are two major downstream targets of mTORC1 signaling. They regulate various aspects of mRNA translation in a phosphorylation-dependent manner. For example, S6Ks phosphorylate ribosomal protein S6, which promotes translation initiation and elongation. 4E-BPs, on the other hand, bind to eIF4E and inhibit its activity, preventing the translation of specific mRNAs. In addition, mTORC1 coordinates translation by phosphorylating other components of the translational machinery, such as eEF2K, eIF2B, and LARP1(10).

The Cullin3-RING E3 (CRL3) ubiquitin ligase complex subfamily is composed of a

catalytic core made up of RBX1 and CUL3, and a substrate-binding adaptor that

contains an interchangeable BTB domain. The CRL3 complex plays a role in regulating

various cellular processes, including cell division, differentiation, and signaling.

Through its E3 ubiquitin ligase activity, the CRL3 complex promotes the degradation

of specific proteins by marking them with ubiquitin. However, the CRL3 complex can

also facilitate non-degradative ubiquitination, which can modulate protein activity,

localization, or interaction(11). Initially characterized as an inhibitor of Bruton's tyrosine

kinase (BTK), the Inhibitor of Bruton's tyrosine kinase (IBTK) contains two BTB

domains and can assemble an active CRL3 complex(12, 13). The downstream targets

and biological processes of the CRL3^{IBTK} complex are still not well understood, but a

limited number of studies suggest that IBTK plays a pro-survival role in cells. IBTK is

preferentially translated in response to eIF2α phosphorylation during endoplasmic

reticulum stress. Knockdown of IBTK expression reduced cell viability and increased

apoptosis in cultured cells(14, 15). A genome-wide RNAi screen identified IBTK as a

synthetic lethal partner of the Ras oncogene in colorectal cancer cells(16). In Eμ-myc

transgenic mice, Ibtk gene knockout delayed the onset of pre-B/B lymphoma and

improved animal survival due to increased apoptosis of pre-cancerous B cells(17). The

expression of IBTK was significantly increased in chronic lymphocytic leukemia (CLL)

progression, but decreased in remission following chemotherapy. Knockdown of IBTK

expression increased spontaneous and chemotherapy agent-induced apoptosis and

impaired cell cycle progression in CLL cell lines(18). To date, only two substrates of

CRL3^{IBTK} have been reported. CRL3^{IBTK} targets PDCD4, an eIF4A1 inhibitor, for

proteasomal degradation(13). Another study showed that CRL3^{IBTK} activates β-

Catenin-dependent transcription of MYC by promoting GSK-3β degradation in

cancerous B cells(19). These findings suggest that CRL3^{IBTK} may play a crucial role in

cancer development and progression, as well as in the regulation of cell survival and

apoptosis.

Through a variety of biochemical and cell biology approaches, we have identified

eIF4A1 as a non-degradative ubiquitinated substrate of the CRL3^{IBTK} complex. Our findings suggest that a mTORC1/S6K1-IBTK-eIF4A1 signaling axis mediates cap-dependent translation, facilitates oncoprotein expression, and promotes tumor cell growth.

Methods and Materials

Cell lines and lentiviral infection

SiHa, HeLa, 293T, H1299, and CT26 cells were acquired from American Type Culture Collection (ATCC). All cell lines were cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS) and 1% penicillin/streptomycin at 37°C with 5% CO_2. We routinely perform DNA fingerprinting and PCR to verify the authenticity of the cell lines and to ensure they are free of mycoplasma infection. We conducted transient transfection using EZ Trans (Shanghai Life-iLab Biotech). For lentiviral transfection, we transfected pLKO shRNA knockdown or pCDH overexpression plasmids and virus packing constructs into 293T cells. The viral supernatant was collected after 48 h. SiHa or HeLa cells were then infected with the viral supernatant in the presence of polybrene (8 μg/ml) and selected in growth media containing puromycin (1.5 μg/ml). The gene-specific shRNA sequences can be found in **Supplementary Table 6.**

Antibodies, recombinant proteins, and chemicals

The information of antibodies, recombinant proteins, and chemicals used in this study is listed in **Supplementary Table 7, 8**.

Gene KO cell line generation

The sgRNAs targeting the IBTK gene were designed using an online CRISPR design tool (http://crispr.mit.edu) and subcloned into the pX459 construct from Dr. Feng Zhang's lab. SiHa, 293T, and H1299 cells were plated and transfected with pX459 constructs overnight. After 24 h of transfection, cells were exposed to puromycin (1.5

µg/ml) for one week. The surviving cells were then seeded in a 96-well plate by limited

dilution to isolate monoclonal cell lines. After ten days, KO cells were screened using

WB analysis and validated through Sanger sequencing. **Supplementary Table 6** lists

the sequences of gene-specific sgRNAs.

Doxycycline inducible expression and protein complex purification

To generate stable cell lines with inducible IBTK gene expression, Flp-In T-REx 293

cells were co-transfected with pOG44 and pcDNA5-FLAG-BirA*-IBTK constructs. After

two days of transfection, the cells were selected with hygromycin (100 µg/ml) for 2

weeks, and then the positive clones were pooled and amplified. To induce exogenous

IBTK expression, doxycycline (10 ng/ml) was added to the stable cell lines. For

purification in the AP-MS pipeline, the sample was lysed in 3 ml (for each plate) of NP-

40 lysis buffer containing fresh protease inhibitor and kept on ice for 2 h. The

homogenate was centrifuged for 30 minutes at 12000 rpm at 4°C. Cleared lysates were

filtered through 0.45 µM spin filters (Millipore) and immunoprecipitated with anti-FLAG

antibody-conjugated M2 agarose (Sigma). The bound polypeptides were eluted with

the FLAG peptide (Sigma). For BioID approach, the cell pellet was thawed in 3 ml of

BC100 lysis buffer (20 mM Tris-Cl, pH 7.9, 100 mM NaCl, 0.2 mM EDTA, 20% glycerol)

containing fresh protease inhibitor and kept on ice for 2 h. The homogenate was

centrifuged for 30 minutes at 12000 rpm at 4°C. Cleared lysates were filtered through

0.45 µM spin filters (Millipore) and immunoprecipitated with Strep-Tactin beads (IBA

lifescience). The bound polypeptides were eluted with 50 nM biotin. Finally, the eluates

were resolved by SDS-PAGE for Coomassie Blue staining. Gel bands were cut out

and subjected to mass-spectrometric sequencing.

Protein half-life assays

To measure the protein half-life, we added cycloheximide (CHX, 100 µg/ml) to the

media of parental and IBTK-KO SiHa cells. At specific time points thereafter, we

prepared whole cell lysates (WCLs) and detected them through WB analysis using the

specified antibodies.

***In vivo* ubiquitination assays**

We transfected 293T cells with HA-ubiquitin and other specified constructs. After 36 h,

we harvested the cells and lysed them in a 1% SDS buffer (20 mM Tris-Cl (pH 7.4),

0.5 mM EDTA, and 1 mM DTT). The lysate was then boiled at 95 °C for 10 minutes.

Next, we added Strep-Tactin beads (IBA lifescience) and a 10-fold volume of lysis

buffer (20 mM Tris-Cl (pH 7.4), 100 mM NaCl, 0.2 mM EDTA, 0.5% NP-40, and 1×

protease inhibitor cocktail) to the lysate, and incubated it overnight at 4 °C with shaking.

The resulting mixture was washed four times with BC100 buffer and eluted with Biotin

(50 mM) for 1 hour at 4 °C. Finally, we detected the ubiquitinated substrates through

WB analysis using an anti-HA antibody.

***In vitro* ubiquitination assays**

In vitro ubiquitination assays were performed using a protocol reported previously with

some modifications(35). Briefly, 2 µg of APP-BP1/Uba3, 2 µg of His-UBE2M, and 5 µg

of NEDD8 were incubated at 30 °C for 2 h in the presence of ATP. The thioester-loaded

His-UBE2M–NEDD8 was further incubated with 3 µg of His-DCNL2 and 6 µg of CUL3–

RBX1 at 4 °C for 2 h to obtain neddylated CUL3–RBX1. The neddylated CUL3–RBX1,

5 µg of GST-IBTK$_{900\text{-}1150aa}$, 5 µg of ubiquitin, 500 ng of E1 enzyme, 750 ng of E2

enzyme (UbcH5a and UbcH5b), and 5 µg of GST-eIF4A1 were incubated with 0.6 µl

of 100 mM ATP, 1.5 µl of 20 µM ubiquitin, 3 µl of 10× ubiquitin reaction buffer (500 mM

Tris-HCl (pH 7.5), 50 mM KCl, 50 mM NaF, 50 mM MgCl2, and 5 mM DTT), 3 µl of

10× energy regeneration mix (200 mM creatine phosphate and 2 µg/µl creatine

phosphokinase), and 3 µl of 10× protease inhibitor cocktail at 30 °C for 2 h, followed

by WB analysis.

IF and confocal microscopy

For IF, SiHa cells were seeded on glass coverslips in 24-well plates and harvested at

80% confluence. The cells were washed with PBS and fixed with 4% paraformaldehyde (PFA) in PBS. After permeabilization with 0.4% Triton X-100 for 10 min and then in the blocking solution (PBS plus 5% donkey serum), for 30 min at room temperature (RT). The cells were then incubated with primary antibodies at 4 °C overnight. After washing with PBST buffer, fluorescence-labelled secondary antibodies were applied. DAPI was utilized to stain nuclei. The glass coverslips were mounted on slides and imaged using a confocal microscope (LSM880, Zeiss) with a 63*/1.4NA Oil PSF Objective. Quantitative analyses were performed using ImageJ software.

For xenograft tumor tissue staining, the tumor tissues were isolated from mice after perfusion with 0.1 M PBS (pH7 .4) and fixed for 3 days with 4% PFA at 4 °C. The tumor tissues were then placed in 30% sucrose solution for 2 days for dehydration. The tumors were embedded into the OCT block and frozen for cryostat sectioning. Cryostat sections (45-μm thick) were washed with PBS, and then incubated in blocking solution (PBS containing 10% goat serum, 0.3% Triton X-100, pH7.4) for 2 h at RT. In antibody reaction buffer (PBS plus 10% goat serum, 0.3% Triton X-100, pH7.4), The samples were stained with primary antibodies against active CD8 (1:100) and Granzyme B (1:200) overnight at 4 °C, followed by Alexa 488 and 647 secondary antibodies (1;2000) at RT for 3 h. DAPI was used for nuclear staining. The sections were then sealed with an anti-fluorescence quencher. The samples were visualized and imaged using a confocal microscope (FV3000, Olympus) along the z-axis with a 40× objective. The number of CD8$^+$ T cells and the area of Granzyme B were quantified using ImageJ by computing corresponding positive staining area. The analysis was performed in eight different units.

Measurement of protein synthesis

We used puromycin labeling to measure nascent protein synthesis. The cells were treated with puromycin (1.5 μM) for 10 minutes before being lysed by 1× SDS sample buffer.   The samples were analyzed by WB using anti-puromycin antibody.

Pulldown assays using m⁷GTP-Sepharose

SiHa cells were washed in PBS and lysed in 1 ml NP-40 (0.1% NP-40) supplemented with protease inhibitor. For each sample, WCLs were incubated with 50 μl 7-methyl-GTP Sepharose 4B beads (GE Healthcare) for 2 h at 4 °C. Then, the resins were washed five times with BC100 buffer, and the proteins bound to the washed Sepharose resins w by using 1× SDS sample buffer and subjected them to SDS-PAGE followed by WB analysis with the indicated antibodies.

Dual-luciferase assays

The bicistronic reporters, pH and pE, were kindly provided by Dr. Peter Sarnow (Stanford University). The cells were seeded onto 24-well plates and co-transfected with each reporter plasmid. After 24 h of transfection, the cells were harvested in passive lysis buffer and immediately quick-frozen at -80°C to fully lyse them. Dual-luciferase assays were performed using a dual-luciferase reporter assay system following the manufacturer's instructions (Promega). The luciferase activity was measured using an Envision Multilabel Plate Reader (PerkinElmer). Three independent experiments were performed.

RNA isolation and quantitative real-time reverse transcription PCR (qRT-qPCR)

Total RNAs were isolated from cells using the TRIzol reagent (Thermo) following the manufacturer's instructions. Concentrations and purity of RNAs were determined by measuring the absorption of ultra-violet lights using a NanoDrop spectrophotometer (Thermo). cDNAs were reversed-transcribed using a HiScript III 1st Strand cDNA Synthesis Kit (Vazyme), followed by amplification of cDNA using ChamQ SYBR qPCR Master Mix (Vazyme). The relative mRNA levels of genes were quantified using the $2^{-\Delta\Delta Ct}$ method, with normalization to Actin. The primer sequences are listed in **Supplementary Table 6**.

CCK-8 assays

The cell proliferation rates of HeLa or SiHa were determined using the Cell Counting Kit-8 (CCK-8) Kit (Beyotime) according to the manufacturer's instructions. Briefly, cells were seeded onto 96-well plates at a density of 1000 cells per well in triplicate. During a 2 to 6 or 7-day culture period, 10 µl of the CCK-8 solution was added to cell culture, and incubated for 2 h. Then samples were measured at an absorbance of 450 nm using a microplate absorbance reader (Bio-Rad).

Colony formation assays

HeLa or SiHa cells were seeded in 6-well plates containing 1000 individual cells per well in triplicate. After incubating for 2 weeks, cells were fixed with 4% PFA for 15 min at 37 °C and then subjected to Giemsa dye (Solarbio) staining for 20 min. Then, the cells were washed with water by dropping gently, and air dried at RT. The number of colonies were photographed using a digital photo camera (Nikon) and quantified using ImageJ.

Migration and invasion assays

Cell migration and invasion were determined using Transwell chamber (Corning). HeLa or SiHa cells were pre-cultured in serum-free media for 24 h. In the case of migration assays, 2×10^5 cells were seeded in serum-free media in the upper chamber, and the lower chamber was filled with DMEM containing 5% FBS. After 48 h, the cells were fixed with 4% PFA for 15 min at 37 °C followed by staining with Giemsa dye (Solarbio) for 20 min. The non-migrating cells on the upper chambers were carefully removed with a cotton swab and migrated cells on the underside of the filter were observed and counted in three different fields. The protocol for invasion assays is similar to the migration assays, with the exception that Matrigel (Corning) is added to the upper chambers before seeding the cells. Three independent experiments were conducted.

Sphere formation assays

SiHa or HeLa cells suspension (2×10^5 cells/well) were mixed with Matrigel and then

plated in 24-well ultra-low attachment plates in DMEM containing 10% FBS. Fresh

media were added every 3 days. The floating spheres that grew in 1-2 weeks were

captured using a digital photo camera, and their number and size were measured using

ImageJ. Three independent experiments were conducted.

Apoptosis detection assays

To measure the apoptosis rates of cells, Annexin V-FITC (Fluorescein isothiocyanate)

and propidium iodide (PI) double staining was employed. SiHa or HeLa cells were

cultured in 6-well plates for 12 hours, followed by the addition of silvestrol (100 nM) for

the specified time period. After that, the suspension and adherent cells were collected

separately and then combined for apoptosis detection using the Apoptosis Detection

Kit from Dojindo. Rocaglamide A (200 nM) was also administered to initiate apoptosis.

All flow cytometric analyses were conducted using a flow cytometer (FACSCalibur, BD

Biosciences). Three independent experiments were performed.

Mouse tumor implantation assays

All experimental protocols (No. TJBG10022102) were approved in advance by the

Ethics Review Committee for Animal Experimentation of Shanghai First Maternity and

Infant Hospital. 4 weeks old BALB/c female mice and athymic nude mice (SLAC

Laboratory) were bred and maintained in our institutional pathogen-free mouse

facilities. Subsequently, 1×10^6 parental or IBTK KD SiHa cells were subcutaneously

(s.c.) injected into female BALB/c mice aged 6-8 weeks. After 20 days of tumor cell

injection, the mice were euthanized, and in vivo solid tumors were excised and

weighed. The growth of tumors was measured every five days in two dimensions using

a digital caliper, with tumor volumes calculated using the ellipsoid volume formula:

$V = (L \times W^2)/2$, where L is the length and W is the width. A portion of tumors was fixed

in formalin and embedded in paraffin for IF analysis. In a similar procedure, 1×10^7

parental and IBTK-KD HeLa cells were s.c. injected into nude mice. After 20 days of

tumor cell injection, the mice were euthanized, and in vivo solid tumors were excised

and weighed.

Recombinant protein production and *in vitro* kinase assays

The pGEX-4T-2 construct containing IBTK$_{900-1150aa}$ was transformed into *E. coli* Rosetta

(DE3) to express the recombinant GST-tagged proteins. These proteins underwent a

two-step purification process on glutathione-agarose beads (Pharmacia), followed by

affinity purification on Amicon Ultra-0.5 Centrifugal Filter Devices (Millipore). IBTK

phosphorylation was assessed using *in vitro* kinase assays. Purified GST-tagged

IBTK$_{900-1150aa}$ at 2 μg was mixed with kinases mTOR/mLST8 or p70S6K1 (Carna

Biosciences) in a reaction mixture that contained 10× kinase buffer (CST), 10mM

ATP(Sigma) for 1 h at 30°C. The reaction was terminated by adding 2× SDS loading

buffer and boiling at 105°C for 5 minutes. SDS-PAGE was used to separate proteins,

with gel bands cut out and subjected to mass-spectrometric sequencing.

IHC for human CESC specimens

All experimental protocols (No. KS2281) were approved in advance by the Ethics

Review Committee of Shanghai First Maternity and Infant Hospital. IBTK protein

expression was detected using the tissue microarray (TMA, HUteS154Su01, Shanghai

Outdo Biotech), which included 152 points comprising 117 CESC tissues and 35

adjacent normal cervical tissues. The TMA slide was first baked at 65 °C for 30 minutes

and then deparaffinized in xylene. The tissue sections were passed through graded

alcohol before being subjected to antigen retrieval with 1 mM EDTA, pH 9.0 (Servicebio)

in a microwave at 50 °C for 10 min and 30 °C for 10 min. Subsequently, the sections

were treated with 3% H_2O_2 for 25 min to quench endogenous peroxidase activity and

washed carefully in phosphate-buffered saline (PBS, pH 7.4) thrice. A solution of 3%

bovine serum albumin was added onto the slide to evenly cover the tissue, which was

then incubated at 37°C for 30 min. Next, the slide was incubated with diluted antibodies

overnight at 4°C. After rinsing with PBS three times, the sections were treated with

horseradish peroxidase-conjugated mouse antibody (Servicebio) for 50 min, followed

by 3,3′-diaminobenzidine incubation. Finally, the slide was counterstained with 0.1%

hematoxylin, dehydrated, and covered before visualization under a confocal

microscope. Each sample was scored based on the intensity of staining (0 = no

staining; 1 = weak staining; 2 = moderate staining; 3 = strong staining) and the

proportion of cells (0=0%; 1=1–25%; 2=25–50%; 3=50–75%; 4=75–100%). All IHC

data were interpreted by the same qualified pathologist for consistency.

Statistical analysis

Kaplan-Meier plots were used to generate survival curves, which displayed *P* values,

fold changes, and ranks. The results of Kaplan-Meier plots were displayed with HR

and P or Cox P values from a log-rank test. Band intensities of WB results were

calculated by ImageJ in accordance with the manufacturer's instructions. Statistical

analysis was performed using GraphPad Prism (GraphPad Software), and the

differences between two groups were analyzed using Student's t-test while the

differences between multiple groups were analyzed using One-way or Two-way

analysis of variance (ANOVA), unless otherwise specified. All data were displayed as

means ± S.D. values for experiments conducted with at least three replicates.

Results

Identification of eIF4A family proteins as IBTK interacting proteins

Affinity purification coupled with mass spectrometry (AP-MS) and proximity-dependent

biotinylation identification (BioID) methods are powerful tools for interrogating protein-

protein interactions(20). Using these techniques, we sought to comprehensively

characterize the interaction partners of IBTK in cells. To accomplish this, we developed

a tag workflow that enabled simultaneous AP-MS and BioID analysis with a single

construct, pcDNA5/FRT vector containing FLAG-BirA*-IBTK. This construct was stably

transfected into Flp-In T-REx 293 cells, resulting in inducible expression of IBTK

(**Supplementary Fig. 1A**). We observed that levels of doxycycline (DOX)-induced

FLAG-BirA*-IBTK protein were comparable to those of endogenous IBTK (**Fig. 1A**). Through our AP-MS and BioID analysis, we identified multiple potential IBTK interactors (**Supplementary Table.1, 2**). One notable finding was the presence of multiple eIFs in the purified IBTK complex, including the subunits of eIF2, eIF3, and eIF4 complex (**Fig. 1B, C, Supplementary Fig. 1B**). Of particular interest, eIF4A1 is a crucial subunit of the eIF4F complex and a promising therapeutic target for cancer. Previous large-scale interactome mapping datasets had suggested a potential interaction between IBTK and eIF4A1, though it had not been analyzed for its biological functions(21). Therefore, we performed additional analyses to explore the pathophysiological significance of the eIF4A1-IBTK interaction.

To validate the interaction between IBTK and eIF4A1, we performed co-immunoprecipitation (co-IP) assays. Our results demonstrated that ectopically expressed IBTK and eIF4A1 interacted with each other (**Fig. 1D**). Moreover, we observed that two eIF4A1 paralogs, eIF4A2 and eIF4A3, also interacted with IBTK (**Fig. 1E, F**). Using FLAG-IBTK as an immunoprecipitation agent, we were able to confirm that endogenous eIF4A1/2/3 and CRL3 subunits (RBX1 and CUL3) associated with IBTK, but not other eIF4 complex subunits (eIF4B, eIF4E, and eIF4G) (**Fig. 1G**). Additionally, we detected endogenous interactions between IBTK and eIF4A1/2/3 (**Fig. 1H-K**). Collectively, these data suggest that IBTK specifically interacts with three members of the eIF4A family in cells.

IBTK mediate non-degradative ubiquitination of eIF4A1

Previous studies have demonstrated that eIF4A1 protein stability can be regulated by the ubiquitin-proteasome pathway, and that deubiquitinase USP9X promotes eIF4A1 stability by facilitating its deubiquitination(22). However, the specific ubiquitin E3 ligases accountable for eIF4A1 proteolysis remain elusive. Given that IBTK is involved in the assembly of a CRL3 ubiquitin ligase complex, we hypothesized that IBTK may target eIF4As for degradation. Nevertheless, overexpressing exogenous IBTK did not influence the levels of endogenous eIF4A1/2/3 (**Fig. 2A**). Moreover, CRISPR/Cas9-

mediated knockout (KO) (**Supplementary Fig. 2A-C**) or shRNA-mediated knockdown

(KD) of IBTK in multiple cell lines did not alter the levels of eIF4A1/2/3 (**Fig. 2B, C**).

Additionally, the protein half-life of eIF4A1/2/3 was unaltered by IBTK depletion

(**Supplementary Fig. 2D, E**). Surprisingly, the protein level of PDCD4, a reported

proteolytic substrate of the CRL3^{IBTK} complex, remained unchanged in IBTK-KO or -

KD cells (**Fig. 2B, C**).

We observed that IBTK-WT co-expression effectively ubiquitinated eIF4A1 and eIF4A2,

but not the ΔBTB mutant (which has a defect in CUL3 binding) (**Fig. 2D**). Consistently,

endogenous eIF4A1/2 ubiquitination was reduced upon IBTK KO (**Fig. 2E**), and we

further demonstrated that the CRL3^{IBTK} complex catalyzed eIF4A1 ubiquitination *in*

vitro (**Fig. 2F**). Intriguingly, neither overexpression nor IBTK KO had any impact on

eIF4A3 ubiquitination (**Fig. 2D, E**), despite a strong interaction between these two

proteins (**Fig. 1J, K**). Notably, eIF4A1 and eIF4A2 share highly similar protein

sequences (>90% identity), but eIF4A1 is generally more abundant than eIF4A2 in

most tissues. Furthermore, eIF4A2 KO did not affect cell viability or global protein

synthesis(23, 24), thus our study primarily focused on the functional consequences of

IBTK regulation of eIF4A1.

After identifying that IBTK-mediated eIF4A1 ubiquitination is non-degradative, we

explored the specificity of polyubiquitin chain linkage on eIF4A1 catalyzed by CRL3^{IBTK}.

To do this, we conducted ubiquitination assays using a panel of ubiquitin mutants with

a single lysine-to-arginine (KR) mutation at seven lysine residues or only one lysine

present with the other six lysine residues mutated to arginine (KO). Additionally, we

included a ubiquitin mutant in which all lysine residues were replaced with arginine (K-

ALL-R). Strikingly, co-expression of any Ub-KR or -KO mutants did not significantly

impact IBTK-mediated eIF4A1 ubiquitination, whereas co-expression of the Ub K-ALL-

R mutant, which is unable to form polyubiquitin chains, formed only a moderate

reduction in IBTK-mediated eIF4A1 ubiquitination (**Fig. 2G**). The C-terminal glycine-

glycine (GG) amino acid residues are essential for Ub conjugation to targeted proteins.

Indeed, co-expression of the Ub-ΔGG mutant abolished IBTK-mediated eIF4A1

ubiquitination (**Fig. 2G**). We then aimed to identify the specific sites on eIF4A1 where ubiquitin was attached. Using MS to analyze the immunoprecipitated eIF4A1-Ub conjugates, we found that eIF4A1 was ubiquitinated at twelve lysine residues (**Supplementary Fig. 2F, Supplementary Table. 3**). By mutating these lysine residues to arginine, we found that IBTK-mediated eIF4A1 ubiquitination was completely terminated (**Supplementary Fig. 2G**), suggesting that at least some of these lysine residues may serve as ubiquitin attachment sites targeted by CRL3^{IBTK}. Collectively, these data suggest that the CRL3^{IBTK} complex mainly catalyzes multi-mono-ubiquitination on eIF4A1, which does not lead to degradation.

IBTK promotes nascent protein synthesis and cap-dependent translation initiation

Given the significance of eIF4A1 in translational regulation, we conducted puromycin incorporation assays to assess the impact of IBTK depletion on global protein synthesis. Our results demonstrated a noticeable decrease in nascent protein synthesis in both IBTK-KO or -KD cells **(Fig. 3A, B)**. However, this effect was reversed by reintroducing IBTK into IBTK-KO cells (**Supplementary Fig. 3A**). Remarkably, the eIF4A1 inhibitor silvestrol also showed a considerable reduction in nascent protein synthesis, particularly in IBTK-KD or KO cells **(Fig. 3A, B)**. A decline in translation initiation is often associated with the formation of cytoplasmic stress granules (SGs), which are aggregates resulting from the phase separation of stalled mRNAs and related factors from the surrounding cytosol(25). We treated cells with sodium arsenite (AS), a conventional SG inducer that produces oxidative stress, and examined SG assembly using eIF4A1 staining as a surrogate. Our data indicated that IBTK-KO cells exhibited significantly higher levels of AS-induced SGs compared to parental cells (**Fig. 3C, D, Supplementary Fig. 3B**). Conversely, overexpression of IBTK-WT resulted in a marked reduction in AS-induced SG assembly, while the ΔBTB mutant had no such effect (**Fig. 3E, F**). In contrast, IBTK overexpression had no influence on P-body assembly, as assessed by EDC4 staining (**Supplementary Fig. 3C**).

We aimed to determine whether IBTK plays a regulatory role in the cap-dependent translation initiation. In mammalian cells, two mechanisms of translation initiation exist: 5' cap-dependent translation initiation and internal ribosomal entry site (IRES)-mediated initiation. To evaluate the effect of IBTK on both mechanisms, we performed dual-luciferase assays. The upstream reporter (Renilla luciferase) of the bicistronic mRNA was translated via a cap-dependent mechanism, whereas the downstream reporter (Firefly luciferase) was under the control of internal initiation from the hepatitis C virus (HCV)-IRES or (EMCV)-IRES(26). Our dual-luciferase assays showed that IBTK depletion caused a significant decrease of Renilla signals by approximately 60% in both systems (**Fig. 3G**), indicating that IBTK has a direct effect on cap-dependent translation initiation. Moreover, we observed that IBTK deficiency reduced the interaction between eIF4A1 and eIF4 subunits (**Fig. 3H**) and the interaction between capped mRNAs and the eIF4 complex in IBTK-KO cells (assessed by m^7GTP-Sepharose pull-down assays, **Fig. 3I**). Collectively, these data suggest that IBTK plays a positive role in regulating the translation initiation activity of the eIF4F complex.

Deficiency of IBTK reduces the expression of oncogenes that are dependent on eIF4A1 and mitigates neoplastic phenotypes in cancer cells

The mRNAs of many oncogenes have long and complex 5'-UTRs, which require high levels of eIF4A1 helicase activity for efficient translation(7). In SiHa cells, the depletion of IBTK resulted in a marked reduction in the protein levels of multiple eIF4A1-regulated oncogenes, such as CDK6, MYC, STAT1, and CCND3 (**Fig. 4A**). Similar results were observed in IBTK-KO H1299 cells and IBTK-KD CT26 cells (derived from a murine colon adenocarcinoma) (**Supplementary Fig. 4A-D**). Importantly, the mRNA levels of oncogenes were largely unaffected by IBTK depletion (**Fig. 4B**), indicating a post-transcriptional regulation exerted by IBTK. However, the overexpression of IBTK only marginally affected the protein levels of oncogenes (**Fig. 4C**), likely due to the constitutive activity of the eIF4 complex in immortalized cell lines.

Given the crucial role of eIF4A1 in cancer biology, we further investigated whether

IBTK promotes neoplastic phenotypes in cancer cells. We observed that IBTK KO significantly decreased SiHa cell proliferation, migration, invasion, and anchorage-independent cell growth (**Fig. 4D-G**). Additionally, IBTK depletion reduced xenograft tumor growth *in vivo* (**Fig. 4H**). Moreover, our data showed that IBTK depletion enhanced apoptosis in response to eIF4A1 inhibitors such as silvestrol or rocaglamide A (**Fig. 4I, J, Supplementary Fig. 5A, B**). The tumor-suppressing effects of IBTK depletion on HeLa cells were also observed (**Supplementary Fig. 5C-H**). Collectively, these data suggest that IBTK plays a critical role in promoting oncogene expression and cancer cell malignancy by modulating eIF4A1 activity.

IBTK facilitates IFN-γ-induced PD-L1 expression and tumor immune escape

Recent study, a control mechanism of tumor immune evasion was demonstrated at the translational level through the eIF4F-STAT1-IRF1-PD-L1 pathway. The eIF4F complex regulates the translation of transcription factor STAT1 and promotes IFN-γ-induced PD-L1 expression(8). As IBTK-KO cells showed compromised eIF4F-initiated translation activity and downregulated STAT1 protein levels, we investigated the potential impact of IBTK on tumor immune escape and IFN-γ-induced PD-L1 expression. Our results demonstrated that the expression of STAT1, IRF1, and PD-L1 induced by IFN-γ was markedly reduced in both IBTK-KO H1299 cells (**Supplementary Fig. 6A-C**) and IBTK-KD CT26 cells (**Supplementary Fig. 6D-F**). To further investigate the effects of IBTK on tumor growth in the presence of a functional immune system, we used a CT26 xenograft tumor model and found that immunocompetent mice implanted with IBTK-KD CT26 cells exhibited reduced tumor growth *in vivo* (**Supplementary Fig. 6G-I**). CD8[+] cytotoxic T lymphocytes (CTLs) play a crucial role in antitumor immunity by secreting granzyme B, which triggers apoptosis. Notably, depletion of IBTK increased the population of CD8[+] CTLs and the release of granzyme B in xenograft tumors (**Supplementary Fig. 6J, K**). Collectively, these data indicate that IBTK promotes IFN-γ-induced PD-L1 expression and tumor immune escape.

IBTK-mediated eIF4A1 ubiquitination is regulated by mTORC1/S6K1 signaling

We investigated whether upstream mTORC1/S6K1 signaling regulates IBTK-mediated eIF4A1 ubiquitination. Three quantitative phosphoproteomic studies revealed that treatment with mTOR inhibitors (rapamycin or Torin 1) led to markedly reduced phosphorylation levels of seven Ser/Thr sites on IBTK(27-29). Notably, these sites are clustered in a short region (990-1068 aa) close to IBTK's last BTB domain (**Fig. 5A**). We established the interaction of IBTK with the mTORC1 complex and S6K1 in cells (**Fig. 5B-E**). We then examined the effect of mTOR signaling on IBTK phosphorylation. To do so, we generated an IBTK deletion mutant (900-1150 aa) spanning the potential mTORC1-regulated phosphorylation sites. Phos-tag gels were utilized for detection of shifts in the electrophoretic mobility of phosphorylated proteins. The results demonstrated that the phosphorylated IBTK$_{900-1150aa}$ was markedly decreased while the non-phosphorylated form was simultaneously increased in amino acid (AA)-deprived or rapamycin-treated cells (**Fig. 5F**). Moreover, phosphorylation levels of IBTK$_{900-1150aa}$ were markedly downregulated in cells deficient in Raptor, an essential subunit of mTORC1 but not mTORC2 (**Fig. 5G**), indicating that mTORC1 signaling mediates IBTK phosphorylation.

In order to investigate whether mTOR or S6K1 directly phosphorylates IBTK, we conducted *in vitro* kinase assays using recombinant IBTK$_{900-1150aa}$ segment as a substrate. The analysis of the phospho-peptides via MS revealed that mTOR phosphorylates residues S911/999/1004/1039/1045/1069/1089/1141, while S6K1 phosphorylates residues S971/990/993/1033/1045/1096 and T1101 (**Fig. 5H, I, Supplementary Fig. 7A, B, Supplementary Table. 4, 5**). Thus, four out of seven sites (excluding S992, T1008, and T1068) were confirmed as mTOR/S6K1 phosphorylation sites *in vitro*, thereby confirming that IBTK is an authentic mTORC1/S6K1 substrate. Additionally, simultaneous mutation of seven Ser/Thr sites to Ala nearly abolished the mobility shift of IBTK$_{900-1150aa}$ (**Fig. 5J**), suggesting that most, if not all, mTOR-regulated sites are localized in this region. We generated a phosphorylation-specific antibody

that targets the combination of phospho-S990/992/993 in IBTK. Using this antibody,

we observed a marked decrease in the phosphorylation levels of three adjacent Ser

residues in IBTK upon AA deprivation or rapamycin treatment (**Fig. 5K**). Furthermore,

we found that IBTK-mediated eIF4A1 ubiquitination was considerably decreased by

AA deprivation or rapamycin treatment (**Fig. 5L**). To investigate the role of

mTORC1/S6K1-regulated phosphorylation in this process, we generated an IBTK

mutant (IBTK-7S/TA) in which seven Ser/Thr residues were replaced with Ala to

abolish phosphorylation regulated by mTORC1/S6K1. We noticed that the IBTK-7S/TA

mutant showed diminished capacity to ubiquitinate eIF4A1 compared to IBTK-WT (**Fig.**

5M). Collectively, these data suggest that IBTK is a phosphorylation substrate for

mTORC1/S6K1 and that the mTORC1/S6K1 signaling pathway positively regulates

IBTK-mediated eIF4A1 ubiquitination.

The phosphorylation of IBTK by mTORC1/S6K1 is crucial for sustaining
oncogenic translation

In order to elucidate the biological significance of IBTK phosphorylation mediated by

the mTOR/S6K1 signaling pathway, we introduced IBTK-WT or IBTK-7S/TA mutants

into IBTK-KO SiHa cells (termed IBTK^{WT} and IBTK$^{7S/TA}$, respectively). Our findings

revealed that IBTK$^{7S/TA}$ cells exhibited lower levels of global protein synthesis and

reduced expression of eIF4A1-regulated oncogenes compared to IBTK^{WT} cells (**Fig.**

6A, B). Additionally, there was a noticeable reduction in the interaction between

eIF4A1 and other eIF4 complex subunits in IBTK$^{7S/TA}$ cells compared to IBTK^{WT} cells

(**Fig. 6C**). The interaction between capped mRNAs and eIF4 complex was also

markedly reduced in IBTK$^{7S/TA}$ cells (**Fig. 6D**). Moreover, IBTK$^{7S/TA}$ cells exhibited

significant reductions in cell growth, migration, and silvestrol-induced apoptosis (**Fig.**

6E-J). Collectively, these data suggest that mTORC1/S6K1-mediated IBTK

phosphorylation favors sustained oncogenic translation mediated by eIF4A1.

Overexpression of IBTK correlates with poor survival in cervical cancer

The aforementioned results established that demonstrated that IBTK promotes oncogenic eIF4A1 activation in cervical cancer cell lines (HeLa and SiHa). In light of these findings, we sought to evaluate the pathological significance of IBTK expression in cervical cancer. The Cancer Genome Atlas (TCGA) dataset revealed that high IBTK expression levels were correlated with poor survival in cervical squamous cell carcinoma (CESC), which is the most prevalent histological subtype of cervical cancer (**Fig. 7A**). Following validation of the antibody specificity for IHC analysis in parental/IBTK-KO cells (**Supplementary Fig. 8**), we conducted an immunohistochemistry (IHC) staining analysis of 117 CESC tissues, including 35 adjacent normal cervical tissues, by using a tissue microarray (**Fig. 7B**). The results showed a marked upregulation of IBTK protein expression in CESC tissues compared with adjacent normal cervical tissues. Moreover, IBTK expression levels were positively associated with advanced pathological grade and predicted poor survival outcomes (**Fig. 7C-E**). Taken together, these data indicate that IBTK protein expression is increased in cervical cancer and correlates with poor prognosis in patients with cervical cancer.

DISCUSSION

Oncogenic signaling appears to dominate translation control at nearly all stage of cancer propagation for very specific and distinct cellular phenotypes. In general, cancer cells upregulate eIF4F activity to promote cap-dependent translation and oncogene expression, leading to rapid cell growth. Specifically, the study reveals a novel signaling axis involving mTORC1/S6K1-IBTK-eIF4A1 that promotes cap-dependent translation and oncoprotein expression, contributing to tumor cell growth. Additionally, we demonstrate that mTORC1 and S6K1 phosphorylate IBTK at multiple sites to boost CRL3[IBTK]-mediated IBTK ubiquitination under nutrient-rich conditions. Overall, this study adds to our understanding of how cancer cells hijack protein synthesis machinery to promote their survival and proliferation, and provides potential avenues for developing new cancer treatments (**Fig. 7E**).

The development of eIF4A inhibitors as therapeutic agents for cancer patients is an exciting area of research. However, the emergence of resistance to these inhibitors highlights the need for a better understanding of the molecular mechanisms underlying drug resistance. The recent genome-wide CRISPR/Cas9 screen has identified three negative NRF2 regulators (KEAP1, CUL3, CAND1) whose inactivation can confer resistance to the silvestrol analogue, providing insights into potential pathways that may contribute to drug resistance(30). Additionally, our study shows that IBTK overexpression can suppress AS-induced SG assembly and that knockdown of IBTK can sensitize cancer cells to eIF4A inhibitor-induced cell death, making IBTK a potential therapeutic target to overcome intrinsic resistance to eIF4A1 inhibitor. Overall, these findings underscore the importance of continued research in the development of eIF4A1 inhibitors and the identification of new therapeutic targets to combat drug resistance in cancer cells.

The tumor suppressor PDCD4 inhibits translation by binding directly to eIF4A. Pisano et al. reported that PDCD4 is the first CRL3^{IBTK} substrate and that CRL3^{IBTK} targets PDCD4 for ubiquitin-dependent degradation(13). Although this study highlighted the potential involvement of IBTK in translation regulation, we did not observe any changes in protein levels of PDCD4 in IBTK-KD or -KO cells. Instead, our study showed that eIF4A1 is a major downstream effector of CRL3^{IBTK}, which has been previously observed to regulate several well-characterized eIF4A targets, such as MYC and MCL1, in Eμ-myc mice(17). Our results also suggest that IBTK may play a role in tumor immune escape through modulating the eIF4F-STAT1–IRF1–PD-L1 axis. This is supported by a recent study indicating that eIF4A1 promotes PD-L1-mediated tumor immune escape by controlling the translation of STAT1(8).

In contrast to eIF4A1/2, IBTK cannot ubiquitinate eIF4A3, despite a strong interaction between the two proteins. It is not currently clear whether IBTK can promote eIF4A3 ubiquitination in certain circumstances or simply modulate eIF4A3 activity through direct binding. eIF4A1 is an essential subunit of EJC. The role of IBTK in post-transcriptional gene regulation through modulation of EJC activity is an area that

warrants further investigation, particularly given that eIF4A3 or IBTK knockdown had

an impact on alternative splicing in HeLa cells(31, 32). Alongside its interaction with

eIF4A1/2, IBTK may also play diverse roles in different stages of the translation

process. The AP-MS/MS data and biochemical validation provided strong evidence for

the interaction between IBTK and eIF2α (**Fig. 1G**). A previous study showed that a

functional uORF in the 5'-UTR sequence allows for preferential translation of IBTK in

response to p-eIF2α and ER stress(33). Our findings suggest that IBTK may interact

with eIF2α and modulate its activity, either positively or negatively, through direct

binding in response to p-eIF2α and ER stress. The identification of IBTK interacting

with multiple subunits of eIF3 complex (**Fig. 1C, G**), which play key roles in different

stages of translation(34), highlights the potential role of IBTK in modulating translation

regulation through eIF3 complex. These findings provide a basis for further

investigations into the precise role of IBTK in controlling the efficiency and fidelity of

protein synthesis.

Several quantitative phosphoproteomic studies have used mTOR inhibitors coupled

with MS to demonstrate that mTOR kinase modulates the phosphorylation of

thousands of currently uncharacterized mTOR substrates, either directly or indirectly

through its downstream kinases(27-29). One such potential mTOR substrate is IBTK.

We observed that seven mTOR-regulated Ser/Thr sites are clustered in a short region

of IBTK and hyperphosphorylation of IBTK$_{900-1150aa}$ was abolished when mTORC1

activity was inhibited. In particular, using *in vitro* kinase assays combined with MS, we

have identified S999/S1004 as sites phosphorylated by mTOR, and S990/993 as sites

phosphorylated by S6K1. However, the kinases responsible for the other three

phosphorylation sites remain unclear and require further investigation. Understanding

the precise mechanisms by which mTORC1 regulates IBTK phosphorylation and

activity may provide insight into the broader functions of mTORC1 signaling in protein

synthesis and cellular homeostasis.

Abbreviations

eIF: eukaryotic initiation factor; CRL3: Cullin 3-RING ubiquitin ligase complex; EJC: exon junction complex; S6Ks: S6 kinasesIBTK: the Inhibitor of Bruton's tyrosine kinase; CCK-8: Cell Counting Kit-8; PI: propidium iodide; BioID: biotinylation identification; DOX: doxycycline; SGs: stress granules; AS: sodium arsenite; WCLs: whole cell lysates; PFA: paraformaldehyde; ATCC: American Type Culture Collection; WT: Wild-Type; RT-qPCR: Quantitative real-time polymerase chain reaction; IHC: Immunohistochemistry; IP: Immunoprecipitation

Acknowledgments

This work was in part supported by the National Natural Science Foundation of China (No. 82272992, 91954106, and 81872109 to K.G.; No. 91957125, 81972396 to C.W.), the State Key Development Programs of China (No. 2022YFA1104200 to C.W.), the Natural Science Foundation of Shanghai (No. 22ZR1449200 to K.G; 22ZR1406600 to C.W.), and the Open Research Fund of State Key Laboratory of Genetic Engineering, Fudan University (No. SKLGE-2111 to K.G.). Science and Technology Research Program of Shanghai (No. 9DZ2282100).

Availability of data

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD028479, PXD028480, and PXD039031.

Author contributions

K.G and C.W. conceived of the project. K.G, C.W., and Y.L. supervised the experiments, analyzed the data, and wrote the manuscript. H.S., D.J., X.Z., Y.C., Q.S., and Z.L. performed experiments and data analyses. All the authors discussed the results and commented on the manuscript.

References

1.   M. Bhat, N. Robichaud, L. Hulea, N. Sonenberg, J. Pelletier, I. Topisirovic, Targeting the
translation machinery in cancer. *Nat Rev Drug Discov* **14**, 261-278 (2015).

2.   L. Fabbri, A. Chakraborty, C. Robert, S. Vagner, The plasticity of mRNA translation during
cancer progression and therapy resistance. *Nat Rev Cancer* **21**, 558-577 (2021).

3.   W. C. Merrick, eIF4F: a retrospective. *J Biol Chem* **290**, 24091-24099 (2015).

4.   W. T. Lu, A. Wilczynska, E. Smith, M. Bushell, The diverse roles of the eIF4A family: you are
the company you keep. *Biochem Soc Trans* **42**, 166-172 (2014).

5.   H. Malka-Mahieu, M. Newman, L. Desaubry, C. Robert, S. Vagner, Molecular Pathways:
The eIF4F Translation Initiation Complex-New Opportunities for Cancer Treatment. *Clin*
Cancer Res **23**, 21-25 (2017).

6.   J. Pelletier, J. Graff, D. Ruggero, N. Sonenberg, Targeting the eIF4F translation initiation
complex: a critical nexus for cancer development. *Cancer Res* **75**, 250-263 (2015).

7.   L. Boussemart, H. Malka-Mahieu, I. Girault, D. Allard, O. Hemmingsson, G. Tomasic, M.
Thomas, C. Basmadjian, N. Ribeiro, F. Thuaud, C. Mateus, E. Routier, N. Kamsu-Kom, S.
Agoussi, A. M. Eggermont, L. Desaubry, C. Robert, S. Vagner, eIF4F is a nexus of resistance
to anti-BRAF and anti-MEK cancer therapies. *Nature* **513**, 105-109 (2014).

8.   M. Cerezo, R. Guemiri, S. Druillennec, I. Girault, H. Malka-Mahieu, S. Shen, D. Allard, S.
Martineau, C. Welsch, S. Agoussi, C. Estrada, J. Adam, C. Libenciuc, E. Routier, S. Roy, L.
Desaubry, A. M. Eggermont, N. Sonenberg, J. Y. Scoazec, A. Eychene, S. Vagner, C. Robert,
Translational control of tumor immune escape via the eIF4F-STAT1-PD-L1 axis in
melanoma. *Nat Med* **24**, 1877-1886 (2018).

9.   J. T. Ernst, P. A. Thompson, C. Nilewski, P. A. Sprengeler, S. Sperry, G. Packard, T. Michels,
A. Xiang, C. Tran, C. J. Wegerski, B. Eam, N. P. Young, S. Fish, J. Chen, H. Howard, J.
Staunton, J. Molter, J. Clarine, A. Nevarez, G. G. Chiang, J. R. Appleman, K. R. Webster, S.
H. Reich, Design of Development Candidate eFT226, a First in Class Inhibitor of Eukaryotic
Initiation Factor 4A RNA Helicase. *J Med Chem* **63**, 5879-5955 (2020).

10.   S. Battaglioni, D. Benjamin, M. Walchli, T. Maier, M. N. Hall, mTOR substrate
phosphorylation in growth control. *Cell* **185**, 1814-1836 (2022).

11.   P. Genschik, I. Sumara, E. Lechner, The emerging family of CULLIN3-RING ubiquitin ligases
(CRL3s): cellular functions and disease implications. *EMBO J* **32**, 2307-2320 (2013).

12.   W. Liu, I. Quinto, X. Chen, C. Palmieri, R. L. Rabin, O. M. Schwartz, D. L. Nelson, G. Scala,
Direct inhibition of Bruton's tyrosine kinase by IBtk, a Btk-binding protein. *Nat Immunol*
2, 939-946 (2001).

13.   A. Pisano, S. Ceglia, C. Palmieri, E. Vecchio, G. Fiume, A. de Laurentiis, S. Mimmi, C. Falcone,
E. Iaccino, A. Scialdone, M. Pontoriero, F. F. Masci, R. Valea, S. Krishnan, M. Gaspari, G.
Cuda, G. Scala, I. Quinto, CRL3IBTK Regulates the Tumor Suppressor Pdcd4 through
Ubiquitylation Coupled to Proteasomal Degradation. *J Biol Chem* **290**, 13958-13971
(2015).

14.   J. A. Willy, S. K. Young, A. L. Mosley, S. Gawrieh, J. L. Stevens, H. C. Masuoka, R. C. Wek,
Function of inhibitor of Bruton's tyrosine kinase isoform alpha (IBTKalpha) in nonalcoholic
steatohepatitis links autophagy and the unfolded protein response. *J Biol Chem* **292**,
14050-14065 (2017).

15.   T. D. Baird, L. R. Palam, M. E. Fusakio, J. A. Willy, C. M. Davis, J. N. McClintick, T. G. Anthony,
R. C. Wek, Selective mRNA translation during eIF2 phosphorylation induces expression of
IBTKalpha. *Mol Biol Cell* **25**, 1686-1697 (2014).

16.   J. Luo, M. J. Emanuele, D. Li, C. J. Creighton, M. R. Schlabach, T. F. Westbrook, K. K. Wong,
S. J. Elledge, A genome-wide RNAi screen identifies multiple synthetic lethal interactions
with the Ras oncogene. *Cell* **137**, 835-848 (2009).

17.   E. Vecchio, G. Golino, A. Pisano, F. Albano, C. Falcone, S. Ceglia, E. Iaccino, S. Mimmi, G.
Fiume, G. Giurato, D. Britti, G. Scala, I. Quinto, IBTK contributes to B-cell lymphomagenesis
in Emu-myc transgenic mice conferring resistance to apoptosis. *Cell Death Dis* **10**, 320
(2019).

18.   F. Albano, F. Chiurazzi, S. Mimmi, E. Vecchio, A. Pastore, C. Cimmino, C. Frieri, E. Iaccino,
A. Pisano, G. Golino, G. Fiume, M. Mallardo, G. Scala, I. Quinto, The expression of inhibitor
of bruton's tyrosine kinase gene is progressively up regulated in the clinical course of
chronic lymphocytic leukaemia conferring resistance to apoptosis. *Cell Death Dis* **9**, 13
(2018).

19.   E. Vecchio, N. Nistico, G. Golino, E. Iaccino, D. Maisano, S. Mimmi, A. Aloisio, M. Renna, A.
Avagliano, A. Arcucci, G. Fiume, I. Quinto, IBtkalpha Activates the beta-Catenin-
Dependent Transcription of MYC through Ubiquitylation and Proteasomal Degradation
of GSK3beta in Cancerous B Cells. *Int J Mol Sci* **23**,   (2022).

20.   X. Liu, K. Salokas, F. Tamene, Y. Jiu, R. G. Weldatsadik, T. Ohman, M. Varjosalo, An AP-
MS- and BioID-compatible MAC-tag enables comprehensive mapping of protein
interactions and subcellular localizations. *Nat Commun* **9**, 1188 (2018).

21.   E. L. Huttlin, R. J. Bruckner, J. Navarrete-Perea, J. R. Cannon, K. Baltier, F. Gebreab, M. P.
Gygi, A. Thornock, G. Zarraga, S. Tam, J. Szpyt, B. M. Gassaway, A. Panov, H. Parzen, S. Fu,
A. Golbazi, E. Maenpaa, K. Stricker, S. Guha Thakurta, T. Zhang, R. Rad, J. Pan, D. P.
Nusinow, J. A. Paulo, D. K. Schweppe, L. P. Vaites, J. W. Harper, S. P. Gygi, Dual proteome-
scale networks reveal cell-specific remodeling of the human interactome. *Cell* **184**, 3022-
3040 e3028 (2021).

22.   Z. Li, Z. Cheng, C. Raghothama, Z. Cui, K. Liu, X. Li, C. Jiang, W. Jiang, M. Tan, X. Ni, A.
Pandey, J. O. Liu, Y. Dang, USP9X controls translation efficiency via deubiquitination of
eukaryotic translation initiation factor 4A1. *Nucleic Acids Res* **46**, 823-839 (2018).

23.   G. Galicia-Vazquez, J. Chu, J. Pelletier, eIF4AII is dispensable for miRNA-mediated gene
silencing. *RNA* **21**, 1826-1833 (2015).

24.   G. Galicia-Vazquez, R. Cencic, F. Robert, A. Q. Agenor, J. Pelletier, A cellular response
linking eIF4AI activity to eIF4AII transcription. *RNA* **18**, 1373-1384 (2012).

25.   H. Mahboubi, U. Stochaj, Cytoplasmic stress granules: Dynamic modulators of cell
signaling and disease. *Biochim Biophys Acta Mol Basis Dis* **1863**, 884-895 (2017).

26.   N. Sonenberg, Measures and countermeasures in the modulation of initiation factor
activities by viruses. *New Biol* **2**, 402-409 (1990).

27.   P. P. Hsu, S. A. Kang, J. Rameseder, Y. Zhang, K. A. Ottina, D. Lim, T. R. Peterson, Y. Choi,
N. S. Gray, M. B. Yaffe, J. A. Marto, D. M. Sabatini, The mTOR-regulated phosphoproteome
reveals a mechanism of mTORC1-mediated inhibition of growth factor signaling. *Science*
332, 1317-1322 (2011).

28.   R. Q. Chen, Q. K. Yang, B. W. Lu, W. Yi, G. Cantin, Y. L. Chen, C. Fearns, J. R. Yates, 3rd, J.

772   D. Lee, CDC25B mediates rapamycin-induced oncogenic responses in cancer cells. *Cancer*
773   *Res* **69**, 2663-2668 (2009).

774 29. Y. Yu, S. O. Yoon, G. Poulogiannis, Q. Yang, X. M. Ma, J. Villen, N. Kubica, G. R. Hoffman,
775   L. C. Cantley, S. P. Gygi, J. Blenis, Phosphoproteomic analysis identifies Grb10 as an
776   mTORC1 substrate that negatively regulates insulin signaling. *Science* **332**, 1322-1326
777   (2011).

778 30. V. R. Sanghvi, P. Mohan, K. Singh, L. Cao, M. Berishaj, A. L. Wolfe, J. H. Schatz, N. Lailler, E.
779   de Stanchina, A. Viale, H. G. Wendel, NRF2 Activation Confers Resistance to eIF4A
780   Inhibitors in Cancer Therapy. *Cancers (Basel)* **13**, (2021).

781 31. G. Fiume, A. Scialdone, F. Rizzo, M. R. De Filippo, C. Laudanna, F. Albano, G. Golino, E.
782   Vecchio, M. Pontoriero, S. Mimmi, S. Ceglia, A. Pisano, E. Iaccino, C. Palmieri, S. Paduano,
783   G. Viglietto, A. Weisz, G. Scala, I. Quinto, IBTK Differently Modulates Gene Expression and
784   RNA Splicing in HeLa and K562 Cells. *Int J Mol Sci* **17**, (2016).

785 32. Z. Wang, V. Murigneux, H. Le Hir, Transcriptome-wide modulation of splicing by the exon
786   junction complex. *Genome Biol* **15**, 551 (2014).

787 33. P. H. Amin, K. R. Carlson, R. C. Wek, An RNA stem-loop functions in conjunction with an
788   upstream open reading frame to direct preferential translation in the integrated stress
789   response. *J Biol Chem* **299**, 102864 (2023).

790 34. A. Gomes-Duarte, R. Lacerda, J. Menezes, L. Romao, eIF3: a factor for human health and
791   disease. *RNA Biol* **15**, 26-34 (2018).

792 35. J. P. Theurillat, N. D. Udeshi, W. J. Errington, T. Svinkina, S. C. Baca, M. Pop, P. J. Wild, M.
793   Blattner, A. C. Groner, M. A. Rubin, H. Moch, G. G. Prive, S. A. Carr, L. A. Garraway, Prostate
794   cancer. Ubiquitylome analysis identifies dysregulation of effector substrates in SPOP-
795   mutant prostate cancer. *Science* **346**, 85-89 (2014).

Conflict of Interest statement

The authors declare no competing interests.

Ethics statement

This study involved human subjects and animal experiments and was approved by the Ethics

Review Committee of Shanghai First Maternity and Infant Hospital.

Figure Legends

Figure 1. IBTK interacts with eIF4A1 in cells.

(**A**) WB analysis of the indicated proteins in WCLs from FLAG-BirA*-IBTK Tet-on-inducible Flp-In T-REx 293 cells treated with DOX (10 ng/ml) for the indicated times.

(**B**) Affinity purification of IBTK-containing protein complex using FLAG-M2 beads was conducted in FLAG-BirA*-IBTK Tet-on-inducible Flp-In T-REx 293 cells. The associated proteins were separated by SDS-PAGE and visualized by Coomassie Blue (CB) staining.

(**C**)The eIF interactome of IBTK identified by AP-MS analysis in (**B**). The full interaction partner list was showed in Supplementary Table 1.

(**D-F**) WB analysis of the indicated proteins in WCLs and co-IP samples of anti-FLAG antibody obtained from 293T cells transfected with the indicated plasmids.

(**G**) FLAG-BirA*-IBTK Tet-on-inducible Flp-In T-REx 293 cells were treated with/without DOX (10 ng/ml) for 12 h then collected and subjected to co-IP with anti-FLAG antibody. WCLs and co-IP samples were prepared for WB analysis with the indicated antibodies.

(**H-K**) Co-IP using anti-eIF4A1(**H**), eIF4A2 (**I**), eIF4A3 (**J**), or IBTK (**K**) antibodies in WCLs prepared from 293T cells, followed by WB analysis with the indicated antibodies.

Figure 2. IBTK promotes non-degradative ubiquitination of eIF4A1.

(**A**) WB analysis of the indicated proteins in WCLs from FLAG-BirA*-IBTK Tet-on-inducible T-REx 293 cells treated with DOX (10 ng/ml) for the indicated times.

(**B**) WB analysis of the indicated proteins in WCLs from SiHa and HeLa cells infected with lentivirus expressing IBTK-specific shRNA (#1, #2) or negative control (NC).

(**C**) WB analysis of the indicated proteins in WCLs from parental/IBTK-KO SiHa, 293T, and H1299 cells.

(**D**) WB analysis of the products of *in vivo* ubiquitination assays from 293T cells transfected with the indicated plasmids.

(**E**) Parental and IBTK-KO 293T cells were transfected with HA-Ub for 24 h, then WCLs were prepared for co-IP with IgG, anti-eIF4A1, eIF4A2, or eIF4A3 antibodies. The polyubiquitinated forms of eIF4A1/2/3 were detected by WB with anti-HA antibody.

(**F**) WB analysis of the products of *in vitro* ubiquitination assays with anti-eIF4A1

834   antibody.

835   (**G**) WB analysis of the products of *in vivo* ubiquitination assays from 293T cells

836   transfected with the indicated plasmids.

Figure 3. IBTK promotes nascent protein synthesis and cap-dependent translation initiation.

(**A**) WB analysis of the indicated proteins in WCLs from parental and IBTK-KO SiHa cells treated with silvestrol (100 nM, 24 h) and puromycin (1.5 µM, 10 min).

(**B**) WB analysis of the indicated proteins in WCLs from parental and IBTK-KD HeLa cells treated with silvestrol (100 nM, 24 h) and puromycin (1.5 µM, 10 min).

(**C**) Representative IF images of parental and IBTK-KO SiHa cells treated with AS (100 µM, 2 h) and then stained with eIF4A1 (red) and DAPI (blue). Scale bar, 20 µm.

(**D**) SGs in (**C**) were quantified by counts of SGs per cell based on eIF4A1 puncta. Data are presented as means ± S.D. (n = 20).

(**E**) Representative IF images of SiHa cells transfected with indicated plasmids, treated

with DMSO or AS (100 μM, 2 h), and then stained with FLAG (green), eIF4A1 (red),

and DAPI (blue). Cells successfully transfected with FLAG-tag plasmids are marked

with white dashed lines. Scale bar, 20 μm.

(**F**) SGs in (**E**) were quantified by counts of SGs per cell (IBTK transfected vs non-

transfected) based on eIF4A1 puncta. Data are presented as means ± S.D. (n = 20).

(**G**) Parental or IBTK-KO SiHa cells were transfected with the reporter pRΔDE·HCVF

(pH) or pRΔDE·EMCVF (pE) for 24 h and the luciferase activities were measured. The

counts of Firefly and Renilla were measured and shown in the graph. Data are

presented as means ± S.D. (n = 3).

(**H**) Co-IP assays using IgG or anti-eIF4A1 antibody in WCLs prepared from parental

and IBTK-KO SiHa cells followed by WB analysis with the indicated antibodies.

(**I**) WCLs of parental and IBTK-KO SiHa cells were incubated with m^{7}GTP-Sepharose

beads, and the pull-down proteins were subjected to WB analysis with the indicated

antibodies.

P values are calculated using One-way ANOVA test in (**D**, **F**, **G**). *p<0.05, **p<0.01,

***p<0.001, ****p<0.0001, n.s. non-significant.

Figure 4. IBTK deficiency reduces eIF4A1-dependent oncoprotein expression and neoplastic phenotypes of cancer cells.

(**A**) WB analysis of the indicated proteins in WCLs from parental and IBTK-KO SiHa cells (left) and WCLs from SiHa cells treated with silvestrol (100 nM) for the indicated times (right).

(**B**) RT-qPCR assessment of the mRNA expression of eIF4A1 target genes in parental and IBTK-KO SiHa cells. The mRNA levels of *Actin* were used for normalization. Data

are shown as means ± S.D. (n = 3).

(**C**) WB analysis of the indicated proteins in WCLs from FLAG-BirA*-IBTK Tet-on-inducible Flp-In T-REx 293 cells treated with DOX (10 ng/ml) for 12 h.

(**D**) CCK-8 cell proliferation analysis of parental and IBTK-KO SiHa cells. Data are shown as means ± S.D. (n=3).

(**E**) Colony formation analysis of parental and IBTK-KO SiHa cells, and the quantitative data is shown (below). Data are shown as means ± S.D. (n=3).

(**F**) Cell migration and invasion analysis of parental and IBTK-KO SiHa cells, and the quantitative data is shown (right). Data are shown as means ± S.D. (n=3). Scale bar, 50 μm.

(**G**) Sphere-formation analysis of parental and IBTK-KO SiHa cells. Representative pictures of SiHa cells after two weeks in three-dimensional culture are shown. Average size per sphere and number of spheres per 1000 cells were calculated by ImageJ. The quantitative data is shown (right). Data are shown as means ± S.D. (n=3). Scale bar, 100 μm.

(**H**) Parental and IBTK-KD HeLa cells were injected s.c. into the right flank of BALB/c mice. The tumors in each group at day 20 were harvested and photographed, the quantitative data of tumor weights is shown (below). Data are shown as means ± SD. (n=9).

(**I**) WB analysis of the indicated proteins in WCLs from parental and IBTK-KO SiHa cells treated with silvestrol (100 nM) for the indicated times.

(**J**) Parental and IBTK-KO SiHa cells were treated with silvestrol (100 nM) for the indicated times. Then, annexin-V-FITC/PI dyes were used to stain the harvested cells, of which later flow cytometry analysis was performed. Data are shown as means ± SD. (n=3).

P values are calculated using unpaired Student's t test in (**H**), One-way ANOVA test in (**E, F, G**), Two-way ANOVA test in (**B, D, J**). *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$, n.s. non-significant.

Figure 5. IBTK-mediated eIF4A1 ubiquitination is regulated by mTORC1/S6K1 signaling.

(**A**) The potential mTOR-regulated phosphorylation sites of IBTK which were sensitive to rapamycin or Torin1 treatment, according to three quantitative phosphoproteomic studies.

(**B-D**) WB analysis of the indicated proteins in WCLs and co-IP samples of anti-FLAG antibody obtained from 293T cells transfected with the indicated plasmids.

(**E**) Co-IP using anti-IBTK antibodies in WCLs prepared from 293T cells, followed by

WB analysis with the indicated antibodies.

(**F**) WB analysis of the indicated proteins in WCLs from 293T cells transfected with FLAG-IBTK$_{900-1150aa}$ and starved of amino acids (AA) or treated with rapamycin (500 nM) for the indicated times. The phosphorylated forms of IBTK$_{900-1150aa}$ were detected by WB using phos-tag gels. The arrow indicates non-phosphorylated IBTK$_{900-1150aa}$.

(**G**) WB analysis of the indicated proteins in WCLs from parental or Raptor-KO 293T cells transfected with FLAG-IBTK$_{900-1150aa}$. The phosphorylated form of IBTK$_{900-1150aa}$ was detected by WB using phos-tag gels.

(**H**) Recombinant GST-IBTK$_{900-1150aa}$ proteins were subjected to phosphorylation by recombinant mTOR/mLST8, or S6K1, as detected using *in vitro* kinase assays. The reaction products were separated by SDS-PAGE and visualized by CB staining.

(**I**) mTOR/mLST8 or S6K1-mediated IBTK *in vitro* phosphorylation sites identified by MS analysis.

(**J**) WB analysis of the indicated proteins in WCLs from 293T cells transfected with FLAG-IBTK$_{900-1150aa}$ (WT or 7S/TA mutant) and starved of amino acids or treated with rapamycin (500 nM) for 2 h.

(**K**) Co-IP using anti-IBTK antibody in WCLs prepared from 293T cells starved of amino acids (AA-) or treated with rapamycin (500 nM) for 2 h, followed by WB analysis with the indicated antibodies.

(**L**) WB analysis of the products of i*n vivo* ubiquitination assays from 293T cells transfected with the indicated plasmids and starved of amino acids or treated with rapamycin (500 nM) for 2 h.

(**M**) WB analysis of the products of *in vivo* ubiquitination assays from 293T cells transfected with the indicated plasmids.

Figure 6. Deficiency in mTORC1/S6K1 signaling-mediated eIF4A1 phosphorylation reduces eIF4A1-dependent oncoprotein expression and neoplastic phenotypes of cancer cells.

(**A**) WB analysis of the indicated proteins in WCLs from parental and IBTK-KO SiHa cells reconstituted with EV, IBTK-WT, or -7S/TA mutant and treated with puromycin (1.5 µM, 10 min).

(**B**) WB analysis of the indicated proteins in WCLs from parental and IBTK-KO SiHa cells reconstituted with EV, IBTK-WT, or -7S/TA mutant.

(**C**) Co-IP assays using anti-eIF4A1 antibody in WCLs prepared from parental and IBTK-KO SiHa cells followed by WB analysis with the indicated antibodies.

(**D**) WCLs of parental and IBTK-KO SiHa cells reconstituted with EV, IBTK-WT, or -7S/TA mutant were incubated with m^7GTP-Sepharose beads, and the pull-down proteins were subjected to WB analysis with the indicated antibodies.

(**E**) CCK-8 cell proliferation analysis of parental and IBTK-KO SiHa cells reconstituted with EV, IBTK-WT, or -7S/TA mutant. Data are shown as means ± S.D. (n=3).

(**F, G**) Colony formation analysis of parental and IBTK-KO SiHa cells reconstituted with EV, IBTK-WT or, -7S/TA mutant, and the quantitative data is shown in (**G**). Data are shown as means ± S.D. (n=3).

(**H, I**) Cell migration analysis of parental and IBTK-KO SiHa cells reconstituted with EV, IBTK-WT or, -7S/TA mutant, and the quantitative analysis is shown on in (**I**). Data are shown as means ± S.D. (n=3). Scale bar, 100 μm.

(**J**) Parental and IBTK-KO SiHa cells were treated with silvestrol (100 nM) for 24 h. Then, annexin-V-FITC/PI dyes were used to stain the harvested cells, of which later flow cytometry analysis was performed. Data are shown as means ± SD. (n=3).

P values are calculated using One-way ANOVA test in (**G**, **I**), Two-way ANOVA test in (**E, J**). *$p<0.05$, **$p<0.01$, ***$p<0.001$, ****$p<0.0001$, n.s. non-significant.

Figure.7 Overexpression of IBTK correlates with poor survival in cervical cancer.

(**A**) Kaplan–Meier survival plots of overall survival (OS) were analyzed according to IBTK mRNA expression in CESC specimens from TCGA cohort (n=295).

(**B, C**) IHC analysis of IBTK protein expression in 35 CESC tissues and 35 adjacent normal cervical tissues. Representative staining results were shown. Scale bar, 30 μm. The relative intensity of IBTK was measured using ImageJ and the quantitative data is shown in (**C**).

(**D**) IBTK expression in samples with different pathological grade.

(**E**) Kaplan–Meier survival curves of overall survival comparing high and low

expression of IBTK in the TMA (n=117).

(**F**) A proposed model depicting that mTORC1-S6K1 signaling promotes translation

initiation by enhancing IBTK-mediated eIF4A1 ubiquitination.

P values in (**A**, **E**) are calculated using Log-rank test while *P* values in (**C**, **D**) are

calculated using unpaired Student's t test. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001,

n.s. non-significant.