Divergent Molecular Pathways for Toxicity of Selected Mutant C9ORF72-derived Dipeptide Repeats

Yuyu Song[1,2,3*], Ming Ying Tsai[4], Bin Wang[4], Priscila Comassio[3,5], Jorge E. Moreira[3,4], Nicola Kriefall[3], Gerardo Morfini[3,4*], and Scott Brady[3,4*]

[1] Department of Neurology, Massachusetts General Hospital, Charlestown, MA 02129, USA

[2] Dept. of Neurology, Harvard Medical School, Boston, MA 02114, USA

[3] Marine Biological Laboratory, Woods Hole, MA 02543, USA

[4] Department of Anatomy, University of Illinois at Chicago, Chicago, IL 60612, USA

[5] Department of Neurology, Neuroscience and Behavior, Ribeirão Preto School of Medicine, University of São Paulo, Ribeirão Preto, SP 14049-900, Brazil

* Corresponding Authors

Yuyu Song

Department of Neurology, Mass General Hospital and Harvard Medical School

114 16th St., Office 2012, Charlestown, MA 02129 USA

Phone: 617-724-8347

Email: ysong13@mgh.harvard.edu

Gerardo A. Morfini

Department of Anatomy and Cell Biology, University of Illinois at Chicago, College of Medicine

808 S Wood St MC512, Chicago, IL 60612 USA.

Phone: (312) 996-6869

Email: gmorfini@uic.edu

Scott T. Brady

Department of Anatomy and Cell Biology, University of Illinois at Chicago, College of Medicine

808 S Wood St MC512, Chicago, IL 60612 USA.

Phone (312) 996-6791

Email: stbrady@uic.edu

ABSTRACT

Expansion of a hexanucleotide repeat in a noncoding region of the C9ORF72 gene is responsible for a significant fraction of Amyotrophic Lateral Sclerosis (ALS) and Frontotemporal Dementia (FTD) cases, but identifying specific toxic gene products and mechanisms has been difficult. Pathogenesis was proposed to involve the production of toxic RNA species and/or accumulation of toxic dipeptide repeats (DPRs), but distinguishing between these mechanisms has been challenging. In this study, we first use complementary model systems for analyzing pathogenesis in adult-onset neurodegenerative diseases to characterize the pathogenicity of DPRs produced by Repeat Associated Non-ATG (RAN) translation of C9ORF72 in specific cellular compartments: isolated axoplasm and giant synapse from the squid. Results showed selective axonal and presynaptic toxicity of GP-DPRs, independent of associated RNA. These effects involved downstream ASK1 signaling pathways that affect fast axonal transport and synaptic function, a pathogenic mechanism shared with other mutant proteins associated with familial ALS, like SOD1 and FUS. These pathways are sufficient to produce the "dying-back" axonopathy seen in ALS. However, other mutant genes (e.g., SOD1) that activate this mechanism rarely produce FTD. When parallel studies in primary motor neurons from rats were conducted, an additional pathogenic mechanism was revealed. The GR- and PR-DPRs, which had no effect on axonal transport or synaptic transmission, were found to disrupt the nuclei of transfected neurons, leading to "dying-forward" neuropathy. All C9-DRP-mediated toxic effects observed here are independent of whether the corresponding mRNAs contained hexanucleotide repeats or alternative codons. These studies establish the divergent toxicity of C9-DPRs that cause neurodegeneration in ALS and FTD, suggesting that these two independent pathogenic mechanisms may contribute to disease heterogeneity and/or synergize on disease progression in C9ORF72 patients with both ALS and FTD symptoms.

Graphic Abstract: Activation of protein kinases and inhibition of axonal transport, synaptic transmission, and nuclear structure are toxic effects common to unrelated FALS-related gene products. FALS-related mutant forms of SOD1 (mSOD1), FUS (mFUS), and C9-GP-DPRs ($GP_{(n)}$) activate specific ASK1-MAPK pathway. Within axons, active ASK1-p38 pathway phosphorylates various substrates, including conventional kinesin, leading to the inhibition of fast axonal transport mediated by the translocation of this motor protein along microtubules. ASK1 can also inhibit synaptic transmission via JNK activation. Both pathways cause reductions in the availability of critical synaptic cargoes, synaptic dysfunction, and "dying-back" degeneration of neurons. On the other hand, C9-PR and GR-DPRs ($PR_{(n)}$ and $GR_{(n)}$) activate other pathways, leading to aberrant alterations in nuclear structure and function and "dying-forward" degeneration of neurons, consistent with reports of transcriptional changes and activation of apoptosis in ALS.

INTRODUCTION

The surprising discovery that mutations in the C9ORF72 gene were involved in a significant number of both familial (~40%) and sporadic (~7%) Amyotrophic Lateral Sclerosis (ALS) cases, as well as ~25% of Frontal Temporal Dementia (FTD) cases [1] presented a conundrum. The demonstration that the mutation involved expansion of a hexanucleotide repeat (GGGGCC) in a non-coding region of a gene of unknown function [2,3] added further complexity to the mystery. How do multiple mutations in a wide range of unrelated genes all render upper and lower motor neurons increasingly vulnerable in ALS[4-6]? Both loss of function and gain of function mechanisms have been proposed to explain pathogenesis in ALS, and C9ORF72 mutations were no exception[7-9]. Loss of C9ORF72 function associated with altered transcription or splicing has been suggested to alter immune function [10] or autophagy pathways[11-13]. However, neither conditional nor ubiquitous deletions of the mouse C9ORF72 ortholog produces motor neuron degeneration[7]. Instead, total knockout of the gene produces a fatal autoimmune disease[14]. Many studies suggesting gain-of-function mechanisms have focused on RNA toxicity associated with the hexanucleotide repeats and based on the RNA foci observed in affected cells [15,16]. Concurrently, other studies have implicated the accumulation of dipeptide repeat proteins (DPRs)[17] produced by repeat-associated non-ATG (RAN) translation[17-19]. GP_n is produced by translation in both forward and reverse directions. GA_n and GR_n DPRs are generated by translation of expanded hexanucleotide sequences in the forward direction, whereas PR_n and AP_n DPRs are produced by translation in the reverse direction. These DPRs have been shown to accumulate in affected cells including cell bodies, distal axons, and synaptic terminals[20,21]. Still, others have suggested alterations on nucleocytoplasmic trafficking, which could be due to either mutant RNAs or DPRs[22,23]. Although these mechanisms are not mutually exclusive, dissecting the relative contribution of these various mechanisms has been problematic, given the lack of model systems that can simultaneously distinguish these mechanisms. Moreover, the molecular basis for the increased vulnerability of motor neurons has remained elusive, making the identification of therapeutic targets challenging.

In this study, we exploited two unique model systems for analyzing pathogenic mechanisms underlying adult-onset neurodegenerative diseases: isolated axoplasm from the squid giant axon [24] and the squid giant synapse[25-27]. Using these systems, we characterized the pathogenicity of five DPRs associated with RAN translation of mutant C9ORF72 (GP, GA, GR, PA and PR) in axons and synapses, major ALS-vulnerable neuronal compartments. Remarkably, GP, but not other DPRs elicited toxic effects on fast axonal transport, a cellular process critical

for sustained axonal health known to be affected in ALS. As reported before with mutant SOD1[28,29], inhibition of an ASK1-P38 MAP kinase cascade mitigated this toxic effect. Given that mutant SOD1 also affects synaptic function[30], the effect of DPRs in the presynaptic terminal of the squid giant synapse was also examined. In this model system, GP, but not GR, caused deficits in synaptic function that were also prevented by ASK1 inhibition.

Concurrently, we evaluated the toxicity of specific DPRs in mammalian models by transfection of rat primary cultured embryonic motor neurons. Consistent with results in squid models, expression of GP-DPR produced a striking axonopathy consistent with the "dying-back" pattern of degeneration observed in ALS-affected neurons[31], which was reduced by inhibition of ASK1 and P38 MAP kinases. As seen in the squid models, neither GA nor PA DPRs affected axonal transport or cell viability.

In contrast with results obtained from isolated squid axons, two additional C9ORF72 DPRs (GR and PR) elicited cytotoxic effects on both fibroblasts and cultured motor neurons. Expression of GR and PR DPRs resulted in somal and nuclear pathology, rather than a distal axonopathy, and the effect was not blocked by inhibition of either ASK1 or P38 MAP kinase, suggesting a "dying-forward" pathology *via* a distinct molecular mechanism. The effects of GP, GR, and PR DPRs in cultured neurons were independent of whether the mRNA encoding these dipeptides contained hexanucleotide repeats or used alternative codons, which would not form atypical secondary structures of RNA, indicating that the observed toxicity was not due to RNA-based mechanisms.

Previous reports demonstrated that activation of P38 MAP kinase is a hallmark of both familial and sporadic forms of ALS[32-36]. Further, activation of ASK1 and P38 MAP kinase is a common feature of multiple gene products implicated in ALS pathology, including mutant and misfolded SOD1[28,29,37,38] and FUS[39]. Here, our results using ASK1 and P38 inhibitors suggest that, for C9ORF72 mutations, the toxicity of GP DPRs may be primarily responsible for motor neuron pathology *via* activation of an ASK1-MAPK pathway commonly shared by all other ALS-associated proteins examined to date. However, PR and GR DPRs may contribute to the development of neurodegeneration *via* an unknown pathway specific for patients with the expansion of the hexanucleotide repeat in C9ORF72.

RESULTS

GP dipeptide repeat protein inhibited axonal transport via the ASK1-P38 pathway.

ALS proceeds as a dying back axonopathy with loss of distal axons and synapses as early pathological events, suggesting that the axonal compartment is a critical pathogenic target in ALS-affected neurons[31]. Previous studies documented inhibition of fast axonal transport by ALS-linked forms of FUS and SOD1 *via* activation of ASK1-P38 MAP kinase and phosphorylation of kinesin heavy chains[28,29,37,39], raising the question of whether axonal P38 MAP kinase activation is a general feature of additional familial ALS (fALS)-linked mutant gene products. C9ORF72 is the gene most frequently mutated in ALS [1,5], but it is unusual in being a hexanucleotide repeat expansion that can produce both multiple distinct dipeptide repeat proteins (DPRs) through RAN translation and abnormal RNA foci.

RAN translation of the hexanucleotide repeat in mutant C9ORF72 resulted in the accumulation of five different DPRs of various lengths (GP, GR, PR, GA, PA) in patient brains[18,40,41]. To evaluate toxic effects of individual DPRs in the axonal compartment, we performed vesicle motility assays following perfusion of synthetic DPRs (16-mer peptides encompassing 8 dipeptide repeats) at the same concentration into isolated squid axoplasm (**Fig. 1**). This *ex vivo* model lacks nuclear transcription, protein synthesis machinery, and glia, allowing testing of axonal transport-specific mechanisms. Since the DPRs were added as peptides, there was no RNA that could generate RNA toxicity. The absence of a plasma membrane allows the introduction of neuropathogenic proteins and pharmacological agents at defined concentrations, and the quantitative analysis of fast axonal transport of membrane-bounded organelles (MBOs) moving in both anterograde (kinesin 1-dependent) and retrograde (cytoplasmic dynein-dependent) directions [24].

Among the five DPRs, only GP_8 (Fig 1A) affected vesicle motility in axoplasm by preferentially inhibiting anterograde transport, an effect similar to that observed for mutant SOD1(G93A and G85R)[28,29,37] (Fig 1F) and FUS[39]. The effects of GA_8, GR_8, PR_8, and AP_8 (Fig 1B-E) were not different from buffer control, with no change in axonal transport rates after perfusion over 50min. Quantitative analysis of these data indicated that only GP_8 and mutant SOD1(G93A) significantly inhibited the anterograde transport of vesicles (Fig 1).

Selective inhibition of anterograde transport by GP_8, an effect similar to that elicited by pathogenic forms of SOD1[28,29,37] and FUS[39] suggested that GP_8 may also activate a similar pathway. Pharmacological and biochemical experiments indicated that mutant SOD1 activates a MAP kinase signaling cascade culminating in phosphorylation of kinesin heavy chain by P38α

MAPK[28,29]. This cascade involved the activation of the MAP kinase kinase kinase (MAP3K) ASK1[29,39], which in turn activated a MAP kinase kinase (MAP2K) leading to p38 MAPK activation[28]. To determine whether GP_8 activated a P38α MAPK-linked cascade in axoplasm, we co-perfused axoplasms with GP_8 and MW069, a selective inhibitor of P38α MAPK[42]. MW069 prevented inhibition of anterograde transport by GP (Fig. 2A), as it did for mutant SOD1 (Fig. 2D), suggesting a critical role for P38α MAPK in axon-specific toxicity of GP. Co-perfusion of GP_8 with a DVD peptide, which prevents a subset of MAP3Ks from docking to and activating downstream MAP2Ks[43], also prevented GP_8 from inhibiting kinesin-based anterograde transport (Fig 2B). Similarly, co-perfusion with NQDI-1, an inhibitor of the MAP3K ASK1[44], prevented inhibition of anterograde transport by GP_8 (Fig 2C), as it did for mutant SOD1[29] and FUS [39]. These studies suggest that pathological forms of SOD1, FUS, and C9ORF72-GP_8 disrupt fast axonal transport through a common pathway: activation of the ASK1-P38 MAP kinase cascade.

To provide additional evidence that inhibition of axonal transport by GP_8 was through specific activation of P38 MAPK pathway, we examined the phosphorylation of kinesin and various candidate kinases in axoplasms perfused with DPRs, as described previously[45] (Fig 3). In the presence of ^{32}P-γ-ATP, GP_8 perfusion increased phosphorylation of Kinesin Heavy Chain (KHC) subunits, as shown before with mutant SOD1[28] (Fig 3A-B). Further, increased levels of activated (phosphorylated) P38 MAPK were observed in axoplasms following GP_8 perfusion, whereas levels of active GSK3 and ERK kinases remained unchanged (Fig 3C-F). Unlike GP_8, GA_8 perfusion had no effect on phospho-P38 levels (Fig 3G-H). Collectively, these results suggested that GP DPRs can activate p38 MAPK and aberrant phosphorylation of the motor protein kinesin in the axonal compartment.

GP dipeptide repeat protein inhibited synaptic transmission via ASK1.

That GP DPRs inhibited axonal transport by activating a MAP kinase cascade, similar to mutant SOD1 that also inhibited synaptic transmission, raised the question of whether GP and other DPRs also affected synaptic transmission[26,27,30]. To address this question, we used the squid giant synapse as a model to mimic mammalian neuromuscular junctions, where synthetic DPRs were microinjected into the presynaptic terminal site, and membrane potentials were measured in both presynaptic and postsynaptic sites to indicate the strength of synaptic transmission (Fig 4A). While presynaptic injection of GR_8 did not affect postsynaptic membrane

potential (PSP) for over 250min after injection (Fig 4B), GP_8 gradually inhibited PSP to the extent that postsynaptic action potential was no longer elicited within 200min after injection (Fig 4C). NQDI, a small molecule inhibitor of ASK1, blocked the inhibitory effect of GP_8 and maintained steady synaptic transmission, similar to the synapses injected with GR_8, as indicated by the constant PSP slope quantified over 250min (Fig 4D).

Mutant SOD1 was previously shown to affect synaptic function by reducing available synaptic vesicle pools[30]. To examine whether GP_8-induced synaptic defects also include an action on synaptic vesicle pool dynamics, trains of high-frequency stimulations (HFS, 50Hz for 5s, with 5s intervals between trains) were applied presynaptically 15min post-injection and postsynaptic depression of EPSPs were measured. Normally there is a two-phase diminution of EPSP slope across the 250 spikes[30]. The first phase is associated with the utilization of the Readily Releasable Pool (RRP) of synaptic vesicles, while the second phase reflects Reserved Pool (RP). GR_8-injected synapses quickly recovered from HFS (Fig 4F), suggesting abundant supplies of both RRP and RP. However, GP_8-injected synapses were significantly inhibited (Fig 4E).

To further illustrate the vesicle pool dynamics, the EPSP slopes were integrated to generate a trace, where the last 50 time points were linearly fitted (Fig 4G). Both the relative size of the RRP, determined by the intersection of the linear fit with the y-axis, and the mobilization rate from the RP to RRP, determined by the slope of the linear fit, showed reductions in GP_8-injected synapses, as compared with the GR_8-injected synapses, which behave similarly to non-injected controls (Fig 4G-H). EM analysis of the active zones confirmed the reduction in total synaptic vesicles, including the large vesicles and docking vesicles, in synapses injected with GP_8, but not in synapses injected with GR_8 (Fig 4I-K). The effects of GP_8 on synaptic physiology and morphology are similar to those observed following mutant G85R-SOD1 injection in the giant synapse[30].

C9ORF72-GP, RP, RG dipeptide repeat proteins induced axonal degeneration in mammalian neurons.

To validate the results showing toxicity of GP DPRs in the axon and the synapse from squid, we evaluated toxicity for all five dipeptides expressed in primary cultured mammalian motor neurons. Transfection of motor neurons with recombinant mCherry constructs that had specific 25-mer DPRs added to their C-terminals showed that mCherry-GP produced axonal

degeneration, while mCherry-GST control had no effect on motor neurons (Fig 5A-B). The effect on motor neurons was the same whether the mRNA sequence encoding for the mCherry-GP dipeptide used the GGGGCC hexanucleotide repeat or used various alternative codons to generate a dipeptide tail on the mCherry without nucleotide repeats at the RNA level (Fig 5C-D). The effects of mCherry-GP were mitigated by treatment with the P38 MAP kinase inhibitor SB203580 (Fig S1) or by treatment with an ASK1 inhibitor (NQDI) (Figs S2-S3), consistent with the rescuing effects of P38 MAP kinase/ASK1 inhibition in the squid axoplasm and giant synapse models. Also as seen in the squid models, neither mCherry-GA nor mCherry-PA affected motor neuron viability regardless of being encoded by either GGGGCC repeats or alternate codons. Contrasting with results in axoplasm and the giant synapse models, transfection of motor neurons with either mCherry-PR or mCherry-RG constructs was toxic to mammalian motor neurons (Fig 5). As with mCherry-GP, toxicity was seen whether GGGGCC repeats or alternative codons were used (5C-D). However, unlike GP, inhibitors of P38 MAP kinase or ASK1 failed to rescue neurons expressing mCherry-PR and mCherry-RG (Figs S1-S3), suggesting that the toxic effects elicited by mCherry-RP and mCherry-RG involve a pathogenic mechanism distinct from that triggered by mCherry-GP.

Although GP, RG, and PR DPRs were all toxic to cultured motor neurons, the phenotype of neurons expressing GP was distinct from that of neurons expressing either PR or RG (Figs 6, S4-S7). Specifically, motor neurons (12 DIV) transfected with mCherry-GP exhibited axonal degeneration with fragmented axons, accumulation of kinesin, and axonal swellings 10 hours after transfection, similar to effects elicited by SOD1(G93A) and FUS (R521G) (Fig. S4). However, the cell bodies and nuclei remained relatively intact with dispersed chromatin (DNA stained with DAPI) and an intact nuclear envelope (labeled by Lamin A/C) (Fig S5). In contrast, axons of motor neurons transfected with either mCherry-RG or mCherry-RP remained largely intact 10 hours after transfection, and kinesin distribution was not altered (Fig. S4). Instead, transfection with mCherry-RG or -RP produced neuronal degeneration in the soma while initially sparing the axonal compartment. Both mCherry-RG and -RP expressing cells exhibited striking changes in nuclear structure, with clumping of chromatin (DNA) and disruption of the nuclear envelope (Lamin A/C) (Figs S6-S7). The effect of transfecting neuron with an empty mCherry vector was minimal after 24 hours. While GP mainly affected axons causing "dying back" degeneration (i.e., from axon to cell body), RP and RG primarily affected nuclear structure consistent with a "dying forward" degeneration (i.e., from cell body to axon).

In addition, mCherry-RP and mCherry-RG exhibited cytotoxicity for both neuronal and non-neuronal cells, whereas mCherry-GP was more toxic for motor neurons than non-neuronal cells. This difference was also observed when non-neuronal mammalian cells (COS 293) were transfected with the same DPR constructs (Figs S8-9). The mCherry-GP transfected cells were not significantly affected, with little evidence of toxicity. Cells transfected with mCherry-RP or mCherry-RG exhibited clear cytotoxicity with nuclear pathology that included aggregation of chromatin (DNA, Fig S8) and disruption of the nuclear envelope (Lamin A/C, Fig S9), similar to the effects of these dipeptides in neurons.

DISCUSSION

ALS is defined by a clinical presentation in which both upper and lower motor neurons are compromised. The reasons for the selective vulnerability of these two neuronal populations continue to be debated and lead to a second question: do all cases of ALS share a common pathogenic mechanism, or is ALS a group of distinct diseases with a common clinical presentation? Given that >90% of ALS cases are idiopathic with no family history and that the remaining familial cases of ALS have been linked to mutations in more than 40 different genes that are otherwise unrelated, answers to this question are critical in the search for therapeutic interventions.

The discovery that expansion of a hexanucleotide repeat in the C9ORF72 gene was associated with a surprisingly large number of both familial and sporadic forms of ALS provided a new challenge as well as an opportunity to understand ALS pathogenesis. The situation is further complicated by the fact that the same hexanucleotide expansion in C9ORF72 can also produce FTD or symptoms of both FTD and ALS in patients[2,3,7]. The frequency of different clinical presentations is roughly 25% ALS, 50% ALS/FTD, and 25% FTD[46,47] and all three phenotypes can be observed in a single family with the same mutation. A single mutation that can variably affect two distinct neuronal populations is unusual, particularly since some other mutant genes associated with ALS, like SOD1, are rarely if ever associated with FTD[4]. The heterogeneity of neurological phenotypes seen with mutant C9ORF72 raises two questions: First, do cases of ALS due to C9ORF72 mutation involve the same molecular mechanisms as cases of ALS associated with other gene mutations or sporadic forms of ALS and, second, are the clinical symptoms of ALS and FTD produced by different pathogenic mechanisms or is there a single shared mechanism?

The C9ORF72 mutation differs from other gene mutations associated with familial ALS in two ways. First, the mutation is an expansion of a hexanucleotide (GGGGCC) repeat that may have hundreds of repeats. Second, the hexanucleotide repeat is in an intron of the gene that is not normally transcribed. Three possible mechanisms for C9ORF72 toxicity were initially proposed: 1) loss of C9ORF72 protein function, 2) mutant C9ORF72 RNA-induced toxicity, and 3) toxicity of DPRs translated from these RNAs. Any mechanism or combination of mechanisms must explain the heterogeneity of disease phenotypes (i.e., ALS, ALS/FTD and FTD) seen with C9ORF72 mutations. ALS and FTD involve two distinct neuronal populations. These two neuronal populations appear to be differentially vulnerable because there are other mutant genes that produce only ALS or only FTD. The variable presentation of disease phenotypes with C9ORF72 mutations raises the possibility that multiple pathogenic mechanisms are involved.

Loss of function was largely eliminated as a significant contributor to the development of either ALS or FTD[9] because deletion of the C9ORF72 gene fails to produce either neurological phenotype but results in a fatal autoimmune disease[7]. This leaves RNA toxicity and/or DPR toxicity as candidate mechanisms. These two mechanisms are not mutually exclusive, so we tested both possibilities, with a focus on DPRs, in this study.

RNA toxicity is dependent on the hexanucleotide repeat forming unique RNA structures that might disrupt RNA processing or nucleocytoplasmic transport. The presence of RNA foci in affected cells was used to support this proposed mechanism, and the focus has been primarily on changes in nucleocytoplasmic trafficking because FUS and TDP43, two other genes associated with familial ALS, have a role in nucleocytoplasmic trafficking of RNA[48]. In addition, cytoplasmic aggregates of wild type TDP43 can be seen in many cases of ALS and FTD in the absence of mutations. However, the question remains as to how RNA toxicity due to hexanucleotide expansion could produce either pure ALS or pure FTD as well as ALS/FTD.

The third candidate mechanism proposes that one or more of the DPR-containing polypeptides generated by RAN translation are toxic[17-19]. RAN translated DPRs have been detected in multiple neurological diseases with repeat expansions and are implicated in pathogenesis[18,19]. Variable levels of the five DPRs resulting from C9ORF72 mutations have been reported in cell bodies, distal axons, and synaptic terminals of affected cells[20,21]. These DPRs could also affect RNA processing or trafficking. Alternatively, the DPRs could affect neuronal signaling pathways important for the regulation of fast axonal transport, as seen with mutations in SOD1[28,29,37] and FUS[39], or synaptic transmission, as shown for mutant SOD1[30].

The possibility that mutant C9ORF72-related DPRs might affect axonal and synaptic signaling is particularly intriguing because one pathological change shared by affected motor neurons in both familial and sporadic cases of ALS is increased activity of the ASK1-MAP kinase pathway[32,33,37]. Our previous studies showed that several seemingly unrelated mutant genes associated with familial ALS activate the same neuronal signaling pathway, ASK1 MAP kinase. These include familial ALS-linked mutant forms of SOD1[28-30,37,38], an oxidized, misfolded form of wild type SOD1[37], and mutant forms of the proteins FUS[39] and profilin 1 (unpublished observations). Aberrant activation of the ASK1-p38/JNK MAP kinase signaling pathway in neurons has a number of potential consequences, including alterations of gene transcription, inhibition of axonal transport[28,29] and impairment of synaptic transmission[30]. These findings raised the question of whether the C9ORF72 mutation could also affect the ASK1 MAP kinase pathway or if a different pathogenic mechanism was involved.

To evaluate axonal and synaptic pathogenicity of DPRs, we tested the pathogenicity of DPR-proteins associated with RAN translation of C9ORF72 in two unique model systems: isolated axoplasm and the giant synapse. These models allow analysis of DPR toxic effects independent of any contributions associated with the nucleus or RNA. Recombinant proteins containing each of the DPRs were perfused into the isolated axoplasm or injected into the giant synapse of the squid. Both axonal transport and synaptic transmission were inhibited by GP-DPR but were unaffected by any of the other four C9ORF72 DPRs. In previous studies with SOD1[28,29,37] and FUS[39], inhibition of axonal transport and of synaptic activity was dependent on activation of the ASK1 dependent kinase pathways. Similarly, the GP-DPR inhibitory effect on transport involved activation of ASK1-P38 kinase signaling as both an ASK1 inhibitor and P38 inhibitor prevented GP-DPR-induced axonal transport impairment. In the squid giant synapse preparation, synaptic inhibition by GP-DPR appears to be mediated by ASK1-JNK pathway as the ASK1 inhibitor, but not the P38 inhibitor, prevented GP-DPR-induced synaptic damage (Fig 4 and unpublished observations). As shown previously, other familial ALS-associated mutant proteins (e.g., mutant SOD1, FUS) and sporadic ALS-associated oxidized SOD1 protein all inhibited axonal transport *via* activation of a ASK1/P38 pathway, and ALS-associated mutant SOD1 protein inhibited synaptic transmission *via* the activation of ASK1-JNK pathway [30,49]. Altogether, these findings suggested a shared molecular component (i.e., ASK1-MAPK pathway) underlying common cellular dysfunctions (i.e., axonal transport inhibition and synaptic transmission deficits) that may lead to axonal degeneration in ALS-affected neurons, regardless of the mutant genes in familial cases and posttranslational modifications in sporadic cases, presenting ASK1 as a potential drug target.

To validate these discoveries in mammalian neurons as well as explore additional pathological mechanisms, we expressed each of the five C9ORF72 DPR-proteins in mouse primary motor neurons. As predicted, GP-DPR induced axonal damage causing dying-back degeneration, similar to the effects seen with mutant SOD1, and initially sparing the neuronal cell body. However, unlike the results seen in isolated axoplasm and the presynaptic terminal of squid that lack cell bodies, two additional C9ORF72 dipeptide repeat proteins, RG-DPR and RP-DPR, were also neurotoxic. Unlike GP-DPR, the RP- and RG-DPRs produced somal defects consistent with a dying-forward degeneration while initially sparing the axons. Significantly, though not surprisingly, inhibition of ASK1 rescued neurons expressing GP-DPR but failed to prevent RG/RP-DPR-mediated somal damage and neurodegeneration, suggesting that a different molecular mechanism for toxicity is involved.

As in the squid axoplasm, PA- and GA-DPRs had no effect on neurons. Since all five of the DPR-proteins included the same hexanucleotide expansion in different reading frames to produce specific DPRs, the lack of toxicity for PA- and GA-DPRs would not be consistent with a role for RNA toxicity in the observed phenotypes for GP-, RP- or RG-DPRs. To confirm that the observed toxicity of the DPR effects were independent of the RNA sequence with the hexanucleotide expansion, we repeated the experiments using transfected recombinant proteins in which the DPR sequence were encoded by alternative codons with no repeats. The RNAs produced by these plasmids would not have the same RNA structure but would express the same DPR. The phenotypes were identical for a given DPR, whether it was produced by plasmids with the GGGGCC hexanucleotide repeats or by alternative codons. The phenotypes seen in this study were the result of the corresponding DPRs, not by RNA. This does not exclude the possibility of an additional toxic gain function for RNAs, but it demonstrates that the DPRs are responsible for at least two distinct toxic gains of function differentially affecting the cell body and axon/synapse.

To determine whether C9ORF72 dipeptide repeat proteins affect non-neuronal cells and characterize the phenotype in greater detail, we expressed the same C9ORF72-DPRs in COS293 cells. RP/RG-DPRs cause cell loss and nuclear envelope damage, similar to effects in neurons, while GP-DPR has no effect on nuclear structure or chromatin and exhibited no cytoxicity in COS293 cells. This is consistent with published literature showing toxicity of RP/RG-DPR in non-neuronal cells [50]. This also raises the possibility of glial toxicity due to RP/RG-induced somal toxicity in ALS, which may affect both motor and frontotemporal cortical neurons.

Neurons are dependent over a lifetime on maintaining active fast axonal transport and synaptic transmission over long distances, which may be as much as a meter or more for motor neurons in an adult human. Similarly, carefully regulated nucleocytoplasmic transport and nuclear function are essential for somal health and protein synthesis in mature neurons. The heterogeneous phenotypes that can be seen with hexanucleotide expansion may result from synergistic toxicity of GP-DPR and RP/RG-DPR. Since RAN translation involves random initiation, the relative levels of each DPR in various subcellular compartments of a given patient may be variable. The GP-DPR would be expected to affect upper and lower motor neurons through an ASK1-mediated pathway and produce a dying back axonopathy, much like mutant and misfolded SOD1, suggesting that the GP-DPR may be sufficient to produce an ALS phenotype. Since RP- or RG-DPRs activate a different pathogenic mechanism that is cytotoxic, drastically disrupting nuclear structure and function in both neuronal and non-neuronal cells, the presence of these DPRs may exacerbate the ALS phenotype as well as affecting additional neuronal populations such as in FTD. RP- and RG-DPRs could also produce dying forward degeneration in neurons by affecting glial cells and vasculature. The ability of mutant C9ORF72 to activate at least two independent, selective DPR-mediated pathogenic mechanisms may explain the existence of multiple distinct neuropathological phenotypes, with one pathogenic pathway contributing to ALS and the other to FTD.

In sum, our study demonstrated a shared ASK1 pathway underlying axonal and synaptic dysfunctions leading to dying back degeneration of motor neurons in ALS with mutations in genes seemingly unrelated, including mutant C9ORF72. This shared pathway is activated by the GP-DPR. In addition, the RP- and RG-DPRs activate a different pathogenic mechanism, which may synergize to render motor neurons selectively vulnerable in ALS as well as the cortical neurons affected in FTD. The nuclear toxicity of RP and RG on neuronal and non-neuronal cells such as glia may further contribute to the disease pathology and include aberrant activation of additional kinase pathways with crosstalk with the ASK1 MAPK signaling. The multiplex disease pathology associated with C9ORF72 may call for a precision-medicine approach and cocktail therapy targeting multiple molecular components in a given patient.

Methods

Preparation of Dipeptides. Dipeptides corresponding to the five identified DPRs produced by RAN translation of expanded C9ORG72 hexanucleotide repeats (16-mer) were chemically

synthesized and sequenced by the UIC Research Resources Center. Stocks of GP(8), GR(8), GA(8), PR(8) and PA(8) DPRs were prepared as a 5 mM (GR) or 10 mM (GP) stock in 50 mM HEPES, ph7.4 or 5 mM DMSO (GA).

Vesicle Motility assays in isolated axoplasm. Axoplasms were extruded from giant axons of the squid Loligo pealii (Marine Biological Laboratory) as described previously [24]. Briefly, the two largest axons are removed from wild-caught squid, and associated tissue is removed. Once cleaned, the axon can be extruded onto a coverslip to leave an intact cylinder axonal cytoplasm (axoplasm), which continues to maintain organelle trafficking in both directions for hours[51,52]. Anterograde and retrograde transport is visualized using video-enhanced contrast differential interference contrast microscopy Zeiss Axiomat with a 100X, 1.3 n.a. objective, and DIC optics. A Hamamatsu Argus 20 and Model 2400 CCD camera system were used for image processing and analysis. Organelle velocities were measured interactively with a Photonics Microscopy C2117 video manipulator (Hamamatsu). Purified proteins (i.e., SOD1), DPRs and kinase inhibitors were all diluted to 1μM final concentration into X/2 buffer (175 mM potassium aspartate, 65 mM taurine, 35 mM betaine, 25 mM glycine, 10 mM HEPES, 6.5 mM $MgCl_2$, 5 mM EGTA, 1.5 mM $CaCl_2$, 0.5 mM glucose, pH 7.2) supplemented with 5 mM ATP) and 25 μl added to perfusion chambers (Fig 1). Both anterograde and retrograde FAT were analyzed by video-enhanced microscopy over 50 minutes, as before[24].

Biochemical experiments in isolated axoplasm. Biochemistry was conducted as described previously[45]. Using two axoplasms from a single squid, one was perfused with control buffer and one was perfused with 1μM GP_8 in Buffer X/2 + 5μM ATP for comparison. For experiments shown in Fig 3A-B, γ-^{32}P-ATP was included in the perfusate. After a 50 min incubation, axoplasms were homogenized and processed for kinesin immunoprecipitation using the H2 antibody against kinesin heavy chain then processed for autoradiography as described previously[45]. Briefly, axoplasm lysates were precleared using a mixture of Protein G agarose beads (Pierce), and non-immune mouse IgG-conjugated Sepharose beads (Jackson Immunoresearch) for 1 hour at RT. After a short spin, supernatants were incubated with H2 antibody and Protein G beads for 4 hours. H2-kinesin immunocomplexes were recovered by centrifugation (3000 g_{max} for 30 seconds), washed four times with 1 ml lysis buffer, once with 50mM HEPES pH 7.4, and resuspended in Laemmli buffer. For immunoblots in Fig 3C, axoplasm lysates were separated by SDS-PAGE on 4-12% Bis-Tris gels (NuPage minigels, Invitrogen), using MOPS Running Buffer (Invitrogen) and transferred to PVDF using Towbin buffer supplemented with 10% (v/v) methanol (90 minutes at 400mA using Hoeffer TE22

apparatus). Immunoblots were blocked with 1% (w/v) non-fat dried milk diluted in TBST (25 mM Tris pH 7.2, 2.68 mM KCl, 136.8 mM NaCl, 0,01% Tween-20, ph: 7.2). Sodium Orthovanadate (1 mM) and Sodium Fluoride (10 mM) was used in all incubation steps involving the use of phosphoantibodies, after correcting for pH. Membranes were incubated with primary antibodies overnight at 4° C in 1% IgG free-BSA (Jackson Immunoresearch), and washed four times with 0.1% Tween-20 in TBS. Primary antibody binding was detected with HRP-conjugated anti-mouse, anti-rabbit or anti-goat antibodies (Jackson Immunoresearch), and visualized by chemiluminescence (ECL, Amersham).

Giant Synapse Electrophysiology. The giant squid synapse is formed between the terminal finger of second order axons of the pre-nerve and the third order axon of the last stellate nerve, which is the giant axon used for axonal transport studies as well as for classical voltage-clamp studies of ion channels[53]. Giant synapse was prepared as described [30] and superfused continuously with oxygenated squid saline (455 mM NaCl, 54 mM $MgCl_2$, 11 mM $CaCl_2$, 10 mM KCl, 3 mM $NaHCO_3$, and 10 mM HEPES, pH 7.2) at 10–15°C. One microelectrode for injecting was inserted in the presynaptic axon to inject current at 0.033 Hz for basal stimulation and at 50 Hz for HFS (each pulse is 2 µA for 2 ms for basal stimulation and 2 µA for 1 ms for HFS), near the palm where the second order axon enters the ganglion to branch and form multiple synapses with postsynaptic axons. At the terminal of the most medial branch, a second microelectrode was inserted to presynaptically inject proteins and reagents of interest as well as recording the presynaptic membrane potentials. Finally, at the postsynaptic terminal of the giant synapse, a third microelectrode was inserted near the medial presynaptic digit to record the postsynaptic membrane potentials. Electrodes 1 and 3 were filled with 3 M KCl while the second microelectrode allowed the injection of 50 µM SOD1 proteins or reagents of interest in 100 mM KCl[54] at 0.1 Hz (each injection was 50 psi for 250 ms). The injection efficiency of peptides was monitored by a fluorescence microscope, which ensures comparable levels of infused peptides. Both presynaptic potential and PSP were recorded using sharp microelectrodes with an Axoclamp-2A amplifier (Axon Instrument) and data analyzed using Labview (National Instruments) software (Yulong Li). Raw waveforms were used to calculate PSP slope using the same parameters for all experiments. For data acquired under HFS, PSP slopes were analyzed, integrated, plotted, and fitted linearly using the last 50 time points to derive the vesicle mobilization rate (slope of the linear fit) and the RRP size (intersection with the *y*-axis).

Electron Microscopy of the giant synapse. After recordings, synapses were prepared and imaged as published [30]. Briefly, synapses were fixed in 4% glutaraldehyde and 2% paraformaldehyde in 0.1 M cacodylate buffer for 12 h at 4°C, then washed and postfixed in 1% osmium tetroxide, followed by blocking impregnation with 2% uranyl acetate in 0.1 M sodium acetate, pH 5.0 for 24 h. After washing, dehydration was performed in ethanol with increasing concentrations, followed by propylene oxide. Resin and propylene infiltration was conducted before embedding in silicone molds in the oven at 62°C for 72 h. Sectioning and imaging were performed following standard EM protocols[55,56]. Vesicle density at the active zones (AZs) was determined as the number of vesicles per square micrometer. All EM samples were numbered, processed, and analyzed under blinded conditions until the last step when data had to be pooled.

Plasmid preparation. Plasmids encoding mCherry-tagged versions of the dipeptides were constructed on the pcDNA3-mCherry vector by insertion of either 25 GGGGCC repeats and adjusting the reading frame to produce specific DPR or DNA encoding the same DPR but using alternative codons so that there were no nucleotide repeats. All constructs were designed to produce 25 repeats of a specific dipeptide repeat following the mCherry. The constructs allow us to compare toxicity with hexanucleotide repeats to that with the same DPR but a different nucleotide sequence. This evaluates the relative contribution of RNA toxicity and dipeptide toxicity to the observed phenotypes.

Primary culture of motor neurons. E15 Spinal motor neurons were isolated from Sprague-Dawley rat embryos (E15) according to published protocols[57]. Briefly, spinal cords were dissected, trypsinized, triturated and then separated on a 9% Optiprep centrifugation. Motor neurons were plated on laminin-coated slides. Primary culture spinal cord motor neurons transfected with mCherry-tagged versions of C9 repeats in Ibidi chambers. At 12 DIV, neurons were transfected using the TransIT-X2 reagent (Mirus #MIR6000). For transfection, DNA (0.5ug) was mixed with 1.5uL X2 in 10uL unsupplemented neurobasal medium and incubated for 15 min at room temperature, and then mixed with media in each culture compartment. Motor neurons were fixed 24hr post-transfection with 4% paraformaldehyde, washed 3x 5 min in PBS, autofluorescence quenched with 50 mM NH4Cl for 15 min, washed 3x 5 min in PBS, permeabilized with 0.1% Triton-X100 for 10 min, washed 3x 5 min in PBS, nonspecific binding blocked with 5% milk for 1hr, washed 2x with PBS, incubated overnight at 4 C with primary antibody (DM1a, 1:1000, Sigma T6199), washed 3x 5 min in PBS, incubated for 45 min at room temperature with secondary antibody (AF488, 1:800, Invitrogen # A11001), washed 3 x 5 min in

PBS, and covered in an aqueous anti-fade mounting medium (Vectashield #H-1000) for storage at 4C until imaged (Zeiss LSM710, 25x).

Neuronal imaging and quantification. Neuronal morphology was examined in fixed neurons transfected with plasmids encoding mCherry versions of C9 dipeptide repeats with or without drug treatment. Degeneration scores were conducted triple-blinded. Images were obtained by one individual from samples, showing neurons with isolated neurites. Twenty or more images were collected for each condition. The second individual placed boxes on all neurite segments in the image that were isolated. The length of neurites in these boxes were measured by a third person and the length showing fragmented, beaded or aggregated phenotypes compared to the total length of the neurite in the box were calculated. All three individuals were blinded as to the experimental condition when selecting images and neurites or while making measurements. Neuritic degeneration was scored as a degeneration score, which is the ratio (degenerating neurite length)/(total neurite length) times 10. Data is shown as a box plot summarizing data from ≥20 images for each condition. (Figs 6 and S1).

Data Availability:

All data generated or analyzed during this study are included in this article and its supplementary information files. All raw datasets and additional information are available from corresponding authors on reasonable request.

References:

Renton, A. E., Chio, A. & Traynor, B. J. State of play in amyotrophic lateral sclerosis genetics. *Nat Neurosci* **17**, 17-23 (2014). https://doi.org/10.1038/nn.3584
DeJesus-Hernandez, M. *et al.* Expanded GGGGCC hexanucleotide repeat in noncoding region of C9ORF72 causes chromosome 9p-linked FTD and ALS. *Neuron* **72**, 245-256 (2011). https://doi.org/10.1016/j.neuron.2011.09.011
Renton, A. E. *et al.* A hexanucleotide repeat expansion in C9ORF72 is the cause of chromosome 9p21-linked ALS-FTD. *Neuron* **72**, 257-268 (2011). https://doi.org/10.1016/j.neuron.2011.09.010
Ghasemi, M. & Brown, R. H., Jr. Genetics of Amyotrophic Lateral Sclerosis. *Cold Spring Harbor perspectives in medicine* **8** (2018). https://doi.org/10.1101/cshperspect.a024125
Taylor, J. P., Brown, R. H., Jr. & Cleveland, D. W. Decoding ALS: from genes to mechanism. *Nature* **539**, 197-206 (2016). https://doi.org/10.1038/nature20413

Kim, G., Gautier, O., Tassoni-Tsuchida, E., Ma, X. R. & Gitler, A. D. ALS Genetics: Gains, Losses, and Implications for Future Therapies. *Neuron* **108**, 822-842 (2020). https://doi.org:10.1016/j.neuron.2020.08.022

Balendra, R. & Isaacs, A. M. C9orf72-mediated ALS and FTD: multiple pathways to disease. *Nat Rev Neurol* **14**, 544-558 (2018). https://doi.org:10.1038/s41582-018-0047-2

Smeyers, J., Banchi, E. G. & Latouche, M. C9ORF72: What It Is, What It Does, and Why It Matters. *Frontiers in cellular neuroscience* **15**, 661447 (2021). https://doi.org:10.3389/fncel.2021.661447

Braems, E., Swinnen, B. & Van Den Bosch, L. C9orf72 loss-of-function: a trivial, stand-alone or additive mechanism in C9 ALS/FTD? *Acta Neuropathol* **140**, 625-643 (2020). https://doi.org:10.1007/s00401-020-02214-x

O'Rourke, J. G. *et al.* C9orf72 is required for proper macrophage and microglial function in mice. *Science* **351**, 1324-1329 (2016). https://doi.org:10.1126/science.aaf1064

Sellier, C. *et al.* Loss of C9ORF72 impairs autophagy and synergizes with polyQ Ataxin-2 to induce motor neuron dysfunction and cell death. *EMBO J* **35**, 1276-1297 (2016). https://doi.org:10.15252/embj.201593350

Nassif, M., Woehlbier, U. & Manque, P. A. The Enigmatic Role of C9ORF72 in Autophagy. *Frontiers in neuroscience* **11**, 442 (2017). https://doi.org:10.3389/fnins.2017.00442

Beckers, J., Tharkeshwar, A. K. & Van Damme, P. C9orf72 ALS-FTD: recent evidence for dysregulation of the autophagy-lysosome pathway at multiple levels. *Autophagy* **17**, 3306-3322 (2021). https://doi.org:10.1080/15548627.2021.1872189

Burberry, A. *et al.* Loss-of-function mutations in the C9ORF72 mouse ortholog cause fatal autoimmune disease. *Science translational medicine* **8**, 347ra393 (2016). https://doi.org:10.1126/scitranslmed.aaf6038

McEachin, Z. T., Parameswaran, J., Raj, N., Bassell, G. J. & Jiang, J. RNA-mediated toxicity in C9orf72 ALS and FTD. *Neurobiol Dis* **145**, 105055 (2020). https://doi.org:10.1016/j.nbd.2020.105055

Zhang, Y. J. *et al.* Heterochromatin anomalies and double-stranded RNA accumulation underlie C9orf72 poly(PR) toxicity. *Science* **363** (2019). https://doi.org:10.1126/science.aav2606

Van't Spijker, H. M. & Almeida, S. How villains are made: The translation of dipeptide repeat proteins in C9ORF72-ALS/FTD. *Gene* **858**, 147167 (2023). https://doi.org:10.1016/j.gene.2023.147167

Cleary, J. D. & Ranum, L. P. Repeat-associated non-ATG (RAN) translation in neurological disease. *Hum Mol Genet* **22**, R45-51 (2013). https://doi.org:10.1093/hmg/ddt371

Cleary, J. D. & Ranum, L. P. New developments in RAN translation: insights from multiple diseases. *Curr Opin Genet Dev* **44**, 125-134 (2017). https://doi.org:10.1016/j.gde.2017.03.006

Freibaum, B. D. & Taylor, J. P. The Role of Dipeptide Repeats in C9ORF72-Related ALS-FTD. *Front Mol Neurosci* **10**, 35 (2017). https://doi.org:10.3389/fnmol.2017.00035

Nishimura, A. L. & Arias, N. Synaptopathy Mechanisms in ALS Caused by C9orf72 Repeat Expansion. *Frontiers in cellular neuroscience* **15**, 660693 (2021). https://doi.org:10.3389/fncel.2021.660693

Kim, H. J. & Taylor, J. P. Lost in Transportation: Nucleocytoplasmic Transport Defects in ALS and Other Neurodegenerative Diseases. *Neuron* **96**, 285-297 (2017). https://doi.org:10.1016/j.neuron.2017.07.029

Zhang, K., Grima, J. C., Rothstein, J. D. & Lloyd, T. E. Nucleocytoplasmic transport in C9orf72-mediated ALS/FTD. *Nucleus* **7**, 132-137 (2016). https://doi.org:10.1080/19491034.2016.1172152

Song, Y., Kang, M., Morfini, G. & Brady, S. T. Fast axonal transport in isolated axoplasm from the squid giant axon. *Methods Cell Biol* **131**, 331-348 (2016). https://doi.org:10.1016/bs.mcb.2015.07.004

Moreno, H. *et al.* Tau Pathology Mediated Presynaptic Dysfunction. *Neuroscience* **325**, 30-38 (2016). https://doi.org:10.1016/j.neuroscience.2016.03.044

Serulle, Y. *et al.* 1-Methyl-4-phenylpyridinium induces synaptic dysfunction through a pathway involving caspase and PKC{delta} enzymatic activities. *Proc Natl Acad Sci U S A* **104**, 2437-2441 (2007).

Llinas, R. *The Squid Giant Synapse: A Model for Chemical Transmission*. (1999).

Morfini, G. A. *et al.* Inhibition of Fast Axonal Transport by Pathogenic SOD1 Involves Activation of p38 MAP Kinase. *PLoS One* **8**, e65235 (2013). https://doi.org:10.1371/journal.pone.0065235

Song, Y. *et al.* Molecular chaperone Hsp110 rescues a vesicle transport defect produced by an ALS-associated mutant SOD1 protein in squid axoplasm. *Proc Natl Acad Sci U S A* **110**, 5428-5433 (2013). https://doi.org:10.1073/pnas.1303279110

Song, Y. Synaptic Actions of Amyotrophic Lateral Sclerosis-Associated G85R-SOD1 in the Squid Giant Synapse. *eNeuro* **7** (2020). https://doi.org:10.1523/ENEURO.0369-19.2020

Fischer, L. R. *et al.* Amyotrophic lateral sclerosis is a distal axonopathy: evidence in mice and man. *Exp Neurol* **185**, 232-240 (2004).

Ackerley, S. *et al.* p38alpha stress-activated protein kinase phosphorylates neurofilaments and is associated with neurofilament pathology in amyotrophic lateral sclerosis. *Mol Cell Neurosci* **26**, 354-364 (2004).

Bendotti, C. *et al.* Activated p38MAPK is a novel component of the intracellular inclusions found in human amyotrophic lateral sclerosis and mutant SOD1 transgenic mice. *J Neuropathol Exp Neurol* **63**, 113-119 (2004).

Bendotti, C. *et al.* Inter- and intracellular signaling in amyotrophic lateral sclerosis: role of p38 mitogen-activated protein kinase. *Neurodegener Dis* **2**, 128-134 (2005).

Tortarolo, M. *et al.* Persistent activation of p38 mitogen-activated protein kinase in a mouse model of familial amyotrophic lateral sclerosis correlates with disease progression. *Mol Cell Neurosci* **23**, 180-192 (2003).

Veglianese, P. *et al.* Activation of the p38MAPK cascade is associated with upregulation of TNF alpha receptors in the spinal motor neurons of mouse models of familial ALS. *Mol Cell Neurosci* **31**, 218-231 (2006).

Bosco, D. A. *et al.* Wild-type and mutant SOD1 share an aberrant conformation and a common pathogenic pathway in ALS. *Nat Neurosci* **13**, 1396-1403 (2010). https://doi.org:nn.2660 [pii]

10.1038/nn.2660

Gibbs, K. L. *et al.* Inhibiting p38 MAPK alpha rescues axonal retrograde transport defects in a mouse model of ALS. *Cell Death Dis* **9**, 596 (2018). https://doi.org:10.1038/s41419-018-0624-8

Sama, R. R. *et al.* ALS-linked FUS exerts a gain of toxic function involving aberrant p38 MAPK activation. *Sci Rep* **7**, 115 (2017). https://doi.org:10.1038/s41598-017-00091-1

Ash, P. E. *et al.* Unconventional translation of C9ORF72 GGGGCC expansion generates insoluble polypeptides specific to c9FTD/ALS. *Neuron* **77**, 639-646 (2013). https://doi.org:10.1016/j.neuron.2013.02.004

Le Ber, I. *et al.* C9ORF72 repeat expansions in the frontotemporal dementias spectrum of diseases: a flow-chart for genetic testing. *J Alzheimers Dis* **34**, 485-499 (2013). https://doi.org:10.3233/JAD-121456

Roy, S. M. *et al.* Targeting human central nervous system protein kinases: an isoform selective p38alphaMAPK inhibitor that attenuates disease progression in Alzheimer's Disease mouse models. *ACS chemical neuroscience* (2015). https://doi.org:10.1021/acschemneuro.5b00002

Takekawa, M., Tatebayashi, K. & Saito, H. Conserved docking site is essential for activation of mammalian MAP kinase kinases by specific MAP kinase kinase kinases. *Mol Cell* **18**, 295-306 (2005).

Volynets, G. P. *et al.* Identification of 3H-naphtho[1,2,3-de]quinoline-2,7-diones as inhibitors of apoptosis signal-regulating kinase 1 (ASK1). *J Med Chem* **54**, 2680-2686 (2011). https://doi.org:10.1021/jm200117h

Kang, M., Baker, L., Song, Y., Brady, S. T. & Morfini, G. Biochemical analysis of axon-specific phosphorylation events using isolated squid axoplasms. *Methods Cell Biol* **131**, 199-216 (2016). https://doi.org:10.1016/bs.mcb.2015.06.003

van Blitterswijk, M. *et al.* Ataxin-2 as potential disease modifier in C9ORF72 expansion carriers. *Neurobiol Aging* **35**, 2421 e2413-2427 (2014). https://doi.org:10.1016/j.neurobiolaging.2014.04.016

van Blitterswijk, M. *et al.* Genetic modifiers in carriers of repeat expansions in the C9ORF72 gene. *Mol Neurodegener* **9**, 38 (2014). https://doi.org:10.1186/1750-1326-9-38

Ajroud-Driss, S. & Siddique, T. Sporadic and hereditary amyotrophic lateral sclerosis (ALS). *Biochim Biophys Acta* **1852**, 679-684 (2015). https://doi.org:10.1016/j.bbadis.2014.08.010

Zhang, Y., Ni, W., Horwich, A. L. & Kaczmarek, L. K. An ALS-Associated Mutant SOD1 Rapidly Suppresses KCNT1 (Slack) Na(+)-Activated K(+) Channels in Aplysia Neurons. *J Neurosci* **37**, 2258-2265 (2017). https://doi.org:10.1523/JNEUROSCI.3102-16.2017

Lin, Y. *et al.* Toxic PR Poly-Dipeptides Encoded by the C9orf72 Repeat Expansion Target LC Domain Polymers. *Cell* **167**, 789-802 e712 (2016). https://doi.org:10.1016/j.cell.2016.10.003

Brady, S. T., Lasek, R. J. & Allen, R. D. Fast axonal transport in extruded axoplasm from squid giant axon. *Science* **218**, 1129-1131 (1982).

Brady, S. T., Lasek, R. J. & Allen, R. D. Video microscopy of fast axonal transport in isolated axoplasm: A new model for study of molecular mechanisms. *Cell Motility* **5**, 81-101 (1985).

Young, J. Z. & Keynes, R. The Functioning of the Giant Nerve Fibres of the Squid. 1938 - J.Z. and the discovery of squid giant nerve fibres. *J Exp Biol* **208**, 179-180 (2005). https://doi.org:10.1242/jeb.01405

Smith, S. J., Buchanan, J., Osses, L. R., Charlton, M. P. & Augustine, G. J. The spatial distribution of calcium signals in squid presynaptic terminals. *J Physiol* **472**, 573-593 (1993). https://doi.org:10.1113/jphysiol.1993.sp019963

Augustine, G. J., Morgan, J. R., Villalba-Galea, C. A., Jin, S., Prasad, K. & Lafer, E. M. Clathrin and synaptic vesicle endocytosis: studies at the squid giant synapse. *Biochem Soc Trans* **34**, 68-72 (2006). https://doi.org:10.1042/BST0340068

Moreno, H. H. *et al.* Synaptic transmission block by presynaptic injection of oligomeric amyloid beta. *Proc Natl Acad Sci U S A* **106**, 5901-5906 (2009).

Camu, W. & Henderson, C. E. Rapid purification of embryonic rat motoneurons: an in vitro model for studying MND/ALS pathogenesis. *J Neurol Sci* **124 Suppl**, 73-74 (1994).

Acknowledgments:

We thank Drs. Bradley Hyman and Merit Cudkowicz for their full support of the Song Laboratory at MGH. We thank Dr. Marty Watterson (NWU) for the generous supply of p38 inhibitors. The research was supported by NIH grants [P30AG062421 (Developmental Project) and AG072516], a Jack Satter Foundation grant, a Pape Adams ALS Transformative Scholar Award, and an AARG grant from the Alzheimer's Association to YS; NIH grants (R21NS120126 and R01NS118177) and a gift from Neurodegenerative Research Inc. to GM; and NIH grants (NS023868 and NS041170) to STB.

Author Contributions:

YS, STB, and GM were responsible for the original conceptualization of the work and experimental design of all studies in this manuscript. YS and STB wrote the manuscript. YS, GM, and STB performed all experiments in the squid axoplasm. YS, PC, and JEM performed all experiments in the squid giant synapse. MYT and BW performed all experiments in cell cultures. NK contributed to experiments in Figure 3C-F. YS, MYT, BW, GM, and STB analyzed all the data. All authors reviewed and edited the final manuscript.

Competing interests:

None

Materials & Correspondence:

Correspondence and material requests should be addressed to Yuyu Song, Gerardo Morfini, and Scott Brady.

Figure Legends:

Figure 1. GP(n$_8$) DRPs selectively inhibits anterograde axonal transport. Plots depict results from vesicle motility assays. Individual velocity (μm/sec) measurements (arrowheads) are plotted as a function of time (minutes) for a whole duration of 50min. Blue arrowheads and lines represent anterograde, kinesin-dependent fast axonal transport rates. Red reverse arrows and lines represent retrograde, cytoplasmic dynein-dependent axonal transport rates. Perfusion of GP(n$_8$) peptide specifically inhibited anterograde, but not retrograde transport (**A**). In contrast, perfusions of GA(n$_8$) (**B**), PR(n$_8$) **(C)**, GR(n$_8$) (**D**), or AP(n$_8$) **(E)** peptides did not affect axonal transport in either direction. The toxic effect of GP(n$_8$) was similar to those elicited by mutant SOD1-G93A (**F**), SOD1-H46R, and SOD1-G85R[28,29]

Figure 2. The inhibition of anterograde transport induced by GP(n$_8$) peptide involves activation of kinases in the ASK1-p38 MAPK pathway. (**A**) MW01-2-069SRM (MW069), a selective inhibitor of p38α (Munoz et al., 2007), blocked the inhibition of anterograde axonal transport induced by GP(n$_8$) peptide repeat. (**B**) Co-perfusion of GP(n$_8$) with 20μM DVD, a peptide inhibitor of a subset of MAPKKK (Morfini et al., 2013), also prevented inhibition of anterograde transport. Interestingly, NQDI, an ASK1 inhibitor, had a similar protective effect **(C)**, suggesting that GP(n$_8$)-induced toxicity on axonal transport is mediated by the ASK1-P38 pathway. Remarkably, our previous studies showed that MW069 (**D**), DVD peptide, and NQDI also prevented the inhibitory effect of mutant SOD1[28,29].

Figure 3. GP(n$_8$) peptide increases kinesin phosphorylation *via* axon-autonomous activation of p38 kinases. Sister axons were extruded and perfused with control buffer (X/2, Ctrl) or with X/2 buffe plus 1μM GP(n$_8$) dipeptide repeat (GP) in the presence of ^{32}P-γ-ATP. After 50 min incubation, an aliquot of lysates was separated by SDS-PAGE and analyzed by autoradiography (**A**). High-Molecular-Weight Neurofilament (HMW) and Neurofilament Heavy Chain (NFH), major axoplasmic phosphoproteins in squid, are indicated. Kinesin was immunoprecipitated from the reminder lysate using H2, (an antibody against Kinesin Heavy Chain). Immunoprecipitated were separated by SDS-PAGE and ^{32}P incorporation quantified by Phosphorimager scanning (KHC IPP) (**B**). Immunoblotting analysis of axoplasms showed

increased activation/phosphorylation of p38 (p-p38) in axoplasms perfused with GP (n_8) dipeptide, compared to "sister' axons perfused with control buffer (Ctrl) (**C**). Anti-kinesin-1 (KHC) antibody served as a loading control for axoplasmic protein. Quantitation of blots revealed no significant differences in levels of active GSK3 (**D**) or ERK kinases (**E**) but a significant increase in p-p38 in axoplasms perfused with GP(n_8) peptide, compared to sister axons perfused with control buffer (**F**). (**G-H**): Perfusion of GA(n_8) peptide did not activate P38, as indicated by comparable phospho-p38 levels in western blots.

Figure 4. GP(n_8) peptide inhibits synaptic transmission in the giant synapse. (**A**) Giant synapses were dissected and continuously superfused with oxygenated squid saline. GP(n_8) or GR (n_8) were injected into the presynaptic terminal (II) at 50 psi for 250ms/injection at 0.1Hz, and membrane potentials were recorded from both pre- (II) and post-synaptic (III) terminals. GR(n_8) had no inhibition of synaptic transmission as the synapse kept firing for 250min (**B**). In contrast, presynaptic injection of GP(n_8) gradually reduced postsynaptic potential and eventually led to the failure in firing within 200min (**C**). Quantitative plots (**D**) showed synaptic actions of GP(n_8) and GR(n_8) and suggested that NQDI, the same ASK1 inhibitor that prevented axonal transport defects in (**Fig. 2C**), also blocked the inhibitory effect of GP(n_8) on synaptic transmission (n = 5). High-frequency stimulation at 50Hz for 5sec didn't affect EPSP slope in GR-injected synapse (**E**) but significantly reduced PSP slope in GP(n_8) injected synapse (**F**). EPSP slopes during the 5sec train stimulation were integrated and plotted with the linear fit (**G**), suggesting reduced RPR size and mobilization rate of synaptic vesicles in the synapses injected with GP(n_8), as compared to the ones injected with GR(n_8) (n = 5) (**H**). Electron Microscopy showed sufficient synaptic vesicles at the active zone of the GR(n_8)-synapses (**I**) but a drastic decrease of available vesicles in GP(n_8)-synapses (**J**). Quantification of various classes of vesicles showed differences in total synaptic vesicles (SVs), large electron lucid vesicles, and docking vesicles, with similar numbers of clathrin-coated vesicles (CCV). Data were plotted from 100 synapses, injected with either GP(n_8) or GR(n_8) across 3 separate biological repeats (**K**).

Figure 5. Toxicity of C9ORF72 Dipeptide Repeats in rat primary cultured motor neurons. Plasmids containing mCherry or mCherry fusion proteins with different dipeptide repeats (n=25 repeats) were used to transfect rat primary cultured spinal cord motor neurons. Neurons were transfected at 6DIV and fixed at 24h posttransfection. (**A**) Transfection with a plasmid expressing mCherry alone had no effect on neuronal viability. Similarly, transfection with mCherry-GA$_{25}$ or -AP$_{25}$ had little or no effect on neuronal morphology. (**B**) In contrast, motor neurons transfected with mCherry-RP$_{25}$, -RG$_{25}$ or -GP$_{25}$ exhibited significant degeneration. This was in contrast to the effects of dipeptide repeats on axonal transport in axonal compartments or presynaptic terminals, where only GP dipeptides had significant effects. Quantitation of neuronal degeneration with each of the dipeptide repeats indicated that the toxicity was comparable whether the dipeptides were the result of adding GGGGCC hexanucleotide repeats to mCherry (**C**) or the dipeptides resulted from using alternate codons for the same dipeptides (**D**). This confirmed that the toxicity of all

three toxic mCherry dipeptide repeats (-RP, -RG, and -GP) were the result of the dipeptide effects rather than RNA toxicity.

Figure 6. **The GP dipeptides produce axonopathy, while RP and RG dipeptides exhibit nuclear changes**. The mCherry-GP affects fast axonal transport, leading to a dying-back axonopathy. The nuclear envelope is intact (Lamin A/C) and chromatin is diffuse (DAPI). Axons are fragmented and truncated. In contrast, axons of neurons transfected with either mCherry-RG or mCherry-RP remain intact, but the chromatin (DAPI) is clumped and the nuclear envelope is breaking down (Lamin A/C). These different phenotypes indicate that there are at least two pathogenic mechanisms involved. Toxicity due to C9-GP dipeptides affect axonal transport and synaptic transmission through a pathway involving ASK1/P38 MAP kinase. In contrast, C9-RG and C9-RP dipeptides damage nuclear structure without affecting fast axonal transport or synaptic function, and their effects are not rescued by inhibitors of P38 or ASK1 (see supplemental figures 1-3).

Statistical Information:

All experiments were repeated at least three times. The data were analyzed by one-way ANOVA followed by the Tukey *post hoc* test (or nonparametric multiple *t* tests, without assuming consistent SD) and plotted in Prism 7 (GraphPad software). Quantitative data were plotted as mean ± SEM, *p* values were calculated. For EM analysis, nested one-way ANOVA was performed.

Song et al. Fig 3

A

I
Pre
III
Post
II

B Presynaptic injection of GR

- - - pre
—— Post (before injection)
—— Post (50min after injection)
—— Post (100min after injection)
—— Post (150min after injection)
—— post (200min after injection)
—— Post (250min after injection)

20mV
2ms

C Presynaptic injection of GP

- - - Pre
—— Post (before injection)
—— Post (50min after injection)
—— Post (100min after injection)
—— Post (150min after injection)
—— Post (200min after injection)

20mV
2ms

D

Normalized EPSP Slope

GP
GR
GP + NQDI

Time (min)

E Presynaptic injection of GR

F Presynaptic injection of GP

20mV
1s

G

Integrated EPSP Slope

GR
GP

RRP (GA)
RRP (GP)

Time (Sec)

H

Normalized Value

GR
GP

***P < 0.00001   ***P < 0.00001

RRP size   Mobilization rate

I

500 nm

GR

J

500 nm

GP

K

Numbers

***P < 0.00001   ***P < 0.00001

GR   GP
GR GP   GR GP   GR GP   GR GP

P > 0.01   *P = 0.01   **P < 0.001

SVs   CCVs   Large SVs   Docking SVs   Total SVs

SVs
CCVs
Large SVs
Docking SVs
Total SVs

n = 102

Song et al. Fig 4

mCherry Control   mCherry-GA   mCherry-AP

mCherry-RP   mCherry-RG   mCherry-GP

Song et al. Fig 5