Title:

A Type VI Secretion System in *Burkholderia* Species *cenocepacia* and *orbicola* Triggers

Distinct Macrophage Death Pathways Independent of the Pyrin Inflammasome

Authors:

Nicole A. Loeven[1,2], Arianna D. Reuven[1], Abigail P. McGee[1], Clarrisa Dabi[1], Bethany W.

Mwaura[1] and James B. Bliska[1,3]

[1]Department of Microbiology and Immunology, Geisel School of Medicine at Dartmouth College,

Hanover, NH, USA

[2]Current address: Department of Pathology and Laboratory Medicine, Hospital of the University

of Pennsylvania, Philadelphia, Pennsylvania, USA

[3]Correspondiing author: james.bliska@dartmouth.edu

Abstract

The *Burkholderia cepacia* complex contains opportunistic pathogens that cause chronic infections and inflammation in lungs of people with cystic fibrosis. Two closely related species within this complex are *Burkholderia cenocepacia* and the recently classified *Burkholderia orbicola*. *B. cenocepacia* and *B. orbicola* encode a type VI secretion system and the effector TecA, which is detected by the pyrin/caspase-1 inflammasome, and triggers macrophage inflammatory death. In our earlier study the pyrin inflammasome was dispensable for lung inflammation in mice infected with *B. orbicola* AU1054*,* indicating this species activates an alternative pathway of macrophage inflammatory death. Notably, *B. cenocepacia* J2315 and K56-2 can damage macrophage phagosomes and K56-2 triggers activation of the caspase-11 inflammasome, which detects cytosolic LPS. Here we investigated inflammatory cell death in pyrin-deficient (*Mefv*$^{-/-}$) mouse macrophages infected with *B. cenocepacia* J2315 or K56-2 or *B. orbicola* AU1054 or PC184. Macrophage inflammatory death was measured by cleavage of gasdermin D protein, release of cytokines IL-1α and IL-1β and plasma membrane rupture. Findings suggest that J2315 and K56-2 are detected by the caspase-11 inflammasome in *Mefv*$^{-/-}$ macrophages, resulting in IL-1β release. In contrast, inflammasome activation is not detected in *Mefv*$^{-/-}$ macrophages infected with AU1054 or PC184. Instead, AU1054 triggers an alternative macrophage inflammatory death pathway that requires TecA and results in plasma membrane rupture and IL-1α release. Amino acid variation between TecA isoforms in *B. cenocepacia* and *B. orbicola* may explain how the latter species triggers a non-inflammasome macrophage death pathway.

Introduction

The *Burkholderia cepacia* complex (Bcc) comprise a group of Gram-negative bacterial species that exist in the environment (1). The Bcc are opportunistic pathogens that cause infections in immunocompromised individuals including people with cystic fibrosis (pwCF) (1). The Bcc typically cause chronic infections associated with inflammation and exacerbations in adult pwCF. Chronic infections are difficult to eliminate due to intrinsic antibiotic resistance in the Bcc. Chronic Bcc infections can progress to an acute disease called cepacia syndrome that is characterized by necrotizing pneumonia, bacteremia, and sepsis. *B. cenocepacia* accounts for ~45% of Bcc isolates from pwCF in the United States and is considered one of the most virulent in this group because of its strong association with cepacia syndrome (2, 3). In addition, *B. cenocepacia* infection is a contraindication for lung transplantation (4). *B. cenocepacia* subgrouping based on *recA* lineages has resulted in classification of genomovars IIIA-D (5) and shown that isolates belonging to genomovar IIIA commonly circulate in the UK, Europe, and Canada, while genomovar IIIB isolates more commonly cause infections in pwCF in the United States (6).

B. cenocepacia encodes numerous virulence factors (1, 7) and preferentially resides within phagocytes in the CF lung (8-10). Residence of *B. cenocepacia* in phagocytes has been observed by in situ imaging of human lung sections from transplant or autopsy, or mouse lung sections from experimental infections (11, 12). *B. cenocepacia* resides in vacuoles in phagocytic cells (11, 12), but may damage phagosomes and escape into the cytosol as well (13, 14). Additionally, *B. cenocepacia* can invade and survive in airway epithelial cells which may play a role in transmigration across the epithelial barrier (7). The intracellular location of *B. cenocepacia* appears to be key to its pathogenesis (8). *B. cenocepacia* can prevent phagosomal maturation and delay co-localization of the NADPH oxidase complex with the vacuolar membrane (15, 16). CF macrophages display defects in oxidative burst (17) and

bacteria-specific autophagy (xenophagy) (9, 10) which may render them permissive to *B.*

cenocepacia. In non-CF macrophages, phagosome damage by *B. cenocepacia* increases

xenophagy (18) but the pathogen may be able to subvert this bactericidal process for

intracellular survival (13). In addition, type I interferon production by non-CF macrophages in

response to cytosolic *B. cenocepacia* infection can act in an autocrine fashion to increase

xenophagy and limit bacterial replication (14).

A *B. cenocepacia* virulence factor implicated in lung infection and intracellular pathogenesis a

type VI secretion system (T6SS) (8, 19). The T6SS contains a tube-like structure that functions

to deliver effectors into bacteria or host cells (20, 21). Upon contact with target cells the tube

comprised of the Hcp protein fires to puncture membranes and release associated effectors.

The effectors have enzymatic activity that modify and inactivate key cellular processes or

structures in target cells. The *B. cenocepacia* T6SS belongs to the first of three classes of these

systems to be characterized (T6SS[i], T6SS[ii] and T6SS[iii]) (21). The T6SS[i] class is common

within Gammaproteobacteria and since the *B. cenocepacia* system is homologous to T6SS-1 of

Burkholderia pseudomallei (22) we hereafter referred to it as T6SS-1. T6SS-1 is required for

intracellular *B. cenocepacia* to inactivate small GTPases including Rac1 and Cdc42, resulting in

disruption of the actin cytoskeleton and preventing assembly of the NADPH oxidase complex in

macrophages (23, 24).

T6SS-1 is detected by inflammasomes in macrophages harboring *B. cenocepacia* (18, 25-27).

Canonical caspase-1 inflammasomes are assembled when sensors such as NLRP3, NLRC4 or

pyrin detect danger signals generated by bacterial pathogens during infection of host cells (28).

Active caspase-1 is produced and cleaves the pro-inflammatory cytokine pro-IL-1β and the

gasdermin-D (GSDMD) protein (29). Cleavage of GSDMD releases a domain that oligomerizes

and forms pores in the plasma membrane (29). Mature IL-1β leaks out of these pores and can

stimulate inflammatory signaling through activation of the IL-1 receptor found on a variety of

host cells (29). Pore formation in plasma membranes can also result in pyroptosis, a pro-

inflammatory form of cell death (29).

Early studies implicated T6SS-1 of *B. cenocepacia* strain K56-2 (genomovar IIIA, Table 1) in

activation of the NLRP3 and NLRC4 inflammasomes in murine macrophages and the pyrin

inflammasome in human macrophages (25, 26). Assembly of the pyrin inflammasome in

response to T6SS-1 was subsequently observed in murine macrophages infected with J2315

(genomovar IIIA, Table 1) (27). Pyrin inflammasome assembly in macrophages infected with

J2315 required inactivation of RhoA by T6SS-1 (27). The T6SS-1 in K56-2 has also been

shown to be detected by the non-canonical caspase-11 inflammasome in murine macrophages

(18, 30), which is consistent with phagosome damage (26) and entry of LPS into the cytosol

(31). Caspase-11 can cleave GSDMD to form pores in macrophages but requires activation of a

caspase-1 inflammasome (e.g. NLRP3) for processing and release of IL-1β (31).

Aubert et al. identified TecA (T6SS effector protein affecting cytoskeletal architecture) as being

responsible for inactivation of RhoA and assembly of the pyrin inflammasome in macrophages

infected with K56-2 or J2315 (32). In silico modeling of *B. cenocepacia* TecA predicted a fold

consistent with a cysteine protease-like hydrolase, and putative catalytic C41, H105 and D149

residues were confirmed by mutagenesis (32). TecA appears to be the founding member of a

T6SS effector superfamily with deamidase activity in bacterial opportunistic pathogens and

symbionts (32). After entry into phagocytes *B. cenocepacia* employs the T6SS-1 to deliver TecA

across the phagosome into the cytosol (32). RhoA is inactivated in response to deamidation of

Asn-41 by TecA (32). Assembly of the pyrin inflammasome in response to deamidation of RhoA

by TecA in macrophages infected with *B. cenocepacia* is an example of a guard mechanism of

effector-triggered immunity (32). More recently TecA has been shown to promote localized

actin polymerization that delays maturation of phagosomes containing J2315 in macrophages

(33).

Previous studies examined the role of pyrin and TecA in lung pathogenesis by intranasally

infecting C57BL/6 mice with J2315 (27, 32). Xu et al. compared C57BL/6 and pyrin knockout

(*Mefv*$^{-/-}$) mice and found that J2315 infection stimulated inflammatory cell infiltration into lungs of

C57BL/6 but not *Mefv*$^{-/-}$ mice (27). Aubert et al. infected C57BL/6 mice with J2315 or a Δ*tecA*

mutant and reported that TecA was important for inflammatory cell infiltration and acute lung

damage (32).

Increasing numbers of whole genome sequences for *Burkholderia* strains are being determined

and analyzed. Results of two recent genomic analysis studies have suggested that genomovar

IIIB isolates be designated as a species separate from *B. cenocepacia* (34, 35). Morales-Ruíz et

al. propose for the new species the name *B. orbicola* (35). This name comes from the Latin

word *orbis* and suffix -*cola* meaning "the whole world" and "dweller" respectively, making

orbicola "inhabitant of the whole world" (35). Here, we follow this convention and hereafter refer

to genomovar IIIB strains as *B. orbicola*.

B. orbicola strain AU1054 (Table 1), isolated from the bloodstream of a CF patient, encodes

T6SS-1 (36) and TecA (37). Results of murine macrophage infections in vitro showed that

AU1054 TecA triggers release of IL-1β in a T6SS-1- and pyrin-dependent manner (37). In an

infection model using C57BL/6 or CF mice (mice with the most common CFTR mutation in

humans, F508del) with AU1054 we confirmed that TecA is important for inflammatory cell

recruitment to lungs (37). Surprisingly, using *Mefv*$^{-/-}$ mice in the CF or non-CF background we

were unable to reproduce a role for pyrin in inflammatory cell accumulation in lungs during

infection with AU1054 or J2315 (37). This finding led us to hypothesize that AU1054 and J2315

can trigger inflammation by a mechanism independent of the pyrin inflammasome. Notably,

J2315 been shown to damage macrophage phagosomes (13), which would allow detection by

the non-canonical inflammasome, similar to its clonal relative K56-2 (18, 30). Activation of the

caspase-11 inflammasome has been shown to contribute to lung inflammation and

immunopathology in mice infected with K56-2 (18).

Here we investigated pyrin-independent macrophage inflammatory cell death in response to

infection with *B. cenocepacia* J2315 or *B. orbicola* AU1054. Results indicate that J2315 and

AU1054 trigger distinct pyrin-independent macrophage inflammatory death pathways with

different inflammasome and TecA requirements. Amino acid differences between TecA protein

sequences of *B. cenocepacia* and *B. orbicola* strains were identified, suggesting distinct effector

isoform activities. These data have important implications for understanding the role of T6SS-1

and TecA in macrophage inflammatory death and the pathogenesis of chronic lung infections

caused by *B. cenocepacia* and *B. orbicola* in pwCF.

159
160   **Results**
161
Pyrin is differentially required for gasdermin-D cleavage in macrophages infected with *B.

cenocepacia* or *B. orbicola*. To determine if AU1054 and J2315 can trigger inflammasome

activation independent of pyrin, bone marrow derived macrophages (BMDMs) from *Mefv*$^{+/+}$ or

Mefv$^{-/-}$ C57BL/6 mice were LPS primed, infected for 90 min and cleavage of GSDMD was

measured by immunoblotting. For control strains we used AU1054 Δ*tecA* and Δ*hcp* mutants which

are unable to trigger the pyrin inflammasome (37) and K56-2 which activates the caspase-11

inflammasome (18, 30) (Table 1). As expected GSDMD cleavage in BMDMs infected with

AU1054 required TecA, Hcp and pyrin (Fig.1, lanes 2-4, 8-10). Interestingly, GSDMD was

cleaved in *Mefv*$^{+/+}$ and *Mefv*$^{-/-}$ BMDMs infected with J2315 (isolate A, Table 1), like K56-2 (Fig.1,

lanes 5,6 and 11,12). These results suggest that J2315 can, like K56-2, activate a pyrin-

independent inflammasome in macrophages.

To determine if the phenotype we observed was specific to the J2315 A isolate from our

laboratory, several additional isolates obtained from other investigators (Isolates B-D, Table 1)

were analyzed in infections with $Mefv^{+/+}$ and $Mefv^{-/-}$ BMDMs. GSDMD was cleaved in $Mefv^{-/-}$

BMDMs infected with all isolates (Fig.2A, lanes 8,9 and Fig.2B, lanes 7-9), indicating that pyrin-

independent inflammasome activation and GSDMD cleavage is a general property of J2315.

We also infected $Mefv^{+/+}$ and $Mefv^{-/-}$ BMDMs with an additional *B. orbicola* isolate, PC184 (Table

1), and found that pyrin was required for cleavage of GSDMD (Fig.2B, lanes 5,10), like AU1054.

These results suggest that *B. orbicola* and *B. cenocepacia* differ in the ability to activate

inflammasomes in macrophages independent of pyrin.

TecA is dispensable for pyrin-independent GSDMD cleavage and IL-1β release in

macrophages infected with J2315. To determine if TecA is required for J2315 to trigger pyrin-

independent GSDMD cleavage and IL-1β release, we obtained an additional isolate (Isolate E)

of this strain and its corresponding Δ*tecA* mutant (32) (Table 1). LPS-primed $Mefv^{+/+}$ or $Mefv^{-/-}$

BMDMs infected with these strains or AU1054 controls were analyzed for GSDMD cleavage as

well as IL-1β release. As expected from our previous ELISA results (37) and the GSDMD

cleavage data presented here (Figs.1, 3A), TecA and pyrin were required for significant release

of IL-1β from BMDMs infected with AU1054 (Fig.3B). However, findings here showed that TecA

and pyrin were not required for GSDMD cleavage (Fig.3A, lane 11) or IL-1β release BMDMs

infected with J2315 (Fig.3C). Additional experiments are needed to solidify these findings

because the IL-1β values for the J2315 samples showed variability. Nevertheless, these results

are consistent with the idea that J2315, like K56-2, can activate both pyrin and the caspase-11

inflammasomes in macrophages (18). This conclusion will need to be verified using *Casp11*[-/-]

BMDMs in future experiments.

TecA triggers pyrin inflammasome-independent IL-1α and LDH release from macrophages

infected with AU1054. To determine if AU1054 triggers another form of pyrin-independent

macrophage inflammatory death, we measured release of IL-1α and LDH. IL-1α can be released

from macrophages by inflammasome-dependent and -independent mechanisms. In the latter

case plasma-membrane rupture (PMR) allows IL-1α to be released from host cells to act as a

danger-associated-molecular pattern (DAMP) (38). LDH at 140 kDa is a marker of PMR. These

experiments used *Mefv*[+/+] or *Mefv*[-/-] BMDMs from mice engineered with a CF genotype (*Cftr*[F508del])

and after LPS priming they were left uninfected or infected with AU1054 or AU1054 Δ*tecA* as

above. For a positive control we treated the BMDMs with *Clostridium difficile* TcdB toxin and the

culture supernatants were analyzed by ELISA for IL-1β, which is dependent upon inflammasome

activation for processing and release. As shown in Fig.4A, significant IL-1β release upon AU1054

infection required TecA and pyrin as expected. However, pyrin was not required for TecA to

trigger release of IL-1α (Fig.4B) or LDH (Fig.4C) from *Mefv*[-/-] BMDMs infected with AU1054. IL-

1α released in response to AU1054 infection was significantly lower in *Mefv*[-/-] as compared to

Mefv[+/+] or BMDMs (Fig.4B), likely because pyrin inflammasome activation contributed to secretion

of the cytokine in the former case. IL-1α and LDH released from *Mefv*[-/-] BMDMs intoxicated with

TcdB was at background levels (Fig.4B, C respectively). These data suggest that AU1054 TecA

triggers an inflammasome-independent mechanism of PMR leading to macrophage inflammatory

death.

The TecA proteins of *B. orbicola* and *B. cenocepacia* exhibit amino acid variation. The

ability of AU1054 TecA to trigger inflammasome-independent PMR in BMDMs (Fig.4) suggested

that the *orbicola* isoform has distinct biological function/activity as compared to the canonical

cenocepacia effector. To examine this possibility, we sequenced *tecA* genes from isolates used

in this study. The resulting predicted TecA protein sequences were aligned using ClustalW

(Fig.5). The *B. cenocepacia* J2315 and K56-2 proteins are 100% identical and the *B. orbicola*

AU1054 and PC184 proteins are 99% identical (Fig.5). In contrast, *B. cenocepacia* TecA

shares only ~90% sequence identity with the *B. orbicola* isoform, due to 15-16 amino acid

differences (Fig.5).

All TecA isoforms have residues C41, H105 and D148 that are important for catalytic activity

(32, 37) (Fig.5). In addition, amino acids immediately flanking the catalytic residues are highly

conserved between *B. cenocepacia* and *B. orbicola* (Fig.5). However, *B. orbicola* TecA has

non-conserved substitutions R38G (charged to non-charged), G101T (non-polar to polar), and

T146A (polar to non-polar) within 3, 4 and 2 amino acids of C41, H105 and D148, respectively

(Fig.5).

Predicted structures of the K56-2 and AU1054 TecA isoforms were determined by AlphaFold2

and displayed by PyMol (Fig.6). Results show that the overall folds and positions of catalytic

residues are highly similar (Fig.6). However, the presence of the non-conserved substitutions

R38G (charged to non-charged), G101T (non-polar to polar), and T146A (polar to non-polar) in

the vicinity of the active site (Fig.6) could impact the biological activity of the *B. orbicola* TecA

protein.

Discussion

The data presented above increase our understanding of the role of T6SS-1 and TecA in the

pathogenesis of *B. orbicola* and *B. cenocepacia* in the following three ways. First, our results

confirm that *B. cenocepacia* triggers inflammasome-dependent macrophage inflammatory death

by two mechanisms, one that is TecA and pyrin dependent and the other likely requires

detection of vacuole damage by caspase-11 (18).   Second, *B. orbicola* via T6SS-1 triggers

TecA-dependent macrophage inflammatory death by two mechanisms, one that requires pyrin

and the other is inflammasome independent. Third, the TecA protein of *B. orbicola* appears to

comprise a separate subgroup of this effector superfamily, with distinct biological activity.

Together, these results indicate that the underlying mechanism of inflammation in chronic *B.*

orbicola and *B. cenocepacia* lung infections is multifactorial and species specific, which has

implications for treating this disease in pwCF.

J2315 and K56-2 are genomovar IIIA CF clinical isolates and members of the epidemic ET12

lineage common in Europe (39). Despite being clonal, the literature indicates that during

macrophage infections the T6SS-1 of J2315 is only detected by the pyrin inflammasome via

TecA (27, 32) while K56-2 is detected by the pyrin and caspase-11 inflammasomes (18, 25, 30).

Our results address this discrepancy by showing that when analyzed in parallel J2315 and K56-

2 both trigger pyrin-independent GSDMD cleavage in infected BMDMs. In addition, J2315

causes low but significant IL-1β release from $Mefv^{-/-}$ BMDMs under our infection conditions. It is

probable that caspase-11 is the proximal inflammasome activated in $Mefv^{-/-}$ BMDMs infected

with J2315 and K56-2. It will be important to carry out *B. cenocepacia* infection experiments with

BMDMs deficient in caspase-11 to confirm this hypothesis. Activation of caspase-11 by J2315

may not have been detected previously (27, 32) due to insufficient lengths of LPS priming of

$Mefv^{-/-}$ BMDMs. The two hour LPS priming used by Xu et al. and Aubert et al. (27, 32) may

have been insufficient to activate expression of caspase-11 (31). The overnight LPS priming we

use to activate expression of pyrin may have fortuitously allowed us to detect activation of

caspase-11 in *Mefv*[-/-] BMDMs infected with J2315 and K56-2.

Xu et al. found that lung inflammation in mice infected with J2315 required pyrin (27). However,

in preliminary experiments we observed only a subtle decrease in lung inflammation in mice

infected with J2315 (37). Experimental variables such as genetic or microbiota differences

between *Mefv*[-/-] mouse lines used by Xu et al. (27) and our study (37) could be responsible for

the discrepancy. However, our result can be explained by the data of Krause et al. (18) who

showed that caspase-11 is important for lung inflammation in mice infected with K56-2 (18).

Presumably, inflammation triggered by caspase-11 could partially compensate for the absence

of pyrin in *Mefv*[-/-] mice infected with J2315. It is possible that pyrin and caspase-11 contribute

additively to inflammation in lungs during *B. cenocepacia,* and this could be studied by directly

comparing responses to *B. cenocepacia* infection in mice that are *Mefv*[-/-] or *Casp11*[-/-] or

deficient in both inflammasomes. Interestingly, *Casp11*[-/-] mice had increased survival compared

to *Casp1*[+/+] mice, even though K56-2 growth was restricted better in the latter animals (18). This

result indicates that lethality in this context is due to immunopathology in the lung rather than

bacterial growth in per se.

The mechanism of caspase-11 activation by J2315 in macrophages is likely the same as K56-2,

where T6SS-1 has been shown to mediate phagosome damage (26). In fact, there is evidence

that J2315 escapes from damaged phagosomes (13, 14). The mechanism of phagosome

damage by *B. cenocepacia* T6SS-1 is unknown. Polymerized actin co-localizes with cytosolic

J2315 in human macrophages (13). However, the role of actin polymerization in phagosome

damage by *B. cenocepacia* is unclear. Additional evidence that *B. cenocepacia* escapes

phagosomes comes from the finding that J2315 activates non-inflammasome PRRs like

mitochondrial antiviral-signaling protein (MAVS) and stimulator of interferon genes (STING) in

macrophages (14). MAVS is activated primarily in response to viral RNAs, where STING is

activated in response to detection of bacterial double stranded DNA (dsDNA) (14). MAVS and

STING activation in macrophages leads to a type I interferon (IFN) response which induces

autophagy/xenophagy and was shown to counteract *B. cenocepacia* J2315 replication in

macrophages (14).

Our data suggest that *B. orbicola* TecA can trigger PMR in infected macrophages by an

inflammasome-independent mechanism because the lack of GSDMD cleavage in *Mefv*[-/-]

macrophages indicates that caspase-1 and caspase-11 are not being activated. *B. orbicola*

TecA can trigger release of LDH from unprimed BMDMs (unpublished observation) further

suggesting this pathway of PMR is inflammasome independent. To confirm this concept, it will

be important to carry out *B. orbicola* infection experiments with BMDMs deficient in caspase-1

and -11. Release of HMGB1 could be measured to confirm that multiple DAMPs are being

released by PMR. It will also be important to determine if *B. orbicola* TecA triggers

inflammasome-independent PMR in human macrophages.

Recently the protein NINJ1 has been shown to be required for PMR downstream of multiple

forms of cell death in BMDMs (40). It is possible that NINJ1 is important for inflammasome-

independent PMR and release of DAMPs from *B. orbicola*-infected macrophages. Microscopy

could be used to confirm that inflammasome-independent DAMP release is associated with

plasma membrane damage in *Mefv*[-/-] macrophages infected with *B. orbicola*. Propidium iodide

uptake measured by fluorescence microscopy and cell rounding and blebbing as detected by

phase microscopy would be indicators of plasma membrane damage (41).

The existence of an inflammasome-independent PMR pathway in macrophages likely explains

why recruitment of inflammatory cells to lungs of mice infected with AU1054 is not decreased in

the absence of pyrin (37). We hypothesize *B. orbicola* TecA triggers redundant pathways of

macrophage inflammatory cell death both in vitro and in vivo. The pyrin-dependent pathway

results in IL-1β release and pyroptosis, while the inflammasome-independent pathway results in

PMR and release of multiple DAMPs. Either pathway can exacerbate inflammatory monocyte

recruitment and persistence in lungs infected with *B. orbicola* (37). For example, IL-1β or

DAMPs released from *B. orbicola*-infected alveolar macrophages could activate airway epithelial

cells to produce chemokines for inflammatory monocytes. Inflammatory monocytes use the

receptor CCR2 to migrate out of the bone marrow into sites of infection in response to the

chemokine CCL2 (42). Inflammatory monocytes can differentiate into macrophages and

depending on the context can cause detrimental immune responses in lung infections (42, 43).

Persistence of excessive numbers inflammatory monocyte-derived macrophages could cause

immunopathology in lungs infected with *B. orbicola* (37). Inflammatory monocyte-derived

macrophages recruited by CCL2 could also become infected with *B. orbicola* and release more

IL-1β or DAMPs, thus initiating a damaging cycle of inflammation.

Our finding that *B. orbicola* TecA triggers inflammasome-independent PMR suggests that the

10% amino acid variation in this isoform imparts different biological activity as compared to *B.*

cenocepacia. Although speculative we suggest that *B. orbicola* TecA deaminates a host protein

target which triggers PMR. To test the above hypothesis, it would be interesting to replace

TecA in AU1054 with the *B. cenocepacia* isoform and determine how this allele swap impacts

macrophage inflammatory death mechanisms upon bacterial infection.

Acknowledgments

We thank George O'Toole, John LiPuma, Feng Shao and Sergio Grinstein for providing

bacterial strains. Ericka A. Tamayo-Guevara helped produce the TecA AlphaFold structures

shown in Fig.6. This work was supported by grants from the Cystic Fibrosis Foundation

(BLISKA22G0 and STANTO19R0) and support by DartCF through NIH grant P30-DK117469.

Materials and Methods

Bacterial strains: Bacterial strains used in this study are shown in Table 1. *B. orbicola* and *B.*

cenocepacia strains were grown at 37°C on Luria-Bertani (LB) agar plates or in LB both with

shaking.

Ethics statement: **I**solation of bone barrow cells from mice was carried out in accordance with a

protocol that adhered to the Guide for the Care and Use of Laboratory Animals of the National

Institutes of Health (NIH) and was reviewed and approved (approval #00002184) by the

Institutional Animal Care and Use Committee at Dartmouth College. The Dartmouth College

animal program is registered with the U.S. Department of Agriculture (USDA) through certificate

number 12-R-0001, operates in accordance with Animal Welfare Assurance (NIH/PHS) under

assurance number D16-00166 (A3259-01) and is accredited with the Association for

Assessment and Accreditation of Laboratory Animal Care International (AAALAC, accreditation

number 398).

Mouse strains: Wild-type (WT) C57BL/6J (stock# 000664) mice were purchased from Jackson

Laboratories. Pyrin-knockout mice ($Mefv^{-/-}$) on the C57BL/6 background were obtained from Jae

Chae and Daniel Kastner at the NIH (44) and bred in the mouse facilities at Dartmouth. Mice

with the $Cftr^{F508del}$ mutation on the C57BL/6 background were obtained from Case Western

Reserve University's Cystic Fibrosis Mouse Models Core and bred at Dartmouth (37). $Mefv^{-/-}$

and $Cftr^{F508del}$ mice were crossed to obtain breeding pairs that were $Cftr^{F508del}$ $Mefv^{+/-}$ and

$Cftr^{F508del}$ $Mefv^{-/-}$. The resulting pups were genotyped as described (37) and used as a source of

bone marrow cells.

Preparation of BMDMs: Bone marrow derived macrophages (BMDMs) were cultured from bone

marrow of mice and cultured as described previously (37). After 7 days of differentiation the

BMDMs were seeded at a density of 0.8×10^6 cells/well in 6-well plates in MGM 10/10 media

containing Dulbecco's Modified Eagle Medium (DMEM) + Glutamax (Gibco®) containing 10%

Fetal Bovine Serum (FBS) (GE®), 10% L929 cell-conditioned media, 1 mM sodium pyruvate

(Gibco®), 10 mM HEPES (Gibco®) and divided into 6-well plates at a density of 0.8×10^6

cells/well in a total volume of 3 mL. The BMDMs were primed with 100 ng/mL O26:B6

Escherichia coli LPS (Sigma®) and incubated overnight at 37°C with 5% CO_2.

<u>BMDM infection assays</u>: Overnight (16-hour) cultures of *B. orbicola* or *B. cenocepacia* were

subcultured 1:100 in fresh LB to OD_{600} = ~0.05 on infection day and shaken at 37°C until the

cultures reached mid-log phase (OD_{600} = 0.300). Cultures were then pelleted, the LB was

removed, and the bacteria were resuspended in PBS to the original volume (~3×10^8 CFU/mL).

The bacterial suspensions were then diluted to an MOI of 20 in warmed, in media lacking 10%

FBS (MGM 0/10). As a positive control for pyrin inflammasome activation, the

glucosyltransferase toxin from *Clostridium difficile* TcdB (List Biological Laboratories, Inc.) was

also diluted to 0.1mg/ml in warmed MGM 10. The BMDMs were washed once in warm 1x PBS

and 3ml fresh MGM 10 was added with bacteria or TcdB to the appropriate wells. The plates

were then centrifuged for 5 minutes at 1,000rpm to bring the bacteria and the cells in contact on

the bottom of the wells. The plates were incubated at 37°C with 5% CO_2 for 90 minutes. Cell

supernatants were collected for cytokine ELISAs and lactate dehydrogenase (LDH) assays. The

BMDMs were lysed using mammalian protein extraction reagent (M-PER, Thermo Scientific®)

with added cOmplete, Mini (Roche®) protease inhibitor and PhosSTOP (Roche®) phosphatase

inhibitor (37).

<u>Western blotting:</u> Five - 10mg of protein from the cell lysates were run on 4-12% NuPAGE Bis-

Tris SDS-PAGE gels (Invitrogen by ThermoFisher Scientific®) and transferred to PVDF

membranes (ThermoFisher Scientific®) using an iBlot 2 Gel Transfer Device (Life Technologies).

Membranes were blocked in 5% non-fat dairy milk and incubated with primary antibody

overnight. The primary antibodies used were rabbit MAb for GSDMD (Abcam®, ab209845), and

rabbit polyclonal for b-actin (Cell Signaling®, #4967). HRP-conjugated anti-rabbit (Jackson

laboratory) was used as a secondary antibody. Protein bands reacting with antibodies were

visualized using chemiluminescent detection reagent (GE Healthcare®) on an iBright FL1500

(ThermoFisher Scientific®).

IL-1β and IL-1α quantification: Murine IL-1β and IL-1α in BMDM supernatants was quantified

using ELISA kits (R&D Systems®, MLB00C and MLA00, DLB50, respectively) following the

manufacturer's instructions.

LDH quantification: LDH in BMDM supernatants was quantified using the CytoTox 96® Non-

Radioactive Cytotoxicity Assay (Promega®) following the manufacturer's instructions.

Statistical analysis of data: Experimental data generated from at least three independent in vitro

experiments are analyzed for significance. Probability (p) values for ELISA and LDH data are

calculated using one-way analysis of variance (ANOVA) or grouped two-way ANOVA with

Tukey's posttest.

TecA gene sequencing and protein sequence analysis: *B. orbicola* and *B. cenocepacia* were

grown on LB agar plates, single colonies were picked, suspended in sterile Elution Buffer

(Qiagen), and boiled for 5 minutes to release bacterial DNA. PCR was performed using the

colony DNA and primers that anneal upstream (Forward 5'-

ACGCCGTGACGGCCTGACGCCGT-3') and downstream (Reverse 5'-

GCGCGACGCCATGCGGAAATCGCCG-3') of all four *tecA* genes analyzed (BCAM1857 in

J2315, WQ49_RS04990 in K56-2, BCEN_RS18170 in AU1054 and EAY64595

in PC184). PCR products were run on a 1% agarose gel and purified using a Qiagen kit.

Purified DNA was sequenced and resulting sequences were analyzed, translated, and aligned

on SnapGene using ClustalW. Percent amino acid sequence identity was determined using

NCBI Protein BLAST. Predicted structures of the AU1054 and K56-2 TecA isoforms were

determined by AlphaFold2 (45) and displayed using PyMol.

Fig.1. Pyrin is differentially required for GSDMD cleavage in BMDMs infected with AU1054, J2315 or K56-2. LPS-primed C57BL/6 *Mefv*$^{+/+}$ or *Mefv*$^{-/-}$ BMDMs were left uninfected or infected with the indicated strains (Table 1) for 90 min at MOI 20 and lysates were analyzed by immunoblotting with antibodies to GSDMD or β-actin. Positions of GSDMD 53 kDa full length (FL) and 32 kDa cleaved forms are indicated. Lane 13 is molecular weight (MW) ladder.

Fig.2. Pyrin is differentially required for GSDMD cleavage in BMDMs infected with *B. cenocepacia* and *B. orbicola*. LPS-primed C57BL/6 *Mefv⁺/⁺* and *Mefv⁻/⁻* BMDMs were left uninfected or infected for 90 min at MOI 20 with the indicated strains (Table 1) and lysates were analyzed by immunoblotting with antibodies to GSDMD or β-actin. The result in (B) with PC184 is representative of two independent experiments. Lanes 11 are a MW ladder.

Fig.3. TecA is dispensable for pyrin-independent GSDMD cleavage and IL-1β release in

BMDMs infected with J2315. LPS-primed C57BL/6 *Mefv*$^{+/+}$ or *Mefv*$^{-/-}$ BMDMs were left

uninfected or infected with the indicated strains (Table 1) at an MOI of 20 for 90 minutes. (A)

BMDM lysates were analyzed by immunoblotting with antibodies to GSDMD or β-actin. Lane 1

is a MW ladder. Cell supernatants were collected and analyzed for IL-1β by ELISA (B,C). Data

shown in (B,C) are from four or five independent experiments, respectively. Error bars represent

standard deviation. Significance determined by two-way ANOVA and Tukey posttest as

compared to AU1054 (B) or J2315 (C) within and between BMDM genotypes. ns, not significant;

* p <0.05, ** p <0.01, *** p <0.001, **** p <0.0001.

Fig.4. TecA triggers pyrin-independent IL-1α and LDH release from BMDMs infected with

AU1054. LPS-primed *Cftr^{F508del} Mefv$^{+/+}$* or *Mefv$^{-/-}$* BMDMs were left uninfected or infected with

AU1054 or AU1054 Δ*tecA* at an MOI of 20 or intoxicated with TcdB for 90 minutes. Cell

supernatants were collected and analyzed for IL-1β (A) or IL-1α (B) by ELISA and PMR by LDH

release (C). Data shown are from at least three independent experiments. Error bars represent

standard deviation. Significance determined by two-way ANOVA as compared to AU1054 within

and between BMDM genotypes. ns, not significant; ** p <0.01, *** p <0.001, **** p <0.0001.

Fig.5. The TecA proteins of *B. cenocepacia* and *B. orbicola* exhibit amino acid variation.

The *tecA* genes of the strains used in this study were sequenced and then translated and aligned

in SnapGene using ClustalW. Amino acids highlighted in yellow indicate differences from the

TecA sequence of J2315.

Fig.6. Comparison of predicted structures of TecA proteins of *B. cenocepacia* and *B.*

***orbicola* and non-conserved amino acids adjacent to catalytic sites.** Predicted structures of

TecA proteins of K56-2 (left) and AU1054 (right) were generated using AlphaFold2 and displayed

using PyMol. Catalytic residues are shown in red (left) or cyan (right). Non-conserved residues

adjacent to catalytic sides are shown in orange (left) or yellow (right).

Species (Genomovar)	Lineage	Strain (Isolate)	Source, Collection, Reference
B. cenocepacia (Genomovar IIIA)	ET-12	J2315 (A)	Bliska strain collection, (37)
		J2315 (B)	George O'Toole
		J2315 (C; H12711)	John LiPuma, BCCM/LMG
		J2315 (D; H12718)	John LiPuma
		J2315 (E)	Feng Shao, BCCM/LMG, (32)
		J2315 ΔtecA	Feng Shao, (32)
		K56-2	George O'Toole
B. orbicola (Genomovar IIIB)	PHDC	AU1054	Peggy Cotter, (36)
		AU1054 Δhcp	Peggy Cotter, (36)
		AU1054 ΔtecA	Peggy Cotter, (37)
	Midwest	PC184	John LiPuma, (46)

Table 1. *B. cenocepacia* and *B. orbicola* isolates used in this study. Letters in parentheses

after J2315 are arbitrary identifiers for isolates used in this study. H12711 and H12718 are

additional identifiers for these isolates obtained from John LiPuma. Belgian Coordinated

Collections of Microorganisms (BCCM/LMG)

REFERENCES

1.   Loutet SA, Valvano MA. A decade of Burkholderia cenocepacia virulence determinant
research. Infect Immun. 2010;78(10):4088-100. doi: 10.1128/IAI.00212-10. PubMed PMID:
20643851; PMCID: PMC2950345.
2.   Hauser AR, Jain M, Bar-Meir M, McColley SA. Clinical significance of microbial infection
and adaptation in cystic fibrosis. Clin Microbiol Rev. 2011;24(1):29-70. Epub 2011/01/15. doi:
10.1128/CMR.00036-10. PubMed PMID: 21233507; PMCID: PMC3021203.
3.   Scoffone VC, Chiarelli LR, Trespidi G, Mentasti M, Riccardi G, Buroni S. Burkholderia
cenocepacia Infections in Cystic Fibrosis Patients: Drug Resistance and Therapeutic
Approaches. Front Microbiol. 2017;8:1592. Epub 2017/09/08. doi: 10.3389/fmicb.2017.01592.
PubMed PMID: 28878751; PMCID: PMC5572248.
4.   De Soyza A, Meachery G, Hester KL, Nicholson A, Parry G, Tocewicz K, Pillay T, Clark
S, Lordan JL, Schueler S, Fisher AJ, Dark JH, Gould FK, Corris PA. Lung transplantation for
patients with cystic fibrosis and Burkholderia cepacia complex infection: a single-center
experience. J Heart Lung Transplant. 2010;29(12):1395-404. Epub 2010/09/03. doi:
10.1016/j.healun.2010.06.007. PubMed PMID: 20810293.
5.   Vandamme P, Holmes B, Coenye T, Goris J, Mahenthiralingam E, LiPuma JJ, Govan
JR. Burkholderia cenocepacia sp. nov.--a new twist to an old story. Res Microbiol.

2003;154(2):91-6. Epub 2003/03/22. doi: 10.1016/S0923-2508(03)00026-3. PubMed PMID:
12648723.
6.   Drevinek P, Mahenthiralingam E. Burkholderia cenocepacia in cystic fibrosis:
epidemiology and molecular mechanisms of virulence. Clin Microbiol Infect. 2010;16(7):821-30.
Epub 2010/10/01. doi: 10.1111/j.1469-0691.2010.03237.x. PubMed PMID: 20880411.
7.   Ganesan S, Sajjan US. Host evasion by Burkholderia cenocepacia. Front Cell Infect
Microbiol. 2011;1:25. Epub 2011/01/01. doi: 10.3389/fcimb.2011.00025. PubMed PMID:
22919590; PMCID: PMC3417383.
8.   Valvano MA. Intracellular survival of Burkholderia cepacia complex in phagocytic cells.
Can J Microbiol. 2015;61(9):607-15. doi: 10.1139/cjm-2015-0316. PubMed PMID: 26220706.
9.   Abdulrahman BA, Khweek AA, Akhter A, Caution K, Kotrange S, Abdelaziz DH, Newland
C, Rosales-Reyes R, Kopp B, McCoy K, Montione R, Schlesinger LS, Gavrilin MA, Wewers MD,
Valvano MA, Amer AO. Autophagy stimulation by rapamycin suppresses lung inflammation and
infection by Burkholderia cenocepacia in a model of cystic fibrosis. Autophagy.
2011;7(11):1359-70. doi: 10.4161/auto.7.11.17660. PubMed PMID: 21997369; PMCID:
PMC3359483.
10.   Assani K, Tazi MF, Amer AO, Kopp BT. IFN-gamma stimulates autophagy-mediated
clearance of Burkholderia cenocepacia in human cystic fibrosis macrophages. PLoS One.
2014;9(5):e96681. Epub 2014/05/07. doi: 10.1371/journal.pone.0096681. PubMed PMID:
24798083; PMCID: PMC4010498.
11.   Schwab U, Abdullah LH, Perlmutt OS, Albert D, Davis CW, Arnold RR, Yankaskas JR,
Gilligan P, Neubauer H, Randell SH, Boucher RC. Localization of Burkholderia cepacia complex
bacteria in cystic fibrosis lungs and interactions with Pseudomonas aeruginosa in hypoxic
mucus. Infect Immun. 2014;82(11):4729-45. doi: 10.1128/IAI.01876-14. PubMed PMID:
25156735; PMCID: PMC4249344.
12.   Sousa SA, Ulrich M, Bragonzi A, Burke M, Worlitzsch D, Leitao JH, Meisner C, Eberl L,
Sa-Correia I, Doring G. Virulence of Burkholderia cepacia complex strains in gp91phox-/- mice.
Cell Microbiol. 2007;9(12):2817-25. doi: 10.1111/j.1462-5822.2007.00998.x. PubMed PMID:
17627623.
13.   Al-Khodor S, Marshall-Batty K, Nair V, Ding L, Greenberg DE, Fraser ID. Burkholderia
cenocepacia J2315 escapes to the cytosol and actively subverts autophagy in human
macrophages. Cell Microbiol. 2014;16(3):378-95. Epub 2013/10/15. doi: 10.1111/cmi.12223.
PubMed PMID: 24119232; PMCID: PMC3927784.
14.   Dorrington MG, Bradfield CJ, Lack JB, Lin B, Liang JJ, Starr T, Ernst O, Gross JL, Sun
J, Miller AH, Steele-Mortimer O, Fraser IDC. Type I IFNs facilitate innate immune control of the
opportunistic bacteria Burkholderia cenocepacia in the macrophage cytosol. PLoS Pathog.
2021;17(3):e1009395. Epub 2021/03/09. doi: 10.1371/journal.ppat.1009395. PubMed PMID:
33684179; PMCID: PMC7971856.
15.   Lamothe J, Huynh KK, Grinstein S, Valvano MA. Intracellular survival of Burkholderia
cenocepacia in macrophages is associated with a delay in the maturation of bacteria-containing
vacuoles. Cell Microbiol. 2007;9(1):40-53. Epub 2006/07/28. doi: 10.1111/j.1462-
5822.2006.00766.x. PubMed PMID: 16869828.
16.   Keith KE, Hynes DW, Sholdice JE, Valvano MA. Delayed association of the NADPH
oxidase complex with macrophage vacuoles containing the opportunistic pathogen Burkholderia
cenocepacia. Microbiology (Reading). 2009;155(Pt 4):1004-15. Epub 2009/04/01. doi:
10.1099/mic.0.026781-0. PubMed PMID: 19332803.
17.   Assani K, Shrestha CL, Robledo-Avila F, Rajaram MV, Partida-Sanchez S, Schlesinger
LS, Kopp BT. Human Cystic Fibrosis Macrophages Have Defective Calcium-Dependent Protein
Kinase C Activation of the NADPH Oxidase, an Effect Augmented by Burkholderia cenocepacia.
J Immunol. 2017;198(5):1985-94. Epub 2017/01/18. doi: 10.4049/jimmunol.1502609. PubMed
PMID: 28093527; PMCID: PMC5322234.

18.   Krause K, Caution K, Badr A, Hamilton K, Saleh A, Patel K, Seveau S, Hall-Stoodley L,
Hegazi R, Zhang X, Gavrilin MA, Amer AO. CASP4/caspase-11 promotes autophagosome
formation in response to bacterial infection. Autophagy. 2018;14(11):1928-42. Epub 2018/09/01.
doi: 10.1080/15548627.2018.1491494. PubMed PMID: 30165781; PMCID: PMC6152495.
19.   Hunt TA, Kooi C, Sokol PA, Valvano MA. Identification of Burkholderia cenocepacia
genes required for bacterial survival in vivo. Infect Immun. 2004;72(7):4010-22. Epub
2004/06/24. doi: 10.1128/IAI.72.7.4010-4022.2004. PubMed PMID: 15213146; PMCID:
PMC427415.
20.   Wood TE, Howard SA, Wettstadt S, Filloux A. PAAR proteins act as the 'sorting hat' of
the type VI secretion system. Microbiology (Reading). 2019;165(11):1203-18. Epub 2019/08/06.
doi: 10.1099/mic.0.000842. PubMed PMID: 31380737; PMCID: PMC7376260.
21.   Hachani A, Wood TE, Filloux A. Type VI secretion and anti-host effectors. Curr Opin
Microbiol. 2016;29:81-93. Epub 2016/01/02. doi: 10.1016/j.mib.2015.11.006. PubMed PMID:
26722980.
22.   Spiewak HL, Shastri S, Zhang L, Schwager S, Eberl L, Vergunst AC, Thomas MS.
Burkholderia cenocepacia utilizes a type VI secretion system for bacterial competition.
Microbiologyopen. 2019:e774. Epub 2019/01/11. doi: 10.1002/mbo3.774. PubMed PMID:
30628184; PMCID: PMC6612558.
23.   Flannagan RS, Jaumouille V, Huynh KK, Plumb JD, Downey GP, Valvano MA, Grinstein
S. Burkholderia cenocepacia disrupts host cell actin cytoskeleton by inactivating Rac and
Cdc42. Cell Microbiol. 2012;14(2):239-54. Epub 2011/10/26. doi: 10.1111/j.1462-
5822.2011.01715.x. PubMed PMID: 22023324.
24.   Rosales-Reyes R, Skeldon AM, Aubert DF, Valvano MA. The Type VI secretion system
of Burkholderia cenocepacia affects multiple Rho family GTPases disrupting the actin
cytoskeleton and the assembly of NADPH oxidase complex in macrophages. Cell Microbiol.
2012;14(2):255-73. Epub 2011/10/26. doi: 10.1111/j.1462-5822.2011.01716.x. PubMed PMID:
22023353.
25.   Gavrilin MA, Abdelaziz DH, Mostafa M, Abdulrahman BA, Grandhi J, Akhter A, Abu
Khweek A, Aubert DF, Valvano MA, Wewers MD, Amer AO. Activation of the pyrin
inflammasome by intracellular Burkholderia cenocepacia. J Immunol. 2012;188(7):3469-77. doi:
10.4049/jimmunol.1102272. PubMed PMID: 22368275; PMCID: PMC3482472.
26.   Rosales-Reyes R, Aubert DF, Tolman JS, Amer AO, Valvano MA. Burkholderia
cenocepacia type VI secretion system mediates escape of type II secreted proteins into the
cytoplasm of infected macrophages. PLoS One. 2012;7(7):e41726. Epub 2012/08/01. doi:
10.1371/journal.pone.0041726. PubMed PMID: 22848580; PMCID: PMC3405007.
27.   Xu H, Yang J, Gao W, Li L, Li P, Zhang L, Gong YN, Peng X, Xi JJ, Chen S, Wang F,
Shao F. Innate immune sensing of bacterial modifications of Rho GTPases by the Pyrin
inflammasome. Nature. 2014;513(7517):237-41. doi: 10.1038/nature13449. PubMed PMID:
24919149.
28.   Zhao Y, Shao F. Diverse mechanisms for inflammasome sensing of cytosolic bacteria
and bacterial virulence. Curr Opin Microbiol. 2016;29:37-42. Epub 2015/11/13. doi:
10.1016/j.mib.2015.10.003. PubMed PMID: 26562791.
29.   Shi J, Gao W, Shao F. Pyroptosis: Gasdermin-Mediated Programmed Necrotic Cell
Death. Trends Biochem Sci. 2017;42(4):245-54. doi: 10.1016/j.tibs.2016.10.004. PubMed PMID:
27932073.
30.   Estfanous S, Krause K, Anne MNK, Eltobgy M, Caution K, Abu Khweek A, Hamilton K,
Badr A, Daily K, Carafice C, Baetzhold D, Zhang X, Li T, Wen H, Gavrilin MA, Haffez H, Soror
S, Amer AO. Gasdermin D restricts Burkholderia cenocepacia infection in vitro and in vivo. Sci
Rep. 2021;11(1):855. Epub 2021/01/15. doi: 10.1038/s41598-020-79201-5. PubMed PMID:
33441602; PMCID: PMC7807041.

31.   Broz P, Monack DM. Noncanonical inflammasomes: caspase-11 activation and effector
mechanisms. PLoS Pathog. 2013;9(2):e1003144. Epub 2013/03/08. doi:
10.1371/journal.ppat.1003144
PPATHOGENS-D-12-02639 [pii]. PubMed PMID: 23468620; PMCID: 3585133.
32.   Aubert DF, Xu H, Yang J, Shi X, Gao W, Li L, Bisaro F, Chen S, Valvano MA, Shao F. A
Burkholderia Type VI Effector Deamidates Rho GTPases to Activate the Pyrin Inflammasome
and Trigger Inflammation. Cell Host Microbe. 2016;19(5):664-74. doi:
10.1016/j.chom.2016.04.004. PubMed PMID: 27133449.
33.   Walpole GFW, Plumb JD, Chung D, Tang B, Boulay B, Osborne DG, Piotrowski JT, Catz
SD, Billadeau DD, Grinstein S, Jaumouille V. Inactivation of Rho GTPases by Burkholderia
cenocepacia Induces a WASH-Mediated Actin Polymerization that Delays Phagosome
Maturation. Cell Rep. 2020;31(9):107721. Epub 2020/06/04. doi: 10.1016/j.celrep.2020.107721.
PubMed PMID: 32492429; PMCID: PMC7315377.
34.   Wallner A, King E, Ngonkeu ELM, Moulin L, Bena G. Genomic analyses of Burkholderia
cenocepacia reveal multiple species with differential host-adaptation to plants and humans.
BMC Genomics. 2019;20(1):803. Epub 2019/11/07. doi: 10.1186/s12864-019-6186-z. PubMed
PMID: 31684866; PMCID: PMC6829993.
35.   Morales-Ruiz LM, Rodriguez-Cisneros M, Kerber-Diaz JC, Rojas-Rojas FU, Ibarra JA,
Estrada-de Los Santos P. Burkholderia orbicola sp. nov., a novel species within the
Burkholderia cepacia complex. Arch Microbiol. 2022;204(3):178. Epub 2022/02/18. doi:
10.1007/s00203-022-02778-0. PubMed PMID: 35174425.
36.   Perault AI, Chandler CE, Rasko DA, Ernst RK, Wolfgang MC, Cotter PA. Host
Adaptation Predisposes Pseudomonas aeruginosa to Type VI Secretion System-Mediated
Predation by the Burkholderia cepacia Complex. Cell Host Microbe. 2020;28(4):534-47 e3.
Epub 2020/08/07. doi: 10.1016/j.chom.2020.06.019. PubMed PMID: 32755549; PMCID:
PMC7554260.
37.   Loeven NA, Perault AI, Cotter PA, Hodges CA, Schwartzman JD, Hampton TH, Bliska
JB. The Burkholderia cenocepacia Type VI Secretion System Effector TecA Is a Virulence
Factor in Mouse Models of Lung Infection. mBio. 2021;12(5):e0209821. Epub 2021/09/29. doi:
10.1128/mBio.02098-21. PubMed PMID: 34579569; PMCID: PMC8546862.
38.   Di Paolo NC, Shayakhmetov DM. Interleukin 1alpha and the inflammatory process. Nat
Immunol. 2016;17(8):906-13. Epub 2016/07/21. doi: 10.1038/ni.3503. PubMed PMID:
27434011; PMCID: PMC5152572.
39.   Holden MT, Seth-Smith HM, Crossman LC, Sebaihia M, Bentley SD, Cerdeno-Tarraga
AM, Thomson NR, Bason N, Quail MA, Sharp S, Cherevach I, Churcher C, Goodhead I, Hauser
H, Holroyd N, Mungall K, Scott P, Walker D, White B, Rose H, Iversen P, Mil-Homens D, Rocha
EP, Fialho AM, Baldwin A, Dowson C, Barrell BG, Govan JR, Vandamme P, Hart CA,
Mahenthiralingam E, Parkhill J. The genome of Burkholderia cenocepacia J2315, an epidemic
pathogen of cystic fibrosis patients. J Bacteriol. 2009;191(1):261-77. Epub 2008/10/22. doi:
10.1128/JB.01230-08. PubMed PMID: 18931103; PMCID: PMC2612433.
40.   Kayagaki N, Kornfeld OS, Lee BL, Stowe IB, O'Rourke K, Li Q, Sandoval W, Yan D,
Kang J, Xu M, Zhang J, Lee WP, McKenzie BS, Ulas G, Payandeh J, Roose-Girma M,
Modrusan Z, Reja R, Sagolla M, Webster JD, Cho V, Andrews TD, Morris LX, Miosge LA,
Goodnow CC, Bertram EM, Dixit VM. NINJ1 mediates plasma membrane rupture during lytic
cell death. Nature. 2021;591(7848):131-6. Epub 2021/01/21. doi: 10.1038/s41586-021-03218-7.
PubMed PMID: 33472215.
41.   Fink SL, Cookson BT. Apoptosis, pyroptosis, and necrosis: mechanistic description of
dead and dying eukaryotic cells. Infect Immun. 2005;73(4):1907-16. Epub 2005/03/24. doi:
73/4/1907 [pii]
10.1128/IAI.73.4.1907-1916.2005. PubMed PMID: 15784530; PMCID: 1087413.

42.   Serbina NV, Jia T, Hohl TM, Pamer EG. Monocyte-mediated defense against microbial
pathogens. Annu Rev Immunol. 2008;26:421-52. Epub 2008/02/29. doi:
10.1146/annurev.immunol.26.021607.090326. PubMed PMID: 18303997.
43.   Heung LJ, Hohl TM. Inflammatory monocytes are detrimental to the host immune
response during acute infection with Cryptococcus neoformans. PLoS Pathog.
2019;15(3):e1007627. Epub 2019/03/22. doi: 10.1371/journal.ppat.1007627. PubMed PMID:
30897162; PMCID: PMC6428256.
44.   Chae JJ, Cho YH, Lee GS, Cheng J, Liu PP, Feigenbaum L, Katz SI, Kastner DL. Gain-
of-function Pyrin mutations induce NLRP3 protein-independent interleukin-1beta activation and
severe autoinflammation in mice. Immunity. 2011;34(5):755-68. doi:
10.1016/j.immuni.2011.02.020. PubMed PMID: 21600797; PMCID: PMC3129608.
45.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool
K, Bates R, Zidek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A,
Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D, Clancy E,
Zielinski M, Steinegger M, Pacholska M, Berghammer T, Bodenstein S, Silver D, Vinyals O,
Senior AW, Kavukcuoglu K, Kohli P, Hassabis D. Highly accurate protein structure prediction
with AlphaFold. Nature. 2021;596(7873):583-9. Epub 2021/07/16. doi: 10.1038/s41586-021-
03819-2. PubMed PMID: 34265844; PMCID: PMC8371605 have filed non-provisional patent
applications 16/701,070 and PCT/EP2020/084238, and provisional patent applications
63/107,362, 63/118,917, 63/118,918, 63/118,921 and 63/118,919, each in the name of
DeepMind Technologies Limited, each pending, relating to machine learning for predicting
protein structures. The other authors declare no competing interests.
46.   Kim JY, Sajjan US, Krasan GP, LiPuma JJ. Disruption of tight junctions during traversal
of the respiratory epithelium by Burkholderia cenocepacia. Infect Immun. 2005;73(11):7107-12.
Epub 2005/10/22. doi: 10.1128/IAI.73.11.7107-7112.2005. PubMed PMID: 16239504; PMCID:
PMC1273860.
677