**Human disturbance increases spatiotemporal associations among mountain forest terrestrial mammal species**

Xueyou Li[1], William V. Bleisch[2], Wenqiang Hu[1], Quan Li[1], Hongjiao Wang[1], Zhongzheng Chen[3], Ru Bai[1], Xuelong Jiang[1]

[1] State Key Laboratory of Genetic Resources and Evolution & Yunnan key laboratory of biodiversity and ecological conservation of Gaoligong Mountain, Kunming Institute of Zoology, Chinese Academy of Sciences; [2] China Exploration and Research Society, 2707-08 SouthMark, Wong Chuk Hang, Hong Kong, China; [3] Anhui Provincial Key Laboratory of the Conservation and Exploitation of Biological Resources, College of Life Sciences, Anhui Normal University, Wuhu, China

**For correspondence:** lixueyou@mail.kiz.ac.cn (XL); jiangxl@mail.kix.ac.cn (XJ)

**Abstract**

Spatial and temporal associations between sympatric species underpin biotic interactions, structure ecological assemblages, and sustain ecosystem functioning and stability. However, the resilience of interspecific spatiotemporal associations to human activity remains poorly understood, particularly in mountain forests where anthropogenic impacts are often pervasive. Here we applied context-dependent Joint Species Distribution Models to a systematic camera-trap survey dataset from a global biodiversity hotspot in eastern Himalaya to understand how prominent human activities in mountain forests influence species associations within terrestrial mammal communities. We categorized anthropogenic pressures into chronic human modification and acute direct human presence and compared their effects on spatiotemporal associations among species. We obtained 10,388 independent detections of 17 focal species (12 carnivores and five ungulates) from 322 stations over 43,163 camera days of effort. We identified a higher incidence of positive associations in habitats with higher levels of human modification (87 %) and human presence (83 %) compared to those located in habitats with lower human modification (64 %) and human presence (65 %) levels. We also detected a significant reduction of pairwise encounter time at increasing levels of human disturbance (human modification: mean = -1.07, 95% CI = -1.39 to -0.73; human presence: mean = -0.69, 95% CI = -0.92 to -0.46), corresponding to more frequent encounters between pairs of species. Our findings indicate that human activities can push mammals together into more frequent encounters and associations, which likely influences the coexistence and persistence of wildlife, with potential far-ranging ecological consequences. With increasing

33  human presence and human modification of areas throughout the world, identifying

34  thresholds of anthropogenic activity that shift species relationships, limiting human activity

35  and increasing landscape connectivity across protected areas may be imperative to maintain

36  interspecific spatiotemporal associations that underpin ecosystem resilience.

## Introduction

Pervasive human activities can disrupt invisible facets of biodiversity such as species associations, with potential cascading ecosystem effects (*Naeem et al., 1994*; *Parsons et al., 2022*). Human encroachment into natural ecosystems squeezes the spatiotemporal niches of wildlife species (*Gilbert et al., 2022*), altering the number and magnitude of associations in a community (*Burkle et al., 2013*), and accelerating species decline and loss (*Jones et al., 2018*). Humans play a role as 'super predators' in shaping the co-occurrence of other species with complicated indirect modifications to multiple interactions between organisms (*Gilbert et al., 2022*; *Moll et al., 2021*). Human disturbance, such as chronic landscape modification and acute direct human presence, may significantly alter spatiotemporal distribution of species and fundamentally change the way that species interact (*Li et al., 2022a*; *Sévêque et al., 2020*). For example, expanding human footprints have compressed the space and time available for mammals to share by restricting animal movements (*Tucker et al., 2018*) and increasing wildlife nocturnality (*Gaynor et al., 2018*). As human and animal activities increasingly overlap in time and space, it is important to assess and quantify the potential for human-induced changes in species association to ecosystem structure and function (*Penjor et al., 2022*).

Species are not distributed independently of each other; rather, they co-occur in time and space and interact (*Tilman, 1994*; *Wisz et al., 2013*). The spatiotemporal co-occurrence of species, termed interspecific associations, provides unique ecological information and has important consequence for ecosystem integrity (*Gorczynski et al., 2022*; *Keil et al., 2021*).

However, segregated species co-occurrence could be generated by processes such as negative

interspecific interaction, distinct environment requirements, and dispersal limitations;

similarly, aggregated species pairs may reflect positive interspecific interaction, but could

also reflect shared environmental preferences (*Kathleen Lyons et al., 2016*; *Song et al., 2020*).

Although interspecific co-occurrence or associations cannot be used to directly estimate

species interactions (*Blanchet et al., 2020*), animals cannot interact if their spatiotemporal

niche does not overlap (*Gilbert et al., 2022*). Thus, interspecific associations convey key

information about interactions between sympatric species (*Boron et al., 2023*).

Disruptions to the spatiotemporal relationships of species can result in serious ecological

consequences including alteration of community structure (*Tulloch et al., 2018*), upsetting the

competitive balance between species (*Boron et al., 2023*), increasing disease transmission

(*Hassell et al., 2017*), and accelerating local extinction (*Fidino et al., 2019*). This can

significantly upset the distribution of ecological functions that species provide, ultimately

influencing ecosystem dynamics (*Gardner et al., 2019*). Taken as a whole, associations

between sympatric species play a fundamental role in community assembly and are intricately

related to ecosystem stability and resilience (*Boron et al., 2023*). Thus, understanding how

interspecific spatiotemporal associations change across human disturbance gradients provides

valuable insight into the long-term implications of human impacts on ecosystem function and

recovery relevant to biodiversity conservation in the Anthropocene.

Species associations are not necessarily fixed as positive or negative, but can vary along

environmental gradients (*Davis et al., 2018*; *Pellissier et al., 2018*; *Perrin et al., 2022*).

Associations between free-ranging species are especially context-dependent (*Chamberlain et*

*al., 2014*). Therefore, a key challenge in community ecology is to identify the conditions

under which negative and positive species associations are more likely to occur (*Beaudrot et*

*al., 2020*). Understanding how species associations vary along environmental conditions is

pivotal in making robust ecological predictions about biodiversity responses to changing

environmental conditions (*Dangles et al., 2018*; *Perrin et al., 2022*). For example, alpine bird

communities have a higher frequency of positive interactions in grasslands with low

productivity compared with in forests with high productivity (*García-Navas et al., 2021*).

However, the resilience of interspecific spatiotemporal associations of terrestrial mammals to

human activity remains poorly understood, particularly in mountain forests where

anthropogenic impacts are often pervasive.

Medium and large-sized terrestrial mammals are a key components of mountain forest

communities. They play crucial roles in maintaining biodiversity and ecosystem functions

(*Lacher et al., 2019*), and are especially vulnerable to anthropogenic activities (*Li et al.,*

*2022b*). Here, we set out to investigate the spatial and temporal patterns of occurrence and the

interspecific associations within a terrestrial mammalian community along human disturbance

gradients. We executed a systematic camera trapping survey spanning 4,100 km$^2$ of the

Yarlung Zangbo Grand Canyon National Nature Reserve in the eastern Himalayas at a total of

322 camera trapping stations (*Supplementary file Figure S1*), obtaining detections of both

wildlife and humans. We classified human disturbances into two groups: human modification

(i.e. relating to habitat modifications by humans) or human presence (i.e. referring to the

direct presence of humans) disturbances. We employed a context-dependent joint species

distribution model (JSDM; *Tikhonov et al., 2017*) to infer pairwise species associations along

human disturbance gradients. We used kernel density distributions of animal diel activity and

time between detections of species pairs (time-to-encounter) to compare temporal

associations at low and high human disturbances. We consider two hypotheses regarding

temporal and spatial interspecific associations. The spatial compress hypothesis posits that

human modification on habitats would suppress space available for animals to share (*Tilman*

*et al., 2017*), resulting in increased positive spatial associations between species. The

temporal compress hypothesis postulates that fear of direct human presence would compress

diurnal activity in animals (*Gaynor et al., 2018*), resulting in reduced encounter time (or

increased encounter rate) between species (*Figure 1*). Our study incorporates two different

types of human disturbances in the analysis to elucidate the effects of humans on

multidimensional (i.e., space and time) species associations. We then consider implications

for conservation.

**Results**

Our camera traps obtained 10,388 independent detections of 17 focal species (12 carnivores

and five ungulates) from 322 stations over 43,163 camera days of effort (*Supplementary file*

*Table S1*). We documented a number of species and subspecies of conservation concern,

including Bengal tiger *Panthera trigris trigris*, clouded leopard *Neofelis nebulosa* and dhole

*Cuon alpinus*. We also captured 2,224 independent detections of humans during the survey

period.

**Species-specific response to habitat covariates**

Species varied in their responses to habitat covariates. Nine out of 17 species showed strong

positive response to forest cover (*Figure 2*). Human presence had significantly negative

effects on the occurrence of clouded leopard (mean = -0.967, 95% CI = -2.35 to -0.025), takin

*Budorcas taxicolor* (mean = -0.449, 95% CI = -0.935 to -0.071) and red goral *Naemorhedus*

*baileyi* (mean = -0.806, 95% CI = -0.935 to -0.071), but was positively associated with the

occurrence of wild boar *Sus scrofa* (mean = 0.521, 95% CI = 0.045 to 1.093) and golden cat

*Catopuma badia* (mean = 0.427, 95% CI = 0.165 to 0.804). Human modification was

negatively associated with the occurrence of dhole (mean = -0.24, 95% CI = -0.493 to -0.027),

red fox *Vulpes vulpes* (mean = -1.588, 95% CI = -2.656 to -0.769), Asiatic black bear *Ursus*

*thibetanus* (mean = -0.234, 95% CI = -0.42 to -0.056), red goral (mean = -0.665, 95% CI =

-1.129 to         -0.238) and Mainland serow *Capricornis sumatraensis* (mean = -0.477, 95%

CI = -0.697 to     -0.26), but was positively associated with muntjac *Muntjac* spp. (mean =

3.632, 95% CI = 0.946 to 9.999), masked palm civet *Paguma larvata* (mean = 2.076, 95% CI

= 1.26 to 3.145), large Indian civet *Viverra zibetha* (mean = 0.782, 95% CI = 0.175 to 1.558)

and marbled cat *Pardofelis marmorata* (mean = 0.689, 95% CI = 0.232 to 1.236; *Figure 2*)).

**Effects of human disturbances on spatial co-occurrence**

Out of the 136 estimated pairwise residual correlation coefficients in occupancy, 87 (64 %)

were positive at lower human modifications (*Figure 3a*). At the moderate and higher

modifications, the species pairs with positive associations increased to 107 (79 %) and 118

(87%) respectively (*Figure 3b,c*). At lower modifications, correlation coefficients for 18

species pairs were positive and had a 95 % CI that did not overlap zero, and the number

increased to 65 in moderate modifications but dropped to 29 at higher modifications

(*Supplementary file Figure S2*).

Along human presence gradients, 88 pairwise residual correlation coefficients (65 %) at

lower human presence habitats were positive or close to neutral (*Figure 4a*). At the moderate

and higher human presence habitats, the species pairs with positive associations increased to

115 (85 %) and 113 (83%) respectively (*Figure 4b,c*). The significant positive associations at

low, moderate and higher human presence habitats were 6 (4%), 76 (56 %), and 44 (32%)

respectively (*Supplementary file Figure S3*).

**Effects of human disturbances on temporal co-occurrence**

Human presence was associated with significantly increased nocturnality of carnivores (mean

= 0.163, 95% CI = 0.089 to 0.236), but showed no significant effects on ungulates (mean =

-0.004, 95% CI = -0.031 to 0.023) and combination of guilds (mean = -0.001, 95% CI =

-0.028 to 0.026; *Figure 5*). Human modification had no strong effects on nocturnality of any

guilds and combination of guilds (i.e. 95 % CIs include zero, *Supplementary file Figure S4*).

Both human modification and human presence were associated with significantly

reduced time between detections of pairs (human modification: mean = -1.07, 95% CI = -1.39

to -0.73; human presence: mean = -0.69, 95% CI = -0.92 to -0.46; *Figure 6*).

**Discussion**

Despite accumulating evidence of widespread impacts of humans on wildlife distribution and

activity patterns, our understanding of how different types of anthropogenic pressures reshape

species associations remains limited (*Van Scoyoc et al., 2023*). Here, we compared the

impacts of human modification (spatial compress effects) and human presence (temporal

compress effects) on spatiotemporal associations among threatened terrestrial mammals based

on systematic camera-trapping data in an understudied Himalayan landscape. Our results

show a strong influence of humans on species co-occurrence patterns. The overall results of

the spatiotemporal associations across levels of human modification and human presence

suggest that humans are associated with increasing positive spatiotemporal associations

among species. Specifically, we detected a higher incidence of positive associations in

habitats with moderate and higher levels of human modification and human presence

compared to those located in habitats with lower disturbance levels (*Figure 3, 4*). On the

temporal axis, we detected a significant reduction of pairwise encounter time at increasing

levels of human disturbance (*Figure 6*). Our results are therefore demonstrate that human

disturbance can upset interspecific associations both on spatial and temporal niche

dimensions.

Interspecific spatial associations can arise from species interaction, response to

environmental covariates and common dispersal barriers (*Blanchet et al., 2020*; d*e Jonge et*

*al., 2021*; *Poggiato et al., 2021*). Our context-dependent JSDM, which accounts for

similarities and disparities in species-specific response to habitat covariates considered

relevant to medium- and large-bodied mammal species, detected a higher prevalence of

interspecific spatial associations in human disturbed habitats. Humans can have "bottom-up"

impacts on animal distribution and associations by land use change and habitat modification

(*Riggio et al., 2020*; *Tucker et al., 2018*). Human activities can also have top-down impacts

on spatiotemporal associations among animals by directly or indirectly changing the

landscape of fear (*Palmer et al., 2022*; *Suraci et al., 2019*). Previous studies suggest that

human disturbance may compress the space and time available for communities to use,

resulting in increased frequency of positive associations (Gilbert et al., 2022; Sévêque et al.,

2022). We interpret our results to mean that anthropogenic presence and disturbance reduce

available habitats for wildlife, causing a greater predominance of positive associations in

anthropogenic landscapes. On the one hand, human presence, such as while grazing livestock

and gathering resource, generates landscapes of fear for wildlife (*Gaynor et al., 2019*; *Palmer*

*et al., 2022*; *Suraci et al., 2019*) and may downgrade habitat quality by overexploitation

(*Filazzola et al., 2020*). On the other hand, human modification such as land-use change often

constrains the realized niche space of wildlife, restricting animal movement (*Smith et al.,*

*2018*; *Tucker et al., 2018*). Overall, our study provides one of the first tests of whether

positive spatiotemporal associations between terrestrial mammals increase along gradients of

different types of anthropogenic pressures. Our results add to the growing body of evidence

that suggests anthropogenic activities reduce available niche space for animals, causing

observed positive spatial associations among species (*Gilbert et al., 2022*; *Gorczynski et al.,*

*2022*; *Murphy et al., 2021*).

Identifying thresholds of anthropogenic activity that shift species behavior and

co-occurrence will be key to drawing useful inference from human impact studies and

improving our knowledge on when altered associations may lead to reverberating impacts on

ecosystems (*Van Scoyoc et al., 2023*; *Wilson et al., 2020*). We note that the number of species

pairs with significant positive associations rapidly increased from lower to moderate levels of

human disturbance, but dropped from mean to higher level of human disturbance

(*Supplementary file Figure S2-3*). Such patterns are consistent across human modification and

human presence levels, indicating a threshold after which the positive effects of human

disturbance on species associations were dampened. Human disturbance is an important

factor in shaping species distributions (*Jones et al., 2018*; *Samia et al., 2015*). Wildlife often

has a limited tolerance threshold to human activity (*Polaina et al., 2018*; *Samia et al., 2015*;

*Smith et al., 2019*). When human pressures reach levels that preclude a species occurring at a

site (*Polaina et al., 2018*), prediction of species associations in an anthropogenic context may

become uninformative. More work needs to be done to further clarify the mechanisms driving

the observed co-occurrence patterns.

Anthropogenic activities can also shift temporal niche of animals (*Gaynor et al., 2018*; *Li*

*et al., 2022a*), possibly altering encounter rates among species and trophic dynamics that

structure communities (*Gilbert et al., 2022*; *Karanth et al., 2017*; *Mills and Harris, 2020*).

Congruent with our hypothesis, we observed significant reduction of pairwise encounter time

at increasing levels of human disturbance, corresponding to more frequent encounters

between the pairs (*Gilbert et al., 2022*). Although we did not find a significant shift in

nocturnality at community level, we detected a significant shift to nocturnal activity of

carnivores, indicating different sensitivities to human presence among the carnivores and

ungulates in our study system. Fear of humans may explain the temporal response of wildlife

to human presence. Numerous studies have shown that fear of humans as "super predators"

can have suppressive effects on widlife activity, which may contribute to increased temporal

overlap among speceis (e.g., *Sévêque et al., 2022*; *Suraci et al., 2019*).

All in all, our results indicate that both human modification and human presence may

rewire species interactions by increasing spatial and temporal co-occurrence. Studies on the

impacts of humans on wildlife communities should explicitly account for different types of

co-occurring disturbances (*Li et al., 2022a*). Our camera trapping survey observed that human

presence is pervasive even inside this remote protected area in the Tibetan Autonomous

Region, indicating that wildlife and their habitats are exposed to frequent human disturbance.

Humans exploit resources in protected areas in many ways, including through livestock

herding, resource gathering, illegal hunting, and recreation, all of which impact wildlife and

their habitats to varying degrees (*Harris et al., 2019*; *Mills and Harris, 2020*). Our results

demonstrate prevalent disruptions to species co-occurrence patterns from humans. If we are to

preserve biodiversity in protected area, we must work to understand the negative effects on

multiple facets of biodiversity from co-occurring anthropogenic pressures. Only then can we

design effective mitigation measures.

Although interspecific associations should not be directly interpreted as a signal of biotic

interactions between pairs of species (*Blanchet et al., 2020*; *Poggiato et al., 2021*), it

describes a unique facet of biodiversity, and can provide important insights into the

coexistence and persistence of wildlife as well as ecosystem function in settings with

anthropogenic activity (*Keil et al., 2021*; *Lai et al., 2020*; *Van Scoyoc et al., 2023*). Since

interspecific associations often increases with the ecological similarity of species involved

(*Gorczynski et al., 2022*), the tendency of many species towards positive associations with

increase in anthropogenic pressures indicates a trend towards homogenization of terrestrial

mammal communities. Such human-mediated changes in species co-occurrence patterns can

have serious ecological consequences at multiple scales (*Gilbert et al., 2022*; *Gorczynski et*

*al., 2022*). For individual species, increased positive associations may affect fitness and

population dynamics, increasing local extinction rates (*Kuussaari et al., 2009*; *Parsons et al.,*

*2022*). Increase in spatial aggregation among functionally similar species may be indicative of

a hidden extinction debt (*Kuussaari et al., 2009*). At the community level, increased positive

associations may depress co-occurrence network complexity and stability, amplify

interactions such as predation, and simplify communities with similar traits or co-occurrence

patterns (*Manlick and Pauli, 2020*; *Mills and Harris, 2020*). Also increased encounter rates

between species may expand disease transmission across communities (*Hassell et al., 2017*).

In addition, wildlife can host a variety of zoonotic disease (*Wicker et al., 2017*) and

populations unnaturally associated with humans are more likely to transmit pathogens (*Jones*

*et al., 2013*). We observed that several species such as masked palm civet and wild boar are

positively associated with humans in our study area, indicating a substantial overlap between

their habitats and human activity. Ongoing habitat modification, livestock grazing and

resource gathering in this region may increase probability of pathogen exchange.

The effects of human disturbance on species associations might be scale-dependent

(*Gilbert et al., 2022*). At present, our sampling design only considers the effects of

cumulative human modification and instantaneous local human presence on spatial

associations and temporal encounter time at each camera trapping station. We are not able to

compare the effects of human disturbance at broader scales (e.g., landscape scale) as such an

analysis requires adequate data from hierarchical samplings. Estimating the effects of human

disturbance at different spatial scale on species associations is a promising approach. Thus,

we encourage future work to further clarify the multiscale ecological effects of human

disturbance on species associations.

In conclusion, our study demonstrated that anthropogenic pressures increase

spatiotemporal associations of terrestrial mammals from low to mean levels of human

disturbances, but the frequency of positive spatial associations dropped from mean to higher

level of human disturbances. Such variations in species associations likely influence the

coexistence and persistence of wildlife, with potentially far-ranging ecological consequences.

Because terrestrial mammals like carnivores and ungulates play fundamental roles in

regulating montane forest ecosystems, the prevalent disruptions to their associations may

precipitate biodiversity loss and impair ecosystem function. With increasing human presence

and human modification of areas throughout the world, identifying thresholds of

anthropogenic activity that shift species relationships, limiting human activity and increasing

landscape connectivity across protected areas may be imperative to maintain interspecific

spatiotemporal associations that underpin ecosystem resilience. Moreover, the methods we

applied highlight the utility of camera trapping surveys in studying the spatiotemporal

relationships among elusive species in settings with anthropogenic activities.

**Materials and methods**

**Study area**

The study was carried out inside the Yarlung Zangbo Grand Canyon National Nature Reserve

(29°05′–30°02′ N, 94°39′–96°6′ E) in Nyingchi City in the southeast Tibet Autonomous

Region of China (*Supplementary file Figure S1*). This area is situated within the Eastern

Himalaya Biodiversity Hotspot, a globally important region for biodiversity conservation (*Li*

*et al., 2021*). The Yarlung Zangbo Grand Canyon is the deepest in the world, with an

elevation drop of more than 7000 m, and has the reputation of "Gene Bank of Mountain

Biological Resources" and "Natural Vegetation Museum" (*Duan et al., 2022*). This region is

characterized by dramatic vertical zonation of vegetation. From the valley bottom to the

mountain peak, the main vegetation types consist of low mountain tropical monsoon rain

forest, subtropical mountain evergreen broad-leaved forest, mid-mountain warm coniferous

forest, sub-alpine cold coniferous forest, alpine subfrigid shrub meadow, and periglacial

alpine vegetation (*Deng et al., 2011*). The complete landscape in the region harbors

endangered species such as Bengal tiger *Panthera tigris tigris* (*Li et al., 2023*). Human

activities such as decentralized residential settlements and free-ranging livestock grazing is

prevalent in the region, even within the protected area (*Li et al., 2021*).

**Camera trap sampling**

Camera trap detections of terrestrial mammals were collected during the dry season from

November 2020 to April 2021 and November 2021 to April 2022 to avoid the heavy rainy

seasons in the region. The mean trapping efforts were 134 days (89 to 147 days range). We

used Yianws L720 camera traps to conduct the survey. To determine how anthropogenic

factors shape spatiotemporal associations among terrestrial mammals, we set up camera

trapping stations within the nature reserve based on intensity of human activities and distance

to nearest human settlement. We deployed 350 camera stations in the reserve with different

degrees of anthropogenic disturbance, maintaining at least 800 m between them (range 886 to

2233 m, median 1219; *Supplementary file Figure S1*). This distance may not satisfy the

assumption of population closure and there may be some degree of pseudo-replication as

observations of wide-ranging animals may not be independent. For these species, the

occupancy estimate can be thought of as an estimate of the probability that the species used

the area where the camera trapping station was located, rather than true occupancy (*Li et al.,*

*2018*). Our camera trapping stations spanned a gradient of forest areas with varying levels of

human activities and habitat modifications in the surrounding area. We affixed camera traps

to trees between 80 and 100 cm off the ground, and they were not baited. Camera sensitivity

was set to "low" to reduce false detections triggered by nonanimal movements. We set

cameras to take three photos per detection event, with 3 s delay between subsequent

detections. Camera trap photos were later identified to species when possible. We combined

all human presence photos into a single "Human" categorization representing a variety of

human activities detected around a camera station (e.g. resource gathering, livestock grazing,

recreation etc.). All photos of the same species (including humans) at the same camera station

were considered independent detections if separated by at least 1 h (*Li et al., 2021*). Several

camera stations were invalid due to camera malfunctions or camera loss. The total valid

sampling effort was 43,163 camera days from 322 camera stations that operated effectively.

The target species consisted of ground-dwelling mammal species observed in more than 10

camera stations and weighing more than 1 kg. Thus, the occurrence dataset consists of

presence-absence information on 17 mammal species at 322 camera stations.

**Anthropogenic and habitat covariates**

We derived two different types of human impacts: chronic human modification (e.g.,

settlement, transportation night-time lights etc.) and acute direct human presence (occurrence

of people and domestic animals detected by camera traps) to address our hypotheses. We

explored the degree of habitat modification based on the Human modification (HM) map

metric (*Kennedy et al., 2019*). The HM metric provides a cumulative estimate of artificial

modification of terrestrial landscapes based on 13 anthropogenic stressors caused by five

human activities (human settlement, agriculture, transportation, energy production and

electrical infrastructure) at a resolution of 1km (*Kennedy et al., 2019*). The metric is based on

both the intensity and extent of impact of each anthropogenic stressor and ranges from 0 (no

human disturbance) to 1 (highest human disturbance). Our sampling stations represent a

varied gradient of human modification from 0.04 in minimum to 0.28 in maximum, with a

mean value of 0.12 (*sd* = 0.05). We quantified the level of acute direct human presence

around each camera trapping station by calculating the independent detections of

human-related activities (e.g., livestock grazing, forest resource collection and tourism) per

100 camera-trap days. The range of human presence recorded by our camera trapping survey

was 0 to 46.81, with a mean value of 6.42 (*sd* = 8.61). We also chose a set of two

environmental covariates known to impact spatiotemporal distribution of mammals (*Li et al.,*

*2018*) and presumed to affect interspecific associations: (1) elevation, which plays a key role

in shaping spatial distribution of many species in mountain forests (*He et al., 2019*; *Li et al.,*

*2018*), and (2) forest cover, which provides food resources, thermal cover and escape shelter

for animals (*Long et al., 2005*). We derived percent of forest cover for each camera station

based on the 250 m Moderate Resolution Imaging Spectroradiometer (MODIS) imagery

(MOD44B Vegetation Continuous Fields (VCF) yearly product) of the study area for the

period of 2021. For each camera trapping station, we derived human modification and percent

of forest cover with a buffer radius of 500 m. This spatial scale should capture the

environment that influences both resident animals with small home ranges and transient

animals moving through the area. Prior to analysis, we log- transformed [log (x + 0.1)] human

presence data to account for its highly skewed distribution. We also standardized human

modification and other environmental covariates by scaling to have mean of zero and unit

variance.

**Spatial co-occurrence analysis**

We employed a context-dependent joint species distribution model (JSDM; *Tikhonov et al.,*

*2017*) to characterize interspecific spatial associations of sympatric species. JSDMs are able

to separate spatial associations between species into shared environmental preferences and

residual correlations that cannot be explained by the environmental factors (Pollock et al.,

2014). The context-dependent JSDM approach allow residual correlations to vary across the

environment by incorporating species- and site-specific latent variables in the model

(*Tikhonov et al., 2017*). We followed this approach and constructed a context-dependent

model by utilizing a latent variable structure, where the factor loadings are modelled as a

linear regression of covariates, allowing species associations to covary with human

disturbance covariates. For our camera trap detections, we modelled the presence-absences of

species $j$ at camera station $i$ as:

$$y_{ij} \sim Bern(\Psi_{ij})$$

381     with $\Psi_{ij} = \varphi^{-1}(\eta_{ij})$, where $\Psi_{ij}$ is the species-specific occurrence probability for each

382     camera trapping station, and $\varphi^{-1}$ is the inverse of a probit link-function. We modelled $\eta_{ij}$

383     as:

$$\eta_{ij} = \sum_{k=1}^{n_c} x_{ik}\beta_{jk} + \varepsilon_{ij}$$

where $n_c$ denotes the number of fixed covariates (i.e., elevation, percent of forest cover,

human modification, and human presence) plus intercept, $\beta_{jk}$ denotes the effect of

environmental covariate $k$ on species $j$, $x_{ik}$ denotes the measured covariates $k = 1 \ldots n_c$ in the

sampling unit $i$. The intercept of the model is included by setting $x_{i1} = 1$ for all sampling

units, so that the number of measured environmental covariates is $n_c - 1$. The species

associations are modelled through the term $\varepsilon_{ij}$, which is defined by a latent factor model:

$$\varepsilon_{ij} = \sum_{h=1}^{n_f} z_{ih}\,\lambda_{jh}(x_{i.}^*)$$

where $z_{ih}$ denotes the value of latent factor $h = 1 \dots n_f$ at the sampling unit $i$, $\lambda_{jh}(x_{i\cdot}^*)$

denotes the response (factor loading) of species $j$ to latent factor $h$, given a vector of

predictors $x_{i\cdot}^*$. The predictors $x_{i\cdot}^* = (x_{i1}^*, \dots, x_{in_c^*}^*)$ on which the species associations are

assumed to depend can be arbitrary, usually a subset of environmental predictors (Tikhonov

et al., 2017). Here we model the factor loadings of species as a function of the two types of

human disturbances (i.e., human modification and human presence):

$$\lambda_{jh}(x_{i\cdot}^*) = \sum_{k=1}^{n_c^*} x_{ik}^* \lambda_{jhk}$$

where $n_c^*$ denotes the number of covariates assumed to impact residual correlations plus

intercept. The intercept is included to the regression part by setting $x_{i1}^* = 1$ for all sampling

units. In this study, assume $n_f = 3$, as our Deviance information criteria did not improve

notably with the addition of more latent variables. We defined the covariance matrix of

species factor loading as a function of human disturbances as $\varepsilon_{i\cdot} \sim N(0, \Omega(x_{i\cdot}^*))$, where

$\Omega(x_{i\cdot}^*) = \Lambda(x_{i\cdot}^*)\Lambda(x_{i\cdot}^*)^{\mathrm{T}}$, and $\Lambda(x_{i\cdot}^*)$ is the matrix of factor loadings, which depends on the

human disturbances. We then scale this covariance matrix $\Omega$ to interspecific correlation

matrices R by defining $R_{j1j2} = \Omega_{j1j2}/\sqrt{\Omega_{j1j1}\Omega_{j2j2}}$ for each pair of species, which

represents disturbance-dependent associations between species that are not explained by fixed

species-specific effects of environmental predictors. These resulted in values between -1 and

1, with negative values representing negative association between species, and positive values

implying the opposite.

We estimated changes in species associations over continuous gradients of human

modification (range: 0.04 to 0.28) and human presence gradients (range: 0.00 to 46.81). After

fitting the model to data, we used the parameterized model to infer how species associations

depend on human presence and human modification and generated predictions at minimum

(lower), mean (moderate) and maximum (higher) conditions of the two variables separately.

We fitted the model based on a Bayesian approach using the greta R-package (Golding,

2019) as described by *Perrin et al. (2022)*. We specified uninformative normally-distributed

priors for all parameters. We made inference from 3000 samples on three chains after a

burn-in of 2000 samples.

**Temporal co-occurrence analysis**

For temporal co-occurrence analysis, we defined "Low" and "High" categories of human

presence and human modification. We ranked camera trapping stations based on human

presence and human modification separately, and filtered detections from the 25% most-

("High" category) and least- ("Low" category) disturbed camera trap stations and pooled

detections within each category. We used kernel density distributions of animal diel activity

and time between detections of species pairs (time-to-encounter) to compare temporal

associations at lower and higher human disturbances. To examine if changes in species diel

activity patterns were a mechanism behind anthropogenic impacts, we computed the

nocturnal probability and time between consecutive detections of species pairs from lower-

and higher-disturbance habitats. We transformed the detection time stamp to "solar time" to

eliminate the impacts of variation in day length (*Nouvellet et al., 2012*). We retained only

those species that had at least 30 independent detections in the low- and high-disturbance

categories. For nocturnality analysis, we excluded detections within the hour around sunrise

and sunset to avoid the effects of crepuscular activities on nocturnal probability. Thus, we

defined nocturnal records as detections 1 h after sunset up to 1 h before sunrise. We

conducted a binomial t-test to evaluate shifts in species nocturnality in the low- and

high-disturbance categories.

To calculate time between detections of species pairs, we filtered camera stations to only

those that detected at least two species for each disturbance category. We then calculated the

time (in days) between successive detections across species pairs at each camera station. For

analysis, we log transformed the time-to-encounter values to account for their markedly

skewed distribution. We used 10,000 bootstrapped samples to evaluate 95% confidence

interval (CI) shifts in time-to-encounter of species pairs from camera stations in the low- and

high-disturbance categories.

**Acknowledgements**

We thank the Tibetan Autonomous Region Forestry and Grassland Department for their

logistic support and their authorities for permitting the study. We thank Minjing Pu, Kang

Luo, Changzhe Pu and other collaborators that helped collect data. We also thank Cheng

Chen for his assistance with data management. The study was supported by the Second

Tibetan Plateau Scientific Expedition and Research Program (STEP, 2019 QZKK0501), the

National Key R&D Program of China (2022YFC2602500, 2022YFC2601200), the West

Light Foundation of The Chinese Academy of Sciences, the Yunnan Provincial Youth Talent

Support Program (YNWR-QNBJ-2020-127), and the China Biodiversity Monitoring and

Research Network (Sino Bon).

**Author contributions**

Xueyou Li, Conceptualization, Funding acquisition, Data curation, Formal analysis,

Investigation, Visualization, Methodology, Writing - original draft, Writing - review and

editing, Processed and curated camera trap images; Wenqiang Hu, Data curation,

Investigation, Processed and curated camera trap images; William V. Bleisch, Writing -

review and editing; Quan Li, Investigation; Hongjiao Wang, Processed and curated camera

trap images; Zhongzheng Chen, Investigation; Ru Bai, Investigation; Xuelong Jiang,

Conceptualization, Resources, Data curation, Supervision, Project administration.

**Data availability**

All data supporting this study are available within the main manuscript and the supplementary

information published online. Any data not shown are available from the corresponding

author upon request.

**References**

Beaudrot, L, Palmer, MS, Anderson, TM, Packer, C. 2020. Mixed-species groups of Serengeti

grazers: a test of the stress gradient hypothesis. *Ecology* **101:** e03163. DOI:

https://esajournals.onlinelibrary.wiley.com/doi/abs/10.1002/ecy.3163.

Blanchet, FG, Cazelles, K, Gravel, D. 2020. Co-occurrence is not evidence of ecological

interactions. *Ecology Letters* **23:** 1050-1063. DOI:

https://onlinelibrary.wiley.com/doi/abs/10.1111/ele.13525.

Boron V, Deere NJ, Hyde M, Bardales R, Stasiukynas D, Payán E. 2023. Habitat

modification destabilizes spatial associations and persistence of Neotropical carnivores.

*Current Biology* In press. DOI:

https://www.sciencedirect.com/science/article/pii/S0960982223010369.

Burkle, LA, Marlin, JC, Knight, TM. 2013. Plant-Pollinator Interactions over 120 Years: Loss

of Species, Co-Occurrence, and Function. *Science* **339:** 1611-1615. DOI:

https://www.science.org/doi/abs/10.1126/science.1232728.

Chamberlain, SA, Bronstein, JL, Rudgers, JA. 2014. How context dependent are species

interactions? *Ecology Letters* **17:** 881-890. DOI: https://doi.org/10.1111/ele.12279.

Dangles, O, Herrera, M, Carpio, C, Lortie, CJ. 2018. Facilitation costs and benefits function

simultaneously on stress gradients for animals. *Proceedings of the Royal Society B:*

*Biological Sciences* **285**: 20180983. DOI: https://doi.org/10.1098/rspb.2018.0983.

Davis, CL, Rich, Lindsey N, Farris, ZJ, Kelly, MJ, Di Bitetti, MS, Blanco, YD, et al. 2018.

Ecological correlates of the spatial co-occurrence of sympatric mammalian carnivores

worldwide. *Ecology Letters* **21**: 1401-1412. DOI:

https://onlinelibrary.wiley.com/doi/abs/10.1111/ele.13124.

de Jonge, MMJ, Benítez-López, A, Hennekens, S, Santini, L, Huijbregts, MAJ, Schipper, AM.

2021. Conditional love? Co-occurrence patterns of drought-sensitive species in European

grasslands are consistent with the stress-gradient hypothesis. *Global Ecology and*

*Biogeography* **30:** 1609-1620. DOI:

https://onlinelibrary.wiley.com/doi/abs/10.1111/geb.13323.

Deng, L, Chen, D, Deng, L. 2011. Ecological evaluation of Yarlung Zangbo Grand Canyon

National Nature Reserve in Tibet. *Scientia Silvae Sinicae* **47:** 1-6.

Duan, S, Han, F, Li, F, Yang, Z. 2022. Spatial evaluation of the ecological value importance

of national park in Yarlung Tsangpo Grand Canyon. *Journal of Environmental*

*Management* **320:** 115943. DOI:

https://www.sciencedirect.com/science/article/pii/S030147972201516X.

Fidino, M, Simonis, JL, Magle, SB. 2019. A multistate dynamic occupancy model to estimate

local colonization–extinction rates and patterns of co-occurrence between two or more

interacting species. *Methods in Ecology and Evolution* **10:** 233-244. DOI:

https://besjournals.onlinelibrary.wiley.com/doi/abs/10.1111/2041-210X.13117.

Filazzola, A, Brown, C, Dettlaff, MA, Batbaatar, A, Grenke, J, Bao, T, et al. 2020. The

effects of livestock grazing on biodiversity are multi-trophic: a meta-analysis. *Ecology*

*Letters* **23:** 1298-1309. DOI: https://onlinelibrary.wiley.com/doi/abs/10.1111/ele.13527.

García-Navas, V, Sattler, T, Schmid, H, Ozgul, A. 2021. Bird species co-occurrence patterns

in an alpine environment supports the stress-gradient hypothesis. *Oikos* **130:** 1905-1918.

DOI: https://onlinelibrary.wiley.com/doi/abs/10.1111/oik.08588.

Gardner CJ, Bicknell JE, Baldwin-Cantello W, Struebig MJ, Davies ZG. 2019. Quantifying

the impacts of defaunation on natural forest regeneration in a global meta-analysis. *Nature*

*Communications* **10:** 4590. DOI: https://doi.org/10.1038/s41467-019-12539-1.

Gaynor, KM, Brown, JS, Middleton, AD, Power, ME, Brashares, JS. 2019. Landscapes of

Fear: Spatial Patterns of Risk Perception and Response. *Trends in Ecology & Evolution* **34:**

355-368. DOI: https://www.sciencedirect.com/science/article/pii/S0169534719300199.

Gaynor, KM, Hojnowski, CE, Carter, NH, Brashares, JS. 2018. The influence of human

disturbance on wildlife nocturnality. *Science* **360:** 1232-1235. DOI:

https://www.science.org/doi/10.1126/science.aar7121.

Gilbert, NA, Stenglein, JL, Pauli, JN, Zuckerberg, B. 2022. Human disturbance compresses

the spatiotemporal niche. *PNAS* **119:** e2206339119. DOI:

https://www.pnas.org/doi/abs/10.1073/pnas.2206339119.

Golding, N. 2019. greta: simple and scalable statistical modelling in R. *Journal of Open*

*Source Software* **4:** 1601. DOI: https://doi.org/10.21105/joss.01601.

Gorczynski, D, Hsieh, C, Ahumada, J, Akampurira, E, Andrianarisoa, MH, Espinosa, S, et al.

2022. Human density modulates spatial associations among tropical forest terrestrial

mammal species. *Global Change Biology* **28:** 7205-7216. DOI:

https://onlinelibrary.wiley.com/doi/abs/10.1111/gcb.16434.

Harris, NC, Mills, KL, Harissou, Y, Hema, EM, Gnoumou, IT, VanZoeren, J, et al. 2019.

First camera survey in Burkina Faso and Niger reveals human pressures on mammal

communities within the largest protected area complex in West Africa. *Conservation*

*Letters* **12:** e12667. DOI:

https://conbio.onlinelibrary.wiley.com/doi/abs/10.1111/conl.12667.

Hassell, JM, Begon, M, Ward, MJ, Fèvre, EM. 2017. Urbanization and Disease Emergence:

Dynamics at the Wildlife–Livestock–Human Interface. *Trends in Ecology & Evolution* **32:**

55-67. DOI: https://www.sciencedirect.com/science/article/pii/S0169534716301847.

He, J, Zhao, W, Li, A, Wen, F, Yu, D. 2019. The impact of the terrain effect on land surface

temperature variation based on Landsat-8 observations in mountainous areas. *International*

*Journal        of        Remote        Sensing*        **40:**        1808-1827.        DOI:

https://doi.org/10.1080/01431161.2018.1466082.

Jones, BA, Grace, D, Kock, R, Alonso, S, Rushton, J, Said, MY, et al. 2013. Zoonosis

emergence linked to agricultural intensification and environmental change. *PNAS* **110:**

8399-8404. DOI: https://www.pnas.org/doi/abs/10.1073/pnas.1208059110.

Jones, KR, Venter, O, Fuller, RA, Allan, JR, Maxwell, SL, Negret, PJ, et al. 2018. One-third

of global protected land is under intense human pressure. *Science* **360:** 788-791. DOI:

https://science.sciencemag.org/content/sci/360/6390/788.full.pdf.

Karanth, KU, Srivathsa, A, Vasudev, D, Puri, M, Parameshwaran, R, Kumar, NS. 2017.

Spatio-temporal interactions facilitate large carnivore sympatry across a resource gradient.

*Proceedings of the Royal Society B: Biological Sciences* **284:** 20161860. DOI:

https://royalsocietypublishing.org/doi/abs/10.1098/rspb.2016.1860 %X.

Kathleen Lyons, S, Amatangelo, KL, Behrensmeyer, AK, Bercovici, A, Blois, JL, Davis, M,

et al. 2016. Holocene shifts in the assembly of plant and animal communities implicate

human impacts. *Nature* **529:** 80-83. DOI: https://doi.org/10.1038/nature16447.

Keil, P, Wiegand, T, Tóth, AB, McGlinn, DJ, Chase, JM. 2021. Measurement and analysis of

interspecific spatial associations as a facet of biodiversity. *Ecological Monographs* **91:**

e01452. DOI: https://esajournals.onlinelibrary.wiley.com/doi/abs/10.1002/ecm.1452.

Kennedy, CM, Oakleaf, JR, Theobald, DM, Baruch-Mordo, S, Kiesecker, J. 2019. Managing

the middle: A shift in conservation priorities based on the global human modification

gradient. *Global Change Biology* **25:** 811-826. DOI:

https://onlinelibrary.wiley.com/doi/abs/10.1111/gcb.14549.

Kuussaari, M, Bommarco, R, Heikkinen, RK, Helm, A, Krauss, J, Lindborg, R, et al. 2009.

Extinction debt: a challenge for biodiversity conservation. *Trends in Ecology & Evolution*

**24:** 564-571. DOI: https://www.sciencedirect.com/science/article/pii/S0169534709001918.

Lacher, TE, Jr, Davidson, AD, Fleming, TH, Gómez-Ruiz, EP, McCracken, GF, Owen-Smith,

N, et al. 2019. The functional roles of mammals in ecosystems. *Journal of Mammalogy* **100:**

942-964. DOI: https://doi.org/10.1093/jmammal/gyy183.

Lai, X-L, Zhou, W-L, Gao, H-L, Wang, M, Gao, K, Zhang, B-W, et al. 2020. Impact of

sympatric carnivores on den selection of wild giant pandas. *Zoological Research* **41:** 273.

DOI: https://www.zoores.ac.cn/en/article/id/c8757c33-ea3b-4c8d-9d03-8ad4b96e40a6.

Li, X, Bleisch, WV, Jiang, X. 2018. Using large spatial scale camera trap data and

hierarchical occupancy models to evaluate species richness and occupancy of rare and

elusive wildlife communities in southwest China. *Diversity and Distributions* **24:**

1560-1572. DOI: https://onlinelibrary.wiley.com/doi/abs/10.1111/ddi.12792.

Li, X, Hu, W, Bleisch, WV, Li, Q, Wang, H, Lu, W, et al. 2022a. Functional diversity loss

and change in nocturnal behavior of mammals under anthropogenic disturbance.

*Conservation        Biology        **36:**        e13839.        DOI:

https://conbio.onlinelibrary.wiley.com/doi/abs/10.1111/cobi.13839.

Li, X, Hu, W, Bleisch, WV, Li, Q, Wang, H, Ti, B, et al. 2022b. Disproportionate loss of

threatened terrestrial mammals along anthropogenic disturbance gradients. *Science of The*

*Total        Environment*        **850:**        158038.        DOI:

https://www.sciencedirect.com/science/article/pii/S0048969722051373.

Li, XY, Bleisch, WV, Liu, XW, Hu, WQ, Jiang, XL. 2021. Human disturbance and prey

occupancy as predictors of carnivore richness and biomass in a Himalayan hotspot. *Animal*

*Conservation*        **24:**        64-72.        DOI:

https://zslpublications.onlinelibrary.wiley.com/doi/abs/10.1111/acv.12600.

Li XY, Hu WQ, Wang HJ, Jiang XL. 2023. Tiger reappearance in Medog highlights the

conservation values of the region for this apex predator. *Zoological Research*, **44**, 747-749.

https://www.zoores.ac.cn/en/article/id/d938ea54-208f-4d79-b4ec-6dc92b57cb2a

Long, ES, Diefenbach, DR, Rosenberry, CS, Wallingford, BD, Grund, MD. 2005. Forest

cover influences dispersal distance of white-tailed Deer. *Journal of Mammalogy* **86:**

623-629. DOI: https://doi.org/10.1644/1545-1542200586[623:FCIDDO]2.0.CO;2.

Manlick, PJ, Pauli, JN. 2020. Human disturbance increases trophic niche overlap in terrestrial

carnivore        communities.        *PNAS*        **117:**        26842-26848.        DOI:

https://www.pnas.org/doi/abs/10.1073/pnas.2012774117.

Mills, KL, Harris, NC. 2020. Humans disrupt access to prey for large African carnivores.

*eLife* **9:** e60690. DOI: https://doi.org/10.7554/eLife.60690.

Moll, RJ, Killion, AK, Hayward, MW, Montgomery, RA. 2021. A framework for the

Eltonian     niche     of     humans.     *BioScience*     **71:**     928-941.     DOI:

https://doi.org/10.1093/biosci/biab055.

Murphy, A, Diefenbach, DR, Ternent, M, Lovallo, M, Miller, D. 2021. Threading the needle:

How humans influence predator–prey spatiotemporal interactions in a multiple-predator

system.     *Journal     of     Animal     Ecology*     **90:**     2377-2390.     DOI:

https://besjournals.onlinelibrary.wiley.com/doi/abs/10.1111/1365-2656.13548.

Naeem, S, Thompson, LJ, Lawler, SP, Lawton, JH, Woodfin, RM. 1994. Declining

biodiversity can alter the performance of ecosystems. *Nature* **368:** 734-737. DOI:

https://doi.org/10.1038/368734a0.

Nouvellet, P, Rasmussen, GSA, Macdonald, DW, Courchamp, F. 2012. Noisy clocks and

silent sunrises: measurement methods of daily activity pattern. *Journal of Zoology* **286:**

179-184.                                             DOI:

https://zslpublications.onlinelibrary.wiley.com/doi/abs/10.1111/j.1469-998.2011.00864.x.

Palmer, MS, Gaynor, KM, Becker, JA, Abraham, JO, Mumma, MA, Pringle, RM. 2022.

Dynamic landscapes of fear: understanding spatiotemporal risk. *Trends in Ecology &*

*Evolution*                 **37:**                 911-925.                 DOI:

https://www.sciencedirect.com/science/article/pii/S0169534722001409.

Parsons, AW, Kellner, KF, Rota, CT, Schuttler, SG, Millspaugh, JJ, Kays, RW. 2022. The

effect of urbanization on spatiotemporal interactions between gray foxes and coyotes.

*Ecosphere*                 **13:**                 e3993.                 DOI:

https://esajournals.onlinelibrary.wiley.com/doi/abs/10.1002/ecs2.3993.

Pellissier, L, Albouy, C, Bascompte, J, Farwig, N, Graham, C, Loreau, M, et al. 2018.

Comparing species interaction networks along environmental gradients. *Biological Reviews*

**93:** 785-800. DOI: https://doi.org/10.1111/brv.12366.

Penjor, U, Astaras, C, Cushman, SA, Kaszta, Ż, Macdonald, DW. 2022. Contrasting effects of

human settlement on the interaction among sympatric apex carnivores. *Proceedings of the*

*Royal     Society     B:     Biological     Sciences*     **289:**     20212681.     DOI:

https://royalsocietypublishing.org/doi/abs/10.1098/rspb.2021.2681.

Perrin, SW, van der Veen, B, Golding, N, Finstad, AG. 2022. Modelling temperature-driven

changes in species associations across freshwater communities. *Global Change Biology* **28:**

86-97. DOI: https://onlinelibrary.wiley.com/doi/abs/10.1111/gcb.15888.

Poggiato, G, Münkemüller, T, Bystrova, D, Arbel, J, Clark, JS, Thuiller, W. 2021. On the

interpretations of joint modeling in community ecology. *Trends in Ecology & Evolution* **36:**

391-401. DOI: https://www.sciencedirect.com/science/article/pii/S0169534721000045.

Polaina, E, González-Suárez, M, Kuemmerle, T, Kehoe, L, Revilla, E. 2018. From tropical

shelters to temperate defaunation: The relationship between agricultural transition stage

and the distribution of threatened mammals. *Global Ecology and Biogeography* **27:**

647-657. DOI: https://onlinelibrary.wiley.com/doi/abs/10.1111/geb.12725.

Pollock, LJ, Tingley, R, Morris, WK, Golding, N, O'Hara, RB, Parris, KM, et al. 2014.

Understanding co-occurrence by modelling species simultaneously with a Joint Species

Distribution Model JSDM. *Methods in Ecology and Evolution* **5:** 397-406. DOI:

https://besjournals.onlinelibrary.wiley.com/doi/abs/10.1111/2041-210X.12180.

Riggio, J, Baillie, JEM, Brumby, S, Ellis, E, Kennedy, CM, Oakleaf, JR, et al. 2020. Global

human influence maps reveal clear opportunities in conserving Earth's remaining intact

terrestrial ecosystems. *Global Change Biology* **26:** 4344-4356. DOI:

https://onlinelibrary.wiley.com/doi/abs/10.1111/gcb.15109.

Samia, DSM, Nakagawa, S, Nomura, F, Rangel, TF, Blumstein, DT. 2015. Increased

tolerance to humans among disturbed wildlife. *Nature Communications* **6:** 8877. DOI:

https://doi.org/10.1038/ncomms9877.

Sévêque, A, Gentle, LK, López-Bao, JV, Yarnell, RW, Uzal, A. 2020. Human disturbance

has contrasting effects on niche partitioning within carnivore communities. *Biological*

*Reviews* **95:** 1689-1705. DOI: https://onlinelibrary.wiley.com/doi/abs/10.1111/brv.12635.

Sévêque, A, Gentle, LK, Vicente López-Bao, J, Yarnell, RW, Uzal, A. 2022. Impact of

human disturbance on temporal partitioning within carnivore communities. *Mammal*

*Review* **52:** 67-81. DOI: https://doi.org/10.1111/mam.12265.

Smith, JA, Duane, TP, Wilmers, CC. 2019. Moving through the matrix: Promoting

permeability for large carnivores in a human-dominated landscape. *Landscape and Urban*

*Planning* **183:** 50-58. DOI:

https://www.sciencedirect.com/science/article/pii/S0169204618312829.

Smith, JA, Thomas, AC, Levi, T, Wang, Y, Wilmers, CC. 2018. Human activity reduces

niche partitioning among three widespread mesocarnivores. *Oikos* **127:** 890-901. DOI:

https://onlinelibrary.wiley.com/doi/abs/10.1111/oik.04592.

Song, WY, Li, XY, Chen, ZZ, Li, Q, Onditi, KO, He, SW, et al. 2020. Isolated alpine habitats

reveal disparate ecological drivers of taxonomic and functional beta-diversity of small

mammal assemblages. *Zoological Research* **41:** 670-683. DOI:

https://www.zoores.ac.cn/article/doi/10.24272/j.issn.2095-8137.2020.085.

Suraci, JP, Clinchy, M, Zanette, LY, Wilmers, CC. 2019. Fear of humans as apex predators

has landscape-scale impacts from mountain lions to mice. *Ecology Letters* **22:** 1578-1586.

DOI: https://onlinelibrary.wiley.com/doi/abs/10.1111/ele.13344.

Tikhonov, G, Abrego, N, Dunson, D, Ovaskainen, O. 2017. Using joint species distribution

models for evaluating how species-to-species associations depend on the environmental

context. *Methods in Ecology and Evolution* **8:** 443-452. DOI:

https://besjournals.onlinelibrary.wiley.com/doi/abs/10.1111/2041-210X.12723.

Tilman, D. 1994. Competition and Biodiversity in Spatially structured habitats. *Ecology* **75:**

2-16. DOI: https://esajournals.onlinelibrary.wiley.com/doi/abs/10.2307/1939377.

Tilman, D, Clark, M, Williams, DR, Kimmel, K, Polasky, S, Packer, C. 2017. Future threats

to biodiversity and pathways to their prevention. *Nature* **546:** 73-81. DOI:

https://doi.org/10.1038/nature22900.

Tucker, MA, Böhning-Gaese, K, Fagan, WF, Fryxell, JM, Van Moorter, B, Alberts, SC, et al.

2018. Moving in the Anthropocene: Global reductions in terrestrial mammalian movements.

*Science* **359:** 466-469. DOI:

https://science.sciencemag.org/content/sci/359/6374/466.full.pdf.

Tulloch, AIT, Chadès, I, Lindenmayer, DB. 2018. Species co-occurrence analysis predicts

management outcomes for multiple threats. *Nature Ecology & Evolution* **2:** 465-474. DOI:

https://doi.org/10.1038/s41559-017-0457-3.

Van Scoyoc, A, Smith, JA, Gaynor, KM, Barker, K, Brashares, JS. 2023. The influence of

human activity on predator-prey spatiotemporal overlap. *Journal of Animal Ecology* **92**,

1124-1134.                               DOI:

https://besjournals.onlinelibrary.wiley.com/doi/abs/10.1111/1365-2656.13892.

Wicker, LV, Canfield, PJ, Higgins, DP. 2017. Potential pathogens reported in species of the

family Viverridae and their implications for human and animal health. *Zoonoses and Public*

*Health* **64:** 75-93. DOI: https://onlinelibrary.wiley.com/doi/abs/10.1111/zph.12290.

Wilson, MW, Ridlon, AD, Gaynor, KM, Gaines, SD, Stier, AC, Halpern, BS. 2020.

Ecological impacts of human-induced animal behaviour change. *Ecology Letters* **23:**

1522-1536. DOI: https://doi.org/10.1111/ele.13571.

Wisz, MS, Pottier, J, Kissling, WD, Pellissier, L, Lenoir, J, Damgaard, CF, et al. 2013. The

role of biotic interactions in shaping distributions and realised assemblages of species:

implications for species distribution modelling. *Biological Reviews* **88:** 15-30. DOI:

https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1469-185X.2012.00235.x.

**Figure legends**

**Figure 1.** Conceptual framework illustrating the community-level effects of human

disturbance on spatiotemporal associations among mountain forest terrestrial mammal

species.

**Figure 2.** Effects of covariates on the occurrence of terrestrial mammals in the Yarlung

Zangbo Grand Canyon, southeast Tibet. * denotes significant positive and negative effects

based on 95 % CIs.

**Figure 3.** Estimates of associations between 17 terrestrial mammals across camera trapping

stations with different human modification in the Yarlung Zangbo Grand Canyon.

Associations are shown for the region's (a) minimum (Lower), (b) mean (Moderate) and (c)

maximum (Higher) human modifications.

**Figure 4.** Estimates of associations between 17 terrestrial mammals across camera trapping

stations with different human presence in the Yarlung Zangbo Grand Canyon. Associations

are shown for the region's (a) minimum (Lower), (b) mean (Moderate) and (c) maximum

(Higher) human presence.

**Figure 5.** Nocturnal shifts of carnivores (a), ungulates (b) and combination of carnivores and

ungulates (c) in the low- and high-human presence habitats.

**Figure 6.** Density distributions of encounter time between successive detections of species

pairs (in log-transformed days) in low- and high-human modification (a) and human presence

(b) habitats, and differences in time-to-encounter between species pairs in low- and

high-human modification (c) and human presence (d) habitats. The solid vertical lines in (c)

and (d) represent mean differences, and the dashed vertical lines indicate 95% confidence

intervals.

**Co-occurrence** (vertical axis)

**Human disturbance** (horizontal axis)

<blue>Spatial associations

**Hypothesis 1:** Human disturbance would compress space available for animals to share resulting in increased spatial associations between species

**Hypothesis 2:** Human disturbance would suppress diurnal activity in animals resulting in reduced encounter time between species pairs

Pairwise encounter time

Human modification and presence

**Lower**

**Moderate**

**Higher**

**Increasing human modification level**

**Lower**

**Moderate**

**Higher**

**Increasing human presence level**