Human disturbance increases spatiotemporal associations among mountain forest

terrestrial mammal species

Xueyou Li[1], William V. Bleisch[2], Wenqiang Hu[1], Quan Li[1], Hongjiao Wang[1], Zhongzheng

Chen[3], Ru Bai[1], Xuelong Jiang[1]

[1] State Key Laboratory of Genetic Resources and Evolution & Yunnan key laboratory of

biodiversity and ecological conservation of Gaoligong Mountain, Kunming Institute of

Zoology, Chinese Academy of Sciences; [2] China Exploration and Research Society, 2707-08

SouthMark, Wong Chuk Hang, Hong Kong, China; [3] Anhui Provincial Key Laboratory of the

Conservation and Exploitation of Biological Resources, College of Life Sciences, Anhui

Normal University, Wuhu, China

For correspondence: lixueyou@mail.kiz.ac.cn (XL); jiangxl@mail.kiz.ac.cn (XJ)

Abstract Spatial and temporal associations between sympatric species underpin biotic

interactions, structure ecological assemblages, and sustain ecosystem functioning and stability.

However, the resilience of interspecific spatiotemporal associations to human activity remains

poorly understood, particularly in mountain forests where anthropogenic impacts are often

pervasive. Here we applied context-dependent Joint Species Distribution Models to a

systematic camera-trap survey dataset from a global biodiversity hotspot in eastern Himalaya

to understand how prominent human activities in mountain forests influence species

associations within terrestrial mammal communities. We obtained 10,388 independent

detections of 17 focal species (12 carnivores and five ungulates) from 322 stations over 43,163

camera days of effort. We identified a higher incidence of positive associations in habitats with

higher levels of human modification (87 %) and human presence (83 %) compared to those

located in habitats with lower human modification (64 %) and human presence (65 %) levels.

We also detected a significant reduction of pairwise encounter time at increasing levels of

human disturbance, corresponding to more frequent encounters between pairs of species. Our

findings indicate that human activities can push mammals together into more frequent

encounters and associations, which likely influences the coexistence and persistence of wildlife,

with potential far-ranging ecological consequences.

Introduction

Pervasive human activities can disrupt invisible facets of biodiversity such as species

associations, with potential cascading ecosystem effects (Naeem et al., 1994; Parsons et al.,

2022). Human encroachment into natural ecosystems squeezes the spatiotemporal niches of

wildlife species (Gilbert et al., 2022), altering the number and magnitude of associations in a

community (Burkle et al., 2013), and accelerating species decline and loss (Jones et al., 2018).

Humans play a role as 'super predators' in shaping the co-occurrence of other species with

complicated indirect modifications to multiple interactions between organisms (Gilbert et al.,

2022; Moll et al., 2021). Human disturbance, such as chronic landscape modification and acute

direct human presence, may significantly alter spatiotemporal distribution of species and

fundamentally change the way that species interact (Li et al., 2022a; Sévêque et al., 2020). For

example, expanding human footprints have compressed the space and time available for

mammals to share by restricting animal movements (Tucker et al., 2018) and increasing wildlife

nocturnality (Gaynor et al., 2018). As human and animal activities increasingly overlap in time

and space, it is important to assess and quantify the potential for human-induced changes in

species association to ecosystem structure and function (Penjor et al., 2022).

Species are not distributed independently of each other; rather, they co-occur in time and

space and interact (Tilman, 1994; Wisz et al., 2013). The spatiotemporal co-occurrence of

species, termed interspecific associations, provides unique ecological information and has

important consequence for ecosystem integrity (Gorczynski et al., 2022; Keil et al., 2021).

However, segregated species co-occurrence could be generated by processes such as negative

interspecific interaction, distinct environment requirements, and dispersal limitations; similarly,

aggregated species pairs may reflect positive interspecific interaction but could also reflect

shared environmental preferences (Lyons et al., 2016; Song et al., 2020). Although interspecific

co-occurrence or avoidance cannot be used to directly estimate species interactions (Blanchet

et al., 2020), animals cannot interact if their spatiotemporal niches do not overlap (Gilbert et

al., 2022) and strong interactions should be expected to lead to significant associations. Thus,

interspecific associations convey key information about interactions between sympatric species

(Boron et al., 2023).

Disruptions to the spatiotemporal relationships of species can result in serious ecological

consequences including alteration of community structure (Tulloch et al., 2018), upsetting the

competitive balance between species (Boron et al., 2023), increasing disease transmission

(Hassell et al., 2017), and accelerating local extinction (Fidino et al., 2019). This can

significantly distort the distribution of ecological functions that species provide, ultimately

influencing ecosystem dynamics (Gardner et al., 2019). Taken as a whole, interactions between

sympatric species play a fundamental role in community assembly and are intricately related to

ecosystem stability and resilience (Boron et al., 2023). Thus, understanding how interspecific

spatiotemporal associations change across human disturbance gradients provides valuable

insight into the long-term implications of human impacts on ecosystem function and recovery

relevant to biodiversity conservation in the Anthropocene.

Species interactions are known to be context-dependent such that they can vary across

space and time, for example along environmental gradients (Chamberlain et al., 2014; Davis et

al., 2018; Pellissier et al., 2018; Perrin et al., 2022). For example, gradients in stress are

associated with variation in the outcomes of pairwise species interactions (Chamberlain et al.,

2014). A key challenge in community ecology is to identify the conditions under which

negative and positive species interactions are more likely to occur. For example, the

controversial stress-gradient hypothesis predicts that positive interactions should increase as

environmental conditions become more severe. Alpine bird communities have been shown to

have a higher frequency of positive associations in grasslands with low productivity compared

with in forests with high productivity (García-Navas et al., 2021) and savanah ungulates are

more likely to form mixed-species groups in areas where there is predation risk (Beaudrot et

al., 2020). Increasingly, major stresses on wild communities derive from human activity, and

understanding how species interactions vary in response to human disturbance is pivotal in

making robust ecological predictions about biodiversity responses to changing environmental

conditions (Dangles et al., 2018; Perrin et al., 2022). However, a review of the literature

indicates that the impact of human activity on interspecific interactions of terrestrial mammals

remains poorly understood (Boron et al., 2023), particularly in mountain forests where

anthropogenic impacts are often pervasive and increasing.

Medium and large-sized terrestrial mammals are a key components of mountain forest

communities. They play crucial roles in maintaining biodiversity and ecosystem functions

(Lacher et al., 2019), and are especially vulnerable to anthropogenic activities (Li et al., 2022b).

Here, we set out to investigate the spatial and temporal patterns of occurrence and the

interspecific associations within a terrestrial mammalian community along human disturbance

gradients. We executed a systematic camera trapping survey spanning $4,100 \ km^2$ of the Yarlung

Zangbo Grand Canyon National Nature Reserve in the eastern Himalayas at a total of 322

camera trapping stations, obtaining detections of both wildlife and humans. We classified

human disturbances into two groups: human modification (i.e. relating to habitat modifications

by humans) or human presence (i.e. referring to the direct presence of humans) disturbances.

We employed a context-dependent joint species distribution model (JSDM; Tikhonov et al.,

2017) to infer pairwise species associations along human disturbance gradients. We used kernel

density distributions of animal diel activity and time between detections of species pairs (time-

to-encounter) to compare temporal associations at low and high human disturbances. We

consider two hypotheses regarding temporal and spatial interspecific associations. The spatial

compress hypothesis posits that human modification on habitats would suppress space available

for animals to share (Tilman et al., 2017), resulting in increased positive spatial associations

between species. The temporal compress hypothesis postulates that fear of direct human

presence would compress diurnal activity in animals (Gaynor et al., 2018), resulting in reduced

encounter time (or increased encounter rate) between species (Figure 1). Our study incorporates

two different types of human disturbances in the analysis to elucidate the effects of humans on

multidimensional (i.e., space and time) species associations. We then consider implications for

conservation.

Results

Our camera traps obtained 10,388 independent detections of 17 focal species (12 carnivores

and five ungulates) from 322 stations over 43,163 camera days of effort (Table 1). We

documented a number of species and subspecies of conservation concern, including Bengal

tiger *Panthera trigris trigris*, clouded leopard *Neofelis nebulosa* and dhole *Cuon alpinus*. We

also captured 2,224 independent detections of humans during the survey period.

Species-specific response to habitat covariates

Species varied in their responses to habitat covariates. Nine out of 17 species showed strong

positive response to forest cover (Figure 2, Figure 2—source data 1). Human presence had

apparent negative effects on the occurrence of clouded leopard (mean = -0.967, 95% CI = -2.35

to -0.025), takin *Budorcas taxicolor* (mean = -0.449, 95% CI = -0.935 to -0.071) and red goral

Naemorhedus baileyi (mean = -0.806, 95% CI = -0.935 to -0.071), but was apparently positively

associated with the occurrence of wild boar *Sus scrofa* (mean = 0.521, 95% CI = 0.045 to 1.093)

and golden cat *Catopuma temminckii* (mean = 0.427, 95% CI = 0.165 to 0.804; Figure 2—

source data 1). Human modification was apparently negatively associated with the occurrence

of dhole (mean = -0.24, 95% CI = -0.493 to -0.027), red fox *Vulpes vulpes* (mean = -1.588, 95%

CI = -2.656 to -0.769), Asiatic black bear *Ursus thibetanus* (mean = -0.234, 95% CI = -0.42 to

-0.056), red goral (mean = -0.665, 95% CI = -1.129 to -0.238) and Mainland serow *Capricornis*

sumatraensis (mean = -0.477, 95% CI = -0.697 to   -0.26), but was positively apparently

associated with muntjac *Muntjac* spp. (mean = 3.632, 95% CI = 0.946 to 9.999), masked palm

civet *Paguma larvata* (mean = 2.076, 95% CI = 1.26 to 3.145), large Indian civet *Viverra*

zibetha (mean = 0.782, 95% CI = 0.175 to 1.558) and marbled cat *Pardofelis marmorata* (mean

= 0.689, 95% CI = 0.232 to 1.236; Figure 2, Figure 2—source data 1).

Effects of human disturbances on spatial co-occurrence

Out of the 136 estimated pairwise residual correlation coefficients in occupancy, 87 (64 %)

were positive at lower human modifications (Figure 3a). At the moderate and higher

modifications, the species pairs with positive associations increased to 107 (79 %) and 118

(87%) respectively (Figure 3b,c). At lower modifications, correlation coefficients for 18 species

pairs were positive and had a 95 % CI that did not overlap zero (Figure 3—figure supplement

1), and the number increased to 65 in moderate modifications (Figure 3—figure supplement 2)

but dropped to 29 at higher modifications (Figure 3—figure supplement 3).

Along human presence gradients, 88 pairwise residual correlation coefficients (65 %) at

lower human presence habitats were positive or close to neutral (Figure 4a). At the moderate

and higher human presence habitats, the species pairs with positive associations increased to

115 (85 %) and 113 (83%) respectively (Figure 4b,c). The significant positive associations at

low, moderate and higher human presence habitats were 6 (4%, Figure 4—figure supplement

1), 76 (56 %, Figure 4—figure supplement 2), and 44 (32%, Figure 4—figure supplement 3)

respectively.

Effects of human disturbances on temporal co-occurrence

Human presence was associated with significantly increased nocturnality of carnivores (mean

= 0.163, 95% CI = 0.089 to 0.236, Figure 5a), but showed no significant effects on ungulates

(mean = -0.004, 95% CI = -0.031 to 0.023, Figure 5b) and combination of guilds (mean = -

0.001, 95% CI = -0.028 to 0.026; Figure 5c). Human modification had no strong effects on

nocturnality of any guilds and combination of guilds (i.e. 95 % CIs include zero, Figure 5—

figure supplement 1).

Both human modification and human presence were associated with significantly reduced

time between detections of pairs (human modification: mean = -1.07, 95% CI = -1.39 to -0.73;

human presence: mean = -0.69, 95% CI = -0.92 to -0.46; Figure 6).

Discussion

Despite accumulating evidence of widespread impacts of humans on wildlife distribution and

activity patterns, our understanding of how different types of anthropogenic pressures reshape

species associations remains limited (Van Scoyoc et al., 2023). Here, we compared the impacts

of human modification (spatial compress effects) and human presence (temporal compress

effects) on spatiotemporal associations among threatened terrestrial mammals based on

systematic camera-trapping data in an understudied Himalayan landscape. Our results show a

strong influence of humans on species co-occurrence patterns. The overall results of the

spatiotemporal associations across levels of human modification and human presence suggest

that humans are associated with increasing positive spatiotemporal associations among species.

Specifically, we detected a higher incidence of positive associations in habitats with moderate

and higher levels of human modification and human presence compared to those located in

habitats with lower disturbance levels (Figure 3, 4). On the temporal axis, we detected a

significant reduction of pairwise encounter time at increasing levels of human disturbance

(Figure 5). Our results are therefore demonstrate that human disturbance can upset interspecific

associations both on spatial and temporal niche dimensions.

Interspecific spatial associations can arise from species interaction, response to

environmental covariates and common dispersal barriers (Blanchet et al., 2020; de Jonge et al.,

2021; Poggiato et al., 2021). Our context-dependent JSDM, which accounts for similarities and

disparities in species-specific response to habitat covariates considered relevant to medium-

and large-bodied mammal species, detected a higher prevalence of interspecific spatial

associations in human disturbed habitats. Humans can have "bottom-up" impacts on animal

distribution and associations by land use change and habitat modification (Riggio et al., 2020;

Tucker et al., 2018). Human activities can also have top-down impacts on spatiotemporal

associations among animals by directly or indirectly changing the landscape of fear (Palmer et

al., 2022; Suraci et al., 2019). Previous studies suggest that human disturbance may compress

the space and time available for communities to use, resulting in increased frequency of positive

associations (Gilbert et al., 2022; Sévêque et al., 2022). We interpret our results to mean that

anthropogenic presence and disturbance reduce available habitats for wildlife, causing a greater

predominance of positive associations in anthropogenic landscapes. On the one hand, human

presence, such as while grazing livestock and gathering resource, generates landscapes of fear

for wildlife (Gaynor et al., 2019; Palmer et al., 2022; Suraci et al., 2019) and may downgrade

habitat quality by overexploitation (Filazzola et al., 2020). On the other hand, human

modification such as land-use change often constrains the realized niche space of wildlife,

restricting animal movement (Smith et al., 2018; Tucker et al., 2018). Overall, our study

provides one of the first tests of whether positive spatiotemporal associations between terrestrial

mammals increase along gradients of different types of anthropogenic pressures. Our results

add to the growing body of evidence that suggests anthropogenic activities reduce available

niche space for animals, causing observed positive spatial associations among species (Gilbert

et al., 2022; Gorczynski et al., 2022; Murphy et al., 2021).

Identifying thresholds of anthropogenic activity that shift species behavior and co-

occurrence will be key to drawing useful inference from human impact studies and improving

our knowledge on when altered associations may lead to reverberating impacts on ecosystems

(Van Scoyoc et al., 2023; Wilson et al., 2020). We note that the number of species pairs with

significant positive associations rapidly increased from lower to moderate levels of human

disturbance, but dropped from mean to higher level of human disturbance (Figure 3—figure

supplement 1-3 and Figure 4—figure supplement 1-3). Such patterns are consistent across

human modification and human presence levels, indicating a threshold after which the positive

effects of human disturbance on species associations were dampened. Human disturbance is an

important factor in shaping species distributions (Jones et al., 2018; Samia et al., 2015).

Wildlife often has a limited tolerance threshold to human activity (Polaina et al., 2018; Samia

et al., 2015; Smith et al., 2019). When human pressures reach levels that preclude a species

occurring at a site (Polaina et al., 2018), prediction of species associations in an anthropogenic

context may become uninformative. More work needs to be done to further clarify the

mechanisms driving the observed co-occurrence patterns.

Anthropogenic activities can also shift temporal niche of animals (Gaynor et al., 2018; Li

et al., 2022a), possibly altering encounter rates among species and trophic dynamics that

structure communities (Gilbert et al., 2022; Karanth et al., 2017; Mills and Harris, 2020).

Congruent with our hypothesis, we observed significant reduction of pairwise encounter time

at increasing levels of human disturbance, corresponding to more frequent encounters between

the pairs (Gilbert et al., 2022). Although we did not find a significant shift in nocturnality at

community level, we detected a significant shift to nocturnal activity of carnivores, indicating different sensitivities to human presence among the carnivores and ungulates in our study system. Fear of humans may explain the temporal response of wildlife to human presence. Numerous studies have shown that fear of humans as "super predators" can have suppressive effects on widlife activity, which may contribute to increased temporal overlap among speceis (e.g., Sévêque et al., 2022; Suraci et al., 2019).

All in all, our results indicate that both human modification and human presence may rewire species interactions by increasing spatial and temporal co-occurrence. Studies on the impacts of humans on wildlife communities should explicitly account for different types of co-occurring disturbances (Li et al., 2022a). Our camera trapping survey observed that human presence is pervasive even inside this remote protected area in the Tibetan Autonomous Region, indicating that wildlife and their habitats are exposed to frequent human disturbance. Humans exploit resources in protected areas in many ways, including through livestock herding, resource gathering, illegal hunting, and recreation, all of which impact wildlife and their habitats to varying degrees (Harris et al., 2019; Mills and Harris, 2020). Our results demonstrate prevalent disruptions to species co-occurrence patterns from humans. If we are to preserve biodiversity in protected area, we must work to understand the negative effects on multiple facets of biodiversity from co-occurring anthropogenic pressures. Only then can we design effective mitigation measures.

Although interspecific associations should not be directly interpreted as a signal of biotic interactions between pairs of species (Blanchet et al., 2020; Poggiato et al., 2021), it describes

a unique facet of biodiversity, and can provide important insights into the coexistence and

persistence of wildlife as well as ecosystem function in settings with anthropogenic activity

(Keil et al., 2021; Lai et al., 2020; Van Scoyoc et al., 2023). Since interspecific associations

often increases with the ecological similarity of species involved (Gorczynski et al., 2022), the

tendency of many species towards positive associations with increase in anthropogenic

pressures indicates a trend towards homogenization of terrestrial mammal communities. Such

human-mediated changes in species co-occurrence patterns can have serious ecological

consequences at multiple scales (Gilbert et al., 2022; Gorczynski et al., 2022). For individual

species, increased positive associations may affect fitness and population dynamics, increasing

local extinction rates (Kuussaari et al., 2009; Parsons et al., 2022). Increase in spatial

aggregation among functionally similar species may be indicative of a hidden extinction debt

(Kuussaari et al., 2009). At the community level, increased positive associations may depress

co-occurrence network complexity and stability, amplify interactions such as predation, and

simplify communities with similar traits or co-occurrence patterns (Manlick and Pauli, 2020;

Mills and Harris, 2020). Also increased encounter rates between species may expand disease

transmission across communities (Hassell et al., 2017). In addition, wildlife can host a variety

of zoonotic disease (Wicker et al., 2017) and populations unnaturally associated with humans

are more likely to transmit pathogens (Jones et al., 2013). We observed that several species

such as masked palm civet and wild boar are positively associated with humans in our study

area, indicating a substantial overlap between their habitats and human activity. Ongoing

habitat modification, livestock grazing and resource gathering in this region may increase

probability of pathogen exchange.

The effects of human disturbance on species associations might be scale-dependent

(Gilbert et al., 2022). At present, our sampling design only considers the effects of cumulative

human modification and instantaneous local human presence on spatial associations and

temporal encounter time at each camera trapping station. We are not able to compare the effects

of human disturbance at broader scales (e.g., landscape scale) as such an analysis requires

adequate data from hierarchical samplings. Estimating the effects of human disturbance at

different spatial scale on species associations is a promising approach. Thus, we encourage

future work to further clarify the multiscale ecological effects of human disturbance on species

associations.

In conclusion, our study demonstrated that anthropogenic pressures increase

spatiotemporal associations of terrestrial mammals from low to mean levels of human

disturbances, but the frequency of positive spatial associations dropped from mean to higher

level of human disturbances. Such variations in species associations likely influence the

coexistence and persistence of wildlife, with potentially far-ranging ecological consequences.

Because terrestrial mammals like carnivores and ungulates play fundamental roles in regulating

montane forest ecosystems, the prevalent disruptions to their associations may precipitate

biodiversity loss and impair ecosystem function. With increasing human presence and human

modification of areas throughout the world, identifying thresholds of anthropogenic activity

that shift species relationships, limiting human activity and increasing landscape connectivity

across protected areas may be imperative to maintain interspecific spatiotemporal associations

that underpin ecosystem resilience. Moreover, the methods we applied highlight the utility of

camera trapping surveys in studying the spatiotemporal relationships among elusive species in

settings with anthropogenic activities.

Materials and methods

Study area

The study was carried out inside the Yarlung Zangbo Grand Canyon National Nature Reserve

(29°05′–30°02′ N, 94°39′–96°6′ E) in Nyingchi City in the southeast Tibet Autonomous Region

of China (Figure 7). This area is situated within the Eastern Himalaya Biodiversity Hotspot, a

globally important region for biodiversity conservation (Li et al., 2021). The Yarlung Zangbo

Grand Canyon is the deepest in the world, with an elevation drop of more than 7000 m, and has

the reputation of "Gene Bank of Mountain Biological Resources" and "Natural Vegetation

Museum" (Duan et al., 2022). This region is characterized by dramatic vertical zonation of

vegetation. From the valley bottom to the mountain peak, the main vegetation types consist of

low mountain tropical monsoon rain forest, subtropical mountain evergreen broad-leaved forest,

mid-mountain warm coniferous forest, sub-alpine cold coniferous forest, alpine subfrigid shrub

meadow, and periglacial alpine vegetation (Deng et al., 2011). The complete landscapes in the

region harbors endangered species such as Bengal tiger *Panthera tigris tigris* (Li et al., 2023).

Human activities such as decentralized residential settlements and free-ranging livestock

grazing is prevalent in the region, even within the protected area (Li et al., 2021).

Camera trap sampling

Camera trap detections of terrestrial mammals were collected during the dry season between

November 2020 and April 2021 and November 2021 and April 2022 to avoid the heavy rainy

seasons in the region. The mean trapping efforts were 134 days (89 to 147 days range). We

used Yianws L720 camera traps to conduct the survey. To determine how anthropogenic factors

shape spatiotemporal associations among terrestrial mammals, we set up camera trapping

stations within the nature reserve based on intensity of human activities and distance to nearest

human settlement. We deployed 350 camera stations in the reserve with different degrees of

anthropogenic disturbance, maintaining at least 800 m between them (range 886 to 2233 m,

median 1219). This distance may not satisfy the assumption of population closure and there

may be some degree of pseudo-replication as observations of wide-ranging animals may not be

independent. For these species, the occupancy estimate can be thought of as an estimate of the

probability that the species used the area where the camera trapping station was located, rather

than true occupancy (Li et al., 2018). Our camera trapping stations spanned a gradient of forest

areas with varying levels of human activities and habitat modifications in the surrounding area.

We affixed camera traps to trees between 80 and 100 cm off the ground, and they were not

baited. Camera sensitivity was set to "low" to reduce false detections triggered by nonanimal

movements. We set cameras to take three photos per detection event, with 3 s delay between

subsequent detections. Camera trap photos were later identified to species when possible. We

combined all human presence photos into a single "Human" categorization representing a

variety of human activities detected around a camera station (e.g. resource gathering, livestock

grazing, recreation etc.). All photos of the same species (including humans) at the same camera

station were considered independent detections if separated by at least 1 h (Li et al., 2021).

Several camera stations were invalid due to camera malfunctions or lost camera. The total valid

sampling effort was 43,163 camera days from 322 camera stations that operated effectively.

The target species consisted of ground-dwelling mammal species observed in more than 10

camera stations and weighing more than 1 kg. Thus, the occurrence dataset consists of presence-

absence information on 17 mammal species at 322 camera stations.

Anthropogenic and habitat covariates

We derived two different types of human impacts: chronic human modification (e.g., settlement,

transportation night-time lights etc.) and acute direct human presence (occurrence of people

and domestic animals detected by camera traps) to address our hypotheses. We explored the

degree of habitat modification based on the Human modification (HM) map metric (Kennedy

et al., 2019). The HM metric provides a cumulative estimate of artificial modification of

terrestrial landscapes based on 13 anthropogenic stressors caused by five human activities

(human settlement, agriculture, transportation, energy production and electrical infrastructure)

at a resolution of 1km (Kennedy et al., 2019). The metric is based on both the intensity and

extent of impact of each anthropogenic stressor and ranges from 0 (no human disturbance) to 1

(highest human disturbance). Our sampling stations represent a varied gradient of human

modification from 0.04 in minimum to 0.28 in maximum, with a mean value of 0.12 (sd = 0.05).

We quantified the level of acute direct human presence around each camera trapping station by

calculating the independent detections of human-related activities (e.g., livestock grazing,

forest resource collection and tourism) per 100 camera-trap days. The range of human presence

recorded by our camera trapping survey was 0 to 46.81, with a mean value of 6.42 (sd = 8.61).

We also chose a set of two environmental covariates known to impact spatiotemporal

distribution of mammals (Li et al., 2018) and presumed to affect interspecific associations: (1)

elevation, which plays a key role in shaping spatial distribution of many species in mountain

forests (He et al., 2019; Li et al., 2018), and (2) percent of forest cover, which provides food

resources, thermal cover and escape shelter for animals (Long et al., 2005). We derived percent

of forest cover for each camera station based on the 250 m Moderate Resolution Imaging

Spectroradiometer (MODIS) imagery (MOD44B Vegetation Continuous Fields (VCF) yearly

product) of the study area for the period of 2021. For each camera trapping station, we derived

human modification and percent of forest cover with a buffer radius of 500 m. This spatial scale

should capture the environment that influences both resident animals with small home ranges

and transient animals moving through the area. Prior to analysis, we log- transformed [log (x +

0.1)] human presence data to account for its highly skewed distribution. We also standardized

human modification and other environmental covariates by scaling to have mean of zero and

unit variance.

Spatial co-occurrence analysis

We employed a context-dependent joint species distribution model (JSDM; Tikhonov et al.,

2017) to characterize interspecific spatial associations of sympatric species. JSDMs are able to

separate spatial associations between species into shared environmental preferences and

residual correlations that cannot be explained by the environmental factors (Pollock et al., 2014).

The context-dependent JSDM approach allow residual correlations to vary across the

environment by incorporating species- and site-specific latent variables in the model (Tikhonov

et al., 2017). We followed this approach and constructed a context-dependent model by utilizing

a latent variable structure, where the factor loadings are modelled as a linear regression of

covariates, allowing species associations to covary with human disturbance covariates. For our

camera trap detections, we modelled the presence-absences of species j at camera station i as:

$$y_{ij} \sim Bern(\Psi_{ij})$$

with $\Psi_{ij} = \varphi^{-1}(\eta_{ij})$, where Ψ_{ij} is the species-specific occurrence probability for each

camera trapping station, and φ^{-1} is the inverse of a probit link-function. We modelled η_{ij} as:

$$\eta_{ij} = \sum_{k=1}^{n_c} x_{ik}\beta_{jk} + \varepsilon_{ij}$$

where n_c denotes the number of fixed covariates (i.e., elevation, percent of forest cover,

human modification, and human presence) plus intercept, β_{jk} denotes the effect of

environmental covariate k on species j, x_{ik} denotes the measured covariates $k = 1 \ldots n_c$ in the

sampling unit i. The intercept of the model is included by setting $x_{i1} = 1$ for all sampling

units, so that the number of measured environmental covariates is $n_c - 1$. The species

associations are modelled through the term ε_{ij}, which is defined by a latent factor model:

$$\varepsilon_{ij} = \sum_{h=1}^{n_f} z_{ih}\,\lambda_{jh}(x_{i\cdot}^*)$$

where z_{ih} denotes the value of latent factor $h = 1 \ldots n_f$ at the sampling unit i, $\lambda_{jh}(x_{i\cdot}^*)$ denotes

the response (factor loading) of species j to latent factor h, given a vector of predictors $x_{i\cdot}^*$. The

predictors $x_{i\cdot}^* = (x_{i1}^*, \ldots, x_{in_c^*}^*)$ on which the species associations are assumed to depend can be

arbitrary, usually a subset of environmental predictors (Tikhonov et al., 2017). Here we model

the factor loadings of species as a function of the two types of human disturbances (i.e., human

modification and human presence):

$$\lambda_{jh}(x_{i\cdot}^*) = \sum_{k=1}^{n_c^*} x_{ik}^* \lambda_{jhk}$$

where n_c^* denotes the number of covariates assumed to impact residual correlations plus

intercept. The intercept is included to the regression part by setting $x_{i1}^* = 1$ for all sampling

units. In this study, assume $n_f=3$, as our Deviance information criteria did not improve notably

with the addition of more latent variables. We defined the covariance matrix of species factor

loading as a function of human disturbances as $\varepsilon_{i\cdot} \sim N(0, \Omega(x_{i\cdot}^*))$, where $\Omega(x_{i\cdot}^*) =$

$\Lambda(x_{i\cdot}^*)\Lambda(x_{i\cdot}^*)^{\mathrm{T}}$, and $\Lambda(x_{i\cdot}^*)$ is the matrix of factor loadings, which depends on the human

disturbances. We then scale this covariance matrix Ω to interspecific correlation matrices R

by defining $R_{j1j2} = \Omega_{j1j2}/\sqrt{\Omega_{j1j1}\Omega_{j2j2}}$ for each pair of species, which represents

disturbance-dependent associations between species that are not explained by fixed species-

specific effects of environmental predictors. These resulted in values between -1 and 1, with

negative values representing negative association between species, and positive values

implying the opposite.

We estimated changes in species associations over continuous gradients of human

modification (range: 0.04 to 0.28) and human presence gradients (range: 0.00 to 46.81). After

fitting the model to data, we used the parameterized model to infer how species associations

depend on human presence and human modification and generated predictions at minimum

(lower), mean (moderate) and maximum (higher) conditions of the two variables separately.

We fitted the model based on a Bayesian approach using the greta R-package (Golding,

2019) as described by Perrin et al. (2022). We specified uninformative normally-distributed

priors for all parameters. We made inference from 3000 samples on three chains after a burn-

in of 2000 samples.

Temporal co-occurrence analysis

For temporal co-occurrence analysis, we defined "Low" and "High" categories of human

presence and human modification. We ranked camera trapping stations based on human

presence and human modification separately, and filtered detections from the 25% most-

("High" category) and least- ("Low" category) disturbed camera trap stations and pooled

detections within each category. We used kernel density distributions of animal diel activity

and time between detections of species pairs (time-to-encounter) to compare temporal

associations at lower and higher human disturbances. To examine if changes in species diel

activity patterns were a mechanism behind anthropogenic impacts, we computed the nocturnal

probability and time between consecutive detections of species pairs from lower- and higher-

disturbance habitats. We transformed the detection time stamp to "solar time" to eliminate the

impacts of day-length variation in day length (Nouvellet et al., 2012). We retained only those

species that had at least 30 independent detections in the low- and high-disturbance categories.

For nocturnality analysis, we excluded detections within the hour around sunrise and sunset to

avoid the effects of crepuscular activities on nocturnal probability. Thus, we defined nocturnal

records as detections 1 h after sunset up to 1 h before sunrise. We conducted a binomial t-test

to evaluate shifts in species nocturnality in the low- and high-disturbance categories.

To calculate time between detections of species pairs, we filtered camera stations to only

those that detected at least two species for each disturbance category. We then calculated the

time (in days) between successive detections across species pairs at each camera station. For

analysis, we log transformed the time-to-encounter values to account for their markedly skewed

distribution. We used 10,000 bootstrapped samples to evaluate 95% confidence interval (CI)

shifts in time-to-encounter of species pairs from camera stations in the low- and high-

disturbance categories. Results were considered to be significant if the key values fell outside

of the 95% Confidence Interval.

Data availability

The authors declare that the data supporting the findings of this study are available within the

article and its supplementary materials. Occupancy model data (independent detections of

wildlife species and human activities histories and model covariates) are openly available in

Science Data Bank: https://doi.org/10.57760/sciencedb.11804.

Acknowledgements

We thank the Tibetan Autonomous Region Forestry and Grassland Department for their logistic

support and their authorities for permitting the study. We thank Minjing Pu, Kang Luo,

Changzhe Pu and other collaborators that helped collect data. We also thank Cheng Chen for

his assistance with data management.

References

Beaudrot L, Palmer MS, Anderson TM, Packer C. (2020). Mixed-species groups of Serengeti

grazers: a test of the stress gradient hypothesis. *Ecology*, **101**, e03163.

https://esajournals.onlinelibrary.wiley.com/doi/abs/10.1002/ecy.3163

Blanchet FG, Cazelles K, Gravel D. (2020). Co-occurrence is not evidence of ecological

interactions. *Ecology Letters*, **23**, 1050-1063.

https://onlinelibrary.wiley.com/doi/abs/10.1111/ele.13525

Boron V, Deere NJ, Hyde M, Bardales R, Stasiukynas D, Payán E. (2023). Habitat modification

destabilizes spatial associations and persistence of Neotropical carnivores. *Current Biology*

In press. DOI: https://www.sciencedirect.com/science/article/pii/S0960982223010369

Burkle LA, Marlin JC, Knight TM. (2013). Plant-Pollinator Interactions over 120 Years: Loss

of Species, Co-Occurrence, and Function. *Science*, **339**, 1611-1615.

https://doi.org/10.1126/science.1232728

Chamberlain SA, Bronstein JL, Rudgers JA. (2014). How context dependent are species

interactions? *Ecology Letters*, **17**, 881-890. https://doi.org/10.1111/ele.12279

Dangles O, Herrera M, Carpio C, Lortie CJ. (2018). Facilitation costs and benefits function

simultaneously on stress gradients for animals. *Proceedings of the Royal Society B:*

Biological Sciences, **285**, 20180983. https://doi.org/10.1098/rspb.2018.0983

Davis CL, Rich Lindsey N, Farris ZJ, Kelly MJ, Di Bitetti MS, Blanco YD, Albanesi S,

Farhadinia Mohammad S, Gholikhani N, Hamel S, Harmsen BJ, Wultsch C, Kane MD,

Martins Q, Murphy AJ, Steenweg R, Sunarto S, Taktehrani A, Thapa K, Tucker JM,

Whittington J, Widodo FA, Yoccoz NG, Miller DAW. (2018). Ecological correlates of the

spatial co-occurrence of sympatric mammalian carnivores worldwide. *Ecology Letters*, **21**,

1401-1412. https://onlinelibrary.wiley.com/doi/abs/10.1111/ele.13124

de Jonge MMJ, Benítez-López A, Hennekens S, Santini L, Huijbregts MAJ, Schipper AM.

(2021). Conditional love? Co-occurrence patterns of drought-sensitive species in

European grasslands are consistent with the stress-gradient hypothesis. *Global Ecology*

and *Biogeography*, **30**, 1609-1620.

https://onlinelibrary.wiley.com/doi/abs/10.1111/geb.13323

Deng L, Chen D, Deng L. (2011). Ecological evaluation of Yarlung Zangbo Grand Canyon

National Nature Reserve in Tibet. *Scientia Silvae Sinicae*, **47**, 1-6.

Duan S, Han F, Li F, Yang Z. (2022). Spatial evaluation of the ecological value importance of

national park in Yarlung Tsangpo Grand Canyon. *Journal of Environmental Management*,

320, 115943. https://www.sciencedirect.com/science/article/pii/S030147972201516X

Fidino M, Simonis JL, Magle SB. (2019). A multistate dynamic occupancy model to estimate

local colonization–extinction rates and patterns of co-occurrence between two or more

interacting species. *Methods in Ecology and Evolution*, **10**, 233-244.

https://besjournals.onlinelibrary.wiley.com/doi/abs/10.1111/2041-210X.13117

Filazzola A, Brown C, Dettlaff MA, Batbaatar A, Grenke J, Bao T, Peetoom Heida I, Cahill Jr

JF. (2020). The effects of livestock grazing on biodiversity are multi-trophic: a meta-

analysis. *Ecology Letters*, **23**, 1298-1309.

https://onlinelibrary.wiley.com/doi/abs/10.1111/ele.13527

García-Navas V, Sattler T, Schmid H, Ozgul A. (2021). Bird species co-occurrence patterns in

an alpine environment supports the stress-gradient hypothesis. *Oikos*, **130**, 1905-1918.

https://onlinelibrary.wiley.com/doi/abs/10.1111/oik.08588

Gardner CJ, Bicknell JE, Baldwin-Cantello W, Struebig MJ, Davies ZG. (2019). Quantifying

the impacts of defaunation on natural forest regeneration in a global meta-analysis. *Nature*

Communications **10:** 4590. DOI: https://doi.org/10.1038/s41467-019-12539-1

Gaynor KM, Brown JS, Middleton AD, Power ME, Brashares JS. (2019). Landscapes of Fear:

Spatial Patterns of Risk Perception and Response. *Trends in Ecology & Evolution*, **34**,

355-368. https://www.sciencedirect.com/science/article/pii/S0169534719300199

Gaynor KM, Hojnowski CE, Carter NH, Brashares JS. (2018). The influence of human

disturbance on wildlife nocturnality. *Science*, **360**, 1232-1235.

https://www.science.org/doi/10.1126/science.aar7121

Gilbert NA, Stenglein JL, Pauli JN, Zuckerberg B. (2022). Human disturbance compresses the

spatiotemporal niche. *Proceedings of the National Academy of Sciences*, **119**,

e2206339119. https://www.pnas.org/doi/abs/10.1073/pnas.2206339119

Golding N. (2019). greta: simple and scalable statistical modelling in R. *Journal of Open*

Source Software, **4**, 1601. https://doi.org/10.21105/joss.01601

Gorczynski D, Hsieh C, Ahumada J, Akampurira E, Andrianarisoa MH, Espinosa S, Johnson

S, Kayijamahe C, Lima MGM, Mugerwa B, Rovero F, Salvador J, Santos F, Sheil D,

Uzabaho E, Beaudrot L. (2022). Human density modulates spatial associations among

tropical forest terrestrial mammal species. *Global Change Biology*, **28**, 7205-7216.

https://onlinelibrary.wiley.com/doi/abs/10.1111/gcb.16434

Harris NC, Mills KL, Harissou Y, Hema EM, Gnoumou IT, VanZoeren J, Abdel-Nasser YI,

Doamba B. (2019). First camera survey in Burkina Faso and Niger reveals human

pressures on mammal communities within the largest protected area complex in West

Africa.   *Conservation*   *Letters*,   **12**,   e12667.

https://conbio.onlinelibrary.wiley.com/doi/abs/10.1111/conl.12667

Hassell JM, Begon M, Ward MJ, Fèvre EM. (2017). Urbanization and Disease Emergence:

Dynamics at the Wildlife–Livestock–Human Interface. *Trends in Ecology & Evolution*,

32, 55-67. https://www.sciencedirect.com/science/article/pii/S0169534716301847

He J, Zhao W, Li A, Wen F, Yu D. (2019). The impact of the terrain effect on land surface

temperature variation based on Landsat-8 observations in mountainous areas.

International Journal of Remote Sensing, **40**, 1808-1827.

Jones BA, Grace D, Kock R, Alonso S, Rushton J, Said MY, McKeever D, Mutua F, Young J,

McDermott J, Pfeiffer DU. (2013). Zoonosis emergence linked to agricultural

intensification and environmental change. *Proceedings of the National Academy of*

Sciences, **110**, 8399-8404. https://www.pnas.org/doi/abs/10.1073/pnas.1208059110

Jones KR, Venter O, Fuller RA, Allan JR, Maxwell SL, Negret PJ, Watson JEM. (2018). One-

third of global protected land is under intense human pressure. *Science*, **360**, 788-791.

https://www.science.org/doi/abs/10.1126/science.aap9565

Karanth KU, Srivathsa A, Vasudev D, Puri M, Parameshwaran R, Kumar NS. (2017). Spatio-

temporal interactions facilitate large carnivore sympatry across a resource gradient.

Proceedings of the Royal Society B: Biological Sciences, **284**, 20161860.

https://royalsocietypublishing.org/doi/abs/10.1098/rspb.2016.1860 %X

Kathleen Lyons S, Amatangelo KL, Behrensmeyer AK, Bercovici A, Blois JL, Davis M, DiMichele WA, Du A, Eronen JT, Tyler Faith J, Graves GR, Jud N, Labandeira C, Looy CV, McGill B, Miller JH, Patterson D, Pineda-Munoz S, Potts R, Riddle B, Terry R, Tóth A, Ulrich W, Villaseñor A, Wing S, Anderson H, Anderson J, Waller D, Gotelli NJ. (2016). Holocene shifts in the assembly of plant and animal communities implicate human impacts. *Nature*, **529**, 80-83. https://doi.org/10.1038/nature16447

Keil P, Wiegand T, Tóth AB, McGlinn DJ, Chase JM. (2021). Measurement and analysis of interspecific spatial associations as a facet of biodiversity. *Ecological Monographs*, **91**, e01452. https://esajournals.onlinelibrary.wiley.com/doi/abs/10.1002/ecm.1452

Kennedy CM, Oakleaf JR, Theobald DM, Baruch-Mordo S, Kiesecker J. (2019). Managing the middle: A shift in conservation priorities based on the global human modification gradient. *Global Change Biology*, **25**, 811-826. https://onlinelibrary.wiley.com/doi/abs/10.1111/gcb.14549

Kuussaari M, Bommarco R, Heikkinen RK, Helm A, Krauss J, Lindborg R, Öckinger E, Pärtel M, Pino J, Rodà F, Stefanescu C, Teder T, Zobel M, Steffan-Dewenter I. (2009). Extinction debt: a challenge for biodiversity conservation. *Trends in Ecology & Evolution*, **24**, 564-571. https://www.sciencedirect.com/science/article/pii/S0169534709001918

Lacher TE, Jr., Davidson AD, Fleming TH, Gómez-Ruiz EP, McCracken GF, Owen-Smith N, Peres CA, Vander Wall SB. (2019). The functional roles of mammals in ecosystems. *Journal of Mammalogy*, **100**, 942-964. https://doi.org/10.1093/jmammal/gyy183

Lai, XL, Zhou, WL, Gao, HL, Wang, M, Gao, K, Zhang, BW, Wei, FW, Nie, YG. (2020).

Impact of sympatric carnivores on den selection of wild giant pandas. *Zoological. Research*,

41, 273. https://www.zoores.ac.cn/en/article/id/c8757c33-ea3b-4c8d-9d03-8ad4b96e40a6.

Li, X, Bleisch, W., Jiang, X. (2018). Using large spatial scale camera trap data and hierarchical

occupancy models to evaluate species richness and occupancy of rare and elusive wildlife

communities in southwest China. *Diversity and Distributions*, **24**, 1560-1572.

https://onlinelibrary.wiley.com/doi/abs/10.1111/ddi.12792.

Li X, Hu W, Bleisch WV, Li Q, Wang H, Lu W, Sun J, Zhang F, Ti B, Jiang X. (2022a).

Functional diversity loss and change in nocturnal behavior of mammals under

anthropogenic disturbance. *Conservation Biology*, **36**, e13839.

https://conbio.onlinelibrary.wiley.com/doi/abs/10.1111/cobi.13839

Li X, Hu W, Bleisch WV, Li Q, Wang H, Ti B, Qin Z, Sun J, Zhang F, Jiang X. (2022b).

Disproportionate loss of threatened terrestrial mammals along anthropogenic disturbance

gradients. Science of The Total Environment, 850, 158038.

https://www.sciencedirect.com/science/article/pii/S0048969722051373

Li XY, Bleisch WV, Liu XW, Hu WQ, Jiang XL. (2021). Human disturbance and prey

occupancy as predictors of carnivore richness and biomass in a Himalayan hotspot. *Animal*

Conservation, **24**, 64-72.

https://zslpublications.onlinelibrary.wiley.com/doi/abs/10.1111/acv.12600

Li XY, Hu WQ, Wang HJ, Jiang XL. (2023). Tiger reappearance in Medog highlights the

conservation values of the region for this apex predator. *Zoological Research*, **44**, 747-

749. https://www.zoores.ac.cn/en/article/id/d938ea54-208f-4d79-b4ec-6dc92b57cb2a

Long ES, Diefenbach DR, Rosenberry CS, Wallingford BD, Grund MD. (2005). Forest Cover Influences Dispersal Distance of White-Tailed Deer. *Journal of Mammalogy*, **86**, 623-629. https://doi.org/10.1644/1545-1542(2005)86[623:FCIDDO]2.0.CO;2

Manlick PJ, Pauli JN. (2020). Human disturbance increases trophic niche overlap in terrestrial carnivore communities. *Proceedings of the National Academy of Sciences*, **117**, 26842-26848. https://www.pnas.org/doi/abs/10.1073/pnas.2012774117

Mills KL, Harris NC. (2020). Humans disrupt access to prey for large African carnivores. *eLife*, **9**, e60690. https://doi.org/10.7554/eLife.60690

Moll RJ, Killion AK, Hayward MW, Montgomery RA. (2021). A Framework for the Eltonian Niche of Humans. *BioScience*, **71**, 928-941. https://doi.org/10.1093/biosci/biab055

Murphy A, Diefenbach DR, Ternent M, Lovallo M, Miller D. (2021). Threading the needle: How humans influence predator–prey spatiotemporal interactions in a multiple-predator system. *Journal of Animal Ecology*, **90**, 2377-2390. https://besjournals.onlinelibrary.wiley.com/doi/abs/10.1111/1365-2656.13548

Naeem S, Thompson LJ, Lawler SP, Lawton JH, Woodfin RM. (1994). Declining biodiversity can alter the performance of ecosystems. *Nature*, **368**, 734-737. https://doi.org/10.1038/368734a0

Nouvellet P, Rasmussen GSA, Macdonald DW, Courchamp F. (2012). Noisy clocks and silent sunrises: measurement methods of daily activity pattern. *Journal of Zoology*, **286**, 179-184. https://zslpublications.onlinelibrary.wiley.com/doi/abs/10.1111/j.1469-

7998.2011.00864.x

Palmer MS, Gaynor KM, Becker JA, Abraham JO, Mumma MA, Pringle RM. (2022). Dynamic

landscapes of fear: understanding spatiotemporal risk. *Trends in Ecology & Evolution*, **37**,

911-925. https://www.sciencedirect.com/science/article/pii/S0169534722001409

Parsons AW, Kellner KF, Rota CT, Schuttler SG, Millspaugh JJ, Kays RW. (2022). The effect

of urbanization on spatiotemporal interactions between gray foxes and coyotes. *Ecosphere*,

13, e3993. https://esajournals.onlinelibrary.wiley.com/doi/abs/10.1002/ecs2.3993

Pellissier L, Albouy C, Bascompte J, Farwig N, Graham C, Loreau M, Maglianesi MA, Melián

CJ, Pitteloud C, Roslin T, Rohr R, Saavedra S, Thuiller W, Woodward G, Zimmermann

NE, Gravel D. (2018). Comparing species interaction networks along environmental

gradients. *Biological Reviews*, **93**, 785-800. https://doi.org/10.1111/brv.12366

Penjor U, Astaras C, Cushman SA, Kaszta Ż, Macdonald DW. (2022). Contrasting effects of

human settlement on the interaction among sympatric apex carnivores. *Proceedings of the*

Royal Society B: Biological Sciences, **289**, 20212681.

https://royalsocietypublishing.org/doi/abs/10.1098/rspb.2021.2681

Perrin SW, van der Veen B, Golding N, Finstad AG. (2022). Modelling temperature-driven

changes in species associations across freshwater communities. *Global Change Biology*,

28, 86-97. https://onlinelibrary.wiley.com/doi/abs/10.1111/gcb.15888

Poggiato G, Münkemüller T, Bystrova D, Arbel J, Clark JS, Thuiller W. (2021). On the

Interpretations of Joint Modeling in Community Ecology. *Trends in Ecology & Evolution*,

36, 391-401. https://www.sciencedirect.com/science/article/pii/S0169534721000045

Polaina E, González-Suárez M, Kuemmerle T, Kehoe L, Revilla E. (2018). From tropical

shelters to temperate defaunation: The relationship between agricultural transition stage

and the distribution of threatened mammals. *Global Ecology and Biogeography*, **27**, 647-

657. https://onlinelibrary.wiley.com/doi/abs/10.1111/geb.12725

Pollock LJ, Tingley R, Morris WK, Golding N, O'Hara RB, Parris KM, Vesk PA, McCarthy

MA. (2014). Understanding co-occurrence by modelling species simultaneously with a

Joint Species Distribution Model (JSDM). *Methods in Ecology and Evolution*, **5**, 397-406.

https://besjournals.onlinelibrary.wiley.com/doi/abs/10.1111/2041-210X.12180

Riggio J, Baillie JEM, Brumby S, Ellis E, Kennedy CM, Oakleaf JR, Tait A, Tepe T, Theobald

DM, Venter O, Watson JEM, Jacobson AP. (2020). Global human influence maps reveal

clear opportunities in conserving Earth's remaining intact terrestrial ecosystems. *Global*

Change   Biology,   **26**,   4344-4356.

https://onlinelibrary.wiley.com/doi/abs/10.1111/gcb.15109

Samia DSM, Nakagawa S, Nomura F, Rangel TF, Blumstein DT. (2015). Increased tolerance

to   humans   among   disturbed   wildlife.   *Nature   Communications*,   **6**,   8877.

https://doi.org/10.1038/ncomms9877

Sévêque A, Gentle LK, López-Bao JV, Yarnell RW, Uzal A. (2020). Human disturbance has

contrasting effects on niche partitioning within carnivore communities. *Biological*

Reviews, **95**, 1689-1705. https://onlinelibrary.wiley.com/doi/abs/10.1111/brv.12635

Sévêque A, Gentle LK, Vicente López-Bao J, Yarnell RW, Uzal A. (2022). Impact of human

disturbance on temporal partitioning within carnivore communities. *Mammal Review*, **52**,

67-81. https://doi.org/10.1111/mam.12265

Smith JA, Duane TP, Wilmers CC. (2019). Moving through the matrix: Promoting permeability

for large carnivores in a human-dominated landscape. *Landscape and Urban Planning*,

183, 50-58. https://www.sciencedirect.com/science/article/pii/S0169204618312829

Smith JA, Thomas AC, Levi T, Wang Y, Wilmers CC. (2018). Human activity reduces niche

partitioning among three widespread mesocarnivores. *Oikos*, **127**, 890-901.

https://onlinelibrary.wiley.com/doi/abs/10.1111/oik.04592

Song, WY., Li, XY., Chen, ZZ., Li, Q, Onditi, KO., He, SW, Jiang, XL. (2020). Isolated alpine

habitats reveal disparate ecological drivers of taxonomic and functional beta-diversity of

small mammal assemblages. *Zoological Research*, **41**, 670-683.

https://www.zoores.ac.cn/article/doi/10.24272/j.issn.2095-8137.2020.085.

Suraci JP, Clinchy M, Zanette LY, Wilmers CC. (2019). Fear of humans as apex predators has

landscape-scale impacts from mountain lions to mice. *Ecology Letters*, **22**, 1578-1586.

https://onlinelibrary.wiley.com/doi/abs/10.1111/ele.13344

Tikhonov G, Abrego N, Dunson D, Ovaskainen O. (2017). Using joint species distribution

models for evaluating how species-to-species associations depend on the environmental

context. *Methods in Ecology and Evolution*, **8**, 443-452.

https://besjournals.onlinelibrary.wiley.com/doi/abs/10.1111/2041-210X.12723

Tilman D. (1994). Competition and Biodiversity in Spatially Structured Habitats. *Ecology*, **75**,

2-16. https://esajournals.onlinelibrary.wiley.com/doi/abs/10.2307/1939377

Tilman D, Clark M, Williams DR, Kimmel K, Polasky S, Packer C. (2017). Future threats to

biodiversity and pathways to their prevention. *Nature*, **546**, 73-81.

https://doi.org/10.1038/nature22900

Tucker MA, Böhning-Gaese K, Fagan WF, Fryxell JM, Van Moorter B, Alberts SC, Ali AH,

Allen AM, Attias N, Avgar T, Bartlam-Brooks H, Bayarbaatar B, Belant JL, Bertassoni

A, Beyer D, Bidner L, van Beest FM, Blake S, Blaum N, Bracis C, … Mueller T. (2018).

Moving in the Anthropocene: Global reductions in terrestrial mammalian movements.

Science, **359**, 466-469. https://science.sciencemag.org/content/sci/359/6374/466.full.pdf

Tulloch AIT, Chadès I, Lindenmayer DB. (2018). Species co-occurrence analysis predicts

management outcomes for multiple threats. *Nature Ecology & Evolution*, **2**, 465-474.

https://doi.org/10.1038/s41559-017-0457-3

Van Scoyoc A, Smith JA, Gaynor KM, Barker K, Brashares JS. (2023). The influence of human

activity on predator-prey spatiotemporal overlap. *Journal of Animal Ecology*, **92**, 1124-

1134. https://besjournals.onlinelibrary.wiley.com/doi/abs/10.1111/1365-2656.13892

Wicker LV, Canfield PJ, Higgins DP. (2017). Potential Pathogens Reported in Species of the

Family Viverridae and Their Implications for Human and Animal Health. *Zoonoses and*

Public Health, **64**, 75-93. https://onlinelibrary.wiley.com/doi/abs/10.1111/zph.12290

Wilson MW, Ridlon AD, Gaynor KM, Gaines SD, Stier AC, Halpern BS. (2020). Ecological

impacts of human-induced animal behaviour change. *Ecology Letters*, **23**, 1522-1536.

https://doi.org/10.1111/ele.13571

Wisz MS, Pottier J, Kissling WD, Pellissier L, Lenoir J, Damgaard CF, Dormann CF,

Forchhammer MC, Grytnes J-A, Guisan A, Heikkinen RK, Høye TT, Kühn I, Luoto M,

Maiorano L, Nilsson M-C, Normand S, Öckinger E, Schmidt NM, Termansen M,

Timmermann A, Wardle DA, Aastrup P, Svenning J-C. (2013). The role of biotic

interactions in shaping distributions and realised assemblages of species: implications for

species distribution modelling. *Biological Reviews*, **88**, 15-30.

https://doi.org/10.1111/j.1469-185X.2012.00235.x

 Table and table legends

690 **Table 1** Independent detection of ground-dwelling medium- and large-bodied mammal

691 species based on camera trapping survey in the Yarlung Zangbo Grand Canyon,

692 southeast Tibet

Order	Family	Genus	Species	Independent detections	IUCN Category
Cetartiodactyla	Bovidae	*Budorcas*	*Budorcas taxicolor*	92	VU
Cetartiodactyla	Bovidae	*Capricornis*	*Capricornis milneedwardsii*	2992	NT
Carnivora	Felidae	*Catopuma*	*Catopuma temminckii*	232	NT
Carnivora	Canidae	*Cuon*	*Cuon alpinus*	256	EN
Carnivora	Mustelidae	*Martes*	*Martes flavigula*	469	LC
Cetartiodactyla	Cervidae	*Muntiacus*	*Muntiacus muntjak*	4696	LC
Cetartiodactyla	Bovidae	*Naemorhedus*	*Naemorhedus baileyi*	254	VU
Carnivora	Felidae	*Neofelis*	*Neofelis nebulosa*	45	VU
Carnivora	Viverridae	*Paguma*	*Paguma larvata*	223	LC
Carnivora	Felidae	*Panthera*	*Panthera tigris*	26	EN
Carnivora	Felidae	*Pardofelis*	*Pardofelis marmorata*	54	NT
Carnivora	Felidae	*Prionailurus*	*Prionailurus bengalensis*	164	LC
Carnivora	Prionodontidae	*Prionodon*	*Prionodon pardicolor*	28	LC
Cetartiodactyla	Suidae	*Sus*	*Sus scrofa*	269	LC
Carnivora	Ursidae	*Ursus*	*Ursus thibetanus*	463	VU
Carnivora	Viverridae	*Viverra*	*Viverra zibetha*	37	LC
Carnivora	Canidae	*Vulpes*	*Vulpes vulpes*	88	LC

Figure legends

Figure 1 Conceptual framework illustrating the community-level effects of human disturbance

on spatiotemporal associations among mountain forest terrestrial mammal species.

Figure 2 The effects of environmental and anthropogenic variables on terrestrial mammals in

the Yarlung Zangbo Grand Canyon.

Figure 2—source data 1 Standardized beta coefficients, and 95 % credible intervals, for the

influence of anthropogenic and environmental covariates on the probability a species used an

area during our camera-trap survey in Medog region. * denotes significant positive and negative

effects based on 95 % CIs.

Figure 3 Estimates of associations between 17 terrestrial mammals across camera trapping

stations with different human modification in the Yarlung Zangbo Grand Canyon. Associations

are shown for the region's (a) minimum (Lower), (b) mean (Moderate) and (c) maximum

(Higher) human modifications.

Figure 3—figure supplement 1 95 % confidence intervals of residual associations between

species pairs at lower human modification level. * denotes significant associations (i.e., $p <$

0.01).

Figure 3—figure supplement 2 95 % confidence intervals of residual associations between

species pairs at moderate modification level. * denotes significant associations (i.e., $p < 0.01$).

Figure 3—figure supplement 3 95 % confidence intervals of residual associations between

species pairs at higher modification level. * denotes significant associations (i.e., $p < 0.01$).

Figure 4 Estimates of associations between 17 terrestrial mammals across camera trapping

stations with different human presence in the Yarlung Zangbo Grand Canyon. Associations are

shown for the region's (a) minimum (Lower), (b) mean (Moderate) and (c) maximum (Higher)

human presence.

Figure 4—figure supplement 1 95 % confidence intervals of residual associations between

species pairs at lower human presence level. * denotes significant associations (i.e., $p < 0.01$).

Figure 4—figure supplement 2 95 % confidence intervals of residual associations between

species pairs at moderate human presence level. * denotes significant associations (i.e., $p <$

0.01).

Figure 4—figure supplement 3 95 % confidence intervals of residual associations between

species pairs at higher presence level. * denotes significant associations (i.e., $p < 0.01$).

Figure 5 Density distributions of encounter time between successive detections of species pairs

(in log-transformed days) in low- and high-human modification (a) and human presence (b)

habitats, and differences in time-to-encounter between species pairs in low- and high-human

modification (c) and human presence (d) habitats. The solid vertical lines in (c) and (d)

represent mean differences, and the dashed vertical lines indicate 95% confidence intervals.

Figure 5—figure supplement 1 Shifts in nocturnality of carnivores (a), ungulates (b) and

combination of carnivores and ungulates (c) in the lower- and higher-human modification

habitats.

Figure 6 Location of study area in the Yarlung Zangbo Grand Canyon National Nature Reserve

in the southeast of Tibetan Autonomous Region of China.

Co-occurrence

Human disturbance

Spatial associations

Hypothesis 1: Human disturbance would compress space available for animals to share resulting in increased spatial associations between species

Hypothesis 2: Human disturbance would suppress diurnal activity in animals resulting in reduced encounter time between species pairs

Pairwise encounter time

Human modification and presence

(a) Elevation (m) (b) Percent of forest cover (%) (c) Human modified area (0−1)

Beta effect size

Lower

Moderate

Higher

Increasing human modification level

Lower
Moderate
Higher

Increasing human presence level