Genotypic and resistance profile analysis of two oat crown rust differential sets urge coordination
and standardisation.

Duong T. Nguyen[1,2*], Eva C. Henningsen[1,3*], David Lewis[1], Rohit Mago[1], Meredith McNeil[4], Radoslaw
Suchecki[2], Scott Boden[5], Jana Sperschneider[1], Shahryar F. Kianian[6], Peter N. Dodds[1], Melania
Figueroa[1] [†]

[1]Commonwealth Scientific and Industrial Research Organisation, Agriculture and Food, Canberra,
ACT 2601, Australia

[2]Commonwealth Scientific and Industrial Research Organisation, Agriculture and Food, Adelaide, SA
5064, Australia

[3]Research School of Biology, The Australian National University, Canberra, ACT 2601, Australia.

[4]Commonwealth Scientific and Industrial Research Organisation, Agriculture and Food, St Lucia, QLD
4067, Australia

[5]School of Agriculture, Food & Wine, Faculty of Sciences, Waite Research Institute, University of
Adelaide, Adelaide, SA 5064, Australia

[6]US Department of Agriculture-Agricultural Research Service, Cereal Disease Laboratory, St. Paul,
MN, USA

*Authors share first authorship
† Corresponding author: M. Figueroa, melania.figueroa@csiro.au

Keywords: KASP, rust, oat, *Pc91*, resistance, virulence

Funding: Grains Research and Development Corporation (GRDC) and CSIRO, project grant CSP2204-
007RTX. ECH was supported by the ANU University Research Scholarship and ANU-CSIRO Digital
Agriculture PhD Supplementary Scholarship.

Abstract

Puccinia coronata f. sp. *avenae* (*Pca*) is the causal agent of the disease known as crown rust which represents a bottleneck in oat production worldwide. Characterisation of pathogen populations often involves race (pathotype) assignments using differential sets, which are not uniform across countries. This study compared virulence profiles of 25 *Pca* isolates from Australia using two host differential sets, one from Australia and one from the USA. These differential sets were also genotyped using DArT sequencing technology. Phenotypic and genotypic discrepancies were detected on eight out of 29 common lines between the two sets, indicating that pathogen race assignments based on those lines are not comparable. To further investigate molecular markers that could assist in the stacking of rust resistance genes important for Australia, four published *Pc91*-linked markers were validated across the differential sets and then screened across a collection of 150 oat cultivars. Drover, Aladdin, and Volta were identified as putative carriers of the *Pc91* locus. This is the first report of the cultivar Volta as the carrier of *Pc91* and demonstrates the value of implementing molecular markers to characterise materials in breeding pools of oat. Overall, our findings highlight the necessity of examining seed stocks using pedigree and molecular markers to ensure seed uniformity and bring robustness to surveillance methodologies.

Introduction

Oat (*Avena sativa* L.) is a grain crop that is in high demand due to its desirable nutritional properties

for humans and animals, as well as for various commercial applications such as cosmetics and fibre

production (Rasane et al. 2015; Suttie and Reynolds 2004). Australia needs to increase oat production

to meet the growing demands from domestic and international markets, especially in Asia and Africa

(Agriculture and Food 2022; Cowman et al. 2021). Crown rust disease, caused by the rust fungus

Puccinia coronata f. sp. *avenae* (*Pca*), severely impacts oat production worldwide (Nazareno et al.

2018; Simons 1985). Crown rust may cause yields to drop by 10% to 40%, and under optimal

environmental conditions, *Pca* infections can destroy entire fields (Behnken et al. 2011; Carson 2017;

Strunk et al. 2022). In Australia, the disease can result in substantial yield losses of up to 50% in

susceptible varieties (Chambers and Thomas 2020).

Oat crown rust disease is managed via two methods, chemical control and planting genetically

resistant cultivars (Barr 1994). Genetic intervention is the preferred method to prevent rust epidemics

as it minimises adverse effects of agrochemicals to the environment, harmful residues in the crop, and

costs associated with treatments (Figueroa et al. 2020). Genetic resistance manifests either as broad

spectrum (e.g., partial or adult plant resistance) or race- specific (Periyannan et al. 2017). Race-specific

resistance, also known as seedling or all-stage resistance (ASR), is advantageous because it is effective

from seedling growth stages onwards and provides full immunity in some cases, while adult plant

resistance (APR) can be more durable. In the best-case scenario, a combination of both types of

resistance is utilised to achieve the best outcomes (Ellis et al. 2014). The underlying mechanism

controlling race-specific resistance involves the recognition of pathogen-derived proteins, known as

avirulence (*Avr*) effectors, by plant immune receptors encoded by resistance (*R*) genes, which trigger

a plant defence response and stop pathogen infection. However, the ability of pathogens to evolve

variants of *Avr* effectors and avoid plant recognition poses a limitation to this strategy (Figueroa et al.

2020). The evolution of *Pca* can also be driven by sexual processes, which contribute to generating

new virulence traits through the reassortment of virulence alleles (Miller et al. 2020; Nazareno et al.

2018). Common buckthorn (*Rhamnus cathartica*) is the sexual host of *Pca* and the lack of this species

in Australia has led researchers to postulate that *Pca* evolution relies solely on asexual propagation in

this part of the world (Harder and Haber 1992; Simons 1985).

As of today, 92 loci (*Pc1-Pc85, Pc91-Pc96, Pc98*) associated with *Pca* resistance are nominated based

on phenotypic data and limited genetic characterisation (Carson 2017; Nazareno et al. 2018). Only a

few of these have been genetically mapped in oats (Carson 2017; Park et al. 2022a). Furthermore,

none of the *Pc* genes have been cloned so far and genetically linked molecular markers have been

developed for only a handful. This makes it difficult to assess whether the same *R* genes or alleles of

the same *R* gene have been independently identified and redeployed under different names.

Unfortunately, reports from the oat global community indicate that most of these genes are now

ineffective in many regions as *Pca* populations have evolved to overcome their resistance. A potential

strategy to deliver rust-durable resistance is the stacking of race-specific and broad-spectrum

resistance genes (Ellis et al. 2014). Such an approach requires the use of molecular markers to select

for each *R* gene in germplasm and effectively incorporate multiple different sources of resistance into

the same cultivar (Figueroa et al. 2020; Nazareno et al. 2018)

At present, rust surveillance programs conducted by pathologists rely on race assignments

(pathotype) to monitor rust evolution and movement and inform breeding programs and growers

about which oat cultivars are suitable for seasonal planting. Races of rust fungi are commonly defined

by a standard nomenclature system that links virulence phenotypes to race-specific genes postulated

in subsets of crop lines, referred to as differential sets (Chong et al. 2000; Figueroa et al. 2018; Leonard

2016). This strategy, although limited to a small set of markers (*R* genes) has provided important

insights into the fast-evolving nature of crown rust populations regardless of geographic origin (Carson

2011). While several *Pc* genes and oat differential sets are shared among international research

organisations, there is no universal collection due to regional differences in pathogen virulence and

host resistance (Chong et al. 2000). The oat differential set used in Australia includes up to 47 lines,

29 of which overlap with lines included in the USA oat differential set of 40 lines (Nazareno et al. 2018).

The oat lines in the differential set are not isogenic and race assignment reports suggest that several

oat differential lines may carry multiple *Pc* genes (Carson 2017; Park 2013), and some *Pc* genes may

be duplicated within sets (Hewitt et al. 2023; Miller et al. 2020).

Pc91 is one of the most recently described race-specific *Pc* genes and was originally identified in the

highly crown rust-resistant oat line Amagalon, which is derived from a cross between tetraploid *A.*

magna and diploid *A. longiglumus*, with *A. magna* being the resistance donor (Rooney et al. 1994).

McCartney et al. (2011) mapped *Pc91* in a recombinant Inbred line (RIL) population (n=100) genotyped

using Diversity Array Technology (DArT) and found several co-segregating DArT markers.

Subsequently, Gnanesh et al. (2013) used the sequence of DArT marker oPt-0350 (McCartney et al.

2011) to design three co-dominant KASP markers based on three separate SNPs within this sequence

(*oPt-0350-KOM4c2*, *oPt-0350-KOM5c1*, and *oPt-0350-KOM6c2*). These markers accurately

distinguished oat lines with and without *Pc91* resistance locus in 16 North American oat breeding lines,

and oPt-0350-KOM4c2 co-segregated with *Pc91* in three mapping populations and was separated by

1.9cM in a fourth population (Gnanesh et al. 2013). In another study, Klos et al. (2017) mapped *Pc91*

to the linkage group Mrg18 through a Genome-Wide Association Study and identified a *Pc91-*

associated SNP marker, *GMI_ES03_c2277_336*. Mrg18 corresponds to chromosome 1A in the

reference genome assembly *Avena sativa* OT3098 v2 (PepsiCo. 2021), while the DArT markers flanking

Pc91 (McCartney et al. 2011, Gnanesh et al. 2013) were correlated with linkage groups from a Kanota

X Ogle map corresponding to a translocation breakpoint in that population involving then defined

chromosomes 7C and 17 (Tinker et al. 2009), with the latter also corresponding to chromosome 1A in

the OT3098 v2 assembly.

In this study, we conducted a genotypic and phenotypic comparison of the widely utilised oat

differential sets in Australia and the USA, which identified substantial discrepancies between lines

supposed to represent the same differential in each set. This suggests the need for care to be taken in

pathogen surveillance and postulation of *R* genes in breeding lines based on comparisons to the

differential sets. Given that *Pc91* remains effective in providing crown rust resistance in Western

Australia (Henningsen et al. 2023), we also tested the aforementioned *Pc91* molecular markers

(Gnanesh et al. 2013; Klos et al. 2017) on a collection of oat cultivars and identified putative carriers

of *Pc91* as a resource for breeding programs seeking to pursue marker-assisted selection to deliver

durable resistance to crown rust.

Materials and methods

Plant material. Two oat crown rust differential sets commonly used in North America (n= 40) and Australia (n=47), respectively, and a collection of oat accessions and cultivars (n=150) were utilised in this study (**Table S1**). The North American differential set is maintained at the US Department of Agriculture-Agricultural Research Service (USDA-ARS) Cereal Disease Laboratory (St. Paul, MN, USA) (Nazareno et al. 2018), and lines of the Australian oat differential set (Cuddy et al. 2016; Park 2013; Park et al. 2022b) were accessed from the Australian Grains Genebank (AGG), the Department of Agriculture and Fisheries (DAF), QLD, Australia, and an *Avena* seed stock at CSIRO. We also built an oat collection that includes cultivars with diverse pedigree information by accessing AGG, DAF, and CSIRO-*Avena* resources. These oat lines were amplified through a single seed descent process. Oats were planted at a depth of 1 cm in an in-house soil blend and grown in a glasshouse with a daily cycle of 16 hours at 23°C light and 8 hours at 18°C dark.

Rust infection assays. A set of 25 *Pca* isolates from major Australian oat-grown regions in 2022 was used (**Table S2,** Henningsen et al. 2023). Infection methods were adapted from Miller et al. (2020), except that rust spores were mixed with talc powder instead of suspended in oil. *Pca* samples were applied at 10 days after sowing by dusting the inoculum onto plants that were pre-misted with water. Plants were kept at high humidity conditions for two days in a sealed inoculation chamber followed by a further 8 days in a growth cabinet (18-23°C, 16h photoperiod). Infection scores were recorded at 10 dpi using a disease rating scale from Murphy (1935) and Nazareno et al. (2018). The scores were subsequently converted to a 0-9 numeric scale for visualisation in heatmaps as described by Miller et al. (2018, 2020) and Henningsen et al. (2023), using ComplexHeatmap (v2.14.0) in R (4.2.2) (Gu, 2022) and comparison of virulent impact using violin plots (ggplot2 package) and Wilcoxon rank sum tests in R v4.2.2.

Genotyping of oat lines and population structure analysis. DNA was extracted from 3-week-old seedlings as described by Ellis et al. (2005) and quality and concentration were assessed using a NanoDrop™ 8000 Spectrophotometer (NanoDrop Technologies Inc., Santa Clara, CA, United States) and fluorescence assay PicoGreen. DArTSeq genotyping of oat lines was performed by Diversity Arrays Technology Pty Ltd. (https://www.diversityarrays.com/) (Canberra, Australia) using restriction enzymes *PstI* and *HpaII* with markers generated in two formats: SilicoDArT (presence/absence variants) and single-nucleotide polymorphisms (SNPs).

Sequences of DArT markers were searched against the reference genome sequence *Avena sativa* OT3098 v2 (PepsiCo. 2021, https://wheat.pw.usda.gov/jb?data=/ggds/oat-ot3098v2-pepsico) using BLAST (Altschul et al. 1990) with an expected value (E) < 5e-7 and sequence identity of > 70%. Only

markers that matched a genomic location were kept. SilicoDArT and SNP markers were separately

filtered using parameters: missing call ≥ 0.5, minimum allele frequency (MAF) < 0.05, and Pearson's

correlation values r > 0.95 (Gezan et al. 2022). After that, SilicoDArT and SNP markers were combined

for genotypic analysis, and markers with high correlation (Pearson's correlation values r > 0.95) were

removed.

CMplot (Yin et al. 2021) was used to visualise the genome-wide distribution of DArTSeq markers across

oat chromosomes of the OT3098 v2 reference genome. The ASRgenomics package (Gezan et al. 2022)

was used in Rstudio (v2022.12.0) to generate a Kinship matrix (G) from DArTSeq genotypic data for

Principal Component Analysis (PCA). Phylogenetic analysis was performed using the maximum

likelihood criterion. The Python script vcf2phylip.py (Ortiz 2019)

(https://github.com/edgardomortiz/vcf2phylip) was used to convert marker vcf files to PHYLIP format

for phylogenetic analysis in RAxML (v8.2.12) (Parameters as follows: raxmlHPC-PTHREADS-AVX -f a -#

500 -m GTRCAT -p 12345 -x 12345 -s differential_set.min4.phy -n ML-BS500) (Stamatakis 2014). The

output phylogenetic tree was midpoint-rooted and visualised using iTOL (v6.7.2) (Letunic and Bork

2021).

Kompetitive Allele Specific PCR (KASP) assays for *Pc91*. In this study, we used three KASP markers

(*oPt-0350-KOM4c2*, *oPt-0350-KOM5c1*, and *oPt-0350-KOM6c2*, that were reported by Gnanesh et al.

(2013). In addition, we converted the GBS-SNP, *GMI_ES03_c2277_336* (Klos et al. 2017) into KASP

maker by using the flanking sequence of the SNP to design primers through 3CR Bioscience Free Assay

Design Service (https://3crbio.com/). This newly designed KASP marker was named

KASP_GMI_ES03_c2277_336, and the sequence of the KASP primers can be found in **Table S3**. The

physical locations of these markers in the OT3098 v2 genome reference were determined by

completing a BLAST search of the flanking sequence of the marker *GMI_ES03_c2277_336* and the

DArT clone oPt-0350 sequence against the reference genome OT3098 v2. The KASP assays were

performed in the SNPline PCR Genotyping System (LGC, Middlesex, United Kingdom), using a method

previously described by Shi et al. (2023).

Pedigree visualisation of oat lines. Pedigree records were obtained from the "Pedigrees of Oat Lines"

POOL database (Fitzsimmons et al. 1983; Tinker and Deyl 2005) and resolved to Parent 1 and Parent

2 to make them compatible with the Helium pedigree viewing software (Shaw et al. 2014) which was

used to visualise relationships between accessions. Previous postulations of *Pc* genes in oat lines based

on phenotypes (Cuddy et al. 2016; Park 2013) were also imported to Helium to be examined in the

context of pedigree.

Results

Comparison of disease resistance phenotypes in oat lines across two host differential sets.

We compared the resistance profiles of 87 oat accessions (**Table S1**) from the crown rust differential sets from USA and Australia using 25 Australian *Pca* isolates (**Figure 1**). Of the 29 oat accessions that overlap between the two oat differential sets, 13 accessions carrying *Pc* genes (Pc35, Pc38, Pc45, Pc50, Pc52, Pc54, Pc55, Pc56, Pc59, Pc63, Pc64, Pc68, and WIX4361-9) showed strong agreement in infection scores with all 25 *Pca* isolates (**Figure 2**), with only 0-2 isolates detecting a different infection outcome, resistant (infection type 0 to 2) vs susceptible (infection type 3-4). These small differences could be due to differences in visual scoring influenced by inoculum load and other environmental factors (e.g., temperature, humidity). However, we detected more substantial phenotypic discrepancies in other lines that should represent the same genotype in each panel. Among these, seven lines (Pc36, Pc39, Pc48, Pc51, Pc58, Pc62, and Pc71) differed in response to 3-4 *Pca* isolates, while nine lines (H548, Pc14, Pc40, Pc46, Pc57, Pc60, Pc61, Pc67, Pc70) showed differences in more than 4 isolates, with some being almost entirely different (e.g., H548, Pc14, Pc60, Pc70).

As reported earlier (Henningsen et al. 2023), the *Pca* subpopulation from Eastern Australia (QLD, NSW, VIC, and SA) exhibited a higher frequency of virulence phenotypes compared to the *Pca* isolates from Western Australia (WA) towards oat differential lines from both the USA (p = 0.00059) and Australian (p = 0.00088) sets (**Figure S1**). Oat cultivars Barcoo, Saia, and the differential line Pc63 exhibited resistance to all 25 *Pca* isolates, whereas certain oat lines, such as Swan, Ukraine, Marvelous, Santa Fe, Trispernia, Landhafer, Bondvic, Pc45, Pc54, and Pc67 displayed susceptibility to the majority or all isolates. Forty oat lines across both differential sets exhibited resistance to all crown rust isolates from WA but were susceptible to one or more isolates from eastern Australia. These lines were Pc36, Pc38, Pc50, Pc52, Pc55, Pc56, Pc52, Pc58, Pc59, Pc63, Pc68, Pc71, Pc91 (ND894904), Pc94, Pc96, Amagalon, Barcoo, Belle, Bettong, Culgoa, Cleanleaf, Genie, Gwydir, H548, HiFi, Legget, Saia, Stainless, TAM-O-0405, Volta, Warrego, X534, and WIX4361-9. Among these, virulence to Pc91, Pc94, and Pc96, as well as Amagalon (carrying *Pc91*) and HiFi (also carrying *Pc91*) has already appeared in NSW and/or QLD, despite containing recently designated *Pc* genes.

Figure 1: Heatmaps showing the virulence profile of 25 Australian *Pca* isolates. High infection scores are shown in red indicating high virulence (susceptibility) and lower scores (yellow/orange) indicate avirulence (resistance), with yellow being the most avirulent. Heatmap using the **(A)** USA oat differential set and **(B)** Australian oat differential set. Dendrograms reflect hierarchical clustering of columns and isolates with similar virulence patterns.

236 **Figure S1:** Violin plots comparing the pooled virulence scores of 25 *Pca* isolates from the Eastern (red)
237 and Western (yellow) regions of Australia on the Australian and US differential sets. The eastern region
238 includes QLD, NSW, VIC, and SA, and west refers to WA. *P*-values shown are from a two-sided Wilcoxon
239 rank sum test between the virulence scores of the eastern and western isolates on the two differential
240 sets.

Figure 2: Heatmap shows comparison of virulence profiles of 29 shared differential lines in the Australian and USA sets. Colour range indicates infection type of isolates (columns) on the host lines (rows): high virulence (susceptibility) in red and low virulence (resistance) in yellow/orange. Asterisks indicate pairs that were found to be genotypically divergent in Figure 4. Dendrograms reflect the hierarchical clustering of columns and isolates with similar virulence patterns.

Genotypic comparison of oat lines included in two differential sets.

To examine the genetic relatedness of oat lines in the two differential sets, genotypes were defined

using DArTSeq, a genome complexity reduction-based sequencing technology. After applying quality

filtering procedures, a total of 48,111 SilicoDArT markers and 17,223 SNPs were retained. Distribution

of markers along chromosomes of the oat reference genome OT3098 v2 (**Figure 3)** indicated complete

coverage across the genome with the number of SNPs on each chromosome ranging from 1,736

(chromosome 3A) to 4,754 (chromosome 4D). The SilicoDArT and SNP markers were then combined,

and highly correlated markers were removed leaving a set of 55,097 markers that was used to examine

the genetic relationships between oat lines in the differential sets. A principal component analysis

(**Figure S2)** showed one group of oat lines in both the USA and Australian sets were closely related and

different from the remainder of the lines which showed no other obvious groupings. This closely

related group is composed of a set of near-isogenic lines (NILs) derived from backcrosses with the oat

cultivar Pendek for various *Pc* genes (*Pc35*, *Pc38*, *Pc39*, *Pc40*, *Pc45*, *Pc46*, *Pc48*, *Pc50*, *Pc54*, *Pc55*, and

Pc56) (Carson, 2017). The differential lines Pc61 (USA) and Pc70 (USA) appeared closely related to the

Pendek NIL group despite their pedigree not indicating that they are part of NIL-derived sets (Carson

2017). A maximum likelihood (ML) phylogenetic tree (**Figure 4**) supported the close relationships of

the Pendek NIL series including Pc61 (USA) and Pc70 (USA). The differential line Pc70 (USA) showed

an identical resistance profile to Pc39 (USA) for 25 *Pca* isolates (**Figure 2**), suggesting that it may in

fact be a repeat of the Pc39 differential in this study. In addition, two small groups of NILs in either

Fraser (Pc62 and Pc63) or Makuru/SunII (Pc64, Pc67, and Pc68) (Carson 2017) were also closely

genetically related as expected (**Figure 4**).

Importantly, most of the 29 lines common to both Australian and USA sets showed a close

phylogenetic relationship, except for eight lines (H548, Pc14, Pc57, Pc58, Pc59, Pc60, Pc61, Pc70).

Seven of these also exhibited substantial differences in their virulence profiles against the set of 25

Pca isolates (**Figure 2**), while the eighth line (Pc59) appeared to be highly similar for resistance

phenotype in both sets. This suggests that while the lines might not be genotypically identical, they

still carry the same *R* gene.

The USA and Australian versions of the Pc40 and Pc46 lines also differed substantially in resistance

profiles, but both of these differentials are part of the closely related Pendek NIL series which makes

it difficult to determine from the phylogeny whether they carry the same R gene or not.

Figure 3. Genome-wide distribution of DArTSeq markers (SNP and SilicoDArT: 65,334) across oat
chromosomes of *OT3098v2* reference genome. The lengths of chromosomes are on the right of the
coloured bars. The colour scale represents marker numbers within a 1Kb window size.

285   **Figure S2.** Principal component analysis of the oat lines from the Australia and USA differential sets.
286   Dots and triangles indicate oat differential lines representing the USA and Australian sets, respectively.
287   Blue colour indicates lines derived from backcrossing of resistance sources to the oat line Pendek
288   (NILs).

Figure 4. Phylogenetic tree of lines/cultivars present in the USA and Australian oat differential set. The phylogeny was generated based on 55,097 markers (SNPs and SilicoDArT) called against the reference genome *OT3098v2* with support values from 500 bootstraps shown near branches. Oat lines were sourced mostly from AGG in Australia (AUS) and the USDA-ARS Cereal Disease Laboratory (USA). The same font colour indicates lines that should be the same but showed genotypic discrepancies. Colour shading indicates groups of near-isogenic lines (NILs) derived from backcrossing sources or resistance with oat cultivar Fraser (green), Pendek (yellow), Makuru/SunII (blue).

Implementation of diagnostic *Pc91*-linked KASP markers for screening oat collections

To determine the possible presence of the *Pc91* gene in oat collections of importance to breeding programs in Australia, four markers previously associated with *Pc91* (Gnanesh et al. 2013; Klos et al. 2017) were tested in the oat differential lines. The markers *oPt-0350-KOM4c2*, *oPt-0350-KOM5c1*, and *oPt-0350-KOM6c2* are all derived from the DArT marker oPt-0350 (Gnanesh et al. 2013), and were found physically located at 419,792,088 bp; 419,808,321 bp; and 419,808,388 bp, respectively, on chromosome 1A of the reference genome OT3098 v2 PepsiCo, while the SNP marker *GMI_ES03_c2277_336* (Klos et al. 2017) occurs at 387,531,571 bp on the same chromosome. The marker *KASP_GMI_ES03_c2277_336* was designed based on the *GMI_ES03_c2277_336* flanking sequence. Amagalon is the original source line used to define *Pc91* and ND94904 and HiFi were both derived from this line (**Figure 5**) and were previously defined as *Pc91* carriers (Gnanesh et al. 2013; McCartney et al. 2011; Rooney et al. 1994). We therefore considered these lines as positive controls to validate KASP markers of *Pc91* in the oat differential lines.

KASP assay results with the *oPt-0350-KOM4c2*, *oPt-0350-KOM5c1*, and *oPt-0350-KOM6c2* markers correlated completely (**Table S4**) and detected four lines containing the resistance-associated alleles in the differential sets (n=87). These include the three positive control lines HiFi, which is homozygous for the resistant alleles, and Pc91 (ND894904) and Amagalon, which appeared heterozygous for the resistance and susceptibility alleles (**Figure 6** and **Table S4**). McCartney et al. (2011) also found that Amagalon and the Pc91 differential line (ND894904) carried both alleles of several markers in this region and suggested this was due to a tandem duplication. The line Volta-1 also appeared heterozygous for the markers and exhibited an identical rust resistance profile to HiFi, ND894904, and Amagalon (**Figure S3**). The pedigree of Volta (**Figure 5**) includes Amagalon as a progenitor, so it could have inherited *Pc91* from this source, including the duplicated marker. In contrast to previous reports (Gnanesh et al. 2013; McCartney et al. 2011), the oat cultivars Stainless, which was derived from a cross of ND931475/AC Assiniboia//HiFi (Mitchell Fetch et al. 2011), was not positive for resistance-associated alleles of all tested markers (**Table S4**) and showed different resistance profiles in comparison to other known *Pc91* carriers (HiFi, ND894904, and Amagalon) (**Figure S3**). Moreover, the cultivar Stainless shows significant divergence in the phylogenetic tree from its parental line HiFi (**Figure 4**). This discrepancy raises the possibility that the current Stainless in the USA differential set differs from the source material used in the studies by Gnanesh et al. (2013), and McCartney et al. (2011).

The marker *KASP_GMI_ES03_c2277_336* detected six differential lines as positive for the resistance-

associated allele, ND94904, HiFi, Cleanleaf, Pc58 (Aus), Warrego, and Leggett, but missed Volta-1 and

Amagalon. No pedigree relationship was evident between Cleanleaf, Pc58 (Aus), Warrego, Leggett,

and the known *Pc91* carriers (ND94904, Amagalon, and HiFi) (**Figure 5** and

https://oat.triticeaetoolbox.org/). These cultivars were previously postulated to carry other *Pc* genes

(Cleanleaf: *Pc38+ Pc39+ Pc52;* Pc58_Aus: *Pc58;* Warrego: *Pc61;* Leggett: *Pc68 + Pc94* by Chong et al.

2000; Mitchell Fetch et al. 2007; Park 2013), and all displayed distinct rust resistance profiles from the

known *Pc91* carriers (**Figure S3**). Klos et al. 2017, previously scored the *GMI_ES03_c2277_336* SNP

marker on 31 lines of the US differential set, with identical results on these lines to those obtained

here using *KASP_GMI_ES03_c2277_336* marker (i.e., only Pc91 was positive).

The *Pc91* markers were further used to screen a collection of 150 oat cultivars (**Table S1**) mostly

sourced from AGG with a few exceptions. The oat cultivars Aladdin, Volta, Genie, and Wizard were

sourced from both AGG and a DAF breeder, representing different generations of selection. For

instance, Aladdin-1 is a single seed stock derived from an AGG seed source representing an F_6-F_7

generation of this line, while Aladdin-2 is a single seed stock derived from a breeder's seed source

representing an F_{15+} generation. Similarly, Volta-1 (in the Australian differential set above) and Volta-

2 (in the oat cultivar set) are separate single seed stocks derived from AGG (F_6-F_7) and Volta-3 is the

single seed stock derived from a Breeder's seed source (F_{15+}).

KASP assays with *oPt-0350-KOM4c2*, *oPt-0350-KOM5c1*, and *oPt-0350-KOM6c2* detected only three

lines (out of 150) containing the resistance-associated alleles: Drover, Aladdin-1, and Volta-2, with the

former homozygous, and the latter two appearing heterozygous (**Figure 6** and **Table S4**). The pedigree

visualisation confirmed that these lines have Amagalon or HiFi as parental lines (**Figure 5**). The oat

cultivar Drover showed the same rust resistance profile as all the known *Pc91* carriers (HiFi, Amagalon,

and ND94904) (**Figure S3**). However, the Aladdin-2 and Volta-3 lines sourced from the breeder were

homozygous for the susceptible genotype, indicating genetic differences between the sources. This

was supported by a comparison of the resistance profiles of these lines (**Figure S3**). Volta-1 (in the

differential set) showed the same resistance profile as Amagalon, Pc91, and HiFi, in contrast, Volta-3

showed a different profile that was identical to Pc50 (USA and Australian sets), while Volta-2 showed

a profile consistent with containing both *Pc91* and *Pc50*. Likewise, the oat cultivars Aladdin-1 and

Aladdin-2 showed different resistance profiles, with the former similar to Amagalon, HiFi, and

ND94904, but with resistance to one additional isolate (**Figure S3**). Five of our lines (HiFi, AC Assiniboia,

Leggett, AC Ronald, and CDC Dancer) were among 16 North American oat breeding lines analysed by

Gnanesh et al. (2013) with these markers and gave identical results.

The marker *KASP_GMI_ES03_c2277_336* detected the resistance allele in 27 of the 150 cultivars

(**Table S4**). Most of these had never been postulated to carry the *Pc91* locus and only Drover, Aladdin

(selections 1 and 2), Volta, and two other oat cultivars, Nugene and Yallara, contained known *Pc91*

sources in their pedigree (**Figure 5**, **Figure S4**, and https://oat.triticeaetoolbox.org/). Like the oPt-0350

markers, *KASP_GMI_ES03_c2277_336* also detected differences between the Volta selections,

although in this case the resistance-associated allele was detected in Volta-3 but not Volta-2.

Figure 5. Pedigree of *Pc91* carrier, Amagalon. A simplified pedigree of the cultivar Amagalon was modified from a Helium output (Shaw et al. 2014). Pedigrees were obtained from the "Pedigrees of Oat Lines" POOL database (Tinker and Deyl 2005) and Fitzsimmons et al. (1983). Red line indicates maternal parent and blue indicates paternal parent. Green circles indicate lines included in the Differential Sets and oat germplasm collection and tested with *Pc91*-linked markers, with lines positive for these markers indicated by "+". ND894904, Amagalon, and HiFi were previously defined as *Pc91* carriers (Gnanesh et al. 2013).

Figure 6. Genotype cluster plots from KASP assays for *Pc91*-linked marker *oPt-0350-KOM4c2* (Gnanesh
et al. 2013). **(A)** Differential Sets, n=88. **(B)** Oat Collection, n =150. The markers *oPt-0350-KOM5c1*,
and *oPt-0350-KOM6c2* displayed a similar pattern as *oPt-0350-KOM4c2*. Blue and red dots show
homozygous genotypes for the resistant (*CC*) or susceptible (*TT*) phenotypes, respectively; green dots
show heterozygous genotypes (*CT*) for disease resistance; black dots represent non-template control.
X- and Y-axes show relative amplification units (ΔRn) for fluorescein amidite (FAM) and Hexachloro-
fluorescein (HEX) signals. Aladdin-1, Volta-1 and Volta-2 were derived from AGG seed source (F_6-F_7),
while Aladdin-2 and Volta-3 were derived from a Breeder's seed source (F_{15+}); DNA from oat lines
Amagalon, HiFi, and Pc50_Aus from the differential sets were used as experimental controls in the
KASP assays of the oat collection.

Figure S3: Heatmap shows virulence profiles of 25 isolates on *Pc91* carrier and non-carrier oat
genotypes. Colour range indicates the infection type of isolate on the host: resistant (0 to 2) to
susceptible (3 to 4). Oat lines in blue font are positive for *Pc91* marker *oPt-0350-KOM4c2*. Postulated
Pc genes based on KASP result, virulence profiles, and previous reports are shown on the right.
Aladdin-1, Volta-1 and Volta-2 were derived from AGG seed source (F_6-F_7), while Aladdin-2 and Volta-
3 were derived from a Breeder's seed source (F_{15+}).

Figure S4. Pedigrees involving the oat differential lines **(A)** Pc50 and **(B)** Pc68. Red lines indicate the
maternal parent connection and blue indicates the paternal parent. The coloured circles are lines
postulated to carry *Pc* genes based on previous historic reports (Brown: *Pc50*, Green: *Pc91*, Blue:
Pc68).

Discussion

Pathogen surveillance programs are key to inform breeders about virulence traits and guide

development of disease resistant cultivars. Differential sets to categorise pathogen collections and

detect changes in virulence are crucial resources for this purpose. In this study, we compared the

performance of two separate oat differential sets used in Australia and the USA. By using phylogenetic

and principal component analyses based on DArTSeq genotypic data, we identified substantial genetic

differences between oat differential lines in the two sets that are supposed to be identical (e.g. H548,

Pc14, Pc57, Pc58, Pc59, Pc60, Pc61, Pc70). The differences between these lines were supported by

disease resistance phenotypic data obtained after infection with a set of 25 *Pca* isolates, except in the

case of Pc59. Thus, these differential lines cannot be considered equivalent when comparing pathogen

race information internationally. These discrepancies may be explained by seed stock heterogeneity

and the outcrossing nature of oat (Coffman and Wiebe 1930; Derick 1933; Fatunla and Frey 1980;

Harrington 1932; Murray et al. 2002), which should be carefully considered when undertaking seed

amplification. Given the need for standard global surveys of *Pca* population for effective pathogen

movement monitoring, we recommend revisiting the core differential set lines and setting a plan for

seed distribution that ensures uniformity. Genotypic data of the differential lines used in this study

provide a starting point for researchers to take a similar approach and check for inconsistencies.

The gene *Pc91* remains effective against *Pca* in Western Australia (Cuddy et al. 2016; Park 2013,

Henningsen et al. 2023). To prolong its effective lifespan, it is critical to combine *Pc91* with other

effective *Pc* genes through strategic breeding efforts. Thus, we tested previously published *Pc91*-

associated markers (Gnanesh et al. 2013; Klos et al. 2017) for future use in national breeding

programs. In our assays, KASP markers derived from DArTSeq marker oPt-0350 were highly specific to

lines known or postulated to contain *Pc91*, detecting resistance-associated alleles in the known

carriers Amagalon, Pc91 (ND894904) and HiFi, as well as in Drover and some accessions of Volta and

Aladdin, which all have *Pc91* carriers in their pedigree and show similar resistance profiles to the

known *Pc91* lines. Drover and Aladdin are grazing oat cultivars and have been previously postulated

to carry *Pc91* based on phenotypic data derived from a pathogen differential set (Cuddy et al. 2016;

Park 2013). Interestingly, Volta and Aladdin have the duplicate alleles of the oPt-0350 markers from

Amagalon, while Drover has the single copy allele of HiFi, which is consistent with the pedigree

relationships among these lines (**Figure 5**). The marker *KASP_GMI_ES03_c2277_336*, derived from

SNP marker *GMI_ES03_c2277_336*, located approximately 32.2 Mb upstream from oPt-0350 showed

lower specificity, with the resistance-associated allele being detected in 6 of the differential lines and

27 of the oat cultivars, most of which do not contain known *Pc91* sources in their pedigree, but not in

the *Pc91* source line Amagalon. This suggests that this marker is less tightly linked to the *Pc91* locus,

and the allele association has been broken by recombination in some lines. Based on these results,

the pPt-0350 markers appear highly diagnostic for *Pc91*.

Importantly, the cultivar Stainless, which originated from the cross ND931475/AC Assiniboia//HiFi

(Mitchell Fetch et al. 2011), was not positive for all four *Pc91* tested markers in our differential set, as

the other carriers (ND894904, Amagalon, and HiFi) and was phylogenetically distinct from the parental

line HiFi (**Figure 4**). Previously, Stainless was defined as a *Pc91* carrier by Gnaneshs et al. (2013) and

McCartney et al. (2011). However, it exhibited different resistance patterns to *Pc91* carriers

(ND894904, Amagalon, and HiFi) in our study and in the studies by Miller et al. (2020) and Hewitt et

al. (2023). These discrepancies suggest that the Stainless line included in these studies either differs

from the ones used by Gnanesh et al. (2013) and McCartney et al. (2011) or has undergone genetic

segregation at the *Pc91* locus after distribution.

Furthermore, we found some discrepancies in genotype and phenotype for different seed stocks of

Volta and Aladdin. Two seed stocks of the cultivar Volta independently amplified by single seed

descent from the same AGG entry (Volta-1 and Volta-2) were positive for *Pc91*, while a third seed

stock derived from material donated by a breeder (Volta-3) was not. While Volta-1 showed a similar

rust resistance profile to ND894904, Amagalon, and HiFi, Volta-3 shared the same resistance profile

as the oat differential line Pc50 and Volta-2 showed a profile that could be explained by the presence

of both *Pc91* and *Pc50*. These results again suggest heterogeneous seed stocks and/or changes as the

cultivar was advanced to the F_{15+} generation by the breeder since being deposited to AGG at the F_{6-7}

generation stage (B. Winter, personal communication). Notably, previous reports based on virulence

phenotyping and race assignment proposed that Volta carried *Pc50* and *Pc68* (Park 2013). However,

pedigree information on the origin of Volta shows ancestors including both *Pc50* and *Pc91* donors, but

not *Pc68* (**Figure S4**). Thus, it is likely that Volta material has segregated for *Pc50* and *Pc91*, although

it is possible that other genotypes have also been present in other stocks used by Park (2013). The

cultivar Aladdin had been proposed to contain *Pc50* and *Pc91* (Park 2013) and our results confirm the

presence of *Pc91* in Aladdin-1 but not Aladdin-2, which contains an unknown *Pc* gene whose resistance

profile is not consistent with *Pc50*. Again, it is possible that other stocks may have included different

genotypes, and these discrepancies interfere with the interpretation of infection data on these lines.

The cultivars Yallara and Nugene, whose pedigree is related to *Pc91* donors, are positive for the

resistance allele of the marker *KASP_GMI_ES03_c2277_336*. However, pPt-0350 markers argue

against the presence of *Pc91* in these cultivars. Given the physical distance between the pPt-0350

markers and *KASP_GMI_ES03_c2277_336,* it is possible that the association between the sequences

recognised by these sets of markers has also been broken by recombination. These cultivars have been

postulated to carry *Pc38* and *Pc48*, respectively, based on resistance phenotyping with an array of *Pca*

isolates (Cuddy et al. 2016; Park 2013).

In conclusion, the rapid host adaptive evolution displayed by *Pca*, challenges the durability of resistant

cultivars once deployed in the field. Therefore, strategic plans to stack genes must be supported by

the development and adoption of molecular markers. This study also reveals the problems in tracking

and maintaining the resistance sources. It is crucial to select and preserve the purity of resistance

sources, with the most effective method also being the use of molecular markers. Standardising oat

crown rust differential lines through collaborative efforts within the oat community is urged for

effective global pathogen monitoring. Ultimately, a systematic approach that incorporates the use of

molecular markers, pedigree analysis, and pathology assays as illustrated in this study is key to

increasing our understanding of the oat crown rust pathosystem.

Funding

This work was jointly funded by the Grains Research and Development Corporation (GRDC) and CSIRO, project grant CSP2204-007RTX.

Acknowledgments

We acknowledge the Australian Grains Genebank, Bruce Winter at the Department of Agriculture and Fisheries (Queensland), and the GRDC-funded oat phenology project team (grant CSP2007) for contributing oat germplasm. We also thank Ben Trevaskis (CSIRO) for helping with pedigree information of several oat cultivars. We acknowledge Liza Apps for technical support at the CSIRO Black Mountain Laboratories Quarantine facility.

Conflict of interest

The authors declare that this study received funding from the GRDC. The funder was not involved in the study design, collection, analysis interpretation of data, the writing of this article, or the decision to submit it for publication.

Author contributions

This study was planned and designed by MF and PND. SFK contributed the US oat differential set. ECH, DL, and MF manage collections of rust isolates, virulence assessments, and seed amplification. SB provided DNA material for a subset of oat lines. DTN performed the genotypic analysis. RM assisted with marker design. MM, RS, and JS supported bioinformatics. DTN, ECH, and MF wrote the first draft of the manuscript. All authors contributed to the interpretation of results, reviewed the manuscript, and approved the final version.

Literature Cited

Agriculture and Food. 2022. *Oats*. https://www.agric.wa.gov.au/hay-production/western-australian-oat-industry

Altschul, S.F., Gish, W., Miller, W., Myers, E.W., Lipman, D.J. 1990. "Basic local alignment search tool." **J. Mol. Biol.** 215:403-410. https://doi.org/10.1016/S0022-2836(05)80360-2

Barr, A. 1994. Strategies for the control of the foliar dieases of oats in South Australia [**University of Adelaide**]. https://digital.library.adelaide.edu.au/dspace/bitstream/2440/21584/2/02whole.pdf

Behnken, L. M., Breitenbach, F. R., Miller, R. P., & Beyer, B. 2011. Impact of foliar fungicide to control crown rust in oats in 2009, 2010, and 2011. https://practicalfarmers.org/wp-content/uploads/previous/2016/05/2011-Crown-Rust-in-Oats-1.pdf

Carson, M. 2011. Virulence in oat crown rust (*Puccinia coronata* f. sp. *avenae*) in the United States from 2006 through 2009. **Plant Dis.** *95*(12): 1528-1534. https://doi.org/10.1094/PDIS-09-10-0639

Carson, M. 2017. *Oat crown rust*. Cereal Disease Lab, Agricultural Research Service, **USDA.** https://www.ars.usda.gov/midwest-area/stpaul/cereal-disease-lab/docs/cereal-rusts/oat-crown-rust/

Chambers, K., & Thomas, G. 2020. Plant diseases impacting oaten hay production in Australia - a review. Department of Primary Industries and Regional Development **(DPIRD).** https://www.agric.wa.gov.au/oats/2020-plant-diseases-impacting-oaten-hay-production-australia

Chong, J., Leonard, K. J., & Salmeron, J. J. 2000. A North American system of nomenclature for *Puccinia coronata* f. sp. *avenae*. **Plant Dis.** *84*(5): 580-585. https://doi.org/10.1094/PDIS.2000.84.5.580

Coffman, F. A., & Wiebe, G. A. 1930. Unusual crossing in oats at Aberdeen, Idaho. **Agron. J.** 22(3), 245-250. https://doi.org/10.2134/agronj1930.00021962002200030005x

Cowman, S., Cox, B., Yamamoto, M., & Kingwell, R. 2021. Opportunities and risks for the Australian oats industry. https://www.aegic.org.au/wp-content/uploads/2021/09/AEGIC-Opportunities-and-risks-for-the-Australian-oat-industry-2021.pdf

Cuddy, W., Park, R., Bariana, H., Bansal, U., Singh, D., Roake, J., & Platz, G. 2016. Cereal rust report 2016. Plant Breeding Institute, **University of Sydney**.

Ellis, M. H., Rebetzke, G. J., Azanza, F., Richards, R. A., & Spielmeyer, W. 2005. Molecular mapping of gibberellin-responsive dwarfing genes in bread wheat. **Theor. Appl. Genet.** *111*(3): 423-430. https://doi.org/10.1007/s00122-005-2008-6

Ellis, J.G., Lagudah, E.S., Spielmeyer, W., Dodds, P.N. 2014. The past, present, and future of breeding for rust resistance in wheat. **Front. Plant Sci.**, 5: 641

Fatunla, T., & Frey, K. J. 1980. Analysis of genetic changes in radiated and non-radiated bulk oat (Avena sativa L.) populations. **Theor. Appl. Genet.** 56(5): 199-202. https://doi.org/10.1007/BF00295449

Figueroa, M., Dodds, P. N., & Henningsen, E. C. 2020. Evolution of virulence in rust fungi - multiple solutions to one problem. **Curr. Opin. Plant Biol.** *56*: 20-27. https://doi.org/10.1016/j.pbi.2020.02.007

Figueroa, M., Hammond-Kosack, K. E., & Solomon, P. S. 2018. A review of wheat diseases-a field
perspective. **Mol. Plant Pathol.** *19*(6): 1523-1536. https://doi.org/10.1111/mpp.12618

Fitzsimmons, R. W., Roberts, G. L., Wrigley, C. W. 1983. Australian oat varieties (Melbourne: CSIRO
Publishing). https://doi.org/10.1071/9780643105447

Gezan, S. A., de Oliveira Giovanni Galli, A. A., Murray, D., Oliveira, de, Gezan salvadorgezan, S. A.,
Amanda de Oliveira, vsnicouk A., & Galli, G. 2022. ASRgenomics: An R package with complementary
genomic functions. Version 1.1.0. **VSN International.** https://vsni.co.uk/free-software/asrgenomics

Gnanesh, B. N., Mitchell Fetch, J., Menzies, J. G., Beattie, A. D., Eckstein, P. E., & McCartney, C. A.
2013. Chromosome location and allele-specific PCR markers for marker-assisted selection of the oat
crown rust resistance gene *Pc91*. **Mol. Breeding.** *32*(3): 679-686. https://doi.org/10.1007/s11032-
013-9900-6

Gu, Z. 2022. Complex heatmap visualization. **IMeta.** 1(3). https://doi.org/10.1002/imt2.43

Harder, D. E., Haber, S., Marshall, H. G., & Sorrels, M. E. 1992. Oat science and technology. **Agron.**
Monogr. *33*: 307-425.

Harrington, J. B. 1932. Natural crossing in wheat, oats and barley at Saskatoon, Saskatchewan. **Sci.**
Agr. 12: 470–483.

Hewitt, T., Henningsen, E. C., Pereira, D., McElroy, K., Nazareno, E. S., Dugyala, S., Nguyen-Phuc, H.,
Li, F., Miller, M. E., Visser, B., Pretorius, Z., Boshoff, W., Sperschneider, J., Stukenbrock, E., Kianian, S.
F., Dodds, P. N., and Figueroa, M. 2023. Genome-enabled analysis of population dynamics and
virulence associated loci in the oat crown rust fungus *Puccinia coronata* f. sp*. avenae*. **bioRxiv**
2023.09.18.557855. https://biorxiv.org/cgi/content/short/2023.09.18.557855v1

Henningsen, E. C., Lewis, D., Nguyen, D. T., Sperschneider, J., Kianian, S., Stone, E., Dodds, P. N., and
Figueroa, M. 2023. Virulence patterns of oat crown rust in Australia - season 2022. **bioRxiv**
2023.09.24.558528. https://doi.org/10.1101/2023.09.24.558528

Klos, K. E., Yimer, B. A., Babiker, E. M., Beattie, A. D., Bonman, J. M., Carson, M. L., Chong, J.,
Harrison, S. A., Ibrahim, A. M. H., Kolb, F. L., McCartney, C. A., McMullen, M., Fetch, J. M.,
Mohammadi, M., Murphy, J. P., & Tinker, N. A. 2017. Genome-wide association mapping of crown
rust resistance in oat elite germplasm. **Plant Genome.** *10*(2).
https://doi.org/10.3835/plantgenome2016.10.0107

Leonard, K. J. 2016. Development of differential varieties - a review. Agricultural Research Service
(ARS) - U.S. Department of Agriculture's. https://www.ars.usda.gov/midwest-area/stpaul/cereal-
disease-lab/docs/cereal-rusts/development-of-differential-varieties-a-review/

Letunic, I., & Bork, P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree
display and annotation. **Nucleic Acids Res.** *49*(W1): W293-W296.
https://doi.org/10.1093/nar/gkab301

McCartney, C.A., Stonehouse, R.G., Rossnagel, B.G. et al. Mapping of the oat crown rust resistance
gene *Pc91*. 2011. **Theor. Appl. Genet.** 122, 317-325. https://doi.org/10.1007/s00122-010-1448-9

Miller, M. E., Nazareno, E. S., Rottschaefer, S. M., Riddle, J., Dos Santos Pereira, D., Li, F., Nguyen-
Phuc, H., Henningsen, E. C., Persoons, A., Saunders, D. G. O., Stukenbrock, E., Dodds, P. N., Kianian,
S. F., & Figueroa, M. 2020. Increased virulence of *Puccinia coronata* f. sp. *avenae* populations
through allele frequency changes at multiple putative *Avr* loci. **PLOS Genet**. *16*(12), e1009291.
https://doi.org/10.1371/journal.pgen.1009291

Mitchell Fetch, J. W., Duguid, S. D., Brown, P. D., Chong, J., Fetch, Jr., T. G., Haber, S. M., Menzies, J.
G., Ames, N., Noll, J., Aung, T. and Stadnyk, K. D. 2007. Leggett oat. **Can. J. Plant Sci**. 87: 509-512.

Mitchell Fetch, J. W., Brown, P. D., AmesN., Chong, J., Fetch, T. G.Jr., Haber, S. M., Menzies, J. G.,
Tekauz, A., Townley-Smith, T. F., and Stadnyk, K. D. 2011. Stainless oat. **Can. J. Plant Sci.** 91(2): 357-
361. https://doi.org/10.4141/CJPS10159

Murray, B., Morrison, I., & Friesen, L. 2002. Pollen-mediated gene flow in wild oat. **Weed Sci**. 50:
321-325.

Murphy, H. C. 1935. Physiologic specialization in *Puccinia coronata avenae.* **Phytopath.** 17: 783-790.
(29)

Nazareno, E. S., Li, F., Smith, M., Park, R. F., Kianian, S. F., & Figueroa, M. 2018. *Puccinia coronata* f.
sp. *avenae*: a threat to global oat production. **Mol. Plant Pathol.** *19*(5): 1047-1060.
https://doi.org/10.1111/mpp.12608

Ortiz E. 2019. vcf2phylip v2.0: convert a VCF matrix into several matrix formats for phylogenetic
analysis. (v2.0). Zenodo. https://doi.org/10.5281/zenodo.2540861

Park, R. 2013. New oat crown rust pathotype with virulence for *Pc91. Cereal Rust Report 2013.* Plant
Breeding Institute, University of Sydney.

Park, R. F., Boshoff, W. H. P., Cabral, A. L., Chong, J., Martinelli, J. A., McMullen, M. S., Fetch, J. W.
M., Paczos-Grzęda, E., Prats, E., Roake, J., Sowa, S., Ziems, L., & Singh, D. 2022a. Breeding oat for
resistance to the crown rust pathogen *Puccinia coronata* f. sp. *avenae*: achievements and prospects.
Theor. Appl. Genet. *135*(11): 3709-3734. https://doi.org/10.1007/s00122-022-04121-z

Park, R., Chhetri, M., Singh, D., & Ding, Y. 2022b. Cereal Rust Report 2022. Plant Breeding Institute,
University of Sydney.

Periyannan, S., Milne, R. J., Figueroa, M., Lagudah, E. S., & Dodds, P. N. 2017. An overview of genetic
rust resistance: From broad to specific mechanisms. **PLOS Pathog.** *13*(7), e1006380.
https://doi.org/10.1371/journal.ppat.1006380

PepsiCo. 2021. *Avena sativa* - OT3098 v2. **GrainGenes: a database for Triticeae and Avena**.
https://wheat.pw.usda.gov/jb?data=/ggds/oat-ot3098v2-pepsico

Rasane, P., Jha, A., Sabikhi, L., Kumar, A., & Unnikrishnan, V. S. 2015. Nutritional advantages of oats
and opportunities for its processing as value added foods - a review. **J. Food Sci. Tech.** *52*(2): 662-
675. https://doi.org/10.1007/s13197-013-1072-1

Rooney, W. L., Rines, H. W., & Phillips, R. L. 1994. Identification of RFLP Markers Linked to Crown
Rust Resistance Genes *Pc* 91 and *Pc* 92 in Oat. **Crop Sci.** *34*(4): 940-944.
https://doi.org/10.2135/cropsci1994.0011183X003400040019x

Shaw, P. D., Graham, M., Kennedy, J., Milne, I., & Marshall, D. F. 2014. Helium: visualization of large
scale plant pedigrees. **BMC Bioinformatics.** *15*(1), 259. https://doi.org/10.1186/1471-2105-15-259

Shi, P., Shen, X., Chen, J., Zhang, Y., Cao, L., Pang, B., Liu, L., Zhang, M., Hoffmann, A. A., & Wei, S.
2023. KASP genotyping and semi-quantitation of G275E mutation in the α6 subunit of *Thrips palmi*
nAChR gene conferring spinetoram resistance. **Pest Manag. Sci.** 79(5): 1777-1782.
https://doi.org/10.1002/ps.7353

Simons, M. D. 1985. Crown rust. Pages 132-172 in: **The Cereal Rusts Vol II: Diseases, distribution,**
epidemiology and control. A. P. Roelfs and W. R. Bushnell eds., Academic Press, Orlando, FL.
https://doi.org/10.1016/B978-0-12-148402-6.50013-4

Stamatakis, A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. **Bioinformatics.** 30: 1312-1313, https://doi.org/10.1093/bioinformatics/btu033

Strunk, C., Kleinjan, J., & Caffe, M. 2022. Crown Rust of Oats. **SDSU Extension.**
https://extension.sdstate.edu/crown-rust-oats-0

Suttie, J. M., & Reynolds, S. G. 2004. Fodder 0ats: A world review (Vol. 33). **FAO** plant production and
protection series.

Tinker, N. A., Deyl, J. K. 2005. A curated internet database of oat pedigrees. **Crop Sci.** 45: 2269-2272.
https://doi.org/10.2135/cropsci2004.0687

Tinker, N.A., Kilian, A., Wight, C.P. et al. 2009. New DArT markers for oat provide enhanced map
coverage and global germplasm characterization. **BMC Genomics** 10, 39.
https://doi.org/10.1186/1471-2164-10-39

Yin, L., Zhang, H., Tang, Z., Xu, J., Yin, D., Zhang, Z., Yuan, X., Zhu, M., Zhao, S., Li, X., Liu, X. 2021.
rMVP: A memory-efficient, visualization-enhanced, and parallel-accelerated tool for genome-wide
association study. **Genomics Proteomics Bioinformatics**. 19(4): 619-628.
https://doi.org/10.1016/j.gpb.2020.10.007