Title: Thermal environment and ecological interactions modulate the importance of evolution in

response to warming

Running Head:

Cara A. Faillace[1,*], Soraya Álvarez-Codesal, Alexandre Garreau, Elvire Bestion, and José M. Montoya

Theoretical and Experimental Ecology Station

French National Centre of Scientific Research (CNRS)

2 Route du CNRS

09200 Moulis, France

[1]**Current institution**: Department of Biological Sciences, University of Pittsburgh, Pittsburgh, PA, USA

***Correspond to**: c.faillace@gmail.com

Abstract

Understanding the interaction between evolutionary history, the abiotic environment, and

biotic interactions is critical for a more nuanced understanding of how communities respond to

anthropogenic stressors. We leveraged a long term experiment manipulating temperature in

mesocosms containing communities of phytoplankton and zooplankton to examine how evolution in

response to long-term community warming could affect consumer-resource dynamics at different

temperatures. We showed that both the current thermal environment, as well as the presence of

interspecific interactions can affect the importance of prior evolution in response to warming, and that

this evolution also influences the outcomes of current ecological dynamics. For each consumer–resource

pair, the effects of evolution were temperature-dependent, but both the effects and the temperature

dependence itself additionally depended upon the identity of evolving species in each pair. The

evolution that we observed in one resource species, *Chlamydomonas reinhardtii*, appeared to confer to

it a double advantage; it was more fit across all temperatures, while simultaneously reducing the

success of the consumer, but with the key result that fitness gains were largest at intermediate

temperatures when the consumer was present. Thus we found support for the "hotter is better"

hypothesis for this resource species, but especially at moderate or intermediate temperatures, and that

hotter was worse for the consumer. In the second species pair, patterns were more complex, but warm-

origin populations of both the resource and the consumer generally failed to show improved fitness.

Overall, our results show that evolution altered resource and consumer fitness, but that these effects

were dependent the current combination of abiotic and biotic conditions.

Key words: consumer-resource interactions, eco-evolutionary dynamics, global climate warming,

predator-prey interactions, temperature dependence, thermal adaptation

Introduction

Global climate change has emerged as a critical challenge to species and ecological communities (Parmesan 2006, Radchuk et al. 2019). Multiple mechanisms by which species and communities respond to increasing temperatures (e.g., range shifts, plastic changes to behavior, phenology, etc.) interact to determine the degree to which species and communities are affected by climate change (O'Connor et al. 2012, Gaitán-Espitia and Hobday 2021). Because temperature represents a strong selective agent for populations, evolution may constitute one important mechanism by which species respond to warming, especially for dispersal-limited populations or species with rapid generation times (O'Donnell et al. 2018). Evolution in response to warming has been experimentally documented in a variety of organisms, including for instance, green algae (Padfield et al. 2016, Schaum et al. 2017), arthropods (Gilchrist and Huey 1999, Geerts et al. 2015, Hangartner and Hoffmann 2016), and larger organisms, such as lizards (Logan et al. 2014). While less frequently studied, evolution in natural populations in response to warming has been observed as well (Higgins et al. 2014, Geerts et al. 2015, Johansson et al. 2016, Johansson and Laurila 2017, Diamond et al. 2017). Because evolution in response to abiotic factors, like climate warming, is frequently explored in single populations, the implications of evolution in communities responding to habitat warming remain understudied. Nonetheless, given that many of the species most likely to evolve in response to warming are critical for ecosystem functioning and services (*e.g.,* water quality, carbon sequestration)(O'Donnell et al. 2018), we can expect that evolutionary responses to warming will have important consequences for both ecological and human systems (Rudman et al. 2017).

In general, intuition suggests that when evolution improves thermal performance at warm temperatures, populations or species should be more successful in warmer environmental conditions (Stoks et al. 2014) relative to colder conditions, although multiple hypotheses exist for how warming

evolution may alter the populations' thermal response curves [e.g., "hotter is better", "biochemical

adaptation", and "partial compensation" (Liu et al. 2022)]. Evidence from natural populations, however,

demonstrates that the performance of populations that evolve in response to warming can be variable.

In some cases responses match theoretical expectations of shifts in thermal tolerance towards warmer

temperatures (Diamond et al. 2017), while in other cases they exhibit unexpected results, which can also

depend upon the life history parameter measured (Johansson et al. 2016). Such unexpected results may,

in part, be explained by life history tradeoffs that can emerge between coping with acute thermal stress

and chronic exposure (Rezende et al. 2014). For instance, evolution may favor increased survival under

chronic warm temperatures at the expense of critical thermal maximum temperature during acute

thermal stress (Johansson and Laurila 2017). Thus, both the type of thermal stress experienced by

populations (i.e., increasing mean temperature vs. frequent acute fluctuations), as well as current

thermal environment (e.g., temperature and duration of thermal experiments) may be important for

interpreting empirical patterns and understanding the implications of evolution that occurs in response

to warming. Physiological responses to temperature (e.g., changes to metabolism) provide an additional

clue to explain why variable responses may be observed in populations evolving to warming. In

particular, many traits exhibit unimodal responses to temperature, such that effects of increased

temperature can appear vastly different depending upon where in the temperature response curve one

examines the response. Thus an additional wrinkle is introduced by the choices of the experimenter.

The presence of additional interacting species may introduce further factors influencing

evolutionary outcomes. The effects of evolution can depend in important ways on the current ecological

conditions in populations experiencing warming. When species are embedded in complex communities,

the opportunities for additional costs, constraints, or benefits of evolution can emerge. For instance,

there can be trait tradeoffs or novel selective pressures present that arise out of additional ecological

(or evolutionary) changes that occur as communities respond to warming. Such trait tradeoffs can drive

differences in evolutionary responses compared to evolution in single populations (Van Doorslaer et al.

2009a, 2010). Selection that occurs within an ecological community is therefore likely to qualitatively

change predictions about thermal adaptation derived from single populations (Angilletta Jr. et al. 2006),

such that individual species' responses to temperature can be insufficient to explain changes to

community composition (Tabi et al. 2020). For example, previous research has demonstrated that

locally adapted populations of consumer and resource species from different latitudes have contrasting

eco-evolutionary dynamics at two temperatures (De Block et al. 2013).

In consequence, we expect that when shifts in thermal performance occur in traits that affect

how species interact, such as resource acquisition or metabolism, it could affect overall fitness of

populations that are simultaneously involved in interspecific interactions and responding to warming.

Specifically, 1) different life history traits within a species can have different thermal dependencies, 2)

consumer and resource species have documented differences in these thermal dependencies, and 3)

evolutionary potential can differ across trophic levels. Therefore, it follows that evolution in response to

warming has the potential to alter consumer-resource dynamics, while also being reciprocally influenced

by them, and that the effects of evolution may additionally depend upon current thermal environment.

If this prediction is correct, we expect more complex outcomes when considering consumer-resource

interactions where changes in thermal performance for both the consumer and the resource are

simultaneously possible. For example, one possibility that emerges is cryptic eco-evolutionary dynamics

where evolution in both species yields a "moving target" that mimics purely ecological dynamics. In this

case, evolution may alter the thermal performance of the interacting species in similar ways, such that

the outcomes of the interaction remain apparently unchanged, but are actually being maintained by the

evolutionary response. A second possibility is that the effects of the interaction on species' fitness may

depend upon the current thermal environment. This might occur if, for example, their thermal

tolerances evolve to different degrees or if only one interacting species evolves. The variety of possible

responses become complex, but studying evolutionary responses in this complexity is necessary to

better understand how evolution can alter the responses of populations and communities to habitat

warming. Additionally, although responses are sometimes examined at multiple temperatures, most

studies are typically of short duration and do not consider the effects of evolution at longer timescales

to quantify changes to fitness. Thus, considering evolutionary responses in more complex communities

acknowledges importance of community ecology in driving evolutionary responses to anthropogenic

environmental change.

To examine both the influence of ecological interactions on evolutionary responses to warming

as well as understanding the long-term effects of evolution on the ecology of interacting species, we

used populations of two resource species (i.e., freshwater green algae, *Chlamydomonas reinhardtii* and

Desmodesmus sp.) and a consumer species (i.e., *Daphnia pulex*) differing in thermal origin. We leveraged

a long-term experiment of manipulated warming in mesocosm plankton communities to understand the

consequences for consumer-resource interactions of evolution in response to long-term warming in

communities. We hypothesized that the fitness effects of this evolution would be dependent upon

current thermal environment and the presence of ecological interactions. Specifically, we asked, 1)

whether biotic interactions affect the importance of evolutionary history across a range of

temperatures, and 2) whether temperature affects the importance of evolutionary history.

128   **Methods**

129   *Source and maintenance of organisms prior to experimentation*

Organisms originated from a long-running mesocosm experiment in Dorset, England. This

mesocosm experiment has been the subject of multiple previous publications (Yvon-Durocher et al.

2015, Schaum et al. 2017). Briefly, at the time of sampling in 2019, open-air experimental ponds of 1 m^3

water of mixed freshwater phytoplankton and zooplankton had been continuously maintained since

2005 (i.e., approximately 14 years). Half of the mesocosms were kept at ambient temperature

conditions, while half experienced a consistent temperature increase of +4 °C above ambient

temperatures (n = 8 for both at the time of our sampling). Temperature in all ponds fluctuated

seasonally. Mesocosm communities initially contained the same mixture of phytoplankton and

zooplankton, but previous work has demonstrated significant changes to community structure in the

warmed mesocosms, as well as evolution in thermal tolerance in warmed populations of at least one

phytoplankton species, *Chlamydomonas reinhardtii* (Yvon-Durocher et al. 2015, Schaum et al. 2017)(Fig.

1).

We sampled two phytoplankton resource species, the algae *C. reinhardtii* and *Desmodesmus* sp.

from each treatment in January 2019, as well as a consumer species, *Daphnia pulex* in September 2019

(for further details see Álvarez-Codesal et al. 2023). The discrepancy in dates occurred because we were

unsuccessful in isolation of consumer populations on the first attempt. Sorting flow cytometry isolated

single algal cells and PacBio Sequel 2 long-read sequencing using full-length 18S fusion primers (forward:

NSF4/18 = CTGGTTGATYCTGCCAGT; reverse: EukR = TGATCCTTCTGCAGGTTCACCTAC) confirmed species

identity and genotypic differences between ambient-origin and warm-origin populations of the algae

(Comeau et al. 2017). We identified one unique clone for each of the two species, *C. reinhardtii* and

Desmodesmus sp. The identity of the consumer populations originating from each of 5 individuals

originally isolated from the experiment was confirmed morphologically by a taxonomist (however,

genotypic differences were not verified by sequencing for the *Daphnia*). All species reproduced

asexually, enabling us to maintain clonal populations of each genotype (algae) or putative clone

(*Daphnia*) in semi-continuous laboratory batch cultures until the start of the experiments. To avoid

confounding effects of plasticity or maternal effects, we maintained populations of these species in

common garden conditions—20 °C with 12:12 light-dark cycle—for over six months (many generations)

prior to subsequent experimentation. All organisms grew in autoclave-sterilized COMBO growth

medium. *Daphnia* grew in modified low-phosphorus COMBO (Kilham et al. 1998) that reduces bacterial

load. Aeration maintained culture health and growth and algal populations were in the exponential

growth phase at the start of the experiment. Additionally, *D. pulex* grew on a laboratory culture of

Ankistrodesmus sp. (sourced from Sciento UK) to ensure that daphnids were not acclimated to the algal

species used in the experiments. *Daphnia* cultures received 50 mL of dense *Ankistrodesmus* once per

week as a food source. Periodic filtering of adult and juvenile daphids into fresh medium removed

metabolic waste products and resting eggs.

165   *Experimental design*

Our experimental design crossed long-term evolutionary history—either to "ambient" or

"warm" temperature conditions— of each consumer–resource pair with 4 growth temperatures (Fig. 1).

In this way, we sought to examine how the effects of evolution in response to warming in a consumer–

resource pair depend upon the current thermal environment. The experiment occurred in 4 incubators

that varied in temperature: 14, 18, 22, and 26 °C. These temperatures encompass much of the observed

growing season activity range for the consumer from this system, with the presumed approximate

thermal optimum of reproduction of the ambient-origin population of 20 °C serving as the mid-point

across the range (Craddock 1976). Non-consumption controls additionally allowed us to verify the

importance of algal evolutionary history in the absence of consumers. We conducted the experiment for

each consumer–resource pair in two temporal blocks (for tractability and to avoid pseudoreplication at

the incubator level). Thus, we had 6 evolutionary history–consumption treatments crossed with 4

growth temperatures in each temporal block for each consumer–resource pair (n = 4 per treatment,

yielding a total of 96 microcosms within temporal block). At the start of each experimental block, we

estimated abundance of the ambient-origin and warm-origin resources using a hemocytometer to count

cells under a compound microscope (NIKON Eclipse Ni). We then standardized approximate starting

abundances of the two populations, however starting densities thus differed somewhat between blocks

within a resource species and between resource species. We randomized microcosm position within

each incubator. The experiment ran a total of 4 times (n = 8 for treatment across 2 temporal blocks,

yielding N = 192 per consumer–resource pair and N = 384 across both pairs).

To each sterile microcosm we added 100 mL of sterile COMBO medium and 100 mL of density-

standardized resource from either the ambient-origin or warm-origin population (either *C. reinhardtii* or

Desmodesmus sp.). To treatments containing *Daphnia*, we then added 5 adult daphnids either of

ambient-origin or warm-origin to each microcosm (previously starved for 24-hours, filtered, and rinsed

in sterile water to remove residual *Ankistrodesmus*), such that the evolutionary origin of the resource

was crossed with that of the consumer. Thus our evolutionary history–consumption treatments

repeated at 4 temperatures were: 1) ambient-origin resource alone, 2) warm-origin resource alone, 3)

both ambient-origin resource and consumer, 4) ambient-origin resource + warm-origin consumer, 5)

warm-origin resource + ambient-origin consumer, and 6) both warm-origin resource and consumer.

We confirmed the survival of the *Daphnia* on the second day of the experiment. The experiment

ran for 3 weeks, or approximately 42–100 generations of algae and 2–4 generations of *Daphnia*. A

failure in the first 24 hours of the experiment for 26 °C incubator resulted in the incubator temperature

declining to 1 °C overnight for the first temporal block using *C. reinhardtii*. We therefore discarded these

microcosms and reassembled them approximately 12 hours after the initial experimental set-up using

the same sources of algae and *Daphnia*.

200   *Estimation of equilibrium densities and interaction strength*

Responses included equilibrium density (per ml) of the resource species, as well as a census of

adult and larval *Daphnia* present in each microcosm. After 3 weeks, we first filtered and preserved all

Daphnia adults and juveniles from a microcosm in 95% ethanol for later enumeration under a dissecting

scope in two size categories, adults and juveniles. We next preserved two 150 µL samples of well-mixed

algae suspension from each microcosm using 15 μL of a 1% sorbitol solution in 96-well plates. Incubation

for 1 hour in the dark ensured sorbitol penetration to prevent cell lysis before sample storage at - 80 °C.

Flow cytometry (BD FACSCanto II high-throughput sampler) then estimated cell densities. We used

averaged log_{10}-transformed densities of the two technical flow cytometry replicates measured from

each microcosm in further analyses (see below). Parameters of flow cytometric lasers were as follows:

FSC: 136, SSC: 258, FITC: 484, PE: 432, PerCP-Cy5: 297, PE-Cy7: 570, APC: 440, APC-H7: 472, V450: 373

and V500: 484. For *Desmodesmus* sp., because flow cytometry measures particles, we report our results

as coenobia (i.e., small colonial cellular aggregates) rather than converting from aggregates to cells.

Coenobia are the units with which consumers interact, making them an appropriate unit for this species.

Safranin (Sigma-Aldrich) stained the *Daphnia* for simpler visualization of the preserved samples during

counting under a stereo microscope (Leica S9i), which ensured that molts were not counted

erroneously.

We used interaction strength to quantify the effect of consumers on resource abundance. To

calculate interaction strength (IS), using Equation 1 we calculated an inverse form of the log-response

ratio (also called the Dynamic Index) (Berlow et al. 2003; Schaum et al. 2018):

$$IS = \frac{ln(\frac{\overline{N}}{D_i})}{t}$$ Equation 1

Where $\overline{N}$ is the final mean algal abundance of the no consumption (algae-alone) controls for each

population, ambient-origin or warm-origin, for each resource species, *C. reinhardtii* or *Desmodesmus*

sp., D_i is the final algal abundance of replicate i from a given population in the presence of consumers,

and t is the length of the experiment (equal to 21 days for our experiment). With the inverse log-

response ratio, larger values of IS indicate stronger effects of consumers on resource abundance.

226   *Analyses*

We log$_{10}$-transformed abundances to better meet assumptions of normality in the subsequent

analyses. In all models, a block effect and all block interactions accounted for differences related to

temporal blocks. We conducted analyses to examine how current thermal environment and biotic

interactions determined the importance of past evolutionary history to warming in governing fitness (as

assessed by abundance across multiple generations of interactions) of resources and consumers. A

single MANOVA including evolutionary history treatments with and without consumers present would

obfuscate differences that might occur in the interaction between evolutionary history and temperature

when the algae grew alone or with consumers (in part through spurious significance related to the

presence or absence of *Daphnia* as a response in the two types of treatments). Therefore, to examine

how current biotic interactions impact the importance of evolutionary history, we conducted separate

analyses on the log$_{10}$-transformed abundance per ml of algae when consumers were present or absent.

In all models, the interaction between the main effects (evolutionary history and temperature) assessed

whether the effects or importance of evolutionary history depend upon current thermal environment.

Thus, we first conducted an analysis of variance (ANOVA) for each algal species grown alone (because

they were tested independently). These analyses tested whether algal evolutionary history interacted

significantly (α = 0.05) with temperature when consumers were absent. The model included the

evolutionary history (2 levels—ambient-origin or warm-origin resources) and temperature (4 levels—

14, 18, 22, and 26 °C test temperature) and the interaction, in addition to the block effect and block

interactions, with final algal density as the response.

We next conducted a separate multivariate analysis of variance (MANOVA) for each consumer–

resource pair. Responses included the log$_{10}$-transformed abundance per ml of the resource, interaction

strength, and the log$_{10}$-transformed abundances per microcosm of adult and juvenile *Daphnia*. Main

effects of the model included evolutionary history treatment (4 levels—ambient-origin resource and

consumers, ambient-origin resource and warm-origin consumers, warm-origin resource + ambient-

origin consumers, and warm-origin resource and consumers), temperature (4 levels—14, 18, 22, and 26

°C test temperature), and the interaction between temperature and evolutionary history. Univariate

tests assessed significance at α = 0.05 for individual responses. Orthogonal contrasts further

disentangled the drivers underlying the evolutionary history treatment and determined specifically if 1)

resource evolution, 2) consumer evolution, or 3) the interaction between evolutionary history of the

resource and consumer (i.e., coevolution) affected responses. All analyses were conducted in SAS 9.4

(SAS Institute 2011).

Results

Evolutionary history in response to warming in complex communities, current temperature, and

their interaction had highly significant effects for both consumer–resource pairs, but the relative

importance of resource, consumer, or coevolution differed between the pairs (see Tables 1 and 2 for

specific results from algae-alone ANOVAs and consumer–resource MANOVAs). We describe specific

effects below using results from ANOVAs for each response variable.

Current thermal environment influences the importance of prior resource evolution in response to

warming

Resource evolution in warmed complex communities affected both algal species when grown

alone, but the effects of evolutionary history were opposite in direction and differed in strength (Table

1). Warm-origin *Chlamydomonas reinhardtii* had higher abundance than the ambient-origin genotype at

all temperatures with no significant interaction between resource evolutionary history and current

thermal environment (effect of evolutionary history in algae-alone ANOVAs: F_1 = 158.47, p < 0.0001;

interaction between evolutionary history and temperature: NS) (Fig. 2A). For *Desmodesmus* sp., the

effect of evolutionary history depended upon temperature, with higher abundance of the warm-origin

genotype at 14 °C, but lower abundance than the ambient-adapted genotype at the remaining three

temperatures (effect of evolutionary history in algae-alone ANOVAs: F_1 = 7.02, p = 0.0109; interaction

between evolutionary history and temperature: F_3 = 5.86, p = 0.0017) (Fig. 2B).

Interspecific interactions modulate how current thermal environment interacts with prior evolution in

response to warming

Effects of interspecific interactions on resource abundance

The presence of the consumer *Daphnia pulex* altered the effects of resource evolutionary

history for both resources, but in opposite directions (Tables 2–4). For *C. reinhardtii*, where no

significant interaction between resource evolutionary history and temperature occurred when grown

alone, the presence of the consumer resulted in a significant evolutionary history by temperature

interaction driven by resource evolution (consumer–resource ANOVA resource evolution contrast: F_1 =

318.47, p < 0.0001). Specifically, when consumers were present, the importance of resource

evolutionary history in driving differences in algal abundance was higher at intermediate temperatures

(18 and 22 °C) than at the two temperature extremes (14 and 26 °C)—in other words, while

temperature did not affect the direction of the response to evolutionary history, it increased the

magnitude of the effect size at intermediate compared to extreme temperatures (effect of evolutionary

history in consumer–resource ANOVA: F_3 = 107.54, p < 0.0001; interaction between evolutionary history

and temperature: F_9 = 4.93, p < 0.0001) (Fig. 2C). On the other hand, for *Desmodesmus* sp., the presence

of consumers rendered the interaction between resource evolutionary history and temperature non-

significant (effect of evolutionary history in consumer–resource ANOVA: F_3 = 3.54, p = 0.0175;

interaction between evolutionary history and temperature: NS) (Fig. 2D). Here, when consumers were

present, resource evolution decreased algal abundance, reversing the pattern seen at 14 °C in the

absence of consumers (consumer–resource ANOVA resource evolution contrast: F_1 = 10.57, p = 0.0016)

(Fig. 2D).

Effects of evolutionary history on interaction strength

The importance of resource evolutionary history on algae abundance was reflected in the

changes to interaction strength for each consumer–resource pair (Tables 3 and 4). For the *D. pulex–C.*

reinhardtii pair, resource evolution reduced interaction strength (consumer–resource ANOVA resource

evolution contrast: F_1 = 54.90, p < 0.0001), but primarily at intermediate temperatures (interaction

between evolutionary history and temperature: F_9 = 5.16, p < 0.0001) (Fig. 2E). For the *D. pulex–*

Desmodesmus sp. pair, however, resource evolution increased interaction strength, independent of

temperature (consumer–resource ANOVA resource evolution contrast: F_1 = 5.86, p = 0.0174; interaction

between evolutionary history and temperature: NS) (Fig 2F).

Consumer responses to evolutionary history and temperature

For the consumer, *Daphnia pulex*, the effects of evolutionary history differed depending upon

with which resource it was paired. When paired with *C. reinhardtii*, resource evolution significantly

reduced the abundance of adult and juvenile *D. pulex* (adult resource evolution contrast: F_1 = 5.50, p =

0.0211; juvenile resource evolution contrast: F_1 = 5.07, p = 0.0266; Table 2, Fig. 3A,B). In this pairing,

neither consumer evolution nor coevolution affected the abundance of *D. pulex* and the effects of

resource evolution were independent of temperature (consumer–resource ANOVA interaction between

evolutionary history and temperature: NS).

When growing instead on the resource *Desmodesmus* sp., the effects of consumer evolution

became apparent. In this case, warm-origin consumers had overall lower abundance than the ambient-

origin population (adult consumer evolution contrast: F_1 = 15.87, p = 0.0001; juvenile consumer

evolution contrast: F_1 = 9.77, p = 0.0024; Table 4, Fig. 3C,D). There was no direct effect of *Desmodesmus*

evolution, but coevolution between the resource and consumer affected consumer abundance (adult

coevolution contrast: $F_1 = 8.52$, $p = 0.0044$; juvenile coevolution contrast: $F_1 = 12.55$, $p = 0.0006$). The

effects of evolutionary history were also dependent upon current thermal environment when growing

on *Desmodesmus* sp. (evolutionary history-temperature interaction for adult and juvenile daphnids

respectively: $F_9 = 2.09$, $p = 0.0380$ and $F_9 = 2.21$, $p = 0.0281$; Table 4, Fig. 3C,D). For *D. pulex* growing on

Desmodesmus sp., the effects of consumer evolution were most evident at 14 °C, where evolved *D.*

pulex adults ("Evolved consumer" and "Both evolved" treatments) had significantly reduced abundance

regardless of *Desmodesmus* evolutionary history compared to the ambient-origin *D. pulex* ("Neither

evolved" and "Evolved resource" treatments). The effects of coevolution between the consumer and

resource were strongest at 26 °C, with warm-adapted adult *D. pulex* abundance being significantly

reduced only in the presence of warm-adapted *Desmodesmus* sp. ("Both evolved" treatment) (Fig. 3B)*.*

In contrast to the temperature-dependence observed for the resource, *C. reinhardtii*, evolutionary

history was less important for consumer abundance at intermediate temperatures.

331   **Discussion**

332   Understanding the interaction between evolutionary history, the abiotic environment, and

biotic interactions is critical for a more nuanced understanding of how communities respond to

anthropogenic stressors. Here we showed that both the current thermal environment, as well as the

presence of interspecific interactions can affect the importance of prior evolution in response to

warming in complex communities, and that this evolution also influences the outcomes of current

ecological dynamics. For each consumer–resource pair, the effects of evolution were temperature-

dependent, but both the effects and the temperature-dependence itself additionally depended upon

the identity of evolving species in each pair.

340   *The importance of evolutionary history depends upon current thermal environment*

Evolution in warmed communities had significant effects on two algal resource species when

grown alone, but these effects were opposite in direction and differed in magnitude. In one species, *C.*

reinhardtii, prior evolution in warmed communities increased fitness (as assessed by abundance) across

all temperatures, a result that is broadly consistent with the "hotter is better" hypothesis (Liu et al.

2022), although our tested temperature range (determined by the thermal tolerance and activity of the

consumer) precluded us from definitively evaluating competing proposed evolutionary responses. The

positive direction of change for *Chlamydomonas reinhardtii* agrees with a previous experiment

examining the short-term effects of thermal evolution of *C. reinhardtii* independently isolated from

these same mesocosms (Schaum et al. 2017). In the second alga, *Desmodesmus* sp., the effects of

evolutionary history were dependent upon the current thermal environment. For this species, fitness

was lower in the warm-origin genotype at three of the four tested temperatures, but the pattern was

reversed at 14 °C, the lowest temperature in our range. This reversal of direction of effect across

temperatures suggests the possibility of a response that is harmful in part of the thermal range, but

possibly advantageous in another. The pattern of the result is surprising however, as the expectation is

that warm-origin populations would perform better at warm temperatures compared to colder

temperatures if the evolutionary response was directly driven by the experimentally imposed

temperature treatment.

Evaluating the mechanisms underlying this result was impossible given that we only successfully

isolated one genotype for each species from each of the ambient and warm mesocosms, however, these

results may be consistent with one or more aspects of our system. First, our algal genotypes were

isolated from the experimental ponds during the winter season, so this particular genotype may be

more successful in the winter conditions experienced by the experimentally warmed ponds. Typical

winter conditions in Dorset ranged from approximately 3–10 °C with a mean of around 6 °C (Yvon-

Durocher et al. 2015), thus the additional 4 °C of warming experimentally imposed upon these ponds

raised the winter temperature to within a few degrees of our 14 °C treatment. We did not examine the

effects of evolutionary history at any temperature lower than 14 °C, however, so we are unable to

compare the relative fitness of each genotype at typical winter temperatures compared to 14 °C.

Conversely, the warm-origin genotype that we isolated may be maladaptive, with lower fitness than the

ambient-origin genotype under most conditions. Anthropogenic warming is anticipated to create an

extinction debt in many communities (Cotto et al. 2017), with poorly adapted species expected to be

lost over time, especially as a result of stochastic processes combined with low dispersal (Kuussaari et al.

2009). It is possible that we have captured part of this process, with a maladapted genotype "trapped"

in the warm mesocosms. Another possibility is that the selection occurred in response to an agent that

emerged secondary to the experimentally imposed temperature treatment. Our manipulative approach

deconfounds other climatic variables that may affect evolutionary trajectory in natural populations, but

our populations were nonetheless potentially evolving in response to a number of factors (e.g.,

community composition) that had diverged over the course of this long-running experiment (Yvon-

Durocher et al. 2011, 2015, Dossena et al. 2012). Thus, while we can ultimately link our responses to the

experimentally-imposed warming treatment, populations are potentially evolving to a number of

simultaneous and related stressors (terHorst et al. 2018). While this situation reflects the expectation

for natural communities responding to habitat warming—namely that responses will encompass altered

biotic interactions as well as direct responses to changed abiotic factors—it is therefore impossible for

us to definitively disentangle whether the lowered fitness that we observed in warm-origin populations

of *Desmodesmus* sp. was the result of maladaptive evolutionary response or evolutionary response to

conditions not captured in our subsequent experiments.

Current ecological context modulates the temperature-dependence of the effects of evolutionary history

Our understanding of how evolution is altered in complex communities is still limited. However,

previous studies suggest that evolution to environmental stressors, like habitat warming, that occurs in

populations embedded within communities can yield different outcomes compared to evolution in

single populations (Van Doorslaer et al. 2009b, 2010, terHorst et al. 2018, Hiltunen et al. 2018, De

Meester et al. 2019). Our experiment complements a growing body of evidence that shows that

evolutionary responses to warming do occur in complex communities (De Meester et al. 2019), and that

such evolution has important consequences for both the direct responses to evolutionary stressors as

well as for the interspecific interactions that ultimately shape communities.

The inclusion of consumer–resource interactions also introduced an additional layer of

complexity necessary to understand how the importance of evolutionary history is affected by

temperature. Organisms exhibit a mixture of unimodal and linear responses of traits to temperature,

and interacting organisms differ in both the ecological and evolutionary responses of these traits. The

interactions themselves may therefore depend upon temperature, influencing the outcomes of

evolutionary change evaluated across temperatures (Barton and Yvon-Durocher 2019). For both

consumer–resource pairs in our study system, the presence of the consumer altered the interaction

between evolutionary history and temperature, indicating that examining the effects of past evolution

in response to environmental change can be highly dependent on the current ecological conditions.

In the case of the *Daphnia pulex–C. reinhardtii* pair, the importance of evolutionary history was

only temperature-dependent when the consumer was present. For this pair, the magnitude of the effect

evolutionary history on algal abundance was higher at intermediate temperatures compared to

temperature extremes, adding nuance to the result that "hotter is better", making it instead "hotter is

best at moderate or intermediate temperatures". Consumer evolution was unimportant when *D. pulex*

was paired with *C. reinhardtii*—warm-origin *C. reinhardtii* significantly reduced the abundance of the

consumer regardless of temperature. Thus, "hotter is better" for this resource translates into "hotter is

worse" for the consumer. We noted that the warm-origin genotype exhibited a mucilaginous extra-

cellular matrix that might be expected to reduce ingestion rates for filter-feeding consumers like *D.*

pulex. These results are also consistent with observations of an increased carbon to phosphorus ratio

(i.e.*,* reduced resource quality) observed previously in warm-adapted *C. reinhardtii* independently

isolated from these experimental mesocosms (Yvon-Durocher et al. 2017). Such a reduction in resource

quality would be expected to negatively affect consumer performance (Leal et al. 2017), but an analysis

of nutrient content of the algae was outside the scope of our experiment. It is noteworthy, however,

that despite the lack of a significant interaction between evolutionary history and temperature, the

importance of resource evolution was strongest at 22 °C, the temperature at which *D. pulex* abundance

was highest, and weakest at 26 °C, where the daphnids struggled to persist. The evolution that we

observed in *C. reinhardtii* thus appeared to confer a double advantage to the resource, making it both

more fit, while simultaneously reducing the success of the consumer, possibly the reason why we

observed larger effect sizes for resource evolutionary history in the presence of the consumer, especially

at intermediate temperatures (i.e., effect size more than doubled on a $\log_{10}$ scale).

The effects of consumers for *Desmodesmus* sp. were opposite to those for *C. reinhardtii*. Here,

the presence of the consumer eliminated the temperature-dependence seen when *Desmodesmus* sp.

grew alone, and even reversed the direction of the effect of evolutionary history at 14 °C. When paired

with the consumer, this subtlety across temperatures was lost, and warm-origin *Desmodesmus* sp.

underperformed relative to ambient-origin *Desmodesmus* regardless of temperature. One possible

explanation for the apparently reduced fitness of the warm-origin genotype could be that it served as a

superior food source for the *Daphnia*. The two *Desmodesmus* sp. lineages have different propensities to

remain as single cells or aggregate into 2, 3, or 4-celled coenobia, with the warm-origin lineage occurring

more frequently as 1 or 2-cell aggregates and the ambient-origin lineage more frequently as larger

aggregates (unpublished data). This difference in average particle size could affect the filtering efficiency

of the *Daphnia*, making the warm-origin genotype potentially more accessible to the consumer and

intensifying the fitness disadvantage of the warm-origin genotype in the presence of consumers. If this

were the case, we would expect to see that the *Daphnia* achieved higher abundances when consuming

the warm-origin algae. In fact, *Desmodesmus* sp. evolutionary history had contrasting effects on the

consumer's fitness depending upon consumer thermal origin (i.e., the effect of coevolution on *D. pulex*

abundance in this pair). Ambient-origin consumers exhibited a trend towards slightly higher fitness with

warm-origin resource, but warm-origin *Daphnia* had lowest fitness when paired with warm-origin

Desmodesmus. Thus, the effect does not appear to be consistent with increased consumption by the

Daphnia. Additionally, in this pair, the *Daphnia* experienced temperature-dependent effects of

evolutionary history. Here, effects were strongest at the two temperature extremes, with warm-origin

Daphnia having lowest abundance at these temperatures, and especially at 26 °C and when paired with

warm-origin *Desmodesmus* sp.

We examined the temperature-dependent consequences of evolution for two consumer–

resource pairs over multiple generations of interactions and found evidence for adaptive evolution in

response to warming in one of the two species pairs examined. For the *D. pulex–C. reinhardtii* pair, we

observed adaptive evolution consistent with the "hotter is better" hypothesis for the resource, but at

the same time observed that hotter was worse for the consumer, likely due to the fact that hotter was

worse for the interaction from the perspective of the consumer. This result may seem surprising, but not

only is evolutionary response to warming not expected in all species (Hoffmann and Sgró 2011), it also

remains possible that the fitness consequences of the response depend upon the length of the

temperature exposure. We focused on the effects of constant temperature over multiple generations,

however, the semi-natural mesocosms from which we isolated these genotypes additionally fluctuate in

temperature both diurnally and seasonally. These fluctuations may be an important component of the

selection imposed by the experimental warming treatment (Diamond 2017), and organisms in warmed

ponds may experience more frequent and intense, but shorter, periods of thermal stress. A more

complete understanding of the interaction between current thermal environment and evolutionary

history, therefore, will require an improved understanding of how populations respond to fluctuating

temperatures.

Conclusions

Our work demonstrates that local thermal environment and biotic interactions can together

modulate the importance of past thermal evolutionary history. For example, in some cases, evolution in

response to warming can be beneficial for resources but detrimental for consumers, suggesting that we

must account for the effects of warming on ingestion rates (see e.g., Barneche et al. 2021). In other

cases, the picture is more complex and changes across a temperature range. This topic deserves further

attention to improve our understanding of how populations and communities evolve to habitat

warming, and how this then feeds back to drive the effects of the evolution. We have built upon

previous studies that have primarily used space-for-time latitudinal substitutions in comparisons of

different thermally adapted populations (e.g., Sanford et al. 2003, De Block et al. 2013, Dinh Van et al.

2013, Tran et al. 2016) and argue that future studies should consider both the current biotic and thermal

context of their experiments, while definitely attributing evolutionary response to a manipulated factor

of interest. These community properties can affect outcomes, not just in terms of the magnitude of the

effect size, but even in direction of the effect. It is therefore of the utmost importance that we account

for these interactions in our collective understanding. Evolution can drive responses in unexpected

directions, but it is essential that we consider evolutionary responses to anthropogenic change in the

context of both the current abiotic and biotic conditions to understand the full extent and direction of

its effects.

Acknowledgements

The French ANR through LabEx TULIP (ANR-10-LABX-41) and the FRAGCLIM Consolidator Grant,

funded by the European Research Council under the European Union's Horizon 2020 research and

innovation programme (Grant Agreement Number 726176) supported this research. We thank X

anonymous referees for their feedback.

Author contributions

CAF and JMM conceived and designed the study. CAF, SAC, AG, and EB isolated organisms and

collected data. CAF analyzed the data. CAF wrote the first draft of the manuscript. CAF, JMM, SAC, and

EB jointly revised the manuscript. The authors declare no competing financial interests.

References

Álvarez-Codesal, S., C. A. Faillace, A. Garreau, E. Bestion, A. D. Synodinos, and J. M. Montoya. 2023.

Thermal mismatches explain consumer–resource dynamics in response to environmental

warming. Ecology and Evolution 13:e10179.

Angilletta Jr., M. J., A. F. Bennett, H. Guderley, C. A. Navas, F. Seebacher, and R. S. Wilson. 2006.

Coadaptation: A unifying principle in evolutionary thermal biology. Physiological and

Biochemical Zoology 79:282–294.

Barneche, D. R., C. J. Hulatt, M. Dossena, D. Padfield, G. Woodward, M. Trimmer, and G. Yvon-Durocher.

2021. Warming impairs trophic transfer efficiency in a long-term field experiment. Nature

592:76–79.

Barton, S., and G. Yvon-Durocher. 2019. Quantifying the temperature dependence of growth rate in

marine phytoplankton within and across species. Limnology and Oceanography 64:2081–2091.

Comeau, A. M., G. M. Douglas, and M. G. I. Langille. 2017. Microbiome Helper: a Custom and

Streamlined Workflow for Microbiome Research. mSystems.

Cotto, O., J. Wessely, D. Georges, G. Klonner, M. Schmid, S. Dullinger, W. Thuiller, and F. Guillaume.

2017. A dynamic eco-evolutionary model predicts slow response of alpine plants to climate

warming. Nature Communications 8:15399.

Craddock, D. R. 1976. Effects of increased water temperature on *Daphnia pulex*. Fishery Bulletin 74:403–

408.

De Block, M., K. Pauwels, M. Van Den Broeck, L. De Meester, and R. Stoks. 2013. Local genetic

adaptation generates latitude-specific effects of warming on predator-prey interactions. Global

Change Biology 19:689–696.

De Meester, L., K. I. Brans, L. Govaert, C. Souffreau, S. Mukherjee, H. Vanvelk, K. Korzeniowski, L.

Kilsdonk, E. Decaestecker, R. Stoks, and M. C. Urban. 2019. Analysing eco-evolutionary

dynamics—The challenging complexity of the real world. Functional Ecology 33:43–59.

Diamond, S. E. 2017. Evolutionary potential of upper thermal tolerance: biogeographic patterns and

expectations under climate change. Annals of the New York Academy of Sciences 1389:5–19.

Diamond, S. E., L. Chick, A. Perez, S. A. Strickler, and R. A. Martin. 2017. Rapid evolution of ant thermal

tolerance across an urban-rural temperature cline. Biological Journal of the Linnean Society

121:248–257.

Dinh Van, K., L. Janssens, S. Debecker, M. De Jonge, P. Lambret, V. Nilsson-Örtman, L. Bervoets, and R.

Stoks. 2013. Susceptibility to a metal under global warming is shaped by thermal adaptation

along a latitudinal gradient. Global Change Biology 19:2625–2633.

Dossena, M., G. Yvon-Durocher, J. Grey, J. M. Montoya, D. M. Perkins, M. Trimmer, and G. Woodward.

2012. Warming alters community size structure and ecosystem functioning. Proceedings of the

Royal Society B: Biological Sciences 279:3011–3019.

Gaitán-Espitia, J. D., and A. J. Hobday. 2021. Evolutionary principles and genetic considerations for

guiding conservation interventions under climate change. Global Change Biology 27:475–488.

Geerts, A. N., J. Vanoverbeke, B. Vanschoenwinkel, W. Van Doorslaer, H. Feuchtmayr, D. Atkinson, B.

Moss, T. A. Davidson, C. D. Sayer, and L. De Meester. 2015. Rapid evolution of thermal tolerance

in the water flea *Daphnia*. Nature Climate Change 5:665.

Gilchrist, G. W., and R. B. Huey. 1999. The direct response of *Drosophila melanogaster* to selection on

knockdown temperature. Heredity 83:15–29.

Hangartner, S., and A. A. Hoffmann. 2016. Evolutionary potential of multiple measures of upper thermal

tolerance in *Drosophila melanogaster*. Functional Ecology 30:442–452.

Higgins, J. K., H. J. MacLean, L. B. Buckley, and J. G. Kingsolver. 2014. Geographic differences and

microevolutionary changes in thermal sensitivity of butterfly larvae in response to climate.

Functional Ecology 28:982–989.

Hiltunen, T., J. Cairns, J. Frickel, M. Jalasvuori, J. Laakso, V. Kaitala, S. Künzel, E. Karakoc, and L. Becks.

2018. Dual-stressor selection alters eco-evolutionary dynamics in experimental communities.

Nature Ecology and Evolution 2:1974–1981.

Hoffmann, A. A., and C. M. Sgró. 2011. Climate change and evolutionary adaptation. Nature 470:479–

485.

Johansson, M. P., F. Ermold, B. K. Kristjánsson, and A. Laurila. 2016. Divergence of gastropod life history

in contrasting thermal environments in a geothermal lake. Journal of Evolutionary Biology

29:2043–2053.

Johansson, M. P., and A. Laurila. 2017. Maximum thermal tolerance trades off with chronic tolerance of

high temperature in contrasting thermal populations of *Radix balthica*. Ecology and Evolution

7:3149–3156.

Kuussaari, M., R. Bommarco, R. K. Heikkinen, A. Helm, J. Krauss, R. Lindborg, E. Öckinger, M. Pärtel, J.

Pino, F. Rodà, C. Stefanescu, T. Teder, M. Zobel, and I. Steffan-Dewenter. 2009. Extinction debt:

a challenge for biodiversity conservation. Trends in Ecology & Evolution 24:564–571.

Leal, M. C., O. Seehausen, and B. Matthews. 2017. The Ecology and Evolution of Stoichiometric

Phenotypes. Trends in Ecology & Evolution 32:108–117.

Liu, K., B. Chen, and H. Liu. 2022. Evidence of partial thermal compensation in natural phytoplankton

assemblages. Limnology and Oceanography Letters 7:122–130.

Logan, M. L., R. M. Cox, and R. Calsbeek. 2014. Natural selection on thermal performance in a novel

thermal environment. Proceedings of the National Academy of Sciences of the United States of

America 111:14165–9.

O'Connor, M. I., E. R. Selig, M. L. Pinsky, and F. Altermatt. 2012. Toward a conceptual synthesis for

climate change responses. Global Ecology and Biogeography 21:693–703.

O'Donnell, D. R., C. R. Hamman, E. C. Johnson, C. T. Kremer, C. A. Klausmeier, and E. Litchman. 2018.

Rapid thermal adaptation in a marine diatom reveals constraints and trade-offs. Global Change

Biology 24:4554–4565.

Padfield, D., G. Yvon-Durocher, A. Buckling, S. Jennings, and G. Yvon-Durocher. 2016. Rapid evolution of

metabolic traits explains thermal adaptation in phytoplankton. Ecology Letters 19:133–142.

Parmesan, C. 2006. Ecological and Evolutionary Responses to Recent Climate Change. Annual Review of

Ecology, Evolution, and Systematics 37:637–669.

Radchuk, V., T. Reed, C. Teplitsky, M. van de Pol, A. Charmantier, C. Hassall, P. Adamík, F. Adriaensen, M.

P. Ahola, P. Arcese, J. Miguel Avilés, J. Balbontin, K. S. Berg, A. Borras, S. Burthe, J. Clobert, N.

Dehnhard, F. de Lope, A. A. Dhondt, N. J. Dingemanse, H. Doi, T. Eeva, J. Fickel, I. Filella, F.

Fossøy, A. E. Goodenough, S. J. G. Hall, B. Hansson, M. Harris, D. Hasselquist, T. Hickler, J. Joshi,

H. Kharouba, J. G. Martínez, J.-B. Mihoub, J. A. Mills, M. Molina-Morales, A. Moksnes, A. Ozgul,

D. Parejo, P. Pilard, M. Poisbleau, F. Rousset, M.-O. Rödel, D. Scott, J. C. Senar, C. Stefanescu, B.

G. Stokke, T. Kusano, M. Tarka, C. E. Tarwater, K. Thonicke, J. Thorley, A. Wilting, P. Tryjanowski,

J. Merilä, B. C. Sheldon, A. Pape Møller, E. Matthysen, F. Janzen, F. S. Dobson, M. E. Visser, S. R.

Beissinger, A. Courtiol, and S. Kramer-Schadt. 2019. Adaptive responses of animals to climate

change are most likely insufficient. Nature Communications 10:3109.

Rezende, E. L., L. E. Castañeda, and M. Santos. 2014. Tolerance landscapes in thermal ecology.

Functional Ecology 28:799–809.

Rudman, S. M., M. Kreitzman, K. M. A. Chan, and D. Schluter. 2017. Evosystem Services: Rapid Evolution

and the Provision of Ecosystem Services. Trends in Ecology & Evolution 32:403–415.

Sanford, E., M. S. Roth, G. C. Johns, J. P. Wares, and G. N. Somero. 2003. Local selection and latitudinal

variation in a marine predator- prey interaction. Science 300:1135–1137.

SAS Institute. 2011. The SAS system for Windows. SAS Institute, Cary, NC.

Schaum, C.-E., S. Barton, E. Bestion, A. Buckling, B. Garcia-Carreras, P. Lopez, C. Lowe, S. Pawar, N.

Smirnoff, M. Trimmer, and G. Yvon-Durocher. 2017. Adaptation of phytoplankton to a decade of

experimental warming linked to increased photosynthesis. Nature Ecology and Evolution 1:94.

Schaum, C.-E., R. Ffrench-Constant, C. Lowe, J. S. Ólafsson, D. Padfield, and G. Yvon-Durocher. 2018.

Temperature-driven selection on metabolic traits increases the strength of an algal-grazer

interaction in naturally warmed streams. Global Change Biology 24:1793–1803.

Stoks, R., A. N. Geerts, and L. De Meester. 2014. Evolutionary and plastic responses of freshwater

invertebrates to climate change: Realized patterns and future potential. Evolutionary

Applications 7:42–55.

Tabi, A., F. Pennekamp, F. Altermatt, R. Alther, E. A. Fronhofer, K. Horgan, E. Mächler, M. Pontarp, O. L.

Petchey, and S. Saavedra. 2020. Species multidimensional effects explain idiosyncratic responses

of communities to environmental change. Nature Ecology and Evolution:1–8.

terHorst, C. P., P. C. Zee, K. D. Heath, T. E. Miller, A. I. Pastore, S. Patel, S. J. Schreiber, M. J. Wade, and

M. R. Walsh. 2018. Evolution in a community context: Trait responses to multiple species

interactions. The American Naturalist 191:368–380.

Tran, T. T., L. Janssens, K. V. Dinh, L. Op de Beeck, and R. Stoks. 2016. Evolution determines how global

warming and pesticide exposure will shape predator–prey interactions with vector mosquitoes.

Evolutionary Applications 9:818–830.

Van Doorslaer, W., R. Stoks, C. Duvivier, A. Bednarska, and L. De Meester. 2009a. Population dynamics

determine genetic adaptation to temperature in *Daphnia*. Evolution 63:1867–1878.

Van Doorslaer, W., R. Stoks, I. Swillen, H. Feuchtmayr, D. Atkinson, B. Moss, and L. De Meester. 2010.

Experimental thermal microevolution in community-embedded *Daphnia* populations. Climate

Research 43:81–89.

Van Doorslaer, W., J. Vanoverbeke, C. Duvivier, S. Rousseaux, M. Jansen, B. Jansen, H. Feuchtmayr, D.

Atkinson, B. Moss, R. Stoks, and L. De Meester. 2009b. Local adaptation to higher temperatures

reduces immigration success of genotypes from a warmer region in the water flea *Daphnia*.

Global Change Biology 15:3046–3055.

Yvon-Durocher, G., A. P. Allen, M. Cellamare, M. Dossena, K. J. Gaston, M. Leitao, J. M. Montoya, D. C.

Reuman, G. Woodward, and M. Trimmer. 2015. Five years of experimental warming increases

the biodiversity and productivity of phytoplankton. PLoS Biology 13:e1002324.

Yvon-Durocher, G., J. M. Montoya, M. Trimmer, and G. Woodward. 2011. Warming alters the size

spectrum and shifts the distribution of biomass in freshwater ecosystems. Global Change

Biology 17:1681–1694.

Yvon-Durocher, G., C.-E. Schaum, and M. Trimmer. 2017. The temperature dependence of

phytoplankton stoichiometry: investigating the roles of species sorting and local adaptation.

Frontiers in Microbiology 8:2003.

Table 1. Results from univariate ANOVAs (n = 8) examining the effects of resource evolution when

growing alone for *Chlamydomonas reinhardtii* and *Desmodesmus* sp. The models included an evolution

treatment and temperature as two main effects, as well as a temporal blocking variable and all

interactions. Here, the evolution treatment included just the evolutionary history of the resource

species, either ambient-origin or warm-origin. The interaction between evolution treatment and

temperature evaluated any temperature-dependence of the effects of evolution. Significant results are

in bold.

C. reinhardtii			
	d.f.	F-value	p-value
Evolution	1	**158.47**	**< 0.0001**
Temperature	3	**12.56**	**< 0.0001**
Temporal block	1	**54.12**	**< 0.0001**
Evolution–temperature	3	0.67	0.5776
Temperature–block	3	**15.43**	**< 0.0001**
Evolution treatment– block	1	**127.63**	**< 0.0001**
Evolution treatment–temperature–block	3	**5.96**	**0.0015**
Desmodesmus sp.			
	d.f.	F-value	p-value
Evolution	1	**7.02**	**0.0109**
Temperature	3	**47.41**	**< 0.0001**
Temporal block	1	3.94	0.0529
Evolution– temperature	3	**5.86**	**0.0017**
Temperature–block	3	**5.19**	**0.0035**
Evolution treatment–block	1	**42.75**	**< 0.0001**
Evolution treatment–temperature–block	3	**3.10**	**0.0353**

Table 2. MANOVA (n = 8) results examining the effects of evolution of both the consumer and the

resource in complex communities in response to warming for two consumer–resource pairs, *C.*

reinhardtii and *D. pulex*, and *Desmodesmus* sp. and *D. pulex*. The models included an evolution

treatment and temperature as two main effects, as well as a temporal blocking variable and all

interactions. Here, the evolution treatment included the four possible combinations crossing the

ambient- or warm-origin populations of both resources and consumers, but these analyses did not

include algae-alone controls. The interaction between the evolution treatment and temperature

evaluated any temperature-dependence of the effects of evolution. Orthogonal contrasts evaluated the

effects of 1) resource evolution, 2) consumer evolution, and 3) coevolution between the resource and

the consumer. Reported are Wilks' Lambda F-values and associated p-values. Significant results are in

bold.

C. reinhardtii–D. pulex			
	d.f.	F-value	p-value
Evolution	12, 243.7	**70495.6**	**< 0.0001**
Temperature	12, 243.7	**45724.4**	**< 0.0001**
Temporal block	4, 92	**1.82E10**	**< 0.0001**
Evolution–temperature	36, 346.5	**1163.05**	**< 0.0001**
Temperature–block	12, 243.7	**51809.9**	**< 0.0001**
Evolution treatment– block	12, 243.7	**63495.4**	**< 0.0001**
Evolution treatment–temperature–block	36, 346.5	**1863.05**	**< 0.0001**
Resource evolution contrast	4, 92	**4.93E10**	**< 0.0001**
Consumer evolution contrast	3, 93	1.72	0.1683
Coevolution contrast	3, 93	1.05	0.3729
Desmodesmus sp.–*D. pulex*			
	d.f.	F-value	p-value
Evolution	12, 243.7	**578.58**	**< 0.0001**
Temperature	12, 243.7	**563.64**	**< 0.0001**
Temporal block	4, 92	**2818757**	**< 0.0001**
Evolution–temperature	36, 346.5	**56.96**	**< 0.0001**
Temperature–block	12, 243.7	**363.01**	**< 0.0001**
Evolution treatment– block	12, 243.7	**633.42**	**< 0.0001**
Evolution treatment–temperature–block	36, 346.5	**61.15**	**< 0.0001**
Resource evolution contrast	4, 92	**55119.7**	**< 0.0001**
Consumer evolution contrast	3, 93	**52771.4**	**< 0.0001**
Coevolution contrast	3, 93	**55945.4**	**< 0.0001**

Table 3. *Chlamydomonas reinhardtii* and *Daphnia pulex* univariate ANOVA (n = 8) results examining the effects of evolution of both the consumer and the resource in complex communities in response to warming. The model included an evolution treatment and temperature as two main effects, as well as a temporal blocking variable and all interactions. Here, the evolution treatment included the four possible combinations crossing the ambient- or warm-origin populations of both resources and consumers, but these analyses did not include algae-alone controls. The interaction between the evolution treatment and temperature evaluated any temperature-dependence of the effects of evolution. Orthogonal contrasts evaluated the effects of 1) resource evolution, 2) consumer evolution, and 3) coevolution between the resource and the consumer. Reported are Type III SS F-values and p-values for the model (due to one missing value causing a slightly unbalanced design). Significant results are in bold.

	C. reinhardtii $\log_{10}$ abundance			Interaction strength			*D. pulex* adult $\log_{10}$ abundance			*D. pulex* juvenile $\log_{10}$ abundance		
	d.f.	F-value	p-value	d.f.	F-value	p-value	d.f.	F-value	p-value	d.f.	F-value	p-value
Evolution	3	**107.54**	**< 0.0001**	3	**19.64**	**< 0.0001**	3	2.35	0.0772	3	1.80	0.1529
Temperature	3	**17.48**	**< 0.0001**	3	**48.61**	**< 0.0001**	3	**6.37**	**0.0006**	3	**7.48**	**0.0001**
Temporal block	1	**162.18**	**< 0.0001**	1	**359.69**	**< 0.0001**	1	0.10	0.7532	1	0.00	0.9543
Evolution–temperature	9	**4.39**	**< 0.0001**	9	**5.16**	**< 0.0001**	9	1.20	0.3067	9	0.53	0.8469
Temperature–block	3	**12.52**	**< 0.0001**	3	**25.83**	**< 0.0001**	3	**5.08**	**0.0026**	3	1.55	0.2062
Evolution–block	3	**4.00**	**0.0099**	3	**11.85**	**< 0.0001**	3	**3.51**	**0.0183**	3	2.31	0.0815
Evolution–temperature–block	9	1.63	0.1170	9	**3.52**	**0.0009**	9	1.40	0.2010	9	1.07	0.3949
Resource evolution contrast	1	**318.47**	**< 0.0001**	1	**54.90**	**< 0.0001**	1	**5.50**	**0.0211**	1	**5.07**	**0.0266**
Consumer evolution contrast	1	1.36	0.2456	1	1.36	0.2456	1	1.48	0.2262	1	0.29	0.5904
Coevolution contrast	1	2.53	0.1150	1	2.53	0.1150	1	0.01	0.9339	1	0.00	0.9740

Table 4. *Desmodesmus* sp. and *Daphnia pulex* univariate ANOVA (n = 8) results examining the effects of evolution of both the consumer and the resource in complex communities in response to warming. The model included an evolution treatment and temperature as two main effects, as well as a temporal blocking variable and all interactions. Here, the evolution treatment included the four possible combinations crossing the ambient- or warm-origin populations of both resources and consumers, but these analyses did not include algae-alone controls. The interaction between the evolution treatment and temperature evaluated any temperature-dependence of the effects of evolution. Orthogonal contrasts evaluated the effects of 1) resource evolution, 2) consumer evolution, and 3) coevolution between the resource and the consumer. Reported are Type III SS F-values and p-values for the model (due to one missing value causing a slightly unbalanced design). Significant results are in bold.

	Desmodesmus sp. $\log_{10}$ abundance			Interaction strength			*D. pulex* adult $\log_{10}$ abundance			*D. pulex* juvenile $\log_{10}$ abundance		
	d.f.	F-value	p-value	d.f.	F-value	p-value	d.f.	F-value	p-value	d.f.	F-value	p-value
Evolution	3	**3.54**	**0.0175**	3	2.50	0.0645	3	**8.26**	**< 0.0001**	3	**8.66**	**< 0.0001**
Temperature	3	**11.26**	**< 0.0001**	3	**13.15**	**< 0.0001**	3	**14.08**	**< 0.0001**	3	**33.57**	**< 0.0001**
Temporal block	1	**47.32**	**< 0.0001**	1	0.88	0.3513	1	**17.64**	**< 0.0001**	1	**83.67**	**< 0.0001**
Evolution–temperature	9	0.73	0.6804	9	0.77	0.6466	9	**2.09**	**0.0380**	9	2.21	0.0281
Temperature–block	3	**5.39**	**0.0018**	3	**5.96**	**0.0009**	3	**4.62**	**0.0046**	3	**54.08**	**< 0.0001**
Evolution–block	3	**14.12**	**< 0.0001**	3	**10.25**	**< 0.0001**	3	2.66	0.0528	3	**4.92**	**0.0032**
Evolution–temperature–block	9	0.45	0.9066	9	0.33	0.9636	9	1.33	0.2309	9	**2.10**	**0.0368**
Resource evolution contrast	1	**10.57**	**0.0016**	1	**5.86**	**0.0174**	1	0.51	0.4752	1	3.61	0.0605
Consumer evolution contrast	1	0.07	0.7925	1	1.16	0.2846	1	**15.87**	**0.0001**	1	**9.77**	**0.0024**
Coevolution contrast	1	0.01	0.9385	1	0.58	0.4496	1	**8.52**	**0.0044**	1	**12.55**	**0.0006**

Figure 1. Experimental design in which populations from two resource species and one consumer species were first collected from a long-running warming experiment using complex communities of plankton in outdoor mesocosms. After isolation and a period of laboratory common garden, for each consumer–resource pair, subsequent experiments examined the interaction between evolutionary history of resources and consumers and current thermal environment on population abundance.

Figure 2A–F. The effects of evolution in warmed communities on the $\log_{10}$-abundance of *C. reinhardtii* and *Desmodesmus* sp. depend upon current thermal environment and whether consumers are absent (A, B) or present (C, D), with related effects to the consumer–resource interaction strength (E, F). Evolutionary history treatments are shown on the x-axis and color of the line indicates the test temperature. Circles are the means (n = 8) of each treatment with the error bars showing ± the SEM.

Figure 3A–D. The effects of evolution in warmed communities on the $\log_{10}$-abundance of *D. pulex* juveniles (A) and adults (B) growing on *C. reinhardtii* and on *Desmodesmus* sp. (C, D) depend upon current thermal environment. In all plots, evolutionary history treatments are shown on the x-axis and color of the line indicates the test temperature. Circles are the means (n = 8) of each treatment with the error bars showing ± the SEM.