Title: Thermal suppression of gametogenesis explains historical collapses in larval recruitment

Authors: Daniel K. Okamoto[1,*], Nathan B. Spindel [1], Brenna Collicutt[2], Maya J. Mustermann[1],

Sam Karelitz[1], Iria Gimenez[2], Kate Rolheiser[2], Evan Cronmiller[2], Megan Foss[2], Natalie

Mahara[2], Dan Swezey[3], Rachele Ferraro[1], Laura Rogers-Bennett[3] , and Stephen Schroeter[4].

Affiliations

1: Department of Biological Science, The Florida State University, Tallahassee, Florida 32303

2: Hakai Institute, Quadra Island, British Columbia, Canada

3: Coastal Marine Science Institute and Bodega Marine Lab, University of California, Davis

4: Marine Science Institute, University of California, Santa Barbara

*: Corresponding Author: dokamoto@bio.fsu.edu

Abstract

Projections for population viability under climate change are often made using estimates

of thermal lethal thresholds. These estimates vary across life history stages and can be valuable

for explaining or forecasting shifts in population viability. However, sub-lethal temperatures can

also lead to declines in vital rates and ultimately shape fluctuations in the reproductive viability

of populations. For example, anomalous climatic events can suppress reproduction and lead to

recruitment failure well before early life stages or adult stages become affected. These sub-lethal

impacts make the effects of climate change more severe than lethal thresholds predict. Despite a

growing awareness of this issue, tying sub-lethal effects to observed recruitment failure remains

a challenge especially in marine environments. Here, we experimentally show that sub-lethal

thermal suppression of female gametogenesis offers a plausible explanation for historical

collapses in sea urchin recruitment during marine heatwaves. These sub-lethal thermal

sensitivities of reproduction can narrow the thermal envelope for population viability compared

to what lethal limits predict.

Introduction

Predicting the impact of warming and extreme heatwaves on ecosystems and populations remains one of the most daunting but important tasks in ecology and conservation. Information about the future dynamics and viability of populations can be gleaned from measuring thermal lethal thresholds (or critical thermal limits - CTLs) in controlled settings, as well as from historical patterns of population performance in nature. For the latter, extrapolating historical patterns without understanding processes that shaped them remains dubious; meanwhile, expectations from CTLs often underpredict the abruptness and magnitude of population responses to warming in nature [1-3]. For example, warming and heatwaves may lead to recruitment failure in populations even when temperatures remain below lethal thresholds [4,5]. Thus, the realized thermal niche of many species (i.e., thermal regimes in which a population exhibits viability over time) is often narrower than expected based on CTLs for individual life stages (gametes, juveniles, adults, etc. - [6]). Combining inference from laboratory studies and historical patterns and analysis can provide a more comprehensive understanding of the processes that shape recruitment and population dynamics.

One pathway by which sub-lethal temperatures can alter population viability is by affecting gametogenesis, maturation, and/or sterility. Reproductive phenology for a diversity of plants and animals responds directly to temperature [7] and as such, temperature can significantly impact reproductive success and thereby fitness and productivity in a warming world. In plants, for example, warmer winters can lead to crop failure by disrupting vernalization pathways. This family of processes, in which plants require specific thermal cues, as well as other environmental patterns or thresholds for induction of flowering and seed/fruit development has long been studied and understood in agricultural and laboratory plant systems e.g., [8,9]. Likewise, a diversity of animal taxa can exhibit infertility or low fecundity at sub-lethal temperatures [10-14].

Yet for animals, and marine organisms with planktonic larvae in particular, it remains unclear if and to what extent patterns of recruitment failure in nature are shaped by sub-lethal thermal suppression of reproduction. This gap remains, in part, for several reasons. First, attributing causality to observed changes in year-class strength remains a challenge for animals with dispersive reproductive propagules. Shifts in recruitment may be explained by variation in

food supply [15], trends in reproductive and mating success [16], shifts in dispersal [17], or a host of

other factors that may also be directly or indirectly related to warming. Second, thermal limits

are often estimated using constant temperatures; yet in the oceans, temperature is highly

dynamic, and those dynamics can shape patterns of reproduction. Thus, while it is well known

that sub-lethal temperature can limit reproduction in marine invertebrates [18,19], it remains less

well understood if dynamic heatwave trends can induce such limitations and/or plausibly explain

collapses in recruitment. If climate change and extreme events in nature lead to suppressed

gametogenesis well before lethal limits of organisms are reached (e.g., thermal stress that

decreases larval, juvenile, or adult survival), then declines in population viability and

contractions in distributions may occur much more quickly than expected from currently

established critical thermal limits.

For populations of purple sea urchins (*Strongylocentrotus purpuratus* - an ecologically

important herbivore) in Southern California, recruitment tends to collapse during warm El Niño

events[20] (Figure 1A). This phenomenon is remarkably consistent over time (extending back at

least six decades), even though temperatures during the reproductive season in southern

California (late fall and early winter [20,21] Figure 1B) are almost always below those considered to

be lethal for adults (~25 °C[22]). Moreover, winter temperatures in the Santa Barbara Channel

when larvae are present (~ 11-16°C) are generally well below the thresholds thought to impede

multiple phases of larval survival[23] (fertilization and early-phase survival of San Diego, Santa

Barbara, and British Columbia animals declines from high to low between 20-22°C - SK,

unpublished data). Specifically, purple urchins tend to spawn in the winter and spring, coinciding

with favorable (low temperatures and high food availability) conditions for larvae [20]. Larvae

persist in the plankton from approximately 28 days to several months depending on food

availability and temperature [24,25]. Adult urchins build gonads over the course of the summer and

fall with gametogenesis occurring in the fall and early winter. El Niño conditions in Southern

California are characterized by warmer water temperatures in summer, fall, and winter than

during neutral years or during La Niña conditions, but still exhibit seasonal cooling (Figure 1B).

One proposed explanation for decreases in recruitment in southern California during El Niño is

that temperatures experienced during these events suppress gametogenesis even though energetic

investment in gonads may be sufficient for reproduction [26-28]. Specifically, Pearse [28]

demonstrated that high temperatures decrease gamete production at sub-lethal levels (i.e., at

21°C vs 14°C), but these high temperatures exceed the extreme values experienced in the

spawning season.

In this study, we used *S. purpuratus* to test whether historical collapses in larval supply

and year-class strength (i.e., abundance for a given young-of-year cohort) can, in part, be

explained by sub-lethal thermal suppression of reproduction. Specifically, we used laboratory

experiments to ask whether dynamic marine heatwaves that have led to collapses in larval supply

also affect reproduction in terms of both gonad production and gametogenesis. We compared

effects of simulated heatwaves to simulated historical cool trends, as well as the outcomes under

a gradient of constant conditions (i.e., the thermal performance under constant temperatures).

Results

Simulated El Niño and warming conditions led to decreased gonad production as well as

suppressed gametogenesis in *S. purpuratus* with no mortalities associated with any treatments.

Declines in gonad production associated with high temperatures were consistent for constant

exposure to 20°C and the El Niño treatment that shifted from 21 to 18°C with a mean of 20°C

(Figure 4). In contrast, the dynamic La Niña treatment (shifting from 18 to 14°C, with a mean of

16°C) led to reduced gonad production compared to a constant mean temperature of 16°C. The

effect of treatment on reproductive status differed depending on whether the thermal

environment varied over time (El Niño and La Niña versus constant temperatures) and by sex.

With a mean of 16°C, we saw no effect of dynamic La Niña or constant temperatures on male or

female reproductive status. For females, both constant high temperatures and the El Niño

treatment led to a substantial reduction in oogenesis; for males, only the dynamic El Niño

treatment affected spermatogenesis, with no change observed across constant temperatures from

10 to 20°C. Below, we detail these results in terms of gonad production and sex-specific

gametogenesis, comparing effects among the constant temperature treatments and dynamic El

Niño and La Niña treatments.

Gonad production

Exposure to the highest temperatures (for both dynamic and constant treatments with

mean of 20°C) negatively affected both male and female gonad production (cumulative gonad

mass, Figure 4) with a 42% decrease (95% highest probability density interval [HPD]: 28-55%)

compared to a constant temperature of 16°C. Gonad production did not differ for the El Niño

(7.03g [95% HPD: 6.20g – 7.96g]) and constant treatment that shared the same mean value of

20°C (7.2g, [95% HPD: 5.97g – 8.50g]). In contrast, the dynamic La Niña treatment (declining

from 18 to 14°C with mean of 16°C) showed a reduction in gonad mass (26% - HPD: 15 – 38%)

compared with a constant value of the same mean (16°C - 8.69g [95% HPD: 7.68g– 9.77g] vs

11.84g [95% HPD 10.71g – 13.15g], respectively). The lowest temperature imposed a smaller,

but still significant reduction (22% reduction, 95% HPD: 4-39%) compared to that at 16°C. The

disparity in gonad production for the El Niño and La Niña was much smaller than when

comparing constant temperature treatments of 20°C and 16°C despite having the same mean

temperature differential. This reduced effect resulted in part from the aforementioned decline in

gonad size in La Niña versus equivalent but constant mean. Model comparison showed >99%

stacking weight for the temperature and treatment effects, and no support for differences among

sexes in the gonad mass overall or interaction among treatment effects with sex.

Reproductive status

The El Niño treatment led to a significant reduction in gametogenesis for both males and

females. Specifically, far fewer females in the El Niño treatment reached Stage IV compared to

La Niña treatment for both males and females (Figure 5). Yet this effect was more extreme for

females compared to males. Female urchins incubated in La Niña treatment actively produced

mature gametes (89% [95% HPD: 77-94%] of females in stage IV, Figure 5 A). In contrast, only

15% [95% HPD: 4-28%] of El Niño females were in stage IV (Figure 5A), while most females

were in stage I. This decline in reproductive status associated with El Niño amounted to an

estimated 74% reduction for females compared to La Niña (Figure 5 A, HPD: 56-91% decline in

females in stage IV). For male urchins, these results were consistent but less extreme. 92% of La

Niña males were stage IV [HPD: 77 - 99%] vs 44% of El Niño females [HPD: 23 -67%]– a

reduction of 48% [HPD: 23 – 72%] - Figure 5 C). Results were consistent across the four

replicate mesocosms for each treatment (Figure 5 C).

Our results also indicate that temperature dynamics, and not just their mean values, can

alter reproductive response depending on sex. Simulating a dynamic, historical El Niño trend

resulted in a reduction in the proportion of males investing in spermatogenesis in contrast to no

shift under a constant mean temperature. Specifically, male animals subjected to the El Niño

treatment (simulated seasonal peak and decrease from 21 to 18°C with a mean of 20°C)

compared to a constant 20°C throughout showed substantially lower rates of full maturity (stage

IV) with a reduction from 81.2 to 47.8% (a difference of 37% [95% HPD: 8% to 65%], Figure 5

C, D). In contrast, we observed similarly low rates of full maturity for females subjected to the El

Niño simulation when compared to constant 20°C (15% for El Niño vs 25% for constant 20°C,

Bayesian $P = 0.28$, (Figure 5A, B). These outcomes are supported by a sex by treatment

interaction on top of the temperature by sex estimates (>99% stacking weight from loo cross

validation for the full model compared to models without any of the treatment, sex, or interaction

terms).

Discussion

Global warming and extreme heatwaves present a significant threat to ecosystems

worldwide. It is well understood that range limits contract and abundances decline well before

lethal limits for entire populations are reached. Yet mechanistically, explaining how and why

such sub-lethal thermal regimes affect populations remains a challenge. In this study, we show

how thermal suppression of reproduction can, in part, provide a plausible explanation for

historical collapses in larval supply associated with warming events. These results provide two

key general conclusions. First, sublethal impacts of warming events on reproduction may lead to

recruitment failure and population declines before lethal upper thermal limits for individual life

stages are experienced. This insight is derived from the results of our lab experiments in the

context of our historical observations of larval supply. Second, dynamic temperature treatments

that reflect real world trends in historically observed heatwaves, in comparison to constant

temperatures, exert a more pronounced effect on reproductive development; thus, ignoring the

dynamics of thermal regimes may underpredict the magnitude of thermal effects. Importantly,

these results raise questions regarding physiological processes by which temperature shapes

gametogenesis, and how such effects may vary in and among populations over time. These

effects will almost certainly interact with other stressors such as food availability for adults sensu

[29] or shifts in physical dispersal, gametogenesis, or heatwave conditions.

Heatwave events in both marine and terrestrial settings are often highly dynamic,
characterized by rapid fluctuations in temperature. Thus, experiments that utilize constant
temperatures ignoring such fluctuations may yield erroneous insights. Here, we show that male
but not female stages of maturity differed between constant versus El Niño temperature profiles
that shared the same mean value (20°C). This outcome also yields insight into a possible fitness
benefit from suppressing gametogenesis during El Niño. Early-stage larval viability (within 48
hours) declines rapidly between 20 to 22°C (SK unpublished data for these subjects as well as
for animals from San Diego). Thus, males in the El Niño treatment incubated briefly at 21°C at
the onset of the experiment may delay production of mature sperm until temperatures have
consistently dropped well below potential thresholds for larval mortality, even though mean
temperatures for all treatments at the end of the experiment, and in nature, lie well below such
values. Because male gametogenesis may be completed more quickly, these disparities may
reflect a more rapid reaction time to decreases in temperature. Females exposed to a constant
18°C proceeded with vitellogenesis and produced mature gametes; those subjected to El Niño
treatment that ended at 18°C did not. In contrast, overall gonad production (mass) differed
between La Niña and constant temperatures but not for El Niño. These disparities in gonad
production appear to result from nonlinear averaging (e.g., Jensen's inequality) associated with
dynamics of feeding, metabolism, and/or structural growth allocation [30]. Importantly, our study
ended in December, when temperatures approach winter lows (Figure 1). Even in El Niño years,
winter temperatures are likely sufficient to facilitate reproduction over the long term (Figure 1).
Thus, rather than inducing skip-spawning, thermal suppression of gametogenesis in fall and early
winter may instead lead to a shortened spawning season. Because urchin settlement timing aligns
with peak phytoplankton production [20], such effects may lead to other negative impacts on
recruitment such as match-mismatch dynamics [31]. Moreover, our dynamic treatments are limited
to a specific set of trajectories and our constant treatments do not include 21°C or above (the
maximum temperature of the El Niño treatment); thus, future work may benefit from assessing
how brief or consistent incursions into the 21°C range, and the timing of such incursions, affect
male sperm production. Thus, while our study highlights a single process that can plausibly
contribute to recruitment failure, it may be one of many interacting factors that shape temporal
trends in larval supply. As a result, future studies may benefit from documenting gonad

production and stages of maturity over time to assess how and when gamete production proceeds
under different heatwave conditions both in the field and in controlled settings.

While our experiments were conducted using sea urchins from British Columbia (where
waters are cooler than in Southern California), the results are consistent with prior insights from
across California. Specifically, field observations in southern California indicated reproduction
(assessed via seasonal gamete extrusion [26] and histology[27]) may have been suppressed during
analogous high temperature periods, and Cochrane and Engelman[26] suggested that 17°C was the
approximate threshold above which gametogenesis may be suppressed. A prior experiment by
Pearse[28] in Monterey Bay, California showed that a constant 21°C suppressed male and female
gametogenesis compared to constant 14°C. We show that for females, this suppression appears
to occur above 18°C but at or below 20°C. Importantly, these results align with the thermal
limits of early-stage larvae for animals from both British Columbia and San Diego, California
(survival of larvae 48 hr. post fertilization, prior to feeding, declines from near 100% to near 0%
in the 20-22°C range - SK unpublished data). Overall, these data highlight a need for further
study focused on the processes that regulate suppressed or delayed reproduction. Specifically, do
such responses vary by population or geographic region? How do depth, signal amplitude, or
acclimatization affect these outcomes? Moreover, how reproductive phenology varies in space
and over time has yet to be examined across the range from Baja California to Alaska. Evidence
for such variation exists in response to other stressors for this species e.g., [32]. Thus, we argue our
experiments demonstrate a thermal sensitivity that represents a plausible explanation for
historical collapses in larval supply. Yet examining the genetic and plastic basis of how
reproduction may respond to changes in temperature regime remains an important and
unanswered question both in this species and more broadly.

These experiments present plausible evidence for how and why historical collapses in
larval supply occur during marine heatwaves. Yet there are likely a myriad of factors that occur
during marine heatwaves that may shape recruitment in addition to, or in synergy with
suppression of reproduction. First, food stress occurs during El Niño events in southern
California [33]. Such trends might exacerbate the effects of temperature on gamete production as
food limitation for adults may alter thermal energetic performance (e.g., via "metabolic
meltdown" [29]). In these experiments, we focused on well-fed adult animals to avoid confounding

effects of food availability. But the results shown here raise the question of whether declines in

availability and productivity of macroalgae can exacerbate responses of gametogenesis. Second,

El Niño events are thought to be coupled with shifts in ocean circulation that might alter patterns

of phytoplankton production (i.e., larval food) as well as the delivery of planktonic larvae to

suitable settlement sites. These trends may also affect reproduction as urchin spawning may be

sensitive to changes in phytoplankton-derived chlorophyll [34,35]. Thus, even in the cases where

offspring are produced from successful maturation and spawning, shifts in larval survival and

delivery may further compound effects of heatwaves. Finally, we conducted these experiments

with healthy animals with no evidence of disease, yet temperature [36] in tandem with density may

increase disease prevalence and disease related mortality [37,38]. Therefore, our results focus on one

key vital rate among many that may simultaneously respond to abiotic and biotic conditions

during extreme events known to alter dynamics of populations.

Marine heatwaves have led to substantial reorganization of ecosystems. This trend has

been particularly apparent in recent years in temperate rocky reefs [39-45] with resulting, negative

effects on ecosystem services and biodiversity [46]. While extreme events can impose direct and

observable effects on mortality, sub-lethal effects, when present, may be far more frequent and

insidious because they occur below lethal thresholds and may have less immediately visible or

observable impacts e.g., [19]. This study is among the first to show that widespread, historical

collapses in larval supply in the field can partially and plausibly be explained by sub-lethal

suppression of gametogenesis resulting from marine heatwaves. Moreover, our study

demonstrates the value of long-term population studies and quantifying non-lethal physiological

and reproductive sensitivities when considering thermal envelopes and the population-level

impacts of global warming and heatwaves.

Methods

To quantify how different thermal regimes affect investment in gonads and development

of gametes in male and female urchins, we conducted a large-scale 10-week experiment in which

300 animals were incubated in replicate 340L mesocosms that simulated dynamic El Niño (N = 4

mesocosms, 60 animals per treatment) or La Niña (N = 4 mesocosms, 60 animals) conditions

based on historical, empirical benthic temperature time series from Scripps Pier in La Jolla,

California[47] (trends: Figure 1B, map: Figure 1C) that coincide with historical collapses in larval

supply in Southern California. We paired these treatments with a range of fixed temperature

incubations (10, 13, 16, 17, 18, 20 °C, N = 2 mesocosms, 30 animals per treatment), two of

which matched the mean temperature of the El Niño (20 °C) and La Niña (16 °C) (Figure 1). We

chose this benthic time series rather than satellite derived sea surface temperature information

because the latter provides high quality spatial representations of sea surface temperature but not

at depths where subtidal animals reside [48]. Experiments were conducted at the Marna Lab at the

Hakai Institute's Quadra Island Ecological Observatory in Heriot Bay, British Columbia due to

availability of sophisticated seawater systems for careful, replicated temperature manipulations.

Field Collections and Acclimation

We collected sea urchins by hand on SCUBA in the vicinity of Ucluelet, British

Columbia, Canada (48.94˚N, 125.56˚ W) from a depth of 20-25 ft relative to mean tide in

September 2021 and transported them to the Marna Lab. We transferred sea urchins to flow-

through sea tables and allowed them to recover for a period of one week before inserting animals

into the mesocosm system. We selected healthy individuals within a constrained size range for

incubations (n = 300, mean test diameter = 56.09 mm, range test diameter = 42.12 – 69.46 mm).

Finally, we assigned animals to mesocosms at random at ambient temperature and exposed each

assigned mesocosm to a temperature ramp, where the ramp reached target temperatures after two

weeks from the initial incoming, ambient temperature (mean across all tanks of 13.3˚C, SD =

0.3˚C) to avoid thermal shock. Once initial target temperatures were reached, they were

maintained or, for the dynamic treatments, were manually adjusted daily in the AM (~8am each

day) as needed by 0.5 °C increments in a scheduled manner to match historical mean El Niño and

La Niña daily temperature conditions.

Mesocosm System

We placed urchins in a custom-built array of twenty replicated 340 L acrylic mesocosms

supplied with flow-through UV sterilized and filtered seawater (Figure 3: , Integrated Aqua

Systems, Inc., Vista, CA, USA). Each mesocosm was capable of independent control of

temperature and animals were provided a lighting regime for all mesocosms using LED fixtures

(Aquamaxx, CA, USA) programmed to provide 10L:14D with two-hour linear light intensity

transition periods for dawn and dusk (0-100% from 07:00 to 09:00 "dawn", and 100-0% from

17:00 to 19:00 "dusk"). Each mesocosm independently maintained temperature treatments using

a heat exchanger fitted with a titanium coil regulated by a dual stage digital temperature

controller (Resolution = 0.1℃, Dwyer Instruments, LLC.©, Michigan City, IN, USA). The

mesocosm system employed central cooling (Aermec Mits Airconditioning Inc., Mississauga,

ON, Canada) and heating (boiler array, Viessmann Manufacturing Company Inc., Warwick, RI,

USA) to supply independent heat exchangers with on-demand cold and warm glycol loops for

down- and up-regulation of water temperature, respectively. We manually checked and re-

calibrated sensors, as needed, using digital traceable thermometers twice daily to control

potential temperature sensor drift. We randomly assigned mesocosms to the specified treatments.

Animal husbandry

We fed individuals uniform dry pellets combining several macroalgal species formulated

for the aquaculture of *S. purpuratus* (Urchinomics Canada Inc., Halifax, NS, Canada). Animals

in mesocosms were fed twice per week, and we removed uneaten food and refuse every 72 h.

Food rations were determined by trial and error in a prior pilot study to ensure all animals had

consistent access to food over time. To optimize access for all mesocosm inhabitants to abundant

food, we enclosed subjects and food in aquaculture baskets (two baskets per mesocosm, 7 or 8

animals per basket, Thunderbird Plastics) such that food was always readily accessible, and

movement was not impeded. Each animal was supplied approximately 2.7 grams of pelleted food

twice per week (either 19 or 21 grams per basket for the baskets with 7 and 8 individuals,

respectively in each mesocosm) for the duration of the experiment.

Histological and gonad assays

At the end of the experiment, we measured all individuals to test for changes in height

and diameter (using precision digital calipers) and wet mass (to the nearest 0.1 g). Animals were

then sacrificed to measure gonad and histological properties. After cracking urchin tests, we

rapidly removed gonads for sampling. We excised a second gonad from each animal, which was

weighed to the nearest 0.01g and categorized the remaining gonads visually for evidence of

gamete production (macroscopic egg extrusion or sperm extrusion). Using a clean, sterile scalpel

we excised an approximately 2mm cross section from one gonad which we immediately placed

in a histological cassette, preserved in Hartmann's fixative for 24 hours, and transferred to 70%

EtOH. Preserved gonads were embedded in paraffin, sliced, stained using eosin and hematoxylin,

and mounted. We assessed gonad samples for sex and developmental stage using four visual

subsections and the entire sample collectively to ensure agreement among subsamples.

Histological slides were scored on a scale of I to IV (sensu [49]), where representative stages are

depicted in Figure 1.

Statistical Analyses

We estimated the probability of individuals in each reproductive stage as a function of

mean temperature by sex and in response to El Niño and La Niña treatments using a Bayesian

regression model with a multinomial likelihood. We modeled the response to mean temperature

using a non-parametric Gaussian process smoother. To account for treatment effects, we

included a categorical effect of treatment (constant, El Niño, La Niña). Gonad mass was modeled

with a Gamma likelihood and log link with the log of test diameter included as a covariate to

account for gonad-body size allometry. Because gonad mass was expected to exhibit a concave

response to temperature, a 3rd order polynomial was used for mean temperature. For both

analyses, intercepts were allowed to vary randomly by mesocosm.

We directly compared parameter posteriors within models and to ensure that comparing

parameters within models was robust to model (structural) uncertainty, we conducted model

comparison using approximate leave-one-out cross validation (loo) and generated weights using

stacking [50-52]. These methods measure the predictive accuracy with a loo approximation and

generate weights by generating weights of models that maximize the predictive accuracy of the

potential models. The selected models represented >99% of posterior model weight and thus

results presented are from these models. We sampled posteriors using brms in R [53,54]. We ran

models with vague priors, and conducted sensitivity analyses to ensure varying prior

distributions and hyperparameters did not qualitatively affect the results. We ran models for

2000 iterations for each of four chains after a 2000 iteration warmup period, and checked

convergence visually and ensuring that that parameters had convergence diagnostics (Rhat) less

than 1.001. Data and code from this study are available at https://www.bco-

dmo.org/project/818918.

Author contributions:

DKO, NBS, BC, LRB and SS designed the research. DKO, NBS, BC, MJM, SK, KR, DS, IG, EC, MF, and NM conducted mesocosm experiments. RF conducted histological analyses. DKO conducted statistical analyses. DKO wrote the initial draft of the manuscript, and all authors contributed to revisions.

Acknowledgements:

This study was funded with generous support from the Hakai Institute, a program of the Tula Foundation, a grant from the US National Science Foundation to DKO and LRB (OCE nos. 2023649 and 2023664), as well as from the Santa Barbara Coastal (SCB) LTER (NSF OCE no.1831937). We thank Eric Peterson, Christina Munck, as well as the facilities staff at the Quadra Island Ecological Observatory for their generous support. We thank the Coastal Marine Science Inst. and Bodega Marine Lab at UC Davis as well as the California Dept. Fish and Wildlife LRB. Rachel Simons, Wiley Evans, and two anonymous reviewers provided useful feedback on prior versions of the paper.

Figure 1: A) Historical mean annual (large points) and monthly mean (small points) larval settlement on standardized brush samplers across six sites in southern California vs Multivariate El Niño Index (MEI) since 1990 (reproduced from Okamoto et al.[20]). Colors correspond to the x-axis representing cooler (negative) and warmer (positive) temperatures associated with the MEI in Southern California. B) Historical benthic temperatures from Scripps Pier [47]. Small points: daily means. Black line: historical seasonal mean. Red and blue lines: historical El Niño & La Niña seasonal trends, respectively, from the data that were simulated in mesocosms (the period simulated in the experiment is indicated by the vertical grey bar). Horizontal dotted lines: constant mean temperatures simulated in replicate mesocosms. C & D) Mean satellite derived sea surface temperature from September through December in four strong (C - 1993, 1996, 1999, & 2010) and poor (D - 1994, 1997, 2004, & 2015) years for larval settlement from 1990 to 2016 [48]. Note that C & D are visualized to show comparative spatial trends, though satellite-derived temperature data may be biased relative to in situ benthic temperature data. Sites in C represent the six historical settlement collection locations that all show negative correlations with the MEI.

Figure 2: Gonad sections stained with hematoxylin and eosin with scale bar representing
500 μm. A-B Female ovaries in Stage I (A) and Stage IV (B) arrows indicate small
previtellogenic oocytes, vitellogenic oocytes in the process of meiosis, and mature ova. C-D.
Females in stage I are largely devoid of developing and vitellogenic oocytes with substantial
reserves invested in nutritive phagocytes (light pink), Stage II has few developing oocytes, Stage
III has some developing oocytes with few, scattered mature eggs, and Stage IV has many mature
eggs (dark, solid circles) and some developing oocytes (dark circles with visible, central
germinal vesicle). Male testes showing stage I (C) with few developing columns of
spermatocytes, and stage IV (D) with large sections of the gonad converted to mature
spermatozoa. Males in stage I have few visible developing spermatocytes (columns of dark
purple around the margins), males in stage II have dense columns of spermatocytes, males in
stage III have few small pockets of spermatozoa, and males in stage IV have large pockets of
spermatozoa and rapidly disappearing somatic tissue (light pink).

Figure 3: A) Mesocosms at the Hakai Institute's Quadra Island Ecological Observatory with
trays holding experimental subjects. B) Experimental subjects consuming controlled,
manufactured Urchinomics algal feed pellets.

400   .

402   Figure 4: Final gonad mass vs mean temperature. Black circles and the smooth trend represent
403   fixed temperature treatments; the red diamond and blue square represent La Niña and El Niño
404   thermal treatments, respectively. Error bars and bands represent the 80, 90, and 95% highest
405   probability density interval of the posterior.

Figure 5: Proportion of adult females (A, B) and males (C, D) in each reproductive stage by
treatment (El Niño (squares), La Niña (diamonds), or constant temperature (circles)) and mean
temperature. A) Proportion of females in stage IV. B) Proportion of females in stages I (dotted
red lines, red points), II (dashed grey lines, grey points, and III (solid purple lines, purple points).
C) Proportion of males in stage IV, and D) proportion of males in stages I, II, and III (same
symbology as females). Circles represent empirical means for constant temperature treatments,
while El Niño and La Niña points represent posterior mean estimates. Constant temperature
treatments had two replicate mesocosms per treatment and 15 animals per mesocosm. El Niño
and La Niña had four replicate mesocosms and 15 animals per mesocosm. Uncertainty
bands/intervals represent 95% (light/thin) and 90% (dark/thick) from the multinomial regression.

References

1 Braschler, B., Chown, S. L. & Duffy, G. A. Sub-critical limits are viable alternatives to
critical thermal limits. *J Therm Biol* **101**, 103106 (2021).
https://doi.org:10.1016/j.jtherbio.2021.103106 PMID - 34879920

2 Kinzner, M. T. *et al.* Is temperature preference in the laboratory ecologically relevant for
the field? The case of *Drosophila nigrosparsa*. *Global Ecol Conservation* **18**, e00638
(2019). https://doi.org:10.1016/j.gecco.2019.e00638

3 Rezende, E. L., Bozinovic, F., Szilágyi, A. & Santos, M. Predicting temperature mortality
and selection in natural *Drosophila* populations. *Science* **369**, 1242-1245 (2020).
https://doi.org:10.1126/science.aba9287 PMID - 32883867

4 Oliver, E. C. *et al.* Marine heatwaves off eastern Tasmania: Trends, interannual
variability, and predictability. *Progress in oceanography* **161**, 116-130 (2018).

5 Munday, P. L., Jones, G. P., Pratchett, M. S. & Williams, A. J. Climate change and the
future for coral reef fishes. *Fish and Fisheries* **9**, 261-285 (2008).

6 Terblanche, J. S., Deere, J. A., Clusella-Trullas, S., Janion, C. & Chown, S. L. Critical
thermal limits depend on methodological context. *Proc Royal Soc B Biological Sci* **274**,
2935-2943 (2007).

7 Inouye, D. W. Climate change and phenology. *Wiley Interdiscip Rev Clim Change* **13**
(2022). https://doi.org:10.1002/wcc.764

8 Amasino, R. Vernalization, Competence, and the Epigenetic Memory of Winter. *Plant
Cell* **16**, 2553-2559 (2004). https://doi.org:10.1105/tpc.104.161070 PMID - 15466409

9 Kim, D.-H. Current understanding of flowering pathways in plants: focusing on the
vernalization pathway in *Arabidopsis* and several vegetable crop plants. *Hortic Environ
Biotechnology* **61**, 209-227 (2020). https://doi.org:10.1007/s13580-019-00218-5

10 Rogers-Bennett, L., Dondanville, R. F., Moore, J. D. & Vilchis, L. I. Response of Red
Abalone Reproduction to Warm Water, Starvation, and Disease Stressors: Implications of
Ocean Warming. *J Shellfish Res* **29**, 599-611 (2010).
https://doi.org:10.2983/035.029.0308

11 Breckels, R. D. & Neff, B. D. The effects of elevated temperature on the sexual traits,
immunology and survivorship of a tropical ectotherm. *J Exp Biol* **216**, 2658-2664 (2013).
https://doi.org:10.1242/jeb.084962 PMID - 23531818

12 Paxton, C. W., Baria, M. V. B., Weis, V. M. & Harii, S. Effect of elevated temperature
on fecundity and reproductive timing in the coral *Acropora digitifera*. *Zygote* **24**, 511-516
(2016). https://doi.org:10.1017/s0967199415000477 PMID - 26349407

13 Hurley, L. L., McDiarmid, C. S., Friesen, C. R., Griffith, S. C. & Rowe, M. Experimental
heatwaves negatively impact sperm quality in the zebra finch. *Proc Royal Soc B
Biological Sci* **285**, 20172547 (2018). https://doi.org:10.1098/rspb.2017.2547 PMID -
29343605

14 Parratt, S. R. *et al.* Temperatures that sterilize males better match global species
distributions than lethal temperatures. *Nat Clim Change* **11**, 481-484 (2021).
https://doi.org:10.1038/s41558-021-01047-0

15 Okamoto, D. K., Schmitt, R. J., Holbrook, S. J. & Reed, D. C. Fluctuations in food
supply drive recruitment variation in a marine fish. *Proc Royal Soc B Biological Sci* **279**,
4542-4550 (2012). https://doi.org:10.1098/rspb.2012.1862 PMID - 23015631

16 Stoner, A. W. & Ray-Culp, M. Evidence for Allee effects in an over-harvested marine
gastropod: density-dependent mating and egg production. *Mar Ecol Prog Ser* **202**, 297-
302 (2000). https://doi.org:10.3354/meps202297

17 Gaines, S. D. & Bertness, M. D. Dispersal of juveniles and variable recruitment in sessile
marine species. *Nature* **360**, 579-580 (1992). https://doi.org:10.1038/360579a0

18 Kelly, M. S. Environmental parameters controlling gametogenesis in the echinoid
Psammechinus miliaris. *Journal of Experimental Marine Biology and Ecology* **266**, 67-
80 (2001).

19 Rogers-Bennett, L., Klamt, R. & Catton, C. A. Survivors of climate driven abalone mass
mortality exhibit declines in health and reproduction following kelp forest collapse.
Frontiers in Marine Science **8**, 725134 (2021).

20 Okamoto, D. K., Schroeter, S. C. & Reed, D. C. Effects of ocean climate on
spatiotemporal variation in sea urchin settlement and recruitment. *Limnol Oceanogr* **65**,
2076-2091 (2020). https://doi.org:10.1002/lno.11440

21 Ebert, T. A., Schroeter, S. C., Dixon, J. D. & Kalvass, P. Settlement patterns of red and
purple sea urchins (*Strongylocentrotus franciscanus* and *S. purpuratus*) in California,
USA. *Mar Ecol Prog Ser*, 41-52 (1994).

22 Farmanfarmaian, A. & Giese, A. C. Thermal tolerance and acclimation in the western
purple sea urchin, Strongylocentrotus purpuratus. *Physiological Zoology* **36**, 237-243
(1963).

23 Azad, A. K., Pearce, C. M. & McKinley, R. S. Influence of stocking density and
temperature on early development and survival of the purple sea urchin,
Strongylocentrotus purpuratus (S timpson, 1857). *Aquaculture Research* **43**, 1577-1591
(2012).

24 Azad, A. K., Pearce, C. M. & McKinley, R. S. Influence of microalgal species and
dietary rations on larval development and survival of the purple sea urchin,
Strongylocentrotus purpuratus (Stimpson, 1857). *Aquaculture* **322**, 210-217 (2011).

25 Strathmann, M. F. *Reproduction and development of marine invertebrates of the northern
Pacific coast: data and methods for the study of eggs, embryos, and larvae.* (University
of Washington Press, 2017).

26 Cochran, R. C. & Engelmann, F. Characterization of spawn-inducing factors in the sea
urchin, *Strongylocentrotus purpuratus*. *Gen Comp Endocr* **30**, 189-197 (1976).
https://doi.org:10.1016/0016-6480(76)90099-x PMID - 992340

27 Basch, L. V. & Tegner, M. J. Reproductive responses of purple sea urchin
(*Strongylocentrotus purpuratus*) populations to environmental conditions across a coastal
depth gradient. *Bulletin of Marine Science* **81**, 255-282 (2007).

28 Pearse, J. S. in *Advances in invertebrate reproduction*   (ed S. Adams W. H. Clark Jr)
(1981).

29 Huey, R. B. & Kingsolver, J. G. Climate warming, resource availability, and the
metabolic meltdown of ectotherms. *The American Naturalist* **194**, E140-E150 (2019).

30 Spindel, N. B. *Impacts of abiotic and biotic stressors on the energetics and nutritional
composition of temperate sea urchins* PhD thesis, Florida State University, ("In Prep").

31 Cushing, D. in *Advances in marine biology* Vol. 26   249-293 (Elsevier, 1990).

32 Kapsenberg, L., Okamoto, D. K., Dutton, J. M. & Hofmann, G. E. Sensitivity of sea
urchin fertilization to pH varies across a natural pH mosaic. *Ecology and Evolution* **7**,
1737-1750 (2017).

33 Tegner, M. & Dayton, P. Sea urchins, El Niños, and the long term stability of Southern
California kelp forest communities. *Marine ecology progress series. Oldendorf* **77**, 49-63
(1991).

34 Seward, L. C. The relationship between green sea urchin spawning, spring phytoplankton
blooms, and the winter-spring hydrography at selected sites in Maine. (2002).

35 Zhadan, P. M., Vaschenko, M. A. & Almyashova, T. N. Spawning failure in the sea
urchin Strongylocentrotus intermedius in the northwestern Sea of Japan: Potential
environmental causes. *Journal of Experimental Marine Biology and Ecology* **465**, 11-23
(2015).

36 Lester, S. E., Tobin, E. D. & Behrens, M. D. Disease dynamics and the potential role of
thermal stress in the sea urchin, *Strongylocentrotus purpuratus*. *Canadian Journal of
Fisheries and Aquatic Sciences* **64**, 314-323 (2007).

37 Behrens, M. D. & Lafferty, K. D. Effects of marine reserves and urchin disease on
southern Californian rocky reef communities. *Mar Ecol Prog Ser* **279**, 129-139 (2004).

38 Lafferty, K. D. Fishing for lobsters indirectly increases epidemics in sea urchins.
Ecological Applications **14**, 1566-1573 (2004).

39 Arafeh-Dalmau, N. *et al.* Extreme marine heatwaves alter kelp forest community near its
equatorward distribution limit. *Frontiers in Marine Science* **6**, 499 (2019).

40 Filbee-Dexter, K. *et al.* Marine heatwaves and the collapse of marginal North Atlantic
kelp forests. *Scientific Reports* **10**, 13388 (2020).

41 McPherson, M. L. *et al.* Large-scale shift in the structure of a kelp forest ecosystem co-
occurs with an epizootic and marine heatwave. *Communications biology* **4**, 298 (2021).

42 Michaud, K. M., Reed, D. C. & Miller, R. J. The Blob marine heatwave transforms
California kelp forest ecosystems. *Communications Biology* **5**, 1143 (2022).

43 Smale, D. A. Impacts of ocean warming on kelp forest ecosystems. *New Phytologist* **225**,
1447-1454 (2020).

44   Thomsen, M. S. *et al.* Local extinction of bull kelp (*Durvillaea* spp.) due to a marine
heatwave. *Frontiers in Marine Science* **6**, 84 (2019).

45   Wernberg, T. in *Ecosystem collapse and climate change*   325-343 (Springer, 2021).

46   Smith, K. E. *et al.* Socioeconomic impacts of marine heatwaves: Global issues and
opportunities. *Science* **374**, eabj3593 (2021).

47   Carter, M. L. *et al.*   (UC San Diego Library Digital Collections, 2022).

48   Reynolds, R. W. *et al.* Daily High-Resolution-Blended Analyses for Sea Surface
Temperature. *J Climate* **20**, 5473-5496 (2007). https://doi.org:10.1175/2007jcli1824.1

49   Byrne, M. Annual reproductive cycles of the commercial sea urchin *Paracentrotus*
lividus from an exposed intertidal and a sheltered subtidal habitat on the west coast of
Ireland. *Marine Biology* **104**, 275-289 (1990).

50   Vehtari, A., Gelman, A. & Gabry, J. Practical Bayesian model evaluation using leave-
one-out cross-validation and WAIC. *Statistics and computing* **27**, 1413-1432 (2017).

51   Yao, Y., Vehtari, A., Simpson, D. & Gelman, A. Using stacking to average Bayesian
predictive distributions (with discussion). (2018).

52   Vehtari A *et al.* loo: Efficient leave-one-out cross-validation and WAIC for Bayesian
models v. R package version 2.6.0. (2023).

53   Bürkner, P.-C. brms: An R package for Bayesian multilevel models using Stan. *Journal*
of statistical software **80**, 1-28 (2017).

54   R Core Team. R: A Language and Environment for Statistical Computing (R Foundation
for Statistical Computing, Vienna, Austria., 2023).