Main Manuscript for

The function of juvenile-adult transition axis in female sexual receptivity of

Drosophila melanogaster

Jing Li[1,2,†,*], Chao Ning[3,4,†], Yaohua Liu[2,5,†], Bowen Deng[6], Bingcai Wang[2,7], Kai Shi[2,7],
Rencong Wang[2,7], Ruixin Fang[1] and Chuan Zhou[1,2,7]

[1]*Institute of Molecular Physiology, Shenzhen Bay Laboratory, Shenzhen 518132, China.*
[2]*State Key Laboratory of Integrated Management of Pest Insects and Rodents, Institute*
of Zoology, Chinese Academy of Sciences, Beijing 100101, China.
[3]*National Laboratory of Biomacromolecules, New Cornerstone Science Laboratory,*
CAS Center for Excellence in Biomacromolecules, Institute of Biophysics, Chinese
Academy of Sciences, 100101, Beijing, China.
[4]*CAS Key Laboratory of Genome Sciences and Information, Beijing Institute of*
Genomics, Chinese Academy of Sciences, Beijing 100101, China.
[5]*Department of Plant Protection, Shanxi Agricultural University, Jinzhong 30801, China*
[6]*Chinese Institute for Brain Research, Peking-Tsinghua Center for Life Sciences,*
Zhongguancun Life Sciences Park, Beijing 102206, China.
[7]*University of Chinese Academy of Sciences, Beijing 100049, China.*
[†]*These authors contributed equally to this work*
----
*Jing Li, Email: lijing@szbl.ac.cn

Author Contributions: Jing Li, Conceptualization, Resources, Data curation, Writing -
original draft, review and editing, Funding acquisition; Chao Ning and Yaohua Liu, Data
curation; Bowen Deng, Resources; Bingcai Wang, Kai Shi and Rencong Wang,
Software; Ruixin Fang, Data curation; Chuan Zhou, Conceptualization, Funding
acquisition.

Competing Interest Statement: The authors declare no competing interest.

Impact statement: Prothoracicotropic hormone and ecdysone belonging to the insect
PG axis modulate virgin female sexual receptivity.

Funding information: This work was supported by the Institute of Molecular
Physiology, Shenzhen Bay Laboratory (NO.21260061 to Chuan Zhou, S239201006 to
Jing Li), National Natural Science Foundation of China (NO.Y711241133 to Chuan
Zhou) and Strategic Priority Research Program of the Chinese Academy of Sciences
(NO.Y929731103 to Chuan Zhou).

Abstract

Female sexual receptivity is essential for reproduction of a species. Neuropeptides play

the main role in regulating female receptivity. However, whether neuropeptides regulate

female sexual receptivity during the neurodevelopment is unknown. Here we found the

peptide hormone prothoracicotropic hormone (PTTH), which belongs to the insect PG

axis, negatively regulated virgin female receptivity through ecdysone during

neurodevelopment in *Drosophila melanogaster*. We identified PTTH neurons as

doublesex-positive neurons, they regulated virgin female receptivity before the

metamorphosis during the 3rd-instar larval stage. PTTH deletion resulted in the

increased EcR-A expression in the whole newly formed prepupae. Furthermore, the

ecdysone receptor EcR-A in pC1 neurons positively regulated virgin female receptivity

during metamorphosis. The decreased EcR-A in pC1 neurons induced abnormal

morphological development of pC1 neurons without changing neural activity. Among all

subtypes of pC1 neurons, the function of EcR-A in pC1b neurons was necessary for

virgin female copulation rate. These suggested that the changes of synaptic

connections between pC1b and other neurons decreased female copulation rate.

Moreover, female receptivity significantly decreased when the expression of PTTH

receptor Torso was reduced in pC1 neurons. This suggested that PTTH not only

regulates female receptivity through ecdysone but also through affecting female

receptivity associated neurons directly. The PG axis has similar functional strategy as

the HPG axis in mammals to trigger the juvenile–adult transition. Our work suggests a

general mechanism underlying which the neurodevelopment during maturation

regulates female sexual receptivity.

Keywords: prothoracicotropic hormone, *Drosophila*, female sexual receptivity,

ecdysone, pC1 neurons

Introduction

The success of copulation is important for the reproduction of a species. *Drosophila*

melanogaster provides a powerful system to investigate the neuronal and molecular

mechanism of sexual behaviors. Females decide to mate or not according to their

physiological status and the environmental condition (Dickson, 2008). Sexually mature

adult virgin females validate males after sensing the courtship song and male-specific

sex pheromone, receive courtship with pausing and opening the vaginal plate (Ferveur,

2010; Greenspan et al., 2000; Hall, 1994; Wang et al., 2021). If the female is not willing

to mate, she may kick her legs, flick her wings, or extrude the ovipositor to deter males

(Connolly et al., 1973). Mated females reject males for several days after mating mainly

through more ovipositor extrusion and less opening the vaginal plate (Fuyama et al.,

1997; Wang et al., 2021). These options need the establishment of neural circuits for

female sexual receptivity. However, the associated mechanism of neural maturation and

the effect of neural maturation on female sexual receptivity is little known.

doublesex (*dsx*) and *fruitless* (*fru*) are the terminal genes in sex determination

regulatory hierarchy. They specify nearly all aspects of somatic sexual differentiation,

including the preparation for sexual behaviors (Dickson, 2008; Manoli et al., 2013;

Manoli et al., 2006; Mellert et al., 2012; Pavlou et al., 2013; Siwicki et al., 2009;

Yamamoto, 2007; Yamamoto et al., 2013). In males, expression of male-specific Fru^M

(Billeter et al., 2006; Demir et al., 2005; Hall, 1978; Manoli et al., 2005; Stockinger et al.,

2005) and male-specific Dsx^M (Kohatsu et al., 2011; Pan et al., 2014; Pan et al., 2011;

Rideout et al., 2010) are important for male courtship behaviors. In females, although

functional Fru protein is not translated, neurons with Dsx^F or *fru* P1 promoter regulate

some aspects of the female sexual behaviors (Kvitsiani et al., 2006; Rideout et al.,

2010). *Fru and dsx* are involved in regulating the sexual dimorphism during

neurodevelopment (Yamamoto et al., 2013). For instance, the sexual dimorphism of P1

and mAL neurons which are all associated with male courtship and aggression

behaviors (Clowney et al., 2015; Hoopfer et al., 2015; Kimura et al., 2008; Kohatsu et

al., 2011; Pan et al., 2012; Sengupta et al., 2022; von Philipsborn et al., 2011) is the

result of regulation by Dsx and/or Fru (Ito et al., 2012; Kimura et al., 2008). In the cis-

vaccenyl acetate (cVA) pathway, which induces the courtship inhibiting in males

(Kurtovic et al., 2007; Wang et al., 2010), the first-order to the fourth-order components

are all fru-Gal4-positive neurons and are either male-specific or sexually dimorphic

(Ruta et al., 2010). However, the role of Dsx^F in neurodevelopment associated with

female sexual behaviors is little understood.

During postembryonic development, the PG axis triggers the juvenile–adult transition,

similar to the function of hypothalamic–pituitary–gonadal (HPG) axis in mammals

(Herbison, 2016; Pan et al., 2019). Hormones of the PG axis act to transform the larval

nervous system into an adult version (Truman et al., 2023). Ecdysone belonging to the

PG axis is the prime mover of insect molting and metamorphosis and is involved in all

phases of neurodevelopment, including neurogenesis, pruning, arbor outgrowth, and

cell death (Truman et al., 2023). The neurons read the ecdysteroid titer through two

isoforms of the ecdysone receptor, EcR-A and EcR-B1, according to spatial and

temporal conditions in the central nervous system (Riddiford et al., 2000; Truman et al.,

1994). EcR-A is required in fru P1-expressing neurons for the establishment of male-

specific neuronal architecture, and ecdysone receptor deficient males display increased

male–male courtship behavior (Dalton et al., 2009; Ganter et al., 2007). However, how

ecdysone regulates the neurodevelopment associated with female sexual receptivity,

especially the fru[+] and dsx[+] neurons, is unknown.

Much of studies to understand female sexual receptivity has focused on its regulation.

How a female respond to males is highly dependent on whether or not she has

previously mated. In virgin females, dsx[+] pCd neurons respond to the cVA, while dsx[+]

pC1 neurons also respond to male courtship song (Zhou et al., 2014). The receptive

females open the vaginal plate (VPO) through activation of the dsx[+] vpoDN neurons

(Wang et al., 2021). After mated, sex peptide in the seminal fluid binds to the fru[+] dsx[+]

sex peptide sensory neurons (SPSNs) in the female uterus. Then neuronal activity in

the dsx[+] sex peptide abdominal ganglion (SAG) neurons of the ventral nerve cord and in

the pC1 neurons is reduced (Avila et al., 2011; Feng et al., 2014; Häsemeyer et al.,

2009; Kubli, 2003; Wang et al., 2020b; Yang et al., 2009; Zhou et al., 2014). Therefore,

the sexual receptivity is reduced with less VPO and more ovipositor extrusion (OE)

which is controlled by dsx[+] DpN13 neurons (Wang et al., 2020a). In addition,

neuropeptides and monoamines play a critical role in regulation of the female

receptivity. The neuropeptides Drosulfakinin, myoinhibitory peptides and SIFamide are

involved in female sexual receptivity (Jang et al., 2017; Terhzaz et al., 2007; Wang et

al., 2022). As monoamines, dopamine, serotonin and octopamine are pivotal to female

sexual behaviors (Ishimoto et al., 2020; Ma et al., 2022; Neckameyer, 1998; Rezával et

al., 2014). So far, the identified neuropeptides and monoamines modulating female

sexual receptivity all function during the adult stage. However, whether neuropeptides

or monoamines regulate the establishment of neural circuits for female sexual

receptivity is unknown.

To explore the factors that regulate *Drosophila* virgin female receptivity especially

during neurodevelopment, we did a knock-out screen including most of CCT members.

We discovered a requirement for the prothoracicotropic hormone (PTTH) during

postembryonic development for virgin female receptivity. We also found that PTTH

neurons expressing PTTH are dsx[+] neurons. PTTH, a brain derived neuropeptide

hormone, is the primary promoter of the synthesis of steroid hormone 20-

hydroxyecdysone (20E) (McBrayer et al., 2007; Rewitz et al., 2009). Indeed, the

enhanced virgin female receptivity due to the loss of PTTH could be rescued through

feeding 20E to the 3[rd]-instar larvae. Because 20E acts through its receptor EcR

(Riddiford et al., 2000), we then tested the function of EcR in pC1 neurons which

encode the mating status of females (Zhou et al., 2014). The reduced EcR-A expression

in pC1 neurons resulted in the abnormal anatomical pattern of pC1 neurons and the

reduced female copulation rate. This may be explained by the increased EcR-A in newly

formed prepupae resulted from the PTTH deletion. Furthermore, the decreased female

copulation rate was due to the reduced EcR-A in pC1b neurons. Besides, we detected

the inhibited female receptivity when PTTH receptor torso was decreased in pC1

neurons, suggested the direct function of PTTH on other dsx$^+$ neurons to regulate

female receptivity. Thus, in addition to demonstrating the function of PTTH in virgin

female receptivity during neurodevelopment, our study identified the necessary role of

the normal pC1b neural morphology in virgin female receptivity.

Results

PTTH modulates virgin female receptivity

In *Drosophila*, neuropeptides and monoamines, belonging to the chemoconnectome

(CCT, the entire set of neurotransmitters, neuromodulators, neuropeptides, and their

receptors underlying chemotransmission) (Deng et al., 2019), play a critical role in

regulation of the female receptivity. To explore the factors that regulate virgin female

receptivity especially during neurodevelopment, we screened 108 chemoconnectome

(CCT) knock-out lines generated by the CRISPR-Cas9 system (Deng et al., 2019)

(unpublished data). The result showed that prothoracicotropic hormone (PTTH) might

regulate virgin female receptivity. The deletion mutant *Ptth^Delete* removed part of the 5'

UTR and almost all coding sequence and is a protein null (**Figure 1A**). We confirmed

the PTTH knock-out flies by using PCR analysis at the PTTH locus in genomic DNA

samples (**Figure 1B**), by using RT-PCR to identify the loss of PTTH transcripts in cDNA

samples (**Figure 1C**) and by detecting the immunoreactivity of PTTH in the central brain

(**Figure 1—figure supplement 1A**). Primers used are listed in **Supplementary File 1**.

PTTH immunoreactivity was found in the brain of wild-type and heterozygous flies

(**Figure 1—figure supplement 1A1 and 1A3**), but was absent in homozygous *Ptth^Delete*

flies (**Figure 1—figure supplement 1A2**). As the previous study, the *Ptth^Delete* larvae

lacking PTTH undergo metamorphosis with about 1 day delay compared with the wild

type control (Shimell et al., 2018) (data not shown). Besides, the *Ptth^Delete* adult male

and female flies had the significant increased weight than wild type flies (**Figure 1—**

figure supplement 1B). This is also consistent with that PTTH regulates developmental

timing and body size in *Drosophila* (Mcbrayer et al., 2007; Shimell et al., 2018).

To confirm the function of PTTH, we tested virgin female receptivity of *Ptth^Delete* female

flies. We found that the virgin female losing PTTH had significantly higher copulation

rate and shorter latency to copulation than wild type flies (**Figure 1D-G**). In addition, the

Ptth^Delete flies had higher copulation rate and lower latency to copulation compared to

heterozygous null mutant females within 2 days (**Figure 1D-E**) and within 3 days,

respectively (**Figure 1D-F**). The enhanced virgin female receptivity had no relationship

either with the attractivity or with the locomotion activity of virgin females (**Figure 1—**

figure supplement 1C-E). These results suggested that PTTH deletion regulates virgin

female receptivity in a dose-dependent manner. Female receptivity increases with the

increase of age after eclosion, not only for wild type flies but also PTTH mutants. At the

first day after eclosion (**Figure 1D**), maybe the loss of PTTH in *PTTH^Delete^/+* flies is not

enough for sexual precocity as *PTTH^Delete^/PTTH^Delete^*. At the second day after eclosion

and after (**Figure 1E-G**), the loss of PTTH in *PTTH^Delete^/+* flies is enough for sexual

precocity compared with wild type flies. However, After the 2^nd^ day of adult, female

receptivity of all genotype flies increases sharply. At the 3^rd^ day of adult and after,

female receptivity of *PTTH^Delete^/PTTH^Delete^* reaches the peak and the receptivity of

PTTH^Delete^/+ reaches more nearly to *PTTH^Delete^/PTTH^Delete^* when flies get older (**Figure**

1F-G). However, the overexpression through PTTH-Gal4>UAS-PTTH is also not

sufficient to change female receptivity (**Figure 1—figure supplement 2A**). Similarly,

decreased expression of PTTH through PTTH-Gal4>UAS-PTTH-RNAi or dsx-

Gal4>UAS-PTTH-RNAi did not result in the similar phenotype to that of

*PTTH^Delete^/PTTH^Delete^* (**Figure 1—figure supplement 2B-C**). It is possible that both

decreasing and increasing PTTH expression are not sufficient to change female

receptivity.

Furthermore, we carried out genetic rescue experiments to further confirm the function

of PTTH in modulating virgin female receptivity. We used the pan-neuronal driver elav-

Gal4 to drive UAS-PTTH expression in PTTH mutant background, although elav-Gal4

did not express in PTTH neurons (**Figure 1—figure supplement 2D**). We detected the

PTTH signals using PTTH antibody in the rescued female brains (**Figure 1—figure**

supplement 1A4). We found that neuron-specific expression of PTTH could restore

the enhanced copulation rate and shorter latency to copulation in $PTTH^{Delete}/PTTH^{Delete}$

virgin females (**Figure 1H**). Except for the projection of axons to PG gland, PTTH also

carries endocrine function to regulate light avoidance of larvae (Yamanaka et al., 2013).

The overexpressed PTTH in other neurons through elav-Gal4>UAS-PTTH may act on

the PG gland through endocrine function and then induce the ecdysone synthesis and

release. In summary, these results suggested that PTTH regulates virgin female

receptivity through ecdysone.

Dsx+ PTTH neurons regulate virgin female receptivity.

We used new Ptth-Gal4 and Ptth-LexA which inserts Gal4 or LexA sequence before the

stop codon of the *Ptth* gene (Deng et al., 2019) to label and manipulate PTTH neurons

expressing PTTH. The labeled neurons were the same as reported before (McBrayer et

al., 2007; Yamanaka et al., 2013), a pair of bilateral neurosecretory cells in the brain

directly innervating the prothoracic gland during the larval stage (**Figure 2A and Figure**

2—figure supplement 1A). The newly emerged flies had the similar anatomical pattern

with that of the larval stage (**Figure 2B and Figure 2—figure supplement 1B**).

However, while the prothoracic gland cells are gradually degenerating during pharate

adult development (Dai et al., 1991; Roy et al., 2018), the pattern of PTTH neurons

labeled by Ptth-Gal4 > UAS-mCD8GFP gradually could not be found before the 10th

hour after eclosion (**Figure 2—figure supplement 2**).

Most identified neurons associated with female sexual behaviors express *doublesex*

gene. We asked whether PTTH neurons are a part of the doublesex circuitry or not.

Double labeling of dsx-LexA and Ptth-Gal4 neurons (LexAop-tomato,UAS-stinger-

GFP/Ptth-LexA;dsx-Gal4/+) revealed that PTTH neurons are all doublesex-positive

(**Figure 2C**). We then used an intersectional strategy to visualize overlapped expression

between dsx-LexA and Ptth-Gal4 (*UAS > stop > myrGFP/+;LexAop2-FlpL,dsx-*

LexA/Ptth-Gal4). We observed all PTTH neurons with GFP signals (**Figure 2D**). These

results suggested that PTTH neurons are dsx$^+$ neurons. Furthermore, we wanted to

know whether Dsx^F regulates female receptivity in PTTH neurons. We decreased the

Dsx^F expression in PTTH neurons and did not detect significantly changed female

receptivity (**Figure 2—figure supplement 1C**). We supposed that PTTH neurons have

some relationship with other Dsx^F-positive neurons which regulate female receptivity.

We then analyzed whether PTTH neurons are involved in the modulation of virgin

female receptivity. First, we activated PTTH neurons transiently in adult virgin females

by driving the temperature-sensitive activator dTrpA1 (Hamada et al., 2008) using Ptth-

Gal4. PTTH neurons were activated at 29^{o}C compared with the control treatment at

23^{o}C. No significantly different copulation rate or latency to copulation was detected

(**Figure 2—figure supplement 3A-C**). This suggested that PTTH neurons do not

regulate virgin female receptivity during the adult stage.

To identify the detail time for the function of PTTH neurons in virgin female receptivity,

we inactivated PTTH neurons through kir2.1 under the control of the temporal and

regional gene expression targeting system (McGuire et al., 2004). When the experiment

was done at the pupal stage is the only situation when the controls were both different

from the experimental. (**Figure 2—figure supplement 1D**). However, when PTTH

neurons were inactivated during whole pupal or whole adult stages, virgin female

copulation rate did not change significantly (**Figure 2—figure supplement 1D**).

Furthermore, we activated PTTH neurons at different stages overlapping the

postembryonic larval developmental time using dTrpA1 (**Figure 3A**). Stage 1 was from

the 1st-instar larvae to six hours before the 3rd-instar larvae. Stage 2 was from six hours

before the 3rd-instar larvae to the end of the wandering larvae (the start of prepupa

stage). Stage 3 was from the start of prepupa stage to the end of the 2nd day of the

pupal stage. Stage 4 was from the end of the 2nd day of the pupal stage to the eclosion

of adults. The copulation rate did not change significantly when activating PTTH

neurons during the stage 1, stage 3 or stage 4 (**Figure 3B, 3D and 3E**). However, we

found the significant lower copulation rate and the longer latency to copulation only

when PTTH neurons were activated during the stage 2 (**Figure 3C and Figure 3F**). The

defected copulation was not due to a lower locomotion activity of virgin females (**Figure**

3G). Taken together, our findings indicated that the activity of dsx$^+$ PTTH neurons

negatively regulate virgin female receptivity during the stage from the start of the 3rd-

instar to the end of wandering stage.

PTTH modulates virgin female receptivity through ecdysone.

The 3rd-intar larval stage is the critical stage for the initiation of metamorphosis involving

the synthesis of ecdysone (Imura et al., 2020; Lavrynenko et al., 2015; Shimell et al.,

2018). To test whether PTTH regulates virgin female receptivity through regulating the

synthesis of ecdysone, we rescued the enhanced female receptivity by feeding 20E to

the 3rd-intar larval $Ptth^{Delete}$ flies. The enhanced copulation rate and shorter latency to

copulation of the $Ptth^{Delete}$ flies were rescued to the comparable level of wild type

females (**Figure 4**). Furthermore, the wild type females fed by 20E had no significantly

different copulation rate and latency to copulation compared with the wild type females

fed by the same volume of 95% ethanol which is the solvent of 20E (**Figure 4**). This

suggested that PTTH regulates virgin female receptivity through the titer of ecdysone.

Ecdysone receptor EcR-A in pC1 neurons regulates virgin female copulation rate.

Given that PTTH regulates virgin female receptivity through ecdysone which acts on its

receptor EcR, we then asked whether ecdysone regulates the function of neurons

associated with virgin female receptivity through EcR. pC1 and vpoDN neurons are two

main dsx$^+$ neurons involved in virgin female receptivity (Wang et al., 2021; Zhou et al.,

2014). pC1 neurons encode the mating status of female flies, vpoDN neurons regulate

the opening of vaginal plate when females attend to accept males. EcR-A and EcR-B1

are the two prominently expressed ecdysone receptors in the CNS (Riddiford et al.,

2000). First, we tested the expression of EcR-A and EcR-B1 in these two neurons on

the 2nd day of the pupal stage when ecdysone functions as the main mover in the

metamorphosis (Dalton et al., 2009; Truman et al., 1994). The GFP signals labeled by

pC1-ss2-Gal4 and vpoDN-ss1-Gal4 were merged well with the signals of both EcR-A

and EcR-B1 antibodies respectively (**Figure 5—figure supplement 1**). This revealed

that EcR-A and EcR-B1 express in both pC1 and vpoDN neurons. We then tested the

function of EcR in pC1 and vpoDN neurons through reducing the expression of EcR-A

and EcR-B1 respectively. We used the split-Gal4 for pC1 and vpoDN neurons to drive

the UAS-EcR-RNAi. First, we reduced the expression of all EcR isoforms in pC1

neurons, this decreased the copulation rate and prolonged the latency to copulation

significantly (**Figure 5—figure supplement 2A**). Furthermore, we reduced the

expression of EcR-A in pC1 neurons. The virgin female had the significant lower

copulation rate and longer latency to copulation (**Figure 5A**). The reduced copulation

rate had no relationship with the attractivity (**Figure 5E**) and the locomotion activity of

virgin females (**Figure 5—figure supplement 2B**). When reducing the expression of

EcR-B1 in pC1 neurons, virgin females had the significant longer latency to copulation

but the comparable copulation rate to controls (**Figure 5B**). However, reducing the

expression of EcR-A (**Figure 5—figure supplement 3A-C**) and EcR-B1 (**Figure 5—**

figure supplement 3D-F) using three split vpoDN-Gal4s in vpoDN neurons all did not

affect virgin female receptivity. This suggested that the expression of EcR-A in pC1

neurons regulates virgin female copulation rate, but EcR isoforms in vpoDN neurons do

not modulate virgin female receptivity.

Two split-Gal4 drivers for pC1 neurons had been obtained previously. pC1-ss1-Gal4

labels pC1-a, c and -e neurons, and pC1-ss2-Gal4 labels all pC1-a, -b, -c, -d and -e

neurons (Wang et al., 2020b). We also tested virgin female receptivity when EcR-A or

EcR-B1 were reduced in pC1-a, -c and -e neurons simultaneously using pC1-ss1-Gal4

respectively. While the copulation rate or the latency to copulation did not change

significantly (**Figure 5C-D**). This suggested that, pC1b only, or both pC1b and pC1d

neurons is necessary for the functions of EcR-A and EcR-B1 in pC1 neurons on virgin

female receptivity. Whether pC1d is involved in the regulation of female receptivity is

uncertain (Deutsch et al., 2020; Schretter et al., 2020; Taisz et al., 2023). However,

when reducing EcR-A in pC1d neurons alone using the specific split-Gal4 SS56987

(Schretter et al., 2020), virgin female receptivity including copulation rate and latency to

copulation did not change significantly compared with controls (**Figure 5—figure**

supplement 4). These results suggested that the function of EcR-A in pC1b neurons is

necessary for virgin female copulation rate.

As recently mated females may reduce sexual receptivity and increase egg laying (Avila

et al., 2011; Kubli, 2003). we asked whether the decreased copulation rate induced by

EcR-A could be a post-mating response and correlate with elevated egg laying. To

address this, we examined the number of eggs laid by virgin females when EcR-A was

reduced in pC1 neurons. We found that manipulation of EcR-A did not enhance egg

laying significantly in virgin females (**Figure 5F**), although the UAS-EcR-A-RNAi control

causes a massive decrease in female fertility. Meanwhile, we further analyzed whether

reduction of EcR-A in pC1 neurons regulates the opening of vaginal plate (VPO) or the

ovipositor extrusion (OE). We found that reducing the EcR-A expression in pC1 neurons

lead to the significantly less VPO and more OE (**Figure 5G-H**). These results suggested

that reduced EcR-A expression in pC1 neurons results in the similar phenotype to that

of mated females.

EcR-A participates in the morphological development of pC1 neurons.

EcR isoforms have distinct temporal and spatial expression patterns in the CNS

(Riddiford et al., 2000; Truman et al., 1994). It is unknown when EcR-A functions in pC1

neurons for virgin female receptivity. Thus, we examined virgin female receptivity when

EcR-A expression was conditionally reduced through RNAi via the pC1-ss2-Gal4 under

the control of the temporal and regional gene expression targeting system (McGuire et

al., 2004). EcR-A was reduced during the whole larval, pupal and adult stage

respectively (**Figure 5I-K**). When the experiment was done at the pupal stage is the

only situation when the controls were both different from the experimental (**Figure 5J**).

The result suggested that EcR-A in pC1 neurons plays a role in virgin female receptivity

during metamorphosis. This is consistent with that PTTH regulates virgin female

receptivity before the start of metamorphosis.

We then tested how EcR-A functions in pC1 neurons to modulate virgin female

receptivity. First, we tested the morphology of pC1 neurons when reducing the

expression of EcR-A in pC1 neurons. We found that the morphology of pC1-ss2-Gal4

expressing neurons appeared after the formation of the white pupa (Figure **6A1**). The

reduced EcR-A expression induced the more elaborated morphologies of the pC1-d/e

cells, especially the extra vertical projection (EVP) near the midline of brains (**Figure 6

B-D**) (Deutsch et al., 2020). These changes exhibited from the second day of the pupal

stage (**Figure 6B1-B2 and 6E**) and maintained at the adult stage (**Figure 6 D1-D2 and

6F**). Meanwhile, the number of pC1 cell bodies in adult flies when EcR-A was reduced

were the same as that of wild type flies (**Figure 6G**). Previous studies suggested that

pC1d cells serve as a hub within the central brain for dsx$^+$ and fru$^+$ neurons (Deutsch et

al., 2020). Thus, the abnormal development of pC1d neurons may induce the changes

between pC1d neurons and other dsx$^+$ and fru$^+$ neurons to affect associated behaviors.

Furthermore, we asked whether reduced female copulation rate was due to that EcR-A

expression affected the activity of pC1 neurons. Because all pC1 cells characterized so

far project to the lateral junction of the lateral protocerebral complex (LPC) (Kimura et

al., 2015; Rezával et al., 2016; Scheffer et al., 2020; Wang et al., 2020b; Wu et al.,

2019; Zhou et al., 2014), we expressed GCamp6s in all pC1 neurons and tested the

calcium signals in the lateral junction of LPC when EcR-A was knocked down (**Figure 6**

D1-D2). Reduced EcR-A did not induce significantly different calcium responses in the

LPC (**Figure 6H**). Thus, our results suggested that the decreased female copulation

rate induced by reduced EcR-A in pC1 neurons was mainly due to the morphological

changes of pC1b neurons, which then modulate the connections of pC1b neurons with

other neurons.

The function of PTTH on EcR-A and pC1 neurons.

As newly formed prepupae, the ptth-Gal4 > UAS-Grim flies display similar changes in

gene expression to the genetic control flies to response to a high-titer ecdysone pulse.

These genes include the repression of EcR (McBrayer et al., 2007). According to the

contradictory functions of PTTH deletion and EcR-A reduction in pC1 neurons, we

wanted to know whether there is a similar feedforward relationship between PTTH and

EcR-A. We quantified the EcR-A expression in the whole pupa body during the start of

prepupa stage, when the pC1-ss2-Gal4 expressing neurons appear. Indeed, PTTH -/-

induced upregulated EcR-A compared with PTTH -/+ flies (**Figure 7A**). This suggested

the feedforward relationship between PTTH and EcR-A expression during the start of

prepupa stage. Consistent with this, when PTTH was deleted, pC1 neurons exhibited

the contradictory pattern to that when EcR-A was decreased in pC1 neurons (**Figure**

7B-C). These suggested the feedforward relationship between PTTH and EcR-A, and

may explain the contradictory functions of PTTH deletion and EcR-A reduction in pC1

neurons for female receptivity.

Furthermore, PTTH neurons are *dsx*-positive, almost all neurons regulating female

receptivity express Dsx^F. We wanted to know whether PTTH neurons affect other dsx$^+$

neurons, including pC1 neurons. Indeed, we detected the slightly overlap of dsx-

LexA>LexAop-RFP and torso-Gal4>UAS-GFP during larval stage (**Figure 7D**).

Furthermore, decreasing torso expression in pC1 neurons significantly inhibit female

receptivity (**Figure 7E**). The inhibited virgin female receptivity had no relationship either

with the locomotion activity of virgin females (**Figure 7F**). These results suggest that,

PTTH regulates female receptivity not only through ecdysone, but also may through

regulating other neurons especially Dsx^F-positive neurons associated with female

receptivity directly.

Discussion

In this study, we found that peptide hormone PTTH negatively modulates virgin female

receptivity through ecdysone. PTTH neurons are doublesex-positive and regulate virgin

female receptivity during neural development. PTTH deletion resulted in the increased

ecdysone receptor EcR-A expression in newly formed prepupae. Furthermore, EcR-A

functions in pC1 neurons to positively regulate virgin female receptivity during

metamorphosis mainly through modulating the anatomical morphology of pC1b

neurons. Additionally, decreasing the expression of PTTH receptor torso in pC1

neurons inhibited female receptivity. Taken together, our results revealed the contrary

functions of PTTH deletion and reduction of EcR-A in pC1 neurons during

neurodevelopment. In addition, EcR-A in pC1 neurons regulates virgin female

copulation rate during metamorphosis mainly through modulating the morphology of

pC1b neurons.

Most of neurons regulating sexual behaviors in female flies are dsx-positive. Our results

showed that PTTH neurons are also dsx^+ neurons. This suggested that PTTH neurons

have relationships with other dsx^+ neurons and the juvenile–adult transition is regulated

by *doublesex* gene. Indeed, we detected the overlap between torso-Gal4 signal and

dsx-LexA signal at the larval stage. Furthermore, when PTTH receptor torso was

decreased in dsx^+ pC1 neurons, female receptivity was inhibited. This suggested that

PTTH functions in pC1 neurons through neuronal projection or endocrine pathway

directly to regulate female receptivity. However, we did not detect the change of female

receptivity when Dsx^F was decreased in PTTH neurons. This suggested that Dsx^F

functions in PTTH neurons on other aspects, such as the development of PTTH

neurons or the synthesis and release of PTTH to regulate development.

PTTH regulates virgin female receptivity in an ecdysone-dependent manner before

metamorphosis. Ecdysone functions through its receptor EcR which is involved in all

phases of the nervous system development. In our study, reduced EcR-A expression in

all pC1 neurons, which encode the mating status of females, lead to the decreased

copulation rate. While, reduced EcR-A in pC1-a, c and e simultaneously did not reduce

the copulation rate significantly. This suggested that EcR-A plays the critical role in pC1-

b or/and -d neurons for regulating virgin female receptivity. Our results revealed that

reduced EcR-A induced the more elaborated morphologies of pC1d neurons. Previous

studies detected the synaptic connections between the axons of pC1d and the dendrites

of DNp13 neurons (Deutsch et al., 2020; Mezzera et al., 2020). DNp13 neurons are

command neurons for ovipositor extrusion. When females extruded, the ovipositor

physically impedes copulation (Mezzera et al., 2020; Wang et al., 2020a). However,

reduced EcR-A expression in pC1d neurons did not affect virgin female receptivity

(**Figure 5—figure supplement 4**). This might be due to three possibilities. First, the

more elaborated morphologies of pC1d neurons did not affect synaptic connections

between pC1d and DNp13 neurons. Second, the unchanged pC1 neural activity could

not affect the neural activity of DNp13 neurons. Third, the morphological change of

pC1d neurons is not sufficient for the decreased copulation rate. To sum, these suggest

that the morphological change of pC1b neurons is necessary for the decreased

copulation rate. However, due to the lack of pC1b drivers, we could not rule out

morphological changes in pC1b neurons when EcR-A was reduced.

In our study, PTTH negatively regulates female receptivity, while EcR-A positively

regulates female receptivity. Previous study revealed that, expression of EcR is

repressed in newly formed prepupae to response to high-titer ecdysone pulse

(Mcbrayer et al., 2007). We detected the similar feedforward relationship between PTTH

and EcR-A in newly formed prepupae. Consistent with this, we detected the contrary

pattern of pC1d neurons when PTTH was deleted, compare with that when EcR-A was

decreased in pC1 neurons. However, it is not sure that PTTH deletion could result in the

increased expression of EcR-A in pC1 neurons. In addition, PTTH deletion must affect

the development of almost other neurons, but not only pC1 neurons. This maybe the

reason for that reduction of Torso in pC1 neurons had contrary effect on female

receptivity to that when PTTH is deleted. So, the feedforward relationship between

PTTH and EcR-A in newly formed prepupae is one possible reason for the contrary

functions of PTTH deletion and reduction of EcR-A in pC1 neurons on female

receptivity.

Previous studies have demonstrated that the development of fru$^+$ neurons need EcR-A

in male *Drosophila melanogaster*. Furthermore, reduced EcR-A in fru$^+$ neurons induced

the male-male courtship (Dalton et al., 2009). We also detected the male-male courtship

behavior when PTTH was deleted (data not shown) as previous study (Mcbrayer et al.,

2007). This remits to the impact of ecdysone on dsx$^+$ or fru$^+$ neurons. Similarly, PTTH

deletion may also affect the development of other neurons and further other behaviors

such as the female fecundity and the body size (Mcbrayer et al., 2007; Rewitz K F 2009;

Shimell et al., 2018), although there is no sufficient evidence for the effects of fecundity

and the body size on female receptivity. So, we could not exclude all effects of other

aspects on female receptivity.

Our results suggested a regulatory role of PTTH in virgin female receptivity. Even

though insects and mammals represent highly diverged classes, insects have evolved a

similar strategy for triggering the juvenile–adult transition (Herbison, 2016; Pan et al.,

2019). The juvenile–adult transition involves the hypothalamic–pituitary–gonadal (HPG)

axis in mammals and the PG axis in insects. Among the neurons belonging to the axis,

PTTH neurons and GnRH neurons have the similar function to stimulate the PG gland

and pituitary gland to release hormones which trigger maturation, respectively. It will be

interesting to study the function of GnRH neurons on the mammal sexual behaviors.

This work extends the understanding of how neurodevelopmental processes regulate

adult sexual behavior.

 Materials and methods

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
antibody	Mouse anti-Bruchpilot antibody (nc82)	Developmental Studies Hybridoma Bank	Cat# nc82, RRID: AB_2314866	IHC (1:40)
antibody	Mouse Anti-Drosophila ecdysone receptor (EcR-A) Monoclonal Antibody	Developmental Studies Hybridoma Bank	Cat# 15G1a (EcR-A), RRID: AB_528214	IHC (1:10)
antibody	mouse Anti-Drosophila ecdysone receptor (EcR-B1) Antibody	Developmental Studies Hybridoma Bank	Cat# AD4.4(EcR-B1), RRID: AB_2154902	IHC (1:10)
antibody	Rabbit polyclonal anti-GFP	Thermo Fisher Scientific	Cat# A-11122, RRID: AB_221569	IHC (1:1000)
antibody	chicken polyclonal anti-GFP	Thermo Fisher Scientific	Cat# A10262, RRID: AB_2534023	IHC (1:1000)
antibody	Goat anti-rabbit, Alexa Fluor 488	Thermo Fisher Scientific	Cat# A-11034, RRID: AB_2576217	IHC (1:500)
antibody	Goat anti-chickent, Alexa Fluor 488	Thermo Fisher Scientific	Cat#A-11039; RRID: AB_2534096	IHC (1:500)
antibody	Goat anti-mouse, Alexa Fluor 488	Thermo Fisher Scientific	Cat# A-11029, RRID: AB_2534088	IHC (1:500)
antibody	Goat anti-rabbit, Alexa Fluor 546	Thermo Fisher Scientific	Cat# A-11010, RRID: AB_2534077	IHC (1:500)
antibody	Rabbit polyclonal anti-RFP	Thermo Fisher Scientific	Cat# R10367, RRID: AB_10563941	IHC (1:500)
antibody	Goat anti-mouse, Alexa Fluor 647	Thermo Fisher Scientific	Cat# A-21235, RRID: AB_2535804	IHC (1:500)
antibody	Rabbit polyclonal anti-PTTH	Zhou Lab, Chinese Academy of Sciences, this paper	N/A	IHC (1:1300)
chemical compound, drug	Paraformaldehyde (PFA)	Electron Microscopy Sciences	Cat#15713	8% PFA diluted in 1XPBS at 1:4 or 1:2
chemical compound, drug	DPX Mountant	Sigma-Aldrich	Cat#44581	
chemical compound, drug	Normal goat serum	Sigma-Aldrich	Cat#G9023	
chemical compound, drug	20-hydroxyecdysone	Cayman	Cat#16145	dissolved in 95% ethanol, 0.2 mg/ml
chemical compound, drug	TRIzol	Ambion	Cat#15596018	
Genetic reagent (*D.melanogaster*)	*LexAop2-mCD8::GFP*	Bloomington Stock Center	BL#32203	
Genetic reagent (*D.melanogaster*)	*;;UAS-mCD8::GFP*	Bloomington Stock Center	BL#32194	
Genetic reagent (*D.melanogaster*)	*;UAS-mCD8::GFP;*	Bloomington Stock Center	BL#5137	
Genetic reagent (*D.melanogaster*)	*UAS-dTrpA1/cyo*	Garrity Lab, Brandeis University	N/A	

Genetic reagent (D.melanogaster)	UAS-Kir2.1	Bloomington Stock Center	BL#6595	
Genetic reagent (D.melanogaster)	Ptth-Gal4	Rao Lab, Peking University	N/A	
Genetic reagent (D.melanogaster)	PtthLexA	Rao Lab, Peking University	N/A	
Genetic reagent (D.melanogaster)	⊿PTTH	Rao Lab, Peking University	N/A	
Genetic reagent (D.melanogaster)	UAS-PTTH	Zhou Lab, Chinese Academy of Sciences, this paper	N/A	
Genetic reagent (D.melanogaster)	isoCS	Rao Lab, Peking University	N/A	
Genetic reagent (D.melanogaster)	elav-Gal4	Rao Lab, Peking University	N/A	
Genetic reagent (D.melanogaster)	UAS-GFPStinger	Janelia Research Campus	N/A	
Genetic reagent (D.melanogaster)	LexAop-tomato	Janelia Research Campus	N/A	
Genetic reagent (D.melanogaster)	LexAop2-FlpL	Janelia Research Campus	N/A	
Genetic reagent (D.melanogaster)	UAS > stop > mCD8-GFP	Janelia Research Campus	N/A	
Genetic reagent (D.melanogaster)	Dsx-Gal4	Janelia Research Campus	N/A	
Genetic reagent (D.melanogaster)	Dsx-LexA	Janelia Research Campus	N/A	
Genetic reagent (D.melanogaster)	tub-Gal80^{ts}	Pan Lab, Southeast University	BL#7018	
Genetic reagent (D.melanogaster)	pC1-ss1-Gal4	Wang Lab, Lingang Laboratory	N/A	
Genetic reagent (D.melanogaster)	pC1-ss2- Gal4	Wang Lab, Lingang Laboratory	N/A	
Genetic reagent (D.melanogaster)	vpoDN-ss1-Gal4	Wang Lab, Lingang Laboratory	N/A	
Genetic reagent (D.melanogaster)	vpoDN-ss2-Gal4	Wang Lab, Lingang Laboratory	N/A	
Genetic reagent (D.melanogaster)	vpoDN-ss3-Gal4	Wang Lab, Lingang Laboratory	N/A	
Genetic reagent (D.melanogaster)	UAS-EcR-RNAi	Bloomington Stock Center	BL#9327	
Genetic reagent (D.melanogaster)	UAS-EcR-A-RNAi	Bloomington Stock Center	BL#9328	
Genetic reagent (D.melanogaster)	UAS-EcR-B1-RNAi	Bloomington Stock Center	BL#9329	
Genetic reagent (D.melanogaster)	pC1d-Gal4	Bloomington Stock Center	BL#86847	

Genetic reagent (*D.melanogaster*)	*UAS-PTTH-RNAi*	VDRC	V102043	
Genetic reagent (*D.melanogaster*)	*UAS-Dsx^F-RNAi*	Pan Lab	N/A	
Genetic reagent (*D.melanogaster*)	*UAS-Torso-RNAi*	Liu Lab	BL#33627	
recombinant DNA reagent	pBSK-attP-3P3-RFP-loxP	Bowen Deng et al., 2019	N/A	
recombinant DNA reagent	pBSK-attB-loxP-myc-T2A-Gal4Gal4-GMR-miniwhite	Bowen Deng et al., 2019	N/A	
recombinant DNA reagent	pBSK-attB-loxP-V5-T2A-LexA::p65-GMR-miniwhite	Bowen Deng et al., 2019	N/A	
software, algorithm	MATLAB	MathWorks, Natick, MA	https://www.mathworks.com/products/matlab.html	
software, algorithm	ImageJ	National Institutes of Health	https://imagej.nih.gov/ij/	
software, algorithm	Prism 7	GraphPad	https://www.graphpad.com/	
software, algorithm	R 4.1.3	RStudio	https://www.r-project.org	

Fly stocks

Flies were reared on standard cornmeal-yeast medium under a 12 hr:12 hr dark:light

cycle at 25°C and 60% humidity. All the knock-out lines in this study for screening have

been published [49]. The following strains were obtained from Dr. Yi Rao: *isoCS* (wild-

type), *ΔPtth*, *Ptth-Gal4, Ptth-LexA, elav-Gal4 and UAS-Kir2.1* (BL#6595). UAS-dTrpA1

was a gift from Dr. Paul Garrity. *UAS-GFPStinger*, *LexAop-tomato, LexAop2-FlpL,*

UAS > stop > mCD8-GFP, *dsx-Gal4* and *dsx-LexA* (Mellert et al., 2010) have been

described previously (Pfeiffer et al., 2008; Pfeiffer et al., 2010) and are obtained from

Janelia Research Campus. *tub-Gal80^{ts}* (BL#7018) was provided by Dr. Yufeng Pan.

pC1-ss1-Gal4, pC1-ss2-Gal4, vpoDN-ss1-Gal4, vpoDN-ss2-Gal4 and *vpoDN-ss3-Gal4*

were provided by Dr. Kaiyu Wang. *Torso-RNAi* (BL#33627) was a gift from Suning Liu.

The following lines were obtained from the Bloomington Drosophila Stock Center: *UAS-*

EcR-RNAi (BL#9327), *UAS-EcR-A-RNAi* (BL#9328), *UAS-EcR-B1-RNAi* (BL#9329),

UAS-mCD8::GFP (BL#32194), *LexAop2-mCD8::GFP* (BL#32203), *UAS-mCD8::GFP*

(BL#5137) and *pC1d-Gal4* (BL#86847). *UAS-PTTH-RNAi* (v102043) was from Vienna

Drosophila Resource Center (VDRC).

Behavioral assays

Flies were reared at 25°C and 60% humidity under a 12 hr light:12 hr dark cycle. Virgin

females and wild-type males were collected upon eclosion, placed in groups of 12 flies

each and aged 4–6 days (except for the assays for PTTH mutant on different days after

eclosion, and the molecular rescue assay for the 24h-old females) before carrying out

behavioral assay except for the transient thermogenetic experiments. Female receptivity

assays were conducted as previously described (Wang et al., 2022; Zhou et al., 2014).

A virgin female of defined genotype and a wild type male were gently cold anesthetized

and respectively introduced into two layers of the courtship chambers separated by a

removable transparent film. The flies were allowed to recover for at least 45 min before

the film was removed to allow the pair of flies to contact. The mating behavior was

recorded using a camera (Canon VIXIA HF R500) for 30 min at 17 fps for further

analysis.

For transient activation experiment by dTrpA1 in adult stage, flies were reared at 23°C.

Flies were loaded into courtship chamber and recovered for at least 30 min at 23°C,

then were placed at 23°C (control group) or 29°C (experimental group) for 30 min prior

to removing the film and videotaping. For activation experiment by dTrpA1 during

development, flies were reared at 29°C during the specific stages compared with the

controls who were reared at 23°C all the time. Flies were loaded into courtship chamber

and recovered for at least 45 min at 23°C prior to removing the film and videotaping.

Quantification and statistical analysis of female receptivity behavior

Two parameters including copulation rate and latency to copulation were used to

characterize receptivity and we got the data sets of two parameters from the same flies.

The time from removing the film to successful copulation was measured for each

female. The number of females that had engaged in copulation by the end of each 1

min interval within 30 min were summed and plotted as a percentage of successful

copulation. The latency to copulation was the time from removing the film to successful

copulation. All the time points that female successfully copulated were manually

analyzed and the data of unhealthy flies were discarded.

Temporally restricted RNAi

tub-Gal80^{ts} crosses were reared at either 18°C for control groups or 30°C for

experimental groups. Virgin females were collected at eclosion and were placed in

groups of 12 flies each and aged 4–6 days before carrying out behavior assay. Assays

were tested at 23°C.

Male courtship index

Courtship index was defined as the proportion of time the male followed, oriented

toward and attempted to copulate the female within 5 min of courtship initiation, marked

by the initial orientation toward and following the female.

Vaginal plate opening and ovipositor extrusion

A virgin female of defined genotype and a wild type male were aspirated into the

courtship chambers and respectively introduced into two layers of the courtship

chambers separated by a removable transparent film. The flies were allowed to recover

for 30 min before the film was removed. To allow visualization of vaginal plate opening,

we recorded uncompressed image sequences at 896 x 896 pixels and 50 frames per

second using a Photron Mini AX camera (Photron) with an AF-S VR Micro-Nikkor

105mm lens (Nikon). Instances of vaginal plate opening and ovipositor extrusion were

scored blind to genotype from frame by-frame playback during the first 5 min of

courtship or until copulation if it occurred within 5 min. Courtship initiation was defined

as the male orienting toward and beginning to follow the female. Rare trials with fewer

than 30 s of total courtship were discarded.

Locomotion assays

The rearing and experimental conditions in locomotion assays were the same as that in

the corresponding female receptivity assays, excepting that individual females were

loaded in the chambers without males. Spontaneous movements of the flies were

recorded with a camera (Canon VIXIA HF R500) for 30 min at 30 fps for further

analysis. The activity of flies during the middle 10 min was analyzed to calculate the

average walking speed using Ctrax software.

Egg Laying

Virgin females were collected upon eclosion and one fly was housed on standard

medium at 25°C, 60% relative humidity, 12 hr light:12 hr dark and allowed to lay eggs in

single vials. Each fly was transferred into new food tube every 24 hr. The number of

eggs was counted at the end of each 24 hr period. The numbers during the 3rd and 4th

day were summed for statistics and plot.

Immunohistochemistry

Whole brains of flies were dissected in 1x PBS and fixed in 2% paraformaldehyde

diluted in 1x PBS for 55 min at room temperature. The samples were washed with PBT

(1x PBS containing 0.3% Triton X-100) for 1 hour (3 x 20 min), followed by blocking in

5% normal goat serum (Blocking solution, diluted in 0.3% PBT) for 1 hour at room

temperature. Then, the samples were incubated in primary antibodies (diluted in

blocking solution) for 18–24 hours at 4°C. Samples were washed with 0.3% PBT for 1

hour (3 x 20 min), then were incubated in secondary antibodies (diluted in blocking

solution) for 18–24 hours at 4°C. Samples were washed with 0.3% PBT for 1 hour (3 x

20 min), then were fixed in 4% paraformaldehyde for 4 hours at room temperature. After

washed with 0.3% PBT for 1 hour (3 x 20 min), brains were mounted on poly-L-lysine-

coated coverslip in 1x PBS. The coverslip was dipped for 5 min with ethanol of

30%→50%→70%→95%→100% sequentially at room temperature, and then dipped for

5 min three times with xylene. Finally, brains were mounted with DPX and allowed DPX

to dry for 2 days before imaging. Primary antibodies used were: chicken anti-GFP

(1:1000; Life Technologies #A10262), rabbit anti-GFP (1:1000; Life Technologies

#A11122), rabbit anti-RFP (1:1000; Life Technologies #R10367), rabbit anti-PTTH

antibody (1:1300), mouse anti-nc82 (1:40; DSHB), mouse anti-EcR-A (1:10;

AB_528214) and mouse anti-EcR-B1 (1:10; AB_2154902). Secondary antibodies used

were: Alexa Fluor goat anti-chicken 488 (1:500; Life Technologies #A11039), Alexa

Fluor goat anti-rabbit 488 (1:500; Life Technologies #A11034), Alexa Fluor goat anti-

rabbit 546 (1:500; Life Technologies #A11010), Alexa Fluor goat anti-mouse 647 (1:500;

Life Technologies #A21235), and Alexa Fluor goat anti-mouse 488 (1:500; Life

Technologies #A11029).

Confocal microscopy and image analysis

Confocal imaging was performed under an LSM 710 inverted confocal microscope

(ZEISS, Germany), with a Plan-Apochromat 20×/0.8 M27 objective or an EC Plan-

Neofluar 40x/1.30 oil DIC M27 objective, and later analyzed using Fiji software.

Generation of anti-PTTH antibody

The antisera used to recognize PTTH peptide were raised in New Zealand white rabbits

using the synthetic peptide N'- TSQSDHPYSWMNKDQPWQFKC -C'. The synthesis of

antigen peptide, the production and purification of antiserum were performed by Beijing

Genomics Institute (BGI).

Generation of UAS-PTTH

pJFRC28-5XUAS-IVS-GFP-p10 (#12073; Fungene Biotechnology, Shanghai, China)

was used for the generation of the pJFRC28-UAS-PTTH construct. The pJFRC28-

10XUAS-IVS-GFP-p10 plasmid was digested with NotI and XbaI to remove the GFP

coding sequence, and then the cDNA of PTTH was cloned into this plasmid by Gibson

Assembly. The Kozak sequence was added right upstream of the ATG. UAS-PTTH

constructs were injected and integrated into the attP2 site on the third chromosome

through phiC31 integrase mediated transgenesis. The construct was confirmed using

DNA sequencing through PCR. The primers used for cloning PTTH cDNA were as

follows:

UAS-PTTH-F

ATTCTTATCCTTTACTTCAGGCGGCCGCAAAATGGATATAAAAGTATGGCGACTCC

UAS-PTTH-R

GTTATTTTAAAAACGATTCATTCTAGATCACTTTGTGCAGAAGCAGCCG

Genomic DNA extraction and RT-PCR

Genomic DNA was extracted from 10 whole bodies of wandering flies using MightyPrep

reagent for DNA (Takara #9182). Whole body RNA was extracted from 10 whole bodies

of wandering flies using TRIzol (Ambion #15596018). cDNA was generated from total

RNA using the Prime Script reagent kit (Takara #RR047A). Candidates of *ΔPtth* were

characterized by the loss of DNA band in the deleted areas through PCR on the

genomic DNA and cDNA. Primer sequences used in Figure 1 are listed in **Table S1**.

Measurements of pupariation timing and adult mass

The flies were reared at 25°C and 60% humidity under a 12 hr light:12 hr dark cycle.

Two-hour time collections of embryos laid on standard food vials. Each vial contained

20-30 eggs. The range of time for pupariation were recorded for each vial. Sexed adults

of 24h-old were weighted in groups of ten flies using a NENVER-TP-214 microbalance

at the same time.

Identification of sex in *Drosophila* larvae

Third instar larvae can be sexed (True et al., 2014). Gonads are translucent and visible

in side view in the posterior third of the larva. The male gonads are about five times

bigger than the female gonads. The identified wandering female and male larvae were

reared in different vials for the subsequent experiments.

Rescue by 20-hydroxyecdysone feeding

Thirty freshly ecdysed *ΔPtth* L3 larvae, grown at 25°C and 60% humidity under a 12 hr

light:12 hr dark cycle, were washed with water and transferred to normal food for

additional aging. After 20 h, larvae were washed and transferred to a vial supplemented

with either 20-hydroxyecdysone (20E, Cayman #16145, dissolved in 95% ethanol, final

concentration 0.2 mg/ml) or 95% ethanol (same volume as 20E). The wild type larvae

were directly transferred to vials supplemented with 20E or 95% ethanol upon L3

ecdysis. Once seeded with L3 larvae, the vials were returned to 25°C and 60% humidity

under a 12 hr light:12 hr dark cycle.

Quantification of fluorescence intensity

The fluorescence intensity was quantified using Fiji software. The areas of interest

(ROI) were marked in the slices including the interested regions and quantified using

the "plot z-axis profile" function. The fluorescence intensity in each slice was summed

for statistics and plot. The parameters used for confocal imaging of each brain were the

same.

Calcium imaging

Flies aged 4-6-days were immobilized on ice for ~30 s. The brain was then dissected

out in extra-cellular solution (ECS) that contains (in millimoles): 108 NaCl, 5 KC1, 5

trehalose, 5 sucrose, 5 HEPES, 26 $NaHCO_3$, 1 NaH_2PO_4, 2 $CaCl_2$, and 1.5 $MgCl_2$ [pH

7.1 – 7.3 when bubbled with 95% (vol/vol) O_2/5% (vol/vol) CO_2, ~290 mOsm] and

mounted on a poly-D-lysine coated coverslip. The samples were continuously perfused

with ECS.

Calcium imaging was performed at 21°C on a customized two-photon microscope

equipped with a resonant scanner (Nikon), a piezo objective scanner (Nikon) and a 40x

water-immersion objective (Nikon). GCaMP6s was excited at 920 nm.

Analysis of calcium imaging data was done offline with NIS-Elements AR 5.30.01.

Briefly, the square region of interest (ROIs), 25 pixels on the pC1 neurons in the center

of lateral junction, was chosen for measurements. For each frame, the average

fluorescence intensity of pixels within ROIs was calculated blind to genotype. The

average fluorescence intensity of ROIs in each frame covering pC1 neurons were

summed for statistics and plot.

qRT-PCR

Total RNA was extracted from about 10 flies using TRIzol (Ambion #15596018). The

cDNA was synthesized using Prime Script reagent kit (Takara #RR047A). Quantitative

PCR was performed on Thermo Piko Real 96 (Thermo) using SYBR Green PCR Master

Mix (Takara #RR820A). The mRNA expression level was calculated by the $2^{-\Delta\Delta Ct}$

method and the results were plotted by using tubulin as the reference gene. Primers are

listed in **Table S1**. All reactions were performed in triplicate. The average of four

biological replicates ± SEM was plotted.

Statistical analysis

Statistical analyses were carried out using R software version 3.4.3 or Prism7

(GraphPad software). For the copulation rate, chi-square test is applied. The Mann-

Whitney U test was applied for analyzing the significance of two columns. Kruskal-

Wallis ANOVA test followed by post-hoc Mann-Whitney U test, was used to identify

significant differences between multiple groups.

Data, Materials, and Software Availability

All study data are included in the main text and supporting information, except for the

sequence data of RNAseq (**Supplementary File 3**) has been deposited in the Genome

Sequence Archive under the accession number CRA012130 in PRJCA018750. This

study does not involve new code. Fly stocks and reagents used in this study are

available from the corresponding author upon reasonable request.

Acknowledgments

We thank Yi Rao (Peking University), Yufeng Pan (Southeast University), Kaiyu Wang

(Lingang Laboratory), Yan Zhu (Chinese Academy of Sciences), Paul Garrity (Brandeis

References

Avila F W, Sirot L K, LaFlamme B A, Rubinstein C D, & Wolfner M F. (2011). Insect seminal fluid
proteins: identification and function. *Annu Rev Entomol, 56*, 21-40.
doi:10.1146/annurev-ento-120709-144823
Billeter J C, Villella A, Allendorfer J B, Dornan A J, Richardson M, Gailey D A, & Goodwin S F.
(2006). Isoform-specific control of male neuronal differentiation and behavior in
Drosophila by the fruitless gene. *Curr Biol, 16*(11), 1063-1076.
doi:10.1016/j.cub.2006.04.039
Clowney E J, Iguchi S, Bussell J J, Scheer E, & Ruta V. (2015). Multimodal chemosensory circuits
controlling male courtship in Drosophila. *Neuron, 87*(5), 1036-1049.
doi:10.1016/j.neuron.2015.07.025
Connolly K, & Cook R. (1973). Rejection responses by female drosophila-melanogaster - their
ontogeny, causality and effects upon behavior of courting male. *Behaviour, 44*(1-2), 142-
166. doi:10.1163/156853973x00364
Dai J D, & Gilbert L I. (1991). Metamorphosis of the corpus allatum and degeneration of the
prothoracic glands during the larval-pupal-adult transformation of Drosophila
melanogaster: a cytophysiological analysis of the ring gland. *Dev Biol, 144*(2), 309-326.
doi:10.1016/0012-1606(91)90424-2
Dalton J E, Lebo M S, Sanders L E, Sun F, & Arbeitman M N. (2009). Ecdysone receptor acts in
fruitless- expressing neurons to mediate drosophila courtship behaviors. *Curr Biol,
19*(17), 1447-1452. doi:10.1016/j.cub.2009.06.063

Demir E, & Dickson B J. (2005). fruitless splicing specifies male courtship behavior in Drosophila.
Cell, 121(5), 785-794. doi:10.1016/j.cell.2005.04.027

Deng B, Li Q, Liu X, Cao Y, Li B, Qian Y, Xu R, Mao R, Zhou E, Zhang W, Huang J, & Rao Y. (2019).
Chemoconnectomics: mapping chemical transmission in Drosophila. *Neuron, 101*(5),
876-893.e874. doi:10.1016/j.neuron.2019.01.045

Deutsch D, Pacheco D, Encarnacion-Rivera L, Pereira T, Fathy R, Clemens J, Girardin C, Calhoun
A, Ireland E, Burke A, Dorkenwald S, McKellar C, Macrina T, Lu R, Lee K, Kemnitz N, Ih D,
Castro M, Halageri A, Jordan C, Silversmith W, Wu J, Seung H S, & Murthy M. (2020). The
neural basis for a persistent internal state in Drosophila females. *Elife, 9*.
doi:10.7554/eLife.59502

Dickson B J. (2008). Wired for sex: the neurobiology of Drosophila mating decisions. *Science,*
322(5903), 904-909. doi:10.1126/science.1159276

Feng K, Palfreyman M T, Hasemeyer M, Talsma A, & Dickson B J. (2014). Ascending SAG neurons
control sexual receptivity of Drosophila females. *Neuron, 83*(1), 135-148.
doi:10.1016/j.neuron.2014.05.017

Ferveur J F. (2010). Drosophila female courtship and mating behaviors: sensory signals, genes,
neural structures and evolution. *Curr Opin Neurobiol, 20*(6), 764-769.
doi:10.1016/j.conb.2010.09.007

Fuyama Y, & Ueyama M. (1997). Ovulation and the suppression of mating in Drosophila
melanogaster females: behavioral basis. *Behav Genet, 27*(5), 483-488.
doi:10.1023/a:1025630602057

Ganter G K, Walton K L, Merriman J O, Salmon M V, Brooks K M, Maddula S, & Kravitz E A.
(2007). Increased male-male courtship in ecdysone receptor deficient adult flies. *Behav*
Genet, 37(3), 507-512. doi:10.1007/s10519-006-9140-1

Greenspan R J, & Ferveur J F. (2000). Courtship in Drosophila. *Annu Rev Genet, 34*, 205-232.
doi:10.1146/annurev.genet.34.1.205

Hall J. (1978). Courtship among males due to a male-sterile mutation inDrosophila
melanogaster. *Behavior Genetics*(8-2).

Hall J C. (1994). The mating of a fly. *Science, 264*(5166), 1702-1714.
doi:10.1126/science.8209251

Hamada F N, Rosenzweig M, Kang K, Pulver S R, Ghezzi A, Jegla T J, & Garrity P A. (2008). An
internal thermal sensor controlling temperature preference in Drosophila. *Nature,*
454(7201), 217-220. doi:10.1038/nature07001

Häsemeyer M, Yapici N, Heberlein U, & Dickson B J. (2009). Sensory neurons in the Drosophila
genital tract regulate female reproductive behavior. *Neuron, 61*(4), 511-518.
doi:10.1016/j.neuron.2009.01.009

Herbison A E. (2016). Control of puberty onset and fertility by gonadotropin-releasing hormone
neurons. *Nat Rev Endocrinol, 12*(8), 452-466. doi:10.1038/nrendo.2016.70

Hoopfer E D, Jung Y, Inagaki H K, Rubin G M, & Anderson D J. (2015). P1 interneurons promote a
persistent internal state that enhances inter-male aggression in Drosophila. *Elife, 4*.
doi:10.7554/eLife.11346

Imura E, Shimada-Niwa Y, Nishimura T, Huckesfeld S, Schlegel P, Ohhara Y, Kondo S, Tanimoto
H, Cardona A, Pankratz M J, & Niwa R. (2020). The Corazonin-PTTH Neuronal Axis
Controls Systemic Body Growth by Regulating Basal Ecdysteroid Biosynthesis in

Drosophila melanogaster. *Curr Biol, 30*(11), 2156-2165 e2155.
doi:10.1016/j.cub.2020.03.050
Ishimoto H, & Kamikouchi A. (2020). A feedforward circuit regulates action selection of pre-
mating courtship behavior in female Drosophila. *Curr Biol, 30*(3), 396-407.e394.
doi:10.1016/j.cub.2019.11.065
Ito H, Sato K, Koganezawa M, Ote M, Matsumoto K, Hama C, & Yamamoto D. (2012). Fruitless
recruits two antagonistic chromatin factors to establish single-neuron sexual
dimorphism. *Cell, 149*(6), 1327-1338. doi:10.1016/j.cell.2012.04.025
Jang Y H, Chae H S, & Kim Y J. (2017). Female-specific myoinhibitory peptide neurons regulate
mating receptivity in Drosophila melanogaster. *Nat Commun, 8*(1), 1630.
doi:10.1038/s41467-017-01794-9
Kimura K, Hachiya T, Koganezawa M, Tazawa T, & Yamamoto D. (2008). Fruitless and doublesex
coordinate to generate male-specific neurons that can initiate courtship. *Neuron, 59*(5),
759-769. doi:10.1016/j.neuron.2008.06.007
Kimura K, Sato C, Koganezawa M, & Yamamoto D. (2015). Drosophila ovipositor extension in
mating behavior and egg deposition involves distinct sets of brain interneurons. *PLoS*
One, 10(5), e0126445. doi:10.1371/journal.pone.0126445
Kohatsu S, Koganezawa M, & Yamamoto D. (2011). Female contact activates male-specific
interneurons that trigger stereotypic courtship behavior in Drosophila. *Neuron, 69*(3),
498-508. doi:10.1016/j.neuron.2010.12.017
Kubli E. (2003). Sex-peptides: seminal peptides of the Drosophila male. *Cell Mol Life Sci, 60*(8),
1689-1704. doi:10.1007/s00018-003-3052
Kurtovic A, Widmer A, & Dickson B J. (2007). A single class of olfactory neurons mediates
behavioural responses to a Drosophila sex pheromone. *Nature, 446*(7135), 542-546.
doi:10.1038/nature05672
Kvitsiani D, & Dickson B J. (2006). Shared neural circuitry for female and male sexual behaviours
in Drosophila. *Curr Biol, 16*(10), R355-356. doi:10.1016/j.cub.2006.04.025
Lavrynenko O, Rodenfels J, Carvalho M, Dye N A, Lafont R, Eaton S, & Shevchenko A. (2015).
The ecdysteroidome of Drosophila: influence of diet and development. *Development,*
142(21), 3758-3768. doi:10.1242/dev.124982
Ma B, Wang R, Liu Y, Deng B, Wang T, Wu F, & Zhou C. (2022). Serotonin signaling modulates
sexual receptivity of virgin female Drosophila. *Neurosci Bull, 38*(11), 1277-1291.
doi:10.1007/s12264-022-00908-8
Manoli D S, Fan P, Fraser E J, & Shah N M. (2013). Neural control of sexually dimorphic
behaviors. *Curr Opin Neurobiol, 23*(3), 330-338. doi:10.1016/j.conb.2013.04.005
Manoli D S, Foss M, Villella A, Taylor B J, Hall J C, & Baker B S. (2005). Male-specific fruitless
specifies the neural substrates of Drosophila courtship behaviour. *Nature, 436*(7049),
395-400. doi:10.1038/nature03859
Manoli D S, Meissner G W, & Baker B S. (2006). Blueprints for behavior: genetic specification of
neural circuitry for innate behaviors. *Trends Neurosci, 29*(8), 444-451.
doi:10.1016/j.tins.2006.06.006
McBrayer Z, Ono H, Shimell M, Parvy J P, Beckstead R B, Warren J T, Thummel C S, Dauphin-
Villemant C, Gilbert L I, & O'Connor M B. (2007). Prothoracicotropic hormone regulates

developmental timing and body size in Drosophila. *Dev Cell, 13*(6), 857-871.
doi:10.1016/j.devcel.2007.11.003
McGuire S E, Mao Z, & Davis R L. (2004). Spatiotemporal gene expression targeting with the
TARGET and gene-switch systems in Drosophila. *Sci STKE, 2004*(220), pl6.
doi:10.1126/stke.2202004pl6
Mellert D J, Knapp J M, Manoli D S, Meissner G W, & Baker B S. (2010). Midline crossing by
gustatory receptor neuron axons is regulated by fruitless, doublesex and the
Roundabout receptors. *Development, 137*(2), 323-332. doi:10.1242/dev.045047
Mellert D J, Robinett C C, & Baker B S. (2012). doublesex functions early and late in gustatory
sense organ development. *PLoS One, 7*(12), e51489. doi:10.1371/journal.pone.0051489
Mezzera C, Brotas M, Gaspar M, Pavlou H J, Goodwin S F, & Vasconcelos M L. (2020). Ovipositor
extrusion promotes the transition from courtship to copulation and signals female
acceptance in Drosophila melanogaster. *Curr Biol, 30*(19), 3736-3748.e3735.
doi:10.1016/j.cub.2020.06.071
Neckameyer W S. (1998). Dopamine modulates female sexual receptivity in Drosophila
melanogaster. *J Neurogenet, 12*(2), 101-114. doi:10.3109/01677069809167259
Naoki Yamanaka, Nuria M Romero, Francisco A Martin, Kim F Rewitz, Mu Sun, Michael B
O'Connor, Pierre Léopold. (2013). Neuroendocrine Control of Drosophila larval light
preference. *Science*, 41(6150), 1113-1116. doi: 10.1126/science.1241210
Pan X, & O'Connor M B. (2019). Developmental Maturation: Drosophila AstA signaling provides
a kiss to grow up. *Curr Biol, 29*(5), R161-R164. doi:10.1016/j.cub.2019.01.040
Pan Y, & Baker B S. (2014). Genetic identification and separation of innate and experience-
dependent courtship behaviors in Drosophila. *Cell, 156*(1-2), 236-248.
doi:10.1016/j.cell.2013.11.041
Pan Y, Meissner G W, & Baker B S. (2012). Joint control of Drosophila male courtship behavior
by motion cues and activation of male-specific P1 neurons. *PNAS, 109*(25), 10065-
10070. doi:10.1073/pnas.1207107109
Pan Y, Robinett C C, & Baker B S. (2011). Turning males on: activation of male courtship
behavior in Drosophila melanogaster. *PLoS One, 6*(6), e21144.
doi:10.1371/journal.pone.0021144
Pavlou H J, & Goodwin S F. (2013). Courtship behavior in Drosophila melanogaster: towards a
'courtship connectome'. *Curr Opin Neurobiol, 23*(1), 76-83.
doi:10.1016/j.conb.2012.09.002
Pfeiffer B D, Jenett A, Hammonds A S, Ngo T T, Misra S, Murphy C, Scully A, Carlson J W, Wan K
H, Laverty T R, Mungall C, Svirskas R, Kadonaga J T, Doe C Q, Eisen M B, Celniker S E, &
Rubin G M. (2008). Tools for neuroanatomy and neurogenetics in Drosophila. *PNAS,*
105(28), 9715-9720. doi:10.1073/pnas.0803697105
Pfeiffer B D, Ngo T T, Hibbard K L, Murphy C, Jenett A, Truman J W, & Rubin G M. (2010).
Refinement of tools for targeted gene expression in Drosophila. *Genetics, 186*(2), 735-
755. doi:10.1534/genetics.110.119917
Rewitz K F Y N, Gilbert L I , et al. (2009). The insect neuropeptide PTTH activates receptor
tyrosine kinase torso to initiate metamorphosis. *Science, 326*(5958), 1403-1405. doi:
10.1126/science.1176450

930 Rewitz K F, Yamanaka N, Gilbert L I, & O'Connor M B. (2009). The insect neuropeptide PTTH
931   activates receptor tyrosine kinase torso to initiate metamorphosis. *Science, 326*(5958),
932   1403-1405. doi:10.1126/science.1176450
933 Rezával C, Nojima T, Neville M C, Lin A C, & Goodwin S F. (2014). Sexually dimorphic
934   octopaminergic neurons modulate female postmating behaviors in Drosophila. *Curr Biol,*
935   *24*(7), 725-730. doi:10.1016/j.cub.2013.12.051
936 Rezával C, Pattnaik S, Pavlou H J, Nojima T, Brüggemeier B, D'Souza L A D, Dweck H K M, &
937   Goodwin S F. (2016). Activation of latent courtship circuitry in the brain of Drosophila
938   females induces male-like behaviors. *Curr Biol, 26*(18), 2508-2515.
939   doi:10.1016/j.cub.2016.07.021
940 Riddiford L M, Cherbas P, & Truman J W. (2000). Ecdysone receptors and their biological
941   actions. *Vitam Horm, 60*, 1-73. doi:10.1016/s0083-6729(00)60016-x
942 Rideout E J, Dornan A J, Neville M C, Eadie S, & Goodwin S F. (2010). Control of sexual
943   differentiation and behavior by the doublesex gene in Drosophila melanogaster. *Nat*
944   *Neurosci, 13*(4), 458-466. doi:10.1038/nn.2515
945 Roy S, Saha T T, Zou Z, & Raikhel A S. (2018). Regulatory pathways controlling female insect
946   reproduction. *Annu Rev Entomol, 63*, 489-511. doi:10.1146/annurev-ento-020117-
947   043258
948 Ruta V, Datta S R, Vasconcelos M L, Freeland J, Looger L L, & Axel R. (2010). A dimorphic
949   pheromone circuit in Drosophila from sensory input to descending output. *Nature,*
950   *468*(7324), 686-690. doi:10.1038/nature09554
951 Scheffer L K, Xu C S, Januszewski M et al.. (2020). A connectome and analysis of the adult
952   Drosophila central brain. *Elife, 9*. doi:10.7554/eLife.57443
953 Schretter C E, Aso Y, Robie A A, Dreher M, Dolan M J, Chen N, Ito M, Yang T, Parekh R, Branson
954   K M, & Rubin G M. (2020). Cell types and neuronal circuitry underlying female
955   aggression in Drosophila. *Elife, 9*. doi:10.7554/eLife.58942
956 Sengupta S, Chan Y B, Palavicino-Maggio C B, & Kravitz E A. (2022). GABA transmission from
957   mAL interneurons regulates aggression in Drosophila males. *PNAS, 119*(5).
958   doi:10.1073/pnas.2117101119
959 Shimell M, Pan X, Martin F A, Ghosh A C, Leopold P, O'Connor M B, & Romero N M. (2018).
960   Prothoracicotropic hormone modulates environmental adaptive plasticity through the
961   control of developmental timing. *Development, 145*(6). doi:10.1242/dev.159699
962 Siwicki K K, & Kravitz E A. (2009). Fruitless, doublesex and the genetics of social behavior in
963   Drosophila melanogaster. *Curr Opin Neurobiol, 19*(2), 200-206.
964   doi:10.1016/j.conb.2009.04.001
965 Stockinger P, Kvitsiani D, Rotkopf S, Tirián L, & Dickson B J. (2005). Neural circuitry that governs
966   Drosophila male courtship behavior. *Cell, 121*(5), 795-807.
967   doi:10.1016/j.cell.2005.04.026
968 Taisz I, Donà E, Münch D, Bailey S N, Morris B J, Meechan K I, Stevens K M, Varela-Martínez I,
969   Gkantia M, Schlegel P, Ribeiro C, Jefferis G, & Galili D S. (2023). Generating parallel
970   representations of position and identity in the olfactory system. *Cell, 186*(12), 2556-
971   2573.e2522. doi:10.1016/j.cell.2023.04.038

Terhzaz S, Rosay P, Goodwin S F, & Veenstra J A. (2007). The neuropeptide SIFamide modulates
sexual behavior in Drosophila. *Biochem Biophys Res Commun, 352*(2), 305-310.
doi:10.1016/j.bbrc.2006.11.030
True, & John. (2014). Atlas of Drosophila Morphology: Wild-type and Classical Mutants.
Quarterly Review of Biology, 89(2), 188-188.
Truman J W, & Riddiford L M. (2023). Drosophila postembryonic nervous system development:
a model for the endocrine control of development. *Genetics, 223*(3).
doi:10.1093/genetics/iyac184
Truman J W, Talbot W S, Fahrbach S E, & Hogness D S. (1994). Ecdysone receptor expression in
the CNS correlates with stage-specific responses to ecdysteroids during Drosophila and
Manduca development. *Development, 120*(1), 219-234. doi:10.1242/dev.120.1.219
von Philipsborn A C, Liu T, Yu J Y, Masser C, Bidaye S S, & Dickson B J. (2011). Neuronal control
of Drosophila courtship song. *Neuron, 69*(3), 509-522.
doi:10.1016/j.neuron.2011.01.011
Wang F, Wang K, Forknall N, Parekh R, & Dickson B J. (2020a). Circuit and Behavioral
Mechanisms of sexual rejection by Drosophila females. *Curr Biol, 30*(19), 3749-3760
e3743. doi:10.1016/j.cub.2020.07.083
Wang F, Wang K, Forknall N, Patrick C, Yang T, Parekh R, Bock D, & Dickson B J. (2020b). Neural
circuitry linking mating and egg laying in Drosophila females. *Nature, 579*(7797), 101-
105. doi:10.1038/s41586-020-2055-9
Wang K, Wang F, Forknall N, Yang T, Patrick C, Parekh R, & Dickson B J. (2021). Neural circuit
mechanisms of sexual receptivity in Drosophila females. *Nature, 589*(7843), 577-581.
doi:10.1038/s41586-020-2972-7
Wang L, & Anderson D J. (2010). Identification of an aggression-promoting pheromone and its
receptor neurons in Drosophila. *Nature, 463*(7278), 227-231. doi:10.1038/nature08678
Wang T, Jing B, Deng B, Shi K, Li J, Ma B, Wu F, & Zhou C. (2022). Drosulfakinin signaling
modulates female sexual receptivity in Drosophila. *Elife, 11*. doi:10.7554/eLife.76025
Wu Y, Bidaye S, & Mahringer D. (2019). Drosophila female-specific brain neuron elicits
persistent position- and direction-selective male-like social behaviors.
doi:10.1101/594960
Yamamoto D. (2007). The neural and genetic substrates of sexual behavior in Drosophila. *Adv
Genet, 59*, 39-66. doi:10.1016/S0065-2660(07)59002-4
Yamamoto D, & Koganezawa M. (2013). Genes and circuits of courtship behaviour in Drosophila
males. *Nat Rev Neurosci, 14*(10), 681-692. doi:10.1038/nrn3567
Yamanaka N, Romero N M, Martin F A, Rewitz K F, Sun M, O'Connor M B, & Léopold P. (2013).
Neuroendocrine control of Drosophila larval light preference. *Science, 341*(6150), 1113-
1116. doi:10.1126/science.1241210
Yang C H, Rumpf S, Xiang Y, Gordon M D, Song W, Jan L Y, & Jan Y N. (2009). Control of the
postmating behavioral switch in Drosophila females by internal sensory neurons.
Neuron, 61(4), 519-526. doi:10.1016/j.neuron.2008.12.021
Zhou C, Pan Y, Robinett C C, Meissner G W, & Baker B S. (2014). Central brain neurons
expressing doublesex regulate female receptivity in Drosophila. *Neuron, 83*(1), 149-163.
doi:10.1016/j.neuron.2014.05.038
1015

Figure Legends

 Figure 1 *Ptth* null mutants have increased virgin female receptivity.

(A-C) Generation and validation of a 974 bp deletion mutant of the *Ptth* gene. The 5'

UTR and almost all coding sequence were deleted. The deletion was confirmed through

PCR analysis at the PTTH locus in genomic DNA samples (B), and through RT-PCR to

identify the loss of PTTH transcripts in cDNA samples of wandering larvae (C). (D-G)

Virgin female receptivity of *Ptth* null mutants on the 1st (D), 2nd (E), 3rd (F) and 6th day

(G) respectively. The comparison referred to $Ptth^{Delete}/Ptth^{Delete}$. (H) Enhanced virgin

female receptivity of *ΔPtth* null mutants was rescued by elav-Gal4 driving UAS-PTTH.

The increased copulation rate and decreased latency to copulation on the 1st day after

eclosion were rescued to the comparable level of control. The comparison referred to

elav-Gal4/+; $Ptth^{Delete}/Ptth^{Delete}$;UAS-PTTH/+. The copulation latency *and copulation rate*

of *elav-Gal4/+; $Ptth^{Delete}/Ptth^{Delete}$* is higher and lower than $Ptth^{Delete}/Ptth^{Delete}$;UAS-

PTTH/+ *respectively*. The number of female flies paired with wild type males is

displayed in parentheses. For the copulation rate, chi-square test is applied. For the

latency to copulation, Kruskal-Wallis ANOVA and post hoc Mann-Whitney U tests are

applied. Error bars indicate SEM. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$, ns

indicates no significant difference.

1035

1036

1037

1038

1039

1040

1041

1042

1043

1044

1045

1046

1047

Figure 2 PTTH neurons are doublesex-positive neurons.

(A-B) Expression pattern of Ptth-Gal4 revealed by anti-GFP in larvae central nervous

system (CNS) (A) and adult brain (B). Representative of five female flies. Scale bars, 50

µm. (C) All PTTH neurons were colabeled by dsx-Gal4 driving UAS-GFP-Stinger (red)

and Ptth-LexA driving LexAop-tomato (green). Representative of five female brains.

Scale bars, 50 µm and 5 µm (zoom-in). (D) All PTTH neurons were *Ptth* and *Dsx* co-

expressing, labeled by intersectional strategy. The larvae flies were the wandering

ones. The adult flies were within-10 hours-old adults. Representative of 5 female brains.

Scale bars, 50 µm.

1057

1058

1059

Figure 3 Activation of PTTH neurons expressing PTTH during the 3rd-instar larvae

inhibits virgin female receptivity.

(A) Four developmental stages of *Drosophila* before eclosion when PTTH neurons were

thermogenetic activated by dTrpA1. L1, L2, and L3: start of three larval stages, W: start

of wandering stage, Pp: puparium formation, P1 and P2: start of the 1st and 2nd day of

pupal stage. (B-E) Ptth-Gal4 driving UAS-dTrpA1 activated PTTH neurons at 29°C.

Activation of PTTH neurons at the stage 2 significantly decreased copulation rate (C),

but not at the stage 1 (B), stage 3 (D) and stage 4 (E). (F) Activation of PTTH neurons

at the stage 2 significantly increased the latency to copulation. (G) Mean velocity had no

significant change when PTTH neurons were activated during the stage 2 compared

with control females (ns = not significant, Kruskal-Wallis ANOVA and post hoc Mann-

Whitney U tests, mean ± SEM, n = 8-12)*. The comparison referred to *Ptth-Gal4/UAS-*

dTrpA1. Female flies were 4-days-old adults. The number of female flies paired with

wild-type males is displayed in parentheses. For the copulation rate, chi-square test is

applied. For the latency to copulation, Mann-Whitney U test is applied. Error bars

indicate SEM. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, ns indicates no

significant difference.

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

1089

1090

1091

Figure 4 Feeding 20E restores virgin female receptivity of *Ptth* null mutant flies.

(A-B) The increased copulation rate and decreased latency to copulation of the 24h-old

ΔPtth flies were rescued to the comparable level of wild type females by feeding 20E to

the 3rd-instar larval *ΔPtth* flies. The wild type larval females fed by 20E had no

significantly different copulation rate and latency to copulation compared with the wild

type females fed by the same volume of 95% ethanol which is the solvent of 20E. The

comparison referred to *PtthDelete/PtthDelete* + 20E. Female flies were 1-day-old. The

number of female flies paired with wild-type males is displayed in parentheses. For the

copulation rate, chi-square test is applied. For the latency to copulation, Kruskal-Wallis

ANOVA and post hoc Mann-Whitney U tests are applied. Error bars indicate SEM. *p <

0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, ns indicates no significant difference.

1114

Figure 5 Virgin females with reduced EcR-A in pC1 neurons have reduced sexual receptivity.

(A) Knock-down of EcR-A in pC1 neurons driven by pC1-ss2-Gal4 significantly decreased the copulation rate and increased the latency to copulation. (B) Knock-down of EcR-B1 in pC1 neurons driven by pC1-ss2-Gal4 significantly prolonged the latency to copulation. (C-D) Knock-down of EcR-A (C) or EcR-B1 (D) in pC1 neurons driven by

pC1-ss1-Gal4 did not affect the copulation rate or the latency to copulation. (E)

Courtship index of wild-type males towards a female with the indicated genotype (n =

8). (F) The number of eggs laid by virgin females during the 3rd - 4th day after eclosion

when EcR-A was knocked down in pC1 neurons (n = 17-36). The UAS-EcR-A-RNAi

control causes a massive decrease in female fertility. (G) Knock-down of EcR-A in pC1

neurons decreased the opening of vaginal plate of virgin females compared with

controls (n = 8). (H) Knock-down of EcR-A in pC1 neurons increased the ovipositor

extrusion of virgin females compared with controls (n = 8). (I-K) Virgin female copulation

rate when EcR-A was knocked down in pC1 neurons temporally restricted by shifts from

18°C to 30°C. EcR-A was knocked down during the whole larval (I), pupal (J) and adult

(K) stages respectively. When the experiment was done at the pupal stage is the only

situation when the controls were both different from the experimental (J). The

comparison referred to flies with decreased EcR isoform in pC1 neurons. Female flies

for behavioral assay were 4-6-days-old adults. The number of female flies paired with

wild-type males is displayed in parentheses. For the copulation rate, chi-square test is

applied. For other comparisons, Kruskal-Wallis ANOVA and post hoc Mann-Whitney U

tests are applied. Error bars indicate SEM. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p <$

0.0001, ns indicates no significant difference.

1139

1140

1141

1142

1143

1144

1145

1146

1147

1148

1149

1150

1151

Figure 6 Reduced EcR-A in pC1 neurons induces the morphological changes.

(A1-A2) pC1-ss2-Gal4 expressing neurons appeared at the start of the pupal stage. (B-D) Reduced EcR-A in pC1 neurons induced more elaborated morphologies of pC1d axons, especially the extra vertical projection (EVP). The EVP regions of pC1d neurons was indicated by arrows. The morphological changes appeared on the 2nd day of the pupal stage (B1-B2) and retained to the adult stage including the 1st day (C1-C2) and the 4th day (D1-D2) of the adult stage. p0, the 1st day of the pupal stage; p2, the 2nd day of the pupal stage; A1, the 1st day of the adult stage; A4, the 4th day of the adult stage. (E-F) Fluorescence intensity of EVP in pC1d neurons on the 2nd day of the pupal stage (E) and the 4th day of the adult stage (F) was quantified when EcR-A was reduced in pC1 neurons (n=7). The quantified EVP regions were marked in (B) and (D) with orange ellipses. (G) pC1 neurons of the 4th day adults had comparable cell body number when EcR-A was reduced in pC1 neurons or not (n = 7). (H) Basal GCaMP6s signals in the LPC region of pC1 neurons when EcR-A was reduced in pC1 neurons (n = 22). LPC regions, the neurites extending from pC1 cell bodies, were marked with orange square in (D1) and (D2). The comparison referred to flies with decreased EcR isoform in pC1 neurons. Female flies in (G-H) were 4-days-old adults. Scale bars are 50 μm. For all comparisons, Mann-Whitney U test is applied. Error bars indicate SEM. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, ns indicates no significant difference.

1171

Figure 7 The function of PTTH on EcR-A and pC1 neurons.

(A) qRT-PCR for EcR-A when PTTH was deleted. Bars represent mean ± SEM. p

values are from Mann-Whitney U test (n = 8 for $Ptth^{Delete}$/+ and n = 6 for

$Ptth^{Delete}/Ptth^{Delete}$, each sample contains about 10 bodies of newly formed prepupae). *p

< 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, ns indicates no significant difference.

(B) Deletion of PTTH induced less elaborated morphologies of pC1d axons, especially

the extra vertical projection (EVP). The EVP regions of pC1d neurons was indicated by

arrows. Female flies were 4-days-old adults. (C) Fluorescence intensity of EVP in pC1d

neurons on the 4th day was quantified when PTTH was deleted (n=7). The quantified

EVP regions were marked in (B1) and (B2) with red ellipses. (D) Torso-Gal4 and Dsx-

LexA were labeled by stinger GFP and stingerRFP respectively. Arrows indicated the

overlap of GFP and RFP signals. Representative of five female brains. Scale bars, 50

μm. (E-F) Knock-down of Torso in pC1 neurons inhibited virgin female copulation rate

and enhanced latency to copulation. Females had similar locomotion speeds between

groups (F). The pC1-ss2-Gal4 control causes a massive decrease in female receptivity.

The comparison referred to pC1-ss2-Gal4/UAS-torso-RNAi. Female flies were 4-6-days-

old adults. The number of female flies paired with wild-type males is displayed in

parentheses. For the copulation rate, chi-square test is applied. For the latency to

copulation, Kruskal-Wallis ANOVA and post hoc Mann-Whitney U tests are applied.

Error bars indicate SEM, *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, ns indicates

no significant difference.

1193

1194

1195

1196

1197

1198

1199

1200

1201

1202

1203

1204

1205

1206

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216

1217

Figure 1—figure supplement 1 PTTH expression, weight, attractiveness and

locomotion behavior of *Ptth* null mutant virgin females.

(A) Brain of indicated genotype, immunostained with anti-PTTH antibody (green) and

counterstained with nc82 (magenta). Arrows show signals (green) stained with anti-

PTTH antibody. Female flies were within 10 hours after eclosion. Scale bars, 50 μm. (B)

The weights of 24h-old adult *Ptth*^{Delete}*/Ptth*^{Delete} null mutant females were significantly

higher than that of wild type females (Mann-Whitney U test, n=8 groups, 10 flies in each

group). (C) Courtship index of wild-type males during the first 5 min of courtship towards

a female with the indicated genotype (Kruskal-Wallis ANOVA and post hoc Mann-

Whitney U tests, n = 7-9). (D-E) Mean velocity had no significant change in

Ptth^{Delete}*/Ptth*^{Delete} null mutant females on the 1st day (D) and the 6th day (E) compared

with control females (Kruskal-Wallis ANOVA and post hoc Mann-Whitney U tests, n = 7-

1230   12). The comparison referred to $Ptth^{Delete}/Ptth^{Delete}$. Female flies for behavioral assay

1231   were 4-6-days-old adults. Error bars indicate SEM. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$,

1232   ****$p < 0.0001$, ns = not significant.

1233

1234

1235

1236

1237

1238

1239

1240

1241

1242

1243

1244

1245

1246

1247

1248

1249

1250

1251

1252

1253

1254

1255

1256

1257

1258

1259

1260

Figure 1—figure supplement 2 Effect of the expression of PTTH on female

receptivity.

(A) Overexpressing PTTH in PTTH neurons. The comparison referred to flies *Ptth-Gal4/+;UAS-PTTH/+.* Female flies for behavioral assay were 4-6 days-old. (B-C) Decreasing PTTH in PTTH neurons. The comparison referred to flies *Ptth-Gal4/+;UAS-PTTH-RNAi/+* and *Dsx-Gal4/+;UAS-PTTH-RNAi/+.* Female flies for behavioral assay were 2-days-old adults. (D) The fly brain of elav-Gal4>UAS-GFP were stained with PTTH antibody. The larvae flies were the wandering ones. The adult flies were within 10 hours after eclosion. Representative of 5 female brains. Scale bars, 50 μm.

1273

Figure 2—figure supplement 1 The function of PTTH neurons in female receptivity.

Expression pattern of Ptth-LexA in the brain revealed by anti-GFP (green) in wandering larvae CNS (A) and within-10-hours adult brain (B). Representative of five female flies. Scale bars, 50 μm. (C) The female receptivity when deceasing the expression of Dsx^F in PTTH neurons. The comparison referred to *Ptth-Gal4/+;UAS-Dsx^F-RNAi/+*. Female flies were 2-days-old adults. (D) PTTH neurons were inactivated during the whole larval, pupal and adult stages respectively by kir2.1, restricted by shifts from 18°C to 30°C. When the experiment was done at the larval stage is the only situation when the controls were both different from the experimental. The comparison referred to *Ptth-Gal4/tub-Gal80ts;kir2.1/+*. Female flies were 2-days-old adults. The number of female flies paired with wild-type males is displayed in parentheses. For the copulation rate, chi-square test is applied. Error bars indicate SEM. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, ns indicates no significant difference.

1288

Figure 2—figure supplement 2 The anatomical pattern of PTTH neurons

expressing PTTH at different developmental stages.

Expression pattern of Ptth-Gal4 in the brain revealed by anti-GFP from the 3rd larval

stage to the 4th day after eclosion. Arrows show PTTH signals (green) stained with anti-

GFP antibody. L3, the 3rd -instar larvae; wander, the wandering larvae; P4, the 4th day

of the pupal stage; A0h, the 1st hour of the adult stage; A3h, the 3rd hour of the adult

stage; A6h, the 6th hour of the adult stage; A9h, the 9th hour of the adult stage; A12h,

the 12th hour of the adult stage; A4d, the 4th day of the adult stage. Representative of

five female flies. Scale bars, 50 μm.

1298

1299

1300

1301

1302

1303

1304

1305

1306

1307

1308

Figure 2—figure supplement 3 PTTH neurons expressing PTTH do not regulate

virgin female copulation rate during adult stage.

PTTH neurons were activated during adult stage by dTrpA1 at 29°C. The female

copulation rate and the latency to copulation did not change significantly. The number of

female flies paired with wild-type males is displayed in parentheses. Female flies were

4-days-old adults. For the copulation rate, chi-square test is applied. For the latency to

copulation, Mann-Whitney U test is applied. Error bars indicate SEM. *p < 0.05, **p <

0.01, ***p < 0.001, ****p < 0.0001, ns indicates no significant difference.

1317

1318

1319

1320

1321

1322

1323

1324

1325

1326

1327

1328

1329

1330

1331

1332

1333

1334

1335

1336

Figure 5—figure supplement 1 Expression of EcR-A and EcR-B1 in pC1 and

vpoDN neurons.

(A-C) pC1 neurons were colabeled by pC1-ss2 driving UAS-mCD8-GFP (green, A) and

EcR-A antibodies (red, B). Magnification of green boxed region in (C) is shown in (D1–

D3). (E-G) pC1 neurons were colabeled by pC1-ss2 driving UAS-mCD8-GFP (green, E)

and EcR-B1 antibodies (red, F). Magnification of green boxed region in (G) is shown in

(H1–H3). (I-K) vpoDN neurons were colabeled by vpo-ss1 driving UAS-mCD8-GFP

(green, I) and EcR-A antibodies (red, J). Magnification of green boxed region in (K) is

shown in (L1–L3). (M-O) vpoDN neurons were colabeled by vpo-ss1 driving UAS-

mCD8-GFP (green, M) and EcR-B1 antibodies (red, N). Magnification of green boxed

region in (O) is shown in (P1–P3). Female flies were during prepupae stage. Scale bars

for magnified regions are 5 μm, for others are 50 μm.

1349

1350

1351

1352

1353

1354

1355

1356

1357

1358

1359

1360

1361

1362

1363

1364

1365

1366

1367

1368

1369

1370

1371

1372

1373

1374

1375

1376

1377

Figure 5—figure supplement 2 Reduced EcR in pC1 neurons reduces virgin female receptivity.

(A) Knock-down of EcR in pC1 neurons driven by pC1-ss2-Gal4 significantly decreased the copulation rate and increased the latency to copulation. (B) Mean velocity had no significant change when EcR-A was knocked down in pC1 neurons compared with controls (Kruskal-Wallis ANOVA and post hoc Mann-Whitney U tests, n = 8-11). The comparison referred to pC1-ss2-Gal4/UAS-EcR-RNAi (A) and pC1-ss2-Gal4/UAS-EcR-A-RNAi (B). Female flies were 4-6-days-old adults. The number of female flies paired with wild-type males is displayed in parentheses. For the copulation rate, chi-square test is applied. For the latency to copulation, Kruskal-Wallis ANOVA and post hoc Mann-Whitney U tests are applied. Error bars indicate SEM. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$, ns indicates no significant difference.

1404

Figure 5—figure supplement 3 Reduced EcR in vpoDN neurons has no effect on

virgin female receptivity.

(A-C) Knock-down of EcR-A in vpoDN neurons driven by vpoDN-ss1-Gal4, vpoDN-ss2-Gal4 and vpoDN-ss3-Gal4 had no effect on virgin female receptivity. (D-F) Knock-down of EcR-B1 in vpoDN neurons driven by vpoDN-ss1-Gal4, vpoDN-ss2-Gal4 and vpoDN-ss3-Gal4 had no effect on virgin female receptivity. The comparison referred to flies with decreased EcR isoform in vpoDN neurons. Female flies were 4-6-days-old adults. The number of female flies paired with wild-type males is displayed in parentheses. For the copulation rate, chi-square test is applied. For the latency to copulation, Kruskal-Wallis ANOVA and post hoc Mann-Whitney U tests are applied. Error bars indicate SEM. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, ns indicates no significant difference.

1416

1417

1418

Figure 5—figure supplement 4 Reduced EcR-A in pC1d neurons has no effect on virgin female receptivity.

(A) Knock-down of EcR-A in pC1d neurons had no effect on virgin female copulation rate and latency to copulation. The comparison referred to pC1d-Gal4/UAS-EcR-A-RNAi. Female flies were 4-6-days-old adults. The number of female flies paired with wild-type males is displayed in parentheses. For the copulation rate, chi-square test is applied. For the latency to copulation, Kruskal-Wallis ANOVA and post hoc Mann-Whitney U tests are applied. Error bars indicate SEM, *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, ns indicates no significant difference.

Supplementary File 1. The primers used for the verification of *Ptth^Delete* null mutant flies and for the real-time quantitative PCR of EcR-A when PTTH is deleted.

Figure 1—Source Data 1. Photo of nucleic acid electrophoresis and copulation time.

Figure 2—Source Data 1. Copulation time.

Figure 3—Source Data 1. Copulation time and walking speed.

Figure 4—Source Data 1. Copulation time.

Figure 5—Source Data 1. Copulation time, courtship index, number of eggs, number of vaginal plate opening (VPO), and number of ovipositor extrusion (OE).

Figure 6—Source Data 1. Fluorescence intensity, cell number and calcium activity.

Figure 7—Source Data 1. Relative mRNA level, fluorescence intensity, copulation time and walking speed.

Figure 1—figure supplement 1—Source Data 1. Body weight, courtship index and walking speed.

Figure 1—figure supplement 2—Source Data 1. Copulation time.

Figure 2—figure supplement 1—Source Data 1. Copulation time.

Figure 2—figure supplement 3—Source Data 1. Copulation time.

Figure 5—figure supplement 2—Source Data 1. Copulation time and walking speed.

Figure 5—figure supplement 3—Source Data 1. Copulation time.

1475

1476   **Figure 5—figure supplement 4—Source Data 1.** Copulation time.

1477