1 **Main Manuscript for**

3 **Prothoracicotropic hormone controls female sexual receptivity through the**

4 **function of ecdysone in pC1 neurons of *Drosophila***

5 Jing Li[1,2,†,*], Chao Ning[3,†], Yaohua Liu[2,4,†], Bowen Deng[5], Bingcai Wang[2,6], Kai Shi[2,6],
6 Rencong Wang[2,6], Ruixin Fang[1] and Chuan Zhou[1,2,6]

7
[1]*Institute of Molecular Physiology, Shenzhen Bay Laboratory, Shenzhen 518132, China*
[2]*State Key Laboratory of Integrated Management of Pest Insects and Rodents, Institute*
of Zoology, Chinese Academy of Sciences, Beijing 100101, China
[3]*CAS Key Laboratory of Genome Sciences and Information, Beijing Institute of*
Genomics, Chinese Academy of Sciences, Beijing 100101, China
[4]*Department of Plant Protection, Shanxi Agricultural University, Jinzhong 30801, China*
[5]*Chinese Institute for Brain Research, Peking-Tsinghua Center for Life Sciences,*
Zhongguancun Life Sciences Park, Beijing 102206, China
[6]*University of Chinese Academy of Sciences, Beijing 100049, China*
[†]*These authors contributed equally to this work*
----
*Jing Li, Email: lijing@szbl.ac.cn

Author Contributions: Jing Li, Conceptualization, Resources, Data curation, Writing -
original draft, review and editing, Funding acquisition; Chao Ning and Yaohua Liu, Data
curation; Bowen Deng, Resources; Bingcai Wang, Kai Shi, Rencong Wang, Software;
Ruixin Fang, Data curation; Chuan Zhou, Conceptualization, Funding acquisition.

Competing Interest Statement: The authors declare no competing interest.

Impact statement: Prothoracicotropic hormone and ecdysone belonging to the insect
PG axis modulate virgin female sexual receptivity through regulating the morphological
development of pC1b neurons.

Funding information: This work was supported by the Institute of Molecular
Physiology, Shenzhen Bay Laboratory (NO.21260061 to Chuan Zhou, S239201006 to
Jing Li), National Natural Science Foundation of China (NO.Y711241133 to Chuan
Zhou) and Strategic Priority Research Program of the Chinese Academy of Sciences
(NO.Y929731103 to Chuan Zhou).

Abstract

Female sexual receptivity is essential for reproduction of a species. Neuropeptides play the main role in regulating female receptivity. However, whether neuropeptides regulate the establishment of neural circuits for female sexual receptivity is unknown. Here we found the peptide hormone prothoracicotropic hormone (PTTH), which belongs to the insect PG axis, regulated virgin female receptivity through ecdysone during neural maturation in *Drosophila melanogaster*. We identified PG neurons expressing PTTH as doublesex-positive neurons, they regulated virgin female receptivity before the metamorphosis during the 3^{rd}-instar larval stage. Furthermore, the ecdysone receptor EcR-A in pC1 neurons regulated virgin female receptivity during metamorphosis. The reduced EcR-A in pC1 neurons induced abnormal morphological development of pC1 neurons without changing neural activity. Among all subtypes of pC1 neurons, the function of EcR-A in pC1b neurons was necessary for virgin female copulation rate. These suggested that the changes of synaptic connections between pC1b and other neurons decreased female copulation rate. Moreover, analysis of brain transcriptomes when EcR-A was reduced in pC1 neurons revealed that, additional genes were regulated downstream of EcR-A function in pC1 neurons. The PG axis has similar functional strategy as the HPG axis in mammals to trigger the juvenile–adult transition. Our work suggests a general mechanism underlying which the neurodevelopment during maturation regulates female sexual receptivity.

Keywords: prothoracicotropic hormone, *Drosophila*, female sexual receptivity,

ecdysone, pC1 neurons

Introduction

The success of copulation is important for the reproduction of a species. *Drosophila*

melanogaster provides a powerful system to investigate the neuronal and molecular

mechanism of sexual behaviors. Females decide to mate or not according to their

physiological status and the environmental condition (Dickson, 2008). Sexually mature

adult virgin females validate males after sensing the courtship song and male-specific

sex pheromone, receive courtship with pausing and opening the vaginal plate (Ferveur,

2010; Greenspan et al., 2000; Hall, 1994; Wang et al., 2021). If female is not willing to

mate, she may kick the legs, flick the wings, or extrude the ovipositor to deter males

(Connolly et al., 1973). Mated females reject males for several days after mating mainly

through more ovipositor extrusion and less opening the vaginal plate (Fuyama et al.,

1997; Wang et al., 2021). These options need the establishment of neural circuits for

female sexual receptivity. However, the associated mechanism of neural maturation and

the effect of neural maturation on female sexual receptivity is little known.

doublesex (*dsx*) and *fruitless* (*fru*) are the terminal genes in sex determination

regulatory hierarchy. They specify nearly all aspects of somatic sexual differentiation,

including the preparation for sexual behaviors (Dickson, 2008; Manoli et al., 2013;

Manoli et al., 2006; Mellert et al., 2012; Pavlou et al., 2013; Siwicki et al., 2009;

Yamamoto, 2007; Yamamoto et al., 2013). In males, male-specific *fru* protein (Fru[M])

(Billeter et al., 2006; Demir et al., 2005; Hall, 1978; Manoli et al., 2005; Stockinger et al.,

2005) and the male-specific *dsx* protein (Dsx[M]) (Kohatsu et al., 2011; Pan et al., 2014;

Pan et al., 2011; Rideout et al., 2010) are important for the male courtship behaviors. In

females, functional Fru protein does not exist, while the neurons in that *fru* P1 promoter

or *dsx* is expressed regulate some aspects of the female sexual behaviors (Kvitsiani et

al., 2006; Rideout et al., 2010). *Fru and dsx* are involved in regulating the sexual

dimorphism during neurodevelopment (Yamamoto et al., 2013). For instance, the sexual

dimorphism of P1 and mAL neurons which are all associated with male courtship and

aggression behaviors (Clowney et al., 2015; Hoopfer et al., 2015; Kimura et al., 2008;

Kohatsu et al., 2011; Pan et al., 2012; Sengupta et al., 2022; von Philipsborn et al.,

2011) is the result of regulation by Dsx and/or Fru (Ito et al., 2012; Kimura et al., 2008).

In the cis-vaccenyl acetate (cVA) pathway, which induces the courtship inhibiting in

males (Kurtovic et al., 2007; Wang et al., 2010), the first-order to the fourth-order

components are all fruGAL4-positive neurons and are either male-specific or sexually

dimorphic (Ruta et al., 2010). However, the role of Dsx[F] in neurodevelopment

associated with female sexual behaviors is little understood.

During postembryonic development, the PG axis triggers the juvenile–adult transition,

similar as the function of hypothalamic–pituitary–gonadal (HPG) axis in mammals

(Herbison, 2016; Pan et al., 2019). Hormones of the PG axis act to transform the larval

nervous system into an adult version (Truman et al., 2023). Ecdysone belonging to the

PG axis is the prime mover of insect molting and metamorphosis and is involved in all

phases of neurodevelopment, including neurogenesis, pruning, arbor outgrowth, and

cell death (Truman et al., 2023). The neurons read the ecdysteroid titer through two

isoforms of the ecdysone receptor, EcR-A and EcR-B1, according to spatial and

temporal conditions in the central nervous system (Riddiford et al., 2000; Truman et al.,

1994). EcR-A is required in fru P1-expressing neurons for the establishment of male-

specific neuronal architecture, and ecdysone receptor deficient males display increased

male–male courtship behavior (Dalton et al., 2009; Ganter et al., 2007). However, how

ecdysone regulates the neurodevelopment associated with female sexual receptivity,

especially the fru$^+$ and dsx$^+$ neurons, is unknown.

Much of studies to understand female sexual receptivity has focused on its regulation.

How a female respond to males is highly dependent on whether or not she has

previously mated. In virgin females, dsx$^+$ pCd neurons respond to the cVA, while dsx$^+$

pC1 neurons also respond to male courtship song (Zhou et al., 2014). The receptive

females open the vaginal plate (VPO) through activation of the dsx$^+$ vpoDN neurons

(Wang et al., 2021). After mated, sex peptide in the seminal fluid binds to the fru$^+$ dsx$^+$

sex peptide sensory neurons (SPSNs) in the female uterus. Then neuronal activity in

the dsx$^+$ sex peptide abdominal ganglion (SAG) neurons of the ventral nerve cord and in

the pC1 neurons is reduced (Avila et al., 2011; Feng et al., 2014; Häsemeyer et al.,

2009; Kubli, 2003; Wang et al., 2020b; Yang et al., 2009; Zhou et al., 2014). Therefore,

the sexual receptivity is reduced with less VPO and more ovipositor extrusion (OE)

which is controlled by dsx$^+$ DpN13 neurons (Wang et al., 2020a). In addition,

neuropeptides and monoamines play a critical role in regulation of the female

receptivity. The neuropeptides Drosulfakinin, myoinhibitory peptides and SIFamide are

involved in female sexual receptivity (Jang et al., 2017; Terhzaz et al., 2007; Wang et

al., 2022). As monoamines, dopamine, serotonin and octopamine are pivotal to female

sexual behaviors (Ishimoto et al., 2020; Ma et al., 2022; Neckameyer, 1998; Rezával et

al., 2014). So far, the identified neuropeptides and monoamines modulating female

sexual receptivity all function during the adult stage. However, whether neuropeptides

or monoamines regulate the establishment of neural circuits for female sexual

receptivity is unknown.

To explore the factors that regulate *Drosophila* virgin female receptivity especially

during neurodevelopment, we did a knock-out screen including most of CCT members.

We discovered a requirement for the prothoracicotropic hormone (PTTH) during

postembryonic development for virgin female receptivity. We also found that PG

neurons expressing PTTH are dsx$^+$ neurons. PTTH, a brain derived neuropeptide

hormone, is the primary promoter of the synthesis of steroid hormone 20-

hydroxyecdysone (20E) (McBrayer et al., 2007a; Rewitz et al., 2009). Indeed, the

enhanced virgin female receptivity due to the loss of PTTH could be rescued through

feeding 20E to the 3rd-instar larvae. Due to that 20E functions through its receptor EcR

(Riddiford et al., 2000), we then tested the function of EcR in pC1 neurons which

encode the mating status of females (Zhou et al., 2014). The reduced EcR-A expression

in pC1 neurons resulted in the unnormal anatomical pattern of pC1 neurons and the

reduced female copulation rate. Furthermore, the decreased female copulation rate was

due to the reduced EcR-A in pC1b neurons. To detect the molecular mechanism for the

regulation of pC1 neurons in virgin female receptivity, we tested the RNAseq when EcR-

A was reduced in pC1 neurons. We found some regulated genes, which provide clues

for our future research in the molecular mechanism by which pC1 neurons regulate

virgin female receptivity. Thus, in addition to demonstrating the function of PTTH in

virgin female receptivity during neural maturation, our study identified the necessary role

of the normal pC1b neural morphology in virgin female receptivity.

Results

PTTH modulates virgin female receptivity

In *Drosophila*, neuropeptides and monoamines, belonging to the chemoconnectome

(CCT) (Deng et al., 2019), play a critical role in regulation of the female receptivity. To

explore the factors that regulate virgin female receptivity especially during

neurodevelopment, we screened 108 chemoconnectome (CCT) knock-out lines

generated by the CRISPR-Cas9 system (Deng et al., 2019) (unpublished data). The

result showed that prothoracicotropic hormone (PTTH) might regulate virgin female

receptivity. The deletion mutant *ΔPtth* removed part of the 5' UTR and almost all coding

sequence and is a protein null (**Figure 1A**). We confirmed the PTTH knock-out flies by

using PCR analysis at the PTTH locus in genomic DNA samples (**Figure 1B**), by using

RT-PCR to identify the loss of PTTH transcripts in cDNA samples (**Figure 1C**) and by

detecting the immunoreactivity of PTTH in the central brain (**Figure 1H**). Primers used

are listed in **Supplementary File 1**. PTTH immunoreactivity was found in the brain of

wild-type and heterozygous flies (**Figure 1 H1 and H3**), but was absent in homozygous

ΔPtth flies (**Figure 1H2**). As the previous study, the *ΔPtth* larvae lacking PTTH undergo

metamorphosis with about 1 day delay compared with the wild type control (Shimell et

al., 2018) (data not shown). Besides, the *ΔPtth* adult male and female flies had the

significant increased weight than wild type flies (**Figure 1—figure supplement 1A**).

This is also consistent with that PTTH regulates developmental timing and body size in

Drosophila (Mcbrayer et al., 2007b; Shimell et al., 2018).

To confirm the function of PTTH, we tested virgin female receptivity of *ΔPtth* female

flies. We found that the virgin female losing PTTH had significantly higher copulation

rate and shorter latency to copulation than wild type flies (**Figure 1 D-G**). In addition, the

ΔPtth flies had higher copulation rate and lower latency to copulation compared to

heterozygous null mutant females within 2 days (**Figure 1 D-E**) and within 3 days,

respectively (**Figure 1 D-F**). The enhanced virgin female receptivity had no relationship

either with the attractivity or with the locomotion activity of virgin females (**Figure 1—**

figure supplement 1B-D). These results suggested that PTTH participates in virgin

female receptivity in a dose-dependent manner.

Furthermore, we carried out genetic rescue experiments to further confirm the function

of PTTH in modulating virgin female receptivity. We used the pan-neuronal driver

elavGAL4 to drive UAS-PTTH expression in PTTH mutant background. We detected the

PTTH signals using PTTH antibody in the rescued female brains (**Figure 1H4**). We

found that neuron-specific expression of PTTH could restore the enhanced copulation

rate and shorter latency to copulation in *ΔPtth* virgin females (**Figure 1I**). These results

confirmed that PTTH modulates virgin female receptivity.

Dsx+ PG neurons regulate virgin female receptivity.

We used new PtthGAL4 and PtthLexA which inserts GAL4 or LexA sequence before the

stop codon of the *Ptth* gene (Deng et al., 2019) to label and manipulate PG neurons

expressing PTTH. The labeled neurons were the same as reported before (McBrayer et

al., 2007a; Yamanaka et al., 2013), a pair of bilateral neurosecretory cells in the brain

directly innervating the prothoracic gland during the larval stage (**Figure 2A and Figure**

2—figure supplement 1A). The newly emerged flies had the similar anatomical pattern

with that of the larval stage (**Figure 2B and Figure 2—figure supplement 1B**).

However, while the prothoracic gland cells are gradually degenerating during pharate

adult development (Dai et al., 1991; Roy et al., 2018), the pattern of PG neurons labeled

by PtthGAL4 > UAS-mCD8GFP gradually could not be found before the 10th hour after

eclosion (**Figure 2—figure supplement 2**).

Most identified neurons associated with female sexual behaviors express *doublesex*

gene. We asked whether PG neurons are a part of the doublesex circuitry or not.

Double labeling of dsxLexA and PtthGAL4 neurons (LexAop-tomato,UAS-

stingerGFP/PtthLexA;dsxGAL4/+) revealed that PG neurons are all doublesex-positive

(**Figure 2C**). We then used an intersectional strategy to visualize overlapped expression

between dsxLexA and PtthGAL4 (UAS > stop > myrGFP/+;LexAop2-

FlpL,dsxLexA/PtthGAL4). We observed all PG neurons with GFP signals (**Figure 2D**).

These results suggested that PG neurons are dsx$^+$ neurons.

We then analyzed whether PG neurons are involved in the modulation of virgin female

receptivity. First, we activated PG neurons transiently in adult virgin females by driving

the temperature-sensitive activator dTrpA1 (Hamada et al., 2008) using PtthGAL4. PG

neurons were activated at 29°C compared with the control treatment at 23°C. No

significantly different copulation rate or latency to copulation was detected (**Figure 2—**

figure supplement 3A-C). This suggested that PG neurons do not regulate virgin

female receptivity during the adult stage.

To identify the detail time for the function of PG neurons in virgin female receptivity, we

inactivated PG neurons through kir2.1 under the control of the temporal and regional

gene expression targeting system (McGuire et al., 2004). The inactivation of PG

neurons during larval stage enhanced virgin female copulation rate significantly (**Figure**

2E). However, when PG neurons were inactivated during pupal or adult stages, virgin

female copulation rate did not change significantly (**Figure 2 F-G**). Furthermore, we

activated PG neurons at different stages overlapping the postembryonic larval

developmental time using dTrpA1 (**Figure 3A**). Stage 1 was from the 1st-instar larvae to

six hours before the 3rd-instar larvae. Stage 2 was from six hours before the 3rd-instar

larvae to the end of the wandering larvae. Stage 3 was from the end of wandering

larvae to the end of the 2nd day of the pupal stage. Stage 4 was from the end of the 2nd

day of the pupal stage to the eclosion of adults. The copulation rate did not change

significantly when activating PG neurons during the stage 1, stage 3 or stage 4 (**Figure**

3B, 3D and 3E). However, we found the significant lower copulation rate and the longer

latency to copulation only when PG neurons were activated during the stage 2 (**Figure**

3C and Figure 3F). The defected copulation was not due to a lower locomotion activity

of virgin females (**Figure 3G**). Taken together, our findings indicated that the activity of

dsx$^+$ PG neurons negatively regulate virgin female receptivity during the stage from the

start of the 3rd-instar to the end of wandering stage.

PTTH modulates virgin female receptivity through ecdysone.

The 3rd-intar larval stage is the critical stage for the initiation of metamorphosis involving

the synthesis of ecdysone (Imura et al., 2020; Lavrynenko et al., 2015; Shimell et al.,

2018). To test whether PTTH regulates virgin female receptivity through regulating the

synthesis of ecdysone, we rescued the enhanced female receptivity by feeding 20E to

the 3rd-intar larval *ΔPtth* flies. The enhanced copulation rate and shorter latency to

copulation of the *ΔPtth* flies were rescued to the comparable level of wild type females

(**Figure 4**). Furthermore, the wild type females fed by 20E had no significantly different

copulation rate and latency to copulation compared with the wild type females fed by the

same volume of 95% ethanol which is the solvent of 20E (**Figure 4**). This suggested

that PTTH regulates virgin female receptivity through the titer of ecdysone.

Ecdysone receptor EcR-A in pC1 neurons regulates virgin female copulation rate.

Given that PTTH regulates virgin female receptivity through ecdysone which acts on its

receptor EcR, we then asked that whether ecdysone regulates the function of neurons

associated with virgin female receptivity through EcR. pC1 and vpoDN neurons are two

main dsx$^+$ neurons involved in virgin female receptivity (Wang et al., 2021; Zhou et al.,

2014). pC1 neurons encode the mating status of female flies, vpoDN neurons regulate

the opening of vaginal plate when females attend to accept males. EcR-A and EcR-B1

are the two prominently expressed ecdysone receptors in the CNS (Riddiford et al.,

2000). First, we tested the expression of EcR-A and EcR-B1 in these two neurons on

the 2nd day of the pupal stage when ecdysone functions as the main mover in the

metamorphosis (Dalton et al., 2009; Truman et al., 1994). The GFP signals labeled by

pC1-ss2-GAL4 and vpoDN-ss1-GAL4 were merged well with the signals of both EcR-A

and EcR-B1 antibodies respectively (**Figure 5—figure supplement 1**). This revealed

that EcR-A and EcR-B1 express in both pC1 and vpoDN neurons. We then tested the

function of EcR in pC1 and vpoDN neurons through reducing the expression of EcR-A

and EcR-B1 respectively. We used the split GAL4 for pC1 and vpoDN neurons to drive

the UAS-EcR-RNAi. First, we reduced the expression of all EcR isoforms in pC1

neurons, this decreased the copulation rate and prolonged the latency to copulation

significantly (**Figure 5—figure supplement 2A**). Furthermore, we reduced the

expression of EcR-A in pC1 neurons. The virgin female had the significant lower

copulation rate and longer latency to copulation (**Figure 5A**). The reduced copulation

rate had no relationship with the attractivity (**Figure 5E**) and the locomotion activity of

virgin females (**Figure 5—figure supplement 2B**). When reducing the expression of

EcR-B1 in pC1 neurons, virgin females had the significant longer latency to copulation

but the comparable copulation rate to controls (**Figure 5B**). However, reducing the

expression of EcR-A (**Figure 5—figure supplement 3 A-C**) and EcR-B1 (**Figure 5—**

figure supplement 3 D-F) using three split vpoDNGAL4s in vpoDN neurons all did not

affect virgin female receptivity. This suggested that the expression of EcR-A in pC1

neurons regulates virgin female copulation rate, but EcR isoforms in vpoDN neurons do

not modulate virgin female receptivity.

Two split-GAL4 drivers for pC1 neurons had been obtained previously. pC1-ss1-GAL4

labels pC1-a, c and -e neurons, and pC1-ss2-GAL4 labels all pC1-a, -b, -c, -d and -e

neurons (Wang et al., 2020b). We also tested virgin female receptivity when EcR-A or

EcR-B1 were reduced in pC1-a, -c and -e neurons simultaneously using pC1-ss1-GAL4

respectively. While the copulation rate or the latency to copulation did not change

significantly (**Figure 5 C-D**). This suggested that, pC1b and/or pC1d neurons are

necessary for the functions of EcR-A and EcR-B1 in pC1 neurons on virgin female

receptivity. Whether pC1d is involved in the regulation of female receptivity is uncertain

(Deutsch et al., 2020; Schretter et al., 2020; Taisz et al., 2023). However, when

reducing EcR-A in pC1d neurons alone using the specific split GAL4 SS56987

(Schretter et al., 2020), virgin female receptivity including copulation rate and latency to

copulation did not change significantly compared with controls (**Figure 5—figure**

supplement 4). These results suggested that the function of EcR-A in pC1b neurons is

necessary for virgin female copulation rate.

As recently mated females may reduce sexual receptivity and increase egg laying (Avila

et al., 2011; Kubli, 2003). we asked whether the decreased copulation rate induced by

EcR-A could be a post-mating response and correlate with elevated egg laying. To

address this, we examined the number of eggs laid by virgin females when EcR-A was

reduced in pC1 neurons. We found that manipulation of EcR-A did not enhance egg

laying significantly in virgin females (**Figure 5F**). Meanwhile, we further analyzed

whether reduction of EcR-A in pC1 neurons regulates the opening of vaginal plate

(VPO) or the ovipositor extrusion (OE). We found that reducing the EcR-A expression in

pC1 neurons lead to the significantly less VPO and more OE (**Figure 5 G-H**). These

results suggested that reduced EcR-A expression in pC1 neurons results in the similar

phenotype to that of mated females.

EcR-A participates in the morphological development of pC1 neurons.

EcR isoforms have distinct temporal and spatial expression patterns in the CNS

(Riddiford et al., 2000; Truman et al., 1994). It is unknown when EcR-A functions in pC1

neurons for virgin female receptivity. Thus, we examined virgin female receptivity when

EcR-A expression was conditionally reduced through RNAi via the pC1-ss2-GAL4 under

the control of the temporal and regional gene expression targeting system (McGuire et

al., 2004). EcR-A was reduced during the larval, pupal and adult stage respectively

(**Figure 5 I-K**). Only during the pupal stage, reducing EcR-A made the significant lower

copulation rate and longer latency to copulation (**Figure 5J**). The result suggested that

EcR-A in pC1 neurons plays a role in virgin female receptivity during metamorphosis.

This is consistent with that PTTH regulates virgin female receptivity before the start of

metamorphosis which is around the puparium formation.

We then tested how EcR-A functions in pC1 neurons to modulate virgin female

receptivity. First, we tested the morphology of pC1 neurons when reducing the

expression of EcR-A in pC1 neurons. We found that the morphology of pC1 neurons

appeared after the formation of the white pupa (Figure **6A1**). The reduced EcR-A

expression induced the more elaborated morphologies of the pC1-d/e cells, especially

the extra vertical projection (EVP) near the midline of brains (**Figure 6 B-D**) (Deutsch et

al., 2020). These changes exhibited from the second day of the pupal stage (**Figure 6**

B1-B2 and 6E) and maintained at the adult stage (**Figure 6 D1-D2 and 6F**). Meanwhile,

the number of pC1 cell bodies in adult flies when EcR-A was reduced were the same as

that of wild type flies (**Figure 6G**). Previous studies suggested that pC1d cells serve as

a hub within the central brain for dsx$^+$ and fru$^+$ neurons (Deutsch et al., 2020). Thus, the

unnormal development of pC1d neurons may induce the changes between pC1d

neurons and other dsx$^+$ and fru$^+$ neurons to affect associated behaviors.

Furthermore, we asked whether reduced female copulation rate was due to that EcR-A

expression affected the activity of pC1 neurons. Because all pC1 cells characterized so

far project to the lateral junction of the lateral protocerebral complex (LPC) (Kimura et

al., 2015; Rezával et al., 2016; Scheffer et al., 2020; Wang et al., 2020b; Wu et al.,

2019; Zhou et al., 2014), we expressed GCamp6s in all pC1 neurons and tested the

calcium signals in the lateral junction of LPC when EcR-A was knocked down (**Figure 6**

D1-D2). Reduced EcR-A did not induce significantly different calcium responses in the

LPC (**Figure 6H**). Thus, our results suggested that the decreased female copulation

rate induced by reduced EcR-A in pC1 neurons was mainly due to the morphological

changes of pC1b neurons, which then modulate the connections of pC1b neurons with

other neurons.

Reduction of EcR-A in pC1 neurons affects gene expression.

To further detect the factors that were regulated downstream of EcR-A in pC1 neurons

for the function in virgin female receptivity, we tested the transcriptome of brain in adult

virgin females when EcR-A was knocked down in pC1 neurons. We focused on the

differently expressed genes of the 4th day after eclosion when mating behaviors were

tested. We identified 527 differentially expressed (p < 0.01) genes, 123 of which passed

a false discovery rate (FDR) cutoff of 0.01 (**Figure 7, Supplementary File 2 and**

Supplementary File 3).

The gene encoding dopamine beta-monooxygenase (DBM) was the top most down-

regulated gene (**Figure 7 A-B**). The mammal homolog of DBM is monooxygenase

DBH-like 1 (Moxd1). Moxd1 has the similar structure as the mammal dopamine β-

hydroxylase (DBH) which is the enzyme for transition from dopamine to norepinephrine

(Park et al., 1976; Prigge et al., 2000; Vendelboe et al., 2016). This implied that DBM is

probably related to the dopamine metabolism. It will be interesting to study the function

of DBM in virgin female receptivity. Besides, CG30428 protein was also down-regulated

(**Supplementary File 2**). It is predicted to be active in nucleus, suggesting its role in

gene expression and cellular metabolism. This is consistent with that reduction of EcR-

A in pC1 neurons induced the unnormal anatomical morphology of pC1 neurons, which

may due to the unnormal cell death and cellular differentiation during development.

Discussion

In this study, we found that peptide hormone PTTH modulates virgin female receptivity

through ecdysone during neural maturation. PG neurons expressing PTTH are

doublesex-positive and regulate virgin female receptivity during the 3rd-instar larval

stage. Furthermore, ecdysone receptor EcR-A functions in pC1 neurons to regulate

virgin female copulation rate during the metamorphosis mainly through modulating the

anatomical morphology of pC1b neurons. Taken together, our results identified the

function of PTTH-ecdysone and their downstream pathway, EcR-A in pC1 neurons, on

regulating virgin female copulation rate during neural maturation mainly through

modulating the morphology of pC1b neurons.

Our results suggested a regulatory role of PTTH in virgin female receptivity. Even

though insects and mammals represent highly diverged classes, insects have evolved a

similar strategy for triggering the juvenile–adult transition (Herbison, 2016; Pan et al.,

2019). The juvenile–adult transition involves the hypothalamic–pituitary–gonadal (HPG)

axis in mammals and the PG axis in insects. Among the neurons belonging to the axis,

PG neurons and GnRH neurons have the similar function to stimulate the PG gland and

pituitary gland to release hormones which trigger maturation, respectively. It will be

interesting to study the function of GnRH neurons in the mammal sexual behaviors.

Most of neurons regulating sexual behaviors in female flies are dsx$^+$ neurons. Our

results showed that PG neurons are also dsx$^+$ neurons. This suggested that PG

neurons have relationships with other dsx$^+$ neurons and the juvenile–adult transition is

regulated by *doublesex* gene.

In our study, PTTH regulates virgin female receptivity in an ecdysone-dependent

manner before the peak of ecdysone for the metamorphosis. Ecdysone functions

through its receptor EcR which is involved in all phases of the nervous system

development. Previous studies have demonstrated that the fru$^+$ neuron development

need EcR-A in male *Drosophila melanogaster*. Furthermore, reduced EcR-A in fru$^+$

neurons induced the male-male courtship (Dalton et al., 2009). In females, we detected

the expression of EcR-A and EcR-B1 in both dsx$^+$ pC1 and dsx$^+$ vpoDN neurons.

However, EcR regulates virgin female receptivity in pC1 neurons but not in vpoDN

neurons. This may due to that EcR does not regulate the morphology of vpoDN neurons

and thus the synaptic connections with other neurons. Alternatively, EcR regulates the

morphology of vpoDN neurons without changing functional connectivity or gene

expressions within associated neurons. It is worth noting that the losing of PTTH

resulted in the enhanced virgin female receptivity, which is contrary to the reduced

receptivity when knocking down EcR-A in pC1 neurons. This may due to that 20E and

EcR regulates multiple neurodevelopment associated with virgin female receptivity

when PTTH is knocked out, but not only functions in pC1 neurons. This may also due to

that PTTH mutant results in increased EcR expression in pC1 neurons.

Reduced EcR-A expression in all pC1 neurons lead to the decreased copulation rate,

while reduced EcR-A in pC1-a, c and e simultaneously did not reduce the copulation

rate significantly. This suggested that EcR-A plays the critical role in pC1-b or/and -d

neurons for regulating virgin female receptivity. Our results revealed that reduced EcR-

A induced the more elaborated morphologies of pC1d neurons. Previous studies

detected the synaptic connections between the axons of pC1d and the dendrites of

DNp13 neurons (Deutsch et al., 2020; Mezzera et al., 2020). DNp13 neurons are

command neurons for ovipositor extrusion. When females extruded, the ovipositor

physically impedes copulation (Mezzera et al., 2020; Wang et al., 2020a). However,

reduced EcR-A expression in pC1d neurons did not affect virgin female receptivity

(**Figure 5—figure supplement 4**). This might be due to three possibilities. First, the

more elaborated morphologies of pC1d neurons did not affect synaptic connections

between pC1d and DNp13 neurons. Second, the unchanged pC1 neural activity could

not affect the neural activity of DNp13 neurons. Third, the morphological change of

pC1d neurons is not sufficient for the decreased copulation rate. To sum, these suggest

that the morphological change of pC1b neurons is necessary for the decreased

copulation rate. However, due to the lack of pC1b drivers, we could not rule out

morphological changes in pC1b neurons when EcR-A was reduced.

We identified DBM as the mostly down-regulated gene when EcR-A was reduced in

pC1 neurons. DBM is the insect homolog of MOXd1 in mammals. However, the function

of MOXd1 has not been identified but is predicted to hydroxylate a hydrophobic

substrate (Xin et al., 2004). The similar structure between MOXd1 and dopamine beta-

monooxygenase in mammals suggests the possible role of DBM in dopamine

metabolism (Cubells et al., 2004; Kim et al., 2002; Timmers et al., 2004). Besides,

previous study identified the Moxd1 as a useful marker for the sexually dimorphic nuclei

and may be involved in the regulation of sex-biased physiology and behaviors in

mammals (Tsuneoka et al., 2017). These make it interesting to further study on the

molecular mechanism by which DBM and Moxd1 regulate sexual behaviors in flies and

mammals respectively.

Taken together, this study demonstrates that the peptide hormone PTTH, expresses in

dsx$^+$ PG neurons, regulates virgin female receptivity during neural maturation. It also

revealed that the morphological development of pC1b neurons regulated by EcR-A is

necessary for virgin female copulation rate. This work extends the understanding of how

neurodevelopmental processes regulate adult sexual behavior.

Materials and methods

Key Resources Table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
antibody	Mouse anti-Bruchpilot antibody (nc82)	Developmental Studies Hybridoma Bank	Cat# nc82, RRID: AB_2314866	IHC (1:40)
antibody	Mouse Anti-Drosophila ecdysone receptor (EcR-A) Monoclonal Antibody	Developmental Studies Hybridoma Bank	Cat# 15G1a (EcR-A), RRID: AB_528214	IHC (1:10)
antibody	mouse Anti-Drosophila ecdysone receptor (EcR-B1) Antibody	Developmental Studies Hybridoma Bank	Cat# AD4.4(EcR-B1), RRID: AB_2154902	IHC (1:10)
antibody	Rabbit polyclonal anti-GFP	Thermo Fisher Scientific	Cat# A-11122, RRID: AB_221569	IHC (1:1000)
antibody	chicken polyclonal anti-GFP	Thermo Fisher Scientific	Cat# A10262, RRID: AB_2534023	IHC (1:1000)
antibody	Goat anti-rabbit, Alexa Fluor 488	Thermo Fisher Scientific	Cat# A-11034, RRID: AB_2576217	IHC (1:500)
antibody	Goat anti-chickent, Alexa Fluor 488	Thermo Fisher Scientific	Cat#A-11039; RRID: AB_2534096	IHC (1:500)
antibody	Goat anti-mouse, Alexa Fluor 488	Thermo Fisher Scientific	Cat# A-11029, RRID: AB_2534088	IHC (1:500)
antibody	Goat anti-rabbit, Alexa Fluor 546	Thermo Fisher Scientific	Cat# A-11010, RRID: AB_2534077	IHC (1:500)
antibody	Rabbit polyclonal anti-RFP	Thermo Fisher Scientific	Cat# R10367, RRID: AB_10563941	IHC (1:500)
antibody	Goat anti-mouse, Alexa Fluor 647	Thermo Fisher Scientific	Cat# A-21235, RRID: AB_2535804	IHC (1:500)
antibody	Rabbit polyclonal anti-PTTH	Zhou Lab, Chinese Academy of Sciences, this paper	N/A	IHC (1:1300)
chemical compound, drug	Paraformaldehyde (PFA)	Electron Microscopy Sciences	Cat#15713	8% PFA diluted in 1XPBS at 1:4 or 1:2
chemical compound, drug	DPX Mountant	Sigma-Aldrich	Cat#44581	
chemical compound, drug	Normal goat serum	Sigma-Aldrich	Cat#G9023	
chemical compound, drug	20-hydroxyecdysone	Cayman	Cat#16145	dissolved in 95% ethanol, 0.2 mg/ml
chemical compound, drug	TRIzol	Ambion	Cat#15596018	
Genetic reagent (*D.melanogaster*)	*LexAop2-mCD8::GFP*	Bloomington Stock Center	BL#32203	
Genetic reagent (*D.melanogaster*)	*;;UAS-mCD8::GFP*	Bloomington Stock Center	BL#32194	
Genetic reagent (*D.melanogaster*)	*;UAS-mCD8::GFP;*	Bloomington Stock Center	BL#5137	
Genetic reagent (*D.melanogaster*)	*UAS-dTrpA1/cyo*	Garrity Lab, Brandeis University	N/A	
Genetic reagent (*D.melanogaster*)	*UAS-Kir2.1*	Bloomington Stock Center	BL#6595	

Genetic reagent (*D.melanogaster*)	*PtthGAL4*	Rao Lab, Peking University	N/A	
Genetic reagent (*D.melanogaster*)	*PtthLexA*	Rao Lab, Peking University	N/A	
Genetic reagent (*D.melanogaster*)	*⊿PTTH*	Rao Lab, Peking University	N/A	
Genetic reagent (*D.melanogaster*)	*UAS-PTTH*	Zhou Lab, Chinese Academy of Sciences, this paper	N/A	
Genetic reagent (*D.melanogaster*)	*isoCS*	Rao Lab, Peking University	N/A	
Genetic reagent (*D.melanogaster*)	*elav-GAL4*	Rao Lab, Peking University	N/A	
Genetic reagent (*D.melanogaster*)	*UAS-GFPStinger*	Janelia Research Campus	N/A	
Genetic reagent (*D.melanogaster*)	*LexAop-tomato*	Janelia Research Campus	N/A	
Genetic reagent (*D.melanogaster*)	*LexAop2-FlpL*	Janelia Research Campus	N/A	
Genetic reagent (*D.melanogaster*)	*UAS > stop > mCD8-GFP*	Janelia Research Campus	N/A	
Genetic reagent (*D.melanogaster*)	*dsxGAL4*	Janelia Research Campus	N/A	
Genetic reagent (*D.melanogaster*)	*dsxLexA*	Janelia Research Campus	N/A	
Genetic reagent (*D.melanogaster*)	*tub-GAL80^{ts}*	Pan Lab, Southeast University	BL#7018	
Genetic reagent (*D.melanogaster*)	*pC1-ss1-GAL4*	Wang Lab, Lingang Laboratory	N/A	
Genetic reagent (*D.melanogaster*)	*pC1-ss2 GAL4*	Wang Lab, Lingang Laboratory	N/A	
Genetic reagent (*D.melanogaster*)	*vpoDN-ss1-GAL4*	Wang Lab, Lingang Laboratory	N/A	
Genetic reagent (*D.melanogaster*)	*vpoDN-ss2-GAL4*	Wang Lab, Lingang Laboratory	N/A	
Genetic reagent (*D.melanogaster*)	*vpoDN-ss3-GAL4*	Wang Lab, Lingang Laboratory	N/A	
Genetic reagent (*D.melanogaster*)	*UAS-EcR-RNAi*	Bloomington Stock Center	BL#9327	
Genetic reagent (*D.melanogaster*)	*UAS-EcR-A-RNAi*	Bloomington Stock Center	BL#9328	
Genetic reagent (*D.melanogaster*)	*UAS-EcR-B1-RNAi*	Bloomington Stock Center	BL#9329	
Genetic reagent (*D.melanogaster*)	*pC1d-GAL4*	Bloomington Stock Center	BL#86847	
recombinant DNA reagent	pBSK-attP-3P3-RFP-loxP	Bowen Deng et al., 2019	N/A	

recombinant DNA reagent	pBSK-attB-loxP-myc-T2A-Gal4-GMR-miniwhite	Bowen Deng et al., 2019	N/A	
recombinant DNA reagent	pBSK-attB-loxP-V5-T2A-LexA::p65-GMR-miniwhite	Bowen Deng et al., 2019	N/A	
software, algorithm	MATLAB	MathWorks, Natick, MA	https://www.mathworks.com/products/matlab.html	
software, algorithm	ImageJ	National Institutes of Health	https://imagej.nih.gov/ij/	
software, algorithm	Prism 7	GraphPad	https://www.graphpad.com/	
software, algorithm	R 4.1.3	RStudio	https://www.r-project.org	

Fly stocks

Flies were reared on standard cornmeal-yeast medium under a 12 hr:12 hr dark:light

cycle at 25°C and 60% humidity. All the knock-out lines in this study for screening have

been published [49]. The following strains were obtained from Dr. Yi Rao: *isoCS* (wild-

type), *ΔPtth*, *PtthGAL4, PtthLexA, elavGAL4 and UAS-Kir2.1* (BL#6595). UAS-dTrpA1

was a gift from Dr. Paul Garrity. *UAS-GFPStinger*, *LexAop-tomato, LexAop2-FlpL,*

UAS > stop > mCD8-GFP, *dsxGAL4* and *dsxLexA* (Mellert et al., 2010) have been

described previously (Pfeiffer et al., 2008; Pfeiffer et al., 2010) and are obtained from

Janelia Research Campus. *tub-GAL80^{ts}* (BL#7018) was provided by Dr. Yufeng Pan.

pC1-ss1-GAL4, pC1-ss2 GAL4, vpoDN-ss1-GAL4, vpoDN-ss2-GAL4 and *vpoDN-ss3-*

GAL4 were provided by Dr. Kaiyu Wang. The following lines were obtained from the

Bloomington Drosophila Stock Center: *UAS-EcR-RNAi* (BL#9327), *UAS-EcR-A-RNAi*

(BL#9328), *UAS-EcR-B1-RNAi* (BL#9329), *UAS-mCD8::GFP* (BL#32194), *LexAop2-*

mCD8::GFP (BL#32203), *UAS-mCD8::GFP* (BL#5137) and *pC1d-GAL4* (BL#86847).

Behavioral assays

Flies were reared at 25°C and 60% humidity under a 12 hr light:12 hr dark cycle. Virgin

females and wild-type males were collected upon eclosion, placed in groups of 12 flies

each and aged 4–6 days (except for the assays for PTTH mutant on different days after

eclosion, and the molecular rescue assay for the 24h-old females) before carrying out

behavioral assay except for the transient thermogenetic experiments. Female receptivity

assays were conducted as previously described (Wang et al., 2022; Zhou et al., 2014).

A virgin female of defined genotype and a wild type male were gently cold anesthetized

and respectively introduced into two layers of the courtship chambers separated by a

removable transparent film. The flies were allowed to recover for at least 45 min before

the film was removed to allow the pair of flies to contact. The mating behavior was

recorded using a camera (Canon VIXIA HF R500) for 30 min at 17 fps for further

analysis.

For transient activation experiment by dTrpA1 in adult stage, flies were reared at 23°C.

Flies were loaded into courtship chamber and recovered for at least 30 min at 23°C,

then were placed at 23°C (control group) or 29°C (experimental group) for 30 min prior

to removing the film and videotaping. For activation experiment by dTrpA1 during

development, flies were reared at 29°C during the specific stages compared with the

controls who were reared at 23°C all the time. Flies were loaded into courtship chamber

and recovered for at least 45 min at 23°C prior to removing the film and videotaping.

Quantification and statistical analysis of female receptivity behavior

Two parameters including copulation rate and latency to copulation were used to

characterize receptivity and we got the data sets of two parameters from the same flies.

The time from removing the film to successful copulation was measured for each

female. The number of females that had engaged in copulation by the end of each 1

min interval within 30 min were summed and plotted as a percentage of successful

copulation. The latency to copulation was the time from removing the film to successful

copulation. All the time points that female successfully copulated were manually

analyzed and the data of unhealthy flies were discarded.

Temporally restricted RNAi

tub-GAL80^{ts} crosses were reared at either 18^{o}C for control groups or 30^{o}C for

experimental groups. Virgin females were collected at eclosion and were placed in

groups of 12 flies each and aged 4–6 days before carrying out behavior assay. Assays

were tested at 23^{o}C.

Male courtship index

Courtship index was defined as the proportion of time the male followed, oriented

toward and attempted to copulate the female within 5 min of courtship initiation, marked

by the initial orientation toward and following the female.

Vaginal plate opening and ovipositor extrusion

A virgin female of defined genotype and a wild type male were aspirated into the

courtship chambers and respectively introduced into two layers of the courtship

chambers separated by a removable transparent film. The flies were allowed to recover

for 30 min before the film was removed. To allow visualization of vaginal plate opening,

we recorded uncompressed image sequences at 896 x 896 pixels and 50 frames per

second using a Photron Mini AX camera (Photron) with an AF-S VR Micro-Nikkor

105mm lens (Nikon). Instances of vaginal plate opening and ovipositor extrusion were

scored blind to genotype from frame by-frame playback during the first 5 min of

courtship or until copulation if it occurred within 5 min. Courtship initiation was defined

as the male orienting toward and beginning to follow the female. Rare trials with fewer

than 30 s of total courtship were discarded.

Locomotion assays

The rearing and experimental conditions in locomotion assays were the same as that in

the corresponding female receptivity assays, excepting that individual females were

loaded in the chambers without males. Spontaneous movements of the flies were

recorded with a camera (Canon VIXIA HF R500) for 30 min at 30 fps for further

analysis. The activity of flies during the middle 10 min was analyzed to calculate the

average walking speed using Ctrax software.

Egg Laying

Virgin females were collected upon eclosion and one fly was housed on standard

medium at 25°C, 60% relative humidity, 12 hr light:12 hr dark and allowed to lay eggs in

single vials. Each fly was transferred into new food tube every 24 hr. The number of

eggs was counted at the end of each 24 hr period. The numbers during the 3rd and 4th

day were summed for statistics and plot.

Immunohistochemistry

Whole brains of flies were dissected in 1x PBS and fixed in 2% paraformaldehyde

diluted in 1x PBS for 55 min at room temperature. The samples were washed with PBT

(1x PBS containing 0.3% Triton X-100) for 1 hour (3 x 20 min), followed by blocking in

5% normal goat serum (Blocking solution, diluted in 0.3% PBT) for 1 hour at room

temperature. Then, the samples were incubated in primary antibodies (diluted in

blocking solution) for 18–24 hours at 4°C. Samples were washed with 0.3% PBT for 1

hour (3 x 20 min), then were incubated in secondary antibodies (diluted in blocking

solution) for 18–24 hours at 4°C. Samples were washed with 0.3% PBT for 1 hour (3 x

20 min), then were fixed in 4% paraformaldehyde for 4 hours at room temperature. After

washed with 0.3% PBT for 1 hour (3 x 20 min), brains were mounted on poly-L-lysine-

coated coverslip in 1x PBS. The coverslip was dipped for 5 min with ethanol of

30%→50%→70%→95%→100% sequentially at room temperature, and then dipped for

5 min three times with xylene. Finally, brains were mounted with DPX and allowed DPX

to dry for 2 days before imaging. Primary antibodies used were: chicken anti-GFP

(1:1000; Life Technologies #A10262), rabbit anti-GFP (1:1000; Life Technologies

#A11122), rabbit anti-RFP (1:1000; Life Technologies #R10367), rabbit anti-PTTH

antibody (1:1300), mouse anti-nc82 (1:40; DSHB), mouse anti-EcR-A (1:10;

AB_528214) and mouse anti-EcR-B1 (1:10; AB_2154902). Secondary antibodies used

were: Alexa Fluor goat anti-chicken 488 (1:500; Life Technologies #A11039), Alexa

Fluor goat anti-rabbit 488 (1:500; Life Technologies #A11034), Alexa Fluor goat anti-

rabbit 546 (1:500; Life Technologies #A11010), Alexa Fluor goat anti-mouse 647 (1:500;

Life Technologies #A21235), and Alexa Fluor goat anti-mouse 488 (1:500; Life

Technologies #A11029).

Confocal microscopy and image analysis

Confocal imaging was performed under an LSM 710 inverted confocal microscope

(ZEISS, Germany), with a Plan-Apochromat 20×/0.8 M27 objective or an EC Plan-

Neofluar 40x/1.30 oil DIC M27 objective, and later analyzed using Fiji software.

Generation of anti-PTTH antibody

The antisera used to recognize PTTH peptide were raised in New Zealand white rabbits

using the synthetic peptide N'- TSQSDHPYSWMNKDQPWQFKC -C'. The synthesis of

antigen peptide, the production and purification of antiserum were performed by Beijing

Genomics Institute (BGI).

Generation of UAS-PTTH

pJFRC28-5XUAS-IVS-GFP-p10 (#12073; Fungene Biotechnology, Shanghai, China)

was used for the generation of the pJFRC28-UAS-PTTH construct. The pJFRC28-

10XUAS-IVS-GFP-p10 plasmid was digested with NotI and XbaI to remove the GFP

coding sequence, and then the cDNA of PTTH was cloned into this plasmid by Gibson

Assembly. The Kozak sequence was added right upstream of the ATG. UAS-PTTH

constructs were injected and integrated into the attP2 site on the third chromosome

through phiC31 integrase mediated transgenesis. The construct was confirmed using

DNA sequencing through PCR. The primers used for cloning PTTH cDNA were as

follows:

UAS-PTTH-F

ATTCTTATCCTTTACTTCAGGCGGCCGCAAAATGGATATAAAGTATGGCGACTCC

UAS-PTTH-R

GTTATTTTAAAAACGATTCATTCTAGATCACTTTGTGCAGAAGCAGCCG

Genomic DNA extraction and RT-PCR

Genomic DNA was extracted from 10 whole bodies of wandering flies using MightyPrep

reagent for DNA (Takara #9182). Whole body RNA was extracted from 10 whole bodies

of wandering flies using TRIzol (Ambion #15596018). cDNA was generated from total

RNA using the Prime Script reagent kit (Takara #RR047A). Candidates of *ΔPtth* were

characterized by the loss of DNA band in the deleted areas through PCR on the

genomic DNA and cDNA. Primer sequences used in Figure 1 are listed in **Table S1**.

Measurements of pupariation timing and adult mass

The flies were reared at 25°C and 60% humidity under a 12 hr light:12 hr dark cycle.

Two-hour time collections of embryos laid on standard food vials. Each vial contained

20-30 eggs. The range of time for pupariation were recorded for each vial. Sexed adults

of 24h-old were weighted in groups of ten flies using a NENVER-TP-214 microbalance

at the same time.

Identification of sex in *Drosophila* larvae

Third instar larvae can be sexed (True et al., 2014). Gonads are translucent and visible

in side view in the posterior third of the larva. The male gonads are about five times

bigger than the female gonads. The identified wandering female and male larvae were

reared in different vials for the subsequent experiments.

Rescue by 20-hydroxyecdysone feeding

Thirty freshly ecdysed *ΔPtth* L3 larvae, grown at 25°C and 60% humidity under a 12 hr

light:12 hr dark cycle, were washed with water and transferred to normal food for

additional aging. After 20 h, larvae were washed and transferred to a vial supplemented

with either 20-hydroxyecdysone (20E, Cayman #16145, dissolved in 95% ethanol, final

concentration 0.2 mg/ml) or 95% ethanol (same volume as 20E). The wild type larvae

were directly transferred to vials supplemented with 20E or 95% ethanol upon L3

ecdysis. Once seeded with L3 larvae, the vials were returned to 25°C and 60% humidity

under a 12 hr light:12 hr dark cycle.

Quantification of fluorescence intensity

The fluorescence intensity was quantified using Fiji software. The areas of interest

(ROI) were marked in the slices including the interested regions and quantified using

the "plot z-axis profile" function. The fluorescence intensity in each slice was summed

for statistics and plot. The parameters used for confocal imaging of each brain were the

same.

Calcium imaging

Flies aged 4-6 days were immobilized on ice for ~30 s. The brain was then dissected

out in extra-cellular solution (ECS) that contains (in millimoles): 108 NaCl, 5 KC1, 5

trehalose, 5 sucrose, 5 HEPES, 26 $NaHCO_3$, 1 NaH_2PO_4, 2 $CaCl_2$, and 1.5 $MgCl_2$ [pH

7.1 – 7.3 when bubbled with 95% (vol/vol) O_2/5% (vol/vol) CO_2, ~290 mOsm] and

mounted on a poly-D-lysine coated coverslip. The samples were continuously perfused

with ECS.

Calcium imaging was performed at 21°C on a customized two-photon microscope

equipped with a resonant scanner (Nikon), a piezo objective scanner (Nikon) and a 40x

water-immersion objective (Nikon). GCaMP6s was excited at 920 nm.

Analysis of calcium imaging data was done offline with NIS-Elements AR 5.30.01.

Briefly, the square region of interest (ROIs), 25 pixels on the pC1 neurons in the center

of lateral junction, was chosen for measurements. For each frame, the average

fluorescence intensity of pixels within ROIs was calculated blind to genotype. The

average fluorescence intensity of ROIs in each frame covering pC1 neurons were

summed for statistics and plot.

RNA isolation and sequencing

About 100 female fly brains were dissected from single individuals and the total RNA

was extracted using TRIzol (Ambion #15596018). Sequencing was performed on an

Illumina Novoseq6000 PE150 (Illumina, CA). Three biological replicates were performed

for each genetic type. The library preparation and sequencing were performed by

Novogene in China.

RNA-seq analyses

RNA-seq reads were mapped to the reference dmel-all-r6.42 (flaybase) using STAR

(v2.7.6a) (Dobin et al., 2013) with default parameters. Aligned reads were assigned to

dmel-all-r6.43.gtf gene annotation model, mapping reads were counted by

featureCounts from subread package with default parameters (Liao et al., 2013), and

then the transcript per millions (TPMs) was calculated. Differential expression was

analyzed with the DESeq2 package (Love et al., 2014) and visualized by volcano plots

using the maximum a posteriori estimated log-fold change using the $\log_2$(fold-change)

calculated from TPMs.

qRT-PCR

Whole brain RNA was extracted from about 100 fly brains using TRIzol (Ambion

#15596018). The cDNA was synthesized using Prime Script reagent kit (Takara

#RR047A). Quantitative PCR was performed on Thermo Piko Real 96 (Thermo) using

SYBR Green PCR Master Mix (Takara #RR820A). The mRNA expression level was

calculated by the $2^{-\Delta\Delta Ct}$ method and the results were plotted by using tubulin as the

reference gene. Primers are listed in **Table S1**. All reactions were performed in

triplicate. The average of four biological replicates ± SEM was plotted.

Statistical analysis

Statistical analyses were carried out using R software version 3.4.3 or Prism7

(GraphPad software). For the copulation rate, chi-square test is applied. The Mann-

Whitney U test was applied for analyzing the significance of two columns. Kruskal-

Wallis ANOVA test followed by post-hoc Mann-Whitney U test, was used to identify

significant differences between multiple groups.

Data, Materials, and Software Availability

All study data are included in the main text and supporting information, except for the

sequence data of RNAseq (**Supplementary File 3**) has been deposited in the Genome

Sequence Archive under the accession number CRA012130 in PRJCA018750. This

study does not involve new code. Fly stocks and reagents used in this study are

available from the corresponding author upon reasonable request.

Acknowledgments

We thank Yi Rao (Peking University), Yufeng Pan (Southeast University), Kaiyu Wang

(Lingang Laboratory), Yan Zhu (Chinese Academy of Sciences), Paul Garrity (Brandeis

University), Wei Zhang (Tsinghua University), Li Liu (Chinese Academy of Sciences)

and Xuan Guo (Jinzhou Medical University), the Bloomington Drosophila Stock Center

and Tsinghua Fly Center for sharing fly strains; Chenzhu Wang (Chinese Academy of

Sciences), Yufeng Pan (Southeast University) and Zhiqiang Yan (Shenzhen Bay

Laboratory) for their comments; Fengming Wu (Chinese Academy of Sciences), Tao

Wang (Chinese Academy of Sciences) and Jin Ge (Chinese Academy of Sciences) for

assistance with behavior assays; Yihui Chen for the maintenance of fly stocks; other

members of the Zhou laboratory for helpful discussions.

References

Avila F W, Sirot L K, LaFlamme B A, Rubinstein C D, & Wolfner M F. (2011). Insect seminal fluid proteins: identification and function. *Annu Rev Entomol, 56*, 21-40. doi:10.1146/annurev-ento-120709-144823

Billeter J C, Villella A, Allendorfer J B, Dornan A J, Richardson M, Gailey D A, & Goodwin S F. (2006). Isoform-specific control of male neuronal differentiation and behavior in Drosophila by the fruitless gene. *Curr Biol, 16*(11), 1063-1076. doi:10.1016/j.cub.2006.04.039

Clowney E J, Iguchi S, Bussell J J, Scheer E, & Ruta V. (2015). Multimodal chemosensory circuits controlling male courtship in Drosophila. *Neuron, 87*(5), 1036-1049. doi:10.1016/j.neuron.2015.07.025

Connolly K, & Cook R. (1973). Rejection responses by female Drosophila-melanogaster - their ontogeny, causality and effects upon behavior of courting male. *Behaviour, 44*(1-2), 142-166. doi:10.1163/156853973x00364

Cubells J F, & Zabetian C P. (2004). Human genetics of plasma dopamine beta-hydroxylase activity: applications to research in psychiatry and neurology. *Psychopharmacology (Berl), 174*(4), 463-476. doi:10.1007/s00213-004-1840-8

Dai J D, & Gilbert L I. (1991). Metamorphosis of the corpus allatum and degeneration of the prothoracic glands during the larval-pupal-adult transformation of Drosophila melanogaster: a cytophysiological analysis of the ring gland. *Dev Biol, 144*(2), 309-326. doi:10.1016/0012-1606(91)90424-2

Dalton J E, Lebo M S, Sanders L E, Sun F, & Arbeitman M N. (2009). Ecdysone receptor acts in fruitless- expressing neurons to mediate drosophila courtship behaviors. *Curr Biol, 19*(17), 1447-1452. doi:10.1016/j.cub.2009.06.063

Demir E, & Dickson B J. (2005). fruitless splicing specifies male courtship behavior in Drosophila. *Cell, 121*(5), 785-794. doi:10.1016/j.cell.2005.04.027

Deng B, Li Q, Liu X, Cao Y, Li B, Qian Y, Xu R, Mao R, Zhou E, Zhang W, Huang J, & Rao Y. (2019). Chemoconnectomics: mapping chemical transmission in Drosophila. *Neuron, 101*(5), 876-893.e874. doi:10.1016/j.neuron.2019.01.045

Deutsch D, Pacheco D, Encarnacion-Rivera L, Pereira T, Fathy R, Clemens J, Girardin C, Calhoun A, Ireland E, Burke A, Dorkenwald S, McKellar C, Macrina T, Lu R, Lee K, Kemnitz N, Ih D, Castro M, Halageri A, Jordan C, Silversmith W, Wu J, Seung H S, & Murthy M. (2020). The neural basis for a persistent internal state in Drosophila females. *Elife, 9*. doi:10.7554/eLife.59502

Dickson B J. (2008). Wired for sex: the neurobiology of Drosophila mating decisions. *Science, 322*(5903), 904-909. doi:10.1126/science.1159276

Dobin A, Davis C A, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, & Gingeras T R. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics, 29*(1), 15-21. doi:10.1093/bioinformatics/bts635

Feng K, Palfreyman M T, Hasemeyer M, Talsma A, & Dickson B J. (2014). Ascending SAG neurons control sexual receptivity of Drosophila females. *Neuron, 83*(1), 135-148. doi:10.1016/j.neuron.2014.05.017

Ferveur J F. (2010). Drosophila female courtship and mating behaviors: sensory signals, genes,
neural structures and evolution. *Curr Opin Neurobiol, 20*(6), 764-769.
doi:10.1016/j.conb.2010.09.007
Fuyama Y, & Ueyama M. (1997). Ovulation and the suppression of mating in Drosophila
melanogaster females: behavioral basis. *Behav Genet, 27*(5), 483-488.
doi:10.1023/a:1025630602057
Ganter G K, Walton K L, Merriman J O, Salmon M V, Brooks K M, Maddula S, & Kravitz E A.
(2007). Increased male-male courtship in ecdysone receptor deficient adult flies. *Behav
Genet, 37*(3), 507-512. doi:10.1007/s10519-006-9140-1
Greenspan R J, & Ferveur J F. (2000). Courtship in Drosophila. *Annu Rev Genet, 34*, 205-232.
doi:10.1146/annurev.genet.34.1.205
Hall J. (1978). Courtship among males due to a male-sterile mutation in Drosophila
melanogaster. *Behavior Genetics,* 8(2), 125-41. doi:10.1007/BF01066870
Hall J C. (1994). The mating of a fly. *Science, 264*(5166), 1702-1714.
doi:10.1126/science.8209251
Hamada F N, Rosenzweig M, Kang K, Pulver S R, Ghezzi A, Jegla T J, & Garrity P A. (2008). An
internal thermal sensor controlling temperature preference in Drosophila. *Nature,
454*(7201), 217-220. doi:10.1038/nature07001
Häsemeyer M, Yapici N, Heberlein U, & Dickson B J. (2009). Sensory neurons in the Drosophila
genital tract regulate female reproductive behavior. *Neuron, 61*(4), 511-518.
doi:10.1016/j.neuron.2009.01.009
Herbison A E. (2016). Control of puberty onset and fertility by gonadotropin-releasing hormone
neurons. *Nat Rev Endocrinol, 12*(8), 452-466. doi:10.1038/nrendo.2016.70
Hoopfer E D, Jung Y, Inagaki H K, Rubin G M, & Anderson D J. (2015). P1 interneurons promote a
persistent internal state that enhances inter-male aggression in Drosophila. *Elife, 4*.
doi:10.7554/eLife.11346
Imura E, Shimada-Niwa Y, Nishimura T, Huckesfeld S, Schlegel P, Ohhara Y, Kondo S, Tanimoto
H, Cardona A, Pankratz M J, & Niwa R. (2020). The Corazonin-PTTH neuronal axis
controls systemic body growth by regulating basal ecdysteroid biosynthesis in
Drosophila melanogaster. *Curr Biol, 30*(11), 2156-2165 e2155.
doi:10.1016/j.cub.2020.03.050
Ishimoto H, & Kamikouchi A. (2020). A feedforward circuit regulates action selection of pre-
mating courtship behavior in female Drosophila. *Curr Biol, 30*(3), 396-407.e394.
doi:10.1016/j.cub.2019.11.065
Ito H, Sato K, Koganezawa M, Ote M, Matsumoto K, Hama C, & Yamamoto D. (2012). Fruitless
recruits two antagonistic chromatin factors to establish single-neuron sexual
dimorphism. *Cell, 149*(6), 1327-1338. doi:10.1016/j.cell.2012.04.025
Jang Y H, Chae H S, & Kim Y J. (2017). Female-specific myoinhibitory peptide neurons regulate
mating receptivity in Drosophila melanogaster. *Nat Commun, 8*(1), 1630.
doi:10.1038/s41467-017-01794-9
Kim C H, Zabetian C P, Cubells J F, Cho S, Biaggioni I, Cohen B M, Robertson D, & Kim K S. (2002).
Mutations in the dopamine beta-hydroxylase gene are associated with human
norepinephrine deficiency. *Am J Med Genet, 108*(2), 140-147.

Kimura K, Hachiya T, Koganezawa M, Tazawa T, & Yamamoto D. (2008). Fruitless and doublesex
832   coordinate to generate male-specific neurons that can initiate courtship. *Neuron, 59*(5),
833   759-769. doi:10.1016/j.neuron.2008.06.007

834   Kimura K, Sato C, Koganezawa M, & Yamamoto D. (2015). Drosophila ovipositor extension in
835   mating behavior and egg deposition involves distinct sets of brain interneurons. *PLoS*
836   *One, 10*(5), e0126445. doi:10.1371/journal.pone.0126445

837   Kohatsu S, Koganezawa M, & Yamamoto D. (2011). Female contact activates male-specific
838   interneurons that trigger stereotypic courtship behavior in Drosophila. *Neuron, 69*(3),
839   498-508. doi:10.1016/j.neuron.2010.12.017

840   Kubli E. (2003). Sex-peptides: seminal peptides of the Drosophila male. *Cell Mol Life Sci, 60*(8),
841   1689-1704. doi:10.1007/s00018-003-3052

842   Kurtovic A, Widmer A, & Dickson B J. (2007). A single class of olfactory neurons mediates
843   behavioural responses to a Drosophila sex pheromone. *Nature, 446*(7135), 542-546.
844   doi:10.1038/nature05672

845   Kvitsiani D, & Dickson B J. (2006). Shared neural circuitry for female and male sexual behaviours
846   in Drosophila. *Curr Biol, 16*(10), R355-356. doi:10.1016/j.cub.2006.04.025

847   Lavrynenko O, Rodenfels J, Carvalho M, Dye N A, Lafont R, Eaton S, & Shevchenko A. (2015).
848   The ecdysteroidome of Drosophila: influence of diet and development. *Development,*
849   *142*(21), 3758-3768. doi:10.1242/dev.124982

850   Liao Y, Smyth G K, & Shi W. (2013). The Subread aligner: fast, accurate and scalable read
851   mapping by seed-and-vote. *Nucleic Acids Res, 41*(10), e108. doi:10.1093/nar/gkt214

852   Love M I, Huber W, & Anders S. (2014). Moderated estimation of fold change and dispersion for
853   RNA-seq data with DESeq2. *Genome Biol, 15*(12), 550. doi:10.1186/s13059-014-0550-8

854   Ma B, Wang R, Liu Y, Deng B, Wang T, Wu F, & Zhou C. (2022). Serotonin signaling modulates
855   sexual receptivity of virgin female Drosophila. *Neurosci Bull, 38*(11), 1277-1291.
856   doi:10.1007/s12264-022-00908-8

857   Manoli D S, Fan P, Fraser E J, & Shah N M. (2013). Neural control of sexually dimorphic
858   behaviors. *Curr Opin Neurobiol, 23*(3), 330-338. doi:10.1016/j.conb.2013.04.005

859   Manoli D S, Foss M, Villella A, Taylor B J, Hall J C, & Baker B S. (2005). Male-specific fruitless
860   specifies the neural substrates of Drosophila courtship behaviour. *Nature, 436*(7049),
861   395-400. doi:10.1038/nature03859

862   Manoli D S, Meissner G W, & Baker B S. (2006). Blueprints for behavior: genetic specification of
863   neural circuitry for innate behaviors. *Trends Neurosci, 29*(8), 444-451.
864   doi:10.1016/j.tins.2006.06.006

865   McBrayer Z, Ono H, Shimell M, Parvy J P, Beckstead R B, Warren J T, Thummel C S, Dauphin-
866   Villemant C, Gilbert L I, & O'Connor M B. (2007a). Prothoracicotropic hormone regulates
867   developmental timing and body size in Drosophila. *Dev Cell, 13*(6), 857-871.
868   doi:10.1016/j.devcel.2007.11.003

869   Mcbrayer Z, Ono H, Shimell M J, Parvy J P, Beckstead R B, Warren J T, Thummel C S, Dauphin-
870   Villemant C, Gilbert L I, & O'Connor M B. (2007b). Prothoracicotropic hormone regulates
871   developmental timing and body size in Drosophila. *Developmental Cell, 13*(6), 857-871.
872   doi:10.1016/j.devcel.2007.11.003

McGuire S E, Mao Z, & Davis R L. (2004). Spatiotemporal gene expression targeting with the TARGET and gene-switch systems in Drosophila. *Sci STKE, 2004*(220), pl6. doi:10.1126/stke.2202004pl6

Mellert D J, Knapp J M, Manoli D S, Meissner G W, & Baker B S. (2010). Midline crossing by gustatory receptor neuron axons is regulated by fruitless, doublesex and the Roundabout receptors. *Development, 137*(2), 323-332. doi:10.1242/dev.045047

Mellert D J, Robinett C C, & Baker B S. (2012). doublesex functions early and late in gustatory sense organ development. *PLoS One, 7*(12), e51489. doi:10.1371/journal.pone.0051489

Mezzera C, Brotas M, Gaspar M, Pavlou H J, Goodwin S F, & Vasconcelos M L. (2020). Ovipositor extrusion promotes the transition from courtship to copulation and signals female acceptance in Drosophila melanogaster. *Curr Biol, 30*(19), 3736-3748.e3735. doi:10.1016/j.cub.2020.06.071

Neckameyer W S. (1998). Dopamine modulates female sexual receptivity in Drosophila melanogaster. *J Neurogenet, 12*(2), 101-114. doi:10.3109/01677069809167259

Yamanaka, N, Romero, N M, Martin, F A, Rewitz, K F, Sun, M, O'Connor, & M B, Léopold, P. (2013). Neuroendocrine control of Drosophila larval light preference. *Science, 341*(6150), 1113-1116. doi: 10.1126/science.1241210

Pan X, & O'Connor M B. (2019). Developmental maturation: Drosophila asta signaling provides a kiss to grow up. *Curr Biol, 29*(5), R161-R164. doi:10.1016/j.cub.2019.01.040

Pan Y, & Baker B S. (2014). Genetic identification and separation of innate and experience-dependent courtship behaviors in Drosophila. *Cell, 156*(1-2), 236-248. doi:10.1016/j.cell.2013.11.041

Pan Y, Meissner G W, & Baker B S. (2012). Joint control of Drosophila male courtship behavior by motion cues and activation of male-specific P1 neurons. *PNAS, 109*(25), 10065-10070. doi:10.1073/pnas.1207107109

Pan Y, Robinett C C, & Baker B S. (2011). Turning males on: activation of male courtship behavior in Drosophila melanogaster. *PLoS One, 6*(6), e21144. doi:10.1371/journal.pone.0021144

Park D H, Kashimoto T, Ebstein R P, & Goldstein M. (1976). Purification and immunochemical characterization of dopamine beta-hydroxylase from human pheochromocytoma. *Mol Pharmacol, 12*(1), 73-81.

Pavlou H J, & Goodwin S F. (2013). Courtship behavior in Drosophila melanogaster: towards a 'courtship connectome'. *Curr Opin Neurobiol, 23*(1), 76-83. doi:10.1016/j.conb.2012.09.002

Pfeiffer B D, Jenett A, Hammonds A S, Ngo T T, Misra S, Murphy C, Scully A, Carlson J W, Wan K H, Laverty T R, Mungall C, Svirskas R, Kadonaga J T, Doe C Q, Eisen M B, Celniker S E, & Rubin G M. (2008). Tools for neuroanatomy and neurogenetics in Drosophila. *PNAS, 105*(28), 9715-9720. doi:10.1073/pnas.0803697105

Pfeiffer B D, Ngo T T, Hibbard K L, Murphy C, Jenett A, Truman J W, & Rubin G M. (2010). Refinement of tools for targeted gene expression in Drosophila. *Genetics, 186*(2), 735-755. doi:10.1534/genetics.110.119917

Prigge S T, Mains R E, Eipper B A, & Amzel L M. (2000). New insights into copper monooxygenases and peptide amidation: structure, mechanism and function. *Cell Mol Life Sci, 57*(8-9), 1236-1259. doi:10.1007/pl00000763

917   Rewitz K F, Yamanaka N, Gilbert L I, & O'Connor M B. (2009). The insect neuropeptide PTTH
918   activates receptor tyrosine kinase torso to initiate metamorphosis. *Science, 326*(5958),
919   1403-1405. doi:10.1126/science.1176450
920   Rezával C, Nojima T, Neville M C, Lin A C, & Goodwin S F. (2014). Sexually dimorphic
921   octopaminergic neurons modulate female postmating behaviors in Drosophila. *Curr Biol,*
922   *24*(7), 725-730. doi:10.1016/j.cub.2013.12.051
923   Rezával C, Pattnaik S, Pavlou H J, Nojima T, Brüggemeier B, D'Souza L A D, Dweck H K M, &
924   Goodwin S F. (2016). Activation of latent courtship circuitry in the brain of Drosophila
925   females induces male-like behaviors. *Curr Biol, 26*(18), 2508-2515.
926   doi:10.1016/j.cub.2016.07.021
927   Riddiford L M, Cherbas P, & Truman J W. (2000). Ecdysone receptors and their biological
928   actions. *Vitam Horm, 60*, 1-73. doi:10.1016/s0083-6729(00)60016-x
929   Rideout E J, Dornan A J, Neville M C, Eadie S, & Goodwin S F. (2010). Control of sexual
930   differentiation and behavior by the doublesex gene in Drosophila melanogaster. *Nat*
931   *Neurosci, 13*(4), 458-466. doi:10.1038/nn.2515
932   Roy S, Saha T T, Zou Z, & Raikhel A S. (2018). Regulatory pathways controlling female insect
933   reproduction. *Annu Rev Entomol, 63*, 489-511. doi:10.1146/annurev-ento-020117-
934   043258
935   Ruta V, Datta S R, Vasconcelos M L, Freeland J, Looger L L, & Axel R. (2010). A dimorphic
936   pheromone circuit in Drosophila from sensory input to descending output. *Nature,*
937   *468*(7324), 686-690. doi:10.1038/nature09554
938   Scheffer L K, Xu C S, Januszewski M, Lu Z, Takemura S Y, Hayworth K J, Huang G B, Shinomiya K
939   et al. & Plaza S M. (2020). A connectome and analysis of the adult Drosophila central
940   brain. *Elife, 9*. doi:10.7554/eLife.57443
941   Schretter C E, Aso Y, Robie A A, Dreher M, Dolan M J, Chen N, Ito M, Yang T, Parekh R, Branson
942   K M, & Rubin G M. (2020). Cell types and neuronal circuitry underlying female
943   aggression in Drosophila. *Elife, 9*. doi:10.7554/eLife.58942
944   Sengupta S, Chan Y B, Palavicino-Maggio C B, & Kravitz E A. (2022). GABA transmission from
945   mAL interneurons regulates aggression in Drosophila males. *PNAS, 119*(5).
946   doi:10.1073/pnas.2117101119
947   Shimell M, Pan X, Martin F A, Ghosh A C, Leopold P, O'Connor M B, & Romero N M. (2018).
948   Prothoracicotropic hormone modulates environmental adaptive plasticity through the
949   control of developmental timing. *Development, 145*(6). doi:10.1242/dev.159699
950   Siwicki K K, & Kravitz E A. (2009). Fruitless, doublesex and the genetics of social behavior in
951   Drosophila melanogaster. *Curr Opin Neurobiol, 19*(2), 200-206.
952   doi:10.1016/j.conb.2009.04.001
953   Stockinger P, Kvitsiani D, Rotkopf S, Tirián L, & Dickson B J. (2005). Neural circuitry that governs
954   Drosophila male courtship behavior. *Cell, 121*(5), 795-807.
955   doi:10.1016/j.cell.2005.04.026
956   Taisz I, Donà E, Münch D, Bailey S N, Morris B J, Meechan K I, Stevens K M, Varela-Martínez I,
957   Gkantia M, Schlegel P, Ribeiro C, Jefferis G, & Galili D S. (2023). Generating parallel
958   representations of position and identity in the olfactory system. *Cell, 186*(12), 2556-
959   2573.e2522. doi:10.1016/j.cell.2023.04.038

960   Terhzaz S, Rosay P, Goodwin S F, & Veenstra J A. (2007). The neuropeptide SIFamide modulates
961   sexual behavior in Drosophila. *Biochem Biophys Res Commun, 352*(2), 305-310.
962   doi:10.1016/j.bbrc.2006.11.030

963   Timmers H J, Deinum J, Wevers R A, & Lenders J W. (2004). Congenital dopamine-beta-
964   hydroxylase deficiency in humans. *Ann N Y Acad Sci, 1018*, 520-523.
965   doi:10.1196/annals.1296.064

966   Sylwester Chyb and Nicolas Gompel. (2014). Atlas of Drosophila morphology: wild-type and
967   classical mutants. "Life stages: third instar larva and pupa". Elsevier, pp. 18-19.

968   Truman J W, & Riddiford L M. (2023). Drosophila postembryonic nervous system development:
969   a model for the endocrine control of development. *Genetics, 223*(3).
970   doi:10.1093/genetics/iyac184

971   Truman J W, Talbot W S, Fahrbach S E, & Hogness D S. (1994). Ecdysone receptor expression in
972   the CNS correlates with stage-specific responses to ecdysteroids during Drosophila and
973   Manduca development. *Development, 120*(1), 219-234. doi:10.1242/dev.120.1.219

974   Tsuneoka Y, Tsukahara S, Yoshida S, Takase K, Oda S, Kuroda M, & Funato H. (2017). Moxd1 is a
975   marker for sexual dimorphism in the medial preoptic area, Bed Nucleus of the Stria
976   Terminalis and Medial Amygdala. *Front Neuroanat, 11*, 26.
977   doi:10.3389/fnana.2017.00026

978   Vendelboe T V, Harris P, Zhao Y, Walter T S, Harlos K, El Omari K, & Christensen H E. (2016). The
979   crystal structure of human dopamine β-hydroxylase at 2.9 Å resolution. *Sci Adv, 2*(4),
980   e1500980. doi:10.1126/sciadv.1500980

981   von Philipsborn A C, Liu T, Yu J Y, Masser C, Bidaye S S, & Dickson B J. (2011). Neuronal control
982   of Drosophila courtship song. *Neuron, 69*(3), 509-522.
983   doi:10.1016/j.neuron.2011.01.011

984   Wang F, Wang K, Forknall N, Parekh R, & Dickson B J. (2020a). Circuit and behavioral
985   mechanisms of sexual rejection by Drosophila females. *Curr Biol, 30*(19), 3749-3760
986   e3743. doi:10.1016/j.cub.2020.07.083

987   Wang F, Wang K, Forknall N, Patrick C, Yang T, Parekh R, Bock D, & Dickson B J. (2020b). Neural
988   circuitry linking mating and egg laying in Drosophila females. *Nature, 579*(7797), 101-
989   105. doi:10.1038/s41586-020-2055-9

990   Wang K, Wang F, Forknall N, Yang T, Patrick C, Parekh R, & Dickson B J. (2021). Neural circuit
991   mechanisms of sexual receptivity in Drosophila females. *Nature, 589*(7843), 577-581.
992   doi:10.1038/s41586-020-2972-7

993   Wang L, & Anderson D J. (2010). Identification of an aggression-promoting pheromone and its
994   receptor neurons in Drosophila. *Nature, 463*(7278), 227-231. doi:10.1038/nature08678

995   Wang T, Jing B, Deng B, Shi K, Li J, Ma B, Wu F, & Zhou C. (2022). Drosulfakinin signaling
996   modulates female sexual receptivity in Drosophila. *Elife, 11*. doi:10.7554/eLife.76025

997   Wu Y, Bidaye S, & Mahringer D. (2019). Drosophila female-specific brain neuron elicits
998   persistent position- and direction-selective male-like social behaviors.
999   doi:10.1101/594960

1000   Xin X, Mains R E, & Eipper B A. (2004). Monooxygenase X, a member of the copper-dependent
1001   monooxygenase family localized to the endoplasmic reticulum. *J Biol Chem, 279*(46),
1002   48159-48167. doi:10.1074/jbc.M407486200

Yamamoto D. (2007). The neural and genetic substrates of sexual behavior in Drosophila. *Adv Genet, 59*, 39-66. doi:10.1016/S0065-2660(07)59002-4

Yamamoto D, & Koganezawa M. (2013). Genes and circuits of courtship behaviour in Drosophila males. *Nat Rev Neurosci, 14*(10), 681-692. doi:10.1038/nrn3567

Yamanaka N, Romero N M, Martin F A, Rewitz K F, Sun M, O'Connor M B, & Léopold P. (2013). Neuroendocrine control of Drosophila larval light preference. *Science, 341*(6150), 1113-1116. doi:10.1126/science.1241210

Yang C H, Rumpf S, Xiang Y, Gordon M D, Song W, Jan L Y, & Jan Y N. (2009). Control of the postmating behavioral switch in Drosophila females by internal sensory neurons. *Neuron, 61*(4), 519-526. doi:10.1016/j.neuron.2008.12.021

Zhou C, Pan Y, Robinett C C, Meissner G W, & Baker B S. (2014). Central brain neurons expressing doublesex regulate female receptivity in Drosophila. *Neuron, 83*(1), 149-163. doi:10.1016/j.neuron.2014.05.038

Figure Legends

Figure 1 *Ptth* null mutants have increased virgin female receptivity.

(A-C) Generation and validation of a 974 bp deletion mutant of the *Ptth* gene. (D-G)

Virgin female receptivity of *Ptth* null mutants on the 1st (D), 2nd (E), 3rd (F) and 6th day

(G) respectively. (H) Brain of indicated genotype, immunostained with anti-PTTH

antibody (green) and counterstained with nc82 (magenta). Arrows show signals (green)

stained with anti-PTTH antibody. Scale bars, 50 µm. (I) Enhanced virgin female

receptivity of *ΔPtth* null mutants was rescued by elavGAL4 driving UAS-PTTH. The

increased copulation rate and decreased latency to copulation on the 1st day after

eclosion were rescued to the comparable level of control. The number of female flies

paired with wild type males is displayed in parentheses. For the copulation rate, chi-

square test is applied. For the latency to copulation, Kruskal-Wallis ANOVA and post

hoc Mann-Whitney U tests are applied. Error bars indicate SEM. $*p < 0.05$, $**p < 0.01$,

$***p < 0.001$, $****p < 0.0001$, ns indicates no significant difference.

1048

1049

1050

1051

1052

1053

1054

1055

1056

1057

1058

1059

Figure 2 Inactivation of doublesex-positive PG neurons expressing PTTH enhances virgin female receptivity during the larval stage.

(A-B) Expression pattern of PtthGAL4 revealed by anti-GFP in larvae central nervous system (CNS) (A) and adult CNS (B). Representative of five female flies. Scale bars, 50 μm. (C) All PG neurons were colabeled by dsxGAL4 driving UAS-GFPStinger (red) and PtthLexA driving LexAop-tomato (green). Representative of five female brains. Scale bars, 50 μm and 5 μm (zoom-in). (D) All PG neurons were *Ptth* and *Dsx* co-expressing, labeled by intersectional strategy. Representative of 5 female brains. Scale bars, 50 μm. (E-G) PG neurons were inactivated during larval (E), pupal (F) and adult (G) stages respectively by kir2.1, restricted by shifts from 18°C to 30°C. The inactivation during the larval stage significantly increased the copulation rate (E). The number of female flies paired with wild-type males is displayed in parentheses. For the copulation rate, chi-square test is applied. Error bars indicate SEM. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, ns indicates no significant difference.

Figure 3 Activation of PG neurons expressing PTTH during the 3rd-instar larvae

inhibits virgin female receptivity.

(A) Four developmental stages of *Drosophila* before eclosion when PG neurons were

thermogenetic activated by dTrpA1. L1, L2, and L3: start of three larval stages, W: start

of wandering stage, Pp: puparium formation, P1 and P2: start of the 1st and 2nd day of

pupal stage. (B-E) PtthGAL4 driving UAS-dTrpA1 activated PG neurons at 29^{o}C.

Activation of PG neurons at the stage 2 significantly decreased copulation rate (C), but

not at the stage 1 (B), stage 3 (D) and stage 4 (E). (F) Activation of PG neurons at the

stage 2 significantly increased the latency to copulation. (G) Mean velocity had no

significant change when PG neurons were activated during the stage 2 compared with

control females (ns = not significant, Kruskal-Wallis ANOVA and post hoc Mann-

Whitney U tests, mean ± SEM, n = 8-12)*. The number of female flies paired with wild-

type males is displayed in parentheses. For the copulation rate, chi-square test is

applied. For the latency to copulation, Mann-Whitney U test is applied. Error bars

indicate SEM. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, ns indicates no

significant difference.

1094

1095

1096

1097

1098

1099

1100

1101

1102

1103

1104

1105

1106

1107

1108

Figure 4 Feeding 20E restores virgin female receptivity of *Ptth* null mutant flies.

(A-B) The increased copulation rate and decreased latency to copulation of the 24h-old
ΔPtth flies were rescued to the comparable level of wild type females by feeding 20E to
the 3rd-instar larval *ΔPtth* flies. The wild type larval females fed by 20E had no
significantly different copulation rate and latency to copulation compared with the wild
type females fed by the same volume of 95% ethanol which is the solvent of 20E. The
number of female flies paired with wild-type males is displayed in parentheses. For the
copulation rate, chi-square test is applied. For the latency to copulation, Kruskal-Wallis
ANOVA and post hoc Mann-Whitney U tests are applied. Error bars indicate SEM. *p <
0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, ns indicates no significant difference.

1132

Figure 5 Virgin females with reduced EcR-A in pC1 neurons have reduced sexual

receptivity.

(A) Knock-down of EcR-A in pC1 neurons driven by pC1-ss2-GAL4 significantly

decreased the copulation rate and increased the latency to copulation. (B) Knock-down

of EcR-B1 in pC1 neurons driven by pC1-ss2-GAL4 significantly prolonged the latency

to copulation. (C-D) Knock-down of EcR-A (C) or EcR-B1 (D) in pC1 neurons driven by

pC1-ss1-GAL4 did not affect the copulation rate or the latency to copulation. (E)

Courtship index of wild-type males towards a female with the indicated genotype (n =

8). (F) The number of eggs laid by virgin females during the 3^{rd} - 4^{th} day after eclosion

when EcR-A was knocked down in pC1 neurons (n = 17-36). (G) Knock-down of EcR-A

in pC1 neurons decreased the opening of vaginal plate of virgin females compared with

controls (n = 8). (H) Knock-down of EcR-A in pC1 neurons increased the ovipositor

extrusion of virgin females compared with controls (n = 8). (I-K) Virgin female copulation

rate when EcR-A was knocked down in pC1 neurons temporally restricted by shifts from

18°C to 30°C. EcR-A was knocked down during larval (I), pupal (J) and adult (K) stages

respectively. Knock-down of EcR-A during pupal stage significantly decreased the

copulation rate (J). The number of female flies paired with wild-type males is displayed

in parentheses. For the copulation rate, chi-square test is applied. For other

comparisons, Kruskal-Wallis ANOVA and post hoc Mann-Whitney U tests are applied.

Error bars indicate SEM. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$, ns indicates

no significant difference.

1154

1155

1156

1157

1158

1159

1160

1161

1162

1163

1164

1165

1166

1167

1168

1169

1170

Figure 6 Reduced EcR-A in pC1 neurons induces the morphological changes.

(A1-A2) pC1 neurons appeared at the start of the pupal stage. (B-D) Reduced EcR-A in pC1 neurons induced more elaborated morphologies of pC1d axons, especially the extra vertical projection (EVP). The EVP regions of pC1d neurons was indicated by arrows. The morphological changes appeared on the 2nd day of the pupal stage (B1-B2) and retained to the adult stage including the 1st day (C1-C2) and the 4th day (D1-D2) of the adult stage. p0, the 1st day of the pupal stage; p2, the 2nd day of the pupal stage; A1, the 1st day of the adult stage; A4, the 4th day of the adult stage. (E-F) Fluorescence intensity of EVP in pC1d neurons on the 2nd day of the pupal stage (E) and the 4th day of the adult stage (F) was quantified when EcR-A was reduced in pC1 neurons (n=7). The quantified EVP regions were marked in (B) and (D) with orange ellipses. (G) pC1 neurons of the 4th day adults had comparable cell body number when EcR-A was reduced in pC1 neurons or not (n = 7). (H) Basal GCaMP6s signals in the LPC region of pC1 neurons when EcR-A was reduced in pC1 neurons (n = 22). LPC regions, the neurites extending from pC1 cell bodies, were marked with orange square in (D1) and (D2). Scale bars are 50 µm. For all comparisons, Mann-Whitney U test is applied. Error bars indicate SEM. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, ns indicates no significant difference.

1189

1190

Figure 7 Reduced EcR-A in pC1 neurons induces the down-regulated dopamine

beta-monooxygenase (DBM) level.

(A) Volcano plot of RNA-seq from virgin female brains in which EcR-A was knocked

down in pC1 neurons or not. Each circle represents a protein-coding gene. Differential

genes with a p-value < 0.01 are highlighted in blue. Differential genes with a fold

change > 4 are highlighted in green. Differential genes with a p-value < 0.01 and fold

change > 4 are highlighted in red. DBM, the most down-regulated genes with

annotation, are indicated by black arrow. Data are from three replicates, each contains

about 100 brains. (B) qRT-PCR for DBM when EcR-A was knocked down in pC1

neurons. Bars represent mean ± SEM. p values are from Mann-Whitney U test (n = 12

based on four replicates for each, each contains about 100 brains). *p < 0.05, **p <

0.01, ***p < 0.001, ****p < 0.0001, ns indicates no significant difference.

1203

1204

1205

1206

1207

1208

1209

1210

Figure 1—figure supplement 1 Weight, attractiveness and locomotion behavior of
***Ptth* null mutant virgin females.**

(A) The weight of 24h-old adult *ΔPtth* null mutant females was significantly higher than
that of wild type females (Mann-Whitney U test, n=8 groups, 10 flies in each group). (B)
Courtship index of wild-type males during the first 5 min of courtship towards a female
with the indicated genotype (Kruskal-Wallis ANOVA and post hoc Mann-Whitney U
tests, n = 7-9). (C-D) Mean velocity had no significant change in *ΔPtth* null mutant
females on the 1st day (C) and the 6th day (D) compared with control females (Kruskal-
Wallis ANOVA and post hoc Mann-Whitney U tests, n = 7-12)*.* Error bars indicate SEM.
*p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, ns = not significant.

A   B

PtthLexA>LexAop-GFP

PtthLexA>LexAop-GFP

1230

1231 **Figure 2—figure supplement 1 PG neurons expressing PTTH labeled by PtthLexA.**

1232 Expression pattern of PtthLexA in the brain revealed by anti-GFP (green) in larvae brain

1233 (A) and adult brain (B). Representative of five female flies. Scale bars, 50 µm.

1234

1235

1236

1237

1238

1239

1240

1241

1242

1243

1244

1245

1246

1247

1248

1249

1250

1251

1252

1253

1254

1255

1256

Figure 2—figure supplement 2 The anatomical pattern of PG neurons expressing PTTH at different developmental stages.

Expression pattern of PtthGAL4 in the brain revealed by anti-GFP from the 3[rd] larval stage to the 4[th] day after eclosion. Arrows show PTTH signals (green) stained with anti-GFP antibody. L3, the 3[rd]-instar larvae; wander, the wandering larvae; P4, the 4[th] day of the pupal stage; A0h, the 1[st] hour of the adult stage; A3h, the 3[rd] hour of the adult stage; A6h, the 6[th] hour of the adult stage; A9h, the 9[th] hour of the adult stage; A12h, the 12[th] hour of the adult stage; A4d, the 4[th] day of the adult stage. Representative of five female flies. Scale bars, 50 μm.

1277

Figure 2—figure supplement 3 PG neurons expressing PTTH do not regulate virgin female copulation rate during adult stage.

PG neurons were activated during adult stage by dTrpA1 at 29°C. The female copulation rate and the latency to copulation did not change significantly. The number of female flies paired with wild-type males is displayed in parentheses. For the copulation rate, chi-square test is applied. For the latency to copulation, Mann-Whitney U test is applied. Error bars indicate SEM. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, ns indicates no significant difference.

1278

1279

1280

1281

1282

1283

1284

1285

1286

1287

1288

1289

1290

1291

1292

1293

1294

1295

1296

1297

1298

1299

1300

1301

1302

1303

1304

Figure 5—figure supplement 1 Expression of EcR-A and EcR-B1 in pC1 and vpoDN neurons.

(A-C) pC1 neurons were colabeled by pC1-ss2 driving UAS-mCD8-GFP (green, A) and EcR-A antibodies (red, B). Magnification of green boxed region in (C) is shown in (D1–D3). (E-G) pC1 neurons were colabeled by pC1-ss2 driving UAS-mCD8-GFP (green, E) and EcR-B1 antibodies (red, F). Magnification of green boxed region in (G) is shown in (H1–H3). (I-K) vpoDN neurons were colabeled by vpo-ss1 driving UAS-mCD8-GFP (green, I) and EcR-A antibodies (red, J). Magnification of green boxed region in (K) is shown in (L1–L3). (M-O) vpoDN neurons were colabeled by vpo-ss1 driving UAS-mCD8-GFP (green, M) and EcR-B1 antibodies (red, N). Magnification of green boxed

1316   region in (O) is shown in (P1–P3). Scale bars for magnified regions are 5 μm, for others

1317   are 50 μm.

1318

1319

1320

1321

1322

1323

1324

1325

1326

1327

1328

1329

1330

1331

1332

1333

1334

1335

1336

1337

1338

1339

1340

1341

1342

1343

1344

1345

1346

1347

Figure 5—figure supplement 2 Reduced EcR in pC1 neurons reduces virgin female receptivity.

(A) Knock-down of EcR in pC1 neurons driven by pC1-ss2-GAL4 significantly decreased the copulation rate and increased the latency to copulation. (B) Mean velocity had no significant change when EcR-A was knocked down in pC1 neurons compared with controls (Kruskal-Wallis ANOVA and post hoc Mann-Whitney U tests, n = 8-11). The number of female flies paired with wild-type males is displayed in parentheses. For the copulation rate, chi-square test is applied. For the latency to copulation, Kruskal-Wallis ANOVA and post hoc Mann-Whitney U tests are applied. Error bars indicate SEM. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, ns indicates no significant difference.

Figure 5—figure supplement 3 Reduced EcR in vpoDN neurons has no effect on virgin female receptivity.

(A-C) Knock-down of EcR-A in vpoDN neurons driven by vpoDN-ss1-GAL4, vpoDN-ss2-GAL4 and vpoDN-ss3-GAL4 had no effect on virgin female receptivity. (D-F) Knock-down of EcR-B1 in vpoDN neurons driven by vpoDN-ss1-GAL4, vpoDN-ss2-GAL4 and vpoDN-ss3-GAL4 had no effect on virgin female receptivity. The number of female flies paired with wild-type males is displayed in parentheses. For the copulation rate, chi-square test is applied. For the latency to copulation, Kruskal-Wallis ANOVA and post hoc Mann-Whitney U tests are applied. Error bars indicate SEM. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, ns indicates no significant difference.

1388

Figure 5—figure supplement 4 Reduced EcR-A in pC1d neurons has no effect on

virgin female receptivity.

(A) Knock-down of EcR-A in pC1d neurons had no effect on virgin female copulation

rate and latency to copulation. The number of female flies paired with wild-type males is

displayed in parentheses. For the copulation rate, chi-square test is applied. For the

latency to copulation, Kruskal-Wallis ANOVA and post hoc Mann-Whitney U tests are

applied. Error bars indicate SEM, *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, ns

indicates no significant difference.

1397

1398

1399

1400

1401

1402

1403

1404

1405

1406

1407

1408

1409

1410

1411

1412   **Supplementary File 1.** The primers used for the verification of *ΔPtth* null mutant flies

1413   and for the real-time quantitative PCR of dopamine beta-monooxygenase (DBM).

1414

1415   **Supplementary File 2.** The differently expressed genes in whole brains between flies

1416   having reduced EcR-A expression in pC1 neurons and the control flies.

1417

1418   **Supplementary File 3.** The raw data of transcriptomes in whole brains when EcR-A

1419   was knocked down in pC1 neurons or not.

1420

1421   **Figure 1—Source Data 1.** Photo of nucleic acid electrophoresis and copulation time.

1422

1423   **Figure 2—Source Data 1.** Copulation time.

1424

1425   **Figure 3—Source Data 1.** Copulation time and walking speed.

1426

1427   **Figure 4—Source Data 1.** Copulation time.

1428

1429   **Figure 5—Source Data 1.** Copulation time, courtship index, number of eggs, number of

1430   vaginal plate opening (VPO), and number of ovipositor extrusion (OE).

1431

1432   **Figure 6—Source Data 1.** Fluorescence intensity, cell number and calcium activity.

1433

1434   **Figure 7—Source Data 1.** Relative mRNA level.

1435

1436   **Figure 1—figure supplement 1—Source Data 1.** Body weight, courtship index and

1437   walking speed.

1438

1439   **Figure 2—figure supplement 3—Source Data 1.** Copulation time.

1440

1441   **Figure 5—figure supplement 2—Source Data 1.** Copulation time and walking speed.

1442

1443   **Figure 5—figure supplement 3—Source Data 1.** Copulation time.

1444

1445   **Figure 5—figure supplement 4—Source Data 1.** Copulation time.

1446

A

PtthLexA>LexAop-GFP

♀

B

PtthLexA>LexAop-GFP

♀