Lysine 36 of *Drosophila* histone H3.3 supports adult longevity

John C. Brown[1,#], Benjamin D. McMichael[1,2,#], Vasudha Vandadi[1,#], Aadit Mukherjee[2], Harmony R. Salzler[1]*, A. Gregory Matera[1,2,3,4]*

[1]Integrative Program for Biological and Genome Sciences, University of North Carolina, Chapel Hill, NC, USA.
[2]Department of Biology, University of North Carolina, Chapel Hill, NC, USA
[3]Department of Genetics, University of North Carolina, Chapel Hill, NC, USA.
[4]RNA Discovery Center, Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, NC, USA.

[#]These authors contributed equally to this work

*Address correspondence to:
      Dr. Harmony Salzler
      Email: hsalzler@email.unc.edu
      Tel: (919) 962-4541

       or

      Dr. Greg Matera
      Email: matera@unc.edu
      Tel: (919) 962-4567

Abstract

Aging is a multifactorial process that disturbs homeostasis, increases disease susceptibility, and ultimately results in death. Although the definitive set of molecular mechanisms responsible for aging remain to be discovered, epigenetic change over time is proving to be a promising piece of the puzzle. Several posttranslational histone modifications (PTMs) have been linked to the maintenance of longevity. Here, we focus on lysine 36 of the replication-independent histone protein, H3.3 (H3.3K36). To interrogate the role of this residue in *Drosophila* developmental gene regulation, we generated a lysine to arginine mutant that blocks the activity of its cognate modifying enzymes. We found that an *H3.3B^{K36R}* mutation causes a significant reduction in adult lifespan, accompanied by dysregulation of the genomic and transcriptomic architecture. Transgenic co-expression of wild-type *H3.3B* completely rescues the longevity defect. Because H3.3 is known to accumulate in non-dividing tissues, we carried out transcriptome profiling of young vs aged adult fly heads. The data show that loss of H3.3K36 results in age-dependent misexpression of NF-κB and other innate immune target genes, as well as defects in silencing of heterochromatin. We propose H3.3K36 maintains the postmitotic epigenomic landscape, supporting longevity by regulating both pericentric and telomeric retrotransposition and by suppressing aberrant immune signaling.

Introduction

Eukaryotic genomes are condensed in the nucleus of each cell as chromatin, a dynamic blend of histones, proteins and DNA. Nucleosomes, the fundamental unit of chromatin, comprise ~150 base pairs of DNA wrapped around an octamer of histones. Nucleosomes not only allow organisms to compress and store vast amounts of genetic information, but they also serve as steric barriers to a variety of DNA-templated processes [1-3]. Accordingly, molecular mechanisms have evolved to alter and regulate genome accessibility. Included among these mechanisms for chromatin remodeling are the covalent post-translational modification (PTM) of histones and the incorporation of histone variants [4-6].

Histone tail residues sustain a remarkable diversity of PTMs, including methylation, acetylation, ubiquitylation, sumoylation, and phosphorylation [7, 8]. Histone PTMs are hypothesized to alter local chromatin states via direct and indirect methods. For example, lysine acetylation alters chromatin directly by neutralizing positively charged lysine residues and weakening electrostatic interactions between histones and DNA [9]. In contrast, other modifications function through recruitment of effector proteins [8].

The incorporation of variant histones further increases regulatory complexity. In *Drosophila*, the histone gene complex (*HisC*) encodes all five canonical histone proteins. It is comprised by a cluster of ~100 tandemly repeated, 5kb repeat units that encode H2A, H2B, H3, H4, as well as the linker histone H1. Canonical (replication-dependent) histones are synthesized and incorporated into chromatin during S phase of the cell cycle. Variant (replication-independent) histones are encoded by genes located outside of *HisC* and are synthesized and incorporated independently of DNA replication. Variant histones typically contain a few amino acid residue changes from their canonical counterparts, which can alter their PTMs, stability within the nucleosome, and interactions with other chromatin components [6].

Here, we focus on lysine 36 of the replication-independent histone, H3.3, and its role in regulating lifespan. H3.3 differs from the canonical H3 (H3.2) by four amino acids [10]. These few differences enable a number of unique protein-protein interactions and PTM preferences [11, 12]. Ascomycete genomes, including budding yeast and other fungi, possess only one non-centromeric H3 protein, which is orthologous to metazoan H3.3. In budding yeast, mutation of either H3K36 or Set2, the enzyme that catalyzes mono-, di- and tri-methylation of H3K36, causes replicative aging defects [13]. In nematodes, inactivation of met-1, the Set2 orthologue, also decreases lifespan, whereas loss of H3.3 decreases the lifespan of long-lived *C. elegans* mutants [14-16]. Although these results suggest that H3.3K36 plays a conserved role in metazoan lifespan regulation, direct genetic tests of a role for this histone residue in aging have been lacking.

To investigate the function of H3.3K36 in *Drosophila* gene expression regulation, we previously generated a lysine to arginine mutant, *H3.3B^{K36R}* [17]. Here, we show that an *H3.3B^{K36R}* mutation (expressed in an *H3.3A^{null}* background) causes age-dependent dysregulation of the transcriptome, resulting in a significantly reduced adult lifespan. This longevity defect can be rescued by ectopic expression of a wild-type *H3.3B* transgene. We

also show that loss of H3.3K36 histones causes overexpression of NF-κB target genes in the adult brain, a disruption in several key aging-related metabolic processes, and progressive deterioration in silencing of heterochromatin. Indeed, we found clear evidence for age-dependent upregulation of repetitive elements, particularly LTR-type retroposons. Remarkably, expression of LINE-like elements specifically involved in telomere maintenance was decreased. These and other findings lead us to propose that H3.3K36 supports longevity by promoting expression of genes involved in cellular homeostasis, by regulating the activity of transposable elements and by suppressing innate immune response genes in the adult brain.

Results

H3.3^{K36R} causes an adult lifespan defect

Drosophila histone H3.3 protein is encoded by two genes, *H3.3A* on the second chromosome and *H3.3B* on the X chromosome [18]. We previously engineered a missense mutation in *H3.3B* (*H3.3B^{K36R}*) using Cas9-mediated homologous DNA repair (for details, see [17]. By combining the *H3.3B^{K36R}* mutation with a deletion of *H3.3A* (*H3.3A^{null}*), we generated *H3.3^{K36R}* animals that completely lack H3.3K36 (Fig. 1A). We also utilized an *H3.3B^{WT}* transgene inserted onto the third chromosome [19] to generate *H3.3A$^{null-WTR}$* and *H3.3$^{K36R-WTR}$* (wild-type rescue) flies (Fig. 1A). As shown in Fig. 1B, the majority *H3.3A^{null}, H3.3^{K36R}*, and *H3.3$^{K36R-WTR}$* animals complete development and eclose as adults at similar frequencies to a wild type Oregon R (Ore-R) control. Thus, arginine substitution of H3.3K36 has no significant effect on organismal viability.

To understand the role that this residue might play in longevity, we carried out adult lifespan assays. We measured the lifespan of *H3.3^{K36R}* mutants and compared the results to those obtained for *H3.3A^{null}* control animals (Fig. 1C, graph). Note that the *H3.3A^{null}* mutation has a small, but significant decrease in median lifespan by comparison to an Ore-R wildtype strain (Fig. S1). As defined by the number of days (post-eclosion) at which 50% of the animals survived, the K36R mutation has a pronounced and highly significant effect on adult lifespan (Fig. 1C, table). Furthermore, ectopic expression of wildtype H3.3 protein fully rescued this defect. That is, the median survival time of *H3.3^{K36R}* mutants was roughly 33 days post-eclosion, whereas that of *H3.3$^{K36R-WTR}$* control animals was 87 days (Fig. 1C, graph). Thus, we conclude that mutation of H3.3 lysine 36 causes a substantial reduction in adult lifespan. Consistent with previous studies showing that H3.3 is essential for lifespan extension in nematodes [15], this work shows that lysine 36 plays a direct role in this process.

Gene expression analysis in young vs. old fly heads from *H3.3^{K36R}* mutants and controls

Organismal lifespan is influenced by multiple interconnected factors, many of which are unknown, and most of which are poorly understood [20-22]. Although cellular senescence is

a natural and beneficial process in certain contexts, the accumulation of senescent cells over time is thought to contribute to aging and age-related diseases [23]. Despite their relatively short lifespan, many fundamental mechanisms underlying aging are conserved between fruit flies and humans, making *Drosophila* a valuable model for studying senescence-related processes. The brain of the adult fly is composed primarily of diploid postmitotic cells that rely on histone variant-specific assembly mechanisms for nucleosome turnover [24]. Indeed, H3.3 is reportedly enriched in the adult fly brain, and its deposition increases with age [25]. Thus, brain cells should be particularly sensitive to potential effects of a histone H3.3 missense mutation.

To gain insight into mechanisms by which H3.3K36 might influence age-dependent regulation of gene expression, we carried out transcriptome profiling of young vs. old fly heads. As detailed in the Materials and Methods, adult head RNA was extracted from six replicates (three male and three female) of each age:genotype pair (young and old $H3.3A^{null}$ and young and old $H3.3^{K36R}$ animals). Libraries were prepared from total Ribo-minus RNA samples and subjected to 100nt, paired-end RNA-sequencing (RNA-seq) using standard procedures (see Methods for details). Principle component analysis of the dataset (Fig. 2A) shows that the samples segregate primarily on the basis of age (PC1) and sex (PC2). Furthermore, the old adult samples can be distinguished by genotype-specific differences (Fig. 2A).

In order to identify differentially expressed genes (DEGs), we plotted the log2 fold-change ($H3.3^{K36R}/H3.3A^{null}$) for each transcript, versus the mean of normalized counts for that transcript. Among the younger flies, we found that there were 1,421 genes (662 up, 759 down) that were differentially expressed (p-adj < 0.05) in the mutant vs. control samples (Fig. 2B). A similar M/A plot comparing the older mutant and control animals showed that there were 2,605 DEGs (1260 up and 1345 down, Fig. 2C). Deposition of H3.3 protein in adult heads has been shown to increase with age from ~17% of total H3 in newly-eclosed animals to ~34% in 20-25 day old flies like the ones used here [25]. Consistent with this finding, we identified nearly twice as many DEGs in the older dataset, 793 of which overlapped with those in the younger animals (Fig. 2D).

To maximize the chances of capturing H3.3-specific effects, we focused our attention on DEGs identified in the older age group, and compared them to an RNA-seq dataset prepared from similarly-aged female *Chd1* (*chromodomain-helicase-DNA binding protein 1*) null mutants, using a separate DESeq2 model only using data from females [25]. Chd1 protein is part of a histone chaperone complex that functions together with Hira (Histone regulator A) to specifically deposit H3.3 into chromatin in the adult brain and other post-mitotic tissues [24]. *Chd1* null animals display a very similar adult longevity phenotype to $H3.3^{K36R}$ (K36R) mutants [25]. Differential expression analysis of *Chd1* null animals identified roughly 12,000 DEGs (p-adj < 0.05), nearly all of which overlapped with the union of our two K36R comparisons (see supplementary Figure S2A). We therefore applied more stringent cutoffs (p-adj < 0.05, log2 FC > |1|) to highlight the genes with the largest magnitude changes in all three groups and found that the overlap was greatly reduced. For

this more stringent comparison, we also restricted it to our female K36R datasets, because the Chd1 DEGs were derived from female-only fly heads (Fig. 2E). Strikingly, the degree of overlap between the old K36R and Chd1 DEGs went from roughly 84% to around 43% (Fig. S2B). Moreover, all of the non-Chd1 overlapping transcripts in the old female K36R dataset remained above the higher stringency threshold (Fig. 2E). Thus, the K36-specific DEGs, generally represented larger magnitude changes.

Interestingly, the female K36R fly head transcriptome appears to be affected earlier, and disrupted to a greater extent, than that of the male (Fig. 3A). Parsing out the longevity data in Fig. 1 by sex revealed that female K36R mutants had a median lifespan of around 30 d, compared to 40 d for males (Fig. S3). Functional enrichment analysis of the old female DEGs produced a number of senescence-associated gene ontology (GO) terms. These include genes involved in metabolic pathways that include cytochrome P450 and other enzymes, such as oxidoreductases (Fig. 3B). In contrast, GO analysis of old male K36R mutants identified detoxification factors ,serine peptidases, and ecdysteroid insect hormone enzymes (Fig. 3B). Interestingly, brain extracellular region proteins were identified in both sexes. This category of genes is among the hallmarks of the senescence-associated secretory phenotype (SASP), which has been implicated as a driver of numerous age-related disorders [26-28]. Another prominent GO term category, identified in both males and females, centers around the oxidization and transport of glutathione (Fig. 3B). In studies from nematodes to mice, glutathione has been broadly implicated in brain disorders and cellular senescence [29-31]. Altogether, these data support the view that H3.3K36 is important for maintaining cellular homeostasis in the adult brain.

A statistical interaction model highlights defects in innate immune signaling.
Though informative, our comparison to the *Chd1* RNA-seq dataset [25] and our original DESeq2 model were unable to take into account the effects of both age and sex. We therefore wanted to design an experiment to interrogate the differential effects that age and sex have on *H3.3^{K36R}* flies, relative to *H3.3A^{null}* controls. As shown in Fig. 4A, we used DESeq2 and R to create a multifactorial statistical model with two genotypes (*H3.3^{K36R}* and *H3.3A^{null}*), four conditions (i.e. with age and sex as covariates) and the interaction between age and genotype (age:genotype). Using this interaction model to measure differential gene expression, we unsurprisingly found that the union set of DEGs (3,256) was very similar in size to that obtained using the non-interaction model (3,233 - i.e. compare the totals from Figs. 4B and 2D). Strikingly, the interaction model reduced the total number of statistically contributing changes (p-adj < 0.05) down to 483 genes (Table S1). Only 69 out of the 793 overlapping DEGs classified in the previous comparison (Fig. 2D) were found to be significant in the interaction model, and 23 new DEGs were uncovered (Fig. 4B).

A volcano plot of all the genes identified using the age:genotype interaction is shown in Fig. 4C. As illustrated, the distribution of the interaction DEGs (p-adj < 0.05, LFC > |1|, red dots) was roughly equal between up- and down-regulated genes. Notable among the genes whose expression went up are are numerous downstream targets of NF-kB signaling.

These include the antimicrobial peptides (AMPs) Drosomycin (Drs), Drosocin (Dro), Attacin A (AttA), Metchnikowin (Mtk) and Bomanins S1, S2, S3 and S5, Bc1, Bc3, T1, T2 T3, (Fig. 4C and Table S1). Other well known immune effectors include IBIN/CG44404, IM4/CG15231, IM14/CG33990, Bbd/CG18067, and SPH93/CG44404. These host defense factors are encoded by well known anti-bacterial and anti-fungal response genes that are targeted by the Toll and IMD signaling pathways [32, 33]. Indeed, the NF-kB ortholog dorsal (dl) is itself among the genes that are differentially upregulated in the mutants (Table S1).

Among the genes that were significantly reduced in expression are several notable transcription factors, including ovarian tumor (otu/CG12743), and caudal (cad/CG1759). Cad and otu are negative regulators of immune (IMD) signaling that, in the adult animal, have been implicated in the maintenance of homeostasis [34-36]. Thus not only are the downstream targets of NF-kB upregulated, but genes involved in dampening or turning off these pathways are also downregulated. Another indicator of a runaway immune response in the brains of aging K36R mutants is highlighted by a 5-fold upregulation of the *Stress-induced DNase* gene, *Sid/CG9989* (Table S1). Sid protein is highly induced by bacterial infection and is thought to protect cells from toxic effects of excess nucleic acids that are released during a vigorous immune challenge [37].

Consistent with the findings outlined above, GO analysis of the interaction DEGs (using a more stringent p-adj < 0.05 and LFC > |1| cutoff) identified these and many other genes implicating a hyperactivation of immune response pathways in the absence of an external infection (Fig. 4D). Although these pathways have been clearly implicated in numerous other studies of senescence and aging, the work here provides the first direct evidence for linking them to H3.3K36-dependent processes in animals (see Discussion).

Age-dependent loss of heterochromatic silencing.
Previously, we showed that *H3.3K36R* mutants eclose at significantly higher frequencies than those that lack both copies of *H3.3*, strongly suggesting that H3.3K36R protein is incorporated into chromatin and is partially functional. Chromatin profiling of wandering third instar wing discs showed that the *H3.3K36R* mutation had little effect on H3K27me3 levels overall, yet the mutant adults displayed mild homeotic transformations [17]. Hence, we were interested in determining which types of chromatin might be affected by loss of H3.3 lysine 36.

Kharchenko and colleagues have classified nine different *Drosophila* chromatin "states" that are based on a wide variety of underlying histone marks and binding factors [38]. We used genome coordinates, derived from profiling data in BG3 larval brain cells, that have been established for each of these states to carry out an analysis of the genes that were differentially expressed in our K36R mutants. Rather than focusing on a small subset of DEGs whose contribution to the phenotype is uncertain, we broadened this analysis to the union set of young and old DEGs (Fig. 2D). Genes whose coordinates did not overlap

with those of any single chromatin state by at least 50% were binned into a tenth (NA) category.

As shown in Fig. 5A, we plotted the mean Log_2 fold-change over time for the DEGs that fell into each of these ten categories (Chromatin States). The plot on the left within each color grouping measures fold-change between young K36R mutants vs controls, whereas the plot on the right does the same for the older animals. The color code and brief descriptions of the characteristics for each chromatin state [38] are provided in Fig. 5B. As illustrated, Chromatin States 1-3, which comprise roughly 30% of the genome and are characterized by 'active' marks, are relatively unchanged from young to old animals (Fig. 5A). DEGs in States 4, 7 and 8 are characterized by less active marks and showed the greatest degree of change over adult lifespan. In contrast, DEGs in States 5 and 6 trended towards lower expression levels over time (Fig. 5A). State 5 is reflective of the dosage-compensated male X chromosome (characterized primarily by high levels of H4K16ac), and the K36R DEGs in this category were highly enriched (245/266 genes) on the X chromsome, Fig. 5C. Base mean expression levels for DEGs in each state are provided for purposes of comparison (Fig. S4A). Surprisingly, the downward trend in expression levels in DEGs from State 5 was apparently not driven by males (Fig. S4B), suggesting that H3.3K36 may not play a direct role in X chromosome dosage compensation.

The DEGs in State 7 showed the greatest degree of expression change in young adults as well as in change over time (Fig. 5A). State 7 is characterized by high levels of heterochromatic H3K9me2/3 marks (Fig. 5B). Although there was no single chromosome arm that harbored the bulk of these genes, 7/30 DEGs were located on the primarily heterochromatic 4th chromosome. Interestingly, 21 of the remaining 23 DEGs were located in centromere-proximal locations along their respective chromosome arms. A metaplot of these 23 genes from State 7 is shown in Fig. 5D, showing that all but one of them were derepressed in the mutant. Note that the genome build for these regions is incomplete, and so the relative position of the centromere is not the same for each arm. Thus the metaplot is not meant to depict the precise location of each DEG, but rather to illustrate its general positioning along the arm. Nevertheless, the results collectively provide strong evidence that loss of H3.3 K36 results in derepression of genes located in heterochromatic regions of the genome.

H3.3[K36R] **mutants exhibit position effects in the adult eye**

In contrast with its typical role as a marker of active genes along chromosome arms, H3.3 is also known to be deposited by the ATRX•Daxx complex to maintain repressive pericentric and (sub)telomeric chromatin [39-41]. In mouse embryonic stem cells, loss of *H3f3*a and *H3f3b* (orthologs of *H3.3A* and *H3.3B* in flies) results in reduced levels of H3K9me3 and ATRX at telomeres [42]. The chromatin state analysis of *H3.3*[K36R] DEGs shown in Fig. 5 clearly implicates a disruption to H3K9-mediated silencing, particularly in centromere-proximal regions. However, due to the paucity of protein coding genes located near

telomeres, this approach is unable to evaluate the transcriptional status of telomere-proximal heterochromatin.

The fruitfly genome is rare among eukyotes because it does not encode a telomerase RNP [43]. Instead, telomere length is maintained by controlled retrotransposition of *HeT-A (Healing Transposon A), TART (Telomere Associated Retrotransposon)* and *TAHRE (Telomere Associated and HeT-A Related)* elements into a telomere-proximal region (Fig. 6A). Telomeric chromatin is comprised of two distinct epigenetic compartments that are believed to perform different functions. The so-called HTT (*HeT-A, TART, TAHRE*) array (Fig. 6A), is primarily marked by H3K9 methylation that is thought to be required for proper expression of the HTT array [43, 44]. Adjacent to the array are the Telomere Associated Sequences (TAS, Fig. 6A), which contain 40-60 copies of telomere-specific satellite repeats per chromosome arm [43, 45]. Interestingly, TAS repeats are marked by repressive H3K27me3 and Polycomb Group (PcG) proteins [46].

In order to assay the functional consequences of *H3.3^{K36R}* mutation at telomeres, we utilized a miniwhite reporter system pioneered by Wallrath and Elgin [47]. These and other reporters have been widely used to analyze position effect variegation (PEV) of the *white* gene in the adult *Drosophila* eye (Fig. 6A). We ingressed six different *hsp70-w* reporter transgenes, located either on 3R or YS, into the background of *H3.3A^{null}* or *H3.3^{K36R}* mutants. Five of these lines contain insertions in telomeric chromatin (two in TAS on 3R; two in the HTT array of the Y chromosome), and one of them (118-E12) is inserted near centric heterochromatin on 3R (Fig. 6B). Surprisingly, none of the HTT reporters on the Y chromosome showed appreciable differences in the intensity or extent of PEV in the K36R mutant eyes, compared to the controls (Fig. 6B). In contrast, each of the TAS reporters on 3R showed an increase in the overall intensity of red eye pigmentation compared to the basal levels of the control animals. Also note that Y chromosome reporters showed significant variegation, whereas the 3R lines had more uniform expression levels. Although the basal level of pigmentation in 118-E6 is quite high, and that of line 118-E12 is rather low, we consistently found slightly higher levels in the K36R animals (Fig. 6B and Fig. S4). This effect is more clearly seen in line 39C-62, which has moderate basal levels and demonstrably higher levels in the presence of the K36R mutation (Fig. 6B). These observations indicate that H3.3K36 may play a role in maintaining a suppressive chromatin environment within TAS as well as centric heterochromatin.

Loss of H3.3 K36 reduces expression of telomere-specific retroposons, but causes derepression of many other transposable elements
The role of ATRX•Daxx-mediated H3.3 deposition in promoting and maintaining the silencing of transposable elements (TEs) is a well documented feature of higher eukaryotic genomes [48-50]. Thus we were curious to directly determine whether transcription of telomeric transposons was altered. Because transposons are repetitive elements, traditional RNA-seq pipelines are not set up to properly apportion short sequencing reads to repeat-rich regions of the genome. Hence, differential analysis of transposable element (TE)

expression was performed using the open source software package, TEtranscripts [51]. A volcano plot of these data is shown in Fig. 6C, revealing a dramatic TE derepression phenotype. Prominent among the classes of *upregulated* TEs are LTR-type retrotransposons (Figs. 6C,D). Increasing mobilization of transposons leading to DNA damage has also been associated with aging [52, 53]. These data suggest that activation of repressed TEs may contribute to the reduced lifespan of *H3.3^K36R* mutant animals.

Conspicuous among the TE families that are *downregulated* in K36R mutants is the *HeT-A* family of LINE-like (non-LTR) retroposons involved in telomere length maintenance. As shown in Fig 6C, expression of transcripts corresponding to the *HeT-A* family is dramatically reduced in old K36R mutant vs. control fly heads. Note that we also found that expression of both *HeT-A* and *TART* elements was significantly reduced in the old K36R mutants if we employed the more commonly-used featureCounts tool instead of TEcounts (see Tables S2 and S3). Among the three telomeric retrotransposons, *HeT-A* elements are particularly important because the *HeT-A* Orf1p/Gag protein recruits its *TART* counterpart to chromosome ends [54, 55]. Thus, reduced expression of *HeT-A* elements would be expected to have serious consequences for telomeric stability [56]. Telomere shortening is a well known hallmark of cellular senescence; these data are consistent with the notion that defects in telomere maintenance contribute to the observed reduced lifespan phenotype in *H3.3^K36R* animals.

Discussion

Aging is a complex process that involves multiple factors contributing to the decline of cellular and organismal functions, ultimately leading to decreased lifespan. Cellular senescence is characterized by numerous features, including irreversible cell cycle arrest, altered gene expression patterns, metabolic changes, and the acquisition of a senescence-associated secretory phenotype (SASP). Senescent cells secrete pro-inflammatory cytokines, growth factors, and matrix remodeling enzymes, contributing to chronic inflammation, tissue dysfunction, and age-related pathologies. Epigenetic changes, including histone modifications, have been identified as crucial contributors to aging-related processes. In this study, we focused on the lysine 36 residue of the replication-independent histone protein, H3.3. Our results demonstrate that expression of H3.3K36R as the sole source of H3.3 protein causes a significant reduction in adult lifespan and dysregulation of the genomic and transcriptomic architecture. Genetic rescue of the longevity defect upon transgenic co-expression of wild-type *H3.3B* suggests a direct role of lysine 36 in supporting longevity. Additionally, our transcriptome profiling of young and aged adult fly heads reveals age-dependent misexpression of innate immune target genes and derepression of heterochromatin in H3.3K36R mutants. Although other studies have implicated innate immune and heterochromatic defects as contributing factors in the aging process, our study provides the first direct evidence linking them to H3.3K36.

Role of Toll/Imd Pathways in Aging and Immuno-senescence
The Toll/Imd pathways are thought to play a vital role in regulating immunity and are intertwined with the aging process. Knockdown of Toll/Imd pathway members extends lifespan, whereas their overexpression shortens it [57]. Immuno-senescence, characterized by age-dependent decline in immune function, is a hallmark of the aging process [58, 59]. Interestingly, both sexes become more susceptible to infection as they age, with different underlying mechanisms. For males, susceptibility is primarily posited to be caused by deterioration in barrier defenses, whereas for females, systemic immune defense senescence is thought to be the major contributor [60].

Our findings are also consistent with observations in mammals, wherein NF-κB activity increases over time in the brains of middle-aged and older mice relative to young mice [61]. The upregulation of AMP expression in aged *Drosophila* [62] further supports the connection between immune signaling and the aging process. Notably, downregulation of NF-κB has been shown to mobilize nutrients and extend lifespan, whereas its overexpression has been linked to neurodegeneration [61, 62]. Our investigation into the molecular consequences of the *H3.3^{K36R}* substitution highlights its involvement in regulating critical aspects of immune signaling and heterochromatic silencing. Using a statistical interaction model, we discerned the effects of age and sex on gene expression in *H3.3^{K36R}* mutants compared to controls. Notably, the expression of downstream targets of NF-κB signaling, such as antimicrobial peptides and other immune effectors, were upregulated in the mutants. This suggests that H3.3K36 may play a role in suppressing immune responses in the absence of external infection, thereby contributing to the maintenance of cellular homeostasis in the adult brain.

H3.3K36, Retrotransposition and Senescence
Understanding the intricate interplay between chromatin dynamics and the aging process should offer valuable insights into mechanisms of aging and age-related diseases. Our findings of disrupted heterochromatic silencing in H3.3K36R mutants align with observations of senescence-associated heterochromatin defects in aging insects [63, 64] as well as in senescent mammalian cells [13, 65]. Moreover, the enrichment of H4K16ac in senescent cells, a mark known to be associated with HIRA-mediated deposition of H3.3, suggests a dynamic and imbalanced chromatin environment during senescence.

Upon loss of H3.3K36, our analysis has revealed a disruption of H3K9-mediated silencing in both centromere- and telomere-proximal regions. This includes a robust downregulation of non-LTR retrotransposons that are critical for maintaining telomere length [54, 55]. In contrast, the observed derepression of genes and LTR-type retroposons in heterochromatic regions provides strong evidence that H3.3K36 is involved in maintaining repressive pericentric and subtelomeric chromatin, possibly through ATRX•Daxx-mediated deposition. Consistent with these observations, Groth and colleages have recently shown that DAXX provides additional functionality to the histone chaperone system by recruiting H3K9 methyltransferases [66]. Even more recently, the same group showed that a failure to

re-establish H3K9me in post-replicative mammalian chromatin causes derepression of LTR-type retroposons [67]. Together, the data suggest that, in certain parts of the genome, H3.3K36 plays a conserved and specific role in silencing these so-called genomic parasites.

We also observed that mutation of H3.3K36 results in reduced repression of reporters in H3K27me3 marked TAS, and reduced expression of chromatin State 6 genes (marked by H3K27me3 and Pc proteins). These findings are in line with our recent work suggesting a role for H3.3K36 in supporting the function of genes marked by H3K27me3 [17]. Studies in many organisms point to gene regulation by PcG proteins as another pathway that influences longevity [14, 68-70]. However, the evidence is inconsistent between organisms, and much remains to be learned mechanistically. Further work clarifying the relationship between H3.3K36, Polycomb mediated gene repression, and aging would be an interesting area of future investigation.

It is intriguing to speculate on mechanisms through which H3.3K36 might support retrotransposon expression at telomeres. One possibility is that this residue facilitates expression of piRNA pathway genes. The piRNA pathway functions to suppress transposon activity in the germline, including activity of the HTT array [71-74]. Consistent with our observation of downregulation of *HeT-A* and widespread derepression of other TE families in *H3.3^K36R* mutants, previous studies have observed differences in regulation of telomeric piRNA clusters compared to those that target non-telomeric TEs [72, 73]. Furthermore, analysis of a dominant negative *H3.3^K36M* model (causing reduction in methylation of both H3.2K36 and H3.3K36), showed that telomeric retrotransposons were downregulated by piRNA pathway genes [75]. In *H3.3^K36M* mutant eye-antennal discs, suppression of *HeT-A* levels was elicited by upregulation of the piRNA promoting gene, *krimp.* Importantly, the suppression of *HeT-A* levels in these mutants could be rescued solely by reducing *krimp* expression [75]. Of note, we also observe upregulation of *krimp* in old *H3.3^K36R* mutants (log2 fold change = ~1.3), as well as another piRNA pathway gene, *Ago3* (log2 fold change = ~1.2), see Table S1 for details.

Another mechanism through which H3.3K36 might sustain expression of the HTT array is by promoting Jil-1 kinase function. Phosphorylation of the Jil-1 target, histone H3 serine 10, antagonizes both methylation of H3K9 and gene repression [76, 77]. Jil-1 is a positive regulator of HTT transcript levels [56, 78]. Moreover, *Jil-1* and *H3.3^K36R* mutants both exhibit repression of *HeT-A* transcription and suppression of TAS-reporter transgenes [56; this work]. Recently, an H3K36me3 reader protein, Jasper, has emerged as an important factor for stabilization and recruitment of Jil-1 [79, 80]. *Jasper* mutants exhibit both an increase in H3K9me2 and a decrease in expression of *HeT-A* and *TART* transposons [79]. However, unlike *H3.3^K36R* mutants, *Jil-1* and *Jasper* mutants do not exhibit widespread derepression of transposons in other parts of the genome [79, 80]. An interesting area of future study would be to determine whether this difference might be due to activities specific to H3.2K36 vs H3.3K36.

In summary, we find that lysine 36 of *Drosophila* histone H3.3 is crucial for supporting adult longevity by regulating immune signaling, heterochromatic silencing, and

retrotransposon repression. The dysregulation of these processes in H3.3K36R mutants likely contributes to age-related changes in gene expression and chromatin organization, ultimately impacting organismal lifespan. Further investigations into the mechanistic details of how H3.3K36 functions in these pathways will shed light on the broader role of histone modifications in the aging process. Additionally, exploring the conservation of these mechanisms in other organisms, including mammals, may have implications for understanding aging and age-related diseases in humans. Future studies of this process in model organisms may also provide potential therapeutic targets to mitigate age-related decline and improve healthy aging.

Materials and Methods

Fly Husbandry and Genetics

We generated *H3.3^{K36R}* mutants through creation of a K36R missense mutation at *H3.3B*, and combining it with an *H3.3A* deletion (*H3.3AΔ*). Both *H3.3A^{null}* and *H3.3^{K36R}* animals are obtained through a cross and negative selection for CyO, twGFP chromosome. *H3.3A^{null}* animals are made by crossing yw; H3.3^{2x1}/CyO, twGFP with yw; Df(2L)Bsc110/CyO, twGFP. *H3.3^{K36R}* animals are made by crossing yw, H3.3B^{K36R}; H3.3^{2x1}/cyO, twGFP with yw, H3.3B^{K36R}; Df(2L)Bsc110/CyO, twGFP. The H3.3A^{2x1} and Df(2L)Bsc110 alleles were obtained from Bloomington Drosophila Stock Center (#68240, #8835), whereas the *H3.3B^{K36R}* mutation was generated as described in [17]. Crosses were carried out in cages sealed with grape juice agar plates with supplemental yeast paste. Plates were replaced daily and GFP-negative larvae were sorted into vials of cornmeal-molasses food. Experimental flies were raised at 25°C, and all experiments were performed at 25°C unless otherwise noted.

Adult lifespan assays

To assess longevity, parental stocks were mated on grape juice agar plates and larvae were sorted into vials of molasses food. Male and Female flies were kept in separate vials. Each vial contained twenty or fewer flies to avoid crowding, and was incubated on its side to reduce accidental deaths from animals getting stuck in the food. Dead flies were removed and survivors flipped into fresh food every 2-3 days to prevent death due to poor conditions. Survivors were re-counted and recorded after each flip. Flies that were killed or escaped during the experiment were removed from the count.

Fly head RNA extraction and RNA Sequencing

H3.3^{K36R} and *H3.3^{null}* animals were generated as above and were separated into groups based on the date of eclosion. Flies that eclosed within 48 hrs of each other were considered an age cohort. Young cohorts (0-2 days post-eclosion) were processed without further aging,

whereas old cohorts were processed after three weeks (21-25 days post-eclosion). Heads were removed from the rest of the carcass and total RNA was extracted. Briefly, 10-25 flies were anesthetized and placed in a 15 mL tube. The tube was submerged in liquid nitrogen for 5-10 seconds and then vortexed for 5-10 seconds to separate heads from carcasses. This process was repeated for a total of 5 times before pouring the contents of the tube into a set of stacked sieves (Cole-Parmer, 3"-Diameter Sieve, Full Height Stainless Steel, No. 25 on top of Cole-Parmer, 3"-Diameter Sieve, Full Height Stainless Steel, No. 40). Fly heads were recovered from the bottom sieve and either stored at -80°C or immediately used for RNA extraction. Fly heads were crushed in TRIzol reagent (Thermo Fisher) and RNA was purified with the RNA clean & concentrator-5 kit (Zymo research). Total (Ribo-minus) RNA-seq libraries, three replicates per genotype, were prepared using a Universal RNA-seq library preparation kit with NuQuant (Tecan). Sequencing was performed on a NovaSeq 6000SP (Illumina) producing 100 nt, paired-end reads.

RNA-seq Analysis
Quality control was performed with FastQC, ver. fastqc/0.11.7 [81], and reads were aligned using STAR [82] to genome build dm6. Aligned reads were counted with featureCounts [83] and differential gene expression analysis was performed with DESeq2 (v1.34.0) [84]. One outlying sample was removed from the downstream analysis. The statistical model used included age, sex, genotype and the interaction between age and genotype (age:genotype). A gene ontology analysis was then performed using g:Profiler [85]. Differential transposon analysis was performed using TEtranscripts [51]. The statistical model used in this case included sex and genotype, as we chose to compare old mutants to old control animals.

Position Effect Variegation Experiments
Position effect variegation (PEV) reporter lines were obtained from the Bloomington Drosophila Stock Center: 118E-12, stock #84015; 39C-62, #44261; 118E-16, #84106; 8-M114, #84095; 8-M76, #84096; and 8-M112, #84093. These PEV lines carry a variegating hsp70::mini-white insertion [47]. To assess the role of H3.3K36 in heterochromatin spreading, PEV males with an H3.3A^{2x1}/CyO 2nd chromosome were crossed to yw, H3.3BK36R; Df(Bsc110)/CyO, twGFP (H3.3BK36R) or yw; Df(Bsc110)/CyO, twGFP (H3.3AΔ) virgin females. Straight-winged male flies were age-matched, and flies that eclosed within 96h of each other were grouped together. Images were taken on the following day, so the flies were 1-5 days post-eclosion.

Figure Legends

Figure 1. *H3.3^{K36R}* mutation reduces adult lifespan relative to *H3.3A^{null}* and wild-type rescue (WTR) controls. **(A)** Detailed genotypes for *H3.3A^{null}*, *H3.3^{K36R}*, *H3.3A^{null-WTR}*, and *H3.3^{K36R-WTR}* animals. The notation *tg:H3.3B^{WT}* corresponds to the rescue transgene. **(B)** Developmental viability assay. For each genotype, pupation and eclosion frequencies (%) were calculated and plotted (50 larvae per biological replicate vial). The mean and SD of these percentages are shown**.** Oregon-R (Ore-R) was used as an additional control. **(C)** Adult longevity assays for *H3.3A^{null}* and *H3.3A^{null-WTR}* controls, and for *H3.3^{K36R}* and *H3.3^{K36R-WTR}* flies. Median lifespan was determined (dotted lines) by identifying the day at which 50% of the animals survived. Statistical comparison of survival curves using Gehan-Breslow-Wilcoxon tests are presented in the accompanying table. A Bonferroni correction for multiple comparisons was employed, resulting in the following adjusted significance values: * $p<0.0125$, ** $p<0.0025$, *** $p<2.5 \times 10^{-4}$, ****$p<2.5 \times 10^{-5}$.

Figure 2. Transcriptomic profiling of young and old adult fly brains. (A) Principle Component Analysis (PCA) of young (0-2 d post-eclosion) and old (21-25 d) *H3.3A^{null}* and *H3.3^{K36R}* fly heads, labeled by sex. **(B)** M/A plot comparing gene expression from mixed sex *H3.3^{K36R}* and *H3.3A^{null}* young fly heads. Red and blue dots represent differentially expressed genes (DEGs) that were significantly (adjusted p-value, p-adj < 0.05) up- or down-regulated, respectively. The number of DEGs in each direction is shown in the upper and lower corners. **(C)** M/A plot of mixed sex *H3.3^{K36R}* vs. *H3.3A^{null}* transcripts from old adults. Labeling and significance as per panel B. **(D)** Venn diagram comparing overall number of DEGs (p-adj <0.05) between *H3.3A^{null}* and *H3.3^{K36R}* genotypes, young and old mixed sex fly heads. **(E)** Venn diagram of DEGs (p-adj < 0.05 and Log$_2$ Fold-change ≥ |1|) from female *H3.3^{K36R}* (young or old) versus aged (20 d) female *Chd1* fly heads. Data for *Chd1* provided by [25].

Figure 3. Sex-specific analysis of significant differentially expressed genes. (A) Venn diagrams of age- and sex-specific differentially expressed genes, DEGs (p-adj < 0.05 and, LFC > |1|) between *H3.3^{K36R}* vs. *H3.3A^{null}* fly heads. Left diagram compares DEGs in young female with young male fly heads (0-2 d post-eclosion), and right diagram compares DEGs in old female with old male fly heads (21-25 d). **(B)** Gene Ontology (GO) analysis of the DEGs identified in panel A (old female and old male). The size of each dot is proportional to the number of genes contained within a given ontology term (gene count), and the fraction of those genes scoring significantly (gene ratio) is represented using a heatmap. Adjusted p-values (–Log$_{10}$ transformed) for each GO term were calculated and plotted separately for both females and males.

Figure 4. Analysis of the interaction between age and genotype in old and young fly heads. (A) Graphical representation of the DESeq2 model including interaction terms (design = sex + age + genotype + age:genotype). Solid arrows represent genotype:condition

terms, dotted arrows illustrate how the model sets reference levels. **(B)** Venn diagram of differentially expressed genes (p-adj < 0.05) in young *H3.3^K36R* vs. *H3.3A^null* (0-2 d post-eclosion) and old *H3.3^K36R* vs. *H3.3A^null* fly heads (21-25 d), compared to those identified using the interaction model (age:genotype). **(C)** Volcano plot of RNA-seq data obtained using the interaction model. Dotted lines represent significance cutoffs for adjusted p-value and $\log_2$ fold-change. Dots represent individual genes, color coded according to the key at right. **(D)** Gene Ontology (GO) analysis of DEGs (padj < 0.05 and LFC > |1|) that were determined using the interaction model. Adjusted p-values ($-\log_{10}$ transformed) for each GO term were calculated and plotted. The size of each dot is proportional to the number of genes contained within a given ontology term (gene count), and the fraction of those genes scoring significantly (gene ratio) is represented using a heatmap.

Figure 5. Chromatin state analysis of differentially expressed genes. (A) The combined set of differentially expressed genes (adjusted p-value < 0.05) from the non-interaction DESeq2 model for both Young and Old conditions for *H3.3^K36R* mutants relative to *H3.3A^null* controls was binned by predominant chromatin state [38]. Genes were binned to a given state if > 50% of the gene was marked by that state. The "NA" category was comprised of genes where no state color was > 50% of gene length (406 genes) or if chromatin states were undetermined when the original chromatin state study was published (55 genes). The mut/ctrl $\log_2$ fold-change value for binned genes was plotted separately for Young and Old conditions. **(B)** A legend for colors corresponding to each chromatin state depicted in panel A, with representative histone marks for each state. **(C and D)** For genes in States 5 and 7, mut/ctrl $\log_2$ fold-change values for the Old condition were plotted versus relative position along chromosome arms for all chromosomes, from centromere (Cen) to telomere (Tel). For State 5, the X chromosome genes are colored green, and autosomal genes, black.

Figure 6. Telomeric position effect and transposable element expression analysis. (A) Cartoon depicting key features of *Drosophila* telomeres. At the distal end of the telomere, a capping complex assembles adjacent to arrays of *Het-A, TAHRE,* and *TART* retrotransposons (HTT array). Proximal to that are is the TAS (Telomere Associated Sequence) Chromatin, which is characterized by the presence of complex satellite repeats that vary by chromosome. **(B)** Images of adult eyes. For each column, representative examples are shown for a particular reporter strain. Rows correspond to genotype (*H3.3A^null* control or *H3.3^K36R* mutant). The location and type of chromatin for each reporter strain is also indicated: Pericentric heterochromatin (CnHc), Telomere Associated Sequences (TAS), YS-TART (Y chromosome TART), and YS-HeTA (Y chromosome HeT-A). n-values for each experiment ranged from 16 to 44, see Fig. S5 for details. **(C)** Volcano plot of annotated transposable elements (TEs) analyzed using the TEcounts tool within TEtranscripts. Each dot or triangle represents the combined reads from all loci for a given TE subtype. Dotted lines represent significance cutoffs for adjusted p-value and $\log_2$ fold-change. Symbols are color coded according to the key in the upper left. LTR-type transposons are represented by

triangles, non-LTR elements are shown as dots. **(D)** Stacked bar graph, binned by whether TEs were upregulated or unchanged in the *H3.3^{K36R}* mutant. The relative proportion of a given type of transposable element (DNA, LINE, LTR, Satellite, and Other) in each bin are shaded as indicated in the legend.

References

1. Bell O, Tiwari VK, Thoma NH, Schubeler D. Determinants and dynamics of genome accessibility. Nat Rev Genet. 2011;12(8):554-64. Epub 2011/07/13. doi: 10.1038/nrg3017. PubMed PMID: 21747402.
2. Lorch Y, LaPointe JW, Kornberg RD. Nucleosomes inhibit the initiation of transcription but allow chain elongation with the displacement of histones. Cell. 1987;49(2):203-10. Epub 1987/04/24. doi: 10.1016/0092-8674(87)90561-7. PubMed PMID: 3568125.
3. Ransom M, Dennehey BK, Tyler JK. Chaperoning histones during DNA replication and repair. Cell. 2010;140(2):183-95. Epub 2010/02/10. doi: 10.1016/j.cell.2010.01.004. PubMed PMID: 20141833; PubMed Central PMCID: PMCPMC3433953.
4. Bannister AJ, Kouzarides T. Regulation of chromatin by histone modifications. Cell Res. 2011;21(3):381-95. Epub 2011/02/16. doi: 10.1038/cr.2011.22. PubMed PMID: 21321607; PubMed Central PMCID: PMCPMC3193420.
5. Clapier CR, Iwasa J, Cairns BR, Peterson CL. Mechanisms of action and regulation of ATP-dependent chromatin-remodelling complexes. Nat Rev Mol Cell Biol. 2017;18(7):407-22. Epub 2017/05/18. doi: 10.1038/nrm.2017.26. PubMed PMID: 28512350; PubMed Central PMCID: PMCPMC8127953.
6. Talbert PB, Henikoff S. Histone variants--ancient wrap artists of the epigenome. Nat Rev Mol Cell Biol. 2010;11(4):264-75. Epub 2010/03/04. doi: 10.1038/nrm2861. PubMed PMID: 20197778.
7. Li B, Carey M, Workman JL. The role of chromatin during transcription. Cell. 2007;128(4):707-19. Epub 2007/02/27. doi: 10.1016/j.cell.2007.01.015. PubMed PMID: 17320508.
8. Rothbart SB, Strahl BD. Interpreting the language of histone and DNA modifications. Biochim Biophys Acta. 2014;1839(8):627-43. Epub 2014/03/19. doi: 10.1016/j.bbagrm.2014.03.001. PubMed PMID: 24631868; PubMed Central PMCID: PMCPMC4099259.
9. Eberharter A, Becker PB. Histone acetylation: a switch between repressive and permissive chromatin. Second in review series on chromatin dynamics. EMBO Rep. 2002;3(3):224-9. Epub 2002/03/08. doi: 10.1093/embo-reports/kvf053. PubMed PMID: 11882541; PubMed Central PMCID: PMCPMC1084017.
10. Elsässer SJ, Goldberg AD, Allis CD. New functions for an old variant: no substitute for histone H3.3. Curr Opin Genet Dev. 2010;20(2):110-7. Epub 2010/02/16. doi: 10.1016/j.gde.2010.01.003. PubMed PMID: 20153629; PubMed Central PMCID: PMCPMC2860041.
11. Filipescu D, Szenker E, Almouzni G. Developmental roles of histone H3 variants and their chaperones. Trends Genet. 2013;29(11):630-40. Epub 2013/07/09. doi: 10.1016/j.tig.2013.06.002. PubMed PMID: 23830582.
12. McKittrick E, Gafken PR, Ahmad K, Henikoff S. Histone H3.3 is enriched in covalent modifications associated with active chromatin. Proc Natl Acad Sci U S A. 2004;101(6):1525-30. Epub 2004/01/21. doi: 10.1073/pnas.0308092100. PubMed PMID: 14732680; PubMed Central PMCID: PMCPMC341768.
13. Sen P, Dang W, Donahue G, Dai J, Dorsey J, Cao X, et al. H3K36 methylation promotes longevity by enhancing transcriptional fidelity. Genes Dev. 2015;29(13):1362-76. Epub 2015/07/15. doi: 10.1101/gad.263707.115. PubMed PMID: 26159996; PubMed Central PMCID: PMCPMC4511212.
14. Ni Z, Ebata A, Alipanahiramandi E, Lee SS. Two SET domain containing genes link epigenetic changes and aging in Caenorhabditis elegans. Aging Cell. 2012;11(2):315-25.

Epub 2012/01/04. doi: 10.1111/j.1474-9726.2011.00785.x. PubMed PMID: 22212395; PubMed Central PMCID: PMCPMC3306474.

15. Piazzesi A, Papic D, Bertan F, Salomoni P, Nicotera P, Bano D. Replication-Independent Histone Variant H3.3 Controls Animal Lifespan through the Regulation of Pro-longevity Transcriptional Programs. Cell Rep. 2016;17(4):987-96. Epub 2016/10/21. doi: 10.1016/j.celrep.2016.09.074. PubMed PMID: 27760329; PubMed Central PMCID: PMCPMC5081402.

16. Pu M, Ni Z, Wang M, Wang X, Wood JG, Helfand SL, et al. Trimethylation of Lys36 on H3 restricts gene expression change during aging and impacts life span. Genes Dev. 2015;29(7):718-31. Epub 2015/04/04. doi: 10.1101/gad.254144.114. PubMed PMID: 25838541; PubMed Central PMCID: PMCPMC4387714.

17. Salzler HR, Vandadi V, McMichael BD, Brown JC, Boerma SA, Leatham-Jensen MP, et al. Distinct roles for canonical and variant histone H3 lysine-36 in Polycomb silencing. Sci Adv. 2023;9(9):eadf2451. Epub 2023/03/02. doi: 10.1126/sciadv.adf2451. PubMed PMID: 36857457; PubMed Central PMCID: PMCPMC9977188.

18. Akhmanova AS, Bindels PC, Xu J, Miedema K, Kremer H, Hennig W. Structure and expression of histone H3.3 genes in Drosophila melanogaster and Drosophila hydei. Genome. 1995;38(3):586-600. Epub 1995/06/01. doi: 10.1139/g95-075. PubMed PMID: 7557364.

19. Penke TJR, McKay DJ, Strahl BD, Matera AG, Duronio RJ. Functional Redundancy of Variant and Canonical Histone H3 Lysine 9 Modification in Drosophila. Genetics. 2018;208(1):229-44. Epub 2017/11/15. doi: 10.1534/genetics.117.300480. PubMed PMID: 29133298; PubMed Central PMCID: PMCPMC5753860.

20. Aron L, Zullo J, Yankner BA. The adaptive aging brain. Curr Opin Neurobiol. 2022;72:91-100. Epub 2021/10/25. doi: 10.1016/j.conb.2021.09.009. PubMed PMID: 34689041; PubMed Central PMCID: PMCPMC8901453.

21. Garschall K, Flatt T. The interplay between immunity and aging in Drosophila. F1000Res. 2018;7:160. Epub 2018/03/01. doi: 10.12688/f1000research.13117.1. PubMed PMID: 29487742; PubMed Central PMCID: PMCPMC5806056.

22. Zhang R, Chen HZ, Liu DP. The Four Layers of Aging. Cell Syst. 2015;1(3):180-6. Epub 2016/05/03. doi: 10.1016/j.cels.2015.09.002. PubMed PMID: 27135911.

23. Childs BG, Durik M, Baker DJ, van Deursen JM. Cellular senescence in aging and age-related disease: from mechanisms to therapy. Nat Med. 2015;21(12):1424-35. Epub 2015/12/10. doi: 10.1038/nm.4000. PubMed PMID: 26646499; PubMed Central PMCID: PMCPMC4748967.

24. Konev AY, Tribus M, Park SY, Podhraski V, Lim CY, Emelyanov AV, et al. CHD1 motor protein is required for deposition of histone variant H3.3 into chromatin in vivo. Science. 2007;317(5841):1087-90. Epub 2007/08/25. doi: 10.1126/science.1145339. PubMed PMID: 17717186; PubMed Central PMCID: PMCPMC3014568.

25. Schoberleitner I, Bauer I, Huang A, Andreyeva EN, Sebald J, Pascher K, et al. CHD1 controls H3.3 incorporation in adult brain chromatin to maintain metabolic homeostasis and normal lifespan. Cell Rep. 2021;37(1):109769. Epub 2021/10/06. doi: 10.1016/j.celrep.2021.109769. PubMed PMID: 34610319; PubMed Central PMCID: PMCPMC8607513.

26. Naylor RM, Baker DJ, van Deursen JM. Senescent cells: a novel therapeutic target for aging and age-related diseases. Clin Pharmacol Ther. 2013;93(1):105-16. Epub 2012/12/06. doi: 10.1038/clpt.2012.193. PubMed PMID: 23212104; PubMed Central PMCID: PMCPMC4051295.

27. Mebratu YA, Soni S, Rosas L, Rojas M, Horowitz JC, Nho R. The aged extracellular matrix and the profibrotic role of senescence-associated secretory phenotype. Am J Physiol Cell

Physiol. 2023;325(3):C565-C79. Epub 2023/07/24. doi: 10.1152/ajpcell.00124.2023. PubMed PMID: 37486065.

28. Mavrogonatou E, Papadopoulou A, Pratsinis H, Kletsas D. Senescence-associated alterations in the extracellular matrix: deciphering their role in the regulation of cellular function. Am J Physiol Cell Physiol. 2023;325(3):C633-C47. Epub 2023/07/24. doi: 10.1152/ajpcell.00178.2023. PubMed PMID: 37486063.

29. Liu H, Wang H, Shenvi S, Hagen TM, Liu RM. Glutathione metabolism during aging and in Alzheimer disease. Ann N Y Acad Sci. 2004;1019:346-9. Epub 2004/07/13. doi: 10.1196/annals.1297.059. PubMed PMID: 15247041.

30. Gusarov I, Shamovsky I, Pani B, Gautier L, Eremina S, Katkova-Zhukotskaya O, et al. Dietary thiols accelerate aging of C. elegans. Nat Commun. 2021;12(1):4336. Epub 2021/07/17. doi: 10.1038/s41467-021-24634-3. PubMed PMID: 34267196; PubMed Central PMCID: PMCPMC8282788.

31. Iskusnykh IY, Zakharova AA, Pathak D. Glutathione in Brain Disorders and Aging. Molecules. 2022;27(1). Epub 2022/01/12. doi: 10.3390/molecules27010324. PubMed PMID: 35011559; PubMed Central PMCID: PMCPMC8746815.

32. Lindsay SA, Lin SJH, Wasserman SA. Short-Form Bomanins Mediate Humoral Immunity in Drosophila. J Innate Immun. 2018;10(4):306-14. Epub 2018/06/20. doi: 10.1159/000489831. PubMed PMID: 29920489; PubMed Central PMCID: PMCPMC6158068.

33. Hanson MA, Cohen LB, Marra A, Iatsenko I, Wasserman SA, Lemaitre B. The Drosophila Baramicin polypeptide gene protects against fungal infection. PLoS Pathog. 2021;17(8):e1009846. Epub 2021/08/26. doi: 10.1371/journal.ppat.1009846. PubMed PMID: 34432851; PubMed Central PMCID: PMCPMC8423362.

34. Ryu JH, Kim SH, Lee HY, Bai JY, Nam YD, Bae JW, et al. Innate immune homeostasis by the homeobox gene caudal and commensal-gut mutualism in Drosophila. Science. 2008;319(5864):777-82. Epub 2008/01/26. doi: 10.1126/science.1149357. PubMed PMID: 18218863.

35. Ryu JH, Nam KB, Oh CT, Nam HJ, Kim SH, Yoon JH, et al. The homeobox gene Caudal regulates constitutive local expression of antimicrobial peptide genes in Drosophila epithelia. Mol Cell Biol. 2004;24(1):172-85. Epub 2003/12/16. doi: 10.1128/MCB.24.1.172-185.2004. PubMed PMID: 14673153; PubMed Central PMCID: PMCPMC303351.

36. Ji S, Luo Y, Cai Q, Cao Z, Zhao Y, Mei J, et al. LC Domain-Mediated Coalescence Is Essential for Otu Enzymatic Activity to Extend Drosophila Lifespan. Mol Cell. 2019;74(2):363-77 e5. Epub 2019/03/19. doi: 10.1016/j.molcel.2019.02.004. PubMed PMID: 30879902.

37. Seong CS, Varela-Ramirez A, Tang X, Anchondo B, Magallanes D, Aguilera RJ. Cloning and characterization of a novel Drosophila stress induced DNase. PLoS One. 2014;9(8):e103564. Epub 2014/08/02. doi: 10.1371/journal.pone.0103564. PubMed PMID: 25083901; PubMed Central PMCID: PMCPMC4118900.

38. Kharchenko PV, Alekseyenko AA, Schwartz YB, Minoda A, Riddle NC, Ernst J, et al. Comprehensive analysis of the chromatin landscape in Drosophila melanogaster. Nature. 2011;471(7339):480-5. Epub 2010/12/24. doi: 10.1038/nature09725. PubMed PMID: 21179089; PubMed Central PMCID: PMCPMC3109908.

39. Drane P, Ouararhni K, Depaux A, Shuaib M, Hamiche A. The death-associated protein DAXX is a novel histone chaperone involved in the replication-independent deposition of H3.3. Genes Dev. 2010;24(12):1253-65. Epub 2010/05/28. doi: 10.1101/gad.566910. PubMed PMID: 20504901; PubMed Central PMCID: PMCPMC2885661.

40. Lewis PW, Elsaesser SJ, Noh KM, Stadler SC, Allis CD. Daxx is an H3.3-specific histone chaperone and cooperates with ATRX in replication-independent chromatin assembly at

telomeres. Proc Natl Acad Sci U S A. 2010;107(32):14075-80. Epub 2010/07/24. doi: 10.1073/pnas.1008850107. PubMed PMID: 20651253; PubMed Central PMCID: PMCPMC2922592.

41. Wong LH, Ren H, Williams E, McGhie J, Ahn S, Sim M, et al. Histone H3.3 incorporation provides a unique and functionally essential telomeric chromatin in embryonic stem cells. Genome Res. 2009;19(3):404-14. Epub 2009/02/07. doi: 10.1101/gr.084947.108. PubMed PMID: 19196724; PubMed Central PMCID: PMCPMC2661805.

42. Udugama M, FT MC, Chan FL, Tang MC, Pickett HA, JD RM, et al. Histone variant H3.3 provides the heterochromatic H3 lysine 9 tri-methylation mark at telomeres. Nucleic Acids Res. 2015;43(21):10227-37. Epub 2015/08/26. doi: 10.1093/nar/gkv847. PubMed PMID: 26304540; PubMed Central PMCID: PMCPMC4666390.

43. Mason JM, Frydrychova RC, Biessmann H. Drosophila telomeres: an exception providing new insights. Bioessays. 2008;30(1):25-37. Epub 2007/12/18. doi: 10.1002/bies.20688. PubMed PMID: 18081009; PubMed Central PMCID: PMCPMC2804870.

44. Cacchione S, Cenci G, Raffa GD. Silence at the End: How Drosophila Regulates Expression and Transposition of Telomeric Retroelements. J Mol Biol. 2020;432(15):4305-21. Epub 2020/06/09. doi: 10.1016/j.jmb.2020.06.004. PubMed PMID: 32512004.

45. Antao JM, Mason JM, Dejardin J, Kingston RE. Protein landscape at Drosophila melanogaster telomere-associated sequence repeats. Mol Cell Biol. 2012;32(12):2170-82. Epub 2012/04/12. doi: 10.1128/MCB.00010-12. PubMed PMID: 22493064; PubMed Central PMCID: PMCPMC3372267.

46. Boivin A, Gally C, Netter S, Anxolabehere D, Ronsseray S. Telomeric associated sequences of Drosophila recruit polycomb-group proteins in vivo and can induce pairing-sensitive repression. Genetics. 2003;164(1):195-208. Epub 2003/05/17. doi: 10.1093/genetics/164.1.195. PubMed PMID: 12750332; PubMed Central PMCID: PMCPMC1462534.

47. Wallrath LL, Elgin SC. Position effect variegation in Drosophila is associated with an altered chromatin structure. Genes Dev. 1995;9(10):1263-77. Epub 1995/05/15. doi: 10.1101/gad.9.10.1263. PubMed PMID: 7758950.

48. Groh S, Milton AV, Marinelli LK, Sickinger CV, Russo A, Bollig H, et al. Morc3 silences endogenous retroviruses by enabling Daxx-mediated histone H3.3 incorporation. Nat Commun. 2021;12(1):5996. Epub 2021/10/16. doi: 10.1038/s41467-021-26288-7. PubMed PMID: 34650047; PubMed Central PMCID: PMCPMC8516933.

49. Sadic D, Schmidt K, Groh S, Kondofersky I, Ellwart J, Fuchs C, et al. Atrx promotes heterochromatin formation at retrotransposons. EMBO Rep. 2015;16(7):836-50. Epub 2015/05/28. doi: 10.15252/embr.201439937. PubMed PMID: 26012739; PubMed Central PMCID: PMCPMC4515123.

50. Elsässer SJ, Noh KM, Diaz N, Allis CD, Banaszynski LA. Histone H3.3 is required for endogenous retroviral element silencing in embryonic stem cells. Nature. 2015;522(7555):240-4. Epub 2015/05/06. doi: 10.1038/nature14345. PubMed PMID: 25938714; PubMed Central PMCID: PMCPMC4509593.

51. Jin Y, Tam OH, Paniagua E, Hammell M. TEtranscripts: a package for including transposable elements in differential expression analysis of RNA-seq datasets. Bioinformatics. 2015;31(22):3593-9. Epub 2015/07/25. doi: 10.1093/bioinformatics/btv422. PubMed PMID: 26206304; PubMed Central PMCID: PMCPMC4757950.

52. Ochoa Thomas E, Zuniga G, Sun W, Frost B. Awakening the dark side: retrotransposon activation in neurodegenerative disorders. Curr Opin Neurobiol. 2020;61:65-72. Epub 2020/02/25. doi: 10.1016/j.conb.2020.01.012. PubMed PMID: 32092528; PubMed Central PMCID: PMCPMC7198348.

53. Cardelli M. The epigenetic alterations of endogenous retroelements in aging. Mech Ageing Dev. 2018;174:30-46. Epub 2018/02/20. doi: 10.1016/j.mad.2018.02.002. PubMed PMID: 29458070.

54. Pardue ML, Rashkova S, Casacuberta E, DeBaryshe PG, George JA, Traverse KL. Two retrotransposons maintain telomeres in Drosophila. Chromosome Res. 2005;13(5):443-53. Epub 2005/09/01. doi: 10.1007/s10577-005-0993-6. PubMed PMID: 16132810; PubMed Central PMCID: PMCPMC1255937.

55. Casacuberta E. Drosophila: Retrotransposons Making up Telomeres. Viruses. 2017;9(7). Epub 2017/07/30. doi: 10.3390/v9070192. PubMed PMID: 28753967; PubMed Central PMCID: PMCPMC5537684.

56. Silva-Sousa R, Lopez-Panades E, Pineyro D, Casacuberta E. The chromosomal proteins JIL-1 and Z4/Putzig regulate the telomeric chromatin in Drosophila melanogaster. PLoS Genet. 2012;8(12):e1003153. Epub 2012/12/29. doi: 10.1371/journal.pgen.1003153. PubMed PMID: 23271984; PubMed Central PMCID: PMCPMC3521665.

57. Khor S, Cai D. Control of lifespan and survival by Drosophila NF-kappaB signaling through neuroendocrine cells and neuroblasts. Aging (Albany NY). 2020;12(24):24604-22. Epub 2020/11/25. doi: 10.18632/aging.104196. PubMed PMID: 33232282; PubMed Central PMCID: PMCPMC7803524.

58. DeVeale B, Brummel T, Seroude L. Immunity and aging: the enemy within? Aging Cell. 2004;3(4):195-208. Epub 2004/07/23. doi: 10.1111/j.1474-9728.2004.00106.x. PubMed PMID: 15268753.

59. Muller L, Fulop T, Pawelec G. Immunosenescence in vertebrates and invertebrates. Immun Ageing. 2013;10(1):12. Epub 2013/04/04. doi: 10.1186/1742-4933-10-12. PubMed PMID: 23547999; PubMed Central PMCID: PMCPMC3637519.

60. Kubiak M, Tinsley MC. Sex-Specific Routes To Immune Senescence In Drosophila melanogaster. Sci Rep. 2017;7(1):10417. Epub 2017/09/07. doi: 10.1038/s41598-017-11021-6. PubMed PMID: 28874758; PubMed Central PMCID: PMCPMC5585412.

61. Zhang G, Li J, Purkayastha S, Tang Y, Zhang H, Yin Y, et al. Hypothalamic programming of systemic ageing involving IKK-beta, NF-kappaB and GnRH. Nature. 2013;497(7448):211-6. Epub 2013/05/03. doi: 10.1038/nature12143. PubMed PMID: 23636330; PubMed Central PMCID: PMCPMC3756938.

62. Kounatidis I, Chtarbanova S, Cao Y, Hayne M, Jayanth D, Ganetzky B, et al. NF-kappaB Immunity in the Brain Determines Fly Lifespan in Healthy Aging and Age-Related Neurodegeneration. Cell Rep. 2017;19(4):836-48. Epub 2017/04/27. doi: 10.1016/j.celrep.2017.04.007. PubMed PMID: 28445733; PubMed Central PMCID: PMCPMC5413584.

63. Wood JG, Hillenmeyer S, Lawrence C, Chang C, Hosier S, Lightfoot W, et al. Chromatin remodeling in the aging genome of Drosophila. Aging Cell. 2010;9(6):971-8. Epub 2010/10/22. doi: 10.1111/j.1474-9726.2010.00624.x. PubMed PMID: 20961390; PubMed Central PMCID: PMCPMC2980570.

64. Wood JG, Jones BC, Jiang N, Chang C, Hosier S, Wickremesinghe P, et al. Chromatin-modifying genetic interventions suppress age-associated transposable element activation and extend life span in Drosophila. Proc Natl Acad Sci U S A. 2016;113(40):11277-82. Epub 2016/09/14. doi: 10.1073/pnas.1604621113. PubMed PMID: 27621458; PubMed Central PMCID: PMCPMC5056045.

65. Dasgupta N, Lei X, Arnold R, Teneche MG, Miller KN, Rajesh A, et al. Histone chaperone HIRA, Promyelocytic Leukemia (PML) protein and p62/SQSTM1 coordinate to regulate inflammation during cell senescence and aging. bioRxiv. 2023;(10.1101):2023.06.24.546372. doi: https://doi.org/10.1101/2023.06.24.546372.

66. Carraro M, Hendriks IA, Hammond CM, Solis-Mezarino V, Volker-Albert M, Elsborg JD, et al. DAXX adds a de novo H3.3K9me3 deposition pathway to the histone chaperone

network. Mol Cell. 2023;83(7):1075-92 e9. Epub 2023/03/04. doi: 10.1016/j.molcel.2023.02.009. PubMed PMID: 36868228; PubMed Central PMCID: PMCPMC10114496.

67. Wenger A, Biran A, Alcaraz N, Redo-Riveiro A, Sell AC, Krautz R, et al. Symmetric inheritance of parental histones governs epigenome maintenance and embryonic stem cell identity. Nat Genet. 2023. Epub 2023/09/05. doi: 10.1038/s41588-023-01476-x. PubMed PMID: 37666988.

68. Siebold AP, Banerjee R, Tie F, Kiss DL, Moskowitz J, Harte PJ. Polycomb Repressive Complex 2 and Trithorax modulate Drosophila longevity and stress resistance. Proc Natl Acad Sci U S A. 2010;107(1):169-74. Epub 2009/12/19. doi: 10.1073/pnas.0907739107. PubMed PMID: 20018689; PubMed Central PMCID: PMCPMC2806727.

69. Ma Z, Wang H, Cai Y, Wang H, Niu K, Wu X, et al. Epigenetic drift of H3K27me3 in aging links glycolysis to healthy longevity in Drosophila. Elife. 2018;7. Epub 2018/05/29. doi: 10.7554/eLife.35368. PubMed PMID: 29809154; PubMed Central PMCID: PMCPMC5991832.

70. Sen P, Shah PP, Nativio R, Berger SL. Epigenetic Mechanisms of Longevity and Aging. Cell. 2016;166(4):822-39. Epub 2016/08/16. doi: 10.1016/j.cell.2016.07.050. PubMed PMID: 27518561; PubMed Central PMCID: PMCPMC5821249.

71. Brennecke J, Aravin AA, Stark A, Dus M, Kellis M, Sachidanandam R, et al. Discrete small RNA-generating loci as master regulators of transposon activity in Drosophila. Cell. 2007;128(6):1089-103. Epub 2007/03/10. doi: 10.1016/j.cell.2007.01.043. PubMed PMID: 17346786.

72. Klattenhoff C, Xi H, Li C, Lee S, Xu J, Khurana JS, et al. The Drosophila HP1 homolog Rhino is required for transposon silencing and piRNA production by dual-strand clusters. Cell. 2009;138(6):1137-49. Epub 2009/09/08. doi: 10.1016/j.cell.2009.07.014. PubMed PMID: 19732946; PubMed Central PMCID: PMCPMC2770713.

73. Mohn F, Sienski G, Handler D, Brennecke J. The rhino-deadlock-cutoff complex licenses noncanonical transcription of dual-strand piRNA clusters in Drosophila. Cell. 2014;157(6):1364-79. Epub 2014/06/07. doi: 10.1016/j.cell.2014.04.031. PubMed PMID: 24906153.

74. Savitsky M, Kwon D, Georgiev P, Kalmykova A, Gvozdev V. Telomere elongation is under the control of the RNAi-based mechanism in the Drosophila germline. Genes Dev. 2006;20(3):345-54. Epub 2006/02/03. doi: 10.1101/gad.370206. PubMed PMID: 16452506; PubMed Central PMCID: PMCPMC1361705.

75. Chaouch A, Berlandi J, Chen CCL, Frey F, Badini S, Harutyunyan AS, et al. Histone H3.3 K27M and K36M mutations de-repress transposable elements through perturbation of antagonistic chromatin marks. Mol Cell. 2021;81(23):4876-90 e7. Epub 2021/11/06. doi: 10.1016/j.molcel.2021.10.008. PubMed PMID: 34739871; PubMed Central PMCID: PMCPMC9990445.

76. Bao X, Deng H, Johansen J, Girton J, Johansen KM. Loss-of-function alleles of the JIL-1 histone H3S10 kinase enhance position-effect variegation at pericentric sites in Drosophila heterochromatin. Genetics. 2007;176(2):1355-8. Epub 2007/04/17. doi: 10.1534/genetics.107.073676. PubMed PMID: 17435241; PubMed Central PMCID: PMCPMC1894597.

77. Zhang W, Deng H, Bao X, Lerach S, Girton J, Johansen J, et al. The JIL-1 histone H3S10 kinase regulates dimethyl H3K9 modifications and heterochromatic spreading in Drosophila. Development. 2006;133(2):229-35. Epub 2005/12/13. doi: 10.1242/dev.02199. PubMed PMID: 16339185.

78. Silva-Sousa R, Casacuberta E. The JIL-1 kinase affects telomere expression in the different telomere domains of Drosophila. PLoS One. 2013;8(11):e81543. Epub 2013/11/19. doi:

10.1371/journal.pone.0081543. PubMed PMID: 24244743; PubMed Central PMCID: PMCPMC3828246.

79. Albig C, Wang C, Dann GP, Wojcik F, Schauer T, Krause S, et al. JASPer controls interphase histone H3S10 phosphorylation by chromosomal kinase JIL-1 in Drosophila. Nat Commun. 2019;10(1):5343. Epub 2019/11/27. doi: 10.1038/s41467-019-13174-6. PubMed PMID: 31767855; PubMed Central PMCID: PMCPMC6877644.

80. Dou K, Liu Y, Zhang Y, Wang C, Huang Y, Zhang ZZ. Drosophila P75 safeguards oogenesis by preventing H3K9me2 spreading. J Genet Genomics. 2020;47(4):187-99. Epub 2020/06/06. doi: 10.1016/j.jgg.2020.02.008. PubMed PMID: 32499180; PubMed Central PMCID: PMCPMC7416749.

81. Andrews S. FastQC: a quality control tool for high throughput sequence data. Babraham Bioinformatics, Babraham Institute, Cambridge, United Kingdom; 2010.

82. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 2013;29(1):15-21. Epub 2012/10/30. doi: 10.1093/bioinformatics/bts635. PubMed PMID: 23104886; PubMed Central PMCID: PMCPMC3530905.

83. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics. 2014;30(7):923-30. Epub 2013/11/15. doi: 10.1093/bioinformatics/btt656. PubMed PMID: 24227677.

84. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550. Epub 2014/12/18. doi: 10.1186/s13059-014-0550-8. PubMed PMID: 25516281; PubMed Central PMCID: PMCPMC4302049.

85. Reimand J, Kull M, Peterson H, Hansen J, Vilo J. g:Profiler--a web-based toolset for functional profiling of gene lists from large-scale experiments. Nucleic Acids Res. 2007;35(Web Server issue):W193-200. Epub 2007/05/05. doi: 10.1093/nar/gkm226. PubMed PMID: 17478515; PubMed Central PMCID: PMCPMC1933153.

A

$$H3.3A^{null} = \frac{+}{+} \; ; \; \frac{H3.3A^{2x1}}{Df(2L)Bsc110} \; ; \; \frac{+}{+}$$

$$H3.3^{K36R} = \frac{H3.3B^{K36R}}{H3.3B^{K36R}} \; ; \; \frac{H3.3A^{2x1}}{Df(2L)Bsc110} \; ; \; \frac{+}{+}$$

$$H3.3A^{null-WTR} = \frac{+}{+} \; ; \; \frac{H3.3A^{2x1}}{Df(2L)Bsc110} \; ; \; \frac{tg:H3.3B^{WT}}{+}$$

$$H3.3^{K36R-WTR} = \frac{H3.3B^{K36R}}{H3.3B^{K36R}} \; ; \; \frac{H3.3A^{2x1}}{Df(2L)Bsc110} \; ; \; \frac{tg:H3.3B^{WT}}{+}$$

B

C

	$H3.3A^{null}$	$H3.3^{K36R}$	$H3.3A^{null-WTR}$	$H3.3^{K36R-WTR}$
$H3.3A^{null}$	---	$< 10^{-6}$ ****	$< 4.2 \times 10^{-3}$ *	---
$H3.3^{K36R}$	$< 10^{-6}$ ****	---	---	$< 10^{-6}$ ****
$H3.3A^{null-WTR}$	$< 4.2 \times 10^{-3}$ *	---	---	ns
$H3.3^{K36R-WTR}$	---	$< 10^{-6}$ ****	ns	---

Figure 1

Figure 2

A

Sex-specific DEGs

p-adj < 0.05, LFC > |1|

B

Gene Ontology Analysis

Figure 3

Figure 4

A Chromatin State Analysis of Differentially Expressed Genes

Log₂ Fold-change: mut/ctrl

Young Old

State # 1 2 3 4 5 6 7 8 9 NA
(Genes) (561) (603) (194) (161) (266) (85) (30) (199) (676) (461)

B

Color	State	Characteristics
	1	H3 K4me2/3, K9ac
	2	H3 K36me3, K79me1
	3	H3 K4me1, K18ac, K27ac
	4	H3K36me1, long introns
	5	H4K16ac
	6	H3K27me2/3
	7	H3K9me2/3 (high)
	8	H3K9me2/3 (mod)
	9	H1, low activity
	NA	< 50% any other state

C X enrichment

Log₂ FC: mut/ctrl

Cen — Tel

D Relative position

Log₂ FC: mut/ctrl

Cen — Tel

Figure 5

Figure 6