Phylogenomics and ancestral reconstruction of Korarchaeota reveals genomic adaptation to habitat switching

Guillaume Tahon[1*], Stephan Köstlbacher[1], Erik A. Pelve[2], Brett J. Baker[3, 4], Jimmy H. Saw[5], Laura Eme[6], Daniel Tamarit[7], Max Emil Schön[8] and Thijs J. G. Ettema[1*]

[1] *Laboratory of Microbiology, Wageningen University, 6708WE Wageningen, The Netherlands*

[2] *Department of Anatomy, Physiology and Biochemistry, Swedish University of Agricultural Sciences, Uppsala, Sweden*

[3] Department of Integrative Biology, University of Texas at Austin, USA

[4] Department of Marine Science, University of Texas at Austin, USA

[5] *Department of Biological Sciences, The George Washington University, Washington, DC 20052, USA*

[6] *Unité d'Ecologie, Systématique et Evolution, CNRS, Université Paris-Saclay, Paris, France*

[7] *Theoretical Biology and Bioinformatics, Utrecht University, 3584CH Utrecht, The Netherlands*

[8] *Department of Biomolecular Mechanisms, Max Planck Institute for Medical Research, 69120 Heidelberg, Germany*

* Correspondence:

Guillaume Tahon

guillaume.tahon@wur.nl

Thijs J. G. Ettema

thijs.ettema@wur.nl

Keywords

Archaea, microbial diversity, phylogeny, hyperthermophiles, hot spring, genome evolution

Abstract

Our knowledge of archaeal diversity and evolution has expanded rapidly in the past decade. However, hardly any genomes of the phylum Korarchaeota have been obtained due to the difficulty in accessing their natural habitats and – possibly – their limited abundance. As a result, many aspects of Korarchaeota biology, physiology and evolution remain enigmatic. Here, we expand this phylum with five high-quality metagenome-assembled genomes. This improved taxon sampling combined with sophisticated phylogenomic analyses robustly places Korarchaeota at the base of TACK and Asgard clades, revisiting the phylum's long-assumed position. Furthermore, we observe a clear split between terrestrial and marine thermal clades. Gene tree-aware ancestral reconstructions suggest that the last Korarchaeota common ancestor was a thermophilic autotroph. In contrast, Korarchaeaceae, the lineage where environmental transitions occurred, shifted towards a heterotrophic lifestyle. Terrestrial Korarchaeota gained many *cas* and CARF genes indicating they may need to manage viral infections. Together, our study provides new insights into these early diverging Archaea and suggests that gradual gene gain and loss shaped their adaptation to different thermal environments.

Importance

Korarchaeota are an ancient group of archaea, but their biology, physiology and evolution have remained obscure. Analysis of five novel Korarchaeota MAGs, and publicly available reference data provides robust phylogenomic evidence that Korarchaeota are placed at the base of Asgard archaea and TACK, revisiting the phylum's long-assumed position. Gene content reconstruction suggests a versatile thermophilic and autotrophic last Korarchaeota common ancestor. Environmental distribution surveying of public databases places all Korarchaeota in thermophilic environments and indicates that their habitat is limited to hydrothermal vents and hot springs. Our modeling indicates at least two transitions linked to habitat switching between these environments in the evolutionary history of Korarchaeota. Both are linked to a significant alteration of the inferred ancestral gene content, including a shift towards a heterotrophic and potential scavenging lifestyle. Furthermore, hot spring Korarchaeota acquired various genes participating in resistance to viruses, suggesting they may need to manage frequent viral threats.

Introduction

Archaea were only recognized as a third domain of life in 1977 (1), and were for a long time made up of only two phyla – Crenarchaeota and Euryarchaeota – represented by very few cultured strains (2). The third archaeal phylum to be discovered and the first one to be proposed without a cultured representative was *Candidatus* Korarchaeota (sometimes referred to as Korarchaeia (3) – hereafter referred to as Korarchaeota) (4). Amplicon sequences of the lineage were first sampled from Obsidian Pool, a near-neutral pH hot spring in the Hayden Valley of Yellowstone National Park (5). Subsequent 16S rRNA gene surveys and phylogenetic analyses supported the separation of this strain from the classical archaeal groups Euryarchaeota and Crenarchaeota, inspiring the name Korarchaeota, for the early divergence of the taxon in archaeal evolution ("koros/kore" referring to a young man/woman in Greek) (4). In 2008, ultrathin filamentous cells isolated from an anaerobic enrichment culture inoculated with Obsidian Pool sediment allowed initial characterization of a Korarchaeon - "*Candidatus* Korarchaeum cryptofilum" OPF8 (6). Its 1.59 Mb complete genome suggested an anaerobic heterotrophic lifestyle relying on a simple mode of peptide fermentation for generating biomass and energy. Also, the inability to synthesize various essential biomolecules indicated a potential dependency on other community members.

With the development of high-throughput DNA sequencing technologies and phylogenetic approaches in the four decades since the recognition of Archaea, the number and diversity of lineages described has increased exponentially. Currently, Archaea comprise four main clades (i.e., TACK, DPANN, Asgard and Euryarchaeota) which encompass over 30 phylum-level lineages and at least 50,000 species (See G. Tahon et al. (7) for a detailed overview). However, Korarchaeota have not undergone nearly the same expansion. Nearly 30 years after its initial discovery, very few Korarchaeota genomes have been sequenced, all of which have been recovered from either marine hydrothermal vents or terrestrial hot springs (Table S2) (8). Furthermore, all but the *Ca.* Korarchaeum cryptofilum OPF8 genome were published after 2018. As a result, many aspects of Korarchaeota biology, physiology and evolution have remained obscure. In addition, their phylogenetic position has been difficult to establish precisely. Initially, the two original environmental korarchaeotal 16S rRNA gene sequences branched deeply within Crenarchaeota, as a sister group

to Eukarya or branching off before the common ancestor of Cren- and Euryarchaeota
(4). Subsequent phylogenomic analyses based on concatenated sets of marker
genes have not managed to unequivocally resolve the position of Korarchaeota.
Depending on the phylogenetic method and dataset used, the phylum was placed
either at the base of the TACK superphylum, as a sister group of Crenarchaeota
inside TACK or as a sister lineage to Asgardarchaeota (9-13). However, the varying
and frequently low-supported position of Korarchaeota within Archaea may be
explained in part because of the use of less sophisticated models of evolution at the
time, as well as by the lack of sequence representative of the korarchaeotal diversity.
The latter led to the existence of a long branch at the base of Korarchaeota, which is
well-known to yield phylogenetic reconstruction artefacts (14).
In this project, we expand the Korarchaeota genomic and taxonomic diversity with
five distinct draft genomes obtained from sediment collected from the Little Hot Creek
terrestrial hot spring (California, USA) and the Taketomi Island shallow submarine
hydrothermal field of the Southern Ryukyu Archipelago (Japan). Using sophisticated
phylogenomic analyses relying on multiple marker protein sets, we do not only
accurately resolve the placement of Korarchaeota within the archaeal domain, but
also the internal topology of the phylum. Furthermore, using gene tree-species tree
reconciliation methods, we infer the evolution gene content along the korarchaeotal
tree and describe how living in different marine and aquatic terrestrial thermal
habitats shaped the genetic composition of the Korarchaeota.
122

Materials and Methods

Sampling

Samples were collected from the Taketomi Island shallow submarine hydrothermal field (TIV) of the Southern Ryukyu Archipelago (N24°20.9' E124°06.10') (15, 16) and from the Little Hot Creek terrestrial hot spring (N37°41.436' W118°50.664') (17, 18). Samples from Taketomi Island were collected as previously described by K. Zaremba-Niedzwiedzka et al. (19). Briefly, a sediment core was collected at 18 m depth, 5 m from the hydrothermal vent outflow by scuba diving, and material from the 10-15 cm section below the seafloor was extracted and stored at –25 °C and –80 °C. The core temperatures at 0, 20 and 30 cm below the sea floor were determined to be 31.0, 57.6 and 59.0 °C, respectively (RMT-2000; RIGO, Tokyo, Japan).

The Little Hot Creek (LHC) sample was collected as previously described (17, 18, 20). Briefly, cells from 80 °C sediment collected from the LHC4 hot spring in the Long Valley Caldera, California, USA, were separated by Nycodenz density centrifugation and stored either on ice (4 °C, in 10% v/v ethanol) or dry ice (80 °C, in 6% w/v betaine).

DNA extraction and sequencing

DNA from the TIV sample was extracted using the Power Max Soil DNA isolation kit (MoBio Labs) and purified with the Aurora system (Boreal Genomics).

Metagenome libraries from the sample were prepared from 1 ng purified DNA using the Nextera XT DNA library kit and sequenced with the Illumina HiSeq 2000 instrument at the Uppsala SNP&SEQ Technology Platform, generating 49 Gb of 2x250 bp paired-end reads, as previously described by Zaremba-Niedzwiedzka et al. (2017).

Cells from the LHC samples were sorted within 1 week of collection and subsequently lysed. The DNA was amplified using multiple displacement amplification (MDA) in a microfluidic device and individual cells were screened with PCR, as previously described by J. A. Dodsworth et al. (20). Subsequently, 1 µl of the original single-amplified genome (SAG) DNA of the Korarchaeon was used in another round of MDA to generate additional DNA of this SAG for library preparation (18). The MDA reaction was performed using the Qiagen REPLI-g Mini kit and purified with the QIAamp DNA Mini kit (Qiagen, Hilden, Germany). Next, a

sequencing library was generated from 1 ng of the resulting SAG DNA with the
NexteraXT library preparation kit (Illumina, San Diego, CA, USA) and the library was
then sequenced on an Illumina MiSeq instrument (2x300 bp), resulting in 4.9 Gbp of
raw data.
Additionally, DNA was extracted from LHC sediment using the FastDNA Spin Kit for
Soil (MP Biomedicals, Solon, Ohio, USA). Two metagenome libraries were generated
using the Nextera DNA Sample Preparation kit (Illumina): a MiSeq and a HiSeq
library which generated 7.3 and 10.8 Gbp, respectively.
165

Assembly and binning
Illumina adapters in the raw reads of each metagenome dataset were removed using
Scythe (https://github.com/vsbuffalo/scythe) with default parameters. Low quality
regions in the sequences were trimmed with Sickle (https://github.com/najoshi/sickle)
with the *-q 30* parameter to only keep reads with a PHRED score above 30.
Afterwards, reads were assembled using SPAdes (version 3.5.0;
https://github.com/ablab/spades) with the following settings *--meta -k 21, 33, 55*; and
IBDA-UA (version 1.1.1; (21)) with the following settings: *--maxk 124*. Assembled
contigs larger than 1 kbp were binned with the Emergent Self-Organizing Maps
approach (ESOM) (22) with the following parameters: *-min 5000 -max 10000 -t*
robust. korarchaeotal bins were visually inspected and manually selected using
Databionic ESOM Tools (23). Bins containing SSU genes displaying high similarity to
the *Ca.* K. cryptofilum SSU gene sequence were selected for further processing. A
description of the *Ca.* Korarchaeota archaeon LHC assembly can be found in J. H.
Saw et al. (18).
181

Cleaning of new and reference Korarchaeota bins
The five korarchaeotal bins (one from LHC and four from TIV) were screened for
non-korarchaeotal contigs using the R package multi-metagenome
(https://github.com/MadsAlbertsen/multi-metagenome) (24). Contigs with deviant
coverage, GC- or kmer content or with marker genes not phylogenetically clustered
to other korarchaeotal genes were manually inspected and removed if deemed not
likely to belong to the korarchaeotal clade.
Together with the five newly assembled Korarchaeota metagenome-assembled
genomes (MAGs), a total of 34 publicly available MAGs classified as Korarchaeota in

the NCBI (8), SILVA (25) and GTDB (3) databases were downloaded from NCBI in
October 2020. The completeness and contamination levels of all MAGs were
determined using CheckM v1.1.3 (26) and CheckM2 v1.0.1 (27). Additionally, MAGs
were screened using an in-house RP15 pipeline, as previously described (28).
Briefly, this pipeline was used to identify contigs that contain between 3 and 15
ribosomal proteins (RPs). The RPs found on these '*ribocontigs'* are then aligned to
the orthologous RPs from a set of reference taxa. Subsequently, the separate
alignments are trimmed, concatenated, and used to infer a phylogenetic tree using
FastTree 2 (*-wag model*) (29) and IQ-TREE 2.1.2 (*LG+C60+F+G4 model*, *-bb 1000 -*
bcor 0.98) (30).
Based on the position of ribocontigs in this tree, we observed that five of the publicly
available MAGs clustered outside of Korarchaeota and we removed them from
subsequent analyses.
204

Phylogeny of 16S rRNA genes
To construct a phylogenetic maximum likelihood (ML) tree of the TACK superphylum,
partial 16S rRNA gene sequences were extracted from Korarchaeota MAGs using
barrnap v0.9 (https://github.com/tseemann/barrnap) with the flags *--kingdom*
arc/bac/euk. Additionally, a total of 305 korarchaeotal sequences (Table S3) were
retrieved from the SILVA Ref NR database SSU r138.1 (25), together with 2367 near
full-length reference sequences (>1300 bp) representing the other taxa
encompassed in the TACK superphylum. Asgardarchaeaota 16S rRNA gene
sequences (>1300 bp; 305 sequences) were retrieved to be used as an outgroup.
Several Korarchaeota 16S rRNA gene sequences retrieved from the MAGs were
already present in the SILVA database and therefore removed from the dataset to
avoid duplicate sequence entries.
The dataset was aligned using MAFFT v7.310 (*--auto*) (31). Subsequently, the
alignment was processed using trimAl v1.4.rev22 to remove all columns with gaps in
more than 80% of the sequences (*-gt 0.2*) (32). The resulting alignment was then
used to construct a ML tree using IQ-TREE 2.1.2 (30), with settings: "*-m GTR+F -bb*
1000 -altr 1000 --bcor 0.98" (33).
The initial phylogenetic tree revealed that several publicly available Korarchaeota
MAGs contained contigs harboring non-korarchaeotal 16S rRNA gene copies. These
contigs were removed from the MAGs before downstream analyses. Additionally, two

Korarchaeota 16S rRNA gene sequences from the SILVA database were shown not

to cluster with this phylum. These non-Korarchaeota 16S rRNA gene sequences

were removed from the dataset after which the above steps were repeated.

If publicly available and documented, the sample origin (i.e., country and

environment) from which the MAGs and 16S rRNA gene sequences originate was

mapped onto the phylogenetic tree to visualize in which environments korarchaeotal

clades have been detected. All trees were visualized using iTOL v6 (34).

Environmental distribution assessment

To detect the presence of Korarchaeota in the environment, the 317 Korarchaeota

16S rRNA gene sequences were subjected to an integrated microbial NGS platform

(IMNGS) search using a similarity threshold of 90% and a minimum size of 200 bp

(35). From the results, sequences from datasets with less than 10.000 reads were

deleted. The remaining Sequence Read Archive (SRA) sequence hits returned by the

analysis were added to the aforementioned 16S rRNA gene alignment using the --

add command in MAFFT. This alignment was then used to construct a ML tree using

the GTR+F model and ultrafast bootstrap support values as described above. Only

SRA sequences clustering with Korarchaeota were retained from the IMNGS results.

The origin of these SRA datasets was retrieved from NCBI and used to create an

overview of the relative abundance of Korarchaeota in the environment (Table S4).

16S rRNA gene primer coverage of Korarchaeota

To analyze how well primers previously used to obtain Korarchaeota 16S rRNA gene

sequences target different Korarchaeota clades, amplification primer information

was, if documented, retrieved for the 294 PCR-amplicon-derived Korarchaeota 16S

rRNA gene sequences in SILVA (Table S3). These primer sequences were then

used in an *in silico* analysis which included all 317 Korarchaeota 16S rRNA gene

sequences, together with 12 recently designed universal primers to determine primer

coverage of different korarchaeotal lineages as described in G. Tahon et al. (7)

(Table S5).

Selection of Korarchaeota MAGs for species tree inference

To investigate the taxonomic diversity of Korarchaeota, average amino acid identity

(AAI) of verified Korarchaeota MAGs was calculated using EzAAI

(https://github.com/endixk/ezaai) (Table S6). Additionally, the 32 Korarchaeota MAGs
were used as input for a phylogenetic analysis pipeline using GToTree (36) with
settings "*-H Archaea -G 0.25",* together with a previous in-house selection of 119
archaeal proteomes spanning archaeal diversity with an emphasis on Asgard
archaeal representatives (Table S1). DPANN archaea were not included because
they are only remotely related to TACK archaea and tend to be long-branching taxa
in phylogenies, which can artefactually distort the inferred topology. The
concatenated alignment obtained by GToTree was subsequently used to construct a
ML phylogenetic tree with IQ-TREE v2.1.2 using 1000 ultrafast bootstraps and aLRT
under the LG+C60+F+G4 model. Using this tree, the 16S rRNA gene tree, CheckM
results, AAI values and the RP15 pipeline results, a subset of 18 MAGs - covering all
major korarchaeotal lineages (Table S2) - was selected for amalgamated likelihood
estimation (ALE) (37).
272

Phylogenetic analyses of concatenated protein alignments for species tree
inference
Two sets of phylogenetic markers were used to infer the species tree. The first one
(RP56) is based on a previously published dataset of 56 RPs (19). The second one
(NM54, for 'new markers') corresponds to 54 proteins extracted from a set of 200
markers previously identified as core archaeal proteins that can be used to robustly
infer the tree of archaea (38, 39).
For both RP56 and NM54 marker sets, we gathered orthologs from all korarchaeotal
predicted proteomes - predicted using prodigal (40) - as well as the other 119
archaeal proteomes. To retrieve orthologs from the set of selected taxa, we used
sequences from the previously published alignments (19, 38) as queries for BLASTp
against all proteomes. For each marker, the best BLAST hit from each proteome was
added to the dataset. For the first iteration, each dataset was aligned using MAFFT-
linsi (41) and ambiguously aligned positions were trimmed using BMGE 1.12 (*-m*
BLOSUM30) (42). ML individual protein phylogenies were inferred using IQ-TREE
v1.5.5 (43) under the LG+C20+F+G4 substitution model with 1000 ultrafast
bootstraps and were manually inspected.
Individual marker phylogenies were inspected for erroneous inclusion of paralogous
or contaminated sequences and to identify putative horizontal gene transfers. After

deleting dubious sequences, datasets were realigned and trimmed, and phylogenies
re-estimated as described above.

Trimmed alignments were then concatenated into two supermatrices (RP56 and
NM54 supermatrices). For each dataset, a phylogeny was inferred using IQ-TREE
under the LG+C60+F+G4 model. Statistical support for branches was calculated
using 1000 replicates for ultrafast bootstrap and aLRT analyses.

The RP56 and NM54 supermatrix alignments were used to create four more
alignments: an SR4-recoded alignment (44), an alignment where 50% of the fastest
evolving sites were removed, a χ^2-trimmed (28, 45) alignment where the 50% most
heterogeneous sites were removed in steps of 1% and a custom recoded alignment.
For the latter we calculated custom reduced amino acid schemes using the minmax
chisq method (https://www.mathstat.dal.ca/~tsusko/doc/minmax-chisq.pdf) (44) and
used these to recode sequence alignments.

For the 8 recoded or trimmed alignments, a phylogenetic ML tree was inferred using
1000 ultrafast bootstraps and aLRT under the GTR+G4+C60SR4 model (SR4-
recoded alignments) or LG+C60+F+G4 model (other alignments). The ten trees (i.e.,
two supermatrices, five alignments each) were subsequently used to infer 100 non-
parametric bootstrap trees using the posterior mean site frequency (PMSF) method
under the transfer bootstrap expectation (TBE) interpretation (*-b 100 --tbe*) (46).
Additionally, we ran four independent PhyloBayes-MPI v1.7a (47) Monte Carlo
Markov Chains (*-cat -gtr*) for each of the alignments. Results were checked to
determine whether convergence (maxdiff <0.3) and/or sufficient effective sample size
(effsize > 300) was reached. Subsequently, posterior predictive checks were
performed with PhyloBayes-MPI to determine whether the inferred phylogenetic
models adequately captured the across-taxa compositional heterogeneity and site-
specific pattern diversity present in the alignments.

Homology detection and protein family phylogenetic analysis
We used the annotation of arCOGs from EggNOG v5.0 (using eggnog-mapper
v2.0.5) (48) to assign all 290,917 proteins from 137 taxa (both MAGs and reference
genomes, Tables S1-2) into orthologous groups. All proteins without arCOG
annotation were subjected to *de novo* clustering using OrthoFinder v2.4.0 *(-os -M*
msa) (49), resulting in a total of 52,724 clusters (9,241 arCOG and 43,483 *de novo*).

For all protein clusters with at least four members (i.e. 9,108) multiple sequence
alignments were computed using MAFFT v7.475 E-INS-I (41) and subsequently
processed with trimAl v1.4.1 to filter gap-rich alignment columns (*-gt 0.01*) (32).
Single-gene trees were inferred from these 9,108 alignments with IQ-TREE v2.0.3
with 1000 ultrafast bootstraps (*-bb 1000 -bnni -wbtl*) (30, 33). A model test was
performed (*-m TESTNEW -mset LG -madd LG+C10, .., LG+C60*) for each tree
inference (50). For clusters with only two or three members, only a single topology is
possible for an unrooted tree and the unrooted gene trees were created for them.
333

Gene tree-aware ancestral reconstruction

ALE v0.4 (37) was applied to reconcile single gene trees with the species trees
based on ribosomal markers (RP56) and the new marker set (NM54). Firstly,
conditional clade probabilities were computed from the bootstrap samples
(*ALEobserve*) and 100 reconciliations with the species tree were sampled
(*ALEml_undated*). We thereby adjusted the reconstructed genome copy number with
the extinction probability per cluster within ALE (see
github.com/maxemil/ALE/commit/136b78e).
We applied a threshold of 0.3 to the raw reconciliation frequencies that are the output
of ALE, meaning that events (loss, transfer, origination or duplication) and
presence/absence (copies) were counted as such if they had a frequency of at least
0.3 (51). Singleton clusters were counted as originations at the corresponding
species node.
347

Annotation of carbohydrate-active enzymes and transporters

Presence and distribution of glycosyl hydrolases, glycosyl transferases,
polysaccharide lyases, carboxyl esterases, carbohydrate-binding modules and
auxiliary activities were identified through the dbCAN2 meta server (52), using the
CAZy database (July 2021). Carbohydrate-active enzymes predicted with at least two
of the integrated tools were used. Transporters were identified using TransAAP (July
2021) (53).
355

Korarchaeota optimal growth temperature

The optimal growth temperature of Korarchaeota MAGs was predicted using
OGT_prediction (54).

Inference of growth temperature of the last Korarchaeota common ancestor was
performed using the alignment and ML tree of the nucleoside diphosphate kinase
sequences from the MAGs, obtained during the protein family phylogenetic analysis.
Ancestral sequences and their unfolding midpoint temperatures (T_m) were predicted
using FireProt^{ASR} (55) and ProTstab2 (56), respectively.

Reverse gyrase phylogeny
To create a phylogenetic tree of reverse gyrase, all corresponding protein sequences
were retrieved from UniProt Release 2023_02 (57). Reverse gyrase sequences from
the 137 MAGs, if not yet present in UniProt, were added to this dataset. Sequences
were aligned with MAFFT v7.310 (*--auto*) after which the alignment was trimmed with
trimAl v1.4.rev22 to remove all columns with gaps in more than 80% of the
sequences (*-gt 0.2*). The ML phylogenetic tree was then generated using IQ-TREE
v2.0.3 (*-bb 1000 -alrt 1000*) with the model LG+C60+F+R8, as chosen by the
Bayesian information criterion (BIC) using ModelFinder Plus *(-m MFP -mset JTT,*
JTT+C10..C60,   WAG,   WAG+C10..C60,   LG,   LG+C10..C60,   Q.pfam,
Q.pfam+C10…+C60 -mrate G4, G6, G8, R4, R6, R8 -mfreq F) (50).

Relative evolutionary divergence
To determine a robust phylogenomic placement of a potential novel family-level
lineage placed at the base of Korarchaeota by F. Vulcano et al. (58), seven MAGs
from lineages Kg_2, Kg_3 and Kg_5, with a completeness estimation above 75%
were added to our dataset containing the original 137 MAGs, and MAGs classified as
Ca. Njordarchaeales. The single MAG from lineage Kg_4 could not be included due
to its low completeness. The expanded dataset was subjected to the same analyses
as before to create a NM54 supermatrix alignment. This alignment was χ^2-trimmed
as mentioned before, after which it was used to create a non-parametric bootstrap
tree using the PMSF method under the TBE interpretation. The resulting ML tree was
used as input for a phylogenomic analysis using the Castor package (59). Relative
evolutionary divergence values were calculated for all nodes of the tree to determine
whether this new lineage was part of the phylum Korarchaeota.

Code availability

All scripts needed to reproduce the protein clustering and running ALE are available

at   https://github.com/maxemil/korarchaeota-scripts   and

https://github.com/maxemil/ALE-pipeline.

Data availability

Sequencing raw reads and draft genome assemblies for the five new Korarchaeota

MAGs have been deposited in the NCBI GenBank database under the accession

numbers PRJNA983117 (TIV) and PRJNA983228 (LHC), respectively.

Results and Discussion

Contamination and misclassification among Korarchaeota

A first round of quality control using all contigs that contained (part of) the ribosomal protein (RP) operon, contigs that harbor partial 16S rRNA genes, and 76 single-copy archaeal marker proteins (see methods for detail) revealed that several of the publicly available metagenome-assembled genomes (MAGs) assigned as Korarchaeota were prone to contamination (Table S2). One MAG contained a 5 kbps contig harboring a bacterial 16S rRNA gene. Another MAG contained a contig harboring bacterial RPs. Finally, one MAG turned out to be a hybrid of bacterial and fungal contigs. In addition, some MAGs did not truly belong to Korarchaeota. Among these, three MAGs were found to represent *Ca.* Njordarchaeales, a recently described clade belonging to the Asgard archaea (39, 60), rather than Korarchaeota. A third MAG (DRAE01.1) was found to represent a yet undescribed lineage in TACK. Although sometimes classified as a lineage at the base of Korarchaeota, this placement was not supported and resulted from inferior phylogenomic analyses (3, 58). These findings underline that automated results generated by commonly used binning software, quality control tools and phylogenomic placement pipelines such as CheckM (26) and GTDB-tk (61) cannot be blindly trusted, because these tools do not perform phylogenetic validation of marker genes. For example, hybrid MAGs may have high completeness and low contamination values and even have robust phylogenetic placement. Therefore, thorough quality control and manual curation of MAGs is needed prior to their public deposit and classification. Phylogenetic analyses of the MAGs verified to be Korarchaeota, based on the 16S rRNA gene, contigs encoding at least 3 ribosomal proteins encoded on of a conserved ribosomal protein operon, and 76 single-copy archaeal marker proteins all revealed that Korarchaeota formed a monophyletic group in Archaea, however, with an unstable phylogenetic position and internal topology (Figs. S1-3). This urged us to perform in-depth phylogenomic analyses to resolve the phylogenetic position of Korarchaeota and its internal topology.

Korarchaeota are placed at the base of the 'ATACK superphylum'

The availability of 32 *bona fide* Korarchaeota MAGs in October 2020 allowed us to robustly place the phylum in the archaeal tree of life. To this end, we subsampled the

dataset to reduce genomic redundancy of highly similar taxa in order to reduce
computational burden. More Korarchaeota MAGs became publicly available after our
initial selection (8, 58, 62). The vast majority of these comprised taxa already
covered by our dataset. Some MAGs, however, represented a novel order at the
base of Korarchaeota (58), as supported by estimation of relative evolutionary
divergence values (Fig. S4). These MAGs became public when our project was in its
final stage and could thus not be included in the analyses anymore. Further analysis
of these MAGs indicated they did not change our findings.

Based on our initial phylogenomic tree including all MAGs (Fig. S3) and average
amino acid identity (AAI) (Table S6) (63), we observed that the 32 MAGs could be
grouped into 20 species, eight genera, four families and one order (Table S2, Fig.
S3). The five MAGs obtained from in-house sequencing efforts all represented
different species – four of which were novel – belonging to different genera. A total of
18 MAGs from all genera were then selected (with completeness >65%) for
downstream analyses. For each taxon, we here propose *Candidatus* names that will
be used throughout the text. Details on the taxonomic descriptions are given in the
Supplementary Note. All names and the relevant nomenclatural types have been
proposed under the SeqCode (64).

A first set of phylogenomic trees was inferred using 56 concatenated conserved
ribosomal proteins (RP56) (39). In addition to the original concatenation, we
performed four treatments aiming to reduce potential phylogenetic reconstruction
artefacts yielded by saturation and compositionally biased sites (39, 65). For this, we
first recoded our alignments into four character-states (SR4-recoding or custom-
recoding) (44). We also removed the 50% fastest-evolving sites. Finally, we used a
χ^2 trimmer to remove the 50% most compositionally heterogeneous sites (28). The
initial alignment and its four derivations (i.e., SR4-recoded, custom recoded, 50%
fastest evolving sites removed, and χ^2-trimmed) were then used to infer maximum
likelihood (ML) trees using the posterior mean site frequency (PMSF) method (Figs.
S5-9), and for phylogenetic reconstruction using PhyloBayes-MPI (Figs. S10-14).
Phylogenies based on all variations of the supermatrix, whether reconstructed in ML
or Bayesian framework, showed Korarchaeota branching at the base of the TACK
superphylum, although statistical support ranged from weak to high (80-99%
bootstrap support; 0.69-1 posterior probability). The only exception was the Bayesian
inference based on the RP56-SR4-recoded dataset, which placed Korarchaeota

nested within TACK. Regarding the internal topology, we consistently recovered two

distinct groups of MAGs originating from hot springs, representing the genera *Ca*.

Korarchaeum and *Ca*. Methanodesulfokora, respectively. However, these were never

found to be monophyletic, since the hydrothermal vent MAG representing the genus

Ca. Fumariihydrokora always placed as sister-lineage to *Ca*. Korarchaeum. The rest

of relationships between deep branches within the Korarchaeota varied depending

on the analysis. In particular, we observed various placements of the hydrothermal

vent genera *Ca*. Calidiprofundikora. and *Ca*. Sedimenticalidikora which are

characterized by high genomic GC content (Fig. 2). This is in part reflected by the

fact that, out of the five RP56 alignments, only the χ^2-trimmed one was appropriately

modelled by the sophisticated CAT-GTR model of evolution, based on bayesian

posterior prediction tests (Table S7). And even in this case, the monophyly between

the two clades from hot spring environments was poorly supported (Figs. S8 and

S13).

As previously shown, RPs can yield phylogenomic reconstruction artefacts

depending on the lineage considered (39). Despite them being highly conserved

across the different domains of life, they are short in length (66) resulting in short

concatenated alignments (i.e. 3575-7158 alignment positions in our case) and thus

limited phylogenetic signal (Table S7). Furthermore, RPs are all part of a unique

multisubunit complex, meaning they have undergone strong co-evolution. As a result,

the assumptions of most evolutionary models of independence of evolution at sites

are violated. Additionally, this implies that any departure from an average amino-acid

composition will be propagated across subunits, and the noise/signal ratio amplified

during phylogenetic reconstruction. This is particularly relevant since the amino-acid

composition of RPs is strongly influenced by adaptation to high temperature

environments, even more than the rest of the proteome (39). All Korarchaeota

genomes were obtained from thermal environments, and using genomic information

we estimated phylum members to have an optimal growth temperature of 70-85 °C

(Table S2). One recently published Korarchaeota MAG was suggested to be

mesophilic (62). However, the authors did not determine any optimal growth

parameters of the organism. Here, we determined the optimal growth temperature of

this MAG to be 79.3 °C, a value similar that of all other known Korarchaeota.

Therefore, it can be assumed that this organism is a thermophile and not a

mesophile. As a result of this adaptation to high temperature, the amino acid
composition of their RPs has evolved strong thermostability-related signatures, such
as a high ratio of charged versus polar amino acids, which can yield phylogenomic
artefacts (39). An illustrative example of how problematic this can be is the recent
case of the so-called *Ca*. Panguiarchaeales (67). Although described as a novel
korarchaeotal lineage, phylogenomic and AAI analyses (Fig. S15, Table S8) show
that all genomes described under this lineage name correspond to a low diverse
taxon that instead belongs to the *Ca.* Njordarchaeales, a hyperthermophilic Asgard
archaeal order (60). A similar problem arises in the work of J. Pan et al. (62), where
several *Ca*. Njordarchaeales MAGs were considered as Korarchaeota (i.e.,
3300022188_28 and GCA_011042815.1). *Ca*. Njordarchaeales have been shown to
artefactually branch outside Asgard archaea as a sister lineage to Korarchaeota
when inferring phylogenies based on RPs (39). The above examples show the
importance of constructing robust phylogenomic trees including a broad diversity of
reference data. Failure to do so (e.g., the work of Y. Liu and M. Li (68)) ultimately
leads to incorrect results and conclusions.

To overcome limitations and inconsistencies specific to RP datasets, a second
supermatrix alignment was constructed based on 54 core archaeal proteins that can
be robustly used to infer the tree of Archaea (39). This new marker (NM54)
supermatrix alignment and its four derivations were subjected to the same analyses
as the RP56 alignments. Contrary to the RP-based phylogenies, all resulting NM54-
based trees showed Korarchaeota not at the base of or within TACK, but instead at
the base of a large superphylum composed of 'TAC' (i.e., TACK clade without
Korarchaeota) and Asgard archaea, with high support (Figs. S16-24); altogether,
they compose the informally named ATACK superphylum. Given this revisited
position of Korarchaeota in the tree of Archaea using robust and sophisticated
phylogenomic analyses, we show Korarchaeota to be a phylum-level lineage and not
a class (i.e., *Ca*. Korarchaeia) of the Thermoproteota, as suggested by D. H. Parks et
al. (3). Additionally, the internal Korarchaeota topology of the NM54 supermatrix
alignments always showed a clear highly supported grouping of MAGs from hot
springs, with the hydrothermal vent lineage *Ca*. Fumariihydrokora interrupting their
monophyly. The χ^2-trimmed alignment was again the only one appropriately modelled
based on posterior prediction tests (Table 1). Contrary to its RP56 counterpart, the
consensus tree of the four PhyloBayes-MPI Monte Carlo Markov Chains now showed

maximum support for branches within Korarchaeota (Fig. 1). Additionally, the

PhyloBayes sampling statistics showed sufficient effective sample size, and posterior

prediction tests indicated that the model adequately captured the site-specific pattern

diversity present in the NM54-χ^2-trimmed alignment (Table 1). Therefore, we opted to

use this alignment and the resulting phylogeny as species tree for the ancestral

reconstruction analyses (below). We also performed phylogenomic analyses of our

dataset (of the untreated and four treated alignments) expanded with *Ca.*

Njordarchaeales (including the so-called *Ca.* Panguiarchaeales) MAGs. In the NM54

phylogenomic trees, contrary to the RP56 marker set, these lineages are not

misplaced outside of the Asgard archaea and next to Korarchaeota but remain in

place within the Asgard archaea (Fig. S25), thus showing that the NM54 marker set

is better suited to resolve deep evolutionary relationships compared to marker sets

based on ribosomal proteins. This highlights the importance of performing such

sophisticated phylogenomic analyses prior to ALE. The different yet more robust and

accurate placement of Korarchaeota in our study, compared to F. Vulcano et al. (58),

J. Pan et al. (62) and Y. Liu and M. Li (68) undoubtedly impacts downstream

analyses. For example, the previously estimated first divergence time of

Korarchaeota (62) – when placed at the base of TACK – will greatly differ. With

Korarchaeota placed at the base of ATACK, their estimated first divergence time

would be in the Archean, long before the Great Oxidation Event. Therefore,

Korarchaeota evolution was very unlikely driven by the availability of oxygen.

559   Korarchaeota have a limited and environmentally confined distribution

Interestingly, all *bona fide* Korarchaeota MAGs originated from either hot springs or

hydrothermal vents (Table S2), as was the case for all korarchaeotal 16S rRNA gene

sequences from SILVA (Table S3). Because of the limited number of MAGs and 16S

rRNA gene sequences, we also explored the phylogenetic diversity of the phylum

using ~280000 16S rRNA gene amplicon datasets included in the IMNGS platform

(35). Only 161 datasets were found to contain Korarchaeota sequences (Table S4),

as verified with a phylogenetic analysis (Fig. S2). Of these, only 5 and 24 had a

Korarchaeota relative abundance of more than 1% and 0.1%, respectively. All but 4

datasets originated from known hydrothermal environments (Fig. S26). Together,

these results suggest that Korarchaeota exhibit a high level of endemism and low

abundance, even in their niche.

To investigate potential explanations for the restricted distribution and limited
abundance of Korarchaeota, metadata was gathered for all MAGs, 16S rRNA gene
sequences and IMNGS datasets. The majority of amplicon-based 16S rRNA gene
sequences were obtained using Korarchaeota-specific and universal primers that
were designed at least one decade ago (Table S5). An *in silico* primer coverage
analysis showed that these primers almost exclusively target *Ca.* Korarchaeaceae
with no or relatively few mismatches (Table S5). At the time of designing, only 16S
rRNA gene sequences from these clades were available, which explains why these
primers fail to target more distant Korarchaeota lineages. Various more recently
developed universal primers, however, target the Archaea much better (Table S5)
(7). Accordingly, the IMNGS datasets containing the highest Korarchaeota relative
abundances were generated using these universal primers. Altogether, this suggests
that the limited availability of Korarchaeota data is the result not only of their true low
abundance and restricted distribution, but also of limited sequencing depth and the
use of inaccurate primers.

All phylogenomic trees (Figs. S3-25) showed the presence of two distinct, yet low
phylogenetically diverse clusters of MAGs originating from hot springs, representing
the genera *Ca.* Korarchaeum and *Ca.* Methanodesulfokora. Each hot spring cluster
was composed of a clade of multiple closely related MAGs from USA hot springs,
with a non-USA (i.e., Russia and New Zealand, respectively) MAG at their base.
Hydrothermal vent MAGs, on the other hand, form six distinct genera, their species
originating either from the Taketomi Island shallow submarine hydrothermal field,
Guaymas Basin or the Mid-Cayman Rise.

We wanted to check if the environmental and geographical split observed on
Korarchaeota MAG phylogenies could be expanded to the larger amount of 16S
rRNA gene sequence data available. The Korarchaeota topology of the 16S rRNA
gene trees (Figs. S1-2) revealed that, compared to hot spring lineages, only few
sequences grouped with the hydrothermal vent clades – especially the basal ones –
which could be explained by primer bias and undersampling of these latter
environments (see above). Overall, the two 16S rRNA gene trees were congruent
with the phylogenomic ones. However, the relationships between hot spring and
hydrothermal vent sequences were less well resolved. The hot spring lineages of *Ca.*
Methanodesulfokora and *Ca.* Korarchaeum were comprised of closely related
subclusters originating from Icelandic, Russian, USA and New-Zealand hot springs,

and Malaysian, New Zealand, Japanese, Russian, Chinese, Bulgarian and USA
locations, respectively (Table S3). Although similar in temperature and pH, these
locations have vastly different chemistries. Therefore, despite their lower
phylogenetic diversity, hot spring Korarchaeota could still be genetically distinct. To
date, however, no MAGs are available from most of these sites. Interestingly, *Ca.*
Methanodesulfokora contained one single hydrothermal vent sequence (accession
number AF411237.1), whereas *Ca.* Fumariihydrokora, close relative of *Ca.*
Korarchaeum, contained a single hot spring sequence (accession number
DQ300332.1). These sequences originated respectively from a coastal shallow water
hydrothermal vent and a geothermal well less than 100m from the beach, thus
possibly representing environments with mixed characteristics. Nevertheless,
biogeographic clustering suggests the existence of geographic variants.

All analyses support a Korarchaeota niche-invasion from marine to terrestrial
geothermal habitats and a high degree of endemism. How this transition exactly took
place and how genetically similar Korarchaeota from distant locations are remains
unclear. Genetically similar korarchaeotal hot spring populations in the USA have
been suggested to follow a spatial distribution resulting from a historical connection
or an invasion-type event (69). Although these hypotheses could explain similar
phylotypes in locations that are geographically close to each other, they fail to explain
the clustering of phylotypes from distant locations. Instead, Korarchaeota probably
adapted to specific environmental conditions of their respective habitats, and spread
globally. For example, many hot springs where Korarchaeota have been detected
share multiple environmental conditions (69, 70) that may have influenced the
evolutionary origin of the two major hot spring phylotypes.

The last Korarchaeota common ancestor: a versatile anaerobic autotrophic
thermophile
To explore the genetic basis of the transitions between hydrothermal vents and hot
springs or *vice versa*, and more generally how ancestral gene content evolved along
the Korarchaeota species tree, we reconciled the histories of gene families and
species tree using ALE (37). First, all proteins from the 137 MAGs were clustered
into gene families, after which single-gene trees were inferred for all gene families.
Gene family histories were then inferred by reconciling gene trees with the species
tree obtained from the NM54-χ^2-trimmed alignment (Fig. 1) to infer four types of

ancestral events (i.e., gene transfer, loss, duplication, and origination) as well as their
resulting gene copy numbers on all ancestral nodes of the species tree (Fig. 2, Table
S9). First, we screened the inferred gene content of the last Korarchaeota common
ancestor (LKorCA).

The ALE analysis suggested that LKorCA produced biomass autotrophically by
assimilating inorganic carbon via the Wood-Ljungdahl pathway, thus turning CO_2 into
acetyl-CoA (Fig. 3). Apart from the carbon monoxide dehydrogenase/acetyl-CoA
synthase, LKorCA also encoded the genetic potential of pyruvate:ferredoxin
oxidoreductase (*porABDG*). This key enzyme in anaerobic metabolism indicates that
LKorCA could further reduce and carboxylate acetyl-CoA to pyruvate and *vice versa*
(71).

LKorCA also contained a single *rbcL* copy, which encodes the archaeal type III
ribulose-1, 5-bisphosphate carboxylase-oxygenase (RuBisCO) of the Calvin-Benson-
Bassham (CBB) cycle. However, a complete CBB pathway was not present, which is
common in archaeal genome sequences (72). Instead, the presence of thymidine
phosphorylase (*deoA*) and eukaryotic translation initiation factor 2B (*e2b2*) hint that
rbcL is involved in the utilization of ribonucleotides and ribonucleosides via the
pentose-bisphosphate pathway (Fig. 3) (72, 73).

Inference of 3-hexulose-6-phosphate synthase (*hps*) and 6-phospho-3-
hexuloisomerase (*phi*) in LKorCA suggests LKorCA could potentially rely on the
reverse ribulose monophosphate pathway for biosynthesizing ribulose 5-phosphate
in absence of the pentose phosphate pathway (74).

The inferred genetic potential of LKorCA also showed evidence for assimilatory
sulfate reduction from sulfate to sulfite, a trait often found in conjunction with the
Wood-Ljungdahl pathway (Fig. 3). LKorCA likely dwelled in deep-sea hydrothermal
ecosystems. In these anoxic sediments niches carbon and sulfur cycling are
intimately linked, and the diffusion of sulfate from seawater makes sulfate reduction a
highly favorable process (75). Therefore, LKorCA might have played an important
role in sulfur cycling. For nitrogen intake, LKorCA relied on a glutamine synthetase
for the conversion of ammonium to glutamine. Phosphate, saccharides, metallic
cations and peptides could be imported using ABC-type transporters. The presence
of a complete branched-chain amino acid transport system (*liv* complex) suggests
LKorCA could take up and utilize protein-degradation products from its environment.
LKorCA also contained genes for the degradation of various amino acids, including

alanine, serine and aspartate. The presence of genes encoding proteins for β-
oxidation indicate at the genetic capacity of LKorCA to degrade fatty acids (Fig. 3).
LKorCA also encoded several cytoplasmic and membrane-bound complexes which
are relevant for energy conservation. Presence of the respiratory complex NADH-
ubiquinone oxidoreductase (*nuo*) would allow reoxidizing reduced ferrodoxin, which
can in turn be used as electron donor, as supported by the absence of *nuoEFG*, the
subunits required for the binding and oxidation of NADH (76). The Mvh/Hdr complex
potentially coupled the endergonic reduction of ferredoxin by H_2 to the exergonic
reduction of a disulfide bond (77). Presence of a FAD-containing hydrogenase (*glcD*)
could also indicate the use of HdrD/GlcD for heterodisulfide reduction (76). Hya and
Mvh [NiFe]-type hydrogenases act in mediating the oxidation of H_2, while an archaeal
ATPase (Ntp) ensures energy conservation (Fig. 3). In addition, LKorCA showed
genetic potential of HypA-F which are involved in the maturation of hydrogenases
(78).
Although a complete *de novo* pyrimidine synthesis pathway was inferred in LKorCA,
it lacked the purine counterpart. Therefore, LKorCA may have relied on transporting
nucleosides across the membrane using the ABC uptake system encoded by
nupABC and *bmpA* (Fig. 3).
Reverse gyrase, an enzyme which induces positive supercoiling in closed circular
DNA in an ATP-dependent manner (79) and unique to (hyper)thermophilic
organisms, was inferred as absent in LKorCA. However, all internal Korarchaeota
ancestors and MAGs have a reverse gyrase. Phylogenetic analysis revealed that
Korarchaeota reverse gyrase sequences are not monophyletic (Fig. S27).
Sequences from *Ca.* Caldabyssikora were distinctly placed from those of the other
Korarchaeota. Since Korarchaeota reverse gyrase sequences are not monophyletic,
we cannot exclude that LKorCA already had it based on our inferences and that it
might have been replaced in the last common ancestor of *Ca.* Caldabyssikora.
Although potential absence of reverse gyrase could indicate that LKorCA was not a
hyperthermophile, this is very unlikely. Firstly, reverse gyrase is not necessarily a
prerequisite for life at high temperatures, because the growth rate of a reverse
gyrase knockout of *Pyrococcus furiosus* was only impacted above 90 °C (80).
Therefore, the enzyme might not have been essential even if LKorCA had a
thermophilic lifestyle. Secondly, the last common ancestors of each of the major
archaeal clades were inferred to be thermophiles (81). Thirdly, nucleoside

diphosphate kinase is a well conserved enzyme of which the stability is directly
related to the host organism's environmental temperature (82). The unfolding
midpoint temperature of the reconstructed korarchaeotal ancestral nucleoside
diphosphate kinase sequence (T_m = 78 °C) was more stable than that of the
hyperthermophilic archaeon *Archaeoglobus fulgidus* DSM4304 (T_m = 62 °C) and thus
indicated LKorCA was a thermophile.

LKorCA contains several genes encoding antioxidant enzymes (e.g., superoxide
reductase, rubrerythrin). Previously, J. Pan et al. (62) suggested these genes to be
absent from the *Ca.* Caldabyssikora (i.e., Kor 7) and *Ca.* Thermotainarokora (i.e., Kor
8). However, our extended Korarchaeota dataset including five newly constructed in-
house MAGs, together with robust sophisticated phylogenomic analyses indicate that
this is not the case.

A first environmental transition is coupled to the switch to a likely
heterotrophic lifestyle and higher virus predation

The NM54-χ^2-trimmed alignment species tree indicated two environmental
transitions, both appearing within the *Ca.* Korarchaeaceae family (Fig. 1). The first
transition – from hydrothermal vents to hot springs – appeared at the branch before
the diversification of the hot spring genera *Ca.* Korarchaeum, *Ca.*
Methanodesulfokora and the hydrothermal vent genus Ca. Fumariihydrokora ('last
Ca. Korarchaeum, Methanodesulfokora and Fumariihydrokora common ancestor' -
LKMFCA). Upon this transition, the genetic repertoire of Korarchaeota was
significantly altered (Fig. 4). Contrary to LKorCA, LKMFCA was not an autotroph. The
Wood-Ljungdahl pathway, apart from *mch*, was lost in the last Korarchaeaceae
ancestor and replaced by the glycolysis pathway. The latter, however, lacks the
genes required for the reversible transformation between glucose and glucose-6-
phosphate. LKMFCA's lifestyle shifted towards that of fermentation, as could be
observed by the gain of the *acs* gene allowing the reversible conversion of acetyl-
CoA into acetate, and formate acetyltransferase.

Likewise, a shift took place in the sulfur metabolism, from assimilatory to dissimilatory
(DsrAB) reduction of sulfur compounds (Fig. 4). This complex, together with the Mvh
hydrogenase, indicates that hydrogen may have been used as an electron donor for
the reduction of sulfite (83). The nitrogen metabolism repertoire was significantly
expanded, including the glutamate dehydrogenase and glutamate synthase allowing

the conversion of ammonia and glutamine, respectively, to glutamate. Additionally,
presence of nitrate reductase suggests the ability to reduce nitrate to nitrite. A
formate dehydrogenase allowed the conversion of formate to carbon dioxide.
Compared to LKorCA, the genetic repertoire of LKMFCA also enlarged with various
glycosyltransferases, mainly of families GT2 and GT4 which are commonly involved
in processes such as N-glycosylation, cell wall biogenesis, cellulose, and sucrose
biosynthesis (84). Where the electron transport chain of LKorCA consisted of only
Complex I, that of LKMFCA was expanded with Complex II, the succinate
dehydrogenase. The environmental switch also coincided with major changes in the
amino acid metabolism of LKMFCA. Many of the biosynthesis pathways gained
genes (e.g., LeuA-D), whereas genes required for degradation were lost. Pathways
related to the metabolism of cofactors and vitamins were also subject to gene gain
and now included a complete NAD$^+$ biosynthesis pathway and greatly expanded
thiamine metabolism. Furthermore, LKMFCA gained multiple transporter systems
including those for arsenite efflux and transport of metal ions, amino acids, cations,
tungstate and nitrate/bicarbonate. As a result, LKMFCA had an increased potential to
scavenge its environment for nutrients and to efflux toxins out of its cell to survive
(85). Interestingly, at the time of environmental transition, LKMFCA gained many
genes participating in resistance to viruses, particularly those of type I and III *cas*,
and CARF systems (86, 87). Given that both environments are hyperthermophilic, it
is unclear exactly why these genes became enriched in hot spring Korarchaeota.
One potential explanation could be a different virome, although further research is
needed to answer this question. Several of the inferred type I and III genes (e.g.
Cmr4g7, Cmr1g7) are known to be associated with hyperthermophiles, indicating
they may be essential for managing viral infections in the hot spring habitats where
many archaeal viruses reside (88, 89). Others, such as the BREX system, might
allow phage adsorption while blocking its DNA replication (90). As such, the phage
could be used as a source of nucleotides in support of growth of LKMFCA (Fig. 4).
769

A reverse transition towards a scavenging or symbiont lifestyle
A second environmental transition, this time from hot spring to hydrothermal vent
adaptation, happened before the diversification of *Ca*. Korarchaeum and Ca.
Fumariihydrokora (LKFCA). While the central carbon metabolism of LKFCA did not
change compared to LKMFCA, many changes, particularly gene losses,

characterized this second transition (Fig. 5). LKFCA lost the ability to perform
dissimilatory reduction of sulfur compounds and to reversibly convert acetyl-CoA into
acetate. Likewise, the inferred genome of LKFCA is devoid of thymidine
phosphorylase (*deoA*), the eukaryotic translation initiation factor 2B (*e2b2*) and the
archaeal type III RuBisCO (*rbcL*) and thus lost the potential to use ribonucleotides
and ribonucleosides via the pentose-bisphosphate pathway. The thiamine
metabolism lost various key genes and was likely no longer functional. Given the
absence of the *acs* and *mobAB* genes, the interconversion of acetate and acetyl-
CoA, and synthesis of enzymes belonging to the DMSO reductase family were not
possible anymore, respectively. Additionally, the repertoire of LKFCA no longer
contained a tungstate ABC transporter. Despite many losses, LKFCA also gained
various novel genes and thus pathways. Like in the transition to LKMFCA, many
genes involved in amino acid biosynthesis were gained. Pathways for the
biosynthesis of valine, leucine, isoleucine, lysine and phenylalanine expanded as
well. A newly obtained spermidine/putrescine ABC transporter allowed regulating
these polyamines which play vital roles in ion homeostasis and cell proliferation.
Taken together, the second transition indicated that LKFCA underwent even more
metabolic changes towards a simplified genomic machinery. Contrary to the once
rather metabolically complex LKorCA, LKFCA transitioned to a simple anaerobic
heterotrophic lifestyle of peptides and amino acids (6). Whether LKFCA is just a very
efficient scavenger or a symbiont, or perhaps both, awaits to be proven with the
cultivation of Korarchaeum strains.
797
The last Methanodesulfokora common ancestor: a virus-combatting
methanogen
The last *Ca*. Methanodesulfokora common ancestor (LMCA) followed a different
evolutionary trajectory. Where LKFCA lost the *rbcL*, *e2be*, *deoA* and *dsrAB* genes,
and several key genes of the thiamine metabolism, LMCA kept this genomic
potential. LMCA also had an expanded amino acid metabolism, yet less extensive
than LKFCA. Several more type I and type III *cas* system genes participating in
resistance to viruses were acquired, indicating that *Ca*. Methanodesulfokora spp.
may need to manage additional viral threats in the hot springs where they reside (Fig.
6). Interestingly, we inferred in LMCA a whole suite of genes encoding methanol
methyltransferase (MtaABC) and methyl-coenzyme M reductase (McrABG) for

methanol reduction and methane production. Although *dsr* and *mcr* genes could hint
at a sulfite-dependent, anaerobic oxidation of methane to methanol (83), recent
cultivation of a strain of a *Ca*. Methanodesulfokora washburnensis has validated its
methylotrophic methanogenic potential (Personal communication with Roland
Hatzenpichler). Using genomic data, only *Ca*. M. washburnensis MAGs – all obtained
from USA hot springs – appear to have the Dsr, Mcr and Mta complexes. Whether
they are truly absent in the other *Ca*. Methanodesulfokora species, currently
comprises a single MAG obtained from a Russian hot spring, needs to be further
investigated by recovery of more complete MAGs and cultivated strains.
818

Conclusion
Despite rapid advancements in metagenomic sequencing analysis which particularly
expanded our view of the archaeal branch of the Tree of Life, our understanding of
Korarchaeota genomics is limited. Here, the addition of five new MAGs improved the
phylogenomic confidence, placing the phylum at the base of TACK and Asgard,
creating an ATACK superphylum, and showed a clear korarchaeotal endemism and
two habitat transition between marine and terrestrial geothermal habitats. The
inclusion of MAGs from all currently known korarchaeotal genera in gene tree-aware
ancestral genome reconstructions revealed that these environmental transitions were
subject to significant metabolic changes, thereby completely reshaping their genetic
composition. Our work provides new insights into the distribution, evolution and
physiology of the Korarchaeota and guides future cultivation-dependent and -
independent studies to further unravel its enigmatic biology and lifestyle.
832

Acknowledgements

The authors would like to express their appreciation to Nina Dombrowski and Kassiani Panagiotou for their support. We thank Peter Vandamme for advice on the etymology of the new names proposed.

Funding

This work was supported by grants from the European Research Council (ERC) (Consolidator grant 817834 to T.E. and Starting grant 803151 to L.E.), the Swedish Research Council (International Postdoc 2018-06609 to D.T.) and the Volkswagen Foundation (grant no. 96725 to T.E.). This work made use of the Dutch national e-infrastructure with the support of the SURF Cooperative using grant no. EINF-2953.

Conflict of interest

The authors declare there is no conflict of interest.

References

1. Woese CR, Fox GE. 1977. Phylogenetic structure of the prokaryotic domain: The primary kingdoms. Proceedings of the National Academy of Sciences 74:5088-5090.
2. Woese CR, Kandler O, Wheelis ML. 1990. Towards a natural system of organisms: proposal for the domains Archaea, Bacteria, and Eucarya. Proceedings of the National Academy of Sciences 87:4576-4579.
3. Parks DH, Chuvochina M, Rinke C, Mussig AJ, Chaumeil P-A, Hugenholtz P. 2021. GTDB: an ongoing census of bacterial and archaeal diversity through a phylogenetically consistent, rank normalized and complete genome-based taxonomy. Nucleic Acids Res 50:D785-D794.
4. Barns SM, Delwiche CF, Palmer JD, Pace NR. 1996. Perspectives on archaeal diversity, thermophily and monophyly from environmental rRNA sequences. Proceedings of the National Academy of Sciences 93:9188-9193.
5. Barns SM, Fundyga RE, Jeffries MW, Pace NR. 1994. Remarkable archaeal diversity detected in a Yellowstone National Park hot spring environment. Proc Natl Acad Sci U S A 91:1609-1613.
6. Elkins JG, Podar M, Graham DE, Makarova KS, Wolf Y, Randau L, Hedlund BP, Brochier-Armanet C, Kunin V, Anderson I, Lapidus A, Goltsman E, Barry K, Koonin EV, Hugenholtz P, Kyrpides N, Wanner G, Richardson P, Keller M, Stetter KO. 2008. A korarchaeal genome reveals insights into the evolution of the Archaea. Proceedings of the National Academy of Sciences 105:8102-8107.
7. Tahon G, Geesink P, Ettema TJG. 2021. Expanding Archaeal Diversity and Phylogeny: Past, Present, and Future. Annu Rev Microbiol 75:359-381.
8. NCBI Resource Coordinators. 2017. Database resources of the National Center for Biotechnology Information. Nucleic Acids Res 46:D8-D13.
9. Spang A, Caceres EF, Ettema TJG. 2017. Genomic exploration of the diversity, ecology, and evolution of the archaeal domain of life. Science 357:eaaf3883.
10. Dombrowski N, Williams TA, Sun J, Woodcroft BJ, Lee J-H, Minh BQ, Rinke C, Spang A. 2020. Undinarchaeota illuminate DPANN phylogeny and the impact of gene transfer on archaeal evolution. Nature Communications 11:3939.
11. Guy L, Ettema TJG. 2011. The archaeal 'TACK' superphylum and the origin of eukaryotes. Trends Microbiol 19:580-587.
12. Shu W-S, Huang L-N. 2022. Microbial diversity in extreme environments. Nat Rev Microbiol 20:219-235.
13. Williams TA, Cox CJ, Foster PG, Szöllősi GJ, Embley TM. 2020. Phylogenomics provides robust support for a two-domains tree of life. Nature Ecology & Evolution 4:138-147.
14. Susko E, Roger AJ. 2021. Long Branch Attraction Biases in Phylogenetics. Syst Biol 70:838-843.
15. Nakamura T, Yamazaki SS, Sakai K, Yamasaki H, Furushima Y, Yamamoto H. 2006. Acroporid corals growing over a methane-bubbling hydrothermal vent, Southern Ryukyu Archipelago. Coral Reefs 25:382-382.
16. Furushima Y, Nagao M, Suzuki A, Yamamoto H, Maruyama T. 2009. Periodic Behavior of the Bubble Jet (Geyser) in the Taketomi Submarine Hot Springs of the Southern Part of Yaeyama Archipelago, Japan. Mar Technol Soc J 43:13-22.

893   17.   Vick TJ, Dodsworth JA, Costa KC, Shock EL, Hedlund BP. 2010. Microbiology and
894   geochemistry of Little Hot Creek, a hot spring environment in the Long Valley
895   Caldera. Geobiol 8:140-154.

896   18.   Saw JH, Spang A, Zaremba-Niedzwiedzka K, Juzokaite L, Dodsworth JA, Murugapiran
897   SK, Colman DR, Takacs-Vesbach C, Hedlund BP, Guy L, Ettema TJG. 2015. Exploring
898   microbial dark matter to resolve the deep archaeal ancestry of eukaryotes.
899   Philosophical Transactions of the Royal Society B: Biological Sciences 370:20140328.

900   19.   Zaremba-Niedzwiedzka K, Caceres EF, Saw JH, Bäckström D, Juzokaite L, Vancaester
901   E, Seitz KW, Anantharaman K, Starnawski P, Kjeldsen KU, Stott MB, Nunoura T,
902   Banfield JF, Schramm A, Baker BJ, Spang A, Ettema TJG. 2017. Asgard archaea
903   illuminate the origin of eukaryotic cellular complexity. Nature 541:353-358.

904   20.   Dodsworth JA, Blainey PC, Murugapiran SK, Swingley WD, Ross CA, Tringe SG, Chain
905   PSG, Scholz MB, Lo C-C, Raymond J, Quake SR, Hedlund BP. 2013. Single-cell and
906   metagenomic analyses indicate a fermentative and saccharolytic lifestyle for
907   members of the OP9 lineage. Nature Communications 4:1854.

908   21.   Peng Y, Leung HCM, Yiu SM, Chin FYL. 2012. IDBA-UD: a de novo assembler for single-
909   cell and metagenomic sequencing data with highly uneven depth. Bioinformatics
910   28:1420-1428.

911   22.   Dick GJ, Andersson AF, Baker BJ, Simmons SL, Thomas BC, Yelton AP, Banfield JF.
912   2009. Community-wide analysis of microbial genome sequence signatures. Genome
913   Biol 10:R85.

914   23.   Ultsch A, Mörchen F. ESOM-Maps : tools for clustering , visualization , and
915   classification with Emergent SOM, p. *In* (ed),

916   24.   Albertsen M, Hugenholtz P, Skarshewski A, Nielsen KL, Tyson GW, Nielsen PH. 2013.
917   Genome sequences of rare, uncultured bacteria obtained by differential coverage
918   binning of multiple metagenomes. Nat Biotechnol 31:533-538.

919   25.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO.
920   2012. The SILVA ribosomal RNA gene database project: improved data processing
921   and web-based tools. Nucleic Acids Res 41:D590-D596.

922   26.   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM:
923   assessing the quality of microbial genomes recovered from isolates, single cells, and
924   metagenomes. Genome Res 25:1043-1055.

925   27.   Chklovski A, Parks DH, Woodcroft BJ, Tyson GW. 2022. CheckM2: a rapid, scalable
926   and accurate tool for assessing microbial genome quality using machine learning.
927   bioRxiv doi:10.1101/2022.07.11.499243:2022.07.11.499243.

928   28.   Martijn J, Vosseberg J, Guy L, Offre P, Ettema TJG. 2018. Deep mitochondrial origin
929   outside the sampled alphaproteobacteria. Nature 557:101-105.

930   29.   Price MN, Dehal PS, Arkin AP. 2010. FastTree 2 – Approximately Maximum-Likelihood
931   Trees for Large Alignments. PLoS One 5:e9490.

932   30.   Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A,
933   Lanfear R. 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
934   Inference in the Genomic Era. Molecular Biology and Evolution 37:1530-1534.

935   31.   Katoh K, Misawa K, Kuma Ki, Miyata T. 2002. MAFFT: a novel method for rapid
936   multiple sequence alignment based on fast Fourier transform. Nucleic Acids Res
937   30:3059-3066.

32. Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: a tool for
automated alignment trimming in large-scale phylogenetic analyses. Bioinformatics
25:1972-1973.
33. Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2017. UFBoot2:
Improving the Ultrafast Bootstrap Approximation. Molecular Biology and Evolution
35:518-522.
34. Letunic I, Bork P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for
phylogenetic tree display and annotation. Nucleic Acids Res 49:W293-W296.
35. Lagkouvardos I, Joseph D, Kapfhammer M, Giritli S, Horn M, Haller D, Clavel T. 2016.
IMNGS: A comprehensive open resource of processed 16S rRNA microbial profiles for
ecology and diversity studies. Sci Rep 6:33721-33721.
36. Lee MD. 2019. GToTree: A user-friendly workflow for phylogenomics. Bioinformatics
35:4162-4164.
37. Szöllősi GJ, Tannier E, Daubin V, Boussau B. 2014. The Inference of Gene Trees with
Species Trees. Syst Biol 64:e42-e62.
38. Petitjean C, Deschamps P, López-García P, Moreira D, Brochier-Armanet C. 2015.
Extending the Conserved Phylogenetic Core of Archaea Disentangles the Evolution of
the Third Domain of Life. Molecular Biology and Evolution 32:1242-1254.
39. Eme L, Tamarit D, Caceres EF, Stairs CW, Anda VD, Schön ME, Seitz KW, Dombrowski
N, Lewis WH, Homa F, Saw JH, Lombard J, Nunoura T, Li W-J, Hua Z-S, Chen L-X,
Banfield JF, John ES, Reysenbach A-L, Stott MB, Schramm A, Kjeldsen KU, Teske AP,
Baker BJ, Ettema TJG. 2023. Inference and reconstruction of the heimdallarchaeial
ancestry of eukaryotes. bioRxiv doi:10.1101/2023.03.07.531504:2023.03.07.531504.
40. Hyatt D, Chen G-L, LoCascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal:
prokaryotic gene recognition and translation initiation site identification. BMC
Bioinformatics 11:119.
41. Katoh K, Standley DM. 2013. MAFFT Multiple Sequence Alignment Software Version
7: Improvements in Performance and Usability. Molecular Biology and Evolution
30:772-780.
42. Criscuolo A, Gribaldo S. 2010. BMGE (Block Mapping and Gathering with Entropy): a
new software for selection of phylogenetic informative regions from multiple
sequence alignments. BMC Evol Biol 10:210.
43. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2014. IQ-TREE: A Fast and
Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies.
Molecular Biology and Evolution 32:268-274.
44. Susko E, Roger AJ. 2007. On Reduced Amino Acid Alphabets for Phylogenetic
Inference. Molecular Biology and Evolution 24:2139-2150.
45. Viklund J, Ettema TJG, Andersson SGE. 2011. Independent Genome Reduction and
Phylogenetic Reclassification of the Oceanic SAR11 Clade. Molecular Biology and
Evolution 29:599-615.
46. Wang H-C, Minh BQ, Susko E, Roger AJ. 2017. Modeling Site Heterogeneity with
Posterior Mean Site Frequency Profiles Accelerates Accurate Phylogenomic
Estimation. Syst Biol 67:216-235.
47. Lartillot N, Rodrigue N, Stubbs D, Richer J. 2013. PhyloBayes MPI: Phylogenetic
Reconstruction with Infinite Mixtures of Profiles in a Parallel Environment. Syst Biol
62:611-615.

48. Huerta-Cepas J, Forslund K, Coelho LP, Szklarczyk D, Jensen LJ, von Mering C, Bork P.
2017. Fast Genome-Wide Functional Annotation through Orthology Assignment by
eggNOG-Mapper. Molecular Biology and Evolution 34:2115-2122.
49. Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for
comparative genomics. Genome Biol 20:238.
50. Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017.
ModelFinder: fast model selection for accurate phylogenetic estimates. Nature
Methods 14:587-589.
51. Schön ME, Martijn J, Vosseberg J, Köstlbacher S, Ettema TJG. 2022. The evolutionary
origin of host association in the Rickettsiales. Nature Microbiology 7:1189-1199.
52. Zhang H, Yohe T, Huang L, Entwistle S, Wu P, Yang Z, Busk PK, Xu Y, Yin Y. 2018.
dbCAN2: a meta server for automated carbohydrate-active enzyme annotation.
Nucleic Acids Res 46:W95-W101.
53. Elbourne Liam DH, Tetu SG, Hassan KA, Paulsen IT. 2016. TransportDB 2.0: a database
for exploring membrane transporters in sequenced genomes from all domains of life.
Nucleic Acids Res 45:D320-D324.
54. Sauer DB, Wang D-N. 2019. Predicting the optimal growth temperatures of
prokaryotes using only genome derived features. Bioinformatics 35:3224-3231.
55. Musil M, Khan RT, Beier A, Stourac J, Konegger H, Damborsky J, Bednar D. 2020.
FireProtASR: A Web Server for Fully Automated Ancestral Sequence Reconstruction.
Brief Bioinform 22.
56. Yang Y, Zhao J, Zeng L, Vihinen M. 2022. ProTstab2 for Prediction of Protein Thermal
Stabilities. Int J Mol Sci 23:10798.
57. The UniProt Consortium. 2022. UniProt: the Universal Protein Knowledgebase in
2023. Nucleic Acids Res 51:D523-D531.
58. Vulcano F, Hribovšek P, Denny EO, Steen IH, Stokke R. 2023. Potential for
homoacetogenesis via the Wood–Ljungdahl pathway in Korarchaeia lineages from
marine hydrothermal vents. Environ Microbiol Rep n/a.
59. Louca S, Doebeli M. 2017. Efficient comparative phylogenetics on large trees.
Bioinformatics 34:1053-1055.
60. Xie R, Wang Y, Huang D, Hou J, Li L, Hu H, Zhao X, Wang F. 2022. Expanding Asgard
members in the domain of Archaea sheds new light on the origin of eukaryotes.
Science China Life Sciences 65:818-829.
61. Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. 2019. GTDB-Tk: a toolkit to classify
genomes with the Genome Taxonomy Database. Bioinformatics 36:1925-1927.
62. Pan J, Zhang X, Xu W, Liu Y, Liu L, Luo Z, Li M. 2023. Wood–Ljungdahl pathway found
in novel marine Korarchaeota groups illuminates their evolutionary history.
mSystems 0:e00305-23.
63. Konstantinidis KT, Rosselló-Móra R, Amann R. 2017. Uncultivated microbes in need of
their own taxonomy. The ISME Journal 11:2399-2406.
64. Hedlund BP, Chuvochina M, Hugenholtz P, Konstantinidis KT, Murray AE, Palmer M,
Parks DH, Probst AJ, Reysenbach A-L, Rodriguez-R LM, Rossello-Mora R, Sutcliffe IC,
Venter SN, Whitman WB. 2022. SeqCode: a nomenclatural code for prokaryotes
described from sequence data. Nature Microbiology 7:1702-1708.
65. Jeffroy O, Brinkmann H, Delsuc F, Philippe H. 2006. Phylogenomics: the beginning of
incongruence? Trends Genet 22:225-231.

66. Yutin N, Puigbò P, Koonin EV, Wolf YI. 2012. Phylogenomics of Prokaryotic Ribosomal
Proteins. PLoS One 7:e36972.

67. Qu Y-N, Rao Y-Z, Qi Y-L, Li Y-X, Li A, Palmer M, Hedlund BP, Shu W-S, Evans PN, Nie G-
X, Hua Z-S, Li W-J. 2023. Panguiarchaeum symbiosum, a potential hyperthermophilic
symbiont in the TACK superphylum. Cell Rep 42:112158.

68. Liu Y, Li M. 2022. The unstable evolutionary position of Korarchaeota and its
relationship with other TACK and Asgard archaea. mLife 1:218-222.

69. Miller-Coleman RL, Dodsworth JA, Ross CA, Shock EL, Williams AJ, Hartnett HE,
McDonald AI, Havig JR, Hedlund BP. 2012. Korarchaeota Diversity, Biogeography, and
Abundance in Yellowstone and Great Basin Hot Springs and Ecological Niche
Modeling Based on Machine Learning. PLoS One 7:e35964.

70. Auchtung TA, Shyndriayeva G, Cavanaugh CM. 2011. 16S rRNA phylogenetic analysis
and quantification of Korarchaeota indigenous to the hot springs of Kamchatka,
Russia. Extremophiles 15:105-116.

71. Blamey JM, Adams MWW. 1993. Purification and characterization of pyruvate
ferredoxin oxidoreductase from the hyperthermophilic archaeon Pyrococcus
furiosus. Biochim Biophys Acta 1161:19-27.

72. Frolov EN, Kublanov IV, Toshchakov SV, Lunev EA, Pimenov NV, Bonch-Osmolovskaya
EA, Lebedinsky AV, Chernyh NA. 2019. Form III RubisCO-mediated transaldolase
variant of the Calvin cycle in a chemolithoautotrophic bacterium. Proceedings of the
National Academy of Sciences 116:18638-18646.

73. Sato T, Atomi H, Imanaka T. 2007. Archaeal Type III RuBisCOs Function in a Pathway
for AMP Metabolism. Science 315:1003-1006.

74. Orita I, Sato T, Yurimoto H, Kato N, Atomi H, Imanaka T, Sakai Y. 2006. The Ribulose
Monophosphate Pathway Substitutes for the Missing Pentose Phosphate Pathway in
the Archaeon *Thermococcus kodakaraensis*. J Bacteriol 188:4698-4704.

75. Frank KL, Rogers DR, Olins HC, Vidoudez C, Girguis PR. 2013. Characterizing the
distribution and rates of microbial sulfate reduction at Middle Valley hydrothermal
vents. The ISME Journal 7:1391-1401.

76. Berghuis BA, Yu FB, Schulz F, Blainey PC, Woyke T, Quake SR. 2019. Hydrogenotrophic
methanogenesis in archaeal phylum Verstraetearchaeota reveals the shared ancestry
of all methanogens. Proceedings of the National Academy of Sciences 116:5037-
5044.

77. Appel L, Willistein M, Dahl C, Ermler U, Boll M. 2021. Functional diversity of
prokaryotic HdrA(BC) modules: Role in flavin-based electron bifurcation processes
and beyond. Biochim Biophys Acta 1862:148379.

78. Thauer RK, Kaster A-K, Goenrich M, Schick M, Hiromoto T, Shima S. 2010.
Hydrogenases from Methanogenic Archaea, Nickel, a Novel Cofactor, and H2 Storage.
Annu Rev Biochem 79:507-536.

79. Kikuchi A, Asai K. 1984. Reverse gyrase—a topoisomerase which introduces positive
superhelical turns into DNA. Nature 309:677-681.

80. Lipscomb GL, Hahn EM, Crowley AT, Adams MWW. 2017. Reverse gyrase is essential
for microbial growth at 95°C. Extremophiles 21:603-608.

81. Williams TA, Szöllősi GJ, Spang A, Foster PG, Heaps SE, Boussau B, Ettema TJG,
Embley TM. 2017. Integrative modeling of gene and genome evolution roots the
archaeal tree of life. Proceedings of the National Academy of Sciences 114:E4602-
E4611.

82. Akanuma S, Nakajima Y, Yokobori S-i, Kimura M, Nemoto N, Mase T, Miyazono K-i,
Tanokura M, Yamagishi A. 2013. Experimental evidence for the thermophilicity of
ancestral life. Proceedings of the National Academy of Sciences 110:11067-11072.

83. McKay LJ, Dlakić M, Fields MW, Delmont TO, Eren AM, Jay ZJ, Klingelsmith KB, Rusch
DB, Inskeep WP. 2019. Co-occurring genomic capacity for anaerobic methane and
dissimilatory sulfur metabolisms discovered in the Korarchaeota. Nature
Microbiology 4:614-622.

84. Amin K, Tranchimand S, Benvegnu T, Abdel-Razzak Z, Chamieh H. 2021. Glycoside
Hydrolases and Glycosyltransferases from Hyperthermophilic Archaea: Insights on
Their Characteristics and Applications in Biotechnology. Biomolecules 11:1557.

85. Saier MH. 2000. A Functional-Phylogenetic Classification System for Transmembrane
Solute Transporters. Microbiology and Molecular Biology Reviews 64:354-411.

86. Steens JA, Salazar CRP, Staals RHJ. 2022. The diverse arsenal of type III CRISPR–Cas-
associated CARF and SAVED effectors. Biochem Soc Trans 50:1353-1364.

87. Makarova KS, Wolf YI, Iranzo J, Shmakov SA, Alkhnbashi OS, Brouns SJJ, Charpentier
E, Cheng D, Haft DH, Horvath P, Moineau S, Mojica FJM, Scott D, Shah SA, Siksnys V,
Terns MP, Venclovas Č, White MF, Yakunin AF, Yan W, Zhang F, Garrett RA, Backofen
R, van der Oost J, Barrangou R, Koonin EV. 2020. Evolutionary classification of
CRISPR–Cas systems: a burst of class 2 and derived variants. Nat Rev Microbiol 18:67-
83.

88. Makarova KS, Wolf YI, Koonin EV. 2003. Potential genomic determinants of
hyperthermophily. Trends Genet 19:172-176.

89. Gudbergsdóttir SR, Menzel P, Krogh A, Young M, Peng X. 2016. Novel viral genomes
identified from six metagenomes reveal wide distribution of archaeal viruses and
high viral diversity in terrestrial hot springs. Environ Microbiol 18:863-874.

90. Goldfarb T, Sberro H, Weinstock E, Cohen O, Doron S, Charpak-Amikam Y, Afik S, Ofir
G, Sorek R. 2015. BREX is a novel phage resistance system widespread in microbial
genomes. The EMBO Journal 34:169-183.

1105

1106

Table Captions

Table 1. Overview of Bayesian phylogenetic analyses results for the five New Marker (NM54) alignments.

Figure Captions

Fig. 1. Phylogenetic tree corresponding to the Bayesian consensus tree (4 chains) reconstructed from the χ2-trimmed alignment (50% most heterogeneous sites removed; 7594 alignment positions) of 54 concatenated new marker proteins (NM54). *Ca*. Korarchaeota clades originating from hot springs and hydrothermal vents have a light and dark blue background shading, respectively. The two transitions between hydrothermal vents and hot springs are indicated with a star symbol. Scale bar indicates 0.1 substitutions per site.

Fig. 2. Overview of the Korarchaeota clade from Fig. 1. Internal nodes are annotated with the ancestor name/number code and the number of inferred coding sequences from the ALE analysis. Branches are annotated with bars representing the number of originations, transfers, duplications and losses. Bar heights in the legend correspond to 1000 such events each. Raw values can be found in Table S9. Terminal branches are labelled with the MAG name and accession number. Additional MAG statistics are given between square brackets: estimated genome size; GC content; number of CDS; estimated number of CDS.

Fig. 3. Ancestral gene content reconstruction of the Last Korarchaeota common ancestor (LKorCA). Genes were considered present if it was inferred by ALE with a copy number ≥ 0.3. For protein complexes, at least half the subunits had to be inferred as mentioned before.

Fig. 4. Ancestral gene content reconstruction of the Last *Ca*. Korarchaeum, *Ca*. Methanodesulfokora and *Ca*. Fumariihydrokora common ancestor (LKMFCA), which highlights the first transition from hydrothermal vents to hot springs. Genes were

1139 considered present if it was inferred by ALE with a copy number ≥ 0.3. For protein

1140 complexes, at least half the subunits had to be inferred as mentioned before.

1141

1142 **Fig. 5**. Ancestral gene content reconstruction of the Last *Ca*. Korarchaeum and *Ca*.

1143 Fumariihydrokora common ancestor (LKFCA), which highlights the second transition

1144 from hydrothermal vents to hot springs. Genes were considered present if it was

1145 inferred by ALE with a copy number ≥ 0.3. For protein complexes, at least half the

1146 subunits had to be inferred as mentioned before.

1147

1148 **Fig. 6**. Ancestral gene content reconstruction of the Last *Ca*. Methanodesulfokora

1149 common ancestor (LMCA). Genes were considered present if it was inferred by ALE

1150 with a copy number ≥ 0.3. For protein complexes, at least half the subunits had to be

1151 inferred as mentioned before.

1152

Table 1. Overview of Bayesian phylogenetic analyses for the five NM54 alignments.

	Untreated	SR4 recoded	Custom recoded	50% least heterogeneous sites (χ^2-trimmed)	50% fastest evolving sites trimmed
General alignment statistics					
No. of sites	15142	15142	15142	7594	7571
Model of evolution	LG+C60+F+G4	GTR+G4+C60SR4	LG+C60+F+G4	LG+C60+F+G4	LG+C60+F+G4
No. of informative sites	13862	11810	13427	6341	6291
Missing data (%)	8.80	8.80	8.80	8.59	8.51
PhyloBayes sampling statistics					
Chains used for optimal Maxdiff	4	2	2	4	2
Cycles	53599 - 54546	69504 - 69901	67722 - 68016	69911 - 70485	68356 - 70038
Burnin	10000	20000	20000	20000	20000
Step	30	30	30	30	30
Effective sampling size	>287	>15702	>16772	>3399	>1510
Optimal Maxdiff	1	0.049	0.217	1	0.053
Maxdiff 4 chains	1	0.397	1	1	1
Posterior predictive tests					
Maximum squared heterogeneity (p-value)	0	0	0	0.36 - 0.41	0
Maximum squared heterogeneity (z-score)	185.41 - 195.53	45.74 - 49.51	36.98 - 39.85	0.06 - 0.13	33.33 - 35.38
Mean squared heterogeneity (p-value)	0	0	0	1	0
Mean squared heterogeneity (z-score)	464.51 - 502.73	97.00 - 100.48	131.87 - 140.52	-7.38 - -7.15	73.08 - 76.04
Mean diversity per site (p-value)	0	0.47 - 0.50	0.04 - 0.06	0	0
Mean diversity per site (z-score)	13.32 - 13.87	0.03 - 0.09	1.56 - 1.68	7.87 – 8.01	6.81 - 7.20

Fig. 1. Phylogenetic tree corresponding to the Bayesian consensus tree (4 chains) reconstructed from the χ2-trimmed alignment (50% most heterogeneous sites removed; 7594 alignment positions) of 54 concatenated new marker proteins (NM54). *Ca.* Korarchaeota clades originating from hot springs and hydrothermal vents have a light and dark blue background shading, respectively. The two

transitions between hydrothermal vents and hot springs are indicated with a star symbol. Scale bar indicates 0.1 substitutions per site.

Fig. 2. Overview of the Korarchaeota clade from Fig. 1. Internal nodes are annotated with the ancestor name/number code and the number of inferred coding sequences from the ALE analysis. Branches are annotated with bars representing the number of originations, transfers, duplications and losses. Bar heights in the legend correspond to 1000 such events each. Raw values can be found in Table S9. Terminal branches

are labelled with the MAG name and accession number. Additional MAG statistics are given between square brackets: estimated genome size; GC content; number of CDS; estimated number of CDS.

Fig. 3. Ancestral gene content reconstruction of the Last Korarchaeota common ancestor (LKorCA). Genes were considered present if it was inferred by ALE with a copy number ≥ 0.3. For protein complexes, at least half the subunits had to be inferred as mentioned before.

Fig. 4. Ancestral gene content reconstruction of the Last *Ca*. Korarchaeum, *Ca*. Methanodesulfokora and *Ca*. Fumariihydrokora common ancestor (LKMFCA), which highlights the first transition from hydrothermal vents to hot springs. Genes were considered present if it was inferred by ALE with a copy number ≥ 0.3. For protein complexes, at least half the subunits had to be inferred as mentioned before.

Fig. 5. Ancestral gene content reconstruction of the Last *Ca*. Korarchaeum and *Ca*. Fumariihydrokora common ancestor (LKFCA), which highlights the second transition from hydrothermal vents to hot springs. Genes were considered present if it was inferred by ALE with a copy number ≥ 0.3. For protein complexes, at least half the subunits had to be inferred as mentioned before.

Fig. 6. Ancestral gene content reconstruction of the Last *Ca*. Methanodesulfokora common ancestor (LMCA). Genes were considered present if it was inferred by ALE with a copy number ≥ 0.3. For protein complexes, at least half the subunits had to be inferred as mentioned before.