Improvisation and Live Accompaniment Increase Motor Response and Reward During a Music Playing Task

Anna Palumbo[1,2,3,4], Karleigh Groves[2,3,4], Eva Luna Muñoz Vidal[1,2,3,4], Alan Turry[5,6], Preeti Raghavan[7], Heidi Schambra[8], Gerald Voelbel[9], Pablo Ripollés[2,3,4]

Abstract

Music is one of the most important sources of abstract reward in humans. While music is often combined with exercise to improve performance and to upregulate mood, the relationship between music-induced reward and motor output is poorly understood. Here, we study music reward and motor output at the same time by capitalizing on music playing. Specifically, we investigate the effects of music improvisation and live accompaniment on motor, autonomic, and affective responses. 30 adults performed a rhythm tapping task while (i) improvising or maintaining the beat and (ii) with live or recorded accompaniment. Motor response was characterized by acceleration and frequency of hand movements (accelerometry), wrist flexor and extensor muscle activation (electromyography), and the number of beats played. Autonomic arousal was measured by tonic response of electrodermal activity (EDA) and heart rate (HR). Affective responses were measured by a 12-item Likert scale. The combination of improvisation and live accompaniment, as compared to all other conditions, significantly increased acceleration of hand movements and muscle activation, as well as participant reports of enjoyment during music-playing. Improvisation, regardless of type of accompaniment, increased the number of beats played and autonomic arousal (including tonic EDA responses and several measures of HR), as well as participant reports of challenge. Importantly, increased motor response was associated with increased enjoyment during music improvisation only and not while participants were maintaining the beat. The increased motor responses achieved with improvisation and live accompaniment have important implications for enhancing dose of movement during music-based interventions for stroke rehabilitation.

Significance Statement: Music has been shown to provide a rewarding stimulus that improves motor performance and learning. However, the relationship between music reward and motor output is poorly understood. Here, we show that music improvisation with live accompaniment increased the intensity and frequency of movement and the level of muscle activation, and that this increase in motor response was associated with increased enjoyment only when improvising. These findings are important for developing interventions that target improved motor performance and learning in exercise and physical rehabilitation.

[1] Rehabilitation Sciences Program, Steinhardt School of Culture, Education, and Human Development, New York University, New York, NY 10003

[2] Department of Psychology, New York University, New York, NY 10003

[3] Music and Audio Research Lab, , New York University, New York, NY 10003

[4] Center for Language, Music, and Emotion (CLaME), New York University, New York, NY 10003

[5] Department of Music and Performing Arts Professions, Steinhardt School of Culture, Education, and Human Development, New York University, New York, NY 10003

[6] Nordoff Robbins Center for Music Therapy, New York University, New York, NY 10003

[7] Department of Physical Medicine and Rehabilitation and Neurology, Johns Hopkins University School of Medicine, Baltimore, MD 21287

[8] New York University Grossman School of Medicine, New York, NY 10016

[9] Department of Occupational Therapy, Steinhardt School of Culture, Education, and Human Development, New York University, New York, NY 10003

Corresponding Authors: Anna Palumbo (ap89@nyu.edu), Pablo Ripollés (pr82@nyu.edu)

Conflict of Interest Statement: The authors declare no competing financial interests.

Acknowledgements: The authors would like to acknowledge Robert Codio, MT-BC, who composed the music for this study and provided the music accompaniment for the task.

Introduction

Reward plays an important role in motor performance and learning (Wulf and Lewthwaite, 2016). Monetary reward during motor tasks increases force production, reduces reaction time, improves recall of motor skills, increases neural activation in reward-related areas and modulates corticospinal excitability (Pessiglione et al., 2007; Widmer et al., 2016; Bundt et al., 2019). Dopaminergic signaling between reward and motor brain areas optimizes motor skill learning in non-human animals (Molina-Luna et al., 2009; Hosp et al., 2011). In everyday life, music provides an abstract reward that is frequently combined with exercise and athletic training to improve mood and performance (Laukka and Quick, 2013; Hallett and Lamont, 2017). Listening to music improves affect, endurance, speed, power output, movement kinematics, and cardiovascular efficiency (Simpson and Karageorghis, 2006; Karageorghis et al., 2009; Terry et al., 2012; Belkhir et al., 2019; Stork et al., 2019; Terry et al., 2020; Bonassi et al., 2023). Music listening activates reward-related brain areas, and musical pleasure is modulated by dopamine release (Blood and Zatorre, 2001; Menon and Levitin, 2005; Salimpoor et al., 2011; Ferreri et al., 2019). Musical groove also activates reward and motor-related brain areas (Matthews et al., 2020). Music playing (i.e., using musical instruments) has been utilized in motor rehabilitation to improve recovery following acquired brain injury, leading to increased motor function, reduced depression and improved positive affect, and increased activation in the affected sensorimotor cortex (Grau-Sanchez et al., 2013; Raghavan et al., 2016; Ripolles et al., 2016; Magee et al., 2017; Palumbo et al., 2022). Critically, music playing supports higher levels of motor recovery among individuals with greater sensitivity to musical reward (Grau-Sanchez et al., 2018). This suggests that increasing reward during music playing improves motor outcomes, but the mechanisms supporting these effects remain unclear. Here we capitalize on music playing to study the putative link between movement and reward. We examine two avenues for enhancing reward and motor output during music playing: music improvisation and live accompaniment.

Music improvisation involves creating or adapting music in the moment, providing choice about what to play (Turry, 2009; Loui, 2018). Choice enhances motor performance and learning, even when the choice is unrelated to the task, and increases motivation and activation in reward-related brain areas (Wulf et al., 2014; Lewthwaite et al., 2015; Murayama et al., 2015; Chua et al., 2018; An et al., 2020). In this vein, choice in combining sounds increases enjoyment and interest (Baur et al., 2018), and music improvisation, compared to playing a pre-defined set of notes, increases activation in the sensorimotor cortex and in brain regions related to executive functions (Beaty, 2015; de Aquino et al., 2019; Vuust et al., 2022). Creativity in music improvisation correlates with grey matter in the hippocampus (Arkin et al., 2017). These findings indicate the potential for music improvisation to influence cognition (e.g., attention, learning), motor responses, and reward during music-playing.

Live accompaniment creates a social experience during music playing. Positive social interactions contribute to reward (Krach et al., 2010), and cooperative learning of motor tasks increase intrinsic motivation, positive affect, self-efficacy, and motor learning as compared to competitive or non-interactive learning (Kaefer and Chiviacowsky, 2022). Live music also modulates heart rate, self-reports of reduced anxiety and improved vigor, and overall mood as compared to recorded music (Bailey, 1983; Bush et al., 2021; but see Belfi et al., 2021). Together, these findings indicate that live music may reduce feelings of anxiety and their associated physiological responses and that social interactions during live music playing may improve reward and motor learning.

Here, we compare the effects of improvisation and live accompaniment in a music-playing task in healthy participants. We predict that improvisation combined with live accompaniment will produce the highest levels of motor activity, positive affect, and autonomic arousal. We also predict that increased motor responses will correlate with increased self-reports of pleasure during music-playing, thus supporting the proposed link between reward and motor performance.

Methods

Participants

Thirty healthy adults were recruited from the New York University community, ages 18-58 (M = 22.9, SD = 8.11), with 19 females. Sample size was based on previous studies with 13-21 participants demonstrating that rhythmic improvisation increases the number of notes played and the activation of motor planning and executive brain areas (Berkowitz and Ansari, 2008; de Aquino et al., 2019). All participants were either enrolled in university (n = 24) or had already earned an undergraduate or graduate degree and were currently employed (n = 4). Participants self-reported their race as Asian (n = 14), White/Caucasian (n=8), Black/African American (n=2), or Mixed (n=2), with 3 participants not reporting their race. Twenty seven percent (n=8) of participants reported Hispanic ethnicity. Inclusion criteria included right-handed with no history of brain injury. Exclusion criteria included musical anhedonia, as indicated by a score of < 65 on the Barcelona Music Reward Questionnaire (Mas-Herrero et al., 2013), or current employment as a professional musician. The participants gave written informed consent as per the institutional review board at New York University and were paid for their participation.

Music Playing Task

Participants played a drum along with piano accompaniment for four one-minute songs, written specifically for this study. These included a March, Waltz, Latin style, and Rock song. Prior to playing, participants listened to each song, during which time they were instructed to remain as still as possible. After listening, participants were provided an opportunity to play along with recorded piano accompaniment and find the beat of the music that felt most natural to them. After this familiarization period, participants were instructed to play the drum along with the piano accompaniment by either (1) maintaining the beat without changing as much as possible or (2) improvising a beat, changing the beat as they desired for personal expression. When participants maintained the beat, they were not provided with a beat to imitate, but instead were asked to identify and use the beat that felt most natural to them as instructed during the familiarization phase. In this way, the "maintain the beat" and "improvisation" conditions both represented personal preference, but only the "maintain the beat" condition was constrained to stay the same throughout the song. Piano accompaniment was either (1) played from a recording or (2) performed live by the same music therapist who made the recordings. The music therapist remained in the room in all conditions, regardless of whether a recording was being played. Piano and drums were played on MIDI devices, including the Roland SPD-One Percussion MIDI drum pad, tuned to the sound of snare drum, and a Roland FP-60x digital keyboard. Both MIDI inputs were recorded to individual tracks in Logic Pro, with real time sound output to a Roland KC-400 150 watt 12-inch keyboard amp. MIDI toolbox 1.1 was used to calculate the number of beats played on the MIDI drum pad during each condition (Eerola and Toiviainen, 2004). Electromyography, accelerometry, electrodermal activity, and electrocardiogram data was collected from all participants while they were playing using a MP160 data acquisition module (BIOPAC Systems Inc.) and AcqKnowledge software. All signal processing for this data was for completed in MATLAB R2022b.

Behavioral Correlates of the Music Playing Experience

After each music-playing task, participants completed a survey with twelve questions that were answered using a 5-point Likert scale and were related to their affective and physical response, as well as their perception of the music (Table 1). Principal component analysis (PCA) was used to identify the main components of participant responses across these questions. The number of main components were determined using parallel analysis. Participant loadings on these main components, rather than responses to each Likert scale question, were analyzed in relation to participant experience of conditions (see Results).

Table 1: Experience of Music Survey Items

How much pleasure did you experience?
How much difficulty did you experience?
How much fatigue did you experience?
How much did you feel in synchrony with the music?
How familiar was the music?
How motivated did you feel to play?
How surprised were you by the music?
How activated did you feel?
How absorbed did you become?
How creative did you feel?
How much momentum did the music provide for you?
How active were you?

Using a Likert Scale of 1 ("none/not at all") to 5 ("a lot/very much") participants were asked to rate their experience after each music playing task.

Electromyography

Electromyography (EMG) was measured via surface electrodes placed over the wrist flexor carpi radialis and extensor carpi radialis longus on the arm used to play the drum. Prior to electrode placement, the skin was cleaned with alcohol. EMG signals were filtered with a 10-400 Hz band-pass filter, constructed by convolving a 2nd order high pass Butterworth filter and 3rd order low pass Butterworth filter (Aluru et al., 2014). The signals were then rectified by taking the root mean square of 10ms windows of the signal. Signals were normalized by dividing by the participant maximum voluntary contraction (MVC; Aluru et al., 2014). The MVC was measured by providing resistance to movement at about 45^o of rotation through wrist extension and flexion, during which participants were asked to exert the maximum possible force against the resistance. MVC signals were filtered and rectified as described above, and the final MVC value was calculated as the average amplitude of the maximum 1 second region of the signal.

To ensure background noise did not influence the overall EMG average, movement related peaks in normalized EMG amplitude were identified by alignment with peaks in the Z component of the accelerometry signal (see below) using custom code. The lowest 10% of peaks in EMG amplitude were dropped to eliminate noise, and plots were visually inspected to ensure proper alignment between EMG and accelerometry signals. The average of movement related peaks in EMG amplitude was calculated. Since participants played the drum with slightly different arm positions, the results present findings for the average wrist flexor and extensor activation, rather than considering each muscle separately. EMG signals were recorded at a sampling rate of 2000 samples/second.

Accelerometry

A triaxial accelerometer was secured to the back of the hand playing the drum using a Velcro strap, with placement approximately under the metacarpophalangeal joints and between the third and fourth fingers. Accelerometry signals were acquired at a sampling rate of 2000 samples/second, except for the first 14 participants, for whom the y component was sampled at 250 samples/sec due to a technical problem. To correct for this, all other accelerometry signals were downsampled to 250 Hz. A 4th order Butterworth high pass filter with cut off at 0.5 Hz was then used to remove the gravitational component (Hurd et al., 2013). Leading and trailing samples were dropped due to filter and resampling artifacts, with an overall sample loss of less than 10% of the signal. The x, y, and z components of the resulting accelerometer signal were then transformed into magnitude of acceleration taking the square root of the sum of the squares of

each component. The total acceleration was calculated as the sum of the magnitude of acceleration over the entire signal (Hurd et al., 2013).

Electrodermal activity
Electrodermal activity (EDA) electrodes were placed on the index and middle finger of the hand not playing the drum, with the hand resting on a table. We instructed participants to move the hand as little as possible to avoid motion artifacts. Tonic response of EDA was calculated as the mean of the signal for each trial (Braithwaite et al., 2013). Music-playing responses were normalized for each participant by subtracting the tonic response to the listen-only condition from each trial. Five participants were dropped from the EDA analysis due to hyperhidrosis that caused ceiling effects in the EDA signal (n = 4) or lack of sufficient signal (n = 1).

Electrocardiography
Electrocardiography (ECG) was recorded using 3 electrodes placed on the torso. Raw ECG signals were detrended, and heart beats were identified as the R wave maxima of the QRS complexes. Noise was removed using custom code, and ECG signals were visually inspected to ensure that QRS complexes were accurately identified. Three participants were dropped from ECG analyses due to insufficient signal-to-noise ratio. Beats per minute (BPM) were calculated as the sampling rate (in samples/sec) divided by the average samples per beat, multiplied by 60 (sec/min; Gorr, 2023). Heart rate variability (HRV) was calculated as the root mean square of successive differences (RMSSD) between R wave maxima (Shaffer et al., 2016; Shaffer and Ginsberg, 2017). ECG signals were recorded at 2000 samples/sec.

Participant Characteristics
Baseline screening included demographic information - including age, sex, race, ethnicity, and education level. Baseline screening also measured factors known to play a role in response to music-playing tasks, including music reward and training. Sensitivity to music reward was measured via the Barcelona music reward questionnaire (BMRQ), a reliable five-factor scale measuring sensitivity to musical reward (Mas-Herrero et al., 2013). Music training was measured via the music training subscale of the Goldsmith Music Sophistication Index (Gold-MSI), a self-report scale with strong internal consistency (Cronbach's alpha = 0.903;Müllensiefen et al., 2014).

Experimental Design and Statistical Analysis
Using a crossover design, all participants completed all four combinations of music playing and accompaniment conditions (Figure 1) with each of the four songs, for a total of 16 music playing trials. Participants were randomized to one of four condition orders (See Fig 1.1 in Extended Materials at the end of the document). The same condition order was repeated for each of the four songs played by each participant. For each outcome, linear mixed models were calculated with fixed effects including the music playing condition (Maintain the Beat vs. Improvise), the music accompaniment condition (Recorded vs. Live), and their interaction, with control variables for musical training, sensitivity to music reward, and song, and a random intercept for subject (Bates et al., 2015). For EDA and ECG, total acceleration was additionally included in the model as a control, to eliminate motion as an explanation for increased autonomic response.

Figure 1: **Experimental Design**
Using a balanced 2x2 design, each participant completed all four combinations of the music playing

Post-hoc comparisons of marginal effects were adjusted for multiple effects using the Tukey method (Lenth et al., 2019).

To evaluate the association between reward-related behavioral measures and motor responses (accelerometry, EMG, and beats played), we again used linear mixed models (Bates et al., 2015). As a measure of reward we used the first principal component of participant self-reports related to the music playing experience. This principal component includes positive loadings for pleasure, as well as other responses related to enjoyment (see results), and therefore more fully accounts for participant responses related to reward as compared to the single rating of pleasure. Fixed predictors included motor response (we computed 3 different models for our 3 different measures: accelerometry, EMG, and number of beats played), music playing condition (improvise vs. maintain the beat), music accompaniment condition (live vs. recorded), and their three-way interaction, as well as control variables for musical training, music reward, and song, with a random intercept for subject. The three-way interaction in this model accounts for potential modulation of the association between motor response and enjoyment by the four music tasks. Post-hoc comparisons of linear trends were evaluated by averaging over other factor level predictors in the model and adjusted for multiple effects using the Tukey method (Lenth et al., 2019).

For all analyses of linear mixed models, Wald Chi Square tests were used to evaluate significance of main effects and interactions. All analyses were done in R (packages: psych, factoextra, lme4, foreign, aod, car, emmeans, emtrends, ggplot2, ggeffects). Custom code for processing of physiological data is available upon request to the first author.

Results

Participant Musical Background
Participant scores on the BMRQ (M = 84.93, SD = 9.08) were in the range of the general population (Mas-Herrero et al., 2013; M = 78.42, SD = 10.47, n = 857). Similarly, participants scores on the Gold-MSI musical training subscale (M = 25.40, SD = 10.82) were in the range of the general population (Müllensiefen et al., 2014; M = 26.52, SD = 11.44, n = 147,633). These findings indicate our sample approximates the general population in relationship to musical reward and training.

Behavioral Correlates of the Music Playing Experience
PCA of responses revealed two main components of music playing experience (Figure 2). The first component (PC1) accounted for 38.05% of the variance in the survey responses and was termed "enjoyment" to summarize positive loadings for participant experiences of feeling activated, motivated, creative, active, absorbed, and for the ratings of pleasure, and momentum. The second component (PC2) accounted for 18.82% of the variance in the survey responses and was termed "challenge" to summarize positive loadings of participant experiences of difficulty, fatigue, and surprise, and negative loadings of participant perception of the music as familiar and in synchrony with their playing.

With Enjoyment as the outcome (see Figure 3), there were significant main effects of music playing condition (Improvisation vs. Maintaining the Beat; χ^2 (1) = 15.18, p = 0.0001) and accompaniment condition (Live vs. Recorded; χ^2 (1) = 7.73, p = 0.005), as well as for song (χ^2 (3) = 33.92, p < 0.00001). In addition, greater sensitivity to musical reward was related to greater Enjoyment (χ^2 (1) = 4.51, p = 0.034). There was also a significant interaction between playing

Figure 2: **Principal Component Analysis of Music Experience Survey**
Biplot of principal components derived from the 12-item music experience survey. The first principal component includes responses related to enjoyment, accounting for 38.41% of the variance in survey responses. The second principal component includes responses related to challenge, and accounted for 19.87% of the variance in participant responses.

condition and accompaniment condition on Enjoyment (Figure 3; χ^2 (1) = 14.53, p = 0.0001). Pairwise comparison for song, corrected for multiple tests, revealed less Enjoyment for the March as compared to the Rock (t(444) = -5.33, p < 0.0001) and Latin (t(444) = -3.20, p = 0.0006) compositions, but not the Waltz (t(444) = -1.57, p = 0.399). Similarly the Waltz was associated with less Enjoyment than the Rock composition (t(444) = -3.76, p = 0.0011) with a trend for less Enjoyment compared to the Latin composition (t(444) = -2.35, p = 0.088). There were no differences in Enjoyment for the Latin and Rock compositions (t(444) = 1.41, p = 0.495). Pairwise comparisons for music conditions, averaged over the levels of song and corrected for multiple tests, revealed improvisation increased enjoyment, as compared to maintaining the beat, for both live accompaniment (t(444) = 9.29, p < 0.00001) and recorded accompaniment (t(444) = 3.90, p = 0.0007). Additionally, live accompaniment increased enjoyment, as compared to recorded accompaniment, for both improvisation (t(444) = 8.17, p < 0.00001) and maintaining the beat (t(444) = -2.78, p = 0.03). Finally, the combination of live accompaniment and improvisation produced the highest levels of enjoyment compared to each of the other conditions, including improvising with a recording (reported above), maintaining the beat with live accompaniment (t(444) = -9.29, p<0.00001), and maintaining the beat with a recording (t(444) = -12.07, p<0.00001).

With Challenge as the outcome (see Figure 3) there was a significant main effect for music playing condition (χ^2 (1) = 50.12, p < 0.00001) and song (χ^2 (3) = 35.42, p < 0.00001). There was no significant interaction between music playing condition and accompaniment condition (χ^2 (1) = 0.011, p = 0.916) or a significant main effect of accompaniment (χ^2 (1) = 0.897, p = 0.344), music training (χ^2 (1) = 0.639, p = 0.424), or sensitivity to musical reward (χ^2 (1) = 1.16, p = 0.281). Pairwise comparisons for song, corrected for multiple tests, revealed greater Challenge for the Latin composition as compared to the March (t(444) = 4.72, p = 0.00002), Waltz (t(444) = 5.39, p = 0.0000007), and Rock (t(444) = 4.14, p = 0.00024) compositions. Pairwise comparisons for music playing condition, averaged over the levels of song and corrected for multiple

Figure 3: **Behavioral Correlates of Music Playing Experience**
There was a significant interaction between music playing condition and music accompaniment condition on enjoyment, such that the combination of improvisation and live accompaniment produced the greatest enjoyment compared to the other three conditions. There was a main effect of music playing condition on challenge, such that challenge was greater during improvisation, as compared to maintaining the beat. * p < 0.05 after correction for multiple tests.

tests, revealed that improvisation produced higher levels of challenge, as compared to maintaining the beat, for both recorded accompaniment (t(444) = -7.08, p < 0.00001) and live piano accompaniment (Figure 3; t(444) = -7.23, p < 0.00001).

Motor Response to Music Conditions
For total acceleration (see Figure 4), there were no main effects for music playing condition (χ^2(1) = 0.766, p = 0.382) or accompaniment condition (χ^2(1) = 0.0005, p = 0.982), music training (χ^2(1) = 0.727, p = 0.394), or sensitivity to musical reward (χ^2(1) = 1.16, p = 0.281), although there was a significant main effect for song (χ^2 (3) = 23.99, p = 0.00003). Importantly, there was a significant interaction between music playing condition and accompaniment condition on total acceleration (Figure 4; χ^2 (1) = 7.54, p = 0.006). Pairwise comparisons for song, controlling for multiple tests, revealed lower total acceleration for the March as compared to the Rock (t(432) = -3.98, p = 0.0005), Latin (t(432) = -4.41, p = 0.0001, and Waltz (t(432) = -3.18, p = 0.0087) compositions. There were no significant differences in total acceleration between the Rock and Latin (t(432) = -0.376, p = 0.982) or Waltz compositions (t(432) = 750, p = 0.877), or between the Latin and Waltz compositions (t(432) = 1.125, p = 0.674). Pairwise comparisons for music playing condition, averaged over the levels of song and controlling for multiple tests, revealed that the combination of improvised playing and live accompaniment produced greater total acceleration than the other three conditions, including maintaining the beat with a recording (t(432) = 4.78, p = 0.00001), improvising with a recording (t(432) = 3.90, p = 0.006), and maintaining the beat with live accompaniment (t(432) = 4.76, p = 0.00002).

For mean EMG activation (see Figure 4), there were main effects of music playing condition ($\chi^2(1)$ = 18.11, p = 0.00002) and song ($\chi^2(3)$ = 82.60, p < 0.00001). There were no significant main effects of accompaniment ($\chi^2(1)$ = 0.13, p = 0.720), musical training ($\chi^2(1)$ = 0.30, p = 0.581), or sensitivity to musical reward ($\chi^2(1)$ = 0.05, p = 0.816). Importantly, there was also an interaction effect between music playing condition and accompaniment condition on mean EMG activation (Figure 4; $\chi^2(1)$ = 6.12, p = 0.013). Pairwise comparisons for song, controlling for multiple tests, revealed less EMG activation for the March as compared to the Rock (t(372.08) = -6.16, p < 0.00001) and Latin (t(372.00) = -6.35, p < 0.00001) compositions, but the Waltz (t(372.12) = 0.461, p = 0.967). Similarly, there was lower EMG activation for the Waltz as compared to the Rock (t(372.04) = -6.50, p < 0.00001) and Latin (t(372.11) = -6.63, p < 0.00001) compositions. There were no significant differences in EMG activation between the Rock and Latin compositions (t(372.08) = 0.097, p = 0.9997). Pairwise comparisons of music conditions, averaged over the levels of song and controlling for multiple comparisons, showed that improvisation produced greater mean EMG activation compared to maintaining the beat for both live accompaniment (t(372) = -7.76, p < 0.00001) and recorded accompaniment (t(372) = 4.26, p = 0.0002). In addition, the combination of improvised playing and live accompaniment produced greater mean EMG activation compared to the other three conditions, including maintaining the beat with a recording (t(372) = 8.11, p < 0.00001), improvising with a recording (t(372) = 3.86, p = 0.0008), and maintaining the beat with live accompaniment (t(372) = 7.76, p < 0.00001).

Figure 4: **Motor Response to Music Conditions**
There was a significant interaction between music playing condition and music accompaniment condition on total acceleration and mean EMG activation, such that the combination of improvisation and live accompaniment produced the greatest motor response. There was a main effect of music playing condition on the number of beats played, such that a greater number of beats were produced during improvisation. * p < 0.05 after correction for multiple tests.

For the number of beats played (see Figure 4), there were main effects of music playing condition ($\chi^2(1)$ = 23.62, p < 0.00001) and song ($\chi^2(3)$ = 61.13, p = 0.00001). There were no significant main effects of accompaniment ($\chi^2(1)$ = 0.20, p = 0.652), musical training ($\chi^2(1)$ = 0.01, p = 0.909), nor a significant interaction between music condition and accompaniment ($\chi^2(1)$ = 1.43, p =

0.232). There was a trend towards a main effect of sensitivity to musical reward on the number of beats played and sensitivity to musical reward ($\chi^2(1)$ = 3.64, p = 0.056). Pairwise comparisons for song, controlling for multiple tests, revealed fewer number of beats played for the March, as compared to the Rock (t(444) = -5.47, p = 0.0000005), Latin (t(444) = -7.39, p < 0.0000001), and Waltz (t(444) = -5.53, p = 0.0000003) compositions. There were no significant differences in the number of beats played between the Rock and Latin (t(444) = -1.92, p = 0.220) or Waltz (t(444) = -0.066, p = 0.9998) compositions, or between the Latin and Waltz compositions (t(444) = 1.86, p = 0.249). Pairwise comparisons in music condition, averaged over the level of song and controlling for multiple tests, revealed that improvisation produced a greater number of beats played compared to maintaining the beat, for both the live accompaniment (t(444) = 4.86, p < 0.00001) and recorded accompaniment (Figure 4; t(444) = 6.55, p < 0.00001).

Autonomic Response to Music Conditions

Figure 5: **Autonomic Response to Music Conditions**
There was a main effect of music playing condition on autonomic response, such that improvisation led to increased autonomic arousal as measured by increased tonic EDA and BPM, and decreased HRV. Pairwise comparisons revealed higher EDA and BPM for improvisation during both recorded and live accompaniment. For HRV, the difference between improvisation and maintain the beat conditions were only significant in the recorded accompaniment condition. * p < 0.05 after correction for multiple tests.

For tonic EDA (see Figure 5), there was a main effect of music playing condition (χ^2 (1) = 5.82, p = 0.016), while controlling for the effects of movement by including total acceleration in the model. There was also a significant main effect of total acceleration (χ^2 (1) = 5.01, p = 0.026) and song (χ^2 (3) = 8.33, p = 0.040). There were no main effects of accompaniment condition ($\chi^2(1)$ = 0.05, p = 0.830), musical training ($\chi^2(1)$ = 0.53, p = 0.467), or sensitivity to musical reward ($\chi^2(1)$ = 0.03, p = 0.874), or any significant interaction between music playing and accompaniment conditions ($\chi^2(1)$ = 0.66, p = 0.418). Pairwise comparisons for song, controlling for multiple testing, revealed lower tonic EDA for the March as compared to the Latin composition (t(361.64) = -2.64, p = 0.043),

but not the Rock (t(363.44) = -0.93, p = 0.788) or Waltz (t(361.42) = -2.04, p = 0.174) compositions. There were no significant differences between the Rock and Latin (t(357.14) = -1.70, p = 0.325) or Waltz (t(357.02) = -1.12, p = 0.675) compositions, or between the Latin and Waltz compositions (t(357.77) = 0.54, p = 0.948). Pairwise comparisons for music conditions, averaged over the levels of song and controlling for multiple testing, revealed that improvisation produced greater tonic EDA, indicating higher autonomic arousal, compared to maintaining the beat for both live accompaniment (t(361.18) = 3.51, p = 0.0005) and recorded accompaniment (t(356.15) = 2.41, p = 0.016).

For BPM (see Figure 5), there was a main effect of music playing condition (χ^2 (1) = 21.34, p < 0.00001), while controlling for the effects of movement by including total acceleration in the model. There were also significant main effects of total acceleration (χ^2 (1) = 13.45, p = 0.0002) and musical training (χ^2 (1) = 4.35, p = 0.037). There were no main effects of accompaniment condition (χ^2(1) = 1.39, p = 0.238), song (χ^2 (3) = 7.48, p = 0.058), or sensitivity to musical reward (χ^2(1) = 0.42, p = 0.519), or any significant interaction between music playing and accompaniment conditions (χ^2(1) = 0.15, p = 0.701). Pairwise comparisons for music condition, averaged over the level of song and controlling for multiple testing, revealed that improvisation produced greater BPM, indicating higher autonomic arousal, compared to maintaining the beat for both live accompaniment (t(398.65) = 3.95, p < 0.00001) and recorded accompaniment (t(397.07) = 4.62, p < 0.00001).

For HRV (see Figure 5), there was a main effect of music playing condition (χ^2 (1) = 4.66, p = 0.041), while controlling for the effects of movement by including total acceleration in the model. There was also a significant main effect of total acceleration (χ^2 (1) = 10.16, p = 0.0014). There were no main effects of accompaniment condition (χ^2(1) = 0.20, p = 0.655), song (χ^2(3) = 3.91, p = 0.271), musical training (χ^2(1) = 3.67, p = 0.055), or sensitivity to musical reward (χ^2(1) = 0.002, p = 0.961), or any significant interaction between music playing and accompaniment conditions (χ^2(1) = 0.66, p = 0.418). Pairwise comparisons for music condition, averaged over the levels of song and controlling for multiple testing, revealed that improvisation produced lower HR, indicating parasympathetic withdrawal (Mackersie and Calderon-Moultrie, 2016), compared to maintaining the beat for recorded accompaniment (t(397.14) = 2.16, p = 0.031). Although HRV was also lower during improvised playing for live accompaniment, the difference from maintaining the beat was not significant in this condition (t(400.12) = 0.976, p = 0.330).

Association Between Enjoyment and Motor Responses
The association between Enjoyment and each of the motor responses (total acceleration, mean EMG activation, and number of beats played) was evaluated by adding motor response to the linear mixed model predicting Enjoyment. The main effect of motor response was considered, as well as the three-way interaction between motor response, music playing condition, and accompaniment condition, as well as the control variables for song, musical training, and sensitivity to musical reward, and a random intercept for subject.

There was no significant main effect of total acceleration on Enjoyment (χ^2 (1) = 0.04, p = 0.841) and no significant triple interaction between total acceleration, music playing condition, and accompaniment on Enjoyment (χ^2 (1) = 0.0004, p = 0.983). However, there was a significant interaction effect between total acceleration and music playing condition on Enjoyment (Figure 6; χ^2 (1) = 12.15, p = 0.0005). There was also a significant main effect of song (χ^2(3) = 28.71, p = 0.000003) and sensitivity to musical reward (χ^2(1) = 3.92, p = 0.048). There were no significant interaction effects for total acceleration and accompaniment (χ^2 (1) = 0.098, p = 0.755) or music playing condition and accompaniment (χ^2 (1) = 3.05, p = 0.081, and no main effects of musical

training (χ^2 (1) = 0.533, p = 0.465), accompaniment (χ^2 (1) = 3.46, p = 0.063), or music playing condition (χ^2 (1) = 3.34, p = 0.841). Notably, the previously significant interaction between music playing condition and accompaniment and the main effect of improvisation on Enjoyment became non-significant after including total acceleration in the model. Pairwise comparisons of song on Enjoyment have been reported above and maintained the same relationships. Simple effects for the association between total acceleration and Enjoyment revealed a positive relationship between total acceleration and Enjoyment during the improvisation condition, using the 95% confidence interval to indicate significance (estimated slope = 0.302; 95% CI = 0.166, 0.438), but not for the maintain the beat condition (estimated slope = -0.003; 95% CI = -0.153, 0.148). The slopes describing the relationship between Enjoyment and total acceleration were significantly different between improvisation and maintain the beat conditions (estimate = 0.305, t(453) = 4.60, p < 0.0001).

Figure 6: **Relationship between reward and motor outputs**.
There was a significant interaction between total acceleration and music playing condition on enjoyment, such that enjoyment was positively associated with total acceleration for the improvisation condition, but not the maintain the beat condition. Similarly there was a significant interaction between mean EMG activation and music playing condition on enjoyment, such that enjoyment was positively associated with mean EMG activation for the improvisation condition, but not the maintain the beat condition.

There was no significant main effect of mean EMG activation on Enjoyment (χ^2 (1) = 0.10, p = 0.753) and no significant triple interaction effect between mean EMG activation, music playing condition, and accompaniment on Enjoyment (χ^2 (1) = 0.30, p = 0.586). However, there was a significant interaction effect between mean EMG activation and music playing condition on Enjoyment (Figure 6; χ^2(1) = 5.55, p = 0.0184), while the interaction between music playing condition and accompaniment approached significance (χ^2 (1) = 3.78, p = 0.052). There was also a significant main effect of music playing condition (χ^2 (1) = 5.54, p = 0.019), song (χ^2(3) = 19.90,

p = 0.0002), and sensitivity to musical reward ($\chi^2(1)$ = 8.89, p = 0.003). There was no significant interaction effect for mean EMG activation and accompaniment (χ^2 (1) = 0.07, p = 0.786) and no main effects of musical training (χ^2 (1) = 0.63, p = 0.426) or accompaniment (χ^2 (1) = 2.40, p = 0.122). Pairwise comparisons of song on Enjoyment have been reported above and maintained the same relationships. Simple effects for the association between mean EMG and Enjoyment revealed a positive relationship between mean EMG activation and Enjoyment during the improvisation condition, using the 95% confidence interval to indicate significance (estimated slope = 1.300; 95% CI = 0.483, 2.117) but not the maintain the beat condition (estimated slope = 0.091; 95% CI = -0.784, 0.967). The slopes describing the relationship between Enjoyment and mean EMG activation were significantly different between improvisation and maintain the beat conditions (estimate = 1.21, t(391) = 3.81, p = 0.0002).

There was no significant main effect of beats played on Enjoyment (χ^2 (1) = 0.84, p = 0.360) and no significant triple interaction effect between beats played, music playing condition, and accompaniment on Enjoyment ($\chi^2(1)$ = 0.0065, p = 0.936). The interaction effect between beats played and music playing condition on Enjoyment approached significance (χ^2 (1) = 3.83, p = 0.0503). There was also a significant main effect of song ($\chi^2(3)$ = 26.08, p = 0.000009), and a trend towards a main effect for sensitivity to musical reward ($\chi^2(1)$ = 3.05, p = 0.081). There were no significant interaction effects for beats played and accompaniment (χ^2 (1) = 0.204, p = 0.651) or music playing condition and accompaniment (χ^2 (1) = 2.04, p = 0.154, and no main effects of musical training (χ^2 (1) = 0.247, p = 0.619), accompaniment (χ^2 (1) = 2.56, p = 0.109), or music playing condition (χ^2 (1) = 2.20, p = 0.138). Notably, the previously significant interaction between music playing condition and accompaniment and the main effect of improvisation on Enjoyment became non-significant after including beats played in the model. Pairwise comparisons of song on Enjoyment have been reported above and maintained the same relationships. Simple effects for the association between beats played and Enjoyment revealed a positive relationship between beats played and Enjoyment during the improvisation condition, using the 95% confidence interval to indicate significance (estimated slope = 0.0133; 95% CI = 0.0078, 0.0189) but not the maintain the beat condition (estimated slope = 0.0046; 95% CI = -0.0012, 0.0104). The slopes describing the relationship between enjoyment and beats played were significantly different between improvisation and maintain the beat conditions (estimate = 0.0087, t(496) = 2.53, p = 0.0118).

Discussion
Here we show that it is feasible to increase enjoyment and motor performance during a music playing task by incorporating improvisation and live accompaniment. The combination of improvisation and live accompaniment led to increased acceleration of hand movements while playing the drum, coupled with increased amplitude of muscle contractions in wrist flexor and extensor muscles. Importantly, increased movement was associated with increased reported enjoyment only when improvising. These findings suggest a mechanism by which combined improvisation and live accompaniment increase enjoyment and motivate increased motor activity. We also show that improvisation leads to increased autonomic arousal, as measured by increased tonic EDA and BPM and decreased HRV. Together, these findings have implications for enhancing reward, engagement, and motor performance during music playing tasks that support exercise and physical rehabilitation.

In our experimental design, we did not provide any instructions to participants regarding how much they should move while playing music in any condition. The increased acceleration and muscle activation during conditions with combined improvisation and live accompaniment might thus reflect participants' intrinsic motivation to move. Supporting this mechanism, participants' self-reports of pleasure, motivation, momentum, and activation loaded onto a principal component

that we termed "enjoyment" and that increased for music playing conditions with improvisation and live accompaniment as compared to the other music-playing conditions. Further, there was a significant association between enjoyment and all three measures of movement which was moderated by the music playing condition, such that increased enjoyment was associated with increased total acceleration of hand movements, average activation in the wrist flexor and extensor muscles, and the number of beats played while improvising, but not while maintaining the beat. This relationship between enjoyment and movement is supported by previous research demonstrating that reward enhances motor performance and learning (Terry et al., 2020) and contributes to motor recovery post-stroke (Grau-Sanchez et al., 2018). An association between enhanced music reward and improved motor learning has also been identified in a music playing task involving playing a learned melody (Bianco et al., 2019). In our task, the flexibility of choice provided during improvisation may have played a key role in enhancing reward and movement. Autonomy of choosing what to do has been shown to enhance reward, motor performance, and learning (Wulf et al., 2014; Lewthwaite et al., 2015; Murayama et al., 2015; Chua et al., 2018; An et al., 2020). However, it is also possible that the limited variation in movement while maintaining the beat inhibited reward-enhanced movement for this condition. Together, findings demonstrate that improvisation and live accompaniment may be used to facilitate increased enjoyment and movement during music playing, and that the association between reward and movement is greater during improvisation, when there is flexibility for how and when to play.

Music playing condition also produced a main effect on average muscle activation, the number of beats played, and enjoyment, with improvisation associated with greater motor response and enjoyment as compared to maintaining the beat. This suggests that improvisation, alone, may increase reward and enhance motor performance. Participants chose their own beat, and therefore set their own pace when improvising and maintaining the beat; however, they increased their frequency of movement while improvising, as demonstrated by the increase in the number of beats played. This increase in frequency during improvisation may be partially due to the freedom from constraints in movement frequency, which necessarily limited the number of beats which could be played during the maintain the beat condition. However, the EMG analysis addressed this limitation by averaging peak EMG activation for each movement, in this way measuring movement intensity independent of movement frequency. This analysis revealed increased muscle activation for improvisation as compared to maintaining the beat, supporting the link between improvisation and enhanced motor performance. Together, these findings suggest an important role of improvisation for enhancing reward and motor performance, with mechanisms that may relate to increased autonomy and agency (Wulf et al., 2014; Lewthwaite et al., 2015; Murayama et al., 2015; Chua et al., 2018; An et al., 2020).

The accompaniment condition produced a main effect on enjoyment, with live accompaniment producing the greater enjoyment compared to recorded accompaniment. However, there were no main effects for accompaniment condition on motor or autonomic responses. This suggests that the benefits of live accompaniment are best realized in combination with improvisation, as the interaction between live accompaniment and improvisation led to the greatest levels of enjoyment and enhanced motor performance. Social cooperation has demonstrated benefits for improving intrinsic motivation and motor learning (Kaefer and Chiviacowsky, 2022), and the present findings suggest that the combined effects of live accompaniment and improvisation supported a cooperative effort to coordinate drum and piano playing. Importantly, the piano accompaniment for this study was provided by a music therapist with specialized training in clinical improvisation, which emphasizes responsiveness to participant playing to facilitate a shared music playing experience (Nordoff et al., 2007). Together, these results suggest that live accompaniment, on its own, does not enhance motor performance, but that live accompaniment can enhance the

benefits of improvisation for motor performance through a mechanism of intrinsic reward that might be related to social cooperation.

We also demonstrated that improvisation during music-playing increased autonomic arousal, as measured via tonic EDA, BPM, and HRV, as well as the experience of challenge. Importantly, the association between improvisation and autonomic arousal remains significant after controlling for movement, suggesting that arousal relates to mechanisms beyond physical exertion (Sara and Bouret, 2012). Given the association between improvisation and challenge, it may be that the increased autonomic arousal during improvisation relates to cognitive control. Supporting this potential mechanism, rhythmic improvisation has previously been linked to increased activation in the dorsal anterior cingulate cortex (dACC), when compared to repeating a rhythm (de Aquino et al., 2019), and previous research links dACC activation to cognitive control (Niendam et al., 2012) and evaluation of creative tasks (Ellamil et al., 2012). However, music improvisation has also been linked to activation in the default mode network and deactivation of the executive network, with the extent of executive network deactivation shown to vary with the level of expertise in music improvisation (Pinho et al., 2014; Beaty, 2015). These findings highlight the complexity of improvisation as a creative task that involves both idea generation and evaluation (Ellamil et al., 2012; Loui, 2018). Our findings suggest that cognitive challenge played a role in rhythmic improvisation, an effect that remain significant while controlling for the level of musical expertise.

Finally, our analyses demonstrate that the song composition significantly modulated Enjoyment, Challenge, motor response, and autonomic response during music playing. Familiarity with song genres and the amount of musical groove may have played a role in modulating responses to each song. Although musical preference is thought to vary between individuals, the effects of song in this study suggest future studies should examine how to manipulate song characteristics to enhance motor and affective outcomes during music playing (Matthews et al., 2020; Park et al., 2019; Periera et al., 2011; Witek et al., 2014).

Together, our findings suggest two distinct mechanisms influencing response to music-playing tasks. First, enjoyment during music-playing is facilitated by the combination of improvisation and live accompaniment. During improvisation, enjoyment is associated with increased acceleration of hand movements, muscle activation, and frequency of movement. Second, improvisation increases challenge and autonomic arousal while controlling for motor output, suggesting increased attentional processes when generating musical ideas. Together, these findings have important implications for increasing reward and attention during music-playing tasks, which may promote engagement, dose of movement and improved outcomes during exercise and physical rehabilitation. Understanding the link between music reward and movement is essential for improving our general understanding of the intricate relationship between motor and reward-related regions and their neural substrates (Aoki et al., 2019). Future studies are needed to investigate functional connectivity between reward and sensorimotor regions during music playing, and to better understand the relationship between improvisation, cognitive challenge, and executive function. Additionally, studies investigating these mechanisms in clinical populations have potential to inform the development of music playing tasks to optimize motor rehabilitation (Braun Janzen et al., 2022).

References
Aluru V, Lu Y, Leung A, Verghese J, Raghavan P (2014) Effect of Auditory Constraints on Motor Performance Depends on Stage of Recovery Post-Stroke. Frontiers in Neurology 5.
An J, Lewthwaite R, Lee S, Wulf G (2020) Choice of practice-task order enhances golf skill learning. Psychology of Sport and Exercise 50:101737.

Aoki S, Smith JB, Li H, Yan X, Igarashi M, Coulon P, Wickens JR, Ruigrok TJH, Jin X (2019) An open cortico-basal ganglia loop allows limbic control over motor output via the nigrothalamic pathway. eLife 8:e49995.

Arkin C, Przysinda E, Pfeifer CW, Zeng T, Loui P (2019). Gray matter correlates of creativity in musical improvisation. Frontiers in human neuroscience 13:169.

Bailey LM (1983) The effects of live music versus tape-recorded music on hospitalized cancer patients. Music Therapy 3:17-28.

Bates D, Mächler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models Using lme4. Journal of Statistical Software 67:1 - 48.

Baur K, Speth F, Nagle A, Riener R, Klamroth-Marganska V (2018) Music meets robotics: a prospective randomized study on motivation during robot aided therapy. J Neuroeng Rehabil 15:79.

Beaty RE (2015) The neuroscience of musical improvisation. Neuroscience & Biobehavioral Reviews 51:108-117.

Belfi AM, Samson DW, Crane J, Schmidt NL (2021) Aesthetic Judgments of Live and Recorded Music: Effects of Congruence Between Musical Artist and Piece. Frontiers in Psychology 12.

Belkhir Y, Rekik G, Chtourou H, Souissi N (2019) Listening to neutral or self-selected motivational music during warm-up to improve short-term maximal performance in soccer players: Effect of time of day. Physiology & Behavior 204:168-173.

Berkowitz AL, Ansari D (2008) Generation of novel motor sequences: the neural correlates of musical improvisation. Neuroimage 41:535-543.

Bianco R, Gold BP, Johnson AP, Penhune VB (2019) Music predictability and liking enhance pupil dilation and promote motor learning in non-musicians. Scientific Reports 9:17060.

Blood AJ, Zatorre RJ (2001) Intensely pleasurable responses to music correlate with activity in brain regions implicated in reward and emotion. Proc Natl Acad Sci U S A 98:11818-11823.

Bonassi G, Lagravinese G, Bove M, Bisio A, Botta A, Putzolu M, Cosentino C, Mezzarobba S, Pelosin E, Avanzino L (2023) How Music Moves Us: Music-induced Emotion Influences Motor Learning. Neuroscience 526:246-255.

Braithwaite JJ, Watson DG, Jones R, Rowe M (2013) A guide for analysing electrodermal activity (EDA) & skin conductance responses (SCRs) for psychological experiments. Psychophysiology 49:1017-1034.

Braun Janzen T, Koshimori Y, Richard NM, Thaut MH (2022) Rhythm and Music-Based Interventions in Motor Rehabilitation: Current Evidence and Future Perspectives. Frontiers in Human Neuroscience 15.

Bundt C, Bardi L, Verbruggen F, Boehler CN, Brass M, Notebaert W (2019) Reward anticipation changes corticospinal excitability during task preparation depending on response requirements and time pressure. Cortex 120:159-168.

Bush HI, LaGasse AB, Collier EH, Gettis MA, Walson K (2021) Effect of Live Versus Recorded Music on Children Receiving Mechanical Ventilation and Sedation. Am J Crit Care 30:343-349.

Chua LK, Wulf G, Lewthwaite R (2018) Onward and upward: Optimizing motor performance. Hum Mov Sci 60:107-114.

de Aquino MPB, Verdejo-Román J, Pérez-García M, Pérez-García P (2019) Different role of the supplementary motor area and the insula between musicians and non-musicians in a controlled musical creativity task. Scientific Reports 9:13006.

Eerola T, Toiviainen P (2004) MIDI toolbox: MATLAB tools for music research.

Ellamil M, Dobson C, Beeman M, Christoff K (2012) Evaluative and generative modes of thought during the creative process. NeuroImage 59:1783-1794.

Ferreri L, Mas-Herrero E, Zatorre RJ, Ripollés P, Gomez-Andres A, Alicart H, Olivé G, Marco-Pallarés J, Antonijoan RM, Valle M, Riba J, Rodriguez-Fornells A (2019) Dopamine modulates the reward experiences elicited by music. Proceedings of the National Academy of Sciences 116:3793-3798.

Gorr H (2023) Calculate heart rate from electrocardiogram data. MATLAB Central File Exchange. Retrieved from: https://www.mathworks.com/matlabcentral/fileexchange/73049-calculate-heart-rate-from-electrocardiogram-data.

Grau-Sanchez J, Amengual JL, Rojo N, Veciana de Las Heras M, Montero J, Rubio F, Altenmuller E, Munte TF, Rodriguez-Fornells A (2013) Plasticity in the sensorimotor cortex induced by Music-supported therapy in stroke patients: a TMS study. Front Hum Neurosci 7:494.

Grau-Sanchez J, Duarte E, Ramos-Escobar N, Sierpowska J, Rueda N, Redon S, Veciana de Las Heras M, Pedro J, Sarkamo T, Rodriguez-Fornells A (2018) Music-supported therapy in the rehabilitation of subacute stroke patients: a randomized controlled trial. Ann N Y Acad Sci.

Hallett R, Lamont A (2017) Music Use in Exercise: A Questionnaire Study. Media Psychology 20:658-684.

Hosp JA, Pekanovic A, Rioult-Pedotti MS, Luft AR (2011) Dopaminergic projections from midbrain to primary motor cortex mediate motor skill learning. J Neurosci 31:2481-2487.

Hurd WJ, Morrow MM, Kaufman KR (2013) Tri-axial accelerometer analysis techniques for evaluating functional use of the extremities. Journal of Electromyography and Kinesiology 23:924-929.

Kaefer A, Chiviacowsky S (2022) Cooperation enhances motor learning. Human Movement Science 85:102978.

Karageorghis CI, Mouzourides DA, Priest DL, Sasso TA, Morrish DJ, Walley CJ (2009) Psychophysical and ergogenic effects of synchronous music during treadmill walking. J Sport Exerc Psychol 31:18-36.

Krach S, Paulus FM, Bodden M, Kircher T (2010) The rewarding nature of social interactions. Front Behav Neurosci 4:22.

Laukka P, Quick L (2013) Emotional and motivational uses of music in sports and exercise: A questionnaire study among athletes. Psychology of Music 41:198-215.

Lenth R, Singmann H, Love J, Buerkner P, Herve M (2019) Package 'emmeans': estimated marginal means, aka least-squares means. Retrieved from: https://cran.r-project.org/web/packages/emmeans/index.html.

Lewthwaite R, Chiviacowsky S, Drews R, Wulf G (2015) Choose to move: The motivational impact of autonomy support on motor learning. Psychon Bull Rev 22:1383-1388.

Loui P (2018) Rapid and flexible creativity in musical improvisation: review and a model. Ann NY Acad Sci 1423:138-145.

Mackersie CL, Calderon-Moultrie N (2016) Autonomic nervous system reactivity during speech repetition tasks: heart rate variability and skin conductance. Ear and Hearing 37:118S-125S.

Magee WL, Clark I, Tamplin J, Bradt J (2017) Music interventions for acquired brain injury. Cochrane Database Syst Rev 1:CD006787.

Mas-Herrero E, Marco-Pallares J, Lorenzo-Seva U, Zatorre RJ, Rodriguez-Fornells A (2013) Barcelona Music Reward Questionnaire. Music Perception.

Matthews TE, Witek MA, Lund T, Vuust P, Penhune VB (2020) The sensation of groove engages motor and reward networks. NeuroImage 214:116768.

Menon V, Levitin DJ (2005) The rewards of music listening: response and physiological connectivity of the mesolimbic system. Neuroimage 28:175-184.

Molina-Luna K, Pekanovic A, Röhrich S, Hertler B, Schubring-Giese M, Rioult-Pedotti MS, Luft AR (2009) Dopamine in motor cortex is necessary for skill learning and synaptic plasticity. PLoS One 4:e7082.

Müllensiefen D, Gingras B, Musil J, Stewart L (2014) Measuring the facets of musicality: The Goldsmiths Musical Sophistication Index (Gold-MSI). Personality and Individual Differences 60:S35.

Murayama K, Matsumoto M, Izuma K, Sugiura A, Ryan RM, Deci EL, Matsumoto K (2015) How self-determined choice facilitates performance: a key role of the ventromedial prefrontal cortex. Cereb Cortex 25:1241-1251.

Niendam TA, Laird AR, Ray KL, Dean YM, Glahn DC, Carter CS (2012) Meta-analytic evidence for a superordinate cognitive control network subserving diverse executive functions. Cognitive, Affective, & Behavioral Neuroscience 12:241-268.

Nordoff P, Robbins C, Marcus D (2007) Creative Music Therapy: A Guide to Fostering Clinical Musicianship, 2nd Edition, Revised. Barcelona Pub.

Palumbo A, Aluru V, Battaglia J, Geller D, Turry A, Ross M, Cristian A, Balagula C, Ogedegbe G, Khatri L, Chao MV, Froemke RC, Urbanek JK, Raghavan P (2022) Music Upper Limb Therapy-Integrated (MULT-I) Provides a Feasible Enriched Environment and Reduces Post Stroke Depression: A Pilot Randomized Controlled Trial. American Journal of Physical Medicine & Rehabilitation 101:937-946.

Park KS, Hass CJ, Fawver B, Lee H, Janelle CM (2019) Emotional states influence forward gait during music listening based on familiarity with music selections. Human movement science 66:53-62.

Pereira CS, Teixeira J, Figueiredo P, Xavier J, Castro SL, et al. (2011) Music and Emotions in the Brain: Familiarity Matters. PLOS ONE 6(11): e27241.

Pessiglione M, Schmidt L, Draganski B, Kalisch R, Lau H, Dolan RJ, Frith CD (2007) How the brain translates money into force: a neuroimaging study of subliminal motivation. Science 316:904-906.

Pinho AL, de Manzano Ö, Fransson P, Eriksson H, Ullén F (2014) Connecting to create: expertise in musical improvisation is associated with increased functional connectivity between premotor and prefrontal areas. Journal of Neuroscience 34:6156-6163.

Raghavan P, Geller D, Guerrero N, Aluru V, Eimicke JP, Teresi JA, Ogedegbe G, Palumbo A, Turry A (2016) Music Upper Limb Therapy-Integrated: An Enriched Collaborative Approach for Stroke Rehabilitation. Front Hum Neurosci 10:498.

Ripolles P, Rojo N, Grau-Sanchez J, Amengual JL, Camara E, Marco-Pallares J, Juncadella M, Vaquero L, Rubio F, Duarte E, Garrido C, Altenmuller E, Munte TF, Rodriguez-Fornells A (2016) Music supported therapy promotes motor plasticity in individuals with chronic stroke. Brain Imaging Behav 10:1289-1307.

Salimpoor VN, Benovoy M, Larcher K, Dagher A, Zatorre RJ (2011) Anatomically distinct dopamine release during anticipation and experience of peak emotion to music. Nature Neuroscience 14:257-262.

Sara SJ, Bouret S (2012) Orienting and reorienting: the locus coeruleus mediates cognition through arousal. Neuron 76:130-141.

Shaffer F, Ginsberg JP (2017) An Overview of Heart Rate Variability Metrics and Norms. Frontiers in Public Health 5.

Shaffer F, Shearman S, Meehan Z (2016) The Promise of Ultra-Short-Term (UST) Heart Rate Variability Measurements. Biofeedback 44:229-233.

Simpson SD, Karageorghis CI (2006) The effects of synchronous music on 400-m sprint performance. Journal of Sports Sciences 24:1095-1102.

Stork MJ, Karageorghis CI, Martin Ginis KA (2019) Let's Go: Psychological, psychophysical, and physiological effects of music during sprint interval exercise. Psychology of Sport and Exercise 45:101547.

Terry PC, Karageorghis CI, Saha AM, D'Auria S (2012) Effects of synchronous music on treadmill running among elite triathletes. J Sci Med Sport 15:52-57.

Terry PC, Karageorghis CI, Curran ML, Martin OV, Parsons-Smith RL (2020) Effects of music in exercise and sport: A meta-analytic review. Psychol Bull 146:91-117.

Turry A (2009) Integrating musical and psychological thinking: The relationship between music and words in clinically improvised songs. Music and Medicine 1:106-116.

Vuust P, Heggli OA, Friston KJ, Kringelbach ML (2022) Music in the brain. Nat Rev Neurosci 23:287–305.

Widmer M, Ziegler N, Held J, Luft A, Lutz K (2016) Rewarding feedback promotes motor skill consolidation via striatal activity. Prog Brain Res 229:303-323.

Witek MAG, Clarke EF, Wallentin M, Kringelbach ML, Vuust P (2014) Syncopation, Body-Movement and Pleasure in Groove Music. PLOS ONE 9(4): e94446.

Wulf G, Lewthwaite R (2016) Optimizing performance through intrinsic motivation and attention for learning: The OPTIMAL theory of motor learning. Psychon Bull Rev 23:1382-1414.

Wulf G, Freitas HE, Tandy RD (2014) Choosing to exercise more: Small choices increase exercise engagement. Psychology of Sport and Exercise 15:268-271.

EXTENDED MATERIALS

Figure 1.1: Condition Orders

Participants were randomized to one of four condition orders. Each condition order alternated between music playing conditions (improvising and maintaining the beat), with half of the condition orders starting with improvisation. Accompaniment conditions (live and pre-recorded) were completed consecutively, such that participants completed both music playing conditions with one accompaniment condition and then repeated both music playing conditions with the other accompaniment condition. The same condition order was repeated for each of the four songs played by each participant.