Differential reorganization of episodic and semantic memory systems in epilepsy-related mesiotemporal pathology

Donna Gift Cabalo[1,2], Jordan DeKraker[1,2], Jessica Royer[1,2,3], Ke Xie[1,2], Shahin Tavakol[1,2], Raúl Rodríguez-Cruces[1,2], Andrea Bernasconi[2,4], Neda Bernasconi[2,4], Alexander Weil[5], Raluca Pana[2], Birgit Frauscher[2,3], Lorenzo Caciagli[6], Elizabeth Jefferies[7], Jonathan Smallwood[8], Boris C. Bernhardt[1,2]

Author affiliations:

Multimodal Imaging and Connectome Analysis Laboratory, McConnell Brain Imaging Centre, Montreal Neurological Institute, McGill University, Montreal, QC, Canada;

Montreal Neurological Institute and Hospital, McGill University, Montreal, QC, Canada;

Analytical Neurophysiology Laboratory, Montreal Neurological Institute, McGill University, Montreal, Quebec, Canada;

Neuroimaging of Epilepsy Laboratory, McConnell Brain Imaging Centre, Montreal Neurological Institute and Hospital, McGill University, Montreal, QC, Canada;

CHU St Justine, Montreal, QC;

Department of Bioengineering, University of Pennsylvania, Philadelphia, USA;

University of York, North Yorkshire, United Kingdom;

Queen's University, Kingston, Ontario, Canada

Correspondence to:

Donna Gift Cabalo or Boris Bernhardt

3801 Rue University, Montréal, QC, Canada, H3A 2B4

donna.cabalo@mail.mcgill.ca | boris.bernhardt@mcgill.ca

Keywords: declarative memory; mesiotemporal lobe; neuroimaging; connectomics; gradients; epilepsy

Abstract

Declarative memory encompasses episodic and semantic divisions. Episodic memory captures singular events with specific spatiotemporal relationships, while semantic memory houses context-independent knowledge. Behavioural and functional neuroimaging studies have revealed common and distinct neural substrates of both memory systems, implicating mesiotemporal lobe (MTL) regions such as the hippocampus and distributed neocortices. Here, we explored declarative memory system reorganization in patients with unilateral temporal lobe epilepsy (TLE), as a human disease model to test the impact of variable degrees of MTL pathology on memory function. Our cohort included 31 patients with TLE as well as 60 age and sex-matched healthy controls and all participants underwent episodic and semantic retrieval tasks during a multimodal MRI session. The functional MRI tasks were closely matched in terms of stimuli and trial design. Capitalizing on non-linear connectome gradient mapping techniques, we derived task-based functional topographies during episodic and semantic memory states, both in the MTL and in neocortical networks. Comparing neocortical and hippocampal functional gradients between TLE patients and healthy controls, we observed a marked topographic reorganization of both neocortical and MTL systems during episodic memory states. In the neocortex, alterations were characterized by reduced functional differentiation in patients in lateral temporal and midline parietal cortices in both hemispheres. In the MTL, on the other hand, patients presented with a more marked functional differentiation of posterior and anterior hippocampal segments ipsilateral to the seizure focus and pathological core, indicating perturbed intrahippocampal connectivity. Semantic memory reorganization was also found in bilateral lateral temporal and ipsilateral angular regions, while hippocampal functional topographies were unaffected. Leveraging MRI proxies of MTL pathology, we furthermore observed alterations in hippocampal

microstructure and morphology are associated with TLE-related functional reorganization during episodic memory. Moreover, correlation analysis and statistical mediation models revealed that these functional alterations contributed to behavioural deficits in episodic, but again not semantic memory in patients. Altogether, our findings suggest that semantic processes rely on distributed neocortical networks, while episodic processes are supported by a network involving both the hippocampus and neocortex. Alterations of such networks can provide a compact signature of state-dependent reorganization in conditions associated with MTL damage, such as TLE.

Introduction

Declarative memory is commonly divided into episodic and semantic memory components. Episodic memories are unique events with specific spatiotemporal associations, while semantic memory refers to context-invariant knowledge and facts about the world.[1,2,3] The relationship between both forms of memory is complex, with behavioural and neuroimaging studies suggesting shared and distinct substrates. Behavioural studies have shown a synergistic interplay between both memory types, where episodic memory for events and words processed with semantic categorization, exceeds those from non-semantic tasks.[4,5,6] Functional and structural neuroimaging evidence in healthy populations supports this finding, suggesting shared neural substrates across both memory types. Shared key networks are situated in the medial and lateral temporal lobes,[7,8, 9,10,11,12,13-16,17,18] alongside posterior parietal regions.[19-21] More broadly, networks involved in both memory types have been shown to engage similar intrinsic functional systems, notably the fronto-parietal network[22-24] (FPN) and the default mode network (DMN).[10,25,26] As such, behavioural and neural evidence collectively implicates shared processes in both forms of declarative memory. On the other hand, findings also point to some divergences.

Generally, episodic processes appear to rely more on medial temporal lobe (MTL) systems, while anterior temporal lobe (ATL) regions are more heavily implicated in semantic processing.[10,16] Similarly, prior work has suggested differential activations in middle frontal *vs* inferior temporal regions in episodic as compared to semantic retrieval.[27] A neural divergence between both memory systems is supported by the *(1)* controlled semantic cognition framework, which stipulates that semantic representations within the ATL 'hub' and modality-specific 'spokes' interact with a 'semantic' control' system that dynamically supervises and adjusts semantic retrieval bilaterally,[1,28-30] and *(2)* hippocampus-neocortical connectivity, such that the hippocampus interacts closely with other regions through process-specific alliances to support episodic memory.[13,31,32] A systematic investigation of neocortical and MTL networks will help to understand substrates contributing to declarative memory function in the human brain.

Temporal lobe epilepsy (TLE) is a common pharmaco-resistant epilepsy in adults and can serve as a human disease model to probe declarative memory system reorganization in young and middle-aged adults.[32,33] Patients typically present with pathology of structures in the MTL, notably the hippocampus[34-37] and prior research has demonstrated marked impairments in episodic memory, in some patients even at a young age.[32,38-40] Episodic memory impairment has been directly related to degrees of hippocampal structural alterations,[41-46] together with widespread structural and functional imbalances in both temporal and extratemporal neocortical networks.[47,48] In contrast to the relatively consistent finding of episodic memory impairment in TLE, semantic memory has been less frequently studied, and reports remain mixed. Semantic memory function appears to be rather mildly affected,[46,49] even in TLE patients with pronounced MTL hypometabolism.[49] Moreover, and in contrast to the frequent finding of episodic memory disruptions in patients undergoing selective MTL resection[50-52] there are less marked semantic

memory impairments seen postsurgically.[53,54] Such resilience suggests a more distributed cortical control network substrate for semantic memory involving both hemispheres,[55,56] in contrast to MTL representations of episodic memory.[57,58] Individuals with TLE may, therefore, have difficulty recalling specific events or episodes from their past while retaining their general knowledge of the world. However, other studies have found significant semantic memory deficits in patients with TLE,[59-61] which have been associated with MTL[41] and ATL lesions.[62] Additionally, poor performance on semantic memory tasks has been demonstrated in TLE patients after unilateral ATL resection.[59,63,64] Nevertheless, as impairments in declarative memory challenge the day-to-day functioning and well-being of patients, it is critical to derive more mechanistic insights into these memory types and their differential vulnerabilities in TLE.

Contemporary systems neuroscience has emphasized a key role of the spatial organization of macroscale functional networks in human cognition.[65-68] Recent analytical and conceptual advances have notably emphasized the existence and utility of spatial *gradients*[65,68-71] as low-dimensional axes of cortical organization, often reflective of established principles of cortical topography and hierarchy.[65,72,73] At the level of intrinsic functional connectivity derived from task-free functional MRI, the first/principal gradient anchors primary sensory and motor areas at one end and transmodal association systems, like the FPN and DMN, at the other end, whereas a second gradient differentiates visual and somatosensory/motor areas.[65] Gradients also compactly describe the subregional hippocampal organization, with a first gradient differentiating anterior-posterior (long axis) divisions and a second gradient the proximal-distal arrangement of different subfields along the infolding of this allocortical structure.[74-77] Beyond the application of gradient mapping techniques to study healthy brain organization,[65,76,78-85] recent studies have interrogated gradient alterations in diseased cohorts, including TLE patients.[86,87] A recent study from our

group demonstrated microstructural gradient contractions in TLE in the neocortex, which were primarily localized in temporo-limbic systems and found to track deficits in episodic memory recall accuracy.[87] Another study demonstrated functional gradient expansion in subcortical structures during resting conditions, with notable differences in ipsilateral hippocampal gradient deviations between left *vs* right-lateralized TLE.[86] Additionally, patterns of cortical asymmetry and atrophy were shown to have been distinctively linked with microstructural and functional gradients.[88] Gradient mapping, therefore, offers a framework to interrogate global memory network topographies and their association with cognition in both health and disease.

The current study investigated declarative memory network organization in healthy individuals and a cohort of TLE patients presenting with variable degrees of MTL pathology. All participants underwent episodic and semantic retrieval tasks inside the MRI scanner, with both tasks being closely matched for stimulus presentation, task design, and task structure.[32,46] For each of the declarative memory states, we mapped spatial gradients of memory state connectivity, both within the MTL and in macroscale neocortical networks. By adopting gradients as an analytical and conceptual framework for task-derived functional MRI data in TLE, we were able to answer the following questions: (*i*) whether networks subserving episodic *vs* semantic memory in patients undergo a shared or selective reorganization relative to controls, (*ii*) whether memory network reorganization reflects TLE-related structural alterations in the MTL, capitalizing on established *in vivo* proxies of mesiotemporal pathology[35,89 90,91] (*iii*) whether these findings reflect behavioural impairment in relational memory seen in patients.

Materials and methods

Participants

We studied 31 patients with drug-resistant TLE (17 females, age=35±11.59 years, left/right handed=1/30), and 60 healthy controls (30 males, age=33± 8 years, left/right handed=2/58). All participants were recruited between May 2018 and March 2023 at the Montreal Neurological Institute (MNI). Controls met the following inclusion criteria: (1) age between 18-65 years; (2) no neurological or psychiatric illness; (3) no MRI contraindication; (4) no drug/alcohol abuse problem; and (5) no history of brain injury and surgery. Patient demographics and clinical features were obtained through interviews with patients and their relatives. Seizure focus lateralization was determined through a comprehensive evaluation of medical history, neurological examination, seizure semiology, video-EEG, and clinical neuroimaging. Detailed clinical data, including location and Engel outcome, is available in **Supplementary Table S1.** Twenty-one patients were diagnosed with left-sided TLE. The mean age of seizure onset was 20.42±10.22 years (range=2-49 years), with a mean epilepsy duration of 14.20±10.97 years (range=1-45 years). Eight patients (26%) had a history of childhood febrile convulsion. Based on quantitative hippocampal MRI volumetry [92], 20/31 patients (65%%) showed marked hippocampal atrophy ipsilateral to the focus (*i.e.*, absolute ipsilateral-contralateral asymmetry z-score > 1.5 and/or ipsilateral volume z-score < -1.5). At the time of the study, 14/31 patients underwent resective temporal lobe surgeries. In those, post-surgical seizure outcome was assessed using Engel's modified classification[93], with an average follow-up duration of 23±15.50 months. Post-surgery, 9 patients (64%) achieved complete seizure freedom (Engel-I), and 5 patients (36%) had recurrent seizures. A total of 9 specimens were available for histopathological

analysis. Here, 4 patients showed mesiotemporal/hippocampal sclerosis, one patient had evidence of mesiotemporal ganglioglioma, and 4 for mild dysplasia in the mesiotemporal structures. There were no significant differences in males/females (χ^2=0.20, p=0.66) and there were no significant differences in age (t=-1.01, p=0.32) between patients and controls. The study was conducted in accordance with the Declaration of Helsinki, and the MRI data acquisition protocols were approved by the Research Ethics Board of the Montreal Neurological Institute and Hospital. All participants provided informed consent.

Declarative memory paradigm

All participants underwent episodic and semantic retrieval fMRI tasks (**Figure 1**). These tasks were carefully designed to be closely matched *i.e.,* they were based on (1) equivalent symbolic stimuli to accommodate a bilingual participant pool in the Montreal area; (2) the same number of trials that were presented in a pseudo-randomized manner; (3) an identical task structure with three alternative forced choice response at the retrieval phases; and (4) two difficulty levels.

A) Episodic Memory. The task had two phases. During the encoding phase, participants were shown paired images of objects and asked to memorize them. To ensure that the images were well-matched, we used the UMBC similarity index[94] to control for semantic relatedness. Specifically, we selected items where the similarity score remained below 0.3 on a scale of 0 to 1. The task comprised 56 pseudo-randomized trials, with 28 difficult (with paired images shown only once) and 28 easy (paired images shown twice) trials. In the retrieval phase, which took place after several minutes of delay, participants were presented with an image of an object at the top of the screen (*prime*) and three different objects at the bottom. From these choices, subjects

were asked to identify the *target* that was paired with the *prime* during encoding. We only analyzed data from the retrieval phase.

B) Semantic Memory. The semantic task involved a retrieval phase only. In each trial, participants were presented with an image of an object at the top of the screen (*prime*) and three different objects at the bottom (1 *target*, 2 *foils*). From these choices, subjects were asked to identify the item (*target*) that was most conceptually related to the *prime*. The 56 pseudo-randomized trials were also modulated for difficulty (*i.e.,* there were 28 difficult trials where prime/target shared a similarity index ≥0.7, whereas prime/foils shared a similarity between 0.3 and 0.7, as well as 28 easy trials, where prime/target shared a similarity ≥0.7 and prime/foils shared a similarity between 0 and 0.3).

C) Performance metrics. Individual episodic and semantic performance was computed by averaging the scores in the difficult and easy trials. We also averaged reaction time.

Figure 1 | Episodic and Semantic memory tasks. **A**. The episodic memory task included encoding **(i)** and retrieval phases **(ii)**, while **(B) the** semantic memory task only included a retrieval phase **(v)**. Both tasks were matched in terms of stimuli, number of trials, difficulty levels, and task structure, such that retrieval phases included three alternative forced-choice responses. Functional connectomes (FC; **iii, vi**) were independently derived from episodic and semantic task FC and affinity matrices (AM) were built based on cosine similarity, followed by gradient decomposition. (**iv, vii**). Neocortical and hippocampal principal gradients (G1) projected to the corresponding surfaces.

Additional screening for executive function and general cognitive impairment

Participants also completed the MoCA[95] tests outside the scanner. The paradigm has been developed to detect mild cognitive impairment and dementia.

MRI Acquisition

MRI data were acquired on a 3T Siemens Magnetom Prisma-Fit with the 64-channel head coil. T1-weighted scans were acquired with a 3D-magnetization-prepared rapid gradient-echo sequence (MPRAGE; 0.8 mm isovoxels, matrix=320×320, 224, sagittal slices, TR=2300 ms, TE=3.14 ms, TI=900 ms, flip=9°, iPAT=2, partial Fourier=6/8, FOV=256×256 mm^2). The DWI data was obtained with a 2D spin-echo echo-planar imaging sequence with multi-band acceleration and consists of three distinct shells, each with different b-values 300, 700, and 2000s/mm^2, with each shell acquired with 10, 40, and 90 diffusion weighting directions, respectively (1.6 mm isotropic voxels, TR=3500 ms, TE=64.40 ms, flip angle=90°, refocusing flip angle=180°, FOV=224×224 mm^2, slice thickness=1.6 mm, multi-band factor=3, echo spacing=0.76 ms). All fMRI were acquired with a 2D/BOLD echo-planar imaging sequence (3.0 mm isovoxels, matrix=80×80, 48 slices oriented to AC-PC-30 degrees, TR=600 ms, TE=30 ms, flip=50°, FOV=240×240 mm^2, slice thickness=3 mm, multi-band

factor=6, echo spacing=0.54 ms). During the resting-state functional MRI (rs-fMRI), participants were instructed to fixate on a grey cross and not think of anything. All fMRI scans were presented via a back-projection system and responses were recorded using an MRI-compatible four-button box. The episodic and semantic retrievals both lasted approximately 6 minutes.

MRI Processing

Scans were pre-processed using *micapipe*[96] *version 0.2.2* (https://github.com/MICA-MNI/micapipe), an open-access multimodal preprocessing, surface-mapping, and data fusion software. T1-weighted data were de-obliqued and reoriented to ensure a consistent orientation across images. Next, a linear coregistration, intensity non-uniformity correction, and skull stripping were applied to extract the brain region from the surrounding non-brain tissues. DWI data underwent denoising, eddy current-induced distortion correction, and motion correction as well as non-uniformity bias field adjustment. The corrected DWI data was fitted with a diffusion tensor model,[97] allowing for the computation of mean diffusivity images (MD).[98] Cortical surfaces were extracted with FreeSurfer *6.0* from each T1w scan, followed by manual surface generation. Resting-state and task-fMRI were preprocessed using a combination of FSL[99] *6.0*, ANTs[100] *2.3.4*, and AFNI[101] *20.3.0* software and involved the following steps: *(1)* removal of first five TRs, *(2)* image reorientation, *(3)* motion correction, *(4)* distortion correction based on AP-PA blip field maps, *(5)* high-pass filtering to be above 0.01Hz, *(6)* MELODIC decomposition, *(7)* Linear/Non-linear co-registration to the T1-weighted scans, *(8)* nuisance signal regression with FMRIB's ICA-based Xnoiseifier[102] (ICA-FIX), *(9)* native cortical surface registration, *(10)* surface-based registration to the Conte69 surface template with 32k surface points (henceforth, *vertices*) per hemisphere, *(11)* surface-based smoothing with a Gaussian

diffusion kernel with a full-width-at-half-maximum, FWHM, of 10 mm, and *(12)* statistical regression of motion spikes. The hippocampus and surrounding structures were automatically segmented using HippUnfold[92] *version 1.0.0,* (https://hippunfold.readthedocs.io/). The task-fMRI and MD data were then sampled along the hippocampal midthickness surface and FWHM=2 mm surface smoothing was applied.

Gradient identification and alignment

For computational efficiency, surface-mapped neocortical time series were downsampled to a mesh with 10k vertices using *connectome workbench.*[103] Intrahemispheric neocortical functional connectomes were generated for each functional state (episodic, semantic, rest) by calculating Pearson correlations between the time series for all vertices for each participant, resulting in a 5k-by-5k connectome per hemisphere (**Figure 1**). As for the neocortex, internal hippocampal connectomes were generated from the extracted timeseries resulting in a 419 by 419 functional connectivity matrix per hippocampus. We applied Fisher r-to-z transformations to both neocortical and hippocampal functional connectomes and retained only the top 10% of row-wise connections. We constructed affinity matrices using a cosine similarity kernel to measure the similarity of connectivity patterns between regions. Diffusion map embedding,[65,104] a non-linear dimensionality reduction technique implemented in *Brainspace*[105] *version 0.1.3* (https://brainspace.readthedocs.io/; **Figure 1**), was applied to identify low-dimensional eigenvectors (*i.e.,* gradients) that accounted for the variance in neocortical and hippocampal connectomes. Diffusion mapping is a powerful and efficient tool to compress large-scale datasets. This algorithm uses the diffusion operator $P\alpha$ to create a new representation of data and is guided by two key parameters: α, which controls the influence of sampling point density on

the manifold, and, *t*, which represents the scale of the data. To maintain global connections among data points in the embedded space, *α* was set to 0.5 (*α=0, maximal influenc*e, *α=1 no influence)* and *t* was set at 0, consistent with previous studies.[65,105] To ensure that results were consistent with the prior literature that derived functional gradients mainly from rs-fMRI data,[65,70] we first generated intra-hemispheric reference gradients by averaging rs-fMRI connectomes across groups and aligning them to a normative gradient template derived from the rs-fMRI data of the Human Connectome Project.[106] We used Procrustes rotation to align individual and state-specific task-fMRI gradients with these normative reference gradients, thereby ensuring that gradients were consistent in ordering and sign across participants and memory states[107]. Similar procedures were performed to align hippocampal gradients. Finally, we normalized neocortical and hippocampal gradient scores relative to the distribution in controls, and sorted gradients in patients relative to the hemisphere ipsilateral to the seizure focus, as in prior work.[87]

Statistical Analysis

After gradient alignment, we inferred between-group differences using surface-based linear models implemented in *Brainstat*[108] *version 0.4.2* (https://brainstat.readthedocs.io). We additionally controlled for age and sex and corrected for vertex-wise multiple comparisons using random field theory for non-isotropic imaging,[109] at a set family-wise error of $p_{FWE}<0.05$. We computed the association between gradient scores in regions showing significant between-group differences and cognitive scores. To investigate the association between gradients and memory, we examined the gradient scores while controlling for MoCA which indicated mild cognitive

impairments. Finally, statistical mediation analyses tested whether the association between group and episodic memory performance is transmitted via gradients.

Morphological and microstructural substrate analysis

To assess structural substrates of functional network changes, we also compared neocortical and hippocampal thickness and mean diffusivity (MD) between controls and TLE patients. These features have previously been utilised to demonstrate both hippocampal as well as neocortical alterations in TLE. [35,110,111] Neocortical thickness was quantified as the Euclidian distance between corresponding pial and white matter vertices, and we used mid-thickness surfaces to sample MD from co-registered diffusion MRI data. To establish an accurate spatial correspondence, cortical thickness and MD maps were registered to the Conte69 surface template and were subsequently downsampled from 32k vertices to 5k vertices/hemisphere. Hippocampal thickness and hippocampal MD was measured using a similar method, using hippocampal surface meshes derived with HippUnfold[92] *v1.0.3,* (https://hippunfold.readthedocs.io/). Multivariate surface-based linear models then assessed morphological and microstructural differences between patients and controls in both hippocampal and neocortical regions, while controlling for effects of age and sex. Finally, we investigated the association between atypical morphology patterns and microstructure and observed TLE-related gradient alterations. Specifically, we computed the mean thickness and MD scores in regions showing significant functional G1 differences between controls and TLE patients, and correlated these measures with G1 changes.

Functional decoding via task-based meta-analysis

We also conducted a functional decoding analysis based on Neurosynth [26] (https://neurosynth.org/), a platform that allows for large-scale *ad hoc* meta-analysis of task-based fMRI data. Here we studied the spatial associations between TLE-related functional gradient changes and previously published fMRI activations to identify cognitive terms that elicit similar activation patterns.

Data availability

The Human Connectome Project dataset used for generating normative resting-state gradients is available at https://db.humanconnectome.org/. Our healthy control cohort is made up of a subset of participants from the MICA-MICs[112] dataset which is openly available on the Canadian Open Neuroscience Platform data portal (https://portal.conp.ca/dataset?id=projects/mica-mics) and Open Science Framework (https://osf.io/j532r/). Functional connectome gradients will be available on osf.io.

Results

Hippocampal and neocortical surface models for each participant were generated using automatic segmentation[92,96] procedures based on T1-weighted MRI. Functional data were co-registered to these surfaces, and resulting neocortical and hippocampal timeseries were used to derive functional connectomes for resting, episodic, and semantic memory states for each participant. Diffusion map embedding,[105] a non-linear dimensionality reduction technique, was applied to estimate low-dimensional eigenvectors explaining spatial gradients of connectivity variance. For consistency, reference gradients were derived from rs-fMRI data[106] and subject- and state-specific neocortical and hippocampal gradients were aligned to these using Procrustes

rotations, as in previous work [105,107] (for pre-alignment gradient patterns, please see *Supplementary Figure S1*).

Declarative memory topographies

Studying neocortical connectivity in healthy individuals, we derived the first eigenvectors (G1) during both episodic and semantic states separately (for findings of subsequent gradients, please see *Supplementary Figure S2*). In episodic/semantic states, G1 explained 22/21% of functional connectome variance and described a sensory-to-transmodal neocortical gradient, as expected from previous work in healthy adults based on rs-fMRI connectivity.[65] Studying hippocampal networks in controls, we also derived the first eigenvectors in episodic and semantic memory states (**Figure 1**). In episodic/semantic states, G1 accounted for 40/38% of functional connectome variance and described a canonical postero-anterior spatial pattern.[76] State-specific neocortical and hippocampal gradients spatially correlated in healthy controls (r=0.99, p=0, n_{perm}=1000), suggesting that both memory systems were captured by both sensory-transmodal neocortical and posterior-anterior MTL trends.

Atypical task-based gradients in patients with MTL pathology

Comparing neocortical topographies in TLE patients to controls, we found reduced G1 scores in the episodic memory state in patients (p_{FWE}<0.05, mean effect sizes in significant regions d>0.50; **Figure 2A**). Functional G1 alterations were primarily localized in the ipsilateral lateral temporal and bilateral posterior cingulate regions. Between-group differences were also seen in the semantic memory state (**Figure 2B**), localized in the bilateral lateral temporal and ipsilateral angular gyrus. The TLE-related functional G1 contractions are indicative of a functional

dedifferentiation between these unimodal and transmodal systems. Consistent with this finding, the histogram depicting overall mean gradient scores, showed an overall reduction in the spread of gradient scores in TLE patients relative to controls (**Figure 2**). Considering the hippocampus, conversely, we observed an expansion of G1 scores in the episodic memory state when comparing patients to controls, which was particularly significant in anterior divisions ipsilaterally (p_{FWE}<0.05, d=-0.84). The expanded functional G1 in the anterior portions reflects increased connectivity profile variability in TLE, indicating a great intra-hippocampal functional disconnection. In other words, the anterior hippocampal gradient scores in TLE are shifted away from the gradient midpoint, further increasing inter-regional differentiation with the posterior hippocampus. While semantic G1 expansion was also observed in TLE, no vertices survived multiple comparison correction.

Figure 2 | Gradient alterations in patients with MTL pathology. Group differences in neocortical (**i,v**) and hippocampal (**ii, vi**) gradient scores for episodic (**A**) and semantic states (**B**). Significant *p*-values were highlighted after correcting for multiple comparisons (p_{FWE}<0.05, black outlines). Histogram plot of neocortical gradient contractions (**iii,vii**) and hippocampal gradient expansions (**iv,viii**) in TLE patients relative to controls in episodic

and semantic states, respectively. **Abbreviations:** HC=healthy controls, TLE= temporal lobe epilepsy, G1 = Gradient 1, G2= Gradient 2.

Relation to temporo-limbic structural alterations

An additional analysis explored how proxies of TLE-related structural pathology[35] contribute to episodic functional memory network reorganization. Proxies were derived from neocortical and hippocampal MRI measures of grey matter (*i.e.,* cortical thickness) and diffusion parameters (*i.e.,* cortical mean diffusivity, MD). Comparing patients and controls using multivariate models based on these parameters, significant alterations were observed in patients ($p<0.05$, **Figure 3A, B;** for univariate findings, see ***Supplementary Figure S3***), mainly localized in ipsilateral tempo-limbic areas and anterior regions of the ipsilateral hippocampus. We next examined whether these structural disruptions are associated with previously observed functional alterations (**Figure 3C**). To this end, we computed the overall temporo-limbic structural changes in TLE patients and controls, both at the level of the neocortex and hippocampus and conducted a correlation analysis. This analysis revealed an association between overall episodic functional G1 changes and structural alterations (r=-0.22, p_{FDR}=0.03). Further tests showed a slightly more selective association with hippocampal (r=-0.22, p_{FDR}=0.04) than with neocortical functional G1 measures (r=0.19, p_{FDR}=0.08).

Figure 3 | Associations to temporo-limbic pathology. A-B) Differences in neocortical **(i)** and hippocampal **(iii)** MRI-proxies of pathology (using a multivariate aggregate of cortical thickness and mean diffusivity) between TLE patients and controls. Regions showing alterations in TLE are highlighted ($p_{FWE}<0.05$, black outline), with mean values represented in **(ii, iv).** **C)** FDR-corrected relationship between overall episodic changes to **(v)** overall structural changes, **(vi)** structural changes in the neocortex, and, **(vii)** hippocampus (HC=grey; TLE=dark purple). **Abbreviations:** ipsi= ipsilateral, contra=contralateral, HC=healthy controls, TLE=temporal lobe epilepsy.

Relation to behavioural indices of declarative memory

We examined the relations between participants' behavioural performance and functional memory network reorganization. Behaviorally, patients showed markedly reduced episodic memory accuracy relative to controls (t=-4.40, $p<0.001$, d=-0.97), and only subthreshold impairment in semantic memory (t=-1.42, p=0.16, d=-0.31). We similarly observed faster reaction times for accurate recall in controls relative to patients (***Supplementary Figure S5***). Notably, functional G1 scores in both neocortical (r_{ipsi}=0.29, p_{FDR}=0.02; r_{contra}=0.27, p_{FDR}=0.02)

and hippocampal (r_{ipsi}=0.29, p_{FDR}=0.02) clusters of significant between-group differences (see *Figure 2*) correlated with episodic memory performance (**Figure 4A**). *Ad hoc* meta-analytical decoding[26] confirmed that TLE-related gradient contractions in the episodic state were enriched for memory-related terms, supporting the behavioural associations (***Supplementary Figure S6***). In semantic memory states, hippocampal functional; G1 (r_{ipsi}=0.21, p_{FDR}=0.06) correlated with semantic task performance at trend levels, but not neocortical functional G1 (r_{ipsi}=0.02, p_{FDR}=0.85; r_{contra}=-0.03, p_{FDR}=0.95; **Figure 4B**).

To assess specificity, we also administered the MoCA test to probe mild cognitive dysfunction. Patients with TLE showed reduced MoCA (t=-3.37, p=0.001, d=-0.74) scores relative to controls. Controlling for individual MoCA scores, significant functional G1 changes persisted in our cohort, in both the hippocampus (p_{FWE}<0.05, d=-0.80, 4% increase in mean effect size) and neocortex (p_{FWE}<0.05, d=-0.50).

In a final analysis, we assessed whether functional G1 alterations mediated the relationship between group (TLE *vs* controls) and episodic memory performance (**Figure 4C,** for individual mediation, see ***Supplementary Table S2***). To this end, we first computed overall episodic alterations by averaging G1 scores in significant regions and submitted this to a statistical mediation analysis. Group and functional G1 alterations correlated strongly (a, p<0.0005), as did functional G1 alterations and episodic memory performance (b, p<0.05), and group and episodic memory performance (c', p<0.05). Importantly, functional G1 alterations in hippocampal and neocortical regions mediated the relationship between group and episodic memory performance (ab, p<0.05). Overall, our statistical mediation analysis, thus, revealed that the relationship

between group and episodic memory performance is transmitted via neocortical and hippocampal functional gradients.

Figure 4 | Behavioral associations. Mean G1 scores of regions showing significant case-control differences in the ipsilateral and contralateral hemispheres for neocortical and hippocampal G1 in episodic **(A)** and semantic **(B)** memory states, correlated with overall behavioural performance. Controlling for the aggregate effects of morphology and superficial white matter properties (mean diffusivity), hippocampal episodic G1 (**ii;** *right panel)* correlated significantly with task performance, while hippocampal semantic G1 (**iv;** *right panel)* correlation to memory performance disappeared. (HC=grey; TLE=blue). **C.** Statistical mediation analysis revealed the relationship between the group and episodic memory performance, transmitted via overall neocortical G1+ hippocampal G1$_{ipsi}$ **(v)** scores. **(vi)** Histogram of effects (nboot=1000). **Abbreviations:** G1 = Gradient 1, Ipsi=

ipsilateral, Contra= contralateral, a = Group ~ Gradient scores, b= Gradient scores ~ Episodic memory performance, c'= Group ~ Episodic memory performance(residual), ab= mediation effect.

Discussion

This study demonstrated *(i)* a selective reorganization of both neocortical and hippocampal episodic functional memory networks in patients with TLE, *(ii)* a close association between functional reorganization and hippocampal, but not neocortical *in vivo* proxies of MTL pathology, and *(iii)* a contribution of atypical temporo-limbic functional organization to behavioural deficits in TLE patients. Our study harnessed several innovative elements, notably closely matched functional MRI tasks of episodic and semantic memory administered in TLE patients and controls, state-of-the-art MRI segmentation and multimodal image preprocessing methods, as well as connectivity gradient mapping techniques to identify functional topographies in a data-driven manner. These techniques were applied to a cohort of TLE patients with variable degrees of MTL pathology and memory impairment, allowing for the study of structure-function relationships in declarative memory systems in the human brain. By identifying the functional topographic alterations of episodic memory networks in TLE that mediate cognitive impairment and their structural underpinnings, our work emphasizes a key role of the MTL in episodic over semantic memory processes.

The segregation of episodic and semantic memory systems[3] supported by our findings is in line with foundational neuropsychological investigations dissociating list recall performance for episodic memory and picture naming for semantic memory.[38,60,61] Moreover, task-based neuroimaging studies have pointed to distinct neural substrates of these task demands.[9,16] The memory tasks[46] we employed were carefully designed to distinctly tap into episodic and

semantic memory while being homogenous in low-level task structure. A recent behavioral study from our group supported the utility of these tasks to assess cognition in both healthy and diseased cohorts, showing reduced episodic, but relatively preserved semantic recall in TLE.[46] Despite these potential differential impairments of TLE patients, mounting evidence suggests that episodic and semantic memory systems may nevertheless share overlapping neural correlates.[9,22,23] By capturing both distinctive features and potential overlaps in the neural underpinnings of episodic *vs* semantic memory, we employed a data driven approach that mapped functional topographic patterns associated with each of these processes. In particular, we analyzed task-based fMRI data across both episodic and semantic memory states, and mapped resulting task-based connectomes into topographic connectivity gradients.[65,70] Gradient mapping techniques have previously been applied to several modalities, in particular resting-state fMRI connectivity as well as structural and microstructural MRI measures.[65,70,72,82,113,114] Instead of performing a conventional functional gradient mapping across task-free fMRI sessions, the current study derived topographic gradients during episodic and semantic memory states separately, and found that they largely followed similar intrinsic organizational axes irrespective of the choice of gradient alignment.[10,23,25] Specifically, gradients associated to both memory states followed a sensory-transmodal neocortical and posterior-anterior hippocampal pattern, in line with previous observations.[65,70,71,74-76] Transmodal core regions (*i.e.,* DMN) are recognized to occupy a cortical territory that is maximally distant and separate from unimodal sensory systems, balancing segregated processing streams and integration.[66,68,65,66,115] This is suggested to help decouple cognition, for instance through self-generated thought processes and episodic representations, from salient sensory information of the immediate external environment.[116-121] Conversely, in the hippocampus, anterior segments are thought to have broader tuning properties

and are suited to generalization, while posterior segments are more narrowly tuned and appear more suited to memorizing specific instances for contextualization.[122,123] This can be supported by the preferential coupling of anterior divisions of the MTL with transmodal systems such as the DMN, while posterior segments become increasingly connected to unimodal and sensory-motor systems.[75,84,124,125] Together, our findings echo this unified account of large-scale hierarchical organization of the brain and bridges the dichotomy between different declarative memory systems with reliable correlates to behavioural phenotypes.[77,126,127]

In TLE, previous work has mainly reported impaired memory performance and atypical functional activation patterns in the MTL and beyond.[128,129] Here, we extend this literature by showing a marked reorganization of task-based functional network topographies during episodic and semantic retrieval states. These effects were strongest in bilateral neocortical regions for both memory states, with selective involvement of the MTL region ipsilateral to the seizure focus on episodic states. Consistent with previous work in TLE based on rs-fMRI data,[128] we found neocortical gradient compressions in regions susceptible to TLE-related pathology.[128,130] Other rs-fMRI studies reported simultaneous increases in short- and decreases in long-range connections in temporo-limbic and dorsomedial regions in TLE[128,129]. Moreover, a large body of structural covariance and diffusion MRI studies has shown reduced connectivity of primarily the temporo-limbic diseased epicentre, but also broader brain networks.[35,128] Studying the hippocampus using gradient mapping techniques, we observed an extended episodic, but not semantic principal gradient in TLE patients relative to controls. These hippocampal gradient alterations are, thus, in an opposing direction than the neocortical findings in TLE. An expanded hippocampal gradient in TLE patients is readily interpretable as an increased functional differentiation between anterior and posterior segments, which could reflect reductions in intra-

hippocampal crosstalk along the long axis. In healthy individuals, long axis organization of the hippocampus has previously been suggested to relate to hierarchy of gist *vs* detailed oriented recollection.[127,131, 132] Studies show that the anterior hippocampus acts as a hub, recruiting transmodal neocortical regions for schema construction, evidenced by direct and reciprocal connections between anterior temporal and dorsal and ventromedial prefrontal cortices.[2,133,134] Furthermore, the coactivation of posterior hippocampal and unimodal sensory regions of the neocortex is thought to allow detailed elaborations. [134] In TLE patients, previous work indicate that patients exhibit a reduced capacity for recalling internal details of specific personal episodes but with relatively preserved narrative components,[135] suggesting that hippocampus is essential for recollecting sensory perceptual aspects of previous experiences. As such, the increasing spatial disconnection between anterior and posterior hippocampus observed in our TLE patients could reflect a compensatory retrieval process emphasizing key elements over extraneous details during the episodic memory task[135]. Together, our findings showing the co-occurrence of gradient contractions in neocortical systems with simultaneous gradient expansion of hippocampal subregions may represent a topographic mechanisms of memory reorganization in TLE, compatible with atypical functional segregation in mesiotemporal and neocortical systems, respectively. [86,128] Accordingly, neocortical gradient contractions indicate that transmodal regions are relatively dedifferentiated from unimodal systems, likely stemming from imbalances in both local and distant connectivity. On the other hand, expansions in hippocampal connectivity gradients are likely indicative of reductions in intra-hippocampal connectivity, which would contribute to marked divergences in signalling between anterior and posterior divisions. As such, our findings echo task-fMRI studies showing widespread extratemporal connectivity reductions and intrinsic MTL connectivity increases,[47] and less concordant fMRI

activations between hippocampal long axis and sensory-transmodal gradients in TLE relative to controls.[32] Collectively, the neocortical gradient contraction and hippocampal gradient expansion reported here provide a compact signature of reorganization of declarative memory systems in TLE.

Prior studies have reported atypical microstructure and morphology in TLE in both the MTL as well as adjacent temporo-limbic neocortical systems, [47, 34,35,89,90,136-138] and have suggested that such changes could alter the spatial configuration of functional networks and influence hierarchical organization supporting episodic memory function[87]. In line with previous work,[35,111,128] white matter disruptions and atrophy were observed in our TLE patients, with most marked findings in temporo-limbic regions and hippocampal anterolateral regions ipsilateral to the seizure focus. These findings could reinforce earlier models that propose increased susceptibility of paralimbic regions to structural and functional changes, potentially due to a greater capacity for plasticity and connectivity rearrangement.[88,139] Interestingly, while neocortical structural changes did not reflect functional changes, morphological and microstructural derangements of the hippocampus correlated with episodic memory network reorganization. Previous studies have consistently revealed anterior hippocampal atrophy[140,141] and alterations in tissue microstructure[128] in TLE patients, aligning marked neuronal loss in the anterior hippocampus.[142] Such structural damage in crucial hubs like the hippocampus extend beyond localized effects and can compromise communication within the interconnected memory network[140,143]. The functional association with pathological markers selective to the hippocampus reported herein, therefore, provides specificity, and reinforces the key role of hippocampal integrity in episodic memory network organization.[144,145]

Previous studies have also linked network alterations in TLE patients to cognitive impairment.[47,87,129,136,138] Consistent with a recent study from our group [46], episodic task impairments were marked in TLE, while semantic task performance remained intact. Additionally, neocortical, and hippocampal gradient alterations were found to relate to episodic impairments in TLE. While we acknowledge that our conservative patient inclusion criteria resulted in a relatively modest sample size of 31 TLE patients, the observed memory deficits and functional perturbations were unlikely influenced by neurodegenerative disorders,[146,46] as functional alterations persisted even after controlling for overall measures of cognitive function (*i.e.* MoCA). Behavioural indices of cognitive impairment in TLE were mediated by functional changes, confirming the already well-established literature on episodic impairment in TLE. In line with the hierarchical models of memory,[147,148] our findings lend support to the idea that semantic memory relies more on extra-hippocampal brain regions compared to hippocampal-dependent episodic systems. This is consistent with evidence from TLE patients who underwent anterior lobe resections,[59] and corroborates the controlled semantic cognition framework with the ATL acting as a central semantic hub.[1,16,29] Additionally, our findings are accounted for by the complementary learning systems framework,[149,150] which stipulates that hippocampus-encoded semantic memories undergo a long-term consolidation process[151,152] that renders the memory trace independent from the hippocampus, and thus more resilient to its structural pathology. Likewise, the multiple trace theory[57,58] specifies that episodic memories are consistently mediated by the hippocampus allowing a vivid recollection of context-rich events,[58,153] and each instance of recalling an episodic memory triggers subsequent re-encoding, leading to the formation of multiple traces or engrams, that are mediated by hippocampo-neocortical neuron

ensembles.[57,58,153,154,155,156] In contrast, successful semantic memory recall does not require rich contextual detail and can, therefore, be solely supported by extra-temporal regions.[157]

Collectively, our work presents a novel decomposition analysis of task-fMRI data revealing an atypical and state-dependent reorganization of declarative memory systems in TLE, with a consistent correlation with behavioural impairments and *in vivo* markers of hippocampal pathology. Our findings support a more selective role of the hippocampus in episodic processes, in line with the episodic theory of memory organization [31,158]. This study also provides insights into mechanisms of cognitive impairments in conditions associated with MTL pathology, showing in particular an opposing pattern of decreased neocortical functional differentiation together with increased intra-hippocampal differentiation. As many drug-resistant patients from this cohort undergo resective MTL surgery, future work could investigate whether gradient patterns will help predict post-operative memory deficits, and thus enrich patient-specific clinical decision-making.

Acknowledgements

We would like to thank all the TLE patients and controls who participated in this study.

Funding

D.G.C. is funded in part by the Canada First Research Excellence Fund and Fonds de recherche du Québec, awarded to the Healthy Brains, Healthy Lives (HBHL) initiative at McGill University and Quebec Bio-Imaging Network (QBIN-RBIQ). J.D. is funded by the Natural Sciences and Engineering Research Council of Canada- Post-Doctoral Fellowship (NSERC-

PDF). J.R. is funded by the Canadian Institutes of Health Research (CIHR). S.T. is funded by the Canadian Open Neuroscience Platform (CONP). K.X. is funded by the China Scholarship Council (CSC: 202006070175). R.R.C. is funded by the Fonds de la Recherche du Québec – Santé (FRQS). L.C. is supported by the Berkeley Fellowship jointly awarded by UCL and Gonville and Caius College, Cambridge, and by Brain Research UK (award 14181). A.B. and N.B. are supported by FRQS and CIHR. B.C.B. acknowledges research support from the NSERC Discovery-1304413, CIHR (FDN-154298, PJT-174995), SickKids Foundation (NI17-039), the Helmholtz International BigBrain Analytics and Learning Laboratory (HIBALL), HBHL, Brain Canada, and the Tier-2 Canada Research Chairs program.

Competing interests

The authors report no competing interests.

Supplementary material

Supplementary material is available at *Brain* online.

References

1.   Ralph MAL, Jefferies E, Patterson K, Rogers TT. The neural and computational bases of semantic cognition. *Nature Reviews Neuroscience*. 2017/01/01 2017;18(1):42-55. doi:10.1038/nrn.2016.150
2.   Patterson K, Nestor PJ, Rogers TT. Where do you know what you know? The representation of semantic knowledge in the human brain. *Nature Reviews Neuroscience*. 2007/12/01 2007;8(12):976-987. doi:10.1038/nrn2277
3.   Tulving E. Episodic and semantic memory. *Organization of memory*. 1972;1(381-403):1.
4.   Brown SC, Craik FIM. Encoding and retrieval of information. *The Oxford handbook of memory*. Oxford University Press; 2000:93-107.
5.   Craik FI, Lockhart RS. Levels of processing: A framework for memory research. *Journal of verbal learning and verbal behavior*. 1972;11(6):671-684.

6.   Craik FI, Tulving E. Depth of processing and the retention of words in episodic memory. *Journal of experimental Psychology: general*. 1975;104(3):268.

7.   Bartha L, Brenneis C, Schocke M, et al. Medial temporal lobe activation during semantic language processing: fMRI findings in healthy left-and right-handers. *Cognitive Brain Research*. 2003;17(2):339-346.

8.   Binder JR, Frost JA, Hammeke TA, Cox RW, Rao SM, Prieto T. Human brain language areas identified by functional magnetic resonance imaging. *Journal of Neuroscience*. 1997;17(1):353-362.

9.   Irish M, Vatansever D. Rethinking the episodic-semantic distinction from a gradient perspective. *Current Opinion in Behavioral Sciences*. 2020;32:43-49. doi:https://doi.org/10.1016/j.cobeha.2020.01.016

10.   Irish M. On the interaction between episodic and semantic representations–constructing a unified account of imagination. *The Cambridge handbook of the imagination*. 2020:447-465.

11.   Palacio N, Cardenas F. A systematic review of brain functional connectivity patterns involved in episodic and semantic memory. *Reviews in the Neurosciences*. 2019;30(8):889-902.

12.   Rajah MN, McIntosh AR. Overlap in the functional neural systems involved in semantic and episodic memory retrieval. *Journal of Cognitive Neuroscience*. 2005;17(3):470-482.

13.   Squire LR, Stark CE, Clark RE. The medial temporal lobe. *Annu Rev Neurosci*. 2004;27:279-306.

14.   Menon V, Boyett-Anderson JM, Schatzberg AF, Reiss AL. Relating semantic and episodic memory systems. *Cognitive brain research*. 2002;13(2):261-265.

15.   Barredo J, Öztekin I, Badre D. Ventral fronto-temporal pathway supporting cognitive control of episodic memory retrieval. *Cerebral Cortex*. 2015;25(4):1004-1019.

16.   Jefferies E. The neural basis of semantic cognition: converging evidence from neuropsychology, neuroimaging and TMS. *Cortex*. 2013;49(3):611-625.

17.   Whitney C, Jefferies E, Kircher T. Heterogeneity of the left temporal lobe in semantic representation and control: priming multiple versus single meanings of ambiguous words. *Cerebral cortex*. 2011;21(4):831-844.

18.   Whitney C, Kirk M, O'Sullivan J, Lambon Ralph MA, Jefferies E. The neural organization of semantic control: TMS evidence for a distributed network in left inferior frontal and posterior middle temporal gyrus. *Cerebral cortex*. 2011;21(5):1066-1075.

19.   Davey J, Cornelissen PL, Thompson HE, et al. Automatic and controlled semantic retrieval: TMS reveals distinct contributions of posterior middle temporal gyrus and angular gyrus. *Journal of Neuroscience*. 2015;35(46):15230-15239.

20.   Ranganath C, Ritchey M. Two cortical systems for memory-guided behaviour. *Nature reviews neuroscience*. 2012;13(10):713-726.

21.   Whitney C, Kirk M, O'Sullivan J, Lambon Ralph MA, Jefferies E. Executive semantic processing is underpinned by a large-scale neural network: revealing the contribution of left prefrontal, posterior temporal, and parietal cortex to controlled retrieval and selection using TMS. *Journal of cognitive neuroscience*. 2012;24(1):133-147.

22.   Ramanan S, Strikwerda-Brown C, Mothakunnel A, Hodges JR, Piguet O, Irish M. Fronto-parietal contributions to episodic retrieval—Evidence from neurodegenerative disorders. *Learning & memory*. 2019;26(7):262-271.

23.   Vatansever D, Smallwood J, Jefferies E. Varying demands for cognitive control reveals shared neural processes supporting semantic and episodic memory retrieval. *Nature communications*. 2021;12(1):2134.

24.   Zhukovsky P, Coughlan G, Buckley R, Grady C, Voineskos AN. Connectivity between default mode and frontoparietal networks mediates the association between global amyloid‐β and episodic memory. *Human Brain Mapping*. 2023;44(3):1147-1157.

25.   Kim H. Default network activation during episodic and semantic memory retrieval: a selective meta-analytic comparison. *Neuropsychologia*. 2016;80:35-46.

26.   Yarkoni T, Poldrack RA, Nichols TE, Van Essen DC, Wager TD. Large-scale automated synthesis of human functional neuroimaging data. *Nature methods*. 2011;8(8):665-670.

27.   Burianova H, Grady CL. Common and unique neural activations in autobiographical, episodic, and semantic retrieval. *Journal of cognitive neuroscience*. 2007;19(9):1520-1534.

28.   Chiou R, Humphreys GF, Jung J, Ralph MAL. Controlled semantic cognition relies upon dynamic and flexible interactions between the executive 'semantic control'and hub-and-spoke 'semantic representation'systems. *cortex*. 2018;103:100-116.

29.   Jefferies E, Wang X. Semantic cognition: semantic memory and semantic control. *Oxford Research Encyclopedia of Psychology*. 2021.

30.   Patterson K, Ralph MAL. The hub-and-spoke hypothesis of semantic memory. *Neurobiology of language*. Elsevier; 2016:765-775.

31.   Moscovitch M, Cabeza R, Winocur G, Nadel L. Episodic memory and beyond: the hippocampus and neocortex in transformation. *Annual review of psychology*. 2016;67:105-134.

32.   Li Q, Tavakol S, Royer J, et al. Atypical neural topographies underpin dysfunctional pattern separation in temporal lobe epilepsy. *Brain*. 2021;144(8):2486-2498.

33.   Giovagnoli AR, Avanzini G. Learning and memory impairment in patients with temporal lobe epilepsy: relation to the presence, type, and location of brain lesion. *Epilepsia*. 1999;40(7):904-911.

34.   Bernasconi N, Duchesne S, Janke A, Lerch J, Collins DL, Bernasconi A. Whole-brain voxel-based statistical analysis of gray matter and white matter in temporal lobe epilepsy. *Neuroimage*. 2004;23(2):717-723.

35.   Bernhardt BC, Bernasconi A, Liu M, et al. The spectrum of structural and functional imaging abnormalities in temporal lobe epilepsy. *Annals of neurology*. 2016;80(1):142-153.

36.   Blümcke I, Thom M, Aronica E, et al. International consensus classification of hippocampal sclerosis in temporal lobe epilepsy: a Task Force report from the ILAE Commission on Diagnostic Methods. *Epilepsia*. 2013;54(7):1315-1329.

37.   Thom M, Eriksson S, Martinian L, et al. Temporal lobe sclerosis associated with hippocampal sclerosis in temporal lobe epilepsy: neuropathological features. *Journal of Neuropathology & Experimental Neurology*. 2009;68(8):928-938.

38.   Barrett Jones S, A. Miller L, Kleitman S, Nikpour A, Lah S. Semantic and episodic memory in adults with temporal lobe epilepsy. *Applied Neuropsychology: Adult*. 2022;29(6):1352-1361.

39.   Helmstaedter C. Effects of chronic epilepsy on declarative memory systems. *Progress in brain research*. 2002;135:439-453.

40.   Viskontas IV, McAndrews MP, Moscovitch M. Remote episodic memory deficits in patients with unilateral temporal lobe epilepsy and excisions. *Journal of Neuroscience*. 2000;20(15):5853-5857.

41.   Messas CS, Mansur LL, Castro LHM. Semantic memory impairment in temporal lobe epilepsy associated with hippocampal sclerosis. *Epilepsy & Behavior*. 2008;12(2):311-316.

42.   McDonald CR, Leyden KM, Hagler DJ, et al. White matter microstructure complements morphometry for predicting verbal memory in epilepsy. *Cortex*. 2014;58:139-150.

43.   Rzezak P, Lima EM, Gargaro AC, et al. Everyday memory impairment in patients with temporal lobe epilepsy caused by hippocampal sclerosis. *Epilepsy & Behavior*. 2017;69:31-36.

44.   Saghafi S, Ferguson L, Hogue O, Gales JM, Prayson R, Busch RM. Histopathologic subtype of hippocampal sclerosis and episodic memory performance before and after temporal lobectomy for epilepsy. *Epilepsia*. 2018;59(4):825-833.

45.   Silvia O, Patricia Ss, Damián C, et al. Mesial temporal lobe epilepsy and hippocampal sclerosis: cognitive function assessment in Hispanic patients. *Epilepsy & Behavior*. 2003;4(6):717-722.

46.   Tavakol S, Kebets V, Royer J, et al. DIFFERENTIAL MEMORY IMPAIRMENT ACROSS RELATIONAL DOMAINS IN TEMPORAL LOBE EPILEPSY. *bioRxiv*. 2022:2022.11. 01.514752.

47.   Fleury M, Buck S, Binding LP, et al. Episodic memory network connectivity in temporal lobe epilepsy. *Epilepsia*. 2022;63(10):2597-2622.

48.   Sidhu MK, Stretton J, Winston GP, et al. A functional magnetic resonance imaging study mapping the episodic memory encoding network in temporal lobe epilepsy. *Brain*. 2013;136(6):1868-1888.

49.   Tramoni E, Felician O, Barbeau EJ, et al. Long-term consolidation of declarative memory: insight from temporal lobe epilepsy. *Brain*. 2011;134(3):816-831.

50.   Boucher O, Dagenais E, Bouthillier A, Nguyen DK, Rouleau I. Different effects of anterior temporal lobectomy and selective amygdalohippocampectomy on verbal memory performance of patients with epilepsy. *Epilepsy Behav*. Nov 2015;52(Pt A):230-5. doi:10.1016/j.yebeh.2015.09.012

51.   Helmstaedter C, Elger C, Hufnagel A, Zentner J, Schramm J. Different effects of left anterior temporal lobectomy, selective amygdalohippocampectomy, and temporal cortical lesionectomy on verbal learning, memory, and recognition. *Journal of Epilepsy*. 1996;9(1):39-45.

52.   Tanriverdi T, Dudley RW, Hasan A, et al. Memory outcome after temporal lobe epilepsy surgery: corticoamygdalohippocampectomy versus selective amygdalohippocampectomy. *J Neurosurg*. Dec 2010;113(6):1164-75. doi:10.3171/2009.10.Jns09677

53.   Fliessbach K, Witt J-A, Packheiser J, von Lehe M, Elger CE, Helmstaedter C. Depth-of-processing effects on memory encoding after selective amygdalohippocampectomy. *Behavioural brain research*. 2011;216(1):402-407.

54.   Mansouri A, Fallah A, McAndrews MP, et al. Neurocognitive and Seizure Outcomes of Selective Amygdalohippocampectomy versus Anterior Temporal Lobectomy for Mesial Temporal Lobe Epilepsy. *Epilepsy Res Treat*. 2014;2014:306382. doi:10.1155/2014/306382

55.   Jung JY, Rice GE, Lambon Ralph MA. The neural bases of resilient semantic system: evidence of variable neuro-displacement in cognitive systems. *Brain Structure and Function*. 2021;226:1585-1599.

56.   Rice GE, Lambon Ralph MA, Hoffman P. The Roles of Left Versus Right Anterior Temporal Lobes in Conceptual Knowledge: An ALE Meta-analysis of 97 Functional Neuroimaging Studies. *Cerebral Cortex*. 2015;25(11):4374-4391. doi:10.1093/cercor/bhv024

57.   Nadel L, Samsonovich A, Ryan L, Moscovitch M. Multiple trace theory of human memory: computational, neuroimaging, and neuropsychological results. *Hippocampus*. 2000;10(4):352-68. doi:10.1002/1098-1063(2000)10:4<352::Aid-hipo2>3.0.Co;2-d

58.   Moscovitch M, Rosenbaum RS, Gilboa A, et al. Functional neuroanatomy of remote episodic, semantic and spatial memory: a unified account based on multiple trace theory. *Journal of anatomy*. 2005;207(1):35-66.

59.   Lambon Ralph MA, Ehsan S, Baker GA, Rogers TT. Semantic memory is impaired in patients with unilateral anterior temporal lobe resection for temporal lobe epilepsy. *Brain*. Jan 2012;135(Pt 1):242-58. doi:10.1093/brain/awr325

60.   Bell BD, Hermann BP, Woodard AR, et al. Object naming and semantic knowledge in temporal lobe epilepsy. *Neuropsychology*. Oct 2001;15(4):434-43. doi:10.1037//0894-4105.15.4.434

61.   Giovagnoli AR, Erbetta A, Villani F, Avanzini G. Semantic memory in partial epilepsy: verbal and non-verbal deficits and neuroanatomical relationships. *Neuropsychologia*. 2005;43(10):1482-92. doi:10.1016/j.neuropsychologia.2004.12.010

62.   Poch C, Toledano R, García-Morales I, Alemán-Gómez Y, Gil-Nagel A, Campo P. Contributions of left and right anterior temporal lobes to semantic cognition: Evidence from patients with small temporopolar lesions. *Neuropsychologia*. Feb 12 2021;152:107738. doi:10.1016/j.neuropsychologia.2020.107738

63.   Glosser G, Salvucci A, Chiaravalloti N. Naming and recognizing famous faces in temporal lobe epilepsy. *Neurology*. 2003;61(1):81-86.

64.   Martin RC, Sawrie SM, Roth DL, et al. Individual memory change after anterior temporal lobectomy: a base rate analysis using regression□based outcome methodology. *Epilepsia*. 1998;39(10):1075-1082.

65.   Margulies DS, Ghosh SS, Goulas A, et al. Situating the default-mode network along a principal gradient of macroscale cortical organization. *Proceedings of the National Academy of Sciences*. 2016;113(44):12574-12579.

66.   Mesulam MM. From sensation to cognition. *Brain*. Jun 1998;121 (Pt 6):1013-52. doi:10.1093/brain/121.6.1013

67.   Paquola C, Amunts K, Evans A, Smallwood J, Bernhardt B. Closing the mechanistic gap: the value of microarchitecture in understanding cognitive networks. *Trends Cogn Sci*. Oct 2022;26(10):873-886. doi:10.1016/j.tics.2022.07.001

68.   Smallwood J, Bernhardt BC, Leech R, Bzdok D, Jefferies E, Margulies DS. The default mode network in cognition: a topographical perspective. *Nature reviews neuroscience*. 2021;22(8):503-513.

69.   Murray JD, Bernacchia A, Freedman DJ, et al. A hierarchy of intrinsic timescales across primate cortex. *Nature neuroscience*. 2014;17(12):1661-1663.

70.   Bernhardt BC, Smallwood J, Keilholz S, Margulies DS. Gradients in brain organization. Elsevier; 2022. p. 118987.

71.   Huntenburg JM, Bazin P-L, Margulies DS. Large-scale gradients in human cortical organization. *Trends in cognitive sciences*. 2018;22(1):21-31.

72.   Paquola C, Vos De Wael R, Wagstyl K, et al. Microstructural and functional gradients are increasingly dissociated in transmodal cortices. *PLoS biology*. 2019;17(5):e3000284.

73.   Vázquez-Rodríguez B, Suárez LE, Markello RD, et al. Gradients of structure–function tethering across neocortex. *Proceedings of the National Academy of Sciences*. 2019;116(42):21219-21227.

74.   Borne L, Tian Y, Lupton MK, et al. Functional re-organization of hippocampal-cortical gradients during naturalistic memory processes. *NeuroImage*. 2023;271:119996.

75.   Strange BA, Witter MP, Lein ES, Moser EI. Functional organization of the hippocampal longitudinal axis. *Nature Reviews Neuroscience*. 2014;15(10):655-669.

76.   Vos de Wael R, Larivière S, Caldairou B, et al. Anatomical and microstructural determinants of hippocampal subfield functional connectome embedding. *Proc Natl Acad Sci U S A*. Oct 2 2018;115(40):10154-10159. doi:10.1073/pnas.1803667115

77.   Genon S, Bernhardt BC, La Joie R, Amunts K, Eickhoff SB. The many dimensions of human hippocampal organization and (dys) function. *Trends in neurosciences*. 2021;44(12):977-989.

78.   Bajada CJ, Jackson RL, Haroon HA, et al. A graded tractographic parcellation of the temporal lobe. *Neuroimage*. Jul 15 2017;155:503-512. doi:10.1016/j.neuroimage.2017.04.016

79.   Cerliani L, Thomas RM, Jbabdi S, et al. Probabilistic tractography recovers a rostrocaudal trajectory of connectivity variability in the human insular cortex. *Human brain mapping*. 2012;33(9):2005-2034.

80.   Guell X, Schmahmann JD, Gabrieli J, Ghosh SS. Functional gradients of the cerebellum. *Elife*. Aug 14 2018;7doi:10.7554/eLife.36652

81.   Haak KV, Marquand AF, Beckmann CF. Connectopic mapping with resting-state fMRI. *Neuroimage*. 2018;170:83-94.

82.   Huntenburg JM, Bazin PL, Goulas A, Tardif CL, Villringer A, Margulies DS. A Systematic Relationship Between Functional Connectivity and Intracortical Myelin in the Human Cerebral Cortex. *Cereb Cortex*. Feb 1 2017;27(2):981-997. doi:10.1093/cercor/bhx030

83.   Paquola C, Bethlehem RA, Seidlitz J, et al. Shifts in myeloarchitecture characterise adolescent development of cortical gradients. *Elife*. Nov 14 2019;8doi:10.7554/eLife.50482

84.   Przeździk I, Faber M, Fernández G, Beckmann CF, Haak KV. The functional organisation of the hippocampus along its long axis is gradual and predicts recollection. *Cortex*. Oct 2019;119:324-335. doi:10.1016/j.cortex.2019.04.015

85.   Tian Y, Zalesky A. Characterizing the functional connectivity diversity of the insula cortex: Subregions, diversity curves and behavior. *Neuroimage*. Dec 2018;183:716-733. doi:10.1016/j.neuroimage.2018.08.055

86.   Lucas A, Mouchtaris S, Cornblath EJ, et al. Subcortical functional connectivity gradients in temporal lobe epilepsy. *NeuroImage: Clinical*. 2023;38:103418.

87.   Royer J, Larivière S, Rodriguez-Cruces R, et al. Cortical microstructural gradients capture memory network reorganization in temporal lobe epilepsy. *Brain*. Apr 21 2023;doi:10.1093/brain/awad125

88.   Park BY, Larivière S, Rodríguez-Cruces R, et al. Topographic divergence of atypical cortical asymmetry and atrophy patterns in temporal lobe epilepsy. *Brain*. May 24 2022;145(4):1285-1298. doi:10.1093/brain/awab417

89.   Hatton SN, Huynh KH, Bonilha L, et al. White matter abnormalities across different epilepsy syndromes in adults: an ENIGMA-Epilepsy study. *Brain*. 2020;143(8):2454-2473.

90.   Bernhardt BC, Bernasconi N, Concha L, Bernasconi A. Cortical thickness analysis in temporal lobe epilepsy: reproducibility and relation to outcome. *Neurology*. 2010;74(22):1776-1784.

91.   Bernhardt BC, Fadaie F, de Wael RV, et al. Preferential susceptibility of limbic cortices to microstructural damage in temporal lobe epilepsy: A quantitative T1 mapping study. *Neuroimage*. 2018;182:294-303.

92.   DeKraker J, Haast RAM, Yousif MD, et al. Automated hippocampal unfolding for morphometry and subfield segmentation with HippUnfold. *eLife*. 2022/12// 2022;11doi:10.7554/eLife.77945

93.   J Jr E. Outocme with respect to epileptic seizures. *Surgical treatment of the epilepsies*. 1993:609-621.

94.   Han L, Kashyap AL, Finin TW, Mayfield J, Weese J. UMBC_EBIQUITY-CORE: Semantic Textual Similarity Systems. 2013:

95.   Nasreddine ZS, Phillips NA, Bédirian V, et al. "The Montreal Cognitive Assessment, MoCA: A brief screening tool for mild cognitive impairment": Corrigendum. *Journal of the American Geriatrics Society*. 2019;67(9):1991-1991. doi:10.1111/jgs.15925

96.   Cruces RR, Royer J, Herholz P, et al. Micapipe: A pipeline for multimodal neuroimaging and connectome analysis. *Neuroimage*. Nov 2022;263:119612. doi:10.1016/j.neuroimage.2022.119612

97.   Basser PJ, Mattiello J, LeBihan D. Estimation of the effective self-diffusion tensor from the NMR spin echo. *J Magn Reson B*. Mar 1994;103(3):247-54. doi:10.1006/jmrb.1994.1037

98.   Veraart J, Sijbers J, Sunaert S, Leemans A, Jeurissen B. Weighted linear least squares estimation of diffusion MRI parameters: strengths, limitations, and pitfalls. *Neuroimage*. Nov 1 2013;81:335-346. doi:10.1016/j.neuroimage.2013.05.028

99.   Jenkinson M, Beckmann CF, Behrens TE, Woolrich MW, Smith SM. FSL. *Neuroimage*. Aug 15 2012;62(2):782-90. doi:10.1016/j.neuroimage.2011.09.015

100.   Avants BB, Tustison NJ, Song G, Cook PA, Klein A, Gee JC. A reproducible evaluation of ANTs similarity metric performance in brain image registration. *Neuroimage*. Feb 1 2011;54(3):2033-44. doi:10.1016/j.neuroimage.2010.09.025

101.   Cox RW. AFNI: software for analysis and visualization of functional magnetic resonance neuroimages. *Comput Biomed Res*. Jun 1996;29(3):162-73. doi:10.1006/cbmr.1996.0014

102.   Salimi-Khorshidi G, Douaud G, Beckmann CF, Glasser MF, Griffanti L, Smith SM. Automatic denoising of functional MRI data: combining independent component analysis and hierarchical fusion of classifiers. *Neuroimage*. Apr 15 2014;90:449-68. doi:10.1016/j.neuroimage.2013.11.046

103.   Marcus DS, Harms MP, Snyder AZ, et al. Human Connectome Project informatics: quality control, database services, and data visualization. *Neuroimage*. Oct 15 2013;80:202-19. doi:10.1016/j.neuroimage.2013.05.077

104.   Coifman RR, Lafon S. Diffusion maps. *Applied and computational harmonic analysis*. 2006;21(1):5-30.

105.   Vos de Wael R, Benkarim O, Paquola C, et al. BrainSpace: a toolbox for the analysis of macroscale gradients in neuroimaging and connectomics datasets. *Commun Biol*. Mar 5 2020;3(1):103. doi:10.1038/s42003-020-0794-7

106.   Van Essen DC, Ugurbil K, Auerbach E, et al. The Human Connectome Project: a data acquisition perspective. *Neuroimage*. Oct 1 2012;62(4):2222-31. doi:10.1016/j.neuroimage.2012.02.018

107.   Langs G, Golland P, Ghosh SS. Predicting activation across individuals with resting-state functional connectivity based multi-atlas label fusion. Springer; 2015:313-320.

108.   Larivière S, Bayrak Ş, Vos de Wael R, et al. BrainStat: A toolbox for brain-wide statistics and multimodal feature associations. *Neuroimage*. Feb 1 2023;266:119807. doi:10.1016/j.neuroimage.2022.119807

109.   Worsley KJ, Taylor J, Carbonell F, et al. A Matlab toolbox for the statistical analysis of univariate and multivariate surface and volumetric data using linear mixed effects models and random field theory. 2009:S102.

110.   Kim H, Bernhardt BC, Kulaga-Yoskovitz J, Caldairou B, Bernasconi A, Bernasconi N. Multivariate hippocampal subfield analysis of local MRI intensity and volume: application to temporal lobe epilepsy. Springer; 2014:170-178.

111.   Liu M, Bernhardt BC, Hong S-J, Caldairou B, Bernasconi A, Bernasconi N. The superficial white matter in temporal lobe epilepsy: a key link between structural and functional network disruptions. *Brain*. 2016;139(9):2431-2440.

112.   Royer J, Rodríguez-Cruces R, Tavakol S, et al. An open MRI dataset for multiscale neuroscience. *Scientific Data*. 2022;9(1):569.

113.   Wagstyl K, Ronan L, Goodyer IM, Fletcher PC. Cortical thickness gradients in structural hierarchies. *Neuroimage*. 2015;111:241-250.

114.   Vos de Wael R, Royer J, Tavakol S, et al. Structural connectivity gradients of the temporal lobe serve as multiscale axes of brain organization and cortical evolution. *Cerebral cortex*. 2021;31(11):5151-5164.

115.   Buckner RL, Krienen FM. The evolution of distributed association networks in the human brain. *Trends in cognitive sciences*. 2013;17(12):648-665.

116.   Smallwood J. Distinguishing how from why the mind wanders: a process–occurrence framework for self-generated mental activity. *Psychological bulletin*. 2013;139(3):519.

117.   Smallwood J, Schooler JW. The science of mind wandering: Empirically navigating the stream of consciousness. *Annual review of psychology*. 2015;66:487-518.

118.   Ho NSP, Poerio G, Konu D, et al. Facing up to the wandering mind: Patterns of off-task laboratory thought are associated with stronger neural recruitment of right fusiform cortex while processing facial stimuli. *Neuroimage*. 2020;214:116765.

119.   Smallwood J, Turnbull A, Wang H-t, et al. The neural correlates of ongoing conscious thought. *Iscience*. 2021;24(3)

120.   Kelley WM, Macrae CN, Wyland CL, Caglar S, Inati S, Heatherton TF. Finding the self? An event-related fMRI study. *Journal of cognitive neuroscience*. 2002;14(5):785-794.

121.   Wang X, Margulies DS, Smallwood J, Jefferies E. A gradient from long-term memory to novel cognition: Transitions through default mode and executive cortex. *NeuroImage*. 2020;220:117074.

122.   Mok RM, Love BC. A multilevel account of hippocampal function in spatial and concept learning: Bridging models of behavior and neural assemblies. *Science Advances*. 2023;9(29):eade6903.

123.   Koster R, Chadwick MJ, Chen Y, et al. Big-loop recurrence within the hippocampal system supports integration of information across episodes. *Neuron*. 2018;99(6):1342-1354. e6.

124.   Paquola C, Benkarim O, DeKraker J, et al. Convergence of cortical types and functional motifs in the human mesiotemporal lobe. *elife*. 2020;9:e60673.

125.   Chase HW, Clos M, Dibble S, et al. Evidence for an anterior–posterior differentiation in the human hippocampal formation revealed by meta-analytic parcellation of fMRI coordinate maps: Focus on the subiculum. *NeuroImage*. 2015;113:44-60.

126.   Sydnor VJ, Larsen B, Bassett DS, et al. Neurodevelopment of the association cortices: Patterns, mechanisms, and implications for psychopathology. *Neuron*. 2021;109(18):2820-2846.

127.   Poppenk J, Evensmoen HR, Moscovitch M, Nadel L. Long-axis specialization of the human hippocampus. *Trends in cognitive sciences*. 2013;17(5):230-240.

128.   Larivière S, Weng Y, Vos de Wael R, et al. Functional connectome contractions in temporal lobe epilepsy: Microstructural underpinnings and predictors of surgical outcome. *Epilepsia*. 2020;61(6):1221-1233.

129.   Xie K, Royer J, Lariviere S, et al. Atypical connectome topography and signal flow in temporal lobe epilepsy. *bioRxiv*. 2023:2023.05. 23.541934.

130.   Morgan VL, Rogers BP, Sonmezturk HH, Gore JC, Abou☐Khalil B. Cross hippocampal influence in mesial temporal lobe epilepsy measured with high temporal resolution functional magnetic resonance imaging. *Epilepsia*. 2011;52(9):1741-1749.

131.   Gutchess AH, Schacter DL. The neural correlates of gist-based true and false recognition. *Neuroimage*. 2012;59(4):3418-3426.

132.   Poppenk J, Moscovitch M. A hippocampal marker of recollection memory ability among healthy young adults: contributions of posterior and anterior segments. *Neuron*. 2011;72(6):931-937.

133.   Van Kesteren MT, Rijpkema M, Ruiter DJ, Fernández G. Retrieval of associative information congruent with prior knowledge is related to increased medial prefrontal activity and connectivity. *Journal of Neuroscience*. 2010;30(47):15888-15894.

134.   McCormick C, St-Laurent M, Ty A, Valiante TA, McAndrews MP. Functional and effective hippocampal–neocortical connectivity during construction and elaboration of autobiographical memory retrieval. *Cerebral cortex*. 2015;25(5):1297-1305.

135.   St-Laurent M, Moscovitch M, Levine B, McAndrews MP. Determinants of autobiographical memory in patients with unilateral temporal lobe epilepsy or excisions. *Neuropsychologia*. 2009;47(11):2211-2221.

136.   Rodríguez-Cruces R, Bernhardt BC, Concha L. Multidimensional associations between cognition and connectome organization in temporal lobe epilepsy. *NeuroImage*. 2020;213:116706.

137.   Dabbs K, Jones J, Seidenberg M, Hermann B. Neuroanatomical correlates of cognitive phenotypes in temporal lobe epilepsy. *Epilepsy & Behavior*. 2009;15(4):445-451.

138.   Doucet G, Osipowicz K, Sharan A, Sperling MR, Tracy JI. Extratemporal functional connectivity impairments at rest are related to memory performance in mesial temporal epilepsy. *Human brain mapping*. 2013;34(9):2202-2216.

139.   García-Cabezas MÁ, Zikopoulos B, Barbas H. The Structural Model: a theory linking connections, plasticity, pathology, development and evolution of the cerebral cortex. *Brain Structure and Function*. 2019;224(3):985-1008.

140.   Barnett AJ, Park MTM, Pipitone J, Chakravarty MM, McAndrews MP. Functional and structural correlates of memory in patients with mesial temporal lobe epilepsy. *Frontiers in neurology*. 2015;6:103.

141.   Bernasconi N, Bernasconi A, Caramanos Z, Antel SB, Andermann F, Arnold DL. Mesial temporal damage in temporal lobe epilepsy: a volumetric MRI study of the hippocampus, amygdala and parahippocampal region. *Brain*. 2003;126(2):462-469. doi:10.1093/brain/awg034

142.   Dam AM. Epilepsy and neuron loss in the hippocampus. *Epilepsia*. 1980;21(6):617-629.

143.   Das SR, Mechanic-Hamilton D, Korczykowski M, et al. Structure specific analysis of the hippocampus in temporal lobe epilepsy. *Hippocampus*. Jun 2009;19(6):517-25. doi:10.1002/hipo.20620

144.   Goldstein FC, Mao H, Wang L, Ni C, Lah JJ, Levey AI. White matter integrity and episodic memory performance in mild cognitive impairment: a diffusion tensor imaging study. *Brain Imaging and Behavior*. 2009;3:132-141.

145.   Witt J-A, Coras R, Schramm J, et al. Relevance of hippocampal integrity for memory outcome after surgical treatment of mesial temporal lobe epilepsy. *Journal of neurology*. 2015;262:2214-2224.

146.   Tromp D, Dufour A, Lithfous S, Pebayle T, Després O. Episodic memory in normal aging and Alzheimer disease: Insights from imaging and behavioral studies. *Ageing research reviews*. 2015;24:232-262.

147.   Vargha-Khadem F, Gadian DG, Watkins KE, Connelly A, Van Paesschen W, Mishkin M. Differential effects of early hippocampal pathology on episodic and semantic memory. *Science*. 1997;277(5324):376-380.

148.   Mishkin M, Suzuki WA, Gadian DG, Vargha–Khadem F. Hierarchical organization of cognitive memory. *Philosophical Transactions of the Royal Society of London Series B: Biological Sciences*. 1997;352(1360):1461-1467.

149.   O'Reilly RC, Norman KA. Hippocampal and neocortical contributions to memory: Advances in the complementary learning systems framework. *Trends in cognitive sciences*. 2002;6(12):505-510.

150.   McClelland JL, McNaughton BL, O'Reilly RC. Why there are complementary learning systems in the hippocampus and neocortex: insights from the successes and failures of connectionist models of learning and memory. *Psychological review*. 1995;102(3):419.

151.   Klinzing JG, Niethard N, Born J. Mechanisms of systems memory consolidation during sleep. *Nature neuroscience*. 2019;22(10):1598-1610.

152.   Squire LR, Genzel L, Wixted JT, Morris RG. Memory consolidation. *Cold Spring Harbor perspectives in biology*. 2015;7(8):a021766.

153.   Moscovitch M, Westmacott R, Gilboa A, et al. Hippocampal complex contribution to retention and retrieval of recent and remote episodic and semantic memories: Evidence from behavioral and neuroimaging studies of healthy and brain-damaged people. *Dynamic cognitive processes*. 2005:333-380.

154.   Eldridge LL, Knowlton BJ, Furmanski CS, Bookheimer SY, Engel SA. Remembering episodes: a selective role for the hippocampus during retrieval. *Nature neuroscience*. 2000;3(11):1149-1152.

155.   Moscovitch DA, McAndrews MP. Material-specific deficits in "remembering" in patients with unilateral temporal lobe epilepsy and excisions. *Neuropsychologia*. 2002;40(8):1335-1342.

156.   Yonelinas AP. The nature of recollection and familiarity: A review of 30 years of research. *Journal of memory and language*. 2002;46(3):441-517.

157.   Rosenbaum RS, Priselac S, Köhler S, et al. Remote spatial memory in an amnesic person with extensive bilateral hippocampal lesions. *Nature neuroscience*. 2000;3(10):1044-1048.

158.   Burgess N, Maguire EA, O'Keefe J. The human hippocampus and spatial and episodic memory. *Neuron*. 2002;35(4):625-641.

Supplementary Materials

Differential reorganization of episodic and semantic memory systems in epilepsy-related mesiotemporal pathology

Donna Gift Cabalo[1,2], Jordan DeKraker[1,2], Jessica Royer[1,2,3], Ke Xie[1,2], Shahin Tavakol[1,2], Raúl Rodríguez-Cruces[1,2], Andrea Bernasconi[2,4], Neda Bernasconi[2,4], Alexander Weil[5], Raluca Pana[2], Birgit Frauscher[2,3], Lorenzo Caciagli[6], Elizabeth Jefferies[7], Jonathan Smallwood[8], Boris C. Bernhardt[1,2]

Author affiliations:

1 Multimodal Imaging and Connectome Analysis Laboratory, McConnell Brain Imaging Centre, Montreal Neurological Institute, McGill University, Montreal, QC, Canada

2 Montreal Neurological Institute and Hospital, McGill University, Montreal, QC, Canada;

3 Analytical Neurophysiology Laboratory, Montreal Neurological Institute, McGill University, Montreal, Quebec, Canada;

4 Neuroimaging of Epilepsy Laboratory, McConnell Brain Imaging Centre, Montreal Neurological Institute and Hospital, McGill University, Montreal, QC, Canada;

5CHU St Justine, Montreal, QC;

6 Department of Bioengineering, University of Pennsylvania, Philadelphia, USA;

7 University of York, North Yorkshire, United Kingdom;

8 Queens University, Kingston, Ontario, Canada

Correspondence to: Donna Gift Cabalo or Boris Bernhardt

3801 Rue University, Montréal, QC, Canada, H3A 2B4

donna.cabalo@mail.mcgill.ca | boris.bernhardt@mcgill.ca

ID	Age	Sex	HD	Laterality	Onset	Duration (years)	Febrile convulsions	Engel outcome
P01	54	M	R	L	temporal	37	0	n/a
P02	26	F	R	L	temporal	10	0	1a
P03	34	M	R	L	temporal	4	0	1a
P04	20	F	R	L	temporal	5	0	1a
P05	25	F	R	L	temporal	12	0	1b
P06	33	F	R	L	temporal	6	0	1a
P07	57	M	R	L	temporal	39	0	1a
P08	41	F	R	R	temporal	25	2	1b
P09	31	M	R	L	temporal	14	1	n/a
P10	52	M	R	L	temporal	3	0	n/a
P11	55	F	R	L	temporal	13	1	1a
P12	26	F	L	L	temporal	9	1	n/a
P13	25	M	R	R	temporal	23	0	n/a
P14	22	F	R	R	temporal	12	0	2b
P15	53	F	R	L	temporal	45	0	n/a
P16	24	M	R	L	temporal	6	0	n/a
P17	38	M	R	L	temporal	12	1	n/a
P18	43	M	R	R	temporal	13	2	1a
P19	32	M	R	R	temporal	14	0	1a
P20	41	M	R	L	temporal	27	0	2b
P21	27	F	R	L	temporal	12	2	n/a

P22	25	F	R	R	temporal	8	0	3a
P23	28	M	R	L	temporal	1	0	n/a
P24	29	M	R	R	temporal	7	0	n/a
P25	27	F	R	L	temporal	14	0	n/a
P26	39	F	R	R	temporal	27	0	1a
P27	36	F	R	L	temporal	9	2	n/a
P28	18	F	R	L	temporal	4	0	n/a
P29	28	F	R	L	temporal	5	0	n/a
P30	29	F	R	R	temporal	11	0	n/a
P31	56	M	R	R	temporal	14	0	n/a

Table S1. Patient's demographics and clinical features. Abbreviations: L=left; R=right, HD= Handedness

	a	b	c'	c	ab
Neocortex					
Episodic G1$_{ipsi}$	0.55(0.12) ***	0.23(0.17)	0.77(0.24) **	0.89(0.21) ***	0.12(0.09)
Episodic G1$_{contra}$	0.52(0.13) ***	0.21(0.15)	0.78(0.23) **	0.89(0.21) **	0.1(0.08)
Semantic G1$_{ipsi}$	0.54(0.16) ***	-0.07(0.11)	0.34(0.28)	0.31(0.25)	-0.06(0.05)
Semantic G1$_{contra}$	0.51(0.13) ***	-0.13(0.1)	0.38(0.28)	0.31(0.25)	-0.07(0.06)
Hippocampus					
Episodic G1$_{ipsi}$	0.78(0.19) ***	0.17(0.14)	0.75(0.2) **	0.89(0.21) ***	0.13(0.12)
Neocortex +	**Hippocampus**				
Episodic G1	0.66(0.1) ***	0.48(0.23) *	0.57(0.23) *	0.89(0.21) ***	0.32(0.16) *

Table S2. **Individual mediation analysis for Episodic and Semantic memory G1**. The mean of significant regions showing significant episodic and semantic gradient scores were used as mediators for each analysis. **Abbreviations:** G1=Gradient1, Ipsi=ipsilateral, Contra=contralateral, a = Group ~ Gradient scores (standard errors in parenthesis), b= Gradient scores ~ Task performance, c'= Group ~ Task performance, c'= Group ~ Task performance (residual), ab= mediation effect

Figure S1. **A,B.** Episodic and semantic neocortical and hippocampal connectivity gradients before alignment to normative gradients derived from resting-state fMRI. **Abbreviations**: G1 =Gradient 1,G2= Gradient 2,G3 = Gradient 3, Ipsi= ipsilateral, Contra=contralateral, TLE=temporal lobe epilepsy

Figure S2. **(i,ii).** Episodic/Semantic G2 describing a visual-to-somatomotor topography and explained 10/9% of variance. No G2 contractions were observed in semantic memory, contrasting the TLE-related reorganization observed in the anterior temporal, insular, parietal, and bilateral frontal regions for episodic memory. Significant *p*-values were highlighted after correcting for multiple comparisons ($p_{FWE} < 0.05$, *black outlines*). In both memory states, hippocampal G2 reflected a proximal-distal representation and explained 10/10% of functional connectome variance. While no significant differences were observed in semantic hippocampal G2, a distinct reorganization in the anterior hippocampal episodic G2 was observed. **(iii, iv).** Episodic/Semantic G3, dominated by networks involved in the declarative memory tasks explained 8% of the total episodic and semantic variance. G3 suppression in TLE for episodic memory state was recorded in prefrontal cortices, ipsilateral to seizure focus. **Abbreviations**: G2= Gradient 2, G3 = Gradient 3, Ipsi= ipsilateral, Contra=contralateral, L=left, R=right.

Figure S3. **Morphology and Microstructure.** Cortex-wide and subregional hippocampal univariate differences in thickness and mean diffusivity (MD) between healthy and TLE patients. Significant *p*-values were highlighted after correcting for multiple comparisons ($p_{FWE} < 0.05$, *black outlines*). **Abbreviations:** TLE=temporal lobe epilepsy, HC= healthy controls, Ipsi= ipsilateral, Contra=contralateral

Figure S4. **Morphology, microstructure, and gradients.** Cortex-wide and subregional hippocampal univariate differences in Episodic G1 **(A)** and Semantic G1 **(B)**, after controlling for thickness and mean diffusivity (MD) between healthy and TLE patients ($p_{FWE} < 0.05$). At the neocortical level, thickness did not have a significant impact in both memory states, while MD significantly weakened episodic G1 and strengthened semantic G1 differences. At the hippocampal level, both morphology and microstructural properties did have a significant impact on episodic G1 differences. **Abbreviations:** TLE=temporal lobe epilepsy, HC= healthy controls, Ipsi= ipsilateral, Contra=contralateral

Figure S5. Differences in total episodic, semantic task performance, MoCA scores, and reaction times. **Abbreviations**: RT=reaction times, TLE=temporal lobe epilepsy, HC=healthy controls.

Functional decoding of regions showing TLE-related Gradient contractions

Figure S6. The associated cognitive terms were extracted using the *Neurosynth* decoder function and represented in a word-cloud plot. The size of the fonts used for each term reflects the correlation between the FDR-corrected maps and the *Neurosynth*-generated meta-analytic maps.