HI-FISH: WHOLE BRAIN IN SITU MAPPING OF NEURONAL ACTIVATION IN DROSOPHILA DURING SOCIAL BEHAVIORS AND OPTOGENETIC STIMULATION

Kiichi Watanabe,[1,3,4] Hui Chiu,[1,5] and David J. Anderson[1,2,*]

[1]Division of Biology and Biological Engineering

Tianqiao and Chrissy Chen Institute for Neuroscience

California Institute of Technology, Pasadena, CA USA

[2]Howard Hughes Medical Institute

[3]Present address: International Center for Cell and Gene Therapy, Fujita Health University, Toyoake, Japan

[4]Present address: Department of Medical Research for Intractable Disease, Fujita Health University, Toyoake, Japan

[5]Present address: Department of Immunobiology, Yale University School of Medicine, New Haven, CT USA

* Authors for correspondence

e-mail: wuwei@caltech.edu

Abstract

Monitoring neuronal activity at single-cell resolution in freely moving *Drosophila* engaged in social behaviors is challenging because of their small size and lack of transparency. Extant methods, such as Flyception, are highly invasive. Whole-brain calcium imaging in head-fixed, walking flies is feasible but the animals cannot perform the consummatory phases of social behaviors like aggression or mating under these conditions. This has left open the fundamental question of whether neurons identified as functionally important for such behaviors using loss- or gain-of-function screens are actually active during the natural performance of such behaviors, and if so during which phase(s). Here we describe a method, called HI-FISH, for brain-wide mapping of active cells expressing the Immediate Early Gene *hr38* using a high-sensitivity/low background amplification method called HCR-3.0. Using double-labeling for hr38 mRNA and for GFP, we describe the activity of several classes of aggression-promoting neurons during courtship and aggression, including P1^a cells, an intensively studied population of male-specific interneurons. Using HI-FISH in combination with optogenetic activation of aggression-promoting neurons (opto-HI-FISH) we identify candidate downstream functional targets of these cells in a brain-wide, unbiased manner. Finally we compare the activity of P1^a neurons during sequential performance of courtship and aggression, using intronic vs. exonic *hr38* probes to differentiate newly synthesized nuclear transcripts from cytoplasmic transcripts synthesized at an earlier time. These data provide evidence suggesting that different subsets of P1^a neurons may be active during courtship vs. aggression. HI-FISH and associated methods may help to fill an important lacuna in the armamentarium of tools for neural circuit analysis in *Drosophila*.

Introduction

Monitoring neuronal activity during specific actions or activities is critical to understanding how the brain controls behavior. Measurements within a local brain region of neural activity in non-transparent awake, behaving animals such as mice or fruit flies can be achieved with electrophysiological and calcium imaging techniques (Buzsáki, 2004; Grewe and Helmchen, 2009). In *Drosophila*, two-photon calcium imaging (TPI) has been widely used (Seelig et al., 2010) because single-unit electrical recording in the fly central brain remains exceedingly difficult (Turner et al., 2008; Wilson et al., 2004). But TPI in *Drosophila* requires head-fixation, which limits the repertoire of behaviors that flies can perform under these conditions. For example, it is difficult to use TPI to image neuronal activity during the consummatory phases of complex social behaviors, such as male-male aggression or male-female courtship (but see Clowney et al., 2015). Recently, a new method called Flyception2 was introduced for brain imaging in freely walking flies (Grover et al., 2020, 2016). However the spatial resolution of this method is limited, it is invasive, and monitoring the activity of single neurons in freely moving animals is still challenging.

In mice, immediate-early genes (IEGs) such as c-fos, Arc, or Egr1 have been used to map the activity of neurons by visualizing their expression histologically using immunohistochemistry (IHC) or in situ hybridization (ISH) (Morgan and Curran, 1991). More recently, this method has been combined with tissue-clearing and light-sheet microscopy to perform systematic and unbiased whole-brain mapping of neurons activated during different behaviors (Kim et al., 2015; Renier et al., 2016). An extension of this approach, called cellular compartment analysis of temporal activity by fluorescence in situ hybridization (catFISH), enables within-animal comparisons of neuronal populations activated during two different sequential behaviors (Guzowski et al., 1999; Lin et al., 2011).

In *Drosophila*, by contrast, the application of IEGs to map whole-brain neuronal activation patterns has been limited. In part this is because c-*fos* is not a particularly good activity marker in flies (Chen et al., 2016). More recently, several studies using fly IEGs such as Hr38 have been reported (Chen et al., 2016; Fujita et al., 2013; Takayanagi-Kiya and Kiya, 2019). Nevertheless, this technique has been relatively under-utilitized in flies. This may be the case, at least in part, because application of FISH to the adult *Drosophila* brain has been technically challenging, due to limited probe penetration and high background caused by traditional amplification methods such as Tyramide (Raap et al., 1995).

To circumvent these problems, we have applied in this study an *in situ* hybridization amplification method called the Hybridization Chain Reaction (HCR), v.3.0 (Choi et al., 2018) to visualize the expression of *Hr38*, one of the few IEGs shown to work in the adult male fly brain (Fujita et al., 2013). The design of HCR, which requires conjoint hybridization to target sequences of adjacent pairs of "half-probes" to achieve amplification, affords both greater sensitivity and specificity, allowing detection of low-abundance transcripts with minimal background noise and off-target hybridization. We call this method HCR3.0-amplified IEG Fluorescent In Situ Hybridization, or "HI-FISH" (Figure 1A).

Here we provide proof-of-concept for three different applications of HI-FISH relevant to understanding the neural circuitry underlying social behaviors in flies. First, we have asked whether specific neuronal populations discovered in functional screens for aggression-promoting neurons (Asahina et al., 2014; Hoopfer et al., 2015; Watanabe et al., 2017) are indeed active during natural aggressive behavior, courtship or both. We have further investigated whether such activity is only observed during the contact-mediated consummatory phase of these behaviors, or can also be detected in freely moving flies in the absence of fighting or mating. Second, we have performed

unbiased, whole-brain functional mapping of the downstream targets activated by optogenetic stimulation of a neuronal population of interest, a variant method we call "opto-HI-FISH.". We have applied this method to P1[a] neurons (Anderson, 2016; Clowney et al., 2015; Hoopfer et al., 2015; Inagaki et al., 2014) a subset of the male-specific P1 class of interneurons originally identified based on their ability to trigger male courtship behavior when stimulated (von Philipsborn et al., 2011). Finally, we have developed a variant of the catFISH technique (Guzowski et al., 1999), using dual-color FISH combining intronic and exonic *Hr38* probes, to compare neuronal populations activated during two different sequential behaviors, a method we call "HI-catFISH." Since experimental activation of P1[a] neurons can promote both aggression and courtship in pairs of male flies (Hoopfer et al., 2015), we have used HI-catFISH to investigate whether separate or overlapping subpopulations of P1[a] neurons are active during natural occurrences of these two social behaviors. Together, the results of these studies illustrate the utility of HI-FISH, opto-HI-FISH and HI-catFISH to detect the activation of both known and previously unidentified neurons during naturalistic social behaviors that are not amenable to head-fixation. The results have also yielded new biological insights into the neural circuit-level control of aggression and courtship.

Results

Brain-wide expression of *Hr38* detected by HI-FISH

The HCR3.0 method provides greater sensitivity, specificity and lower background than traditional FISH using Tyramide signal amplification (TSA). Application of older FISH methods in the *Drosophila* adult whole brain has been limited, due to low sensitivity and high background (Wilkie and Davis, 1998). For example, the detection by FISH of low-abundance transcripts such

as those encoding GPCRs has been virtually impossible in adult flies up to now. Application of the HCR-FISH method, by contrast, was able to detect mRNA encoding an octopamine receptor Oct-TyrR (Arakawa et al., 1990; Saudou et al., 1990), as well as Dh44, a high-abundance neuropeptide-encoding transcript (Figure 1-figure supplement A and B) (Choi et al., 2018). Background in controls lacking the HCR initiator probes was essentially undetectable.

Next, we investigated the ability of HCR to detect brain-wide expression of *Hr38* following artificial (optogenetic) activation of fruitless-GAL4 (Stockinger et al., 2005) neurons expressing the red-shifted opsin Chrimson::tdT. Fru-GAL4 labels ~2,000 neurons in the adult male brain, providing a large number of potentially activatable cells. Following photostimulation *Hr38* signals were detected in virtually all Chrimson::tdT$^+$ neurons as well as in other tdT$^-$ neurons (Figure 1B). The latter may be indirectly activated via synaptic inputs from Fru-GAL4$^+$ neurons. Control flies of the same genotype but without photostimulation, or photostimulated flies with an "empty" GAL4 driver (BDPG4uw; Pfeiffer et al., 2008) displayed much lower levels of *Hr38* expression.

While performing these experiments, we observed that simply transferring flies to a new arena induced a significant level of *Hr38* expression throughout the brain (Figure 1-figure supplement C and D), likely reflecting the stress caused by the manipulation and new environment. Based on this observation, we conducted subsequent experiments after flies were habituated to the testing arena before starting the assays (see Methods). This minor modification proved essential to distinguishing behavior-specific HI-FISH signals from the widespread *Hr38* induction caused by transfer of the flies from their home vial to an experimental apparatus.

Mapping of neurons activated by social behaviors

As an initial proof-of-concept application of HI-FISH, we asked whether neuronal subsets initially identified in functional screens for aggression-promoting neurons (Asahina et al., 2014; Hoopfer et al., 2015; Watanabe et al., 2017) were actually active during natural aggressive behavior. These included P1[a], Tachykinin-FruM[+] (Tk[FruM]) and aSP2 neurons. Despite the fact that activity in these neuron subsets was shown to be essential for normal aggression, it was formally possible that the behavioral phenotypes caused by their artificial stimulation were due to their supra-physiological activation. Therefore, we performed HI-FISH on fly lines carrying a myr:GFP reporter in the cell populations of interest in males paired with different types of target flies under different conditions. Thirty to 45 minutes after a 20 min interaction with the target fly, we fixed the brains and performed double-label HI-FISH together with fluorescent antibody staining of myr::GFP-expressing neurons (Figure 1A; see Methods).

In the control condition, where the experimental flies were kept alone during the observation period, there was little or no detectable *Hr38* expression in P1[a], Tk, or aSP2 neurons (Figure 2A-D, "Control"). In contrast, we observed clear *Hr38* signals in GFP[+] P1[a], Tk, and aSP2 neurons after a male-male interaction that included fighting behavior (Aggression: Figure 2A-D, Figure 2-figure supplement). In the case of courtship with a female, comparably strong *Hr38* induction was observed in P1[a] and aSP2 neurons, while there was no *Hr38* expression in Tk neurons (Figure 2A-D, Figure 2-figure supplement; Courtship). These data are consistent with prior observations indicating that the functional manipulation of P1[a] or aSP2 neurons in the male flies affects both male-male aggression and male-female courtship behavior (Hoopfer et al., 2015; Takayanagi-Kiya and Kiya, 2019; Watanabe et al., 2017), while the manipulation of Tk neurons affects only aggression (Asahina et al., 2014).

Social interactions in flies consist of multiple phases: approach, investigation, initial contact and progression to consummatory behaviors (Chiu et al., 2021; Yamamoto and Koganezawa, 2013). While electrophysiological recording or calcium imaging can identify neuronal activation that is time-locked to specific behavioral actions, the detection of *Hr38* (or any IEG) induction may reflect activation at any or all of these sequential phases due to the long-lived, accumulated expression of the IEG mRNA. To identify the phase(s) of social behavior wherein *Hr38* induction occurred, we separated the signals or interactions characteristic of these sequential phases in space rather than time. First, we asked whether the activation of these neurons during a male-male social interaction requires actual fighting behavior, or simply close contact with the target male. Initially, we paired the experimental male fly with a dead male, as a source of chemosensory cues. Because some male-derived cues (e.g., visual motion) may require a live conspecific, we also placed the experimental and tester males in an arena with an agarose substrate. Under these conditions, physical contact but not fighting occurs (Lim et al., 2014). In both "touching but no fighting" conditions, the fraction of Hr38[+] cells among P1[a], Tk and aSP2 neurons was slightly lower but not statistically significantly different from that measured during actual aggression, while the intensity of *Hr38* mRNA expression in positive cells was slightly lower in the No Food condition (Dead Male and No Food: Figure 2B-D, Figure 2-figure supplement). Thus the activation of P1[a], Tk, or aSP2 neurons does not require consummatory aggressive actions such as lunging, tussling or boxing (Chen et al., 2002).

Interacting male flies can, in principle, exchange visual, auditory, mechanosensory or chemosensory cues. In the case of chemosensory signals, inter-male aggression has been shown to require both detection of volatile cues by the olfactory system, and of non-volatile cues by the gustatory system (Fernández and Kravitz, 2014; Wang et al., 2011; Wang and Anderson, 2010).

While both types of cues are present when an experimental fly is exposed to an intact or a dead male fly, the detection of non-volatile cues requires contact chemosensation (Vosshall and Stocker, 2007). We therefore exposed the experimental males to a target fly separated by a mesh filter which prevented physical contact between the freely moving flies. Interestingly, *Hr38* induction in all 3 cell populations under this condition was significantly weaker that that measured during free interaction with a male on agarose, where contact but no fighting occurred (Mesh: Figure 2B-D, Figure 2-figure supplement). Taken together with the relatively strong *Hr38* induction observed following contact with a dead male (which presumably provide a source of both olfactory and gustatory pheromones), these data suggest that strong activation of P1ᵃ, Tk, and aSP2 neurons requires the integration of both volatile and non-volatile male-derived chemosensory cues. However the results do not exclude the possibility that these neurons exhibit additional activity during contact-mediated aggression, but which is not statistically detectable by this method in comparison to "no fighting" controls.

Detection of *Hr38*⁺ cells induced by optogenetic stimulation of P1ᵃ neurons in the male brain

Optogenetic and thermogenetic stimulation experiments have shown that that P1ᵃ interneurons can promote both male-directed aggression and male- or female-directed courtship (Hoopfer et al., 2015; von Philipsborn et al., 2011). While the neural circuitry downstream of P1 neurons that promotes female-directed courtship behavior has been well-characterized (von Philipsborn et al., 2011), the pathway by which P1ᵃ neurons can promote aggression is poorly understood. As a first step to identify systematically P1ᵃ functional downstream targets, we optogenetically activated P1ᵃ neurons using Chrimson:tdT and mapped the brain-wide pattern of *Hr38* induction in males (Figure 3A-D), a method we call "opto-HI-FISH (Figure 1A)." To analyze

these data, images of *Hr38* expression from multiple individuals were registered to a reference brain using antibody labeling of Bruchpilot (Brp), a reference neuropil marker, to generate an average induction pattern (Figure 3E) (Cachero et al., 2010). We then created a voxel-based heat map for the *Hr38* expression from the average image. The heat map indicated that neurons around the posterior dorsal region of the central brain were strongly activated by P1[a] stimulation (Figure 3F).

To validate this approach, we first asked whether we could detect *Hr38* induction in pCd neurons, which were previously shown by calcium imaging to be (indirect) targets of P1[a] neurons . Double-labeling in pCd-GAL4; UAS-GFP flies using HI-FISH and immunostaining for GFP (see Methods) confirmed that *Hr38* was strongly induced in pCd neurons by P1[a] photostimulation (Figure 3G, pCd). Next, we asked whether we could identify any of the specific cellular targets of P1[a] neurons in brain regions that exhibited strong induction of *Hr38*. These regions included Kenyon cells (KCs) and PAM dopaminergic neurons in the mushroom body (MB; Figure 3F). We used the GAL4 drivers OK107 (Connolly et al., 1996) to label KCs, and 0273 and Ddc-GAL4 to label PAM neurons. Following P1[a] optogenetic activation, brains were processed for double-labeling with HI-FISH and anti-GFP antibody staining to identify the cells of interest. We observed evidence of *Hr38* induction in all three cell populations, although the signal was difficult to quantify due to the high density of GFP[+] cells (Figure 3G).

To confirm the results of the opto-HI-FISH experiments, we next asked whether PAM neurons and/or Kenyon cells were activated in response to P1[a] stimulation using *in vivo* calcium imaging in living, head-fixed flies. We expressed a calcium indicator, jGCaMP7b (Dana et al., 2019) in either Kenyon cells or PAM neurons using the GAL4/UAS system and Chrimson::tdT in P1[a] neurons using the LexA/LexAop system (Brand and Perrimon, 1993; Lai and Lee, 2006). The

neurites of both KCs and PAM neurons showed significant time-locked responses to optogenetic activation of P1^a neurons (Figure3-figure supplement 3A-F). We also imaged jGCaMP7b signals from cell bodies of PAM neurons while photo-stimulating P1^a neurons, and observed clear responses in both cases as well (Figure 3-figure supplement 3G-J). Although we could not identify the specific activated cell populations among the KCs (data not shown), P1^a photo-stimulation evoked a significant rise in $\Delta F/F$ in ~50 % of PAM neurons (Figure 3-figure supplement 3I). These data validate the results from brain-wide *Hr38* induction mapping and provide proof-of-concept that brain regions containing functional targets of a cell population of interest (here P1^a neurons), as well as specific neuron subtype targets within these regions (KCs and PAM neurons), can be identified by opto-HI-FISH. These results are also consistent with a recent report showing that subsets of PAM dopaminergic neurons receive excitatory inputs from P1 neurons, and assign to KCs a positive valence associated with P1 activation (Shen et al., 2023)

HI-catFISH reveals distinct fighting- and mating-activated P1^a neuron subsets

Previous reports showed that artificial activation of P1^a neurons promoted both aggression and courtship . The foregoing data indicated that *Hr38* was induced in P1^a neurons after both natural aggressive and courtship behavior (Figure 2, Figure 2-figure supplement). An important outstanding question is whether the aggression-and courtship-activated P1^a populations are the same or different (Anderson, 2016). Because individual P1 neuron subtypes cannot be distinguished by FISH, between-animal comparisons of *Hr38* induction following fighting or mating cannot distinguish these two possibilities. To compare directly which P1^a neurons are active during naturalistic mating vs. aggression, we developed an *Hr38* analog of catFISH (HI-catFISH) to compare neurons activated during aggression and courtship in the same animal (Figure

4A). The catFISH method exploits the different kinetics of nuclear-cytoplasmic transport and of IEG mRNA stability (Guzowski et al., 1999) to distinguish neurons activated during a behavior performed just prior to analysis (nuclear pre-mRNA), from those activated during a different behavior performed 30 min before the second (cytoplasmic mRNA); cells activated during both behaviors contain both cytoplasmic and nuclear signals (Figure 4B). In mice, we have used catFISH with c-*fos* to compare hypothalamic neurons activated during aggression vs. mating in males (Lin et al., 2011).

We detected neuronal populations activated during two consecutive social behaviors using HI-FISH containing *Hr38* intronic and exonic probes amplified using different fluorophores. In two color FISH, unspliced pre-mRNAs were detected as nuclear dots labeled by both probes, while mRNA was detected as a more diffuse cytoplasmic signal detected by only the exonic probe (Figure 4B). A time course study using opto-HI-FISH with these two different *Hr38* probes indicated that the primary RNA transcripts first appeared in P1[a] nuclei ≤15 min after optogenetic activation, and disappeared by 60 min. Conversely, cytoplasmic processed mRNAs started to accumulate in the cytoplasm 30 min after activation and disappeared by 2-3 hours after stimulation (Figure 4B, top two panels and Figure 4-figure supplement 4C).

Having established this calibration of the temporal resolution of HI-catFISH, we compared labeling during free social interactions with a male or a female target. Following pre-habituation (see above), a naïve experimental male fly was placed into a behavioral arena, and a male or female fly was introduced as a first target. After 20 minutes of free social interactions the target fly was removed from the arena and, after a further 30 min recovery period, replaced by a second target fly (male or female; Methods). All four combinations of sex of the 1st and 2nd target fly were tested (Fig. 4C). Ten minutes after exposure to the second target fly (and ~1 hr after exposure to the first

target fly), the experimental fly was sacrificed and analyzed by dual-color HI-FISH *Hr38* (Figure 4A and B).

As expected, many of the neurons activated by the first target were re-activated by the second target (nuclear and cytoplasmic signal), when the sex of the two target flies was the same (Figure 4C, Male -> Male, Female -> Female, arrowheads and Figure 4D, "Both" vs. "Cyto"). By contrast, many neurons activated by the first target fly (cytoplasmic signal) were not activated by the second target fly (nuclear signal), when the target sexes were different (Figure 4C, Female -> Male, Male -> Female, arrows and Figure 4D, "Cyto" vs. "Both"). Responses to males were clearly detected in a subset of P1[a] neurons whether the male was presented first (Figure 4C, Male→Female; GFP[+]; dashed oval outlines containing blue diffuse cytoplasmic signal) or second (Figure 4C, Female→Male; magenta nuclear dots). In addition, there were male-activated cells nearby that were not labeled by the P1[a]-split GAL4 driver. In general, these male HI-FISH signals did not overlap with the HI-FISH signals evoked by female targets. (We note, however, that relatively few neurons were activated by the female fly when the sex of the two targets was different, regardless of whether the female was presented first or second; the reasons for this are currently unclear but may reflect a suppressive effect of aggression neuron activation on courtship neuron activity, as suggested by earlier studies (Hoopfer et al., 2015).) These data suggest that there may be distinct P1[a] subpopulations that are activated during either male-male aggression or male-female courtship.

Discussion

Drosophila is a powerful model system for identifying and characterizing neural circuits that control innate behaviors. With recent progress in two-photon imaging techniques using

genetically encoded calcium sensors, it has become possible to monitor activity from genetically labeled neurons at single-cell resolution in head-fixed flies (Seelig et al., 2010). Although single-neuron imaging in unrestrained behaving animals has been achieved in mammalian studies (Flusberg et al., 2008; Remedios et al., 2017; Ziv et al., 2013) most imaging studies with single-cell resolution in *Drosophila* require head-fixed conditions. Therefore, the repertoire of naturalistic behaviors that can studied using this approach – in particular social behaviors dependent on chemosensory cues (but see Hindmarsh Sten et al., 2021)– remains limited.

In mammalian systems, the discovery that IEGs such as c-*fos* are induced by neuronal activity (Greenberg et al., 1985; Greenberg and Ziff, 1984) has enabled brain-wide functional mapping of neuronal activation following specific behaviors (Morgan and Curran, 1991; Renier et al., 2016). This method and its derivatives have had a transformative impact on the identification of behaviorally relevant neural circuits in rodents. In *Drosophila*, by contrast, few if any studies describe the application of IEGs for brain-wide activity mapping. In this report, we applied the HCR3.0 RNA-FISH technique (Choi et al., 2018) to visualize the expression of a fly IEG, *Hr38* in the entire adult male brain following different manipulations (see also Shao et al., 2019). We present data providing proof-of-concept that HI-FISH can be used to detect the activation by naturalistic stimuli of neurons previously identified in thermogenetic behavioral activation screens, as well as to identify candidate downstream targets of these neurons following their optogenetic activation (opto-HI-FISH). We also compared neuronal subsets within a small identified population (P1[a] neurons) that is activated during two different social behaviors, using HI-catFISH.

Monitoring Neuronal Activities during Social Behaviors

Previous studies reported that male-specific P1[a], Tk, and aSP2 neurons promote male-male aggression (Asahina et al., 2014; Hoopfer et al., 2015; Watanabe et al., 2017). In addition, P1[a] and aSP2 neurons promote male-female courtship (Hoopfer et al., 2015; Takayanagi-Kiya and Kiya, 2019). Artificial manipulation of the activity of these neurons affected social behaviors and demonstrated their causal roles in these behaviors. However, there was no direct evidence showing that these neurons are normally activated during the specific behaviors they can promote.

Our HI-FISH data demonstrated that P1[a] and aSP2 neurons are active during male-male aggression and male-female courtship, consistent with the behavioral effects of activating these neurons (Figure 2) (Hoopfer et al., 2015; Watanabe et al., 2017). In contrast, Tk[FruM] neurons were only active during male-male aggression (Figure 2), consistent with their lack of a functional role in courtship (Asahina et al., 2014). Furthermore, we show that P1[a], Tk, and aSP2 neurons can apparently be activated by chemosensory cues from an opponent male fly in the absence of attack behavior (Figure 2). These data suggest that P1[a], Tk, and aSP2 neurons may serve to process and integrate sensory cues from the opponent and relay that information to the downstream neurons that encode the consummatory phase of aggression (Chiu et al., 2021). However they do not exclude that these neurons may play a role during overt attack.

Although the time resolution of IEG-based activity monitoring falls far short of that of calcium imaging, our data show that one can nevertheless use HI-FISH to separate activation that occurs during the appetitive vs. consummatory phases of male-male aggression, by judicious manipulation of experimental conditions.

Identification of downstream target neurons of genetically labeled neurons by opto-HI-FISH

Artificial activation of P1[a] neurons induces male-female courtship and an internal state that promotes male-male aggression (Bath et al., 2014; Hoopfer et al., 2015; Inagaki et al., 2014). Furthermore, artificial activation of male specific P1 neurons mimics courtship-induced place preference as well as supporting appetitive olfactory conditioning (Shen et al., 2023). It is critical to identify neurons that are functionally downstream of P1[a] cells to understand how these neurons can promote this state. In this study, using opto-HI-FISH we identified pCd neurons, Kenyon cells and PAM neurons as downstream targets of P1[a] neurons. These results were confirmed by two-photon calcium imaging analysis (Figure 3, Figure 3-figure supplement). Importantly, recent studies have reached the same conclusions regarding neuronal populations using different approaches (Jung et al., 2020; Shen et al., 2023). These findings provide additional support for our methodology. The heat map illustrates the presence of additional potential P1[a] -follower populations beyond the three interrogated here (Figure 3F). Further analysis will unveil the detailed characteristics of these potential downstream neurons of P1[a] neurons. An important caveat is that not all active *Drosophila* neurons may express *Hr38*. Act-seq data from mice indicates that different neuron subsets activated during social behaviors express different IEGs to different extents (Kim et al., 2019).

Functional heterogeneity of P1[a] populations during Social Behaviors

In addition to identifying functional downstream targets of P1[a] neurons, characterizing cellular diversity among P1[a] neurons and how this functional heterogeneity maps onto their connectivity with different downstream targets is critical to understanding how these cells can induce both courtship and aggressive behavior. Optogenetic activation of P1[a] neurons at a low frequency promotes aggression towards males, while high-frequency stimulation promotes male-

directed courtship during stimulation, followed by aggression in the inter-stimulation intervals (Hoopfer et al., 2015). P1[a] neurons may control these two behaviors according to their activity level and/or the subtype of neuron activated in the population. Alternatively, P1[a] neurons may indirectly promote fighting by inhibiting aggression-promoting neurons during courtship (Koganezawa et al., 2016); post-inhibitory rebound in such neurons following the offset of P1[a] optogenetic stimulation may trigger aggressive behavior (Anderson, 2016).

Our preliminary *Hr38* catFISH results suggest the presence of male- or female-responsive subsets within the P1[a] population, which consists of 8-10 neurons/hemibrain (Figure 4). To our knowledge, this is the first evidence that P1[a] neurons are activated during male-male social interactions. Previous studies using calcium imaging of head-fixed flies suggested that P1[a] neurons were inhibited by the male-specific pheromone, 11-cis-vaccenyl acetate (cVA; Clowney et al., 2015). The reason for this apparent discrepancy is not yet clear. The molecular identity of the male- vs. female-responsive P1[a] subsets remains to be determined. It has been suggested that FruM[+] P1 neurons exclusively control courtship, while Dsx[+] P1 neurons (also called pC1 neurons; Koganezawa et al., 2016)) control aggression. However triple-intersection experiments have demonstrated that thermogenetic (Hoopfer et al., 2015) or optogenetic (Watanabe et al., 2017) activation of FruM[+] P1[a] neurons, which comprise only 1-4 neurons/hemibrain, can also promote aggression. Therefore, while our results support the idea that different P1 subsets control aggression vs. courtship, they do not support the view that FruM[+] P1 neurons only control the latter and not the former social behavior.

In mice, distinct subpopulations of estrogen receptor 1 (Esr1)-expressing neurons in the ventrolateral subdivision of the ventromedial hypothalamus (VMHvl) are activated during male-directed aggression versus female-directed mating (Karigo et al., 2021; Remedios et al., 2017).

Our observation that apparently distinct subpopulations of P1ᵃ neurons are activated during aggression versus courtship in *Drosophila* strengthens the analogy between these two control nodes for social behavior (Anderson, 2016). Application to P1ᵃ neurons of Act-Seq, a method that uses scRNAseq to identify transcriptomic cell types that are active during different behaviors as determined by IEG expression (Kim et al., 2019; Wu et al., 2017), may help to identify transcriptomic and functional diversity within the P1ᵃ population. Such correlative studies may be extended to functional perturbations by using transcriptomic data to identify more specific intersectional GAL4 drivers for distinct P1ᵃ subsets. Together, our studies illustrate how HI-FISH can provide a systematic, unbiased and brain-wide approach to identifying entry points to circuits controlling specific social behaviors, which can be integrated eventually with imaging, connectomic, transcriptomic and functional approaches to circuit analysis.

Note

While this paper was in the final stages of preparation, a manuscript by Takayanagi-Kiya et al. appeared that described a method for labeling neurons activated during innate behaviors using a different immediate early gene, *stripe/egr-1* (Takayanagi-Kiya et al., 2023). That study used a GAL4 driver for the *stripe/egr-1* gene to label and functionally manipulate activated neurons. In contrast, our approach is based purely on detecting expression of the IEG mRNA using a state-of-the-art amplification method, HCR v3.0 (Figure 1-figure supplement B) (Choi et al., 2018). The HI-FISH method enables us to test essentially any IEG of interest, rather than only those IEGs for which GAL4 drivers are available. Furthermore, by using direct mRNA detection rather than an IEG-GAL4 driver, a series of GAL4 or split-GAL4 lines can be combined with our

approach to characterize the identity of IEG-labeled neurons. Thus, the two methods complement each other.

Materials and methods

Fly stocks

A list of the full genotypes for the flies used in this research is in Supplementary file 1.

R15A01-iLexA (improved LexA) was generated for this study. Briefly, the R15A01 enhancer fragment (Jenett et al., 2012) was cloned into pBPnlsLexA::p65::GADUw (Chiu et al., 2021).

20xUAS-IVS-Syn21-Chrimson::tdTomato3.1 (attP2) and 13xLexAop2-IVS-Syn21-Chrimson::tdTomato3.1 (su(Hw)attP5) were obtained from G. Rubin (Janelia Research Campus).

0273-GAL4 (Burke et al., 2012) was obtained from S. Waddell (The University of Oxford).

fru[GAL4] (Stockinger et al., 2005) was from B. Dickson (Janelia Research Campus).

The following strains were obtained from Bloomington Drosophila Stock Center (Indiana University); R15A01-AD (attP40), R71G01-DBD (attP2), R41A01-GAL4 (attP2), OK107, Ddc-GAL4, 20xUAS-IVS-jGCaMP7b (VK00005).

Optogenetic activation in behaving flies

A detailed description of the setup used for optogenetic activation was described previously (Inagaki et al., 2014; Watanabe et al., 2017). Flies were raised at 25℃ and collected on the day of eclosion and kept in the dark for 7-10 days on the fly media containing 0.4 mM all *trans*-Retinal (Merck, St. Louis, MO). The experiments were performed with the "8-well" chamber (16 mm diameter x 10 mm height) with IR backlight . (855 nm, SOBL-6x4-850, SmartVision Lights,

Norton Shores, MI). A 655 nm 10 mm Square LED (Luxeon Star LEDs Quadica Developments, Brantford, Canada) was used for photostimulation. The stimulation protocols for each experiment are described in the figures or figure legends.

Behavioral Assays

Flies were maintained at 9AM:9PM Light:Dark cycle. For male-male aggression assays and male-female courtship assays, the "8-well" acrylic chamber was used as described previously (Inagaki et al., 2014; Watanabe et al., 2017). The walls of the chamber were coated with a fluoropolymer resin (PTFE-30, Insect-a-Slip) (BioQuip Products, Rancho Dominguez, CA). The clear top ceiling plate was coated with Sigmacote (Merck, St. Louis, MO) to prevent flies from walking on the surfaces. The floor was covered with a layer of apple juice/sucrose-agarose food. The arena was illuminated with IR backlighting.

Two-photon calcium imaging

Calcium imaging with optogenetic activation was performed as described (Inagaki et al., 2014; Jung et al., 2020). A deep red (660 nm) fiber-coupled LED (Thorlabs, Newton, NJ) with a band-pass filter (660 nm, Edmund Optics, Barrington, NJ) was used for light source to activate Chrimson.

Flies were collected on the day of eclosion and kept in the dark on the fly media containing 0.4 mM all *trans*-Retinal (Merck, St. Louis, MO). 7-10 days old flies were cold-anesthetized using a temperature-controlled stage and mounted on a plastic plate with a small window using wax. The cuticle of the dorsal side of the head was removed by forceps. Flies were then placed beneath the objective.

Fly saline (108 mM NaCl, 5 mM KCl, 4 mM NaHCO$_3$, 1 mM NaH$_2$PO$_4$, 5 mM trehalose, 10 mM sucrose, 5 mM HEPES, 0.5 mM CaCl$_2$, 2 mM MgCl$_2$, pH = 7.5) was used to bathe the brain during imaging.

Immunohistochemistry and HCR fluorescence in situ hybridization (FISH)

In situ HCR was performed as previously described with minor modifications (Choi et al., 2018, also see https://www.moleculartechnologies.org/supp/HCRv3_protocol_generic_solution.pdf). For the hybridization buffer, formamide was substituted with an equal volume of 8M urea solution for better preservation of tissue morphology (Sinigaglia et al., 2018).

To process multiple individuals, we collected flies (~ 8 individuals at once), soaked them in ice-cold ethanol to remove the wax on the cuticle, and opened the head cuticle of each fly with minimal time-lag among samples in the ice-cold fixative (4% paraformaldehyde in PBS). After brief fixation, the brain from each individual was dissected and fixed for 2 hours at 4°C.

For combination of immunohistochemistry (IHC) and HCR FISH, we performed IHC as described previously before HCR FISH procedure in RNase free condition (Watanabe et al., 2017). For antibody incubation, we added RNasin Plus (5% of final volume, Promega, Madison, WI). After immunostaining, the samples were post-fixed with 4% formaldehyde in PBS and proceeded for HCR. HCR probes were purchased from Molecular Technologies (https://www.moleculartechnologies.org/). The samples were mounted in 88% (w/v) Histodenz in 0.02 M Phosphate buffer (pH 7.5 with NaOH) with 0.1% Tween-20 and 0.01% sodium azide (Yang et al., 2014).

The antibodies used were as follows: rat anti-DN Cadherin (1:50, DN-EX #8, developed by T. Uemura, obtained from Developmental Studies Hybridoma Bank, University of Iowa, Iowa City, IA), mouse anti-Bruchpilot (1:50, nc82, developed by E. Buchner, obtained from Developmental Studies Hybridoma Bank, University of Iowa, Iowa City, IA), rabbit anti-GFP (1:1,000, A11122, Thermo Fisher Scientific, Waltham, MA), Goat anti-Mouse IgG (H+L) Alexa Fluor 488 (1:1,000, A-11001, Thermo Fisher Scientific, Waltham, MA), Goat anti-Mouse IgG (H+L) Alexa Fluor 568 (1:1,000, A-11004, Thermo Fisher Scientific, Waltham, MA), Goat anti-Mouse IgG (H+L) Alexa Fluor 633 (1:1,000, A-21050, Thermo Fisher Scientific, Waltham, MA), Goat anti-Rabbit IgG (H+L) Alexa Fluor 488 (1:1,000, A-11008, Thermo Fisher Scientific, Waltham, MA), Goat anti-Rabbit IgG (H+L) Alexa Fluor 488 (1:1,000, A-11011, Thermo Fisher Scientific, Waltham, MA), Goat anti-Rabbit IgG (H+L) Alexa Fluor 633 (1:1,000, A-21070, Thermo Fisher Scientific, Waltham, MA)

Confocal serial optical sections were obtained with a Fluoview FV3000 Confocal Microscope (Olympus, Tokyo, Japan).

For the Hr38 exonic probe, we first measured average fluorescence intensity within the cell boundary using Fiji (Schindelin et al., 2012; Schneider et al., 2012) and calculated the relative intensity to the background signals and then defined the positive cells whose signals were above 3σ.

For image registration, brain images were registered to T1 template brain (Yu et al., 2010) using the CMTK registration plugin of Fiji (https://flybrain.mrc-lmb.cam.ac.uk/dokuwiki/doku.php?id=warping_manual:registration_gui). To create average images for each condition, MATLAB was used to generate averaged images of each z-plane from different individuals. For voxel-based heat map analysis of the *Hr38* expression, the median filter

(pixel size = 10) in Fiji was applied to the averaged images. Fiji and Fluorender software (Yong Wan et al., 2009) were used to create z stack images.

catFISH

To prepare target flies, small iron filings were attached on the dorsal side of the thorax of 6-8 days old male or female flies using UV-activated glue. This allowed us to remove the target fly from a behavioral arena through a small hole using a neodymium magnet.

6-8 days single-housed male flies were used for the experimental flies. A male fly was introduced in the behavioral arena and allowed to habituated to the new environment for ~ 3 hours. The first target fly was introduced into the arena, and the experimental fly was exposed to the target for 20 mins. After the exposure, the target fly was removed from the arena as describe above. The experimental fly was given a 30-min interval before the second target was introduced. After 20 min exposure, the second target was removed, and *Hr38* expression was analyzed 10 min after the target fly removal (Figure 4A and B). Neurons with a clear dot signal were counted as positive for the Hr38 intronic probe.

Statistical Analysis

Statistical analyses were performed using MATLAB (MathWorks, Natick, MA) and Python (Python Software Foundation). The detailed information including the number of samples, the statistical method, and the p value for each experiment is indicated in the figure legends.

Calcium imaging data analysis

ROIs (region of interest) corresponding to nerve bundle or cell bodies of labeled neurons were manually selected and the fluorescence intensity in the ROIs was measured using Fiji. The average fluorescence intensity during the first 15 s of image acquisition was used as the baseline to calculate the $\Delta F/F$. The mean of $\Delta F/F$ during the indicated periods was calculated and visualized using Python. For cell body analysis, we defined the positive response if the average $\Delta F/F$ during the photostimulation was statistically higher than the baseline ($p < 0.01$). The strong response neurons ($\Delta F/F > 0.4$) and the intermediate neurons ($0.15 < \Delta F/F \leq 0.4$) were defined based on the average $\Delta F/F$ value during the stimulation, respectively.

Author Contributions

Kiichi Watanabe, Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Resource, Software, Visualization, Writing - original draft, Writing - review and editing; Hui Chiu, Resource; David J. Anderson, Conceptualization, Funding acquisition, Methodology, Project administration, Supervision, Writing - original draft, Writing - review and editing.

Acknowledgements

We thank N.A. Pierce and M. Schwarzkopf for advice on HCR techniques; A. Sanchez for fly stock maintenance; S. Cao for advice on imaging experiments; B. Weissbourd for advice on catFISH experiments; G.M. Rubin, B.D. Pfeiffer, and S. Waddell for reagents; C. Chiu, G. Mancuso, and L. Chavarria for the laboratory management and administrative assistance; and members of the Anderson laboratory for valuable comments on this work. This work was

supported by NIDA Grant R01 DA031389. D.J.A. is an Investigator of the Howard Hughes Medical Institute.

FIGURE LEGENDS

Figure 1. Mapping neuronal activities using Hr38 FISH.

(A) Illustrated summary of this study for mapping Hr38 expression induced by social behavior or artificial activation of specific neuronal populations. (B) Representative images of Hr38 expression before (fru-No PS) and after photo-stimulation of fru-GAL4>Chrimson neurons (fru-PS). BDPG4uw-PS represents the images after Photo-stimulation of empty-GAL4>Chrimson neurons. magenta: Chrimson::tdT, green: Hr38 HCR signals. (C) High magnification images of Hr38 expression after photo-stimulation of fru-GAL4>Chrimson neurons. SMP, superior medial protocerebrum; LH, lateral horn; KC, Kenyon cells.

Figure 2. Investigation of neurons activated by social behavior with Hr38 FISH.

(A) Hr38 expression in the neurons labeled with P1[a]-split GAL4, Tk-GAL4, and aSP2-split GAL4 drivers. magenta: myr::GFP, yellow: Hr38 HCR signals. Dotted lines depicted the outlines of myr::GFP labeled cell bodies. Arrows: cell bodies of Hr38 positive neurons. (B) - (D) Expression level of Hr38 HCR signals (top) and percentage of Hr38 positive neurons (bottom) in P1[a] (B), Tk (C), and aSP2 (D) after different behavioral episodes. Aggression: aggressive behavior against another male, Dead: interaction with a dead male, No Food: interaction with another male without food (no fighting), Mesh: interaction with another male, separated with mesh (no physical contact), Courtship: courtship behavior with a virgin female. Bottom: They are grouped in categories according to the characters (a, b, c). In each group, the behavioral conditions do not have a

statistically significant difference (p < 0.05, chi-square test). Number of neurons analyzed; P1[a]-Control: 135, P1[a]-Aggression: 84, P1[a]-Dead Male: 126, P1[a]-No Food: 115, P1[a]-Mesh: 87, P1[a]-Courtship: 114, Tk-Control: 45, Tk-Aggression: 69, Tk-Dead Male: 61, Tk-No Food: 63, Tk-Mesh: 82, Tk-Courtship: 66, aSP2-Control: 20, aSP2-Aggression: 36, aSP2-Dead Male: 23, aSP2-No Food: 49, aSP2-Mesh: 36, aSP2-Courtship: 24.

Figure 3. Investigation of downstream neurons activated by optogenetic activation of P1[a] neurons.

(A) Illustration of the experimental set up. (B) Expression pattern of Chrimson::tdT driven by P1[a] split GAL4. (C) Scheme of the data analysis procedure. (D) Time-course study of Hr38 HCR expression after P1[a] photo-stimulation in P1[a] labeled neurons and the surrounding area. The expression level reaches its peak at 30-60 mins. (E) Average image of Hr38 expression of control (BDPG4Uw) and P1[a]-activated brains. 5 individual brains from each conditions were registered into a template brain and averaged. Green: Hr38 HCR signals, Magenta: Brp. (F) Heat map analysis of Hr38 expression. Areas depicted by dotted lines represent the areas of PAM neusons (PAM), mushroom bodies (MB), and pC neurons (pC). (G) Confocal images of Hr38 expressions in pCd neurons (R41A01), Kenyon cells (OK107), and PAM neurons (0273 or Ddc) of control and P1[a] activated brains. Green: myr::GFP, Magenta: Hr38. Dotted lines depicted the outlines of cell bodies (pCd) or cluster of neurons (KC, PAM).

Figure 4. Drosophila catFISH reveals P1[a] subpopulations activated by two different social behaviors.

(A) Experimental design of the fly catFISH. Each fly experienced the 1st behavior (aggression or courtship) followed by 2nd behavior (aggression or courtship), and the HCR was performed with the Hr38 probes targeted for the exon or the intron sequence. (B) Time-course of Hr38 signals detected by the exon- or the intron-targeted Hr38 HCR probes and expected Hr38 signals in each neurons. (Top) The intron-targeted Hr38 HCR signals (magenta) induced by 1st behavior disappear at the timing of HCR analysis while the exon-targeted Hr38 signals (blue) remain in the cytoplasmic area. (Middle) Both intron-targeted and exon-targeted signals are present in the nucleus. (Bottom) Both intron-targeted and exon-targeted signals induced by the 2nd behavior are present in the nucleus, and exon-targeted Hr38 signals induced by the 1st behavior are present in the cytoplasmic area. In 'Neurons active during both', the exon probe signals from the 1st behavior decline (dotted blue line *1) after 60 mins (also see Fig. 4B-supplement). However, it is canceled out as the signals from 2nd behavior are increased (dotted blue line *2). As a result, the combined Hr38 exon probe signals (solid blue line) remain high at the time point for Hr38 HCR. (C) Representative images of Hr38 HCR signals in P1^a neurons after four different behavioral conditions. Green: myr::GFP, Magenta: intron-targeted Hr38 HCR signals: exon-targeted Hr38 signals. (D) Fraction of responsive P1^a neurons across indicated conditions. N.R.: not responded, Nuc: nuclear signal only (responded to only second target), Cyto: cytoplasmic signal only (responded to only first target), Both: nuclear and cytoplasmic signal (responded to both first and second target).

Figure 1 - figure supplement. HCR FISH and Hr38 signals in the adult fly brain.

(A) and (B) Examples of whole mount FISH on adult fly brains. (A) Dh44 from neuropeptides, high copy number genes. (B) Octopamine-Tyramine receptor from G protein-coupled receptors,

low copy number gene. (C) Representative images of Hr38 expression with or without habituation after transfer procedure. (D) Comparison of the relative expression of Hr38 signals between samples with or without habituation.

Figure 2 - figure supplement. Complete data set of Hr38 expression after social behaviors.

(A) Hr38 expression in the neurons labeled with P1[a]-split GAL4, Tk-GAL4, and aSP2-split GAL4 drivers. magenta: myr::GFP, yellow: Hr38 HCR signals. Dotted lines depicted the outlines of myr::GFP labeled cell bodies. Arrows: cell bodies of Hr38 positive neurons. (B) - (D) Expression level of Hr38 HCR signals in Hr38 positive neurons in P1[a] (B), Tk (C), and aSP2 (D) after different behavioral episodes being statistically different from the control condition in % Hr38[+] neurons (see Figure 2B-D, bottom). Mann-Whitney U test with Bonferroni correction for multiple comparisons. $*p < 0.05$, $**p < 0.01$. Aggression: aggressive behavior against another male, Dead: interaction with a dead male, No Food: interaction with another male without food (no fighting), Mesh: interaction with another male, separated with mesh (no physical contact), Courtship: courtship behavior with a virgin female. Number of neurons analyzed; P1[a]-Aggression: 44, P1[a]-Dead Male: 62, P1[a]-No Food: 44, P1[a]-Mesh: 12, P1[a]-Courtship: 74, Tk-Aggression: 53, Tk-Dead Male: 41, Tk-No Food: 35, aSP2-Aggression: 28, aSP2-Dead Male: 15, aSP2-No Food: 33, aSP2-Courtship: 23.

Figure 3 - figure supplement. Calcium imaging of PAM neurons and Mushroom body neurons combined with P1[a] photostimulation.

(A)-(C) Calcium imaging of PAM neurons, labeled with 0273 GAL4 driver while P1a neurons were optogenetically activated. (A) Representative image of GCaMP signals in neuronal bundles

of PAM neurons before, during, and after P1^a activation. dF/F (B) and average dF/F (C) of PAM neurons in response to photostimulation of P1^a neurons (n = 3 individuals; mean ± SEM (B) and a box and whisker plot (C)). (D)-(F) Calcium imaging of nerve bundle of mushroom body kenyon cells, labeled with OK107 GAL4 driver while P1^a neurons were optogenetically activated. (D) Representative image of GCaMP signals in mushroom body peduncle before, during, and after P1^a activation. dF/F (E) and average dF/F (F) of Kenyon cells in response to photostimulation of P1^a neurons. (n = 3 individuals; mean ± SEM (E) and a box and whisker plot (F)) (G)-(J) Calcium imaging of cell bodies of PAM neurons while P1^a neurons were optogenetically activated. (G) Representative image of GCaMP signals in PAM neurons averaged from three consecutive trials of P1^a stimulation. Arrowheads: cell bodies shows strong responses (dF/F > 0.4) to P1^a stimulation. (H) Distribution of responsive PAM neurons. Colored by the strength of responses (Dark blue: strong response, light blue: intermediate response, grey: weak response, white: no response). (I) Temporal profiles of responses evoked by P1 photostimulation. Average dF/F during P1^a stimulation are higher than that before the stimulation (Mann-Whitney U test, p < 0.05, red line segregates between responded and non-responded groups). (J) Fraction of PAM neurons responsive to P1^a stimulation (n = 175 cells from 3 individuals).

Figure 4 - figure supplement. Time-course study of the expression of Hr38 with the exon or intron targeted probes after photostimulation of P1^a neurons.

(A) Confocal images of Hr38 expression detected by the exon-targeted (Hr38^{Ex}, blue) and the intron-targeted (Hr38^{INT}, magenta) probe in the area including P1^a cell body (myr::GFP, green). Arrows: Cells positive for both Hr38^{INT} and cytoplasmic Hr38^{Ex} signals. (B) Relative expression of Hr38 RNA detected by the exon-targeted probe over time. Red: positive (the expres(Morgan

and Curran, 1991)sion level > +3σ), blue: borderline (+3σ > the expression level > +2σ), grey: negative (< +2σ). (C) Percentage of Hr38 positive neurons detected by the intron-targeted probe (green) and the exon-targeted probe (blue) over time.

References

Anderson DJ. 2016. Circuit modules linking internal states and social behaviour in flies and mice. *Nat Rev Neurosci* **17**:692–704. doi:10.1038/nrn.2016.125

Arakawa S, Gocayne JD, McCombie WR, Urquhart DA, Hall LM, Fraser CM, Venter JC. 1990. Cloning, localization, and permanent expression of a Drosophila octopamine receptor. *Neuron* **4**:343–354. doi:10.1016/0896-6273(90)90047-J

Asahina K, Watanabe K, Duistermars BJ, Hoopfer E, González CR, Eyjólfsdóttir EA, Perona P, Anderson DJ. 2014. Tachykinin-Expressing Neurons Control Male-Specific Aggressive Arousal in Drosophila. *Cell* **156**:221–235. doi:10.1016/j.cell.2013.11.045

Bath DE, Stowers JR, Hörmann D, Poehlmann A, Dickson BJ, Straw AD. 2014. FlyMAD: Rapid thermogenetic control of neuronal activity in freely walking Drosophila. *Nat Methods* **11**:756–762. doi:10.1038/nmeth.2973

Brand a H, Perrimon N. 1993. Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Development (Cambridge, England)* **118**:401–15.

Burke CJ, Huetteroth W, Owald D, Perisse E, Krashes MJ, Das G, Gohl D, Silies M, Certel S, Waddell S. 2012. Layered reward signalling through octopamine and dopamine in Drosophila. *Nature* **492**:433–7. doi:10.1038/nature11614

Buzsáki G. 2004. Large-scale recording of neuronal ensembles. *Nat Neurosci* **7**:446–451. doi:10.1038/nn1233

Cachero S, Ostrovsky AD, Yu JY, Dickson BJ, Jefferis GSXE. 2010. Sexual Dimorphism in the Fly Brain. *Current biology : CB* **20**:1589–1601. doi:10.1016/j.cub.2010.07.045

Chen S, Lee AY, Bowens NM, Huber R, Kravitz E a. 2002. Fighting fruit flies: a model system for the study of aggression. *Proc Natl Acad Sci USA* **99**:5664–8. doi:10.1073/pnas.082102599

Chen X, Rahman R, Guo F, Rosbash M. 2016. Genome-wide identification of neuronal activity-regulated genes in drosophila. *eLife* **5**:1–21. doi:10.7554/eLife.19942

Chiu H, Hoopfer ED, Coughlan ML, Pavlou HJ, Goodwin SF, Anderson DJ. 2021. A circuit logic for sexually shared and dimorphic aggressive behaviors in Drosophila. *Cell* **184**:507-520.e16. doi:10.1016/j.cell.2020.11.048

Choi HMT, Schwarzkopf M, Fornace ME, Acharya A, Artavanis G, Stegmaier J, Cunha A, Pierce NA. 2018. Third-generation in situ hybridization chain reaction: multiplexed, quantitative, sensitive, versatile, robust. *Development* **145**:dev165753–dev165753. doi:10.1242/dev.165753

Clowney EJ, Iguchi S, Bussell JJ, Scheer E, Ruta V. 2015. Multimodal Chemosensory Circuits Controlling Male Courtship in Drosophila. *Neuron* **87**:1036–1049. doi:10.1016/j.neuron.2015.07.025

Connolly JB, Roberts IJH, Armstrong JD, Kaiser K, Forte M, Tully T, O'Kane CJ. 1996. Associative Learning Disrupted by Impaired G $_s$ Signaling in *Drosophila* Mushroom Bodies. *Science* **274**:2104–2107. doi:10.1126/science.274.5295.2104

Dana H, Sun Y, Mohar B, Hulse BK, Kerlin AM, Hasseman JP, Tsegaye G, Tsang A, Wong A, Patel R, Macklin JJ, Chen Y, Konnerth A, Jayaraman V, Looger LL, Schreiter ER, Svoboda K, Kim DS. 2019. High-performance calcium sensors for imaging activity in neuronal populations and microcompartments. *Nat Methods* **16**:649–657. doi:10.1038/s41592-019-0435-6

Fernández MP, Kravitz EA. 2014. Aggression and Courtship in Drosophila: Pheromonal Communication and Sex Recognition. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* **119**:1065–76. doi:10.1007/s00359-013-0851-5

Flusberg BA, Nimmerjahn A, Cocker ED, Mukamel EA, Barretto RPJ, Ko TH, Burns LD, Jung JC, Schnitzer MJ. 2008. High-speed, miniaturized fluorescence microscopy in freely moving mice. *Nat Methods* **5**:935–938. doi:10.1038/nmeth.1256

Fujita N, Nagata Y, Nishiuchi T, Sato M, Iwami M, Kiya T. 2013. Visualization of neural activity in insect brains using a conserved immediate early gene, Hr38. *Current Biology* **23**:2063–2070. doi:10.1016/j.cub.2013.08.051

Greenberg ME, Greene LA, Ziff EB. 1985. Nerve growth factor and epidermal growth factor induce rapid transient changes in proto-oncogene transcription in PC12 cells. *J Biol Chem* **260**:14101–14110.

Greenberg ME, Ziff EB. 1984. Stimulation of 3T3 cells induces transcription of the c-fos proto-oncogene. *Nature* **311**:433–438. doi:10.1038/311433a0

Grewe BF, Helmchen F. 2009. Optical probing of neuronal ensemble activity. *Current Opinion in Neurobiology* 520–529. doi:10.1016/j.conb.2009.09.003

Grover D, Katsuki T, Greenspan RJ. 2016. Flyception: Imaging brain activity in freely walking fruit flies. *Nat Methods* **13**:569–572. doi:10.1038/nmeth.3866

Grover D, Katsuki T, Li J, Dawkins TJ, Greenspan RJ. 2020. Imaging brain activity during complex social behaviors in Drosophila with Flyception2. *Nature Communications* **11**. doi:10.1038/s41467-020-14487-7

Guzowski JF, McNaughton BL, Barnes CA, Worley PF. 1999. Environment-specific expression of the immediate-early gene Arc in hippocampal neuronal ensembles. *Nat Neurosci* **2**:1120–1124. doi:10.1038/16046

Hindmarsh Sten T, Li R, Otopalik A, Ruta V. 2021. Sexual arousal gates visual processing during Drosophila courtship. *Nature* **595**:549–553. doi:10.1038/s41586-021-03714-w

Hoopfer ED, Jung Y, Inagaki HK, Rubin GM, Anderson DJ. 2015. P1 interneurons promote a persistent internal state that enhances inter-male aggression in Drosophila. *eLife* **4**:1–27. doi:10.7554/eLife.11346

Inagaki HK, Jung Y, Hoopfer ED, Wong AM, Mishra N, Lin JY, Tsien RY, Anderson DJ. 2014. Optogenetic control of Drosophila using a red-shifted channelrhodopsin reveals experience-dependent influences on courtship. *Nat Methods* **11**:325–332. doi:10.1038/nmeth.2765

Jenett A, Rubin GM, Ngo T-TB, Shepherd D, Murphy C, Dionne H, Pfeiffer BD, Cavallaro A, Hall D, Jeter J, Iyer N, Fetter D, Hausenfluck JH, Peng H, Trautman ET, Svirskas RR, Myers EW, Iwinski ZR, Aso Y, DePasquale GM, Enos A, Hulamm P, Lam SCB, Li H-H, Laverty TR, Long F, Qu L, Murphy SD, Rokicki K, Safford T, Shaw K, Simpson JH,

Sowell A, Tae S, Yu Y, Zugates CT. 2012. A GAL4-driver line resource for Drosophila neurobiology. *Cell Rep* **2**:991–1001. doi:10.1016/j.celrep.2012.09.011

Jung Y, Kennedy A, Chiu H, Mohammad F, Claridge-Chang A, Anderson DJ. 2020. Neurons that Function within an Integrator to Promote a Persistent Behavioral State in Drosophila. *Neuron* **105**:322-333.e5. doi:10.1016/j.neuron.2019.10.028

Karigo T, Kennedy A, Yang B, Liu M, Tai D, Wahle IA, Anderson DJ. 2021. Distinct hypothalamic control of same- and opposite-sex mounting behaviour in mice. *Nature* **589**:258–263. doi:10.1038/s41586-020-2995-0

Kim D-W, Yao Z, Graybuck LT, Kim TK, Nguyen TN, Smith KA, Fong O, Yi L, Koulena N, Pierson N, Shah S, Lo L, Pool A-H, Oka Y, Pachter L, Cai L, Tasic B, Zeng H, Anderson DJ. 2019. Multimodal Analysis of Cell Types in a Hypothalamic Node Controlling Social Behavior. *Cell* **179**:713-728.e17. doi:10.1016/j.cell.2019.09.020

Kim Y, Venkataraju KU, Pradhan K, Mende C, Taranda J, Turaga SC, Arganda-Carreras I, Ng L, Hawrylycz MJ, Rockland KS, Seung HS, Osten P. 2015. Mapping Social Behavior-Induced Brain Activation at Cellular Resolution in the Mouse. *Cell Reports* **10**:292–305. doi:10.1016/j.celrep.2014.12.014

Koganezawa M, Kimura K, Yamamoto D. 2016. The Neural Circuitry that Functions as a Switch for Courtship versus Aggression in Drosophila Males. *Current Biology* **26**:1395–1403. doi:10.1016/j.cub.2016.04.017

Lai S-L, Lee T. 2006. Genetic mosaic with dual binary transcriptional systems in Drosophila. *Nat Neurosci* **9**:703–9. doi:10.1038/nn1681

Lim RS, Eyjólfsdóttir E, Shin E, Perona P, Anderson DJ. 2014. How Food Controls Aggression in Drosophila. *PLoS ONE* **9**:e105626. doi:10.1371/journal.pone.0105626

Lin D, Boyle MP, Dollar P, Lee H, Lein ES, Perona P, Anderson DJ. 2011. Functional identification of an aggression locus in the mouse hypothalamus. *Nature* **470**:221–226. doi:10.1038/nature09736

Morgan JI, Curran T. 1991. Stimulus-transcription coupling in the nervous system: involvement of the inducible proto-oncogenes fos and jun. *Annu Rev Neurosci* **14**:421–451. doi:10.1146/annurev.ne.14.030191.002225

Pfeiffer BD, Jenett A, Hammonds AS, Ngo T-TB, Misra S, Murphy C, Scully A, Carlson JW, Wan KH, Laverty TR, Mungall C, Svirskas R, Kadonaga JT, Doe CQ, Eisen MB, Celniker SE, Rubin GM. 2008. Tools for neuroanatomy and neurogenetics in Drosophila. *Proc Natl Acad Sci USA* **105**:9715–20. doi:10.1073/pnas.0803697105

Raap AK, van de Corput MPC, Vervenne RAM, van Gijlswijk RPM, Tanke HJ, Wiegant J. 1995. Ultra-sensitive FISH using peroxidase-mediated deposition of biotin- or fluorochrome tyramides. *Hum Mol Genet* **4**:529–534. doi:10.1093/hmg/4.4.529

Remedios R, Kennedy A, Zelikowsky M, Grewe BF, Schnitzer MJ, Anderson DJ. 2017. Social behaviour shapes hypothalamic neural ensemble representations of conspecific sex. *Nature* **550**:388–392. doi:10.1038/nature23885

Renier N, Adams EL, Kirst C, Wu Z, Azevedo R, Kohl J, Autry AE, Kadiri L, Umadevi Venkataraju K, Zhou Y, Wang VX, Tang CY, Olsen O, Dulac C, Osten P, Tessier-Lavigne M. 2016. Mapping of Brain Activity by Automated Volume Analysis of Immediate Early Genes. *Cell* **165**:1789–1802. doi:10.1016/j.cell.2016.05.007

Saudou F, Amlaiky N, Plassat JL, Borrelli E, Hen R. 1990. Cloning and characterization of a Drosophila tyramine receptor. *The EMBO Journal* **9**:3611–3617. doi:10.1002/j.1460-2075.1990.tb07572.x

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**:676–682. doi:10.1038/nmeth.2019

Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* **9**:671–675. doi:10.1038/nmeth.2089

Seelig JD, Chiappe ME, Lott GK, Dutta A, Osborne JE, Reiser MB, Jayaraman V. 2010. Two-photon calcium imaging from head-fixed Drosophila during optomotor walking behavior. *Nat Methods* **7**:535–540. doi:10.1038/nmeth.1468

Shao L, Chung P, Wong A, Siwanowicz I, Kent CF, Long X, Heberlein U. 2019. A Neural Circuit Encoding the Experience of Copulation in Female Drosophila. *Neuron* **102**:1025-1036.e6. doi:10.1016/j.neuron.2019.04.009

Shen P, Wan X, Wu F, Shi K, Li J, Gao H, Zhao L, Zhou C. 2023. Neural circuit mechanisms linking courtship and reward in Drosophila males. *Current Biology* **33**:2034-2050.e8. doi:10.1016/j.cub.2023.04.041

Sinigaglia C, Thiel D, Hejnol A, Houliston E, Leclère L. 2018. A safer, urea-based in situ hybridization method improves detection of gene expression in diverse animal species. *Developmental Biology* **434**:15–23. doi:10.1016/j.ydbio.2017.11.015

Stockinger P, Kvitsiani D, Rotkopf S, Tirián L, Dickson BJ. 2005. Neural circuitry that governs Drosophila male courtship behavior. *Cell* **121**:795–807. doi:10.1016/j.cell.2005.04.026

Takayanagi-Kiya S, Kiya T. 2019. Activity-dependent visualization and control of neural circuits for courtship behavior in the fly *Drosophila melanogaster*. *Proc Natl Acad Sci USA* **116**:5715–5720. doi:10.1073/pnas.1814628116

Takayanagi-Kiya S, Shioya N, Nishiuchi T, Iwami M, Kiya T. 2023. Cell assembly analysis of neural circuits for innate behavior in *Drosophila melanogaster* using an immediate early gene *stripe / egr-1*. *Proc Natl Acad Sci USA* **120**:e2303318120. doi:10.1073/pnas.2303318120

Turner GC, Bazhenov M, Laurent G. 2008. Olfactory Representations by Drosophila Mushroom Body Neurons. *Journal of Neurophysiology* **99**:734–746. doi:10.1152/jn.01283.2007

von Philipsborn AC, Liu T, Yu JY, Masser C, Bidaye SS, Dickson BJ. 2011. Neuronal Control of Drosophila Courtship Song. *Neuron* **69**:509–522. doi:10.1016/j.neuron.2011.01.011

Vosshall LB, Stocker RF. 2007. Molecular architecture of smell and taste in Drosophila. *Annual review of neuroscience* **30**:505–33. doi:10.1146/annurev.neuro.30.051606.094306

Wang L, Anderson DJ. 2010. Identification of an aggression-promoting pheromone and its receptor neurons in Drosophila. *Nature* **463**:227–31. doi:10.1038/nature08678

Wang L, Han X, Mehren J, Hiroi M, Billeter J-C, Miyamoto T, Amrein H, Levine JD, Anderson DJ. 2011. Hierarchical chemosensory regulation of male-male social interactions in Drosophila. *Nat Neurosci* **14**. doi:10.1038/nn.2800

Watanabe K, Chiu H, Pfeiffer BD, Wong AM, Hoopfer ED, Rubin GM, Anderson DJ, Watanabe K, Chiu H, Pfeiffer BD, Wong AM, Hoopfer ED, Rubin GM. 2017. A Circuit Node that Integrates Convergent Input from Neuromodulatory and Social Behavior-Promoting Neurons to Control Aggression in Drosophila Article A Circuit Node that Integrates Convergent Input from Neuromodulatory and Social Behavior-Promoti. *Neuron* **95**:1112-1128.e7. doi:10.1016/j.neuron.2017.08.017

Wilkie GS, Davis I. 1998. Visualizing mRNA by in situ hybridization using "high resolution" and sensitive tyramide signal amplification. *Technical Tips Online* **3**:94–97. doi:10.1016/S1366-2120(08)70110-7

Wilson RI, Turner GC, Laurent G. 2004. Transformation of Olfactory Representations in the Drosophila Antennal Lobe. *Science* **303**:366–370. doi:10.1126/science.1090782

Wu YE, Pan L, Zuo Y, Li X, Hong W. 2017. Detecting Activated Cell Populations Using Single-Cell RNA-Seq. *Neuron* **96**:313-329.e6. doi:10.1016/j.neuron.2017.09.026

Yamamoto D, Koganezawa M. 2013. Genes and circuits of courtship behaviour in Drosophila males. *Nat Rev Neurosci* **14**:681–92. doi:10.1038/nrn3567

Yang B, Treweek JB, Kulkarni RP, Deverman BE, Chen C-K, Lubeck E, Shah S, Cai L, Gradinaru V. 2014. Single-Cell Phenotyping within Transparent Intact Tissue through Whole-Body Clearing. *Cell* **158**:945–958. doi:10.1016/j.cell.2014.07.017

Yong Wan, Otsuna H, Chi-Bin Chien, Hansen C. 2009. An interactive visualization tool for multi-channel confocal microscopy data in neurobiology research. *IEEE Trans Visual Comput Graphics* **15**:1489–1496. doi:10.1109/TVCG.2009.118

Yu JY, Kanai MI, Demir E, Jefferis GSXE, Dickson BJ. 2010. Cellular Organization of the Neural Circuit that Drives Drosophila Courtship Behavior. *Current biology : CB* **20**:1602–1614. doi:10.1016/j.cub.2010.08.025

Ziv Y, Burns LD, Cocker ED, Hamel EO, Ghosh KK, Kitch LJ, Gamal AE, Schnitzer MJ. 2013. Long-term dynamics of CA1 hippocampal place codes. *Nat Neurosci* **16**:264–266. doi:10.1038/nn.3329

Figure 3

A. optoHI-FISH / 50Hz / 30s x 2

B. P1a > Chrimson

C. Photostimulation of P1ª Neurons → Counter staining / Hr38 HCR → Registration & Averaging

D. Time after Photostimulation (min)
P1ª / Hr38
0, 5, 15, 30, 60

E. Hr38 Brp
Control / P1a > Chrimson

F. Hr38
Anterior View / Posterior View
PAM, MB, pC
Low — High

G. P1 activated / Control
pCd: 41A01 Hr38
Kenyon cells: OK107 Hr38
PAM Neurons: 0273 Hr38
Ddc Hr38

Figure 4

A. HI-catFISH

Experimental Fly → 1st Behavior: Male (Aggression) / Female (Courtship) → 2nd Behavior: Male (Aggression) / Female (Courtship) → HCR Analysis: Hr38 Expression Intron / Exon Probes

B. Neuronal Types | Hr38 Expression Level | Hr38 signal pattern

C. Behavior Order: Female→Male, Male→Female, Male→Male, Female→Female

D.
- Female→Male: (112 cells / 7 individuals)
- Male→Female: (183 cells / 10 individuals)
- Male→Male: (109 cells / 7 individuals)
- Female→Female: (55 cells / 5 individuals)

N.R.: Not Responded
Nuc: Nuclear
Cyto: Cytoplasmic
Both: Nuclear & Cytoplasmic