Title: Articular cartilage corefucosylation regulates tissue resilience in osteoarthritis

Short title: Articular cartilage corefucosylation and tissue resilience in OA

Kentaro Homan[1], Tomohiro Onodera[1]*, Hisatoshi Hanamatsu[2], Jun-ichi Furukawa[1, 2],

Daisuke Momma[3], Masatake Matsuoka[1], Norimasa Iwasaki[1]

[1] Department of Orthopaedic Surgery, Faculty of Medicine and Graduate School of Medicine,

Hokkaido University, Kita 15, Nishi 7, Kita-ku, Sapporo 060-8638, Japan

[2] Institute for Glyco-core Research (iGCORE), Nagoya University, Nagoya 464-8601, Japan

[3] Center for Sports Medicine, Hokkaido University Hospital, Kita 14, Nishi 5, Sapporo 060-

8638, Japan.

*Corresponding author: Tomohiro Onodera

Department of Orthopaedic Surgery, Faculty of Medicine and Graduate School of Medicine

Hokkaido University, Kita 15, Nishi 7, Kita-ku, Sapporo 060-8638, Japan

Telephone: +81-11-706-2111

Email: tomozou@med.hokudai.ac.jp

Abstract

This study aimed to investigate the glycan structural changes that occur before histological

degeneration in osteoarthritis (OA) and to determine the mechanism by which these glycan

conformational changes affect cartilage degeneration. An OA model was established in

rabbits using mannosidase injection, which reduced high-mannose type N-glycans and led to

cartilage degeneration. Further analysis of glycome in human OA cartilage identified specific

corefucosylated N-glycan expression patterns. Inhibition of N-glycan corefucosylation

resulted in unrecoverable cartilage degeneration, while cartilage-specific blocking of

corefucosylation led to accelerated development of aging-associated and instability-induced

OA models. We conclude that α1,6 fucosyltransferase is required postnatally to prevent

preosteoarthritic deterioration of articular cartilage. These findings provide a novel definition

of early OA and identify glyco-phenotypes of OA cartilage, which may distinguish

individuals at higher risk of progression.

Keywords: early osteoarthritis, glyco-phenotypes, α1,6 fucosyltransferase, cartilage glycome

INTRODUCTION

Osteoarthritis (OA), a degeneration of joint components, such as proteoglycans (PGs) and

collagen, destroys cartilage and impairs joint function (1). The global prevalence of OA is

estimated to be >500 million, which is a significant socioeconomic burden (2). Therefore,

preventive measures must be implemented to reduce the future disease burden and disability.

Once the cartilage is destroyed, it cannot be repaired homogeneously and progresses

irreversibly thereafter (3, 4). Since articular cartilage has no blood vessels and is not

innervated, it is difficult for symptoms to appear. By the time clinical symptoms appear, the

condition is often already considerably advanced (5). Therefore, from a preventive

perspective, it is necessary to define and detect "pre-OA," the stage before the onset of early

OA (6, 7). Pre-OA is a knee with no pain, no motor function problems, and no arthroscopic or

structural changes on radiography, only emerging cellular effects. Despite many studies, there

is no gold standard for OA biomarkers (8). It is important to characterize the disease state

before OA to shift the paradigm from mitigation to late-onset disease prevention.

In the very early stages of OA, before cartilage matrix degeneration, cartilage destruction by

enzymatic processes that degrade glycoproteins, PGs, and collagen, in that order, is underway

(9). Collagenolytic activity is largely unchanged on intact articular surfaces (10, 11), and

glycoprotein degradation is thought to be an early transient event in the enzymatic process of

cartilage destruction because hexose degradation of glycoproteins promotes degradation of

the protein portion of the molecule by proteinases (12). Additionally, cartilage damage is

reversible if type 2 collagen breakdown is mild (13). Structural change analysis in

glycoproteins is necessary for the early stages of OA but has not yet been performed

sufficiently.

Matsuhashi *et al.*'s study showed that structural changes in N-glycans occurred in

degenerated cartilage in a rabbit OA model before tissue changes occurred (14). Urita *et al.*

demonstrated that these N-glycans in degenerated mice and human cartilage are high-

mannose type N-glycans (15). Furthermore, among several glycosidases, mannosidase

activity was very high in OA joints, and its activity was high before the progression of

cartilage fibrosis (16). This suggests that high-mannose type N-glycan plays an important

role in OA pathogenesis. However, whether changes in the glycan structure are involved in

OA pathogenesis is unclear.

This study analyzed the glycan structural changes that preceded OA degeneration and

elucidated the mechanism of glycan structural changes in cartilage degeneration. For this

purpose, the effects of mannosidase, a specific degrading enzyme of high-mannose type N-

glycans associated with OA, on cartilage tissues and its molecular mechanisms were

analyzed.

RESULTS

Mannosidase injection caused early OA-like change in vivo

A decrease in Safranin O staining in the right femoral condyle of the α-mannosidase-injected

knee after 4 weeks was observed (Fig. 1A, B). Terminal deoxynucleotidyl transferase dUTP

nick end labeling (TUNEL) staining was enhanced in the superficial layer compared with that

in control. Articular cartilage chondrocytes exposed to α-mannosidase were not arranged in

columns in the middle zone. The surfaces of each group were smooth and had a good

macroscopic appearance (Fig. 1C). None of the groups showed a decrease in type 2 collagen

staining, and the orientation was maintained (Fig. 1D). To achieve optimal activity, the pH

range of 4 to 5 is ideal for mannosidase. In a physiological saline solution with a pH of 7 and

a concentration of 100 μg/ml, it exhibited specific activity. Nevertheless, at a concentration of

1.0 μg/ml, this activity was no longer present (S1 Figure). The mean histological score of the

α-mannosidase group was 0.83, indicating mild OA (Fig. 1E). The control group was

unchanged. Cartilage degeneration during intermittent intra-articular administration of

mannosidase was reversed upon discontinuation (Fig. 1F).

Depletion of high-mannose type N-glycans in chondrocytes leads to cartilage

degradation ex vivo

To quantify the degeneration and resilience promoted by the injection of α-mannosidase, the

cultured cartilage was harvested at different time points (Fig. 2A). Concanavalin A (Con A)

reactivity declined in articular cartilage treated with α-mannosidase (Fig. 2B). The

application of α-mannosidase stimulated the release of PG from cultured cartilage in

association with extracellular matrix degradation (Fig. 2C) and showed significantly higher

spontaneous nitric oxide (NO) release than the control over time (Fig. 2F). Mannosidase-

exposed cartilage showed an increase in TUNEL-positive cells in chondrocytes in the deep

zone and type 10 collagen-positive cells in superficial chondrocytes; however, these changes

did not occur in cytokine-stimulated cartilage (S2 Figure). Mannosidase exposure increased

Adamts5 expression and suppressed *Mmp13* expression and anabolic factors (Fig. 3E). In the

articular cartilage that was exposed to freeze-thaw treatment before culture, no differences in

PG loss were observed between the α-mannosidase-stimulated cartilage and non-stimulated

cartilage (Fig. 2E). Excluding mannosidase loading, PG leakage and Safranin O staining were

restored (Fig. 2D), but Con A staining loss was not. Sialidase addition did not cause PG

leakage or NO production (S3 Figure), and the aggrecan lost in the catabolic phase was not

replenished in the subsequent anabolic phase.

Corefucosylated N-glycan was formed in resilient cartilage

Glycoblotting analysis revealed a significant difference in the N-glycan profiles (Fig. 3A and

S1 Table). Plotting m/z after mannosidase exposure against m/z before mannosidase exposure

demonstrated that only N-glycans were affected by enzyme dissociation from the y = x line

(Fig. 3B, S2 Table). In both denatured and reverted cartilages, Man8-9GlcNAc2 was

decreased, and Man5-6GlcNAc2 was increased by the action of mannosidase, which has a

high specificity for the terminal mannosidic residues of glycans (Fig. 3C, D). In contrast,

corefucosylated N-glycans, which were formed in the same biosynthesis pathway, were also

found (Fig. 3D). Similarly, the N-glycan profiles of isolated chondrocytes showed the

presence of this corefucosylated structure (S4 Figure). The expression of *Fut8*, the only

enzyme involved in corefucosylation, increased significantly in mannosidase-treated cartilage

(Fig. 3E).

Deficiency of corefucosylated glycans in articular cartilage inhibits recovery from cartilage damage and promotes cartilage degeneration

We first examined whether the absence of core fucose affects the resilience of cartilage

degeneration. Mice lacking *Fut8* exhibited severe growth retardation and died shortly after

birth. Therefore, we decided to generate *Fut8* conditional knockout (cKO) mice (Fig. 4A, B).

The tissue specificity of recombination was confirmed by polymerase chain reaction (PCR)

analysis of mouse genomic DNA that detected either the intact $Fut8^{loxP}$ allele or the Cre-

recombined $Fut8^{\Delta}$ allele in repeated crosses; N-glycan analysis of cartilage demonstrated the

loss of corefucosylated sugar chains (Fig. 4C). The proportion of high-mannose type N-

glycan content was unchanged in cartilage lacking the *Fut8* gene (S5 Figure). Next, we used

real-time (RT)-PCR to determine relative *Fut8* messenger (m)-RNA levels in the cartilage of

Col2-$Fut8^{-/-}$ mice and their wild-type (WT) littermates ($Fut8^{loxP/loxP}$). Real-time polymerase

chain reaction (RT-PCR) analysis showed that *Fut8* mRNA expression was reduced by >99%

in the articular cartilage of Col2-$Fut8^{-/-}$ mice compared to that in their WT littermates (P <

0.01; n = 3 mice per group) (Fig. 4D). Chondrocyte-surface corefucosylation levels were

analyzed by lectin staining using *Pholiota squarrosa* lectin (PhoSL), which specifically

recognizes α1-6 core fucose (17). PhoSL lectin staining was completely absent in the articular

cartilage of *Fut8* cKO mice, in contrast to the positive staining observed in *Fut8* flox mice

under physiological conditions (Fig. 4E). Enzymatic degradation of high-mannose type N-

glycans on the cartilage in *Fut8*-deficient mice exacerbated cartilage degeneration (Fig. 4E,

F). In addition, reversible matrix repair by enzyme removal observed in control cartilage was

not observed in *Fut8*-deficient cartilage explant cultures.

Loss of corefucosylated glycans in articular cartilage leads to premature OA

Col2-$Fut8^{-/-}$ mice developed and grew normally without major organ abnormalities of major organs and could not be distinguished from their WT littermates. The whole skeletons of newborn mice stained with alcian blue and alizarin red did not differ in appearance between the genotypes (Fig. 5A). Col2-$Fut8^{-/-}$ mice showed a lower body weight than their WT littermates, as indicated by the growth curve (Fig. 5B). We analyzed the development of instability-induced OA changes in $Fut8^{loxP/loxP}$ and $Fut8$ cKO mice. Sham operations produced no significant changes in OA in either mouse genotype. Eight weeks after instability-inducing surgery, the joints of WT mice developed changes in OA, such as cartilage erosion and a reduction in Safranin O staining and chondrocyte number. Deletion of FUT8 results in more severe changes in OA. In the joints of $Fut8$ cKO mice, 8 weeks after surgery, the noncalcified zone (upper layer above the tidemark) was almost completely lost. The number of chondrocytes was significantly reduced (Fig. 5C). Quantitative assessment using the OA Research Society International (OARSI) score supported these histological findings (difference: -2.900; $Fut8^{flox/flox}$/destabilization of the medial meniscus (DMM) versus $Col2a1$-Cre; $Fut8^{flox/flox}$/DMM, 95% confidence interval [CI] for the difference: -5.027 to -0.7734) (Fig. 5D). Changes in OA progressed earlier in the aging models of $Fut8$-deficient mice than in control mice (WT and floxed mice). The mice were followed up until 15 months of age to assess the spontaneous development of OA with aging (Fig. 5E). There were no apparent changes of OA in the knee joints of either mouse genotype at 3 months of age (Fig. 5F). At 4 months of age, significant changes in OA were detected in the knee joints of the $Fut8$ cKO mice. The cartilage surface integrity was no longer maintained at 9 months. These OA changes progressed more in $Fut8$ cKO mice than in their WT littermates at 15 months. These histological findings were quantitatively confirmed by the Osteoarthritis Research Society International (OARSI) scores (difference: -2.000; WT versus $Col2a1$-$Cre; Fut8^{flox/flox}$,

95% CI for the difference: -3.723 to -0.2771 at 4 months, difference: -2.700; WT versus

Col2a1-Cre; Fut8^{flox/flox}, 95% CI for the difference: -4.423 to -0.9771 at 9 months, and

difference: -6.700; WT versus *Col2a1-Cre; Fut8^{flox/flox}*, 95% CI for the difference: -8.423 to -

4.977 at 15 months).

Altered glycosylation of human OA cartilage based on comprehensive glycan analysis

Total glycome profiling of human OA cartilage is shown in Fig. 6A and S3 Table. The

amount of high-mannose type N-glycan was significantly decreased, the same as previously

reported in OA cartilages (Fig. 6A) (15). Focusing on fucosylation, most of the

complex/hybrid type glycans were modified with fucose. The fucosylation that occurs at the

innermost N-acetylglucosamine (GlcNAc) of N-glycans could be recognized with a lectin

from the mushroom Pholiota squarrosa (PhoSL) and was able to detect the expression of the

corefucose structure. Therefore, we suspected that the stainability of PhoSL is increased in

OA cartilage. As expected, the PhoSL staining of OA cartilage was significantly increased

relative to that of a healthy control (Fig. 6B). We performed principal component analysis

(PCA) on 1110 glycomic data, discriminating OA from healthy cartilages along the first

principal component (PC1) axis (Fig. 6C). Hierarchical cluster analysis showed that clades

containing corefucosylated glycans were clustered (Fig. 6D), and N-13 ((Hex)1 (HexNAc)1

(Fuc)1 + (Man)3(GlcNAc)2) has the largest squared correlation with its cluster component.

DISCUSSION

To test the hypothesis that structural changes in high-mannose type N-glycans may induce

OA, we tested the effect of mannosidase loading on the structure of high-mannose type N-

glycans. Consequently, modifying the structure of high-mannose type N-glycans by

mannosidase loading induced OA-like histological alterations. Furthermore, the OA-like

changes induced by mannosidase were recovered by enzyme removal. These results suggest

that structural alterations of high-mannose type N-glycans lead to OA-like histological

changes in articular cartilage but do not necessarily cause the pathogenesis of eventual OA.

Subsequently, glycoblotting analysis revealed that core-fucose-containing glycans underwent

complementary structural alterations per high-mannose type N-glycans. This suggests that the

structural change of core-fucose-containing glycans may have some compensatory

mechanism for OA-like alterations caused by the structural change of high-mannose type N-

glycans. Furthermore, under the regulation of *Fut8* expression, which regulates the synthesis

of core fucose, mannosidase was found to induce irreversible OA. These results provide the

first evidence that structural alterations in glycans in the articular cartilage regulate eventual

OA development.

Two key targets of cartilage degeneration during OA are type II collagen, a major substrate of

matrix metalloproteinase (MMP)-13 (18), and aggrecan, a PG with glycosaminoglycan side

chains of chondroitin and keratin sulfate (19, 20). The degradation of non-collagenous

molecules, such as aggrecan, occurs before the degradation of type II collagen in the early

stages of OA (20, 21). The histological analysis in this study revealed that the alteration in the

structure of high-mannose type N-glycans resulted in a decrease in Safranin O staining (a

decrease in glycosaminoglycans and/or PGs), whereas there was no change in collagen

staining. These findings are consistent with alterations in the early stages of OA.

Furthermore, we observed that removing mannosidase could cause PG loss to the control

level. Karsdal *et al.* demonstrated that articular cartilage degradation is reversible in the

presence of aggrecanase and concluded that the repair ability is not impaired before the

degradation of aggrecan and type II collagen by MMPs (22). Reimann *et al.* observed that the

glycosaminoglycans loss in articular cartilage is reversible and an early sign of OA before the

"state of no return." (23) The present results are consistent with findings of previous reports

on early OA and suggest that the mannosidase-loaded model can be considered an early OA-

like alteration.

Since the structural change in high-mannose type N-glycans alone did not cause irreversible

OA, we investigated other regulatory mechanisms related to high-mannose type N-glycans

that could induce irreversible OA. The expression of corefucosylated N-glycans on a series of

biosynthetic pathways was upregulated with structural changes in high-mannose type N-

glycans. FUT8 is known to corefucosylate small oligomannose N-glycans (Man4-

Man5GlcNAc2) but not large oligomannose N-glycans, such as Man8-Man9GlcNAc2 (24).

This suggests that small oligomannose N-glycans trimmed by mannosidase could activate

Fut8. *Fut8* is also upregulated in human OA chondrocytes (25, 26), and increased expression

of these corefucosylated N-glycans has been observed during late chondrocyte differentiation

(27). These findings suggested that these structural changes are related to OA-related

glycosylation.

To clarify the relationship between the alteration of corefucosylated N-glycans and cartilage

degeneration, *Fut8* cKO mice were used. Since the results of the ex vivo freeze-thaw model

showed that cartilage degeneration was induced by changes in the glycan structure of

chondrocytes, we used chondrocyte-specific KO mice. As a result, the reversibility of the

cartilage degeneration model was diminished. This finding suggests that *Fut8*-mediated

corefucosylation in chondrocytes plays a function in promoting glycan repair. Furthermore, to

directly determine whether *Fut8* was involved in the inhibition of OA progression, an aging

model was investigated. The results showed that conditional KO of *Fut8* in chondrocytes

hastened the onset of OA. This suggests that cartilage corefucosylation associated with FUT8

plays a protective role against the attenuation of type II collagen in the OA process.

Finally, the expression of α1,6-linked core fucose on N-glycans in human cartilage is

associated with OA, and corefucosylation was found to be one of the characteristics in OA

cartilage. Total cellular glycome analysis, including N- and O-linked glycans derived from

glycoproteins, GSL-glycans, GAGs, and fOSs, is informative for analyzing cell-specific

characteristics (28−30). Glycome profiling enabled the detection of corefucosylated N-

glycans by sequential clustering of degenerated cartilage characteristics. Core fucose

deficient mice generated by ablating the α1,6 fucosyltransferase enzyme, FUT8, were

reported to have suppressed phosphorylation of Smad and increased expression of MMPs

(matrix metalloproteinases) due to decreased binding of TGF-β ligand caused by the lack of

core fucose addition to the TGF-β type II receptor (31, 32). The present study shows that

treatment of cartilage with mannosidase caused corefucosylation, suppression of *Mmp13*, and

increased expression of *Tgf-β*. Moreover, treatment of FUT8-knockout cartilage with

mannosidase resulted in the suppression of *Tgf-β* expression and increased *Mmp13*

expression. These results suggest that corefucosylation following the loss of high-mannose

type N-glycans may provide chondroprotective effects via the TGF-β signaling pathway,

which may have been lost by FUT8-knockout.

One limitation of this study was that we did not have access to cartilage samples from

immediately before the onset of OA; thus, we do not know the expression of Fut8 in human

cartilage in the early stage of OA onset and its influence on extracellular matrix degradation

under homeostatic conditions. Obtaining cartilage specimens continues to be challenging,

especially for those with early stages of OA.

In summary, we show for the first time a key role of FUT8 and glycan-dependent signaling

mediators in extracellular matrix resilience associated with cartilage corefucosylation during

early OA.

METHODS

Intra-articular injection of α-mannosidase in the knee

Japanese White rabbits weighing 3.10 ± 0.15 kg purchased from a professional breeder (Japan SLC Inc., Hamamatsu, Japan) were used for this study according to established ethical guidelines approved by the local animal care committee. Animals were anesthetized with 10 mg/kg of intravenous ketamine, followed by isoflurane in oxygen gas. Both knees in each rabbit were shaved, prepared, and draped in a sterile fashion. α-Mannosidase from Canavalia ensiformis (Jack bean) (Sigma, St Louis, MO, USA) was dissolved in saline. Five hundred microliter volumes containing 1.9 units per ml (0.05 mg) of mannosidase solution were injected into the joint cavity of the right knee. α-Mannosidase from Canavalia ensiformis (Jack bean) has been widely used as a tool for glycan analysis, which has a high specificity towards terminal mannosidic residues of glycans and participates in mannose trimming reactions (33). The measurement of its performance in a saline solution was evaluated using the QuantiChrom α-Mannosidase Assay Kit (BioAssay Systems, Hayward, CA, USA). This colorimetric assay uses 4-nitrophenyl-α-d-mannopyranoside as a substrate to identify the generation of 4-nitrophenol. The following doses of α-Mannosidase were administered: 0.5 ml at 0.01 μg/ml, 1.0 μg/ml, or 100 μg/ml. Simultaneously, the contralateral knee joint received an equal volume of saline solution. Each joint was injected once or four times in alternate weeks. The animals were euthanized at 4 and 16 weeks postoperatively. Each section of the knee joint was stained with hematoxylin and eosin or Safranin O/fast green, as well as with immunohistochemical staining for type II collagen and TUNEL. Two investigators quantified the OA severity using the OARSI scoring method.

Evaluation of collagen orientation by a polarized light microscope

To evaluate the collagen orientation of the repaired tissue at 16 weeks postoperatively, a

supplemental evaluation using a polarized light microscope (PLM) (ECLIPSE E600 POL;

Nikon, Tokyo, Japan) was conducted (34). This evaluation was performed on HE-stained

sections from each group. Two cross polarizers were used so that highly ordered collagen

perpendicular to the articular surface appeared bright, while collagen that was not ordered

(nonbirefringent) appeared dark. The contrast of fibrils more parallel to the articular surface

was darker than collagen oriented perpendicular to the articular surface. To observe the

predominant direction of the birefringent regions and confirm the lack of orientation in the

nonbirefringent regions, the sections were rotated at 0°, 45°, and 90° for the fixed filters.

Microscopic images were acquired using a digital camera (DS-5M-L1, Nikon, Tokyo, Japan).

Each sample was examined independently by an experienced PLM user.

Ex vivo analysis of deletion of high-mannose type N-glycan in cartilage

According to the established ethical guidelines approved by the local animal care committee,

cartilage catabolism was analyzed by culturing femoral head cartilage of 4-week-old C57BL6

male mice with α-mannosidase (1.9 U/mL) ex vivo (35, 36). Mannosidase was removed on

day 3, and organ culture was continued until day 6. To evaluate cartilage degradation,

quantification of PG release from cartilage explants was performed using the

dimethylmethylene blue assay (36, 37) and exposure to sialidase (1.9 U/mL) and interleukin-

1 beta (Cat #. 300-01A; BioLegend, San Diego, CA, USA) (10 ng/mL) was also used as

another control. The NO concentration in the culture medium was measured using the Griess

reagent system (Promega, Tokyo, Japan). To determine the direct effects of mannosidase on

the matrix, some cartilage explants were subjected to freeze-thaw cycles in liquid nitrogen to

destroy chondrocytes present in the superficial layer of the cartilage (22, 38). After the

procedure, the cells were replaced with the same culture medium described above. Each

tissue sample was stained with Safranin O. Lectin staining with Con A (Wako Pure Chemical

Industries, Richmond, VA, USA) and PhoSL (J-Oil Mills Inc., Tokyo, Japan) was used to

detect the high-mannose type N-glycans. Staining was performed using the ABC Kit (Vector

Labs, Burlingame, CA, USA), and the sections were counterstained with HE.

RNA extraction and quantitative RT-PCR analysis

Total RNA was extracted from the samples using an RNeasy Mini Kit (Qiagen, Hilden,

Germany). For complementary DNA synthesis, 1.0 μg of RNA was reverse transcribed using

random hexamer primers (Promega, Tokyo, Japan) and ImProm II reverse transcriptase

(Promega). Quantitative RT-PCR was performed using a Thermal Cycler Dice Real Time

System II (Takara Bio Inc., Otsu, Japan) and SYBR Premix Ex Taq II (Takara, Shiga, Japan)

with gene-specific primers (S4 Table) in 15 μl of the mixture, following the manufacturer's

instructions. Quantitative data were normalized using peptidylprolyl isomerase A as an

endogenous reference gene and were calculated using the ΔΔCt method (39).

Generation of $Fut8^{loxp/loxp}$ Col2a1-Cre cKO mice

$Fut8^{loxp/loxp}$ mice were provided by Tohoku Pharmaceutical University (Sendai, Japan). To

interrupt the synthesis of corefucosylated N-glycans in cartilage, we generated mice with

knockout of the chondrocyte-specific Fut8 gene (Col2-$Fut8^{-/-}$ mice) by crossing $Fut8^{loxp/lox}$

mice with Col2a1-Cre–transgenic mice (40), in which Cre recombinase is expressed

specifically in chondrocytes under the Col2a1 promoter. Transgenic mice carrying the

Col2a1-Cre transgene (strain B6; SJL-Tg [Col2a1-Cre] 1Bhr/J; stock number 003554) were

obtained from the Jackson Laboratory (Bar Harbor, ME, USA). Homozygous $Fut8^{loxp/loxp}$

mice were bred with $Col2a1$-Cre mice, and offspring with the genotype $Fut8^{loxp/+}$ Col2a1-

Cre were bred with $Fut8^{loxp/loxp}$ mice to obtain the $Fut8^{loxp/loxp}$ Col2a1-Cre cKO experimental

group and $Fut8^{loxp/loxp}$ (flox) littermate controls. All animals were maintained at Hokkaido

University under the regulations of the Institutional Animal Care and Use Committee.

Skeletal examination

To observe the skeletal systems, whole skeletons of WT and cKO littermates were stained

with alcian blue and alizarin red, as previously described (41, 42). The newborns were

skinned, eviscerated, and fixed in 95% ethanol. Intact skeletons were stained with 150 mg/ml

of alcian blue 8GX in 75% (vol/vol) ethanol/20% acetic acid for 24–48 hours, and excess

tissues were removed by 1–2% KOH digestion for 2–3 days. The skeletal preparations were

stained with 75 mg/ml of alizarin red S in 1% KOH for 24 hours and cleared in graded

glycerol solutions (43).

Instability-induced OA model

An OA model was created in 10-week-old mice by destabilizing the knee joint as previously

described (44). With the mice under general anesthesia, we destabilized the right knee joint

by transection of the medial collateral ligament and removing the cranial half of the medial

meniscus using a microsurgical technique (DMM). A sham operation was performed on the

left knee joint using the same approach, without ligament transection and meniscectomy. For

histological assessment, the mice were sacrificed, and the entire knee joint was dissected 8

weeks postoperatively.

Age-associated OA model

At 3, 4, 9, and 15 months of age, the mice were sacrificed, and their entire knee joints were

dissected to assess the spontaneous development of OA as a model of age-associated OA (44,

45).

Human Cartilage Preparation

Acquisition and use of patient tissues were approved by the institutional review board (IRB) of Hokkaido University (approval number: 014-0144), and informed consent was obtained in advance. Samples were from patients undergoing total knee arthroplasty for clinically and radiologically diagnosed OA (n=5; age range 61–84 years, mean 72.4; male: female ratio 1:4). Age-matched control articular cartilage was from patients undergoing total hip arthroplasty for femoral neck fracture with no history of joint disease (n=5; age range 45–88 years, mean 61.2 years; male: female ratio 3:2). Each cartilage specimen was evaluated macroscopically by at least two veteran doctors to ensure that it had not suffered damage or degeneration and was treated within 9 hours of harvest in the operating room. Immediately upon receipt, isolated human cartilage was homogenized with a Polytron blender (Kinematica, Luzern, Switzerland), and then cold ethanol was added to separate the protein and lipid fractions.

Total cartilage glycomic analysis

N-glycans were released directly using cartilage lysates following deglycosylation by overnight treatment with peptide N-glycanase F (PNGase F, 2U) (Roche, Switzerland). The supernatants containing GSLs and fOSs were dried with a centrifugal evaporator. GSL-glycans were isolated by enzymatic digestion using EGCase I (Takara Bio Inc. Japan), whereas fOSs were recovered from an EGCase I-free fraction (46). Cartilage lysate was delipidated and digested by GAG disaccharides (47). Extracted glycoproteins were subjected to BEP for O-glycan analysis. Detailed procedures and materials are provided elsewhere (48). N-glycans, fOSs, GAGs, and GSL-glycans were subjected to glycoblotting (49). Purified N-glycans, fOSs, O-glycans, and GSL-glycan solutions were mixed with 2,5-dihydrobenzoic acid and subjected to Matrix-assisted laser desorption/ionization-time of flight mass

spectrometry (MALDI-TOF MS). 2AB-labeled GAG disaccharides were analyzed by HPLC.

The glycan compositions were manually determined by conducting database searches (i.e., a

compositional search of the UniCarbKB database (http://www.unicarbkb.org/query) for fOSs

and N- and O-glycans and of the SphinGOMAP database (http://www.sphingomap.org/) for

GSL-glycans). All previously deposited GSL-glycans in the SphinGOMAP database were

extracted and compiled as an in-house database to allow searching by the m/z value and/or

composition. The absolute quantification was obtained by comparative analyses between the

MS signal areas derived from each glycan and the internal standard.

Statistics

All data in this study are presented as the mean ± standard deviation and were repeated at

least three times unless otherwise indicated. Data analysis was performed using GraphPad

Prism 9 software version 9.4.1 (GraphPad Software, Inc., San Diego, CA, USA), and the

Welch t-test/Welch analysis of variance with subsequent use of the Tukey–Kramer multiple

comparison tests were used to determine significant differences between the groups. P-values

<0.05 were considered significant. JMP Pro version 16.2.0 (SAS Institute, Cary, NC, USA)

was used to reveal the relationships between each subject of sub glycomes; we performed

principal component analysis (PCA) and hierarchical cluster analysis.

Study approval

Acquisition and use of patient tissues were approved by the institutional review board (IRB)

of Hokkaido University (approval number: 014-0144), and informed consent was obtained in

advance. All animal studies were performed according to established ethical guidelines

approved by the local animal care committee.

Data availability

All data reported in this paper is available from the corresponding author upon request. This

paper does not report original code. Any additional information required to reanalyze the data

reported in this paper is available from the lead contact upon request.

ACKNOWLEDGMENTS

We thank the Iwasaki laboratory members for technical support, J. Gu (Tohoku

Pharmaceutical University) for the Flox-α1,6 fucosyltransferase (*Fut8*) mouse, Y. Kobayashi

(J-OIL MILLS, Inc.) for PhoSL-lectin. This work was funded by the Japan Society for the

Promotion of Science KAKENHI (Grant-in-Aid for Challenging Exploratory Research:

funding numbers: 16K15650, 19K22672 [to NI]; Grant-in-Aid for Research (B): funding

numbers: 22H03917 [to TO]; and Grant-in-Aid for Early-Career Scientists: funding number:

19K18516 [to KH]) and AMED (Grant Number JP22zf01270004h0002 [to NI, TO, KH]).

AUTHOR CONTRIBUTIONS

Conceptualization: NI

Methodology: KH, TO

Investigation: KH, TO, HH, JF, DM, MM, NI

Visualization: KH, HH

Funding acquisition: KH, NI, TO

Project administration: KH, TO, NI

Supervision: TO, NI

Writing – original draft: KH, TO

Writing – review & editing: KH, TO, HH, JF, DM, MM, NI

CONFLICT-OF-INTEREST STATEMENT: The authors have declared that no conflict of

interest exists.

References

1. Grillet B, et al. Matrix metalloproteinases in arthritis: towards precision medicine. *Nat.*

Rev. Rheumatol. [published online ahead of print: May 2023]; doi:10.1038/s41584-023-

00966-w

2. Safiri S, et al. Global, regional and national burden of osteoarthritis 1990-2017: a

systematic analysis of the Global Burden of Disease Study 2017. *Ann. Rheum. Dis.*

2020;79(6):819−828.

3. Thorstensson CA, et al. Natural course of knee osteoarthritis in middle-aged subjects with

knee pain: 12-year follow-up using clinical and radiographic criteria. *Ann. Rheum. Dis.*

2009;68(12):1890−1893.

4. Huey DJ, Hu JC, Athanasiou KA. Unlike bone, cartilage regeneration remains elusive.

Science 2012;338(6109):917−921.

5. Glyn-Jones S, et al. Osteoarthritis. *Lancet* [published online ahead of print: March 3,

2015]; doi:10.1016/S0140-6736(14)60802-3

6. Chu CR. Defining Pre-Osteoarthritis Is Key to Prevention. *Cartilage* 2016;7(2):204.

7. Mahmoudian A, et al. Early-stage symptomatic osteoarthritis of the knee - time for

action. *Nat. Rev. Rheumatol.* 2021;17(10):621−632.

8. Henrotin Y. Osteoarthritis in year 2021: biochemical markers. *Osteoarthritis Cartilage*

2021;S1063-4584(21)00960−2.

467 9. Vignon E, et al. Screening of degradative enzymes from articular cartilage in

468  experimental osteoarthritis. *Clin. Rheumatol.* 1987;6(2):208–214.

469 10. Brew CJ, et al. Gene expression in human chondrocytes in late osteoarthritis is changed

470  in both fibrillated and intact cartilage without evidence of generalised chondrocyte

471  hypertrophy. *Ann. Rheum. Dis.* 2010;69(1):234–240.

472 11. Otero M, et al. Human chondrocyte cultures as models of cartilage-specific gene

473  regulation. *Methods Mol. Biol. Clifton NJ* 2012;806:301–336.

474 12. Goldring MB. Chondrogenesis, chondrocyte differentiation, and articular cartilage

475  metabolism in health and osteoarthritis. *Ther. Adv. Musculoskelet. Dis.* 2012;4(4):269–

476  285.

477 13. van Meurs JB, et al. Kinetics of aggrecanase- and metalloproteinase-induced

478  neoepitopes in various stages of cartilage destruction in murine arthritis. *Arthritis*

479  *Rheum.* 1999;42(6):1128–1139.

480 14. Matsuhashi T, et al. Alteration of N-glycans related to articular cartilage deterioration

481  after anterior cruciate ligament transection in rabbits. *Osteoarthritis Cartilage*

482  2008;16(7):772–778.

483 15. Urita A, et al. Alterations of high-mannose type N-glycosylation in human and mouse

484  osteoarthritis cartilage. *Arthritis Rheum.* 2011;63(11):3428–3438.

485 16. Snelling S, et al. A gene expression study of normal and damaged cartilage in

486  anteromedial gonarthrosis, a phenotype of osteoarthritis. *Osteoarthr. Cartil. OARS*

487  *Osteoarthr. Res. Soc.* 2014;22(2):334–343.

488 17. Kobayashi Y, et al. A novel core fucose-specific lectin from the mushroom Pholiota

489  squarrosa. *J. Biol. Chem.* 2012;287(41):33973–33982.

18. Wu W, et al. Sites of collagenase cleavage and denaturation of type II collagen in aging

and osteoarthritic articular cartilage and their relationship to the distribution of matrix

metalloproteinase 1 and matrix metalloproteinase 13. *Arthritis Rheum.*

2002;46(8):2087−2094.

19. Song R-H, et al. Aggrecan degradation in human articular cartilage explants is mediated

by both ADAMTS-4 and ADAMTS-5. *Arthritis Rheum.* 2007;56(2):575−585.

20. Malfait A-M, et al. Inhibition of ADAM-TS4 and ADAM-TS5 prevents aggrecan

degradation in osteoarthritic cartilage. *J. Biol. Chem.* 2002;277(25):22201−22208.

21. Aigner T, et al. Anabolic and catabolic gene expression pattern analysis in normal versus

osteoarthritic cartilage using complementary DNA-array technology. *Arthritis Rheum.*

2001;44(12):2777−2789.

22. Karsdal MA, et al. Cartilage degradation is fully reversible in the presence of

aggrecanase but not matrix metalloproteinase activity. *Arthritis Res. Ther.*

2008;10(3):R63.

23. Reimann I, Christensen SB, Diemer NH. Observations of reversibility of

glycosaminoglycan depletion in articular cartilage. *Clin. Orthop.* 1982;(168):258−264.

24. Yang Q, Wang L-X. Mammalian α-1,6-Fucosyltransferase (FUT8) Is the Sole Enzyme

Responsible for the N-Acetylglucosaminyltransferase I-independent Core Fucosylation

of High-mannose N-Glycans. *J. Biol. Chem.* 2016;291(21):11064−11071.

25. Toegel S, et al. Glycophenotyping of osteoarthritic cartilage and chondrocytes by RT-

qPCR, mass spectrometry, histochemistry with plant/human lectins and lectin

localization with a glycoprotein. *Arthritis Res. Ther.* 2013;15(5):R147.

26. Yu H, et al. Characterization of aberrant glycosylation associated with osteoarthritis

based on integrated glycomics methods. *Arthritis Res. Ther.* 2023;25(1):102.

27. Homan K, et al. Alteration of the Total Cellular Glycome during Late Differentiation of

Chondrocytes. *Int. J. Mol. Sci.* 2019;20(14):E3546.

28. Fujitani N, et al. Total cellular glycomics allows characterizing cells and streamlining

the discovery process for cellular biomarkers. *Proc. Natl. Acad. Sci. U. S. A.*

2013;110(6):2105–2110.

29. Yoshida Y, et al. Quantitative analysis of total serum glycome in human and mouse.

Proteomics 2016;16(21):2747–2758.

30. Furukawa J-I, et al. Impact of the Niemann-Pick c1 Gene Mutation on the Total Cellular

Glycomics of CHO Cells. *J. Proteome Res.* 2017;16(8):2802–2810.

31. Gao C, et al. Sensitivity of heterozygous α 1,6-fucosyltransferase knock-out mice to

cigarette smoke-induced emphysema: implication of aberrant transforming growth

factor-β signaling and matrix metalloproteinase gene expression. *J. Biol. Chem.*

2012;287(20):16699–16708.

32. Wang X, et al. Dysregulation of TGF-beta1 receptor activation leads to abnormal lung

development and emphysema-like phenotype in core fucose-deficient mice. *Proc. Natl.*

Acad. Sci. U. S. A. 2005;102(44):15791–15796.

33. Gonzalez DS, Jordan IK. The alpha-mannosidases: phylogeny and adaptive

diversification. *Mol. Biol. Evol.* 2000;17(2):292–300.

34. Ross KA, et al. Comparison of Three Methods to Quantify Repair Cartilage Collagen

Orientation. *Cartilage* 2013;4(2):111–120.

35. Glasson SS, et al. Deletion of active ADAMTS5 prevents cartilage degradation in a

murine model of osteoarthritis. *Nature* 2005;434(7033):644–648.

36. Stanton H, et al. Investigating ADAMTS-mediated aggrecanolysis in mouse cartilage.

Nat. Protoc. 2011;6(3):388–404.

37. Estes BT, et al. Isolation of adipose-derived stem cells and their induction to a

chondrogenic phenotype. *Nat. Protoc.* 2010;5(7):1294–1311.

38. Patwari P, et al. Ultrastructural quantification of cell death after injurious compression

of bovine calf articular cartilage. *Osteoarthr. Cartil. OARS Osteoarthr. Res. Soc.*

2004;12(3):245–252.

39. Rao X, et al. An improvement of the 2ˆ(-delta delta CT) method for quantitative real-

time polymerase chain reaction data analysis. *Biostat. Bioinforma. Biomath.*

2013;3(3):71–85.

40. Ovchinnikov DA, et al. Col2a1-directed expression of Cre recombinase in

differentiating chondrocytes in transgenic mice. *Genes. N. Y. N 2000* 2000;26(2):145–

146.

41. Yamada T, et al. Carminerin contributes to chondrocyte calcification during

endochondral ossification. *Nat. Med.* 2006;12(6):665–670.

42. Nakamichi Y, et al. Chondromodulin I is a bone remodeling factor. *Mol. Cell. Biol.*

2003;23(2):636–644.

43. Lufkin T, et al. Homeotic transformation of the occipital bones of the skull by ectopic

expression of a homeobox gene. *Nature* 1992;359(6398):835–841.

44. Matsui Y, et al. Accelerated development of aging-associated and instability-induced

osteoarthritis in osteopontin-deficient mice. *Arthritis Rheum.* 2009;60(8):2362–2371.

45. Stoop R, et al. Type II collagen degradation in spontaneous osteoarthritis in C57Bl/6 and

BALB/c mice. *Arthritis Rheum.* 1999;42(11):2381−2389.

46.   Fujitani N, et al. Qualitative and quantitative cellular glycomics of glycosphingolipids

560   based on rhodococcal endoglycosylceramidase-assisted glycan cleavage, glycoblotting-

561   assisted sample preparation, and matrix-assisted laser desorption ionization tandem

562   time-of-flight mass spectrometry analysis. *J. Biol. Chem.* 2011;286(48):41669−41679.

563   47.   Takegawa Y, et al. Simultaneous analysis of heparan sulfate, chondroitin/dermatan

564   sulfates, and hyaluronan disaccharides by glycoblotting-assisted sample preparation

565   followed by single-step zwitter-ionic-hydrophilic interaction chromatography. *Anal.*

566   *Chem.* 2011;83(24):9443−9449.

567   48.   Furukawa J, et al. Quantitative O-Glycomics by Microwave-Assisted β-Elimination in

568   the Presence of Pyrazolone Analogues. *Anal. Chem.* 2015;87(15):7524−7528.

569   49.   Furukawa J, et al. Comprehensive approach to structural and functional glycomics based

570   on chemoselective glycoblotting and sequential tag conversion. *Anal. Chem.*

571   2008;80(4):1094−1101.

FIGURES

Fig. 1. Effects of α-mannosidase injection in the knee joint in vivo.

(A) Time course of intra-articular α-mannosidase injection and unloading. Red arrows

indicate the one-shot, intra-articular injection that was performed once a week. **(B)**

Histological evaluation of knee articular cartilage 4 weeks after intermittent α-mannosidase

infusion and 16 weeks after load removal. Scale bars, 50 μm. **(C)** Representative macroscopic

assessment of the articular cartilage using mannosidase concentration. Scale bar, 1 cm. **(D)**

Evaluation of the collagen orientation of reparative tissues at 4 weeks postoperatively.

Sections stained with HE was viewed under a polarized light microscope at multiple angles

(0°, 45°, and 90°). Scale bar, 200 μm. **(E)** OARSI scores at 4 weeks postoperatively on

sections from the knee joint (n = 6). **(F)** Tissue degeneration scores at each time point are

based on mannosidase concentration. Data are shown as mean ± standard deviation. *$P <$

0.05, **$P < 0.01$ versus the saline group in (E), and versus 4 weeks in (F). In (E), the Welch

t-test was used for statistical analysis. In (F), n = 6 rabbits per group. One-way analysis of

variance with Tukey's multiple comparison test was used to perform statistical analysis. HE,

hematoxylin and eosin; OARSI, Osteoarthritis Research Society International; TUNEL,

terminal deoxynucleotidyl transferase dUTP nick end labeling.

Fig. 2. Cartilage degradation induced by mannosidase stimulation.

(A) Schematic diagram of the procedures used to establish the early OA model before

irreversible and progressive destruction of articular cartilage. **(B)** Mannosidase-treated

cartilage stained with Safranin O and Con A. The distribution of high-mannose type N-

glycans is decreased in mannosidase-induced degraded mouse cartilage. Scale bars, 50 μm.

(C), (D) PG release after exposure to mannosidase (C) and recovery after its removal from

cultured cartilage grafts (D). **(E)** To investigate the non-chondrocyte-mediated release of

fragments, PG loss was measured in cartilage explants that had undergone freeze-thaw

cycles. **(F)** Sequential NO released into media measured as concentrations of nitrite for

cartilage explants stimulated with mannosidase. Data are shown as mean ± standard deviation

(s.d.). *$P < 0.05$, **$P < 0.01$ versus the control group in **c, e,** and (F), and versus day 3 in (D).

In (C) and (E), n = 3 samples (six mice) per group. One-way ANOVA with the Tukey

multiple comparison test was used to perform statistical analysis. In (D), n = 12 mice in each

group and n = 3 samples (six mice) at each time point. In (F), n = 16 mice at each time point

and n = 4 samples (eight mice) per group. In (D) and (F), two-way ANOVA with the Sidak

multiple-comparisons test was used to perform statistical analysis. Con A, concanavalin A;

NO, nitric oxide; ANOVA, analysis of variance; PG, proteoglycan.

Fig. 3. Cartilage N-glycomes.

(A) MALDI-TOF MS spectra showing the quantitative N-glycan profiles of cartilage before

(upper) and after (lower) mannosidase treatment. **(B)** Scatter plot of the changes in *m/z* per

protein abundance in the mannosidase (x-axis) and control (y-axis) N-glycan structures. The

red dots indicate the top 10 N-glycan structures based on the deviation from the y = x line.

(C) Heatmap of N-glycome showing differential N-glycan expression in the mannosidase

(man+) group versus the control (man-) group. **(D)** N-glycans altered by mannosidase were

corefucosylated glycans and high-mannose type N-glycans. (**E**) RT-PCR expression analysis

of marker genes in organ culture of cartilage with mannosidase. Data are shown as mean ±

standard deviation. *$P < 0.05$, **$P < 0.01$ versus the control group in (D), and versus 0 hours

in (E). In (D) and (E), n = 3 samples (six mice) per group. In (D), unpaired t-tests were used

to perform statistical analyses. In (E), one-way ANOVA with the Dunnett multiple

comparison test was used to perform statistical analysis. MALDI-TOF MS, matrix-assisted

laser desorption/ionization-time of flight mass spectrometry; ANOVA, analysis of variance;

RT-PCR, real-time polymerase chain reaction.

Fig. 4. Loss of resilience due to FUT8 deficiency in cartilage.

(A) Biological reaction of FUT8. FUT8 transfers fucose to the innermost GlcNAc residue of

complex N-glycans via α1,6-linkage (corefucosylation). **(B)** Targeted disruption of *Fut8*

locus. The *Fut8* gene (WT allele; top), targeting vector (middle), and disrupted *Fut8* locus

(mutant allele; bottom). Schematic representation of the *Fut8*-targeting strategy and Cre-

mediated recombination of the *Fut8*^loxP allele. **(C)** MALDI-TOF MS mass spectra of N-

glycans from WT and *Fut8* cKO mice. The corefucosylation levels in cartilage were

decreased and undetectable in *Fut8* cKO mice. (**D**) Gene profile in chondrocytes isolated

from *Fut8* cKO mice. The expression levels of these genes in WT cells were set to 1. PPIA,

peptidylprolyl isomerase A. (**E**) Histological findings in cartilage explants from *Fut8* cKO

mice and their floxed littermates, cultured with mannosidase and subjected to Safranin O

staining and PhoSL lectin staining. Scale bar, 50 μm. (**F**) PG release in cultured cartilage

explants from *Fut8* cKO mice and their floxed littermates. Data are shown as mean ±

standard deviation. In (E), n = 3 samples (six mice) per group. *$P < 0.05$, **$P < 0.01$ versus

the control group. One-way ANOVA, with the Tukey multiple comparison test, was used to

perform statistical analysis. MALDI-TOF MS, matrix-assisted laser desorption/ionization-

time of flight mass spectrometry; WT, wild-type; cKO, conditional knockout; PG,

proteoglycan; ANOVA, analysis of variance.

Fig. 5. OA acceleration in *Fut8* cKO mice.

(A) Double staining with alizarin red and alcian blue of the whole skeleton of wild-type and

Col2a1-Cre;Fut8^{flox/flox} cKO littermate embryos (newborn). Scale bars, 1 cm (left). Weight

and body length of wild-type and cKO littermate embryos (right). **(B)** Growth curves

determined by body weight in male (left) and female (right) wild-type mice and their cKO

littermates. **(C)**, **(D)** Features of instability-induced OA in *Fut8^{flox/flox}* (flox) mice and their

cKO littermates at 8 weeks after surgery. Safranin O staining is shown for each mouse

genotype. Scale bar, 100 μm (C). Summed histological scores for OA severity in the knee

cartilage from flox and cKO mice, as determined using the OARSI scoring system, are shown

(D). **(E)**, **(F)** Features of age-associated osteoarthritis in wild-type mice and their flox and

cKO littermates. Safranin O staining of the knee joint is shown for each mouse genotype at 3,

4, 9, and 15 months of age. Scale bar, 100 μm (E). The summed OARSI scores are shown (F).

Data are shown as mean ± standard deviation. In (A), the Welch t-test was used to perform

statistical analyses (n = 6 mice per group). In (B), n = 15 mice per group at each time point.

In (D) and (F), n = 5 mice per group at each time point. *$P < 0.05$, **$P < 0.01$ versus the

wild-type group in (B) and (F), and versus flox mice in (D). One-way ANOVA with the

Tukey multiple comparisons test (B), (D) and two-way ANOVA with the Tukey multiple

comparisons test (F) were used to perform statistical analysis. OA, osteoarthritis; cKO,

conditional knockout; OARSI, Osteoarthritis Research Society International. ANOVA,

analysis of variance; ns, not significant.

Fig. 6. Altered glycosylation of human Osteoarthritis (OA) cartilage based on

comprehensive glycan analysis.

(A) Total glycome profiling of human OA cartilage. Pie charts at the vertices of the pentagon

correspond to the glycan expression profiles of N-glycans, O-glycans, GSL-glycans, free

oligosaccharides (fOS), and glycosaminoglycan (GAG). The size of each circle and its

constituent colors reflect the absolute quantity of glycans (pmol/100 µg protein) and the

glycan substructures, respectively. The sizes of the circles representing the GAG contents are

increased by 10-fold. Each color indicates the estimated glycan structure and corresponds to

the respective glycan number listed in the S3 Table. **(B)** Expression of core fucose in healthy

and OA cartilage. Scale bar, 100 μm. **(C)** Principal component analysis (PCA) of a glycan

expression data set. Data points represent individual samples. The first principal component

(PC1) distinguishes healthy and OA samples. **(D)** Hierarchical cluster analysis results

showing cluster image display for total glycans with color gradient for relative glycan

expression and dendrogram for each glycan structure. Cluster summary of PCA on the

glycome are shown on the right. The most representative variable is (Hex)1 (HexNAc)1

(Fuc)1 + (Man)3(GlcNAc)2 that means the largest squared correlation with its cluster

component.