1   Full title: Eye-specific active zone clustering underlies synaptic competition in the

2   developing visual system

5   Short title: Active zone clustering during eye-specific synaptic competition

7   Author list: Chenghang Zhang[1], Tarlan Vatan[1], and Colenso M. Speer[1,*]
8

9   Author Affiliations: [1] Department of Biology, University of Maryland, College Park,

10   Maryland, USA 20742.

11   [*]Corresponding author

12   E-mail: cspeer@umd.edu

Abstract

Spatially clustered synaptic inputs enable local dendritic computations important for learning, memory, and sensory processing. In the mammalian visual system, individual retinal ganglion cell (RGC) axons form clustered terminal boutons containing multiple active zones onto relay cell dendrites in the dorsal lateral geniculate nucleus (dLGN). This mature architecture arises through the addition of release sites, which strengthens selected afferents while weaker inputs are pruned. Following eye-opening, spontaneous activity and visual experience promote synaptic refinement and bouton clustering after binocular inputs have segregated. However, anatomical changes in release site addition and spatial patterning during earlier stages of eye-specific competition are not well understood. To investigate this, we examined the spatial organization of eye-specific active zones in wild type mice and a mutant line with disrupted cholinergic retinal waves. Using volumetric super-resolution single-molecule localization microscopy and electron microscopy, we found that individual retinogeniculate boutons begin forming multiple nearby presynaptic active zones during the first postnatal week. Both eyes generate these "multi-active-zone" (mAZ) inputs throughout refinement, but the dominant-eye forms more numerous mAZ contacts, each with more active zones and larger vesicle pools. At the height of competition, the non-dominant-eye projection adds many single active zone (sAZ) synapses. Mutants with abnormal cholinergic retinal waves still form mAZ inputs, but develop fewer sAZ synapses and show reduced synapse clustering in projections from both eyes. Together, these findings reveal activity-dependent, eye-specific differences in release site addition during synaptic competition in circuits essential for visual perception and behavior.

Introduction

A key mechanism for neuronal signal processing is the formation of spatially clustered synapses that facilitate local computations within individual dendrites [1-3]. Through biophysical signal integration mechanisms, neighboring synaptic inputs play critical computational roles in learning, memory, and sensory processing underlying cognition and behavior [4-7]. During circuit development, both spontaneous and sensory-driven neural activity regulate synaptic clustering, stabilizing some synapses and eliminating others to establish mature connectivity patterns [6-8].

The refinement of retinal inputs to the dorsal lateral geniculate nucleus (dLGN) of the thalamus is a model example of activity-dependent synaptic clustering [9]. Over development, boutons within the terminal arbors of individual retinogeniculate axons cluster progressively [10-13], creating multiple synaptic contacts onto individual postsynaptic targets [14-16]. Concurrently, developing RGC boutons add presynaptic release sites such that some mature retinogeniculate terminals contain several dozen active zones (AZs) [16, 17]. The formation of clustered boutons with multiple release sites subserves a critical "driver" function of retinogeniculate input [18, 19].

Retinogeniculate bouton clustering depends on visual experience, and dark-rearing after eye-opening reduces clustering within individual axon arbors [12]. However, less is known about synaptic spatial relationships and activity-dependent release site

refinement during eye-specific competition before eye-opening. During early axon

ingrowth, individual RGCs extend sparse side branches in the incorrect eye-specific

territory [20-22]. These transient branches form synapses that contain fewer presynaptic

vesicles than those made by the same axon in the correct eye-specific domain [23].

Functional recordings after eye-opening show that while many RGC inputs are pruned,

the remaining inputs strengthen by adding release sites [24]. This leads to glutamate

spillover and cross talk between RGC inputs that increases postsynaptic relay neuron

excitability [25]. Thus, early eye-specific differences in release site number or spacing

may contribute to binocular input refinement prior to eye-opening.

In a previous study from our laboratory, we used volumetric stochastic optical

reconstruction microscopy (STORM), anterograde tract tracing, and

immunohistochemical labeling of synaptic proteins to show that dominant-eye synapses

contain more total vesicles and have more vesicles near the AZ (putative docking)

compared with non-dominant-eye synapses [26]. However, we did not examine whether

eye-specific inputs differ in AZ number or AZ spatial proximity (clustering) during

synaptic competition. To address this gap, we reanalyzed our published dataset,

focusing on eye-specific AZ spatial relationships. We found that projections from each

eye form a subset of retinogeniculate inputs in which several (2-4) AZs share a common

cluster of presynaptic vesicles. We refer to these synapses as multi-active-zone (mAZ)

inputs. All other retinogeniculate inputs contained one single active zone and its

associated vesicle cluster, and we term these single-active-zone (sAZ) synapses.

During eye-specific synaptic competition, the dominant-eye projection formed more

mAZ inputs, each with more AZs and a larger presynaptic vesicle pool compared to the

non-dominant eye projection. Similarly, the dominant-eye had higher vesicle signal at

sAZ inputs. At the peak of synaptic competition midway through the first postnatal week,

the non-dominant-eye formed numerous sAZ inputs, equalizing the global synapse

density between the two eyes. These eye-specific AZ patterns were disrupted in a

mutant mouse line with abnormal stage II cholinergic retinal waves and retinogeniculate

segregation defects. Our results demonstrate that eye-specific presynaptic release site

clustering arises before eye-opening and is impacted by defects in spontaneous retinal

activity.

Results

Retinogeniculate inputs form multiple active zones during eye-

specific competition

To investigate active zone refinement during eye-specific segregation, we reanalyzed a

volumetric super-resolution imaging dataset previously published by our laboratory [26].

We used volumetric STochastic Optical Reconstruction Microscopy (STORM) [27] to

image the dLGNs of wild-type (WT) mice at three postnatal ages (P2, P4, and P8) (Fig.

1A). We labeled eye-specific inputs by monocular injection of Alexa Fluor-conjugated

cholera toxin subunit B tracer (CTB) together with immunostaining for presynaptic

Bassoon, postsynaptic Homer1, and presynaptic vesicular glutamate transporter 2

(VGluT2) proteins (Fig. 1B). We collected separate image volumes (~45K μm^3 each)

from three biological replicates at each age. To assess the impact of spontaneous

retinal activity on synaptic development across the same time period, we performed

identical experiments in β2-knockout (β2KO) mice lacking the beta 2 subunit of the

nicotinic acetylcholine receptor, a mutation that disrupts spontaneous cholinergic retinal

wave activity, eye-specific segregation, and retinogeniculate synapse development [22, 26,

28-36] (Fig. 1A). Because eye-specific segregation is incomplete until ~P8, we limited our

re-analysis to the future contralateral eye-specific region of the dLGN, which is reliably

identified across all stages of postnatal development (Fig. 1A, see also Materials and

Methods).

By analyzing synapses in the contralateral dLGN from eighteen mice across three ages

and two genotypes (Table S1), STORM revealed two classes of retinogeniculate inputs

distinguished by active zone (AZ) number (Fig. 1B). We defined each retinogeniculate

input as a single contiguous VGluT2 cluster together with all its associated presynaptic

(Bassoon) and postsynaptic (Homer1) paired synaptic labels. Using this definition,

inputs that had multiple (2-4) Bassoon AZs were classified as multi-active-zone (mAZ)

inputs, while those with a single Bassoon AZ were designated single-active-zone (sAZ)

inputs (Fig. 1B). Most mAZ inputs contained two AZs (~70-90%, varying with age,

genotype, and eye-of-origin); smaller proportions contained three AZs (~10-20%) or four

or more AZs (<5%) (Table S1).

Each mAZ input could be a single terminal bouton with several AZs, or a cluster of sAZ

synapses within separate boutons [10, 11, 15, 16, 37]. To address this, we used electron

microscopy (EM) to image retinogeniculate terminals in the dLGN at P8. We generated

a transgenic line expressing mitochondrial matrix-targeted dimeric dAPEX2 reporter [38]

in ipsilaterally-projecting RGCs [39-41], providing unambiguous mitochondrial labeling in

ipsiRGC axons. EM images confirmed the presence of individual retinogeniculate

boutons with multiple active zones, consistent with our STORM data (Fig. 1C). Previous

EM reconstructions of retinogeniculate inputs reported no evidence of RGC bouton

convergence at the end of the first postnatal week [11]. Together, these results suggest

that mAZ inputs in STORM images are single RGC terminals that house several closely

spaced release sites. Hereafter, we use the word "synapse" only to refer to each

partnered active zone (Bassoon/Homer1 pairs); mAZ inputs contain several synapses,

while each sAZ input is one synapse.

138 Changes in eye-specific input density during synaptic

139 competition

In our previous analysis, we reported global eye-specific synapse densities that

reflected the combined mAZ and sAZ inputs. Dominant-eye synapse density was

greater than that of the non-dominant eye at P2 and P8, but eye-specific synapse

densities were equivalent at P4 during the peak of synaptic competition [26]. To determine

how these two input classes evolve during competition, we quantified the densities and

percentages of mAZ and sAZ inputs in WT and β2KO mice (Fig. 2A). Eye-of-origin for

each retinogeniculate input was assigned by colocalizing CTB signal with VGluT2 (Fig.

2B). Binocular CTB control injections showed that anterograde tracing labeled >97% of

retinogeniculate synapses at P4 and P8, ensuring accurate eye-specific assignment [26].

Within the contralateral eye-specific region of the dLGN, CTB(+) VGluT2 clusters were

classified as "dominant-eye" inputs and CTB(-) VGluT2 clusters as "non-dominant eye"

inputs (Fig. 2B).

After assigning eye-of-origin to each retinogeniculate synapse, we found that the

density of CTB(+) mAZ inputs was significantly higher than CTB(-) mAZ inputs at every

age in WT mice (Fig. 2C, top left; Fig. S1, top). β2KO mice also developed eye-specific

differences in mAZ input density (Fig. 2C, bottom left; Fig. S1, bottom). For WT sAZ

synapses, the dominant-eye had a higher synapse density at P2 and P8. At P4,

however, sAZ synapse density was equivalent between the eyes (Fig. 2C, top right).

This pattern resulted from non-dominant eye sAZ synapse addition, which equalized the

global input density between the eyes at P4 as we reported previously (5/95%

confidence interval, -0.014-0.011 synapses / μm^3). In β2KO mice, the non-dominant eye

formed fewer sAZ synapses, trending toward an eye-specific difference. (Fig. 2C,

bottom right; $p < 0.05$, power=0.62; see Tables S2/S3). Thus, β2KO mice with disrupted

retinal activity maintain a higher mAZ input fraction in the dominant-eye projection, but

sAZ synapse addition is reduced at the peak of competition.

Multi‐ and single‐active‐zone inputs from the dominant eye

show increased vesicle pool size and vesicle proximity to the

active zone

We previously reported a dominant-eye bias in VGluT2 volume when considering all

retinogeniculate inputs [26]. Here, we assessed presynaptic maturation separately in sAZ

and mAZ inputs by measuring their total VGluT2 volume. In WT mice, both mAZ (Fig.

3A, left) and sAZ (Fig. 3B, left) inputs showed significant eye-specific volume

differences at each age. In the middle of eye-specific competition at P4 in WT mice, the

median VGluT2 cluster volume in dominant-eye mAZ inputs was 3.72 fold larger than

that of non-dominant-eye inputs (Fig. 3A, left). In contrast, β2KO mice showed a smaller

1.1 fold difference at the same age (Fig. 3A, right panel). For sAZ synapses at P4, the

magnitudes of eye-specific differences in VGluT2 volume were smaller: 1.35-fold in WT

(Fig. 3B, left) and 0.41-fold in β2KO mice (Fig. 3B, right). Thus, both mAZ and sAZ input

size favors the dominant eye, with larger eye-specific differences seen in WT mice (see

Table S3).

In addition to total vesicle pool volume, we quantified VGluT2 volume within a 70 nm

shell around each AZ in mAZ and sAZ inputs, considering this a proxy for docked

vesicles (Fig. S2A-C) [26]. In WT mice at P4, dominant-eye inputs showed greater per-AZ

vesicle volume than non-dominant inputs in both mAZ and sAZ categories (Fig. S2B,

left). These eye-specific differences were absent in β2KO mice (Fig. S2B, right).

Moreover, at all ages and in both genotypes, vesicle volume within the 70 nm shell did

not differ between mAZ and sAZ inputs (Fig. S2A-C). This shows that vesicle docking at

release sites favors the dominant-eye as we previously reported [26], but is similar for like-

eye type inputs regardless of AZ number.

Vesicle pool size scales with active zone number

Because mAZ inputs showed greater total VGluT2 volume than sAZ synapses, yet

exhibited comparable vesicle docking, the disparity could reflect a scaling effect of

vesicle pool size with increased AZ number. To evaluate how presynaptic vesicle pool

volume scales with AZ number, we compared the Bassoon cluster number to VGluT2

volume for every retinogeniculate input. In both WT (Fig. S2D) and β2KO mice (Fig.

S2E), mAZ inputs contained an average of 2-3 Bassoon clusters (separate AZs) in the

first postnatal week. In WT mice, CTB(+) dominant-eye mAZ inputs contained more AZs

than CTB(-) non-dominant-eye mAZ inputs at P4 and P8 (Fig. S2D). This maturation

was delayed until P8 in β2KO mice (Fig. S2E). For CTB(+) dominant-eye inputs in both

genotypes at P4, vesicle pool volume correlated positively with AZ number (Fig. 3C;

Pearson correlation coefficients: WT [0.99] and β2KO [0.95]). A similar, but weaker

correlation was observed for CTB(-) non-dominant-eye inputs (Pearson correlation

coefficients: WT [0.97] and β2KO [0.90]. Dividing the total presynaptic VGluT2 volume

by the AZ number revealed a consistent vesicle volume per AZ for both sAZ and mAZ

inputs (Fig. 3D/S3, Table S3). Collectively, these results indicate that presynaptic

vesicle pool volume scales with AZ number for each eye-specific input, while a

dominant-eye bias persists throughout development.

212   Eye-specific synapse clustering before eye-opening

Eye-specific competition is thought to involve stabilization of co-active, neighboring

inputs from the same eye and elimination of out of sync inputs from the opposite eye [42-

44]. Because axon refinement depends on the relative strength of neurotransmission

between competing inputs [39, 45-51], mAZ inputs with multiple release sites could help

stabilize nearby like-eye inputs [48, 52-54].

To quantify synaptic clustering patterns, we measured the distance from every eye-

specific sAZ synapse to all other sAZ and mAZ inputs within each image field. sAZ

synapses were often found nearby other inputs from the same eye (Fig. 4A). To define

clustering, we searched volumetrically around each mAZ input and measured the

fraction of sAZ synapses that were nearby at increasing distances (Fig. S4A/B). We

then compared the observed percentages with datasets in which sAZ synapse positions

were randomly shuffled within each neuropil volume. The largest differences occurred

within a 1-2 μm search distance (Fig. S4A/B), and so we chose a 1.5 μm cutoff from the

edge of each mAZ input to designate it "clustered" if at least one neighboring sAZ

synapse lay within this distance and "isolated" if none did (Fig. 4B). No significant

clustering was detected when sAZ and mAZ inputs originated from opposite eyes (Fig.

S4C/D).

At the peak of competition (P4) in WT mice, more than 65% of both mAZ and sAZ

inputs were clustered for both eyes, while this proportion fell to ~50% in β2KO mice (Fig.

4B). In the WT dominant-eye projection, the fractions of clustered mAZ and sAZ inputs

were similar at each age (Fig. 4C, top left panel). For the non-dominant eye projection,

however, clustered mAZ inputs outnumbered clustered sAZ inputs at P4 (Fig. 4C,

bottom left panel), the age when this eye adds sAZ synapses (Fig. 2C). β2KO mice

showed no difference between mAZ and sAZ clustering at any age (Fig. 4C, right

panels). Additionally, in WT mice at P4 and P8, clustered mAZ inputs from both eyes

had more neighboring sAZ synapses than did clustered sAZ synapses (Fig. 4D, left);

this enrichment was absent in β2KO mice (Fig. 4D, right). Thus, while most

retinogeniculate synapses lie within 1.5 μm of another like-eye input, WT mice show a

relative increase in neighboring synapses near mAZ inputs during synaptic competition.

Clustered mAZ inputs in P4 WT mice were also closer together than isolated mAZ

inputs. Clustered mAZ inputs formed by the dominant-eye were ~32% closer to the

nearest like-eye clustered mAZ input compared to isolated mAZ inputs (Fig. 5A, left

panel). In the non-dominant eye projection, clustered mAZ inputs were ~55% closer

together compared to isolated mAZ inputs (Fig. 5B, left panel). Once segregation was

complete at P8, distances between clustered and isolated mAZ inputs were more

similar (Fig. S5). In β2KO mice, distances between isolated and clustered mAZ inputs

and their nearest clustered mAZ neighbor did not differ for either eye at P4 or P8 (Fig.

5/S5, right panels). These patterns are consistent with sAZ synapse addition when mAZ

inputs are in close proximity during competition.

Finally, we tested whether sAZ vesicle pool volume varies with distance to mAZ inputs.

We classified each sAZ as "near" (≤1.5 μm) or "far" (>1.5 μm) relative to the nearest

like-eye mAZ input. Across ages and genotypes, vesicle pool volume was similar for

near and far sAZ synapses in both eyes (Fig. S6). Thus, vesicle pool size in sAZ

synapses appears independent of their proximity to mAZ inputs.

Discussion

Spontaneous retinal activity guides eye-specific refinement by strengthening dominant eye inputs in the correct territory and pruning weaker inputs from the competing eye. Using volumetric super-resolution microscopy and eye-specific synaptic immunolabeling, we previously identified activity-dependent, eye-specific differences in presynaptic vesicle pool size and vesicle association with the active zone (AZ), both favoring the dominant eye during synaptic competition [26]. Here, we reanalyzed this dataset to measure synaptic spatial arrangements during normal development and how these are affected in a mutant model with abnormal eye-specific segregation.

Early input maturation and addition of release sites

Before eye-opening, retinal ganglion cell (RGC) axons from both eyes formed terminals with either multiple active zones (mAZ inputs) or a single active zone (sAZ synapses). Electron microscopy of dAPEX2-labeled ipsilateral RGC inputs revealed individual boutons with multiple active zones. Prior electron microscopy studies in mice showed no clustering from neighboring RGC boutons at postnatal day 8 [10, 11], suggesting the mAZ inputs seen in STORM images are single boutons containing multiple release sites. During competition, the dominant eye projection generated more of these mAZ inputs, each with more active zones and larger vesicle pools than the non-dominant eye. We observed a linear relationship between presynaptic vesicle signal volume and active zone number, suggesting that release site addition and vesicle pool expansion may be linked during input maturation. Functional recordings indicate that release site addition is the primary driver of retinogeniculate input strengthening after eye-opening [24]. Our

findings reveal that eye-specific differences in release site addition at individual

terminals emerge at the earliest stages of binocular refinement.

Spatial interactions during competition

Ipsilateral RGC axons are delayed in entering the dLGN until after contralateral inputs

have already innervated the nucleus [55]. Despite their late arrival, these axons competed

by forming numerous sAZ inputs, which equalized the overall input density between the

eyes in the future contralateral eye domain at postnatal day 4 (P4). Most of these

synapses formed nearby other like-eye neighbors, and were enriched near mAZ inputs,

supporting the idea of release site addition at strong synapses. Neighboring sAZ

synapses could ultimately mature into mAZ inputs as release sites are added and

vesicle pools expand.

Distance analysis showed that clustered mAZ boutons were closer together than

isolated mAZ boutons, further supporting release site addition near strong neighboring

inputs. It remains unknown, however, whether nearby mAZ inputs originate from the

same RGC. Nearby mAZ inputs within individual RGC arbors could promote nearby

release site addition via intrinsic mechanisms, such as increased presynaptic calcium

entry or surface delivery of synaptogenic molecules. Boutons with multiple release sites

could also trigger non-cell-autonomous signaling mechanisms that stabilize neighboring

inputs from the same eye [48, 53, 54]. Release site addition increases extracellular

glutamate concentration and promotes spillover onto adjacent synapses, which

enhances the excitability of developing relay cells [25] and could contribute to long term

synaptic plasticity evoked by high frequency spiking in retinal wave bursts [56-59].

Consequently, release site addition and local bouton clustering may act in tandem to

stabilize coactive inputs.

Competitive refinement also involves synapse elimination and axonal retraction through

punishment signals. Genetic deletions of VGluT2 or RIM1 proteins in ipsilaterally-

projecting RGCs decreased presynaptic vesicle release and prevented retraction of

contralateral RGC axons from the ipsilateral territory [39, 45]. One downstream mediator of

synaptic punishment is JAK2 kinase, which is phosphorylated in less active synapses [52].

Similar to neurotransmission mutant phenotypes, disruption of JAK2 signaling prevents

axon retraction (punishment) during competition [52]. The spatial analyses we developed

here will enable future mapping of input-specific punishment signals during synaptic

competition. This includes phospho-JAK2 as well as molecular tags for glial pruning of

weak inputs during eye-specific segregation [60-62].

Requirement for spontaneous retinal activity

β2KO mice showed significant defects in synapse addition during the first postnatal

week. mAZs still formed with similar overall ratios as in wild type controls (Fig. S1, Table

S3) and eye-specific differences in vesicle pool size still emerged. However, β2KO mice

failed to form many sAZ synapses at the height of competition, particularly in the late-

arriving non-dominant (ipsilateral) eye projection. The failure to add synapses reduced

synaptic clustering and more inputs formed in isolation in the mutants compared to

controls.

While our findings support a role for spontaneous retinal activity in presynaptic release

site addition and clustering, activity-dependent postsynaptic mechanisms also influence

input refinement at later stages. Retinogeniculate synapses undergo postsynaptic

strengthening and weakening through potentiation and depression mediated by

AMPARs and NMDARs [56-59]. After eye-specific segregation, spontaneous retinal activity

is required for postsynaptic AMPAR insertion, synaptic strengthening, and elimination of

weaker inputs [63]. Continued maintenance of segregation depends on calcium influx into

relay neurons via L-type calcium channels, further implicating postsynaptic signaling in

late-stage refinement [64, 65]. Input targeting may also be guided by molecular cues to

form non-random, eye-specific connections with postsynaptic targets. Ipsilaterally-

projecting RGCs have distinct gene expression profiles that specify axon guidance and

may further support eye-specific synaptic targeting [66, 67]. Spontaneous retinal activity

may permit axons to read out molecular regulators of synaptogenesis, as previously

shown for RGC axon retraction [68].

Release site addition as a general mechanism underlying synaptic competition

Early synaptic clustering during retinogeniculate development resembles other circuits

where neural activity guides competitive synaptic and axonal remodeling. At the

neuromuscular junction (NMJ), motor neuron terminals compete for control of a

postsynaptic muscle fiber; a single motor axon input strengthens while competing axons

are eliminated [69-71]. Competition at the NMJ depends on inter-synaptic distance, with

motor axons losing connections located near stronger competing synapses [69, 70]. As

winning terminals are enlarged, presynaptic release site addition maintains a consistent

density of Bassoon clusters (~2-3/μm^2) [72]. Neighboring inputs with weaker synaptic

transmission are selectively eliminated [73, 74]. Competing motor axons differ in their

release probabilities early in development [75], suggesting that presynaptic efficacy

triggers local "stabilization" signals in winners and "punishment" signals in losers [76].

Presynaptic agrin release stabilizes postsynaptic receptor clusters by counteracting the

destabilizing, dispersal effects of acetylcholine release, emphasizing the importance of

early presynaptic transmission in postsynaptic stabilization [77, 78].

Similarly, in the developing cerebellum, Purkinje cells initially receive inputs from

multiple climbing fibers (CFs) during the first postnatal week. Subsequently, one winning

CF emerges and consolidates its synaptic inputs, while losing CFs are pruned [79, 80].

Here again, the winning input strengthens by adding presynaptic release sites, which

increases multivesicular release and elevates glutamate concentration in the synaptic

cleft [81-83]. Postsynaptic ultrastructural and molecular changes occur several days later

as dominant CF inputs potentiate and losing CF inputs depress [82, 84, 85] through spike-

timing dependent remodeling [86, 87]. Across these models, the earliest distinguishing

feature between competing inputs is relative presynaptic transmission strength. Thus,

release-site addition may be a conserved mechanism that biases synaptic refinement

outcomes across developing neural circuits.

Materials and methods:

The raw imaging data in this paper were previously reported [26]. Materials and methods

below are adapted from this work. All Matlab and Python code used in the work is

available on GitHub (https://github.com/SpeerLab). Raw STORM images of the full data

are available on the open-access Brain Imaging Library [88]. These images can be

accessed here: https://doi.brainimagelibrary.org/doi/10.35077/g.1164.

Animals

Wild-type C57BL/6J mice (Stock Number 000664) and APEX2 mice (Stock Number

032764) used in this study were purchased from the Jackson Laboratory. ET33^{Cre} mice

and β2KO were generously gifted by Drs. Eric M. Ullian (University of California, San

Francisco) and Michael C. Crair (Yale School of Medicine), respectively. All

experimental procedures were performed in accordance with an animal study protocol

approved by the Institutional Animal Care and Use Committee (IACUC) at the University

of Maryland. Neonatal male and female mice were used interchangeably for all

experiments. Tissue from biological replicates (N=3 animals) was collected for each age

(P2/P4/P8) from each genotype (WT and β2KO) (18 animals total). Primers used for

genotyping β2KO mice are: forward: CAGGCGTTATCCACAAAGACAGA; reverse:

TTGAGGGGAGCAGAACAGAATC; mutant reverse: ACTTGGGTTTGGGCGTGTTGAG.

Primers used for genotyping ET33Cre mice are: Forward:

GGTCCTTGGCAGATGGGCAT; reverse: CGGCAAACGGACAGAAGCATT.

Electron microscopy tissue preparation

Animals were deeply anesthetized with ketamine/xylazine and transcardially perfused

with 5-10 mls of 37°C 0.9% sterile saline (pH 7.2) followed by 20-30 mls of 37°C 4%

EM-Grade paraformaldehyde (PFA, Electron Microscopy Sciences), 2% EM-Grade

glutaraldehyde (GA, Electron Microscopy Sciences), 4mM calcium chloride ($CaCl_2$,

Sigma-Aldrich) in 0.2M cacodylate buffer (pH 7.4). Brains were postfixed in the same

perfusion fixative solution overnight at 4°C. Brains were vibratome sectioned in 0.2M

cacodylate buffer (pH 7.4) at 100 μm. A circular tissue punch (~500 μm diameter)

containing the dLGN was microdissected from each section using a blunt-end needle.

dLGN sections were washed in 0.2M cacodylate buffer (pH 7.4) at 4°C on a rotator (4x

20 minutes each). Sections were incubated in 20mM glycine (Sigma-Aldrich) in 0.2M

cacodylate buffer (pH 7.4) at 4°C on a rotator for 30 minutes, followed by 5 x 20 minutes

washes in 0.2M cacodylate buffer (pH 7.4) at 4°C on a rotator. dLGN sections were

immersed in 0.5mg/mL DAB solution, covered with foil, for 30 minutes at 4°C on a

rotator. 10 μL of 0.03% hydrogen peroxide (Sigma-Aldrich) was mixed into the DAB

solution (1 mL) and incubated for 10 minutes at 4°C on a rotator (light protected). The

reaction was quenched with 0.2M cacodylate buffer (pH 7.4) washes (3 x 1-minute

washes followed by 2 x 20 minutes washes).

416   Tissue preparation for scanning electron microscopy

dLGN sections were fixed in 2% osmium tetroxide (Electron Microscopy Sciences) in

0.15M cacodylate buffer (pH 7.4) for 45 minutes at room temperature. Sections were

reduced with 2.5% potassium ferricyanide (Electron Microscopy Sciences) in 0.15M

cacodylate buffer (pH 7.4) for 45 minutes at room temperature in the dark, followed by 2

x 10-minute washes in double distilled water. Sections were incubated in 1% aqueous

thiocarbohydrazide (Electron Microscopy Sciences) for 20 minutes, followed by 2 x 10-

minute washes in double distilled water. Samples were fixed in 1% aqueous osmium

tetroxide for 45 minutes at room temperature, followed by 2 x 10-minute washes in

double distilled water. Sections were postfixed in 1% uranyl acetate (Electron

Microscopy Sciences) in 25% ethanol in the dark for 20 minutes at room temperature

and washed and stored in double distilled water overnight. The next day, samples were

stained with aqueous lead aspartate at 60°C for 30 minutes and washed with double

distilled water 2 x 10 minutes. Tissues were dehydrated in a graded dilution series of

100% ethanol (35%; 50%; 70%; 95%; 100%; 100%; 100% EtOH) for 10 minutes each at

room temperature. Samples were immersed in propylene oxide (Electron Microscopy

Sciences) 3 x 10 minutes at room temperature, followed by a series of epon

resin/propylene oxide (812 Epon Resin, Electron Microscopy Sciences) exchanges with

increasing resin concentrations (50% resin/50% propylene oxide; 65% resin/35%

propylene oxide; 75% resin/25% propylene oxide; 100% resin; 100% resin) for 90

minutes each. Tissues were transferred to BEEM capsules (Electron Microscopy

Sciences) that were filled with 100% resin and polymerized for at least 48 hours at 60°C.

Transmission electron microscopy image acquisition

Plasticized sections were cut at 70 nm with a Histo Jumbo diamond knife (DiATOME)

using a Leica UC7 ultramicrotome. Sections were decompressed using chloroform

vapor and collected onto 3.05mm 200 mesh Gilder thin bar hexagonal mesh copper

grids (T200H-Cu, Electron Microscopy Sciences). Sections were imaged unstained on a

Hitachi HT7700 transmission electron microscope (HT7700, Hitachi High-Tech America,

Inc.) at 80kV.

Eye injections

Intraocular eye injections were performed one day before tissue collection. Briefly, mice

were anesthetized by inhalant isoflurane and sterile surgical spring scissors were used

to gently part the eyelid to expose the corneoscleral junction. A small hole was made in

the eye using a sterile 34-gauge needle and ~0.5 μL of cholera toxin subunit B

conjugated with Alexa Fluor 488 (CTB-488, ThermoFisher Scientific, Catalogue Number:

C34775) diluted in 0.9% sterile saline was intravitreally pressure-injected into the right

eye using a pulled-glass micropipette coupled to a Picospritzer (Parker Hannifin).

dLGN tissue preparation for STORM imaging

Animals were deeply anesthetized with ketamine/xylazine and transcardially perfused

with 5-10 mls of 37°C 0.9% sterile saline followed by 10 mls of room temperature 4%

EM-Grade paraformaldehyde (PFA, Electron Microscopy Sciences) in 0.9% saline.

Brains were embedded in 2.5% agarose and vibratome sectioned in the coronal plane

at 100 μm. From the full anterior-posterior series of dLGN sections (~6-8 sections) we

selected the central two sections for staining in all biological replicates. These sections

were morphologically consistent with Figs. 134-136 (5.07-5.31 mm) of the postnatal day

6 mouse brain from Paxinos's "Atlas of the developing mouse brain" Academic Press,

2020 [89]. Selected sections were postfixed in 4% PFA for 30 minutes at room

temperature and washed for 30-40 minutes in 1X PBS. The dLGN was identified by the

presence of CTB-488 signals using a fluorescence dissecting microscope. A circular

tissue punch (~500 µm diameter) containing the dLGN was microdissected from each

section using a blunt-end needle. A small microknife cut was made at the dorsal edge of

the dLGN which, together with the CTB-488 signal, enabled us to identify the dLGN

orientation during image acquisition.

Immunohistochemistry

We used a serial-section single-molecule localization imaging approach to prepare

samples and collect super-resolution fluorescence imaging volumes as previously

described [26]. dLGN tissue punches were blocked in 10% normal donkey serum

(Jackson ImmunoResearch, Catalogue Number: 017-000-121) with 0.3% Triton X-100

(Sigma-Aldrich Inc.) and 0.02% sodium azide (Sigma-Aldrich Inc.) diluted in 1X PBS for

2-3 hours at room temperature and then incubated in primary antibodies for ~72 hours

at 4°C. Primary antibodies used were Rabbit anti-Homer1 (Synaptic Systems,

Catalogue Number: 160003, 1:100) to label postsynaptic densities (PSDs), mouse anti-

Bassoon (Abcam, Catalogue Number AB82958, 1:100) to label presynaptic active

zones (AZs), and guinea pig anti-VGluT2 (Millipore, Catalogue Number AB251-I, 1:100)

to label presynaptic vesicles. Following primary antibody incubation, tissues were

washed in 1X PBS for 6 x 20 minutes at room temperature and incubated in secondary

antibody solution overnight for ~36 hours at 4°C. The secondary antibodies used were

donkey anti-rabbit IgG (Jackson ImmunoResearch, Catalogue Number 711-005-152,

1:100) conjugated with Dy749P1 (Dyomics, Catalogue Number 749P1-01) and Alexa

Fluor 405 (ThermoFisher, Catalogue Number: A30000), donkey anti-mouse IgG

(Jackson ImmunoResearch, Catalogue Number 715-005-150, 1:100) conjugated with

Alexa Fluor 647 (ThermoFisher, Catalogue Number: A20006) and Alexa Fluor 405, and

donkey anti-guinea pig IgG (Jackson ImmunoResearch, Catalogue Number 706-005-

148, 1:100) conjugated with Cy3b (Cytiva, Catalogue Number: PA63101). Tissues were

washed 6 x 20 minutes in 1X PBS at room temperature after secondary antibody

incubation.

Postfixation, dehydration, and embedding in epoxy resin

Tissue embedding was performed as previously described [26]. Tissues were postfixed

with 3% PFA + 0.1% GA (Electron Microscopy Sciences) in PBS for 2 hours at room

temperature and then washed in 1X PBS for 20 minutes. To plasticize the tissues for

ultrasectioning, the tissues were first dehydrated in a graded dilution series of 100%

ethanol (50%/70%/90%/100%/100% EtOH) for 15 minutes each at room temperature

and then immersed in a series of epoxy resin/100% EtOH exchanges (Electron

Microscopy Sciences) with increasing resin concentration (25% resin/75% ethanol; 50%

resin/50% ethanol; 75% resin/25% ethanol; 100% resin; 100% resin) for 2 hours each.

Tissues were transferred to BEEM capsules (Electron Microscopy Sciences) that were

filled with 100% resin and polymerized for 16 hours at 70°C.

Ultrasectioning

510 Plasticized tissue sections were cut using a Leica UC7 ultramicrotome at 70 nm using a

511 Histo Jumbo diamond knife (DiATOME). Chloroform vapor was used to reduce

512 compression after cutting. For each sample, ~100 sections were collected on a

513 coverslip coated with 0.5% gelatin and 0.05% chromium potassium (Sigma-Aldrich Inc.),

514 dried at 60 degrees for 25 minutes, and protected from light prior to imaging.

Imaging chamber preparation

Coverslips were chemically etched in 10% sodium ethoxide for 5 minutes at room

temperature to remove the epoxy resin and expose the dyes to the imaging buffer for

optimal photoswitching. Coverslips were then rinsed with ethanol and dH_2O. To create

fiducial beads for flat-field and chromatic corrections, we mixed 715/755nm and

540/560nm, carboxylate-modified microspheres (Invitrogen, Catalogue Numbers F8799

and F8809, 1:8 ratio respectively) to create a high-density fiducial marker and then

further diluted the mixture at 1:750 with Dulbecco's PBS to create a low-density bead

solution. Both high- and low-density bead solutions were spotted on the coverslip (~0.7

µL each) for flat-field and chromatic aberration correction respectively. Excess beads

were rinsed away with dH2O for 1-2 minutes. The coverslip was attached to a glass

slide with double-sided tape to form an imaging chamber. The chamber was filled with

STORM imaging buffer (10% glucose, 17.5µM glucose oxidase, 708nM catalase, 10mM

MEA, 10mM NaCl, and 200mM Tris) and sealed with epoxy.

Imaging setup

Imaging was performed using a custom single-molecule super-resolution imaging

system. The microscope contained low (4x/10x air) and high (60x 1.4NA oil immersion)

magnitude objectives mounted on a commercial frame (Nikon Ti-U) with back optics

arranged for oblique incident angle illumination. We used continuous-wave lasers at

488nm (Coherent), 561nm (MPB), 647nm (MPB), and 750nm (MPB) to excite Alexa 488,

Cy3B, Alexa 647, and Dy749P1 dyes respectively. A 405 nm cube laser (Coherent) was

used to reactivate Dy749P1 and Alexa647 dye photoswitching. The microscope was

fitted with a custom pentaband/pentanotch dichroic filter set and a motorized emission

filter wheel. The microscope also contained an IR laser-based focus lock system to

maintain optimal focus during automatic image acquisition. Images were collected on

640*640-pixel region of an sCMOS camera (ORCA-Flash4.0 V3, Hamamatsu Photonics)

with a pixel size of ~155 nm.

Automated image acquisition

Fiducials and tissue sections on the coverslip were imaged using the low magnification

objective (4X) to create a mosaic overview of the specimen. Beads/sections were then

imaged at high magnification (60X) to select regions of interest (ROIs) in the Cy3B and

Alexa 488 channels. Before final image acquisition, laser intensities and the incident

angle were adjusted to optimize photoswitching for STORM imaging and utilize the full

dynamic range of the camera for conventional imaging.

Low-density bead images were taken in 16 partially overlapping ROIs. 715/755nm

beads were excited using 750 nm light and images were collected through Dy749P1

and Alexa 647 emission filters. 540/560nm beads were excited using a 488 nm laser

and images were collected through Alexa 647, Cy3B, and Alexa 488 emission filters.

These fiducial images were later used to generate a non-linear warping transform to

correct chromatic aberration. Next, ROIs within each tissue section were imaged at

conventional (diffraction-limited) resolution in all four-color channels sequentially.

Following conventional image acquisition, a partially overlapping series of 9 images

were collected in the high-density bead field for all 4 channels (Dy749P1, Alexa 647,

Cy3B, and Alexa 488). These images were later used to perform a flat-field image

correction of non-uniform laser illumination across the ROIs. Another round of bead

images was taken as described above in a different ROI of the low-density bead field.

These images were later used to confirm the stability of chromatic offsets during

imaging. All ROIs within physical sections were then imaged by STORM for Dy749P1

and Alexa 647 channels. Images were acquired using a custom progression of

increasing 405nm laser intensity to control single-molecule switching. 8000 frames of

Dy749P1 channel images were collected (60 Hz imaging) followed by 12000 frames of

Alexa 647 channel images (100 Hz). In a second imaging pass, the same ROIs were

imaged for Cy3B and Alexa 488 channels, each for 8000 frames (60 Hz).

We imaged the ipsilateral and contralateral ROIs separately in each physical section of

the dLGN. For consistency of ROI selection across biological replicates at each age, we

identified the dorsal-ventral (DV) axis of the dLGN and selected ROIs within the center

(core region) at 2/5 (ipsilateral ROI) and 4/5 (contralateral ROI) of the full DV length.

Image processing

Single-molecule localization was performed using a previously described DAOSTORM

algorithm modified for use with sCMOS cameras [90, 91]. Molecule lists were rendered as

8-bit images with 15.5 nm pixel size where each molecule is plotted as an intensity

distribution with an area reflecting its localization precision. Low-density fiducial images

were used for chromatic aberration correction. We localized 715/755 beads in Dy749P1

and Alexa 647 channels, and 540/560 beads in Alexa 647, Cy3B, and Alexa 488

channels. A third-order polynomial transform map was generated by matching the

positions of each bead in all channels to the Alexa 647 channel. The average residual

error of bead matching was <15 nm for all channels. The transform maps were applied

to both 4-color conventional and STORM images. Conventional images were upscaled

(by 10X) to match the STORM image size. The method to align serial sections was

previously described [26]. STORM images were first aligned to their corresponding

conventional images by image correlation. To generate an aligned 3D image stack from

serial sections, we normalized the intensity of all Alexa 488 images and used these

normalized images to generate both rigid and elastic transformation matrices for all four-

color channels of both STORM and conventional data. The final image stack was then

rotated and cropped to exclude incompletely imaged edge areas. To further confirm that

the processed region corresponds to the contralateral dLGN, conventional CTB(+)

signals of the labeled contralateral projection were thresholded to create a polygonal

mask (a convex hull analysis linking the outermost CTB signals in the image volume).

The mask was then applied to STORM images to exclude peripheral areas where CTB

signals were absent or faint.

Cell body filter

The aligned STORM images had non-specific labeling of cell bodies in Dy749P1 and

Alexa 647 channels corresponding to Homer1 and Bassoon immunolabels. To limit

synaptic cluster identification to the neuropil region we identified cell bodies based on

their Dy749P1 signal and excluded these regions from further image processing.

STORM images were convolved with a Gaussian function (σ=140 nm) and then

binarized using the lower threshold of a two-level Otsu threshold method. We located

connected components in the thresholded images and generated a mask based on

components larger than e^{11} voxels. Because cell body clusters were orders of

magnitude larger than synaptic clusters, the cell body filter algorithm was robust to a

range of size thresholds. The mask was applied to images of all channels to exclude

cell body areas.

Eye-specific synapse identification and quantification

To correct for minor variance in image intensity across physical sections, we normalized

the pixel intensity histogram of each section to the average histogram of all sections.

Image histograms were rescaled to make full use of the 8-bit range. Using a two-level

Otsu threshold method, the conventional images were thresholded into three classes: a

low-intensity background, low-intensity signals above the background representing non-

synaptic labeling, and high-intensity signals representing synaptic structures. The

conventional images were binarized by the lower two-level Otsu threshold, generating a

mask for STORM images to filter out background signals. STORM images were

convolved with a Gaussian function (σ= 77.5 nm) and thresholded using the higher two-

level Otsu threshold. Following thresholding, connected components were identified in

three dimensions using MATLAB 'conncomp' function. A watershedding approach was

applied to split large clusters that were improperly connected. Clusters were kept for

further analysis only if they contained aligned image information across two or more

physical sections. We also removed all edge synapses from our analysis by excluding

synapses that did not have blank image data on all adjacent sides.

To distinguish non-specific immunolabeling from true synaptic signals, we quantified two

parameters for each cluster: cluster volume and cluster signal density calculated by the

ratio of within-cluster pixels with positive signal intensity in the raw STORM images.

Two separate populations were identified in 2D histograms plotted from these two

parameters. We manually selected the population with higher volumes and signal

densities representing synaptic structures. To test the robustness of the manual

selection, we performed multiple repeated measurements of the same data and

discovered a between-measurement variance of <1% (data not shown).

To identify paired pre- and postsynaptic clusters, we first measured the centroid-

centroid distance of each cluster in the Dy749P1 (Homer1) and Alexa 647 (Bassoon)

channels to the closest cluster in the other channel. We next quantified the signal

intensity of each opposing synaptic channel within a 140 nm shell surrounding each

cluster. A 2D histogram was plotted based on the measured centroid-centroid distances

and opposing channel signal densities of each cluster. Paired clusters with closely

positioned centroids and high intensities of apposed channel signal were identified

using the OPTICS algorithm. Retinogeniculate synapses were identified by pairing

Bassoon (Alexa 647) clusters with VGluT2 (Cy3B) clusters using the same method as

pre/post-synaptic pairing. Synapses from the right eye were identified by pairing VGluT2

clusters with CTB (Alexa 488) clusters. The volume of each cluster reflected the total

voxel volume of all connected voxels, and the total signal intensity was a sum of voxel

intensity within the volume of the connected voxels.

Multi-AZ synapse identification and quantification

To determine whether an eye-specific VGluT2 cluster is a mAZ synapse or a sAZ

synapse, we measured the number of active zones (defined by individual Bassoon

clusters) associated with each VGluT2 cluster in the dataset. A 3D shell was extended

140 nm from the surface voxels of each VGluT2 cluster and any Bassoon clusters that

fell within the shell were considered to be associated with the target VGluT2 cluster.

The number of active zones (AZs) associated with each VGluT2 cluster was then

measured. VGluT2 clusters associated with more than 1 AZ were defined as mAZ

synapses, while those associated with only 1 AZ were defined as sAZ synapses.

Quantification of mAZ and sAZ synapse VGluT2 cluster volume was performed using

the "regionprops" function in MATLAB, which provided the voxel size and weighted

centroid of each VGluT2 cluster. The search for sAZ synapses adjacent to mAZ

synapses (synaptic clustering analysis) was conducted using a similar search approach

as for associated Bassoon clusters, with expansion shell sizes ranging from 1 μm to 4

μm from the surface voxels of each mAZ synapse. The main Figs. in the study utilized

an expansion distance of 1.5 µm. An eye-specific sAZ synapse was considered to be

near a mAZ synapse if its weighted centroid fell within the expanded volume.

Quantification and statistical analysis

Statistical analysis was performed using SPSS. Plots were generated by SPSS or R

(ggplot2). Statistical details are presented in the figure legends and supplemental tables

S2/S3. For all measurements in this paper, we analyzed N = 3 biological replicates

(individual mice) for each genotype (WT and β2KO) at each age (P2, P4, and P8).

Cluster densities, synapse AZ number, average VGluT2 cluster volume, and all

fractional measurements were presented as mean ± SEM values in paired bar graphs

and statistical analysis was performed using two-tailed paired-T tests. Nonparametric

Kolmogorov-Smirnov tests were used in all cumulative histogram comparisons. We

used a linear mixed model to compare VGluT2 cluster volumes (Figs. 3 and S2) and

distance measurements (Figs. 5 and S5). For VGluT2 cluster volume comparisons, the

age or eye-of-origin was the fixed main factor and biological replicate IDs were nested

random factors. In distance measurement comparisons, the mAZ synapse AZ number

was the fixed main factor and biological replicate IDs were nested random factors. In

violin plots, each violin showed the distribution of grouped data from all biological

replicates from the same condition. Each black dot represents the median value of each

biological replicate, and the horizontal black line represents the group median. Black

lines connect measurements of CTB(+) and CTB(-) populations from the same

biological replicate. Asterisks in all figures indicate statistical significance: *P<0.05,

**P<0.01, ***P<0.001. For VGluT2 volume comparisons, we calculated 5/95%

confidence intervals based on the linear mixed model. 5/95% confidence intervals in

synapse densities, AZ numbers, and distances were calculated for paired or unpaired

data comparisons using SPSS.

All statistical results are listed in Table S2. All conclusions we draw in the main text from

the data have corresponding confidence intervals listed in both the figure legends and

Table S3.

Acknowledgements:

We thank Drs. Michael C. Crair (Yale University) and Eric M. Ullian (University of

California, San Francisco) for generously sharing the β2KO and ET33[Cre] mouse lines

used in this work. We are grateful to Tim Maugel in the Laboratory for Biological

Ultrastructure (UMD) for his assistance with transmission electron microscopy

experiments. The research was funded by National Institutes of Health

grant DP2MH125812 (C.M.S.).

Author contributions:

Conceptualization, C.Z. and C.M.S.; data curation, C.Z., T.V., C.M.S.;

formal analysis, C.Z. and C.M.S.; funding acquisition, C.M.S.; investigation,

C.Z., T.V., C.M.S.; methodology, C.Z., T.V., C.M.S.; project administration,

C.Z. and C.M.S.; resources, C.Z., T.V., C.M.S.; software, C.Z.

and C.M.S.; supervision, C.M.S.; validation, C.Z., T.V., C.M.S.; visualization,

C.Z., T.V., C.M.S.; writing – original draft preparation, C.Z. and C.M.S.; writing –

review & editing, C.Z., T.V., C.M.S.

Figure legends:

Fig. 1: Retinogeniculate boutons form multiple active zones during eye-specific competition

(A) Experimental design. CTB-Alexa 488 was injected into the right eye of wild-type and β2KO mice. One day after the treatment, tissue was collected from the left dLGN at P2, P4, and P8. Red squares indicate the STORM imaging regions that were analyzed. (B) Representative examples of individual sAZ and mAZ inputs, with corresponding active zone counts ranging from one to three. Upper panels show Z-projections of inputs and lower panels show the corresponding 3D volume. Arrowheads point to individual Bassoon clusters (active zones) paired with postsynaptic Homer1 labels within each input. All examples are from a WT P8 sample. (C) Electron micrographs of mAZ retinogeniculate inputs in a P8 ET33^{Cre}::ROSA26$^{LSL-Matrix-dAPEX2}$ mouse. Darkly-stained dAPEX2(+) mitochondria are present within ipsilaterally-projecting RGC terminals. Arrowheads point to electron-dense material at the postsynaptic density, apposed to individual active zones with clustered presynaptic synaptic vesicles.

Fig. 2: Changes in eye-specific input density during synaptic competition

(A) Representative Z-projection images of mAZ and sAZ inputs across ages and genotypes. Arrowheads point to individual Bassoon/Homer1 cluster pairs indicating release sites. (B) Representative CTB(+) dominant-eye (top panels) and CTB(-) non-

dominant-eye (bottom panels) mAZ inputs in a WT P8 sample, showing synaptic (left

panels), CTB (middle panels), and merged labels (right panels). Arrowheads point to

individual Bassoon/Homer1 paired clusters. (C) Eye-specific mAZ (left) and sAZ (right)

input density across development in WT (top panels) and β2KO mice (bottom panels).

Black dots represent mean values from separate biological replicates and black lines

connect eye-specific measurements within each replicate (N=3 for each age and

genotype). Error bars represent group means ± SEMs. Statistical significance between

eye-specific measurements was assessed using two-tailed paired T-tests at each age.

*p<0.05, **p<0.01.

Fig. 3: Dominant-eye inputs show larger vesicle pools that scale with active zone

number

(A and B) Violin plots showing the distribution of VGluT2 cluster volume for (A) mAZ and

(B) sAZ inputs in WT (filled) and β2KO mice (striped) at each age. The width of each

violin plot reflects the relative synapse proportions across the entire grouped data set at

each age (N=3 biological replicates) and the maximum width was normalized across all

groups. The black dots represent the median value of each biological replicate (N=3),

and the black horizontal lines represent the median value of all inputs grouped across

replicates. Black lines connect measurements of CTB(+) and CTB(-) populations from

the same biological replicate. Statistical significance was determined using a linear

mixed model ANOVA with a post-hoc Bonferroni correction. Black asterisks indicate

significant eye-specific differences at each age. *p<0.05, **p<0.01. ***p<0.001. (C) Eye-

specific VGluT2 signal volume for all inputs separated by number of AZs in WT (left

panel) and β2KO mice (right panel) at P4. (D) Average VGluT2 volume per AZ for all

inputs separated by number of AZs in WT (left panel) and β2KO mice (right panel) at P4.

In panels (C) and (D), error bars indicate group means ± SEMs (N=3 biological

replicates for each age and genotype). Black dots represent mean values from separate

biological replicates and black lines connect eye-specific measurements within each

replicate. Statistical significance between eye-specific measurements was assessed

using two-tailed paired T-tests: *$p<0.05$; **$p<0.01$.

Fig. 4: Eye-specific synapse clustering before eye-opening

(A) Representative mAZ (left panels) and sAZ (right panels) inputs in a WT P8 sample

with nearby sAZ synapses (arrowheads) clustered within 1.5 µm (dashed yellow ring).

Arrows point to the centered mAZ or sAZ inputs. (B) Ratio of clustered and isolated

mAZ and sAZ inputs for CTB(+) (upper panels) and CTB(-) (lower panels) inputs in WT

and β2KO mice at P4. (C) Comparison of the clustered input ratio between mAZ and

sAZ inputs across different ages, genotypes, and eyes of origin. (D) Comparison of the

average number of nearby sAZ synapses for clustered mAZ and sAZ inputs across

different ages, genotypes, and eyes of origin. In panels B-D, black dots represent mean

values from separate biological replicates and black lines connect measurements within

each replicate (N=3 for each age and genotype). Error bars represent group means ±

SEMs. Two-tailed paired T-tests were used to test statistical significance between mAZ

and sAZ inputs at each age. *$p<0.05$; **$p<0.01$.

Fig. 5: Clustered mAZ inputs are closer than isolated inputs during competition

(A-B) Distance between clustered and isolated mAZ inputs and the closest like-eye

clustered mAZ input, shown for (A) CTB(+) and (B) CTB(-) projections at P4 in WT and

β2KO mice. Boxes indicate the 25%-75% distribution of input measurements from N=3

biological replicates and whiskers extend to 1.5 times the interquartile range. Gray dots

represent individual distance measurements for all mAZ inputs. Black and red dots

represent mean values from separate biological replicates and black lines connect

measurements within each replicate (N=3 for each age and genotype). Statistical

significance was determined using a linear mixed model ANOVA with post-hoc

Bonferroni correction. Black asterisks indicate significant differences. *$p<0.05$,

***$p<0.001$.

Fig. S1: Fraction of mAZ inputs across development, related to Fig. 2

Eye-specific mAZ input fraction across development in WT (top panel) and β2KO mice

(bottom panel). Black dots represent mean values from separate biological replicates

and black lines connect measurements within each replicate (N=3 for each age and

genotype). Error bars represent group means ± SEMs. Statistical significance between

eye-specific measurements was assessed using two-tailed paired T-tests. **$p<0.01$,

***$p<0.001$

Fig. S2: Quantification of docked vesicle pool volume and AZ number in mAZ and

sAZ inputs, related to Fig. 2.

(A-C) Violin plots show the distribution of VGluT2 cluster volume within a 70 nm shell

surround individual Bassoon clusters at (A) P2, (B) P4, and (C) P8. The width of each

violin plot reflects the relative synapse proportions at each volume across the entire

grouped data set (N=3 biological replicates). The maximum width of the violin plots was

normalized across all groups. Black horizontal lines represent the median value of all

inputs grouped across replicates. The black dots represent the median value of each

biological replicate and black lines between dots connect measurements from the same

biological replicate. Statistical significance was determined using a linear mixed model

ANOVA with post-hoc Bonferroni correction. See Table S3 for 5/95% confidence

intervals. (D-E) Average AZ number per mAZ input in (D) WT and (E) β2KO mice. Black

dots represent mean values from separate biological replicates (N = 3) and black lines

connect measurements within each replicate. Error bars represent means ± SEMs.

Statistical significance was assessed using two-tailed paired T-tests. *$p<0.05$.

Fig. S3: Relationship between vesicle pool volume and active zone number,

related to Fig. 3.

(A) Total VGluT2 volume per input and (B) average VGluT2 volume per AZ for all inputs

separated by number of AZs in WT (filled bars) and β2KO mice (striped bars) at P2.

(C) Total VGluT2 volume per input and (D) average VGluT2 volume per AZ for all inputs

separated by number of AZs in WT (filled bars) and β2KO mice (striped bars) at P8. In

all panels, error bars indicate group means ± SEMs (N=3 biological replicates for each

age and genotype). Black dots represent mean values from separate biological

replicates and black lines connect measurements within each replicate. Statistical

significance between eye-specific measurements was assessed using two-tailed paired

T-tests: *$p<0.05$; **$p<0.01$.

Fig. S4: sAZ synapse clustering near like-eye mAZ inputs, related to Fig. 4

(A) The ratio of CTB(+) dominant-eye and (B) CTB(-) non-dominant-eye sAZ synapses

nearby like-eye mAZ inputs within increasing distance cutoffs in WT samples across

ages. At distances of 1-2 µm across all ages, the observed ratios (filled bars) are higher

than a reshuffling of the data (open bars) where the position of sAZ inputs was

randomized within the neuropil. (C-D) Percentage of sAZ synapses within 1.5 µm of

opposite-eye mAZ inputs across ages in WT and β2KO mice. Data are compared

against a randomization of sAZ positions (open bars) as in panels A/B. For all panels,

error bars represent group means ± SEMs (N=3 biological replicates for each age and

genotype). Black dots represent mean values from separate biological replicates and

black lines connect measurements within each replicate. Statistical significance was

assessed using two-tailed paired T-tests: *p<0.05; **p<0.01; ***p<0.001.

Fig. S5: Clustered and isolated mAZ inputs show similar spacing after

competition, related to Fig. 5.

(A-B) Distance between clustered and isolated mAZ inputs and the closest like-eye

clustered mAZ input, shown for (A) CTB(+) and (B) CTB(-) projections at P8 in WT and

β2KO mice. Boxes indicate the 25%-75% distribution of input measurements from N=3

biological replicates and whiskers extend to 1.5 times the interquartile range. Gray dots

represent individual distance measurements for all mAZ inputs. Black and red dots

represent mean values from separate biological replicates and black lines connect

measurements within each replicate (N=3 for each age and genotype). Statistical

significance was determined using a linear mixed model ANOVA with post-hoc

Bonferroni correction (see Table S3 for 5/95% confidence intervals).

Fig. S6: sAZ synapse vesicle pool volume is independent of distance to mAZ

inputs, related to Fig. 5.

(A) Cumulative distributions of VGluT2 volume for CTB(+) dominant-eye or (B) CTB(-)

non-dominant-eye sAZ synapses near (<1.5 μm) or far from (>1.5 μm) like-eye mAZ

inputs. The distributions show merged data across all ages (P2/P4/P8; N=3 biological

replicates at each time point). A nonparametric Kolmogorov-Smirnov test was used for

statistical analysis (see Table S3 for 5/95% confidence intervals).

References:

1.   Rall W. Electrophysiology of a dendritic neuron model. Biophys J. 1962;2(2 Pt 2):145-67.
doi: 10.1016/s0006-3495(62)86953-7. PubMed PMID: 14490040; PMCID: PMC1366481.
2.   Mel B, editor. The Clusteron: Toward a Simple Abstraction for a Complex Neuron1991
1991: Morgan-Kaufmann.
3.   Poirazi P, Mel BW. Impact of active dendrites and structural plasticity on the memory
capacity of neural tissue. Neuron. 2001;29(3):779-96. doi: 10.1016/s0896-6273(01)00252-5.
PubMed PMID: 11301036.
4.   Mel BW, Schiller J, Poirazi P. Synaptic plasticity in dendrites: complications and coping
strategies. Curr Opin Neurobiol. 2017;43:177-86. Epub 20170425. doi:
10.1016/j.conb.2017.03.012. PubMed PMID: 28453975.
5.   Kastellakis G, Poirazi P. Synaptic Clustering and Memory Formation. Front Mol Neurosci.
2019;12:300. Epub 20191206. doi: 10.3389/fnmol.2019.00300. PubMed PMID: 31866824;
PMCID: PMC6908852.
6.   Winnubst J, Lohmann C. Synaptic clustering during development and learning: the why,
when, and how. Front Mol Neurosci. 2012;5:70. Epub 20120531. doi:
10.3389/fnmol.2012.00070. PubMed PMID: 22666187; PMCID: PMC3364493.

7.   Leighton AH, Lohmann C. The Wiring of Developing Sensory Circuits-From Patterned
Spontaneous Activity to Synaptic Plasticity Mechanisms. Front Neural Circuits. 2016;10:71. Epub
20160905. doi: 10.3389/fncir.2016.00071. PubMed PMID: 27656131; PMCID: PMC5011135.
8.   Kirchner JH, Gjorgjieva J. Emergence of synaptic organization and computation in
dendrites2022;28(1):21-30. doi: doi:10.1515/nf-2021-0031.
9.   Liang L, Chen C. Organization, Function, and Development of the Mouse
Retinogeniculate Synapse. Annu Rev Vis Sci. 2020;6:261-85. doi: 10.1146/annurev-vision-
121219-081753. PubMed PMID: 32936733.
10.   Bickford ME, Slusarczyk A, Dilger EK, Krahe TE, Kucuk C, Guido W. Synaptic development
of the mouse dorsal lateral geniculate nucleus. J Comp Neurol. 2010;518(5):622-35. doi:
10.1002/cne.22223. PubMed PMID: 20034053; PMCID: PMC4278806.
11.   Monavarfeshani A, Stanton G, Van Name J, Su K, Mills WA, 3rd, Swilling K, Kerr A,
Huebschman NA, Su J, Fox MA. LRRTM1 underlies synaptic convergence in visual thalamus. Elife.
2018;7. Epub 20180209. doi: 10.7554/eLife.33498. PubMed PMID: 29424692; PMCID:
PMC5826289.
12.   Hong YK, Park S, Litvina EY, Morales J, Sanes JR, Chen C. Refinement of the
retinogeniculate synapse by bouton clustering. Neuron. 2014;84(2):332-9. Epub 20141002. doi:
10.1016/j.neuron.2014.08.059. PubMed PMID: 25284005; PMCID: PMC4322918.
13.   Mason CA. Development of terminal arbors of retino-geniculate axons in the kitten--I.
Light microscopical observations. Neuroscience. 1982;7(3):541-59. doi: 10.1016/0306-
4522(82)90063-x. PubMed PMID: 7070666.
14.   Hamos JE, Van Horn SC, Raczkowski D, Sherman SM. Synaptic circuits involving an
individual retinogeniculate axon in the cat. J Comp Neurol. 1987;259(2):165-92. doi:
10.1002/cne.902590202. PubMed PMID: 3584556.
15.   Hammer S, Monavarfeshani A, Lemon T, Su J, Fox MA. Multiple Retinal Axons Converge
onto Relay Cells in the Adult Mouse Thalamus. Cell Rep. 2015;12(10):1575-83. Epub 20150828.
doi: 10.1016/j.celrep.2015.08.003. PubMed PMID: 26321636; PMCID: PMC5757867.
16.   Morgan JL, Berger DR, Wetzel AW, Lichtman JW. The Fuzzy Logic of Network
Connectivity in Mouse Visual Thalamus. Cell. 2016;165(1):192-206. doi:
10.1016/j.cell.2016.02.033. PubMed PMID: 27015312; PMCID: PMC4808248.
17.   Budisantoso T, Matsui K, Kamasawa N, Fukazawa Y, Shigemoto R. Mechanisms
Underlying Signal Filtering at a Multisynapse Contact. J Neurosci. 2012;32(7):2357. doi:
10.1523/JNEUROSCI.5243-11.2012.
18.   Sherman SM, Guillery RW. On the actions that one nerve cell can have on another:
distinguishing "drivers" from "modulators". Proc Natl Acad Sci U S A. 1998;95(12):7121-6. doi:
10.1073/pnas.95.12.7121. PubMed PMID: 9618549; PMCID: PMC22761.
19.   Guillery RW, Sherman SM. Thalamic Relay Functions and Their Role in Corticocortical
Communication: Generalizations from the Visual System. Neuron. 2002;33(2):163-75. doi:
10.1016/S0896-6273(01)00582-7.
20.   Sretavan D, Shatz CJ. Prenatal development of individual retinogeniculate axons during
the period of segregation. Nature. 1984;308(5962):845-8. doi: 10.1038/308845a0. PubMed
PMID: 6201743.

21.   Sretavan DW, Shatz CJ. Prenatal development of retinal ganglion cell axons: segregation
into eye-specific layers within the cat's lateral geniculate nucleus. J Neurosci. 1986;6(1):234-51.
doi: 10.1523/JNEUROSCI.06-01-00234.1986. PubMed PMID: 3944621; PMCID: PMC6568631.
22.   Dhande OS, Hua EW, Guh E, Yeh J, Bhatt S, Zhang Y, Ruthazer ES, Feller MB, Crair MC.
Development of single retinofugal axon arbors in normal and beta2 knock-out mice. J Neurosci.
2011;31(9):3384-99. doi: 10.1523/JNEUROSCI.4899-10.2011. PubMed PMID: 21368050; PMCID:
PMC3060716.
23.   Campbell G, Shatz CJ. Synapses formed by identified retinogeniculate axons during the
segregation of eye input. J Neurosci. 1992;12(5):1847-58. doi: 10.1523/JNEUROSCI.12-05-
01847.1992. PubMed PMID: 1578274; PMCID: PMC6575897.
24.   Chen C, Regehr WG. Developmental remodeling of the retinogeniculate synapse.
Neuron. 2000;28(3):955-66. doi: 10.1016/s0896-6273(00)00166-5. PubMed PMID: 11163279.
25.   Hauser JL, Liu X, Litvina EY, Chen C. Prolonged synaptic currents increase relay neuron
firing at the developing retinogeniculate synapse. J Neurophysiol. 2014;112(7):1714-28. Epub
20140625. doi: 10.1152/jn.00451.2014. PubMed PMID: 24966302; PMCID: PMC4157180.
26.   Zhang C, Yadav S, Speer CM. The synaptic basis of activity-dependent eye-specific
competition. Cell Rep. 2023;42(2):112085. Epub 20230207. doi: 10.1016/j.celrep.2023.112085.
PubMed PMID: 36753422.
27.   Vatan T, Minehart JA, Zhang C, Agarwal V, Yang J, Speer CM. Volumetric super-
resolution imaging by serial ultrasectioning and stochastic optical reconstruction microscopy in
mouse neural tissue. STAR Protocols. 2021;2(4):100971. doi: 10.1016/j.xpro.2021.100971.
28.   Muir-Robinson G, Hwang BJ, Feller MB. Retinogeniculate axons undergo eye-specific
segregation in the absence of eye-specific layers. J Neurosci. 2002;22(13):5259-64. doi:
10.1523/jneurosci.22-13-05259.2002. PubMed PMID: 12097474; PMCID: PMC6758243.
29.   Xu HP, Burbridge TJ, Chen MG, Ge X, Zhang Y, Zhou ZJ, Crair MC. Spatial pattern of
spontaneous retinal waves instructs retinotopic map refinement more than activity frequency.
Dev Neurobiol. 2015;75(6):621-40. Epub 20150330. doi: 10.1002/dneu.22288. PubMed PMID:
25787992; PMCID: PMC4697738.
30.   Xu HP, Furman M, Mineur YS, Chen H, King SL, Zenisek D, Zhou ZJ, Butts DA, Tian N,
Picciotto MR, Crair MC. An instructive role for patterned spontaneous retinal activity in mouse
visual map development. Neuron. 2011;70(6):1115-27. doi: 10.1016/j.neuron.2011.04.028.
PubMed PMID: 21689598; PMCID: PMC3119851.
31.   Xu HP, Burbridge TJ, Ye M, Chen M, Ge X, Zhou ZJ, Crair MC. Retinal Wave Patterns Are
Governed by Mutual Excitation among Starburst Amacrine Cells and Drive the Refinement and
Maintenance of Visual Circuits. J Neurosci. 2016;36(13):3871-86. doi:
10.1523/JNEUROSCI.3549-15.2016. PubMed PMID: 27030771; PMCID: PMC4812142.
32.   Rossi FM, Pizzorusso T, Porciatti V, Marubio LM, Maffei L, Changeux JP. Requirement of
the nicotinic acetylcholine receptor beta 2 subunit for the anatomical and functional
development of the visual system. Proc Natl Acad Sci U S A. 2001;98(11):6453-8. Epub
20010508. doi: 10.1073/pnas.101120998. PubMed PMID: 11344259; PMCID: PMC33489.
33.   Grubb MS, Rossi FM, Changeux JP, Thompson ID. Abnormal functional organization in
the dorsal lateral geniculate nucleus of mice lacking the beta 2 subunit of the nicotinic
acetylcholine receptor. Neuron. 2003;40(6):1161-72. doi: 10.1016/s0896-6273(03)00789-x.
PubMed PMID: 14687550.

34. Sun C, Warland DK, Ballesteros JM, van der List D, Chalupa LM. Retinal waves in mice
lacking the beta2 subunit of the nicotinic acetylcholine receptor. Proc Natl Acad Sci U S A.
2008;105(36):13638-43. Epub 20080829. doi: 10.1073/pnas.0807178105. PubMed PMID:
18757739; PMCID: PMC2527347.
35. Stafford BK, Sher A, Litke AM, Feldheim DA. Spatial-temporal patterns of retinal waves
underlying activity-dependent refinement of retinofugal projections. Neuron. 2009;64(2):200-
12. doi: 10.1016/j.neuron.2009.09.021. PubMed PMID: 19874788; PMCID: PMC2771121.
36. Bansal A, Singer JH, Hwang BJ, Xu W, Beaudet A, Feller MB. Mice lacking specific
nicotinic acetylcholine receptor subunits exhibit dramatically altered spontaneous activity
patterns and reveal a limited role for retinal waves in forming ON and OFF circuits in the inner
retina. J Neurosci. 2000;20(20):7672-81. doi: 10.1523/jneurosci.20-20-07672.2000. PubMed
PMID: 11027228; PMCID: PMC6772851.
37. Hammer S, Carrillo GL, Govindaiah G, Monavarfeshani A, Bircher JS, Su J, Guido W, Fox
MA. Nuclei-specific differences in nerve terminal distribution, morphology, and development in
mouse visual thalamus. Neural Dev. 2014;9:16. Epub 20140710. doi: 10.1186/1749-8104-9-16.
PubMed PMID: 25011644; PMCID: PMC4108237.
38. Zhang Q, Lee WA, Paul DL, Ginty DD. Multiplexed peroxidase-based electron microscopy
labeling enables simultaneous visualization of multiple cell types. Nat Neurosci. 2019;22(5):828-
39. Epub 20190318. doi: 10.1038/s41593-019-0358-7. PubMed PMID: 30886406; PMCID:
PMC6555422.
39. Koch SM, Dela Cruz CG, Hnasko TS, Edwards RH, Huberman AD, Ullian EM. Pathway-
specific genetic attenuation of glutamate release alters select features of competition-based
visual circuit refinement. Neuron. 2011;71(2):235-42. doi: 10.1016/j.neuron.2011.05.045.
PubMed PMID: 21791283; PMCID: PMC3375067.
40. Su J, Byer L, Liang Y, Fox MA. Distribution, development, and identity of retinal ganglion
cells labeled in the Sert-Cre reporter mouse. Journal of Comparative Neurology.
2024;532(3):e25606. doi: https://doi.org/10.1002/cne.25606.
41. Johnson KP, Fitzpatrick MJ, Zhao L, Wang B, McCracken S, Williams PR, Kerschensteiner
D. Cell-type-specific binocular vision guides predation in mice. Neuron. 2021;109(9):1527-39.e4.
Epub 20210329. doi: 10.1016/j.neuron.2021.03.010. PubMed PMID: 33784498; PMCID:
PMC8112612.
42. Assali A, Gaspar P, Rebsam A. Activity dependent mechanisms of visual map formation--
from retinal waves to molecular regulators. Semin Cell Dev Biol. 2014;35:136-46. Epub
20140822. doi: 10.1016/j.semcdb.2014.08.008. PubMed PMID: 25152335.
43. Fassier C, Nicol X. Retinal Axon Interplay for Binocular Mapping. Front Neural Circuits.
2021;15:679440. Epub 20210604. doi: 10.3389/fncir.2021.679440. PubMed PMID: 34149367;
PMCID: PMC8213063.
44. Penn AA, Riquelme PA, Feller MB, Shatz CJ. Competition in retinogeniculate patterning
driven by spontaneous activity. Science. 1998;279(5359):2108-12. doi:
10.1126/science.279.5359.2108. PubMed PMID: 9516112.
45. Assali A, Le Magueresse C, Bennis M, Nicol X, Gaspar P, Rebsam A. RIM1/2 in retinal
ganglion cells are required for the refinement of ipsilateral axons and eye-specific segregation.
Sci Rep. 2017;7(1):3236. Epub 20170612. doi: 10.1038/s41598-017-03361-0. PubMed PMID:
28607399; PMCID: PMC5468276.

46.   Fredj NB, Hammond S, Otsuna H, Chien C-B, Burrone J, Meyer MP. Synaptic Activity and
Activity-Dependent Competition Regulates Axon Arbor Maturation, Growth Arrest, and
Territory in the Retinotectal Projection. J Neurosci. 2010;30(32):10939. doi:
10.1523/JNEUROSCI.1556-10.2010.
47.   Hua JY, Smear MC, Baier H, Smith SJ. Regulation of axon growth in vivo by activity-based
competition. Nature. 2005;434(7036):1022-6. doi: 10.1038/nature03409. PubMed PMID:
15846347.
48.   Rahman TN, Munz M, Kutsarova E, Bilash OM, Ruthazer ES. Stentian structural plasticity
in the developing visual system. Proc Natl Acad Sci U S A. 2020;117(20):10636-8. Epub
20200504. doi: 10.1073/pnas.2001107117. PubMed PMID: 32366647; PMCID: PMC7245132.
49.   Munz M, Gobert D, Schohl A, Poquérusse J, Podgorski K, Spratt P, Ruthazer ES. Rapid
Hebbian axonal remodeling mediated by visual stimulation. Science. 2014;344(6186):904-9. doi:
10.1126/science.1251593. PubMed PMID: 24855269.
50.   Zhang LI, Tao HW, Holt CE, Harris WA, Poo M-m. A critical window for cooperation and
competition among developing retinotectal synapses. Nature. 1998;395(6697):37-44. doi:
10.1038/25665.
51.   Matsumoto N, Barson D, Liang L, Crair MC. Hebbian instruction of axonal connectivity by
endogenous correlated spontaneous activity. Science. 2024;385(6710):eadh7814. Epub
20240816. doi: 10.1126/science.adh7814. PubMed PMID: 39146415.
52.   Yasuda M, Nagappan-Chettiar S, Johnson-Venkatesh EM, Umemori H. An activity-
dependent determinant of synapse elimination in the mammalian brain. Neuron.
2021;109(8):1333-49 e6. Epub 20210325. doi: 10.1016/j.neuron.2021.03.006. PubMed PMID:
33770504; PMCID: PMC8068677.
53.   Louail A, Sierksma MC, Chaffiol A, Baudet S, Assali A, Couvet S, Nedjam M, Roche F,
Zagar Y, Duebel J, Nicol X. cAMP-Dependent Co-stabilization of Axonal Arbors from Adjacent
Developing Neurons. Cell Rep. 2020;33(1):108220. doi: 10.1016/j.celrep.2020.108220. PubMed
PMID: 33027659.
54.   Kutsarova E, Schohl A, Munz M, Wang A, Zhang YY, Bilash OM, Ruthazer ES. BDNF
signaling in correlation-dependent structural plasticity in the developing visual system. PLoS
Biol. 2023;21(4):e3002070. Epub 20230403. doi: 10.1371/journal.pbio.3002070. PubMed PMID:
37011100; PMCID: PMC10101647.
55.   Godement P, Salaün J, Imbert M. Prenatal and postnatal development of
retinogeniculate and retinocollicular projections in the mouse. J Comp Neurol.
1984;230(4):552-75. doi: 10.1002/cne.902300406. PubMed PMID: 6520251.
56.   Mooney R, Madison DV, Shatz CJ. Enhancement of transmission at the developing
retinogeniculate synapse. Neuron. 1993;10(5):815-25. doi: 10.1016/0896-6273(93)90198-z.
PubMed PMID: 8388224.
57.   Ziburkus J, Dilger EK, Lo FS, Guido W. LTD and LTP at the developing retinogeniculate
synapse. J Neurophysiol. 2009;102(6):3082-90. Epub 20090923. doi: 10.1152/jn.90618.2008.
PubMed PMID: 19776360; PMCID: PMC2804430.
58.   Butts DA, Kanold PO, Shatz CJ. A burst-based "Hebbian" learning rule at retinogeniculate
synapses links retinal waves to activity-dependent refinement. PLoS Biol. 2007;5(3):e61. doi:
10.1371/journal.pbio.0050061. PubMed PMID: 17341130; PMCID: PMC1808114.

59.   Lee H, Brott BK, Kirkby LA, Adelson JD, Cheng S, Feller MB, Datwani A, Shatz CJ. Synapse
elimination and learning rules co-regulated by MHC class I H2-Db. Nature. 2014;509(7499):195-
200. Epub 20140330. doi: 10.1038/nature13154. PubMed PMID: 24695230; PMCID:
PMC4016165.
60.   Chung WS, Clarke LE, Wang GX, Stafford BK, Sher A, Chakraborty C, Joung J, Foo LC,
Thompson A, Chen C, Smith SJ, Barres BA. Astrocytes mediate synapse elimination through
MEGF10 and MERTK pathways. Nature. 2013;504(7480):394-400. Epub 20131124. doi:
10.1038/nature12776. PubMed PMID: 24270812; PMCID: PMC3969024.
61.   Stevens B, Allen NJ, Vazquez LE, Howell GR, Christopherson KS, Nouri N, Micheva KD,
Mehalow AK, Huberman AD, Stafford B, Sher A, Litke AM, Lambris JD, Smith SJ, John SW, Barres
BA. The classical complement cascade mediates CNS synapse elimination. Cell.
2007;131(6):1164-78. doi: 10.1016/j.cell.2007.10.036. PubMed PMID: 18083105.
62.   Schafer DP, Lehrman EK, Kautzman AG, Koyama R, Mardinly AR, Yamasaki R, Ransohoff
RM, Greenberg ME, Barres BA, Stevens B. Microglia sculpt postnatal neural circuits in an activity
and complement-dependent manner. Neuron. 2012;74(4):691-705. doi:
10.1016/j.neuron.2012.03.026. PubMed PMID: 22632727; PMCID: PMC3528177.
63.   Hooks BM, Chen C. Distinct roles for spontaneous and visual activity in remodeling of
the retinogeniculate synapse. Neuron. 2006;52(2):281-91. doi: 10.1016/j.neuron.2006.07.007.
PubMed PMID: 17046691.
64.   Dilger EK, Krahe TE, Morhardt DR, Seabrook TA, Shin HS, Guido W. Absence of plateau
potentials in dLGN cells leads to a breakdown in retinogeniculate refinement. J Neurosci.
2015;35(8):3652-62. doi: 10.1523/jneurosci.2343-14.2015. PubMed PMID: 25716863; PMCID:
PMC4339365.
65.   Cork RJ, Namkung Y, Shin HS, Mize RR. Development of the visual pathway is disrupted
in mice with a targeted disruption of the calcium channel beta(3)-subunit gene. J Comp Neurol.
2001;440(2):177-91. doi: 10.1002/cne.1378. PubMed PMID: 11745616.
66.   Fernández-Nogales M, López-Cascales MT, Murcia-Belmonte V, Escalante A, Fernández-
Albert J, Muñoz-Viana R, Barco A, Herrera E. Multiomic Analysis of Neurons with Divergent
Projection Patterns Identifies Novel Regulators of Axon Pathfinding. Adv Sci (Weinh).
2022;9(29):e2200615. Epub 20220821. doi: 10.1002/advs.202200615. PubMed PMID:
35988153; PMCID: PMC9561852.
67.   Wang Q, Marcucci F, Cerullo I, Mason C. Ipsilateral and Contralateral Retinal Ganglion
Cells Express Distinct Genes during Decussation at the Optic Chiasm. eneuro.
2016;3(6):ENEURO.0169-16.2016. doi: 10.1523/ENEURO.0169-16.2016.
68.   Nicol X, Voyatzis S, Muzerelle A, Narboux-Nême N, Südhof TC, Miles R, Gaspar P. cAMP
oscillations and retinal activity are permissive for ephrin signaling during the establishment of
the retinotopic map. Nat Neurosci. 2007;10(3):340-7. Epub 20070128. doi: 10.1038/nn1842.
PubMed PMID: 17259982.
69.   Balice-Gordon RJ, Chua CK, Nelson CC, Lichtman JW. Gradual loss of synaptic cartels
precedes axon withdrawal at developing neuromuscular junctions. Neuron. 1993;11(5):801-15.
doi: 10.1016/0896-6273(93)90110-d. PubMed PMID: 8240805.
70.   Gan WB, Lichtman JW. Synaptic segregation at the developing neuromuscular junction.
Science. 1998;282(5393):1508-11. doi: 10.1126/science.282.5393.1508. PubMed PMID:
9822385.

71.   Wyatt RM, Balice-Gordon RJ. Activity-dependent elimination of neuromuscular synapses.
J Neurocytol. 2003;32(5-8):777-94. doi: 10.1023/b:Neur.0000020623.62043.33. PubMed PMID:
15034267.
72.   Chen J, Mizushige T, Nishimune H. Active zone density is conserved during synaptic
growth but impaired in aged mice. J Comp Neurol. 2012;520(2):434-52. doi: 10.1002/cne.22764.
PubMed PMID: 21935939; PMCID: PMC3731066.
73.   Balice-Gordon RJ, Lichtman JW. Long-term synapse loss induced by focal blockade of
postsynaptic receptors. Nature. 1994;372(6506):519-24. doi: 10.1038/372519a0. PubMed PMID:
7990923.
74.   Buffelli M, Burgess RW, Feng G, Lobe CG, Lichtman JW, Sanes JR. Genetic evidence that
relative synaptic efficacy biases the outcome of synaptic competition. Nature.
2003;424(6947):430-4. doi: 10.1038/nature01844. PubMed PMID: 12879071.
75.   Kopp DM, Perkel DJ, Balice-Gordon RJ. Disparity in neurotransmitter release probability
among competing inputs during neuromuscular synapse elimination. J Neurosci.
2000;20(23):8771-9. doi: 10.1523/jneurosci.20-23-08771.2000. PubMed PMID: 11102485;
PMCID: PMC6773052.
76.   Sanes JR, Lichtman JW. Development of the vertebrate neuromuscular junction. Annu
Rev Neurosci. 1999;22:389-442. doi: 10.1146/annurev.neuro.22.1.389. PubMed PMID:
10202544.
77.   Misgeld T, Kummer TT, Lichtman JW, Sanes JR. Agrin promotes synaptic differentiation
by counteracting an inhibitory effect of neurotransmitter. Proc Natl Acad Sci USA.
2005;102(31):11088-93. doi: 10.1073/pnas.0504806102.
78.   Lin W, Dominguez B, Yang J, Aryal P, Brandon EP, Gage FH, Lee KF. Neurotransmitter
acetylcholine negatively regulates neuromuscular synapse formation by a Cdk5-dependent
mechanism. Neuron. 2005;46(4):569-79. doi: 10.1016/j.neuron.2005.04.002. PubMed PMID:
15944126.
79.   Hashimoto K, Kano M. Synapse elimination in the developing cerebellum. Cell Mol Life
Sci. 2013;70(24):4667-80. Epub 20130628. doi: 10.1007/s00018-013-1405-2. PubMed PMID:
23811844; PMCID: PMC3830199.
80.   Bosman LW, Konnerth A. Activity-dependent plasticity of developing climbing fiber-
Purkinje cell synapses. Neuroscience. 2009;162(3):612-23. Epub 20090123. doi:
10.1016/j.neuroscience.2009.01.032. PubMed PMID: 19302832.
81.   Hashimoto K, Kano M. Functional differentiation of multiple climbing fiber inputs during
synapse elimination in the developing cerebellum. Neuron. 2003;38(5):785-96. doi:
10.1016/s0896-6273(03)00298-8. PubMed PMID: 12797962.
82.   Nitta A, Yamasaki M, Miyazaki T, Konno K, Yoshimura H, Watanabe M. Molecular and
Anatomical Strengthening of "Winner" Climbing Fiber Synapses in Developing Mouse Purkinje
Cells. J Neurosci. 2025. Epub 20250227. doi: 10.1523/JNEUROSCI.2156-24.2025. PubMed PMID:
40015986.
83.   Wilson AM, Schalek R, Suissa-Peleg A, Jones TR, Knowles-Barley S, Pfister H, Lichtman
JW. Developmental Rewiring between Cerebellar Climbing Fibers and Purkinje Cells Begins with
Positive Feedback Synapse Addition. Cell Rep. 2019;29(9):2849-61.e6. doi:
https://doi.org/10.1016/j.celrep.2019.10.081.

84.   Bosman LWJ, Takechi H, Hartmann J, Eilers J, Konnerth A. Homosynaptic Long-Term
Synaptic Potentiation of the "Winner" Climbing Fiber Synapse in Developing Purkinje Cells. J
Neurosci. 2008;28(4):798. doi: 10.1523/JNEUROSCI.4074-07.2008.
85.   Ohtsuki G, Hirano T. Bidirectional plasticity at developing climbing fiber-Purkinje neuron
synapses. Eur J Neurosci. 2008;28(12):2393-400. Epub 20081121. doi: 10.1111/j.1460-
9568.2008.06539.x. PubMed PMID: 19032589.
86.   Lorenzetto E, Caselli L, Feng G, Yuan W, Nerbonne JM, Sanes JR, Buffelli M. Genetic
perturbation of postsynaptic activity regulates synapse elimination in developing cerebellum.
Proc Natl Acad Sci USA. 2009;106(38):16475-80. doi: 10.1073/pnas.0907298106.
87.   Kawamura Y, Nakayama H, Hashimoto K, Sakimura K, Kitamura K, Kano M. Spike timing-
dependent selective strengthening of single climbing fibre inputs to Purkinje cells during
cerebellar development. Nat Commun. 2013;4(1):2732. doi: 10.1038/ncomms3732.
88.   Benninger K, Hood G, Simmel D, Tuite L, Wetzel A, Ropelewski A, Watkins S, Watson A,
Bruchez M. Cyberinfrastructure of a Multi-Petabyte Microscopy Resource for Neuroscience
Research. Practice and Experience in Advanced Research Computing; Portland, OR, USA:
Association for Computing Machinery; 2020. p. 1–7.
89.   Paxinos G. Atlas of the developing mouse brain at E17.5, P0 and P6. 1st ed. Amsterdam ;
Boston: Elsevier; 2007. xi, 353 p. p.
90.   Babcock H, Sigal YM, Zhuang X. A high-density 3D localization algorithm for stochastic
optical reconstruction microscopy. Optical Nanoscopy. 2012;1(6). doi: 10.1186/2192-2853-1-6.
91.   Babcock HP, Huang F, Speer CM. Correcting Artifacts in Single Molecule Localization
Microscopy Analysis Arising from Pixel Quantum Efficiency Differences in sCMOS Cameras. Sci
Rep. 2019;9(1):18058. Epub 20191202. doi: 10.1038/s41598-019-53698-x. PubMed PMID:
31792238; PMCID: PMC6889274.
1167

A

Wild-type   β2KO

versus

STORM imaging in the
contralateral dLGN

P2   P4   P8

B

| | Single-AZ | Multiple-AZ | |
| | One AZ | Two AZs | Three AZs |

Bassoon / Homer1 / VGluT2

1 µm

0.42, 2.33, 2.33 (µm)

2.33, 2.33, 1.26 (µm)

2.33, 1.75, 2.33 (µm)

C

Mitochondria
Vesicle pool
PSD
Terminal

A mAZ inputs

CTB(+) dominant-eye / CTB(-) non-dominant-eye

B sAZ inputs

CTB(+) dominant-eye / CTB(-) non-dominant-eye

C CTB(+) dominant-eye / CTB(-) non-dominant-eye

D CTB(+) dominant-eye / CTB(-) non-dominant-eye

A

Clustered mAZ | Clustered sAZ

Bassoon / Homer1 / VGluT2

2 μm

B

Multi-AZ input

1.5 μm cutoff

Clustered vs **I**solated

Single-AZ input

1.5 μm cutoff

Clustered vs **I**solated

WTP4 | WTP4 | β2P4 | β2P4

CTB(+) — Input ratio (% of total)

C I | C I | C I | C I

WTP4 | WTP4 | β2P4 | β2P4

CTB(−) — Input ratio (% of total)

C I | C I | C I | C I

C

Multi-AZ / Single-AZ

CTB(+) — Clustered input ratio (% of total)

P2 | P4 | P8   P2 | P4 | P8

CTB(−) — Clustered input ratio (% of total)

P2 | P4* | P8   P2 | P4 | P8

Wild-type   β2 KO

D

Multi-AZ / Single-AZ

CTB(+) — sAZ synapses per clustered input

P2 | P4* | P8**   P2 | P4 | P8

CTB(−) — sAZ synapses per clustered input

P2 | P4** | P8**   P2 | P4 | P8

Wild-type   β2 KO

A Distance between all mAZ inputs and nearest like-eye clustered mAZ input

CTB(+) dominant-eye

Clustered   Isolated

WT P4 CTB(+)

β2KO P4 CTB(+)

B Distance between all mAZ inputs and nearest like-eye clustered mAZ input

CTB(-) non-dominant-eye

Clustered   Isolated

WT P4 CTB(-)

β2KO P4 CTB(-)

A Distance between all mAZ inputs and nearest like-eye clustered mAZ input

CTB(+) dominant-eye

Clustered   Isolated

WT P8 CTB(+)

β2KO P8 CTB(+)

B Distance between all mAZ inputs and nearest like-eye clustered mAZ input

CTB(-) non-dominant-eye

Clustered   Isolated

WT P8 CTB(-)

β2KO P8 CTB(-)