Title

Characterization of Z chromosomal inversion and lineage-specific sweeps reveal

widespread local adaptation in *Helicoverpa armigera*

Fushi Ke[a]

[a] School of Biological Sciences, The University of Hong Kong, Pok Fu Lam Road, Hong Kong,

China.

Corresponding author

Fushi Ke: fjkfs@163.com

Abstract

Efficient pest management requires comprehensive knowledge of its biology, ecology, and

evolution, particularly adaptation that exacerbating pest damage. *Helicoverpa armigera* (Hübner)

is a notorious pest that attacks more than 200 species, adapts to diverse environments, and has

developed resistance to almost all applied insecticides. However, local adaptation of this species

especially acclimatization was less investigated. Here, we analyzed genomic variation of

Helicoverpa armigera in China and Oceania to identify adaptative loci in different lineages.

Despite intensive gene flow, two differentiated Z chromosomal lineages in Central China (i.e.,

northern and southern lineages) were formed by chromosome inversion and local adaptation.

Based on population genomic analysis, we identified genes related to circadian clock (*period*,

cycle, *cyr2* and *daywake*), insulin secretion (*IGF2BP2*) and energy metabolism (*SLC25A25*,

CCG7766, *D11DS*, *TSTP* and *CPQ*) were under selection in northern lineage. These genes may

have contributed to its adaptation to high latitudes. In the southern lineage however, the Z

chromosome inversion harbours alleles of two clock genes (*period* and *cycle*) introgressed from

Oceanian lineage (*H. a. conferta*) and may have disrupted clock oscillator for acclimatization. We

further conducted selective sweep analysis in lineages from Northwestern China, Central China,

and Oceania, and found several insecticide resistant genes that shared among different lineages

were likely contributed by introgression. Nevertheless, many candidates, including a cadherin that

involved in resistance to *Bacillus thuringiensis* protein in Oceanian populations, are

lineage-specific. Our results highlight the importance of investigating local adaptation in effective

pest control under globalization of agriculture.

Key words

34   Cotton bollworm; local adaptation; clock genes; energy metabolism; Z chromosome inversion;

35   lineage-specific sweep

Introduction

Agricultural insect pests cause substantial losses of crop production each year and pose a threat to global food security (Paini et al., 2016). It's difficult to implement effective pest control strategies without a thorough understanding of pest biology, ecology and evolution, particularly the adaptive evolution that exacerbates the pest damage (Roderick & Navajas, 2003; Renou, 2022). Investigating the ability and genetic basis of pests adapting to local habitats, in terms of acclimatization and pesticide resistance thus, would benefit successful pest control. For most insect pests, environmental factors play a crucial role in their performance. Identifying genotypes of higher fitness and selective agents causing acclimatization could provide important information for their interaction with local environments, prediction of their further distribution (Pélissié et al., 2018), and implementation of efficient molecular techniques (e.g., through gene editing) for pest control. In the case of resistance to applied pesticides and toxins expressed by transgenic crops, local adaptation could imply that control strategies might need to be tuned according to which populations and regions that are targeted in order to keep their efficiency (Alyokhin et al., 2015; Pélissié et al., 2018; Szendrei et al., 2012).

For insect species, forecasting seasonal changes and responding to them in an appropriate way are species' specific requirements to live in high latitudes or altitudes (Hoikkala & Poikela, 2022). Since temperature and photoperiod are generally reliable indicators of seasonality that have shaped life histories of many temperate climate organisms (Fält-Nardmann et al., 2016), efficient adaptations to thermal stress and photoperiod such as diapause ensure survival and succession of local populations during long overwintering periods (Toepfer et al., 2014). Several pathways have been identified to be underlying the genetic and physiological mechanisms of diapause regulation and evolution (Ragland et al., 2019). Associated genes involved in the circadian clock with diapause were uncovered, with examples showing strong connections between seasonal adaptations and adaptative alleles of clock genes (Ragland et al., 2019; Kozak et al., 2019; Hasebe & Shiga, 2022; Peng et al., 2023). In addition, pathways that insulin involved (Williams et al. 2006; Paaby et al., 2010) and energy metabolism related could also facilitate adaptation of insects to high latitude habitats (Hahn & Denlinger, 2011). Identification of adaptative genes/alleles in the

lineages of high latitude thus could give us a comprehensive knowledge of adaptative evolution

(e.g., diapause) that could be employed in successful management of insect pests, especially those

from temperate regions.

Recombination of genotypes from different genetic background would generate new genetic

combinations that facilitate local adaptation of descendants (Ortiz-Barrientos et al., 2016). It could

also increase interchange of maladaptative alleles that restrict adaptation of local lineages (Crispo

et al., 2011; Arnold et al., 2017). Structural variations such as chromosome inversion that inhibit

recombination would restrict gene reshuffling in heterozygotes and maintain adaptative or

maladaptative genotypes with linkage (Kirkpatrick & Barton 2006; Roesti et al., 2022).

Meanwhile, biotic and abiotic stress that mediated local adaptation would be barrier of gene flow

in adaptative loci during homogenization of global genetic variation in differentiated lineages

(Tigano et al., 2016). Therefore, investigating the interplay of gene flow, local adaptation and

chromosome structural variation would benefit identification of adaptative genes in local

populations.

The cotton bollworm, *Helicoverpa armigera* Hübner (Lepidoptera: Noctuidae), is a worldwide

crop pest that attacks over 200 species (Pratissoli et al., 2015; Riaz et al., 2021) and shows its

capability of adaptation to diverse habitats and hosts. It has also developed resistance to many

applied insecticides with a high number of reported cases of insecticide resistance worldwide

(Klai et al., 2020). Meanwhile, it could migrate over 2,000 km and mediated intensive gene flow

among different populations (Nibouche et al., 1998; Jones et al., 2015; Riaz et al., 2021). Recently,

this pest has invaded to South America, hybridized with its sibling species and introduced

insecticide-resistant genes into local colonies (Valencia-Montoya et al., 2020). Diapause is an

important strategy by which insects, such as *H. armigera*, avoid unfavorable environmental

conditions (Danks, 2000). Successful range expansion, particularly at high latitudes, thus requires

that species should be able to coordinate their phenology and synchronize life cycles using local

temperature and/or day length as cues (Fält-Nardmann et al., 2016; Saikkonen et al., 2012).

Investigating phenological adaptation of source populations to environmental cues would help in

predicting sites of their invasion and implementing effective pest control. While difference in

response to temperature and day length in *H. armigera* lineages have been explored based on

molecular experiments (Wu et al., 1997; Wu & Guo 1997; Wu & Guo 2007; Chen et al., 2013;

Chen et al., 2016; Virachack et al., 2018), less analysis was conducted to investigate the genetic

basis of its acclimation (but see Jin et al., 2023) in the presence of widespread gene flow among

geographically distinct lineages.

In this study, we revisited resequencing data of *H. armigera* samples from Asia and Oceania

(Zhang et al., 2022), and investigated adaptative genes that have contributed to local adaptation of

different lineages. When constructing phylogeny based on Z chromosome polymorphism, we

found two differentiated lineages of *H. armigera* in Central China (i.e., the southern (CC-S) and

northern (CC-N) lineages) despite widespread gene flow across this region. We further verified

intensive introgression between *H. a. conferta* in Oceania and *H. armigera* in Central China, while

gene flow of adaptative loci into CC-N was restricted due to local adaptation, representing a

model of divergent selection with gene flow (Pinho & Hey, 2010). In addition, our results support

Z chromosome inversion harboring *H. a. conferta* alleles of two linked clock genes may have

disrupted acclimation in CC-S, and contributed to differential gene flow of *H. a. conferta* into

northern and southern *H. armigera* lineages in Central China. Finally, we identify several

lineage-specific signals of selection that may be related to insecticide resistance of local

populations. We further discuss our findings under current global climate change scenario and

severe pest damage to agricultural industry.

Materials & Methods

Polymorphic sites and genome resequencing data

We downloaded polymorphic SNPs from figshare (https://figshare.com/articles/dataset/population
_genomics_of_cotton_bollworm/17008922). This dataset includes 141 *Helicoverpa armigera*
individuals from China, 19 individual *H. a. conferta* from Australia, and nine *H. zea* from North
and South America. Samples of China collected from sites at Yellow River Region (YRR),
Changjiang River Region (CRR), and Northwestern Region (NR) during 2016-2018. Genomic
analysis shows high similarity between samples from YRR and CRR (Zhang et al., 2022), we thus
reassigned their geographical origin as Central China (CC). A total of 100 individuals and 41
individuals from CC and NR regions were employed, respectively. Among them, 47, 33 and 10
male individuals with ZZ genotype of sex chromosome, were found in CC, NR *H. armigera*
lineages, and *H. a. conferta*, respectively. See Zhang et al., (2022) for detailed information of
these samples.

To further calculate population genetic parameters, such as d_{xy}, we downloaded sequencing reads
of 141 *H. armigera* individuals (BioProject No. PRJNA731848) and six *H. zea* individuals NCBI
(SRR15526515-20), and called SNPs by using Genome Analysis Toolkit 3.8 (GATK) (DePristo et
al. 2011). Sequencing reads of each individual were filtered by using fastq
(https://github.com/OpenGene/fastp) and mapped to HaSCD2 assembly (Zhang et al., 2022) using
BWA (Li, 2013). The standard GATK pipeline (DePristo et al. 2011) was employed to generate
raw sites from bam files. We further conducted filterings in GATK with default settings to exclude
low-quality sites, and removed loci with mean coverage depth lower than 3 or higher than 30, or
with missing rate higher than 0.5 using VCFtools (Danecek et al. 2011). Unless stated, this dataset
with high-quality non-variants was only used for π and d_{xy} calculation.

Phylogeny tree and genetic structure

Phylogeny trees were reconstructed using three methods: neighbour-joining (NJ) method,
maximum likelihood (ML) method based on concatenated SNPs, and species tree inferred from
windowed trees based on ASTRAL-III (Zhang et al., 2018). For NJ phylogeny, genetic distance

matrix of each individual was calculated using VCF2Dis (https://github.com/hewm2008/VCF2Dis)

and used for phylogeny construction in phylip-3.697

(http://evolution.genetics.washington.edu/phylip). We further reconstructed the phylogeny

relationship based on maximum likelihood (ML) method implemented in IQ-TREE-2.1.3 (Minh et

al. 2020). Bi-allelic SNPs thinned by an interval of 100bp were concatenated and used in ML tree

construction. We employed +ASC model in model testing to include ascertainment bias correction

as SNP data do not contain constant sites. Following the manual, we also appended -bnni to the

regular UFBoot command to reduce the risk of overestimating branch support (Hoang et al. 2018).

The topologies inferred by NJ and ML methods based on autosomal polymorphism were similar,

while topology difference was found when using polymorphic data of Z chromosome. To be

specific, we found two differentiated lineages (CC-S, southern lineage, and CC-N, northern

lineage) in Central China based on ML tree. Thus, we further constructed the individual phylogeny

based on ASTRAL-III using Z chromosomal SNPs. We first divided Z chromosome into 100-kb

windows according to coordinates and constructed window tree with SNPs of each window using

IQTREE (Minh et al. 2020). A summary tree was then generated based on the constructed 179

window trees in ASTRAL-III (Zhang et al., 2018). Five regions in Z chromosome that were

divided by Fst values were further used for tree construction in IQ-TREE.

We further investigated population genetic structure of *H. armigera* (including 19 *H. a.*

conferta individuals) based on autosomal SNPs. We generated SNP datasets by pruning SNPs

based on LD value of 0.2 with an interval of 50 loci and included only SNP loci at a missing rate $\leq$

20% within the intergenic regions 20kb away from the genic regions. We used ADMIXTURE for

genetic structure analyses and performed each K from 1 to 20 with 20 replicates. The best K value

was identified according to cross-entropy criterion. Employing the same dataset, we conducted

Principal Component Analysis (PCA) based on PLINK2 (Chang et al., 2015).

Inversion identification and genome-wide association analysis

We used Asaph (Nowling et al., 2022) to detect chromosomal inversion in samples collected from

Central China as we found two differentiated lineages in phylogeny based on Z chromosomal

polymorphims. Asaph could efficiently detect inversion when samples are all from a single

population and SNPs are all from a single chromosome (Nowling et al., 2022). Two first PC

coordinates of Z chromosomal polymorphism were generated to separate samples in Central China.

Further, single-SNP association tests using genotypes of each SNP against the samples' first PC

coordinate were performed. Without long sequencing reads to identify the accurate boundary of

the inversion, we set an arbitrary boundary of the inversion by picking SNPs with a $-\log_{10}(P) > 15$.

Within the inversion, haplotypes possessing derived alleles in over half of the differentiated SNPs,

we treated them as haplogroups of inversion. To further identify loci that may have contributed to

differentiation of two lineages in China, the GWAS analysis was extended to polymorphic SNPs

in autosomal SNPs of each chromosome. We set an arbitrary while widely used P value (5×10^{-8})

as the threshold, and selected SNPs with a P value lower than the threshold as outliers.

Calculation of population genetic parameters

We calculated nucleotide diversity (π), genetic differentiation (F_{ST}) and genetic divergence (d_{xy})

by using Simon Martin's script (https://github.com/simonhmartin/genomics_general). In addition,

we calculated window-based parameter f_{dM} and D based on Dsuite (Malinsky et al., 2021) to

investigate genome-wide introgression between different lineages. Only male individuals were

included for analysis. Note that we have regenerated a genotype matrix including non-variant sites

for accurate calculation of π and d_{xy}. We further calculated r^2 of SNP pairs in Z chromosome based

on PopLDdecay (https://github.com/BGI-shenzhen/PopLDdecay) and compared linkage

disequibrilum (LD) between lineages including *H. zea*, CC-N *H. armigera* and CC-S *H. armigera.*

To account for different sampling size, we randomly sampled eight individuals in CC-N *H.*

armigera and *H. zea* ten times and used these replicates for further comparisons. In addition, to

compared LD of different regions, we further divided Z chromosome into "inversion" (identified

by Asaph) and "the other" regions, and calculated r^2 of SNP pairs in each region.

Selective sweep analysis

To identify genetic loci underlying local adaptation in each geographically distinct lineage,

selective sweep analysis was performed using site frequency spectrum (SFS) in SWEEPFINDER2

(DeGiorgio et al., 2016). We first assigned *H. zea* as the outgroup and calculated the derived allele

frequency spectrum in each population. We further ran selective sweep analysis in each

chromosome in a step of 1000 nucleotides. Regions with a CLR value at the top 0.05% of all

autosomal regions in each population were identified as outliers. Further, we used a

homozygosity-based method and unphased multilocus genotype (MLG) to calculate G12

parameter in each population (Harris et al., 2018). G12 could employ unphased genotype data and

is robust similar to H12 (using phased genotype, Garud et al., 2015) in identifying selective

sweeps. We don't intend to separate soft and hard sweeps in each population, but to identify

genomic regions under positive natural selection. For *H. armigera* has a similar effective

population size while much higher recombination rate (4.86 x 10^{-6} cM/bp, Zhang et al., 2022)

compared with *Drosophila melanogaster* (5 x 10^{-7}cM/bp, Garud et al., 2015), we used a smaller

window of 100 SNPs in a step of 20 SNPs. Windows with G12 values over 99.95% of autosomal

windows in each population were identified as outliers.

Genetic data illustration

For the phylogeny tree, we visualized the tree by using Itol (https://itol.embl.de/). Several R

packages, including rworldmap (South, 2011), reshape (Wickham, 2007) and rworldxtra (South,

2012) were used for illustrating inversion and ancestral haplogroups on the map. Pearson

correlation was conducted in R and plotted using ggpubr (Kassambara & Kassambara, 2020) and

ggplot2 (Wickham, 2011). Package CMplot (Yin, 2020) was used for generation of Manhattan plot.

In addition, RectChr (https://github.com/BGI-shenzhen/RectChr) were employed for generating

genotype matrix plot.

Results

Two differentiated lineages identified by Z chromosomal variation

The phylogeny reconstructed using Z chromosomal variation identified two differentiated lineages

in Central China, of which all samples in the southern lineage (CC-S) were from sites near

Changjiang River Region (lower latitudes, Figure 1A). This lineage (CC-S) clustered with samples

from Oceania (*H. a. conferta*) and differentiated with other *H. armigera* samples in China (Figure

1A). This topology was further supported by summary tree generated by ASTRAL-III based on

179 window (100kb) trees using Z chromosomal polymorphisms (Supplementary Figure 1). The

genetic pattern however, was differentiated with the neighbour-joint (NJ) phylogeny (Zhang et al.,

2022) and genetic structure (Figure 1B; Supplementary Figure 2) based on autosomal SNPs. To

further characterize regional variation across Z chromosome, we calculated genetic differentiation

between CC-S and CC-N (CC-N vs. AUS for comparison), and divided Z chromosome into five

regions. Of these five regions, two major topologies were identified, with topology one consistent

with the autosomal phylogeny (Zhang et al., 2022) and topology two clustering CC-S *H. armigera*

lineage with *H. a. conferta*. Tree topology in region 4 was consistent with topology two.

Topologies in region 1, region 2 and region 5 that show similarity with topology two have

individuals from CC-N or NR clustering with *H. a. conferta* individuals. Only region 3 has a tree

topology consistent with topology one based on autosomal SNPs (Supplementary Figure 3). In

addition, we found high correlation between $Fst_{(CC-N\ vs.\ CC-S)}$ and $Fst_{(CC-N\ vs.\ AUS)}$ (Figure 1C), which

support widespread introgression of *H. a. confera* into *H. armigera*. This was further supported by

window-based introgression analysis (Figure 1D; Supplementary Figure 4).

Z chromosomal inversion in *H. armigera*

The first principal component (PC) of the Z chromosomal variation divided samples in Central

China into two distinguished clusters (Supplementary Figure 5), which is consistent with two

differentiated lineages in Central China (Figure 1A). PC1 further identified three individuals (Aq4,

Jz42 and DF44) with mixed genetic background. Similar to samples from CC-S, these three

individuals were all from the Changjiang River region. Based on the identified PC, we further

conducted single-SNP association tests using genotypes each SNP against the samples' first PC

coordinate in Asaph. This analysis identified one distinguished peak at Z chromosome, with the

boundary ranges from 6.7 - 12.4 Mb, by setting a threshold of -log10(P) > 15 (Figure 2A). We

further isolated SNPs with $-\log_{10}(P) > 15$ in the inversion region and constructed genotype matrix

of these differentiated SNPs in male individuals of Central China (Figure 2B, see Supplementary

Figure 6 for genotype matrix of all individuals). This inversion region in CC-S lineage was with

higher linkage disequilibrilum (LD) compared with other regions of CC-S as well as CC-n lineage

and AUS (*H. a. conferta*) (Figure 2C). The genotype matrix identified nine individuals with

derived alleles and were with Z chromosomal inversion. The geographical distribution of this

inversion showed all individuals possessing this inversion are restricted to Changjiang River

region (lower latitudes, Figure 2D).

Genetic variation of CC-N was higher than CC-S, the regions with lower π ratio (π_{CC-N}/π_{CC-S})

and higher -$\log_{10}(P)$ thus would be indicatives of selection signals on CC-N lineage (i.e., reduced

genetic variation). We treated regions with π ratio lower than 99% of Z chromosome regions (a

total of 37 20kb-windows) as candidates and further considered the -$\log_{10}(P)$ more than 7.3 ($P = 5$

$\times 10^{-8}$). Interestingly, we identified several genes that maybe potentially important in adaptation of

CC-N and differentiation of CC-N and CC-S lineages. Of which, *SLC25A25* and *IGF2BP2* were

located outside the inversion, while period circadian protein (*per*) and Aryl hydrocarbon receptor

nuclear translocator–like protein 1 (*ARNTL*, or *cycle*) were within the inversion (Figure 2A).

Genotype matrix of *per* and *cycle* show that the male individuals in CC-S have similar derived

genotype with those in *H. a. conferta* males (Figure 3A). In addition, three male individuals

including Aq4, DF44, and Jz22 that are collected from Changjiang River region have the same

derived genotypes at both *per* and *cycle*. Several additional individuals from both CC-N (Ay13,

DF28, DF46, Xj7, Kf2, Kf21, Kf6) and NR (Hami12, Hami7, Hami9, and Kel26) have only

derived genotype in *cycle* (Figure 3A). We also found the genetic differentiation (Fst) and

divergence (dxy) of these two genes elevated compared with surrounding regions, and nucleotide

diversity of two genes were lower in CC-N lineage compared with those in CC-S lineage (Figure

3B). When conducted protein modeling of two divergent haplotypes dominated in each group, we

found difference not only in amino acid but also in protein secondary structure (Figure 3C). In

CC-S lineage, these two clock genes, although separated by a distance of around 4 Mb, show

highly linkage within and between genes (Figure 3D).

Genes contributed to differentiation of southern and northern lineages

In addition to genes in Z chromosome that differentiated between southern and northern *H.*

armigera lineages. Genome-wide association analysis of the autosomal SNP genotype against first

principal component (of Z chromosome variation) further isolated several regions with significant

-$\log10(P)$ value (> 7.3) and elevated genetic differentiation (Fst). These regions spread across the

genome and distributed at more than five chromosomes (Figure 4A). We further checked the

annotated genes in these regions, and identified acyl-CoA Delta (11) desaturase (*D11DS*) and

bifunctional trehalose-6-phosphate synthase/phosphatase (*TPSP*) in Chr12, phosphorylase b

kinase regulatory subunit (*CG7766*), carboxypeptidase Q (*CPQ*) of Chr 13, and cryptochrome-2

(*Cry2*) from Chr15, and circadian clock-controlled protein *daywake* (*annon-3B1.2*) in Chr23.

Interestingly, these genes all have elevated Fst and dxy, which supported a divergent selection

with gene flow in formation of these divergent peaks. In addition, decreased π ratio (π_{CC-N}/π_{CC-S})

of these genes compared with nearby regions supported local adaptation of CC-N population may

have contributed to this divergence.

Widespread selective sweep signals in different *H. armigera* lineages

Several selective sweeps were identified based on strict threshold in different populations (Figure

5). Compared with SWEEPFINDER2, haplotype-based parameter G12 generally identified more

regions with sweep signals. Among them, several linked P450 genes identified including

CYP337B3V1, CYP6B2, and CYP6a21 were shared across three populations (Figure 5). As we

found many regions contributed to differentiation of southern and northern *H. armigera* lineages

of Central China, and intensive while diverse application of insecticide for controlling different *H.*

armigera populations (Wu et al., 2005; Li et al., 2010; Zhang et al., 2012; Yang et al., 2013; Lu et

al., 2021; Downes et al., 2017), different *H. armigera* lineages should have adapted to local

climates (Chen et al., 2013) and regionally specific insecticides. Indeed, we found many

lineage-specific signals of selective sweeps in populations in Central China, Northwestern China,

and Oceania (*H. a. conferta*) based on G12 parameter (Figure 5). In Central China, we found

several GSTs on Chr7, carboxylic ester hydrolase (CCE001c) and Juvenile hormone esterase

(JHER) of Chr 15, and carboxyl/choline esterase (CCE014a) on Chr30 were under selection. In

addition, trehalose-6-phosphate synthase/phosphatase (TPSP) located at Chr12, and Cytochrome

P450 12a4 (CYP12A4) and CPQ of Chr13 that were identified as outliers based on G12. In

Northwestern China, we found several genes with signals of sweeps except CYP337B3 (Figure 5).

In Oceania, cadherin (CadN, Chr 4), choline O-acetyltransferase (ChAT, Chr6), cadherin-89D

(Cad89D, Chr6), GST(Chr7), carboxylic ester hydrolase (CCE001c, Chr15), and juvenile

hormone esterase (JHER, Chr15) showed selective signals. SWEEPFINDER2 identified less but

unique selective signals among different populations. For example, in Central China lineage,

PTP10D on Chr13 showed higher CLR values compared with those of other regions. Similarly,

two genes (D11DS, and Scd1) on Chr12 have higher CLR values in NC lineage.

Discussion

Understanding the genetic basis of local adaptation in different insect pest lineages is important in effective pest control and management. To investigate genetic differentiation and local adaptation of two *H. armigera* lineages in Central China identified by Z chromosome polymorphism, we combined genetic differentiation and gene flow analysis to disentangle effects of introgression, local adaptation and chromosome structure on genome-wide genetic variation. Compared with southern lineage in Central China, we identified several genes in northern lineage of Central China with sweep signals (e.g., low nucleotide diversity and high differentiation), restricting introgression from the other subspecies *H. a. conferta*, and may have contributed to adaptation of this lineage to high latitudes. On the contrary, a Z chromosome inversion in southern lineage, keeping divergent alleles of two circadian clock genes introgressed from *H. a. conferta*, have contributed to differentiation of these two lineages in Central China, and may have led to disorder of clock oscillator for acclimation. Further, we conducted selective sweep analysis based on two parameters and identified several genes that may have contributed to local adaptation in different *H. armigera* lineages (Northwestern China, Central China, and Oceania). Overall, our results reveal widespread local adaptation of different *H. armigera* lineages that should be considered for predicting *H. armigera* range expansion and executing effective pest management.

Regional climate contributed to differentiation of southern and northern *H. armigera* lineages in Central China

We conducted analysis on *H. armigera* resequencing data and identify two differentiated lineages in Central China that are differentiated at Z chromosome. Based on population genetic differentiation analysis and gene flow, we found a pattern of divergent selection with gene flow in Central China that was contributed by local adaptation of northern lineage from the high latitudes. A total of four clock genes (Z chromosome: *per* and *cycle*; autosome chromosome: *cry2* and *daywake*) were under selection in the northern lineage. While in the southern lineage, a Z chromosome inversion locked alleles of two core clock genes (*per* and *cycle*) from *H. a. conferta* that further boost divergence of these two lineages. Introduction divergent alleles of these two clock genes into local population may have disrupted sensing system to temperature and

photoperiod. In addition, apart from southern lineage, linked introgressed alleles of *per* and *cycle*

were only identified in three individuals (Aq4, DF44, and Jz22) from Changjiang River Region

without an Z chromosome inversion. These results suggest a combination of certain genotypes in

clock genes may be needed to maintain normal function of seasonal sensing system.

Studies shows acclimation to high latitudes through diapause was contributed by multiple loci

with minor effects and support a polygenic model (Ragland et al., 2019). In our study, in addition

to four clock genes, we further isolated several genes under selection in northern lineage and may

have contributed to local adaptation of this lineage to temperature and photoperiod of high

latitudes. In Z chromosome, we found *SLC25A25*, a Ca^{2+}-sensitive ATP carrier in the inner

mitochondrial membrane, fluctuates in a circadian manner in mouse skeletal muscle and might be

involve in muscle thermogenesis (Nakao et al., 2017). The other gene on Z chromosome is

IGF2BP2, which is from insulin-like growth factor2 mRNA binding protein family in human and

may associated with impaired insulin secretion (Dai et al., 2020). Phosphorylation of this gene is

critical for coordinating cellular function and nutrient metabolism (Dai et al., 2020). In the

autosomal GWAS analysis by correlation analysis with each SNP and PCA1 identified by Z

chromosome polymorphism, *CCG7766*, a Phosphorylase b kinase regulatory subunit that involved

in carbohydrate metabolic process and glycogen metabolic process was identified as an outlier and

show local adaptation in northern lineage. The other gene that appears on this gene panel was

Acyl-CoA Δ^{11}Z-desaturases (D11DS) that occurred ubiquitously in animal and fungal kingdoms

and plays essential roles in fatty acid metabolism and response to temperature fluctuations by

regulation of cell membrane fluidity (Knipple et al., 1998). In addition, bifunctional

trehalose-6-phosphate synthase/phosphatase (TSTP) that shows selective sweep signal in northern

lineage is important in energy production and other biological processes of invertebrates (Tang et

al., 2018). Additionally, we found an adaptative CPQ gene that encodes carboxypeptidase Q,

which is involved in proteolytic functions and resistant to deficit water status (Gheyas et al., 2021;

Mohamadipoor Saadatabadi et al., 2021), such as diapause in long overwintering periods.

Overall, our studies identified candidate genes in northern lineage of Central China that involved

in circadian clock and could sense seasonal variation of temperature and photoperiod, and in

energy metabolism that may have contributed to local adaptation during stressful and long

overwintering periods. However, although verified by bioinformatic analysis, additional evidence

from transcriptome data and molecular experiment are needed to depict a more robust and

comprehensive understanding of adaptation to high latitude of this pest. We also propose a more

intensive sampling in southern China as well as southeast Asia to investigate the distribution of Z

chromosome inversion and identify genetically distinct lineages that may have adapted to different

temperature and photoperiod (Chen et al., 2013; Virachack et al., 2018).

Adaptation to insecticide leads to widespread local adaptation in different *H. armigera*

lineages

Up to now, *H. armigera* shows the highest number of reported cases of insecticide resistance

worldwide and have developed resistance against most of applied chemicals and several *Bacillus*

thuringiensis (Bt) toxins (Valencia-Montoya et al., 2020). Resistance to insecticides such as

carbamates, organophospates, pyrethroids families, or to Bt has been reported all over the world

for *H. armigera*, while others including spinosyns, indoxacarb, emamectin, and benzoate were

reported regionally (https://irac-online.org/). The differentiation in resistance in regional

populations indicate widespread local adaptation to applied insecticides of different *H. armigera*

lineages.

CYP337B3 shows selection signals in all populations, which is consistent with widespread

distribution of this P450 gene in field that was resistant to fenvalerate (Joußen et al., 2012; Xu et

al., 2014; Han et al., 2015; Rasool et al., 2015). The resistant genotype of *CYP337B3* was also

introduced along with populations of *H. armigera* into Brazil (Durigan et al., 2017) and

introgressed to its sibling species (i.e., *H. zea*) in South America. In addition, we found selection

signals in several lineage-specific genes that indicate local adaptation by development of

insecticide resistance. For example, in Central China, the glutathione S-transferase genes that were

also identified in previous study (Zhang et al., 2022) took part in insecticide and plant secondary

metabolites (Francis et al., 2005; Gawande et al., 2011; Rane et al., 2019). This gene, not

identified in Northwestern China (e.g., Xinjiang) but in Oceania, may indicate introgression

between *H. armigera* and *H. a. conferta*. In addition, one esterase (*CCE001*) that have involved in

fenvalerate resistance was also identified in these two regions (Central China and Oceania), which

support widespread introgression of insecticide-resistant genes.

In Central China, one lineage-specific gene encodes carboxyl esterase (*CCE014a*) that could

mediate metabolic resistance to organophosphates and carbamates (Venkatesan et al., 2022) and

Cytochrome P450 (*CYP12A4*) that could facilitate adaptation to lufenuron (Bogwitz et al., 2005)

were identified, while no lineage-specific insecticide resistant genes were isolated in Northwestern

China. This may be due to shorter insecticide applied history in Northwestern China compared

with that in Central China, differentiation of cropping pattern or geographical

and climatic environment (Li et al., 2010; Lu, 2021). On the contrast, we found many

population-specific genes (over the threshold) that may have contributed to insecticide resistance

in Oceania (i.e., *H. a. conferta* lineage). For example, two cadherins (*CadN* and *Cad89D*) could

have contributed to *Bt* resistance (Fabrick & Wu 2015; Du et al., 2019) of transgenic crops in this

region were outliers. In addition, one gene that encodes choline acetyltransferase (*ChAT*) could be

underlying resistance of this lineage to insecticides that targeted on neurons (Liu et al., 2022). We

noted that *Cad89D* and *ChAT* were with weak peaks in Central China (although not being outliers

base on threshold), while *CadN* sweep was strong in Oceania and not identified in Central China.

The time to start Bt transgeneic crop cultivation in China (since 1997, Huang et al., 2002) and

Australia (since 1996, Fitt, 2002) were similar, and the difference in identification of

lineage-specific Bt genes could be due to sampling bias or due to the difference in development of

Bt resistance, which needs further investigation.

Management of agricultural insect pest required more comprehensive understandings of

local adaptation and introgression

For most insect pests, environmental factors are important for successful reproduction and

succession of local populations (Fält-Nardmann et al., 2016). Among the genetic agents

underlying adaptation of lineages to high latitudes, circadian clock genes could influence insect

behaviors such as locomotion, courtship, and photoperiodism should receive more attention

(Khyati et al., 2017). In addition, due to more hostile environments of high latitudes, successful

range expansion of agricultural pests requires that species are able to use local temperature and/or

day length as cues to coordinate their phenology (Saikkonen et al., 2012), pre-adaptation of insects

to similar environments thus would facilitate rapid colonization to new habitats (Li et al., 2015).

For their importance in viability of insects, techniques such as molecular editing and sterile insect

technique (SIT) that involved clock genes (and other genes related to phenological adaptation)

should benefit integrated pest management to minimize pesticide use and prevent rapid resistance

to insecticides (Khyati et al., 2017).

Addressing the genetic basis of pesticide resistance is the best way to find a sustainable crop plant

protection strategy (Barzman et al., 2015). Our analysis of adaptative genes that are related to

different insecticides not only identify target genes underlying molecular basis of insecticide

resistance, but also revealed that gene flow may have contributed to widespread and rapid

development of insecticide resistance of *H. armigera*. This information would help putting

forward efficient pest management strategy according to the resistant list of local/regional pest

populations. Further analysis to investigate the frequency and distribution of insecticide resistant

alleles (introgressed or locally adaptative) across Southeast Asia and Oceania, as well as other

regions, are needed for a more successful regional integrated pest management strategy.

Conclusion

We combined association analysis and population genetic parameters to isolate adaptative agents

underlying local adaptation of different *H. armigera* lineages. Genes related acclimation and

insecticide resistance were identified despite widespread gene flow. In addition, Z chromosome

inversion could disrupt local adaptation by locking maladaptive alleles of introgressed genes. Our

results show widespread local adaptation of different *H. armigera* populations, which should

further be implemented in integrated pest management and accounted for efficient regional

management of insecticide resistance.

References

Alyokhin A, Mota☐Sanchez D, Baker M, et al. The Red Queen in a potato field: integrated pest management versus chemical dependency in Colorado potato beetle control. Pest Manag. Sci., 2015, 71(3): 343-356.

Arnold M L, Kunte K. Adaptive genetic exchange: a tangled history of admixture and evolutionary innovation. Trends Ecol. Evol., 2017, 32(8): 601-611.

Minh BQ, H.A. Schmidt, O. Chernomor, D. Schrempf, M.D. Woodhams, A. von Haeseler, R. Lanfear (2020) IQ-TREE 2: New models and efficient methods for phylogenetic inference in the genomic era. Mol. Biol. Evol., 37:1530-1534.

Barzman, M., Bàrberi, P., Birch, A.N.E. et al. Eight principles of integrated pest management. Agron. Sustain. Dev. 2015, 35: 1199–1215.

Bogwitz MR, Chung H, Magoc L, Rigby S, Wong W, O'Keefe M, McKenzie JA, Batterham P, Daborn PJ. Cyp12a4 confers lufenuron resistance in a natural population of Drosophila melanogaster. Proc Natl Acad Sci U S A. 2005, 102(36):12807-12.

Chang C C, Chow C C, Tellier L C A M, et al. Second-generation PLINK: rising to the challenge of larger and richer datasets. Gigascience, 2015, 4(1): s13742-015-0047-8.

Crispo E, Moore J S, Lee☐Yaw J A, et al. Broken barriers: Human☐induced changes to gene flow and introgression in animals: An examination of the ways in which humans increase genetic exchange among populations and species and the consequences for biodiversity. BioEssays, 2011, 33(7): 508-518.

Dai, N. The diverse functions of IMP2/IGF2BP2 in metabolism. Trends Endocrinol. Metab., 2020, 31(9): 670-679.

Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., Handsaker, R.E., Lunter, G., Marth, G.T., Sherry, S.T. and McVean, G., 2011. The variant call format and VCFtools. Bioinformatics, 27(15): 2156-2158.

Danks, H. V. (2000) Measuring and reporting life-cycle duration in insects and arachnids. Eur. J. Entomol. 97: 285–303.

DePristo, M.A., Banks, E., Poplin, R., Garimella, K.V., Maguire, J.R., Hartl, C., Philippakis, A.A., Del Angel, G., Rivas, M.A., Hanna, M. and McKenna, A., 2011. A framework for variation discovery

and genotyping using next-generation DNA sequencing data. Nat. Genet., 43(5): 491-498.

Downes S, Kriticos D, Parry H, et al. A perspective on management of *Helicoverpa armigera*: transgenic Bt cotton, IPM, and landscapes. Pest Manag. Sci., 2017, 73(3): 485-492.

Du, L., Chen, G., Han, L. et al. Cadherin CsCad plays differential functional roles in Cry1Ab and Cry1C intoxication in *Chilo suppressalis*. Sci. Rep., 2019, 9: 8507.

Durigan MR, Corrêa AS, Pereira RM, Leite NA, Amado D, de Sousa DR, Omoto C. High frequency of CYP337B3 gene associated with control failures of *Helicoverpa armigera* with pyrethroid insecticides in Brazil. Pestic Biochem Physiol. 2017, 143:73-80.

Fabrick, J.A., Wu, Y. and Wu, Y., 2015. Roles of insect midgut cadherin in Bt intoxication and resistance. *Bt resistance: characterization and strategies for GM crops producing Bacillus thuringiensis toxins*, 2015 pp.69-86.

Fält-Nardmann, J., Klemola, T., Roth, M., Ruohomäki, K., & Saikkonen, K. (2016). Northern geometrid forest pests (Lepidoptera: Geometridae) hatch at lower temperatures than their southern conspecifics: Implications of climate change. Eur. J. Entomol., 2016: 113.

Fitt G. Deployment and impact of transgenic Bt cotton in Australia. Econ. Environ. Impacts Agrobiotechnol., 2003:141-161.

Francis F, Vanhaelen N, Haubruge E. Glutathione S-transferases in the adaptation to plant secondary metabolites in the *Myzus persicae* aphid. Arch. Insect Biochem. Physiol., 2005, 58(3):166-74.

Garud N R, Messer P W, Buzbas E O, et al. Recent selective sweeps in North American Drosophila melanogaster show signatures of soft sweeps. PLoS Genet., 2015, 11(2): e1005004.

Gawande ND, Subashini S, Murugan M, Subbarayalu M. Molecular screening of insecticides with sigma glutathione S-transferases (GST) in cotton aphid Aphis gossypii using docking. Bioinformation. 2014: 10(11):679-83.

Gheyas A A, Vallejo-Trujillo A, Kebede A, et al. Integrated environmental and genomic analysis reveals the drivers of local adaptation in African indigenous chickens. Mol. Biol. Evol., 2021, 38(10): 4268-4285.

Han Y, Yu W, Zhang W, Yang Y, Walsh T, Oakeshott JG, Wu Y. Variation in P450-mediated fenvalerate resistance levels is not correlated with *CYP337B3* genotype in Chinese populations of *Helicoverpa armigera*. 2015, 121:129-35.

Hahn, D.A. and Denlinger, D.L., 2011. Energetics of insect diapause. Annu. Rev. Entomol., *56*,

pp.103-121.

Harris, AM, Garud NR, DeGiorgio M. Detection and classification of hard and soft sweeps from

unphased genotypes by multilocus genotype identity. Genetics, 2018: 1429-1452.

Hasebe M, Shiga S. Clock gene-dependent glutamate dynamics in the bean bug brain regulate

photoperiodic reproduction. PLoS Biol., 2022, 20(9): e3001734.

Hoang D.T., O. Chernomor, A. von Haeseler, B.Q. Minh, L.S. Vinh. UFBoot2: Improving the ultrafast

bootstrap approximation. Mol. Biol. Evol., 2018, 35:518–522.

Hoikkala, A. and Poikela, N., 2022. Adaptation and ecological speciation in seasonally varying

environments at high latitudes: *Drosophila virilis* group. Fly, 16(1): 85-104.

Huang JK, RF Hu, CH Fan, CE Pray, S Rozelle, Bt cotton benefits, costs, and impacts in

China. AgBioForum 2002, 5: 153–166.

Jin, MH, Henry L. North, Yan Peng, Hangwei Liu, Ruiqing Pan, Yan Zhou, Weigang Zheng et al.

Adaptive evolution to the natural and anthropogenic environment in a global invasive crop pest,

the cotton bollworm. The Innovation (2023), 4(4): 100454.

Jones, C. M., Papanicolaou, A., Mironidis, G. K., Vontas, J., Yang, Y., Lim, K. S., Oakeshott, J. G.,

Bass, C., & Chapman, J. W. Genome wide transcriptional signatures of migratory flight activity in

a globally invasive insect pest. Mol. Ecol., 2015, 24: 4901– 4911.

Joußen N, Agnolet S, Lorenz S, et al. Resistance of Australian *Helicoverpa armigera* to fenvalerate is

due to the chimeric P450 enzyme CYP337B3. Proc. Natl. Acad. Sci. U.S.A., 2012, 109(38):

15206-15211.

Kassambara, A. and Kassambara, M.A. Package 'ggpubr'. R package version 0.1, 2020, 6(0).

Khyati, Malik I, Seth R K. Insect clocks: implication in an effective pest management. Biol. Rhythm

Res., 2017, 48(5): 777-788.

Kirkpatrick M, Barton N. Chromosome inversions, local adaptation and speciation. Genetics, 2006,

173(1): 419-434.

Klai, K., Chenais, B., Zidi, M., Djebbi, S., Caruso, A., Denis, F., Confais, J., Badawi, M., Casse, N. and

Mezghani Khemakhem, M., 2020. Screening of *Helicoverpa armigera* mobilome revealed

transposable element insertions in insecticide resistance genes. Insects, 2020, 11(12): 879.

Knipple D C, Rosenfield C L, Miller S J, et al. Cloning and functional expression of a cDNA encoding

a pheromone gland-specific acyl-CoA Δ11-desaturase of the cabbage looper moth, *Trichoplusia ni*.

Proc. Natl. Acad. Sci. U.S.A., 1998, 95(26): 15287-15292.

Kozak G M, Wadsworth C B, Kahne S C, et al. Genomic basis of circannual rhythm in the European

corn borer moth. Curr. Biol., 2019, 29(20): 3501-3509. e5.

Li S, Cadotte M W, Meiners S J, et al. The effects of phylogenetic relatedness on invasion success and

impact: deconstructing Darwin's naturalisation conundrum. Ecol. Lett., 2015, 18(12): 1285-1292.

Li, G., Feng, H., Gao, Y., Wyckhuys, K. A., & Wu, K. Frequency of Bt resistance alleles in *Helicoverpa*

armigera in the Xinjiang cotton-planting region of China. Environ. Entomol., 2010, 39(5),

1698-1704.

Li, H., 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. arXiv

preprint arXiv:1303.3997.

Liu J, Gao S, Wei L, et al. Choline acetyltransferase and vesicular acetylcholine transporter are required

for metamorphosis, reproduction, and insecticide susceptibility in *Tribolium castaneum*. Gene,

2022, 842: 146794.

Lu YH. Ever-evolving advances in the researches of cotton insect pest management in China. Acta

Phytophylacica Sin., 2021, 48(5): 937-939.

Malinsky, M, Matschiner, M, Svardal, H. Dsuite - Fast D-statistics and related admixture evidence

from VCF files. Mol Ecol Resour. 2021; 21: 584– 595.DeGiorgio M, Christian D. Huber, Melissa

J. Hubisz, Ines Hellmann, Rasmus Nielsen, Sweep Finder 2: increased sensitivity, robustness and

flexibility. Bioinformatics, 2016, 32(12): 1895-1897.

Mohamadipoor Saadatabadi L, Mohammadabadi M, Amiri Ghanatsaman Z, et al. Signature selection

analysis reveals candidate genes associated with production traits in Iranian sheep breeds. BMC

Vet. Res., 2021, 17(1): 1-9.

Nakao R, Shimba S, Oishi K. Ketogenic diet induces expression of the muscle circadian gene Slc25a25

via neural pathway that might be involved in muscle thermogenesis. Sci. Rep., 2017, 7(1): 1-11.

Nibouche, S., Buès, R., Toubon, J.- F., & Poitout, S. Allozyme polymorphism in the cotton bollworm

Helicoverpa armigera (Lepidoptera: Noctuidae): Comparison of African and European

populations. Heredity, 1998, 80: 438– 445.

Ortiz-Barrientos D, Engelstädter J, Rieseberg L H. Recombination rate evolution and the origin of

species.Trends Ecol. Evol., 2016, 31(3): 226-236.

Paaby AB, Blacket MJ, Hoffmann AA, Schmidt PS: Identification of a candidate adaptive

polymorphism for Drosophila life history by parallel independent clines on two continents. Mol. Ecol., 2010, 19:760-774.

Paini D R, Sheppard A W, Cook D C, et al. Global threat to agriculture from invasive species. Proc. Natl. Acad. Sci. U.S.A., 2016, 113(27): 7575-7579.

Pélissié B, Crossley M S, Cohen Z P, et al. Rapid evolution in insect pests: the importance of space and time in population genomics studies. Curr. Opin. Insect. Sci., 2018, 26: 8-16.

Peng, Yan, Minghui Jin, Zhimin Li, Hongran Li, Lei Zhang, Songmiao Yu, Zhuting Zhang et al. Population-genomics provide insights into the evolution and adaptation of the Asia corn borer. Mol. Biol. Evol., 2023, 40(5): msad112.

Pinho, C. and Hey, J. Divergence with gene flow: models and data. Annu. Rev. Ecol. Evol. S., 2010, 41: 215-230.

Pratissoli, D., Lima, V. L., Pirovani, V. D., & Lima, W. L. Occurrence of Helicoverpa armigera (Lepidoptera: Noctuidae) on tomato in the Espírito Santo state. Hortic. Bras., 2015, 33: 101– 105.

Ragland G J, Armbruster P A, Meuti M E. Evolutionary and functional genetics of insect diapause: a call for greater integration. Curr. Opin. Insect. Sci., 2019, 36: 74-81.

Rane, R.V., Ghodke, A.B., Hoffmann, A.A., Edwards, O.R., Walsh, T.K. and Oakeshott, J.G., 2019. Detoxifying enzyme complements and host use phenotypes in 160 insect species. Curr. Opin. Insect. Sci., 2019 31: 131-138.

Rasool A, Joußen N, Lorenz S, Ellinger R, Schneider B, Khan SA, Ashfaq M, Heckel DG. An independent occurrence of the chimeric P450 enzyme CYP337B3 of *Helicoverpa armigera* confers cypermethrin resistance in Pakistan. Insect Biochem. Mol. Biol. 2014, 53:54-65.

Renou M. Is the evolution of insect odorscapes under anthropic pressures a risk for herbivorous insect invasions?. Curr. Opin. Insect. Sci., 2022: 100926.

Riaz S, Johnson J B, Ahmad M, et al. A review on biological interactions and management of the cotton bollworm, *Helicoverpa armigera* (Lepidoptera: Noctuidae). J. Appl. Entomol., 2021, 145(6): 467-498.

Nowling RJ, F Fallas-Moya, et al. Fast, low-memory detection and localization of large, polymorphic inversions from SNPs. PeerJ, 2022: e12831.

Roderick GK and Navajas M.Genes in new environments: genetics and evolution in biological control. Nat. Rev. Genet., 2003, 4: 889-899.

620   Roesti M, Gilbert KJ, Samuk K. Chromosomal inversions can limit adaptation to new environments.

621   Mol. Ecol., 2022, 31(17):4435-4439.

622   Saikkonen K, Taulavuori K, Hyvönen T, et al. Climate change-driven species' range shifts filtered by

623   photoperiodism. Nat. Clim. Chang., 2012, 2(4): 239-242.

624   South, A. rworldmap: A new R package for mapping global data. R J., 2011 3(1).

625   South, A. rworldxtra: Country boundaries at high resolution. R package version 1.01. 2012.

626   Szendrei Z, Grafius E, Byrne A, et al. Resistance to neonicotinoid insecticides in field populations of

627   the Colorado potato beetle (Coleoptera: Chrysomelidae). Pest Manag. Sci., 2012, 68(6): 941-946.

628   Tang B, Wang S, Wang S G, et al. Invertebrate trehalose-6-phosphate synthase gene: genetic

629   architecture, biochemistry, physiological function, and potential applications. Front. Physiol.,

630   2018, 9: 30.

631   Tigano, Anna, and Vicki L. Friesen. Genomics of local adaptation with gene flow. Mol. Ecol.,

632   2016, 25(10): 2144-2164.

633   Toepfer, S., Li, H., Pak, S.G., Son, K.M., Ryang, Y.S., Kang, S.I., Han, R. and Holmes, K., 2014. Soil

634   insect pests of cold temperate zones of East Asia, including DPR Korea: A review. J. Pest Sci., 87:

635   567-595.

636   Valencia-Montoya W A, Elfekih S, North H L, et al. Adaptive introgression across semipermeable

637   species boundaries between local *Helicoverpa zea* and invasive *Helicoverpa armigera* moths. Mol.

638   Biol. Evol., 2020, 37(9): 2568-2583.Venkatesan, T., B. R. Chethan, and M. Mani. Insecticide

639   resistance and its management in the insect pests of horticultural crops. Trends Hortic.

640   Entomol., 2022: 455-490.

641   Virachack, Hatsada, et al. Temperature effects on the development of Vietnamese cotton bollworm

642   Helicoverpa armigera (Hübner). J. Fac. Agr., Kyushu Univ., 2018, 63 (2): 303-309.

643   Wickham, H. Reshaping data with the reshape package. J. Stat. Softw., 2007, 21:1-20.

644   Wickham, H., 2011. ggplot2. Wiley Interdiscip. Rev. Comput. Stat., 3(2): 180-185.

645   Williams KD, Busto M, Suster ML, So AKC, Ben-Shahar Y, Leevers SJ, Sokolowski MB: Natural

646   variation in Drosophila melanogaster diapause due to the insulin-regulated PI3- kinase. Proc. Natl.

647   Acad. Sci. U. S. A., 2006, 103:15911-15915.

648   Wu K, Mu W, Liang G, et al. Regional reversion of insecticide resistance in *Helicoverpa armigera*

649   (Lepidoptera: Noctuidae) is associated with the use of Bt cotton in northern China. Pest Manag.

Sci., 2005, 61(5): 491-498.

Wu KM, Guo YY. Genotype differentiation and regional migratory regularity of *Helicoverpa armigera*

in China. Plant Prot., 2007, 33(5):5-11.

Wu KM, Guo YY. Investigations on the migration and diapause in *Helicoverpa armigera* -Diapause

responses of different geographical populations of cotton bollworm *Helicoverpa armigera* Hübner.

Zhongguo Nong Ye Ke Xue, 1997, 30 (3): 1-6.

Wu Kongming, Guo Yuyuan, Wei Jianfu, Sun Fuzai, On the cold hardiness of cotton bollworm

Helicoverpa armigera Hübner. Sheng Tai Xue Bao, 1997, 17(3): 298-302.

Xu L, Li D, Qin J, Zhao W, Qiu L. Over-expression of multiple cytochrome P450 genes in

fenvalerate-resistant field strains of *Helicoverpa armigera* from north of China. Pestic. Biochem.

Physiol., 2016, 132:53-8.

Yang Y, Li Y, Wu Y. Current status of insecticide resistance in *Helicoverpa armigera* after 15 years of

Bt cotton planting in China. J. Econ. Entomol., 2013, 106(1): 375-381.

Yin, L. CMplot: circle manhattan plot. R package version, 2020, 3(2).

Chen YS and Tu XY, Geographic variation in developmental duration of the cotton bollworm ,

Helicoverpa armigera Hübner (Lepidoptera: Noctuidae). J. Environ. Entomol., 2016 ,38(4):771 -

778.

Chen, Y.S., Chen, C., He, H.M., Xia, Q.W. and Xue, F.S. Geographic variation in diapause induction

and termination of the cotton bollworm, *Helicoverpa armigera* Hübner (Lepidoptera:

Noctuidae). J. Insect Physiol., 2013, *59*(9):855-862.

Zhang C, Rabiee M, Sayyari E, et al. ASTRAL-III: polynomial time species tree reconstruction from

partially resolved gene trees. BMC Bioinform., 2018, 19(6): 15-30.

Zhang H, Tian W, Zhao J, et al. Diverse genetic basis of field-evolved resistance to Bt cotton in cotton

bollworm from China. Proc. Natl. Acad. Sci. U.S.A., 2012, 109(26): 10275-10280.

Zhang J, Zhang F, Tay W T, et al. Population genomics provides insights into lineage divergence and

local adaptation within the cotton bollworm. Mol. Ecol. Resour., 2022, 22(5): 1875-1891.

Figure Legends

Figure 1. Z chromosome divergence of *H. armigera* despite widespread gene flow. (A) Maximum

likelihood phylogeny based on concatenated SNPs from Z chromosome. Central China southern

lineage (CC-S, with arrows) and Central China northern lineage (CC-N) were labeled with blue

and cyan, respectively. Similar topology was reconstructed based on Astral (Supplementary Figure

1) (B) Principal component analysis based on autosomal SNPs. Note the CC-S and CC-N samples

were clustered together. (C) Genetic differentiation of *H. armigera* males among different lineages.

Two major topologies were illustrated, with topology one (①) consistent with the species tree,

while topology two (②) gathering *H. a. conferta* (AUS) and CC-S individuals. "②※" indicates a

window tree similar to topology two, but with CC-S lineage including more individuals from

CC-N lineage. See supplementary Figure 2 for more information. The lower panel shows

significant correlation between genetic differentiations of CC-S vs. CC-N and AUS vs. CC-N in Z

chromosome. (D) Widespread gene flow between *H. a. conferta* from Oceania and *H. armigera*

populations in Central China.

Figure 2. Detection and characterization of putative Z chromosome inversion in *Helicoverpa*

armigera. (A) Manhattan plot of GWAS analysis between Z chromosome polymorphism and

principal component 1 (PC1). The break point of the inversion was set at the site with $-\log_{10}(p)$

value more than 25. The central panel is π ratio (π_{CC-N}/π_{CC-S}) calculated in 5kb window, with

labeled genes identified in 1% lowest π ratio regions. The lower panel is *fdM* distribution of CC-N

vs. AUS and CC-S vs. AUS calculated with 100-SNP window and 25-SNP step. (B) Genotype

matrix of highly differentiated SNPs ($-\log_{10}(p) > 15$) in the putative chromosome inversion. A total

of 3,237 SNPs were used for analysis. (C) Linkage disequilibrium analysis in different population

and genomic regions; Inversion indicates the putative inversion region, and other regions are the

rest of Z chromosome excluding the inversion. AUS are male individuals sampled from *H. a.*

conferta. (D) Geographical distribution of two haplogroups (ancestral and inversion) in Central

China. Only CC-S and CC-N individuals included for analysis as inversion identified in Central

China (see Supplementary Figure 4 for the genotype matrix of all individuals).

Figure 3. Characterization of *per* and *ARNTL* (*cycle*) in the inversion region. (A) Genotype matrix

of polymorphism at CDS regions of *per* and *cycle* genes. (B) Population genetic statistics of *per*

and *cycle*. The red dashed line in each panel was the mean value of each statistic calculated by Z

chromosomal genetic polymorphism. (C) Protein structures of two differentiated haplotypes in *per*

and *cycle*. Cyan structure was constructed from the CC-N dominant haplotype, and additional blue

structure that shows the difference was constructed with the CC-N dominant one. (D) Patterns of

LD blocks between and within genes (*per* and *cycle*). Red and purple rectangles denote CDS and

UTR, respectively. Only male individuals were used in Z chromosome-related analysis.

Figure 4. Autosomal genome-wide association tests show differentiation between two

Helicoverpa armigera lineages. (A) Manhattan plot of genome-wide polymorphisms. Peak regions

with more than one highly differentiated SNPs (*p* value lower than 5.0×10^{-8}) were used for gene

identification. The lower panel is genetic differentiation (Fst) between CC-S and CC-N lineages.

(B) Characterization of six highly differentiated regions. π ratio indicates π_{CC-N}/π_{CC-S}, and d_{xy} is

nucleotide divergence between CC-S and CC-N.

Figure 5. Widespread local adaptation in *Helicoverpa armigera* lineages. Selective sweep signals

in different populations from Central China (CC), Northwest China (NC), and Australia (*H. a.*

conferta, AUS) based on G12 (A) and CLR (B) statistics. Horizontal dash lines denote threshold

values of G12/CLR over 99.95% autosomal windows in each lineage.

731
732

737
738

739 **Supplementary Figure 1.** Phylogenetic tree of Z chromosome based on
740 ASTRAL-III.
741

742
743

Supplementary Figure 2. Genetic structure of all *H. armigera* individuals based on

autosomal polymorphisms.

Supplementary Figure 3. Tree topologies of five regions in Z chromosome.

Supplementary Figure 4. Genome-wide calculation of D value between two *H.*

armigera lineages (CC-N and CC-S) in Central China with Oceanian *H. a. conferta*

lineage.

Supplementary Figure 5. Principal component analysis of *H. armigera* individuals

sampled from Central China based on Z chromosomal polymorphisms.

762
763

764   **Supplementary Figure 6.** Genotype matrix of all sampled individuals in the putative

765   inversion region of Z chromosome (6.7 - 12.4 Mb).

768