# Curcumin and turmeric extract inhibited SARS-CoV-2 pseudovirus cell entry and Spike mediated cell fusion

Authors: Endah Puji Septisetyani[1]*, Dinda Lestari[1,2], Komang Alit Paramitasari[1], Pekik Wiji Prasetyaningrum[1], Ria Fajarwati Kastian[1], Adi Santoso[1], Kartini Eriani[2]

Authors Affiliation:

[1]Mammalian Cell Engineering Research Group, Research Center for Genetic Engineering, National Research and Innovation Agency (BRIN), Jl. Raya Bogor KM. 46, Cibinong, Bogor, West Java, 16911, Indonesia

[2] Universitas Syiah Kuala, Faculty of Mathematics and Natural Sciences, Department of Biology, Banda Aceh, Indonesia

Corresponding Author:

Name   : Dr. Endah Puji Septisetyani

Address: Research Center for Genetic Engineering, National Research and Innovation Agency, Gedung Admin OR Hayati dan Lingkungan, BRIN, Jl. Raya Bogor KM 46, Cibinong, Bogor, 16911, Indonesia

email: enda041@brin.go.id

**AUTHORS' CONTRIBUTION**

Conceptualization: Septisetyani EP. Data curation: Lestari D, Prasetyaningrum PW, Paramitasari KA, Kastian RF, Septisetyani EP. Writing - original draft: Septisetyani EP,

Lestari D, Paramitasari KA. Writing - review & editing: Septisetyani EP, Prasetyaningrum PW, Santoso A, Eriani K.

## ACKNOWLEDGMENTS

We are grateful for the financial support from LPDP/BRIN and Research Organization for Life Sciences and Environment, BRIN and the proofreading service by BRIN.

## FINANCIAL SUPPORT

We are grateful for the financial support from Educational Fund Management Institution/National Research and Innovation Agency (LPDP/BRIN) (grant no. 102/FI/P-KCOVID-19.2B3/IX/2020 and RIIM No. KEP-5/LPDP/LPDP.4/2022) and Research Organization for Life Sciences and Environment, BRIN (Research Program (DIPA Rumah Program) 2022).

## CONFLICT OF INTEREST

The authors declared no potential conflict of interests.

## ETHICAL APPROVAL

Not applicable.

## DATA AVAILABILITY

All data and materials are provided within the research article.

**ORCID IDs**

Endah Puji Septisetyani ORCID 0000-0002-6764-9686

Dinda Lestari ORCID 0009-0001-9293-4139

Komang Alit Paramitasari ORCID 0000-0002-4054-3398

Pekik Wiji Prasetyaningrum ORCID 0000-0001-6075-9850

Ria Fajarwati Kastian ORCID 0000-0003-3199-547X

Adi Santoso ORCID 0000-0002-7105-8483

Kartini Eriani ORCID 0000-0002-9063-5944

**Curcumin and turmeric extract inhibited SARS-CoV-2 pseudovirus cell entry and**

**Spike mediated cell fusion**

**ABSTRACT**

Turmeric extract (TE) with curcumin as its main active ingredient has been studied as a

potential COVID-19 therapeutic. Curcumin has been studied in silico and in vitro against a

naive SARS-CoV-2 virus, yet little is known about TE's impact on SARS-CoV-2 infection.

Moreover, no study reveals the potentials of both curcumin and TE on the inhibition of SARS-

CoV-2 cell-to-cell transmission. Here, we investigated the effects of both curcumin and TE on

the inhibition of SARS-CoV-2 entry and cell-to-cell transmission using pseudovirus (PSV) and

syncytia models. We performed PSV entry assay in 293T or 293 cells expressing hACE2. The

cells were pretreated with curcumin or TE, then treated with PSV with or without the test

samples. Next, we carried out syncytia assay by co-transfecting 293T cells with plasmids

encoding Spike, hACE2, and TMPRSS2 to be treated with the test samples. The results showed

that in PSV entry assay on 293T/hACE/TMPRSS2 cells, both curcumin and TE inhibited PSV

entry at concentrations of 1 µM and 10 µM for curcumin and 1 µg/ml and 10 µg/ml for TE.

Moreover, both curcumin and TE also reduced syncytia formation compared to control cells.

Based on our study, both TE and curcumin are potential inhibitors of SARS-CoV-2 infection

at entry points, either by direct or indirect infection models.

**Keywords:** COVID-19, curcumin, pseudovirus, SARS-CoV-2, syncytia, turmeric extract.

**GRAPHICAL ABSTRACT:**

## 1. INTRODUCTION

SARS-CoV-2 can infect the target cells by direct viral infection and cell-to-cell transmission (Jackson et al., 2021; Rajah et al., 2021). The first mode of viral infection has been widely explored by studying cell and organ tropisms. SARS-CoV-2 infection toward various cell types has been performed, which includes studies using pulmonary cancer cells Calu-3, kidney cells 293FT, liver cells Huh7, colon cancer cells CaCO2, and Vero (Hoffmann et al., 2020; Katopodis et al., 2022; Saccon et al., 2021). In addition, organ tropisms have been studied using post-mortem tissue samples from COVID-19 patients and organoid models (Peng et al., 2022). Based on these studies, SARS-CoV-2 tropisms are strongly related to hACE2 expression as its target receptor and TMPRSS2 protease, which enhances the viral entry process. SARS-CoV-2 infection involves the interaction of Spike glycoprotein with hACE2, with or without Spike priming by TMPRSS2, to facilitate cell entry through endocytosis or membrane fusion mechanisms (Hirano et al., 2020; Jackson et al., 2021).

In addition, SARS-CoV-2 can transmit through a cell-to-cell mechanism, which induces cell-to-cell fusion and syncytia formation (Zheng et al., 2021). Syncytia has been found in the lungs of patients infected with SARS-CoV-2 (Lin et al., 2021). SARS-CoV-2 infection induces spike glycoprotein expression. Eventually, as a transmembrane protein, Spike will be transported to the plasma membrane with the receptor binding domain (RBD) at the extracellular position, ready for receptor binding. Similar to the virally integrated Spike as the outer protein, cellular Spike protein can also bind to the adjacent cells' hACE2 receptor, which further induces cell-to-cell fusion and finally forms a multinucleated giant cell or syncytium (Zheng et al., 2021; Lin et al., 2021). Syncytia formation has been implicated with the severity of COVID-19 prognosis by several mechanisms, including immune cell phagocytosis, antibody evasion, and induction of syncytial cell death (Rajah et al., 2022).

Due to the availability of registered COVID-19 drugs and the continuous emergence of daily COVID-19 cases, alternative treatments that can be easier to obtain for preventing and treating future COVID-19 cases are essential to be developed. Turmeric (Curcuma longa L.) is a WHO-selected medicinal plant that grows throughout tropical areas (WHO, 1999). Turmeric rhizome has been processed into turmeric decoction and turmeric extract, and some has been developed as a standardized medicinal extract. Turmeric extract contains curcuminoid active compounds, with curcumin as its main component (Sharifi-Rad et al., 2020). Clinical studies have demonstrated the effect of nano-encapsulated curcumin in reducing clinical manifestations of COVID-19, such as fever, cough, and dyspnea, compared to the control group (Tahmasebi et al., 2021; Valizadeh et al., 2020).

In silico analysis showed that curcumin has a high affinity for binding to the spike glycoprotein through the formation of six hydrogen bonds (Maurya et al., 2020), thus, curcumin has the potential to prevent the binding of the viral spike protein to the ACE2 receptor and inhibit the initiation of the host cell infection process. Moreover, curcumin forms four hydrophobic interactions via hydrogen bonds with TMPRSS2, which may inhibit cellular entry through the cell fusion mechanism (Motohashi et al., 2020).

A plaque assay study in Vero cells has shown that curcumin inhibits SARS-CoV-2 infection in pre- and post-treatment of the D614 strain and Delta variant (Marin-Palma et al., 2021). A study by Bormann *et al.* (2021) shows that turmeric root juice and curcumin showed inhibition of SARS-CoV-2 infection in Vero and Calu-3 cells determined by plaque assay and in-cell ELISA to detect the signal of SARS-CoV-2 nucleocapsid protein (N). Nonetheless, the effect of turmeric extract (TE) on SARS-CoV-2 infection is less studied than curcumin. Moreover, there is no report about the effects of both curcumin and TE on SARS-CoV-2 cell-to-cell transmission. In the present study, we examined the effects of TE and curcumin on

inhibiting SARS-CoV-2 infection using the pseudovirus (PSV) and syncytia models for

targeting viral entry points and cell-to-cell transmission.

**2. MATERIALS AND METHODS**

**2.1. Cell culture and reagents**

The 293T (ECACC 12022001), CHO-K1 (ECACC 85051005), and 293 (ECACC 85120602)

cell lines are collections of the Research Center for Genetic Engineering, National Research

and Innovation Agency (BRIN, Indonesia). 293T cells were cultured in High-Glucose

Dulbecco's-modified Eagle's medium (Gibco, Billings USA) supplemented with 10% (v/v) of

heat-inactivated fetal bovine serum (Sigma-Aldrich, St. Louis USA) and antibiotics (100 μg/ml

streptomycin and 100 U/ml penicillin). CHO-K1 cells were cultured in F-12 medium (Sigma-

Aldrich, St. Louis USA) supplemented with 10% (v/v) of heat-inactivated fetal bovine serum

and antibiotics (100 μg/ml streptomycin and 100 U/ml penicillin). 293 cells were cultured in

MEM medium (Sigma-Aldrich, St. Louis USA) supplemented with 10% (v/v) of heat-

inactivated fetal bovine serum, 1% NEAA (Gibco, Billings USA), and antibiotics (100 μg/ml

streptomycin and 100 U/ml penicillin). Cells were grown inside a 37 °C tissue culture incubator

maintained at 5% $CO_2$. The pcDNA3.1-SARS2-Spike (#145032), pcDNA3.1-hACE

(#145033), and TMPRSS2 (#53887) plasmids were obtained from Addgene. Cells were

transfected with expression vectors using polyethylenimine (PEI-Max, Polysiences).

**2.2. Turmeric extract and curcumin preparation**

Turmeric extract (TE) was purchased in the form of soft capsules from PT. Industri Jamu dan

Farmasi Sido Muncul Tbk (Semarang, Indonesia). Each capsule was standardised to contain

350 mg concentrated liquid extract, equal to 100 mg of curcuminoids. The TE was dissolved

in DMSO at 50 mg/mL, and the stock was stored at -20 °C. The working solution was freshly

prepared by serially diluting the stock in culture media. Curcumin powder was obtained from

Sigma (C1386), Sigma-Aldrich, St. Louis, USA), reconstituted in DMSO at 50 mM, and

maintained as the stock at -20 °C. For every treatment involving curcumin, the working

solution was made at 10 and 100 µM concentrations by diluting the stock with complete culture

media with a maximum final concentration of DMSO less than 1%.

## 2.3. MTT assay

MTT cell viability assay was carried out in CHO-K1 and 293T cells. One day prior to the MTT

assay, $8 \times 10^4$ 293T or $7 \times 10^4$ CHO-K1 cells per ml were seeded onto a 96-well plate. The next

day, the cells were treated and incubated with culture media containing various concentrations

of TE or curcumin and incubated for about 24 h. A set of wells containing cells without any

treatment and another set of wells were prepared with medium only for background subtraction

(blank). Following incubation, the culture media containing TE or curcumin were discarded

and cells were washed with 1X PBS. The cells were then incubated for 2 hours in the presence

of 0.5 mg/mL MTT solution (3-(4,-5-dimethylthiazo-2-yl)-2,5-diphenyltetrazolium bromide).

At the end of incubation, MTT reagents were discarded and 100 µl of DMSO were added to

each well and agitated at 100 rpm for 10 minutes to ensure complete formazan crystals

solubilisation. Finally, the absorbance of each well was recorded at 570 nm wavelength. Cell

viability was calculated according to the formula: $(OD_{treated}-OD_{blank})/(OD_{untreated}-OD_{blank})$ X

100%

## 2.4. Western blot

Cell lysates were prepared using ice-cold RIPA buffer (Abcam, USA) with the addition of a

protease inhibitor cocktail, then total protein concentration was determined by BCA assay

(Thermo Scientific). About 10-40 µg protein was subjected for SDS-PAGE and then the

resolved protein was transferred onto an activated PVDF membrane. The membrane was then

incubated in blocking buffer (5% skim milk in TBS/0.05% tween-20) followed by blotting with

primary antibodies (anti-SARS-CoV-2 spike (Abcam, USA) 1:2,000 or anti-$\alpha$-actin (Sigma,

USA) 1:4,000) for 2 h at room temperature or overnight at 4°C. After washing with TBS/T, the

membrane was incubated in secondary antibodies (ALP-conjugated antibodies (Abcam or

Invitrogen), 1:4,000 or IR-Dye conjugated antibodies (LI-COR), 1:10,000) in blocking buffer

for about 2 h at room temperature or overnight at 4°C. Western blot signal was detected by

incubating the membrane with NBT-BCIP substrate (Thermo Scientific) or observed by a LI-

COR Odyssey CLx instrument.

## 2.5. Immunofluorescence staining

The 293T cells were seeded at a density of $3 \times 10^4$ cells/well on gelatin-coated cover glass placed

inside a 24-well plate. After about 3 h incubation, the cells were transfected overnight with the

pcDNA3.1-hACE2 expression vector. Transfected cells were washed with 1X PBS and fixed

with 4% paraformaldehyde for 10 min. Upon fixation, cells were permeabilised with 0.2 %

Triton-X, then incubated for 30 min at RT with blocking buffer (1% Bovine Serum Albumin

in 1X PBS) and further incubated for at least 1 hour at RT with rabbit anti-hACE2 antibody

(SAB 3500978, Sigma-Aldrich) or rabbit anti-TMPRSS2 antibody (BS-6285R, Bioss) diluted

in blocking buffer (1:250). Following primary antibody staining, cells were washed three times

with 1X PBS and incubated with goat anti-rabbit Alexa Fluor™ 594 or 488 secondary antibody

(1:1,000) for 1 hour at RT. The secondary antibody was then washed with 1X PBS three times

and the nuclei were stained with DAPI (4′,6-diamidino-2-phenylindole)-containing mountant

(Abcam Ab104139). Samples were imaged with a motorized fluorescence microscope

(Olympus IX83, Tokyo, Japan).

## 2.6. Preparation of SARS-CoV-2 pseudovirus

Pseudotyping was performed in 293T cells previously transfected with a plasmid encoding SARS-CoV-2 Spike (293T/Spike). Briefly, 293T/Spike cells were incubated with pseudotyped G*ΔG-GFP rVSV (Kerafast EH1024-PM) at MOI ~3 for 1 h. Then, the medium was replaced with a fresh medium containing anti-VSV-G antibody 1:2,000 to neutralize excess of G*ΔG-GFP rVSV and left overnight in the $CO_2$ incubator. A conditioned medium (CM) containing pseudotyped Spike*ΔG-GFP rVSV (SARS-CoV-2 pseudovirus) was collected and spun the next day to remove cell debris. The supernatant was aliquoted and stored at -80°C prior to being used for PSV entry assay.

## 2.7. Pseudovirus entry assay

PSV entry assay study was performed in the target cells of 293T transiently overexpressing hACE2 and TMPRSS2 (293T/hACE2/TMPRSS2) or 293 cells stably expressing hACE2 (293/hACE2) prepared using lentivirus system. Briefly, a total of $4x10^5$ 293T cells were grown in each well of an 8-well chamber slide (SPL Life Sciences, Pyeongtaek, South Korea) that was previously coated with 2% gelatin. Upon overnight incubation, the cells were transfected with vectors harboring hACE2 and TMPRSS2 by using PEI. On the next day, the medium was aspirated from the cell monolayers and cells were subjected to pretreatment with a complete medium containing TE at 10 and 100 µg/ml or curcumin at 10 and 100 µM for 30 min. Then, the medium was removed and the PSV was added at 1:2 ratios in 300 µl of medium containing TE or curcumin. After 16 h of viral infection, the cells were fixed and mounted. Finally, the images of GFP dots that represented the internalization of pseudovirus were acquired using a motorized fluorescence microscope (Olympus IX83). The GFP dots were counted from 8 different areas and analyzed by Fiji software (National Institute of Health).

293/hACE2 stable cells were generated by transduction of 293 cells using lentivirus produced in 293T cells utilizing plasmids: pCMV-dR8.2-dvpr (Addgene #8455); pCAGGS-G-Kan (Kerafast EH1017); and (RRL.sin.cPPT.SFFV/Ace2.IRES-neo.WPRE(MT129)(Addgene#145840). 293/hACE2 population were then maintained in culture medium containing G418 to obtain 293 cells that stably expressing recombinant hACE2. For the PSV entry assay using 293/hACE2, the cells were grown onto gelatin-coated 8-chamber slides and incubated overnight before PSV treatment. The GFP dots were counted from 5 different areas and analyzed by Fiji software.

**2.8. Syncytia inhibition assay**

293T cells were seeded in a 24-well plate at $1.4 \times 10^5$ cells/ml density in DMEM supplemented with 10% FBS. After overnight incubation, the cells were co-transfected with plasmids bearing SARS-CoV-2 Spike, hACE2, and TMPRSS2 and incubated for about 6 h. Subsequently, 500 μl/well of dilutions of curcumin at 1 and 10 μM or turmeric extract at 1 μg/ml and 10 μg/ml was added to cell monolayers in duplo and incubated for about 16 h. The following day, the syncytia formation was observed using an inverted microscope (Olympus CKX53) and 10 images were acquired to represent each well. The total number of syncytia was calculated with Fiji software and then sorted based on the number of nuclei using 4 categories: (i) <5 nuclei, (ii) 6-10 nuclei, (iii) 11-15 nuclei, and (iv) >15 nuclei.

**2.9. Statistical analysis**

Data were presented as mean ± SD (standard deviation) as indicated in each figure. Student's t-test calculated statistical differences and a $p$-value $<0.05$ was considered significant.

**3. RESULTS**

**3.1. Cell viability assay of curcumin and TE on CHO-K1 and 293T cells**

Curcumin and TE have been known to show cytotoxic effects in various cancer cells for instance breast cancer cells, colorectal cancer cells, and brain cancer cells (Tomeh et al., 2019). In addition, even though curcumin and TE show less cytotoxicity against normal cells at high concentrations, the toxic effects could still be observed (Li et al., 2022). Therefore, before we investigated their antiviral activities, we evaluated the cytotoxicity of curcumin and TE on CHO-K1 and 293T cells to determine the non-toxic concentration to be used in PSV entry and syncytia fusion assays.

We applied 1, 5, 10, 25, and 50 µM curcumin serial concentrations and 1, 5, 10, 25, and 50 µg/ml TE for MTT assay in CHO-K1 cells. The results showed that following a 24 h treatment, CHO-K1 cell viability declined according to increase in curcumin concentrations which were 95.82, 91.08, 72.61, 41.66, and 26.27 %, respectively. While after 24 h of TE treatment, the cell viabilities were 91.70, 96.09, 92.64, 39.06, and 35.19 % as the concentration increased. Moreover, we tested the effect of curcumin and TE in 293T cells using serial concentrations of 1, 10, and 100 µM curcumin or 1, 10, and 100 µg/ml TE for 24 h. The results indicated that the cell viabilities of 293 cells after curcumin treatment were 86.52, 74.18, and 11.09 %. In addition, the cell viabilities after TE treatment were 81.52, 76.54, and 5.33 %, respectively (**Fig 1**). Higher concentrations of tested samples significantly reduced cell viability, with the 293T cells showing more susceptibility to curcumin and TE treatment than CHO-K1 cells. Therefore, to minimize the cytotoxic effect of tested samples in further assays, we applied lower concentrations of curcumin (1 and 10 uM) and TE (1 and 10 ug/ml) and treated the cells for only about 16-18 h.

**Fig 1. CHO-K1 (a-b) cell viability profiles and 293T cells (c-d) after treatment with**

**curcumin and turmeric extract.** Cells were plated onto a 96-well plate and treated with the

tested samples in triplicate for 24 sh. The medium was then replaced with MTT-containing

medium and incubated for 2 h. At the end of incubation, the formazan generated was dissolved

with DMSO and the optical density was measured at 570 nm.

**3.2. Pseudovirus entry assay**

We prepared SARS-CoV-2 PSV with a VSV backbone and GFP reporter in Spike-

transfected 293T cells (pseudotyped Spike*ΔG-GFP rVSV). Spike expression in 293T/Spike

cells used for pseudotyping was confirmed by Western blot. Moreover, we also detected Spike

in the CM obtained after pseudotyping, indicating the formation of SARS-CoV-2 pseudovirus

and its release into the culture medium (**Fig 2**).

Furthermore, we used the CM to perform PSV entry assay in 293T overexpressing

hACE2/TMPRSS2 and in 293/ACE2 stable cells. The 293T/hACE2/TMPRSS2 cells were

pretreated with curcumin or TE for 30 min, then treated with PSV/curcumin or PSV/TE and

followed by 16-18 h incubation in a $CO_2$ incubator. GFP expression indicating PSV

internalization and viral genome release was observed by fluorescence microscopy with 60x

magnification to expose the GFP dots. To confirm that GFP dots were formed within the cells,

we also observed the cells in bright field mode and used DAPI to stain the nuclei. As the results

showed, nontreated cells showed a higher number of GFP dots compared to the treated cells

with fluorescence focus units (FFU) 296$\pm$2. The FFU of cells treated with curcumin 1 and 10

µM were 141$\pm$8.76 and 56$\pm$6.52. Whereas, FFU of cells treated with TE 1 and 10 µg/ml were

81+5.93 and 49$\pm$3.31. Cells treated with curcumin and TE significantly reduced the FFU

number, especially at higher concentrations, indicating the potential inhibitory effect on SARS-

CoV-2 viral cell entry (**Fig 3**).

Next, we clarified the inhibitory effect of curcumin as the active compound to inhibit

SARS-CoV-2 PSV entry by performing an entry assay in 293 cells that stably expressed ACE2

(293/ACE2). Besides treating the 293/ACE2 cells with curcumin prior to PSV addition, the

curcumin was also combined with PSV after incubation for 30 min. The number of GFP

dots/cell for non-treated cells was 1.45. The number of GFP dots per cell in curcumin and

PSV/curcumin-treated cells was 0.99, while the number of GFP dots per cell in curcumin-

treated cells was 0.887. The results indicated that curcumin reduced PSV entry, especially for

curcumin pretreatment before the addition of PSV (P = 0.035) (**Fig 4**).

**Fig 2. Observation of Spike, hACE2, and TMPRSS2 expression.** a. Detection of Spike

expression in 293T cells transfected with Spike plasmid. b. Detection of Spike in the

conditioned medium, which represented SARS-CoV-2 PSV. c-d. Detection of hACE2 and

TMPRSS2 expression by immunofluorescence staining.

**Fig 3. Effect of turmeric extract and curcumin on PSV entry in 293T/hACE2/TMPRSS2**

**cells.** a. Schematic representation of PSV entry assay in 293T/hACE2/TMPRSS2 cells. b.

Representative image showing GFP dots to represent PSV internalization. c. Graph

representing quantification of GFP dots (n = 8 microscope fields). TE: turmeric extract; Cur:

curcumin; NC: nontreated cells.

**Fig 4. Effect of turmeric extract and curcumin on PSV entry in 293/ACE2 cells.** a. Observation of hACE2 expression in 293/hACE2 cells after transduction, before and after selection with antibiotic G418. b. Schematic representation of PSV entry assay in 293/hACE2 stable cells. c. Representative image showing GFP dots to represent PSV internalization. d. Graph representing quantification of GFP dots (n = 5 microscope fields). $P<0.05$ = *. NC: nontreated cells.

**3.3. Syncytia formation assay**

This assay represents the cells infected with SARS-CoV-2 and expressing its membrane protein Spike which has the capability to bind hACE2 receptor of the adjacent cell. Instead of using the original virus, we transfected the cells directly with Spike plasmid. We used 293T cells transfected with plasmids encoding SARS-CoV-2 Spike, hACE2, and TMPRSS2 to perform the syncytia assay. The cells expressing Spike will locate the Spike in the transmembrane region, enabling the interaction between Spike with the hACE2 receptor and TMPRSS2 of the adjacent cells. As a result, the two interacting cells will form an intercellular bridge that eventually will fuse and form multinucleated cells (Buchrieser et al., 2020). The more cells interact and fuse, the more multinucleated cells or syncytia will be generated.

Here we show that 293T cells after cotransfection and treatment with DMSO 0.1% and 1% formed syncytia as much as $157.5\pm5.9$ and $314\pm11.32$ per field. In addition, the treatment of cells with curcumin and TE reduced the number of syncytia. The number of syncytia formed after cell treatment with curcumin 1 and 10 µM was 158.5 (P<0.0001 vs DMSO 0.1%) and 55 (P<0.0001 vs DMSO 1%), while the number of syncytia formed after cell treatment with TE 1 and 10 µg/ml was 140.5 (P<0.0001 vs DMSO 0.1%) and 66.5 (P<0.0001 vs DMSO 0.1%) (**Fig 5a-b**).

Furthermore, we analyzed the distribution of nuclei numbers present within the syncytia and categorised it into nuclei number <5, 5-10, 11-15, and <15. The syncytia with nuclei number <5 for DMSO 0.1, curcumin 1 µM and TE 1 µg/ml were 36.5, 20, 42.5, while nuclei number 5-10 were 98, 35.5, 45, nuclei no 11-15 were 59.5, 43.5, 33, and nuclei number <15 were 117.5, 42.5, 28. The syncytia with nuclei number <5 for DMSO 1, curcumin 10 µM and TE 10 µg/ml were 20, 18, 20, while nuclei number 5-10 were 60.5, 23, 17.5, nuclei no 11-15 were 38.5, 18.5, 9, and nuclei no <15 were 38.5 ,18.5, 4. These data indicated that treatment of

the cells with curcumin and TE reduced the nuclei number within syncytia, representing lower

fusion events compared to the DMSO treatment (**Fig 5c**).

**Fig 5. Effect of curcumin and turmeric extract on syncytia formation.** a. Representative

image showing the syncytia formed after curcumin, TE, and DMSO treatment. b. Graph

representing quantification of total syncytia (n = 10 microscopic fields). c. Graph representing

quantification of nuclei number per categories of syncytia (n = 10 microscopic fields). Red

area: syncytia. $P<0.05 = *$, $P<0.01 = **$, $P<0.001 = ***$, ns = not significant.

**4. DISCUSSION**

Curcumin has been tested for its anti-SARS-CoV-2 activities by plaque assay in Vero cells. Using the original virus, curcumin inhibits SARS-CoV-2 infection (Marin-Palma et al., 2021). However, surface receptors in Vero cells are different from human cells, and the study may not correlate well with actual events (Pandamooz et al., 2022). Thus, antiviral study of curcumin has also been carried out using human Calu-3 cells, which endogenously express hACE2 and TMPRSS2. A study using these cells demonstrated that curcumin may inhibit SARS-CoV-2 viral replication as indicated by reduced N protein expression following viral infection (Bormann et al., 2021). Besides investigating curcumin's antiviral effect, Bormann et al. (2021) also tested the effect of turmeric juice (water extract). However, antiviral study of turmeric extract (TE) has not been widely reported compared to its active compound, curcumin.

In this manuscript, we reported the SARS-CoV-2 antiviral activities of curcumin and TE especially at the entry point, by utilising the pseudovirus model, and also the syncytia model to represent the cell-to-cell transmission. Here we showed that curcumin and TE reduced PSV entry in 293T/hACE2/TMPRSS2 cells in which 10 µM curcumin and 10 µg/ml TE significantly affected the number of GFP dots. The effect of TE was comparable to the curcumin effect on PSV cell entry in which 10 µg/ml TE was equal to 2.86 µg/ml curcuminoid with the majority of curcumin content and 10 µM curcumin was equal to 3.68 µg/ml of curcumin.

From the previous studies, Marin-Palma et al. (2021) reported that 10 µM curcumin is capable of inhibiting SARS-CoV-2 infection in Vero E6 cells. Wen et al. (2007) also reported that curcumin inhibits SARS-CoV-2 infection at concentrations of 3-10 µM. Furthermore, in A549 cells, curcumin shows SARS-CoV-2 antiviral concentration of 5 µM (Goc et al., 2021). We also showed that curcumin inhibited PSV entry in 293/hACE2 cells. These results

corroborate curcumin effects against SARS-CoV-2 infection with our data representing curcumin inhibition at PSV cell entry point. Previous reports by Mounce et al. (2017). Chen et al. (2010) and Chen et al. (2013) suggest that curcumin affects the early stages of viral replication cycles including viral-receptor attachment, internalization and fusion that have been studied against several types of viruses which involve influenza, dengue, zika, chikungunya, pseudorabies, and VSV.

Moreover, both curcumin and TE inhibit secondary infection via cell-to-cell transmission in a syncytia formation model mediated by SARS-CoV-2 Spike expression. Cells treated with curcumin and TE showed smaller syncytia with fewer nuclei than control cells. The more cells fused to generate syncytia, the larger syncytia with more nuclei will be formed and vice versa.

Based on in silico data, curcumin can interact with SARS-CoV-2 Spike RBD (Shanmugarajan et al., 2020), hACE2 (Subbaiyan et al., 2020), and TMPRSS2 as well (Motohashi et al., 2020). These data align with our results that curcumin inhibited PSV entry and syncytia formation. Curcumin is capable of interfering with the Spike-receptor binding either during direct viral infection or intercellular transmission, which hinders viral infection and cell fusion (Marin-Palma et al., 2021). In addition, TE as the crude extract which contains curcumin also has the potential to inhibit SARS-CoV-2 infection and potentially to be developed as an independent herbal-derived product for prevention of viral infection with curcuminoids used as identity compounds for TE standardization.

## 5. CONCLUSION

Our in vitro study using PSV and syncytia models revealed that both curcumin and TE are potential inhibitors of SARS-CoV-2 infection, especially at the entry points either by direct infection or cell-to-cell transmission mediated by Spike induced cell fusion.

## REFERENCES

Bormann M, Alt M, Schipper L, van de Sand L, Le-Trilling VTK, Rink L, et al (2021) Turmeric Root and Its Bioactive Ingredient Curcumin Effectively Neutralize SARS-CoV-2 In Vitro. Viruses 13, 1914. https://doi.org/10.3390/v13101914.

Buchrieser J, Dufloo J, Hubert M, Monel B, Planas D, Rajah MM, et al (2020) Syncytia formation by SARS-CoV-2-infected cells. The EMBO Journal 39(23), e106267. https://doi.org/10.15252/embj.2020106267.

Chen TY, Chen DY, Wen HW, Ou JL, Chiou SS, Chen JM, et al (2013) Inhibition of Enveloped Viruses Infectivity by Curcumin. Plos One 8(5), e62482. https://doi.org/10.1371/journal.pone.0062482.

Chen DY, Shien JH, Tiley L, Chiou SS, Wang SY, Chang TJ, et al (2010) Curcumin inhibits influenza virus infection and haemagglutination activity. Food Chemistry 119(4), 1346–1351. https://doi.org/10.1016/j.foodchem.2009.09.011.

Goc A, Sumera W, Rath M, Niedzwiecki A (2021) Phenolic compounds disrupt spike-mediated receptor-binding and entry of SARS-CoV-2 pseudo-virions. Plos One 16(6), e0253489. https://doi.org/10.1371/journal.pone.0253489.

Hirano T, Murakami M (2020) COVID-19: A New Virus, but a Familiar Receptor and

Cytokine Release Syndrome. Immunity 52(5):731-733.

https://doi.org/10.1016/j.immuni.2020.04.003

Hoffmann M, Kleine-Weber H, Schroeder S, Krüger N, Herrler T, Erichsen S, et al (2020)

SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a

Clinically Proven Protease Inhibitor. Cell 181:271-280.e8.

Jackson CB, Farzan M, Chen B, Choe H (2021) Mechanisms of SARS-CoV-2 entry into cells.

Nat Rev Mol Cell Biol 3-20. https://doi.org/10.1038/s41580-021-00418-x.

Katopodis P, Randeva HS, Spandidos DA, Saravi S, Kyrou I, Karteris E (2022) Host cell entry

mediators implicated in the cellular tropism of SARS-CoV-2, the pathophysiology of

COVID-19 and the identification of microRNAs that can modulate the expression of

these mediators (Review). Int J Mol Med 49(2):20.

https://doi.org/10.3892/ijmm.2021.5075.

Li P, Pu S, Lin C, Liu H, Zhao H, Yang C, et al (2022) Curcumin selectively induces colon

cancer cell apoptosis and S cell cycle arrest by regulates Rb/E2F/p53 pathway. J. Mol.

Struct, 1263: 133180. https://doi.org/10.1016/j.molstruc.2022.133180.

Lin L, Li Q, Wang Y, Shi Y (2021) Syncytia formation during SARS-CoV-2 lung infection: a

disastrous unity to eliminate lymphocytes. Cell Death Differ 28, 2019–2021.

https://doi.org/10.1038/s41418-021-00795-y.

Marín-Palma D, Tabares-Guevara JH, Zapata-Cardona MI, Flórez-Álvarez L, Yepes LM,

Rugeles MT, et al. (2021) Curcumin Inhibits In Vitro SARS-CoV-2 Infection In Vero E6

Cells through Multiple Antiviral Mechanisms. J Molecules 26(22) :6900.

https://doi.org/10.3390/molecules26226900.

Maurya VK, Kumar S, Prasad AK, Bhatt MLB, Saxena SK (2020) Structure-based drug

designing for potential antiviral activity of selected natural products from Ayurveda

against SARS-CoV-2 spike glycoprotein and its cellular receptor. Virusdisease

31(2):179–193. https://doi.org/10.1007/s13337-020-00598-8.

Motohashi N, Vanam A, Gollapudi R (2020) In Silico Study of Curcumin and Folic Acid as

Potent Inhibitors of Human Transmembrane Protease Serine 2 in the Treatment of

COVID-19.    INNOSC    Theranostics    and    Pharmacological    Sciences,    3–9.

https://doi.org/10.36922/itps.v3i2.935.

Mounce BC, Cesaro T, Carrau L, Vallet T, Vignuzzi M (2017) Curcumin inhibits Zika and

chikungunya virus infection by inhibiting cell binding. Antiviral Research 142, 148–157.

https://doi.org/10.1016/j.antiviral.2017.03.014.

Pandamooz S, Jurek B, Meinung CP, Baharvand Z, Shahem-abadi AS, Haerteis S, et al (2022)

Experimental Models of SARS-CoV-2 Infection: Possible Platforms to Study COVID-

19 Pathogenesis and Potential Treatments. Annu Rev Pharmacol Toxicol 62 :25-53.

https://doi.org/10.1146/annurev-pharmtox-121120-012309.

Peng L, Gao L, Wu X, Fan Y, Liu M, Chen J, et al (2022) Lung Organoids as Model to Study

SARS-CoV-2 Infection. Cells 11(17):2758. https://doi.org/10.3390/cells11172758.

Rajah MM, Bernier A, Buchrieser J, Schwartz O (2022) The Mechanism and Consequences of

SARS-CoV-2 Spike-Mediated Fusion and Syncytia Formation. Journal of Molecular

Biology 2022;434 :167280. https://doi.org/10.1016/j.jmb.2021.167280.

Rajah MM, Hubert M, Bishop E, Saunders N, Robinot R, Grzelak L, et al (2021) SARS-CoV-

2 Alpha, Beta, and Delta variants display enhanced Spike-mediated syncytia formation.

The EMBO Journal 40:e108944. https://doi.org/10.15252/embj.2021108944 .

Saccon E, Chen X, Mikaeloff F, Rodriguez JE, Szekely L, Vinhas BS, et al (2021) Cell-type-

resolved quantitative proteomics map of interferon response against SARS-CoV-2.

iScience 24:102420. https://doi.org/10.1016/j.isci.2021.102420.

Shanmugarajan D, Prabitha P, Kumar BRP, Suresh B (2020) Curcumin to inhibit binding of

spike glycoprotein to ACE2 receptors: computational modelling, simulations, and

ADMET studies to explore curcuminoids against novel SARS-CoV-2 targets, RSC Adv

10: 31385-31399. https://doi.org/10.1039/D0RA03167D.

Sharifi-Rad J, Rayess YE, Rizk AA, Sadaka C, Zgheib R, Zam W, et al (2020) Turmeric and

Its Major Compound Curcumin on Health: Bioactive Effects and Safety Profiles for Food,

Pharmaceutical, Biotechnological and Medicinal Applications. Front Pharmacol

11:01021. https://doi.org/10.3389/fphar.2020.01021.

Subbaiyan A, Ravichandran K, Singh SV, Sankar M, Thomas P, Dhama, K, et al (2020) In

silico Molecular Docking Analysis Targeting SARS-CoV-2 Spike Protein and Selected

Herbal Constituents. J Pure Appl Microbiol 14(suppl 1):989-998.

https://doi.org/10.22207/JPAM.14.SPL1.37.

Tahmasebi S, El-Esawi MA, Mahmoud ZH, Timoshin A, Valizadeh H, Roshangar L, et al

(2021) Immunomodulatory effects of nanocurcumin on Th17 cell responses in mild and

severe COVID-19 patients. J Cell Physiol 236(7), 5325–5338.

https://doi.org/10.1002/jcp.30233.

Tomeh MA, Hadianamrei R, Zhao X (2019) A Review of Curcumin and Its Derivatives as

Anticancer Agents. Int. J. Mol. Sci 20(5):1033. https://doi.org/10.3390/ijms20051033.

Valizadeh H, Abdolmohammadi-Vahid S, Danshina S, Ziya GM, Ammari A, Sadeghi A, et al

(2020) Nano-curcumin therapy, a promising method in modulating inflammatory

cytokines in COVID-19 patients. Int Immunopharmacology 89(PtB),107088.

https://doi.org/10.1016/j.intimp.2020.107088.

Wen CC, Kuo YH, Jan JT, Liang PH, Wang SY, Liu HG et al (2007) Specific plant terpenoids

and lignoids possess potent antiviral activities against severe acute respiratory syndrome

coronavirus. J Med Chem 50(17), 4087–4095. https://doi.org/10.1021/jm070295s.

WHO (1999). Monographs on Selected Medicinal Plants. (Geneva: World Health

Organization; ), 1, 295.

Zheng Y, Zhou LL, Su Y, Sun Q (2021) Cell fusion in the pathogenesis of COVID-19. Mil

Med Res 8(1):68. https://doi.org/10.1186/s40779-021-00348-x.

**ABBREVIATIONS**

**ACE2**          **:** angiotensin converting enzyme-2

**COVID-19**      : coronavirus disease-2019

**GFP**          : green fluorescence protein

**PSV**          : pseudovirus

**RBD**          : receptor binding domain

**SARS-CoV-2** : severe acute respiratory syndrome coronavirus-2

**TE**            : turmeric extract

**VSV**          : vesicular stomatitis virus