Elasticity of the HIV-1 Core Facilitates Nuclear Entry and Infection

Akshay Deshpande[1,#], Alexander J. Bryer[2,#], Jonathan Andino[3], Jiong Shi[4], Jun Hong[4], Cameron

Torres[4], Shimon Harel[1], Ashwanth C. Francis[3,5], Juan R. Perilla[2,*], Christopher Aiken[4,*], Itay

Rousso[1,*]

[1]Ben-Gurion University of the Negev, Department of Physiology and Cell Biology, Beer Sheva, Israel;
[2]University of Delaware, Department of Chemistry and Biochemistry, Newark DE, USA; [3]Florida State
University, Institute of Molecular Biophysics, Tallahassee, FL, USA; [4]Vanderbilt University Medical
Center, Department of Pathology, Microbiology and Immunology, Nashville, TN, USA; [5]Florida State
University, Department of Biological Sciences, Tallahassee, FL, USA

*Corresponding authors

#Co-first authors

HIV-1 infection requires passage of the viral core through the nuclear pore of the cell, a process that depends on functions of the viral capsid [1,2]. Recent studies have shown that HIV-1 cores enter the nucleus prior to capsid disassembly [3-5]. Interactions with the nuclear pore complex are necessary but not sufficient for nuclear entry, and the mechanism by which the viral core traverses the comparably sized nuclear pore is unknown. Here we show that the HIV-1 core is highly elastic and that this property is linked to nuclear entry and infectivity. Using atomic force microscopy-based approaches, we found that purified wild type cores rapidly returned to their normal conical morphology following a severe compression. Results from independently performed molecular dynamic simulations of the mature HIV-1 capsid also revealed its elastic property. Analysis of four HIV-1 capsid mutants that exhibit impaired nuclear entry revealed that the mutant viral cores are brittle. Suppressors of the mutants restored elasticity and rescued infectivity and nuclear entry. Elasticity was also reduced by treatment of cores with the capsid-targeting compound PF74 and the antiviral drug lenacapavir. Our results indicate that capsid elasticity is a fundamental property of the HIV-1 core that enables its passage through the nuclear pore complex, thereby facilitating infection. These results provide new insights into the mechanisms of HIV-1 nuclear entry and the antiviral mechanisms of HIV-1 capsid inhibitors.

The HIV-1 capsid, which forms the shell of the core and acts as a protective barrier, encapsidates the enzymes reverse transcriptase and integrase and the viral genomic RNA. During infection, the HIV-1 genome is transported into the nucleus and integrates into host cell chromosomal DNA. Recent studies suggest that the core remains fully or mostly intact during nuclear entry, thereby ensuring completion of reverse transcription within the nucleus [3,4]. It is presently unclear how the core passes through the nuclear pore [6-8] while remaining intact. Docking

of the HIV-1 core at the NPC and subsequent nuclear entry are effected by interactions between the capsid and cellular factors, including cyclophilin A, CPSF6, and nucleoporins (e.g. Nup358 and Nup153) [6,9-16]. Treatment with low concentrations of the capsid-binding compounds PF-3450074 (PF74) [17] and Lenacapavir (LCV) [18] reduces HIV-1 infectivity by inhibiting nuclear entry [19-27]. Nuclear entry can also be reduced by mutations in the capsid protein (CA). For example, the capsid-stabilizing CA mutation E45A markedly reduces nuclear entry and consequently infectivity [28,29] (Fig. 1 A-D and Fig. 2). However, position 45 in CA resides at an inter-subunit interface in the capsid that is not a binding site for known host proteins that promote nuclear entry [2]. In addition, E45 lies outside the binding site for PF74 and LCV, yet the mutant virus is resistant to these compounds [20,30]. Remarkably, the suppressor mutation R132T rescues nuclear entry and infectivity of the E45A mutant and restores its sensitivity to PF74 without reversing the hyperstability of the capsid [29].

In earlier work, we employed atomic force microscopy (AFM) to show that CA mutations can affect the physical property of the capsid known as stiffness, defined as the resistance to an applied force, To determine whether capsid stiffness plays a role in HIV-1 nuclear entry and sensitivity to capsid inhibitors, we quantified the stiffness of purified HIV-1 cores using AFM operated in the nano-indentation mode (Fig. 1F). Untreated WT cores exhibited a mean stiffness value of 0.12 ± 0.01 N/m, and addition of the capsid-stabilizing host cell metabolite inositol hexakisphosphate (IP6) [31] (100 μM) increased the stiffness to a mean value of 0.24 ± 0.02 N/m. Cores treated with both IP6 and the antiviral compounds PF74 (1.25 μM) or IP6 and LCV (500 pM) exhibited a comparable increase in stiffness (value of 0.20 ± 0.02 N/m or 0.30 ± 0.03 N/m, respectively). Thus, addition of the capsid-targeting antiviral compounds at concentrations that inhibit HIV-1 nuclear entry did not further increase the stiffness of HIV-1 cores that were treated

with IP6. We analyzed three additional CA mutants Q63A/Q67A, E212A/E213A and E180A. Like

E45A, these mutants exhibited low infectivity, impaired nuclear entry, and cell-cycle-dependent

infection yet underwent efficient reverse transcription in target cells (Fig. 1A-D and Fig. 2).

Moreover, infection by these mutants was minimally affected by depletion of Nup153 (Fig. 1E) [32,]

[33]. Purified cores from these mutants also exhibited increased stiffness. Q63A/Q67A cores showed

the highest mean stiffness value of 0.31 ± 0.03 N/m, whereas the value for E180A and

E212A/E213A cores was 0.26 ± 0.03 N/m. The pseudorevertant Q63A/Q67E emerged during

prolonged culture of the Q63A/Q67A mutant in T cells and exhibited nuclear entry and infectivity

similar to the wild type (Fig. 1 and 2). The mean stiffness value of this mutant was reduced relative

to its parent mutant Q63A/Q67A (0.23 ± 0.03 vs. 0.31 ± 0.03, respectively; P= 0.0455). As

previously reported, the suppressor mutation R132T partially rescued E45A infectivity and nuclear

entry (Fig. 1A, B and Fig. 2), though it does not restore normal capsid stability [29]. These variants

exhibited similar mean stiffness values (0.20 ± 0.03 N/m for E45A and 0.21 ± 0.02 N/m for

E45A/R132T; p > 0.05). Collectively, these observations indicate that rescue of E45A and

Q63A/Q67A nuclear entry by R132T and Q67E, respectively, does not involve restoration of

normal core stiffness, suggesting that another property of the core promotes nuclear entry and

infectivity.

Studies of the HIV-1 capsid have demonstrated that the inter-subunit interfaces are

structurally variable, indicating that the capsid is flexible [34,35]. We hypothesized that capsid

elasticity is required for HIV-1 nuclear entry. To quantify elasticity, we exploited the ability of the

AFM probe to induce structural deformation in the viral core and analyzed the consequences on

the mechanical behavior and structural integrity of the capsid. For this purpose, the AFM was

operated in the nano-indentation mode but with high maximal loading forces (≥ 5 nN, vs. the 1–

1.5 nN range used in stiffness measurements) to explore the mechanical tolerance limits of the

cores and quantify their critical loading force: the force at which there is an abrupt drop in the

force-distance curve, representing the failure and collapse of the structure [36]. We observed no

evidence for structural failure in the force–distance curves, even at a maximal loading force of 20

nN (Fig. 3A). Morphological analysis by scanning AFM further demonstrated that all cores (n=72)

showed no evidence for structural damage following perturbation (Extended Data Fig. 1). For one

core in the 10 nN applied force group, we observed a drop in the force–distance curve (Fig. 3A

inset). Morphological analysis of this core structure showed that it remained intact and non-

deformed. Therefore, this core likely underwent reversible buckling rather than breakage. The

results show that HIV-1 cores can withstand marked deformation without undergoing irreversible

structural damage, indicating that the HIV-1 capsid is highly elastic. This finding is in sharp

contrast to other viruses, such as bacteriophage HK97, hepatitis B virus, and herpes simplex virus,

which have brittle capsids that undergo structural failure at forces as low as 1-6 nN [37].

To complement the AFM measurements, we performed coarse grained *in silico*

computational nanoindentation simulations at four distinct regions along the wild-type capsid

surface to model the resistance of the capsid to deformation. After placing four spherical AFM

probes equidistant along the capsid's surface (Fig. 3 B-E), we modeled a constant pushing velocity

to the tip (3.125 nm/μs). While the simulated tip velocity is higher than what is achievable in

physical experiments, simulations can provide a high temporal and spatial resolution view of how

the capsid deforms and responds to mechanical stress. Indentation of the broader region of the

capsid initially showed the highest stiffness among the four probed regions, with the capsid

appearing to buckle after 3 nm of indentation demonstrating a change in the elasticity of the

capsid's mechanical properties at indentation depths beyond 3 nm (Fig. 3B). Indentations in the

relatively flat, middle regions of the capsid revealed the lowest stiffness (Fig. 3C, D) and the most

linearity in FD-curves. The narrow end of the capsid showed high stiffness through the indentation

profile, until contact with the baseplate is made (Fig. 3E). Our simulations support our

experimental AFM data and show that the wild-type capsid is an elastic container, able to withstand

deformations without failure of the intermolecular interfaces comprising the CA lattice. The

different stiffness values observed at different positions further suggests that the HIV-1 capsid has

a material heterogeneity that is determined by its variable curvature.

Simulated nanoindentation of E45A and E45A/R132T capsids showed that the mutations

produced a significant increase in stiffness (Extended Data Fig. 2) and, additionally, affect capsid

deformability (Extended Data Figs. 2-5), with E45A capsids in particular rupturing at relatively

smaller indentation distances compared to wild type and E45A/R132T while absorbing more

power through deformation. Importantly, when we considered simulated probe locations, and thus

regions of the capsid, independently, we found that wild type, E45A and E45A/R132T capsids

have similar stiffness in the low curvature mid-regions of the capsid (Extended Data Fig. 6). At

the highly curved broad end, the wild type was slightly stiffer than either mutant, further

demonstrating that the capsid's material response is predicated by its shape and location of

indentation. The simulations show that the wild-type capsid is a robust and elastic container, able

to sustain extreme deformations and recover (Extended Data Fig. 7) prior to failure of the

intermolecular interfaces comprising the CA lattice. The simulations further suggest that the capsid

has a material heterogeneity that is dictated by its curvature.

To physically assess the elasticity of HIV-1 cores, we analyzed the shape of the viral core

soon after release of strong applied force. In this approach, a core was first imaged by scanning at

a low loading force of 300 pN (Figure 3A). We then selected a region of interest (Extended Data

Fig. 8) located in the main body of the core and rescanned it. This region was then scanned at a

high loading force (5 nN) that caused the core to compress dramatically at this region. The force

was immediately reduced to 300 pN and the compressed region rapidly imaged repeatedly over

time (Fig. 4B). The volume under the scanned region was then calculated as a function of time.

Fig. 4C shows two representative trajectories in which the volume was compressed to 50% and

30% of the initial volume and recovered to nearly 100% and 80%, respectively. Mean values of

the compressed volume (pressed: 39 ± 3%), the immediate recovery volume (Immediate: 92 ±

1%), and the volume 5–7 minutes after compressing the core (later: 92 ± 1%) were determined

from analysis of 36 individual cores (Fig. 4D). Wild type HIV-1 cores displayed nearly complete

recovery after being strongly compressed. The measured recovery volume was highly consistent

individual core volume trajectories. Consequently, the averaged immediate and late volumes were

identical, indicating that the core recovered its initial shape in a time shorter than the temporal

resolution of the assay (10–15 seconds) (Fig. 4C and D). These results further demonstrate that

HIV-1 cores are highly elastic and rapidly recover nearly their full initial volume after being

strongly compressed. For the majority of cores, we observed no structural damage in the

compressed region of the CA-lattice. In 14% of cores analyzed, (5 of 36), structural damage was

apparent near the end of the core following compression and recovery (Fig. 5A). We previously

reported that capsid disassembly begins at the end of isolated cores during *in vitro* reverse

transcription. This highly curved and pentamer-rich section has been proposed to be the least stable

region in the HIV-1 capsid [38]. Our data are consistent with this model, in which the weakest region

in the structure collapses as the core is compressed.

To investigate the potential role of elasticity in HIV-1 infection, we measured the structural

recovery of cores from CA mutants that exhibit impaired nuclear entry and quantified the

percentage of broken cores following compression as a measure of inelasticity/brittleness (Fig.

5B). Representative AFM images of intact and broken cores following compression are shown in

Extended Data Fig. 9. In contrast to the wild type, E45A mutant cores were more brittle (41% of

cores were broken following compression). By contrast, E45A/R132T cores were less brittle,

exhibiting breakage (21%) similar to wild type cores, indicating that the addition of the suppressor

mutation restored elasticity to the E45A mutant capsid narrow end. Cores from the nuclear entry-

impaired mutants E212A/E213A, Q63A/Q67A, and E180A mutant cores were also brittle,

exhibiting 40%, 39%, and 32% breakage upon compression, respectively. Substitution of glutamic

acid for alanine at position 67 (Q63A/Q67E) restored elasticity to the wild type level (21% of

broken cores). Moreover, addition of PF74 or LCV to wild type cores rendered them brittle as

reflected by increased breakage upon compression (to 39% and 36% for PF74 and LCV,

respectively). Thus, capsid-targeting small molecules reduce HIV-1 core elasticity at

concentrations that inhibit nuclear entry. As previously shown for E45A and Q63A/Q67A [30],

infection by the nuclear entry-impaired mutants E180A and E212A/E213A was resistant to the

inhibitors. Like E45A/R132T, the pseudorevertant Q63A/Q67E was sensitive (Extended Data Fig.

10).

Our finding that the CA substitutions reduced HIV-1 core elasticity provides a compelling

mechanistic explanation for the nuclear entry defect associated with the E45A and Q63A/Q67A

mutant virus and their suppression by R132T and Q67E. By restoring elasticity, the R132T or

Q67E mutations may permit penetration of the nuclear pore complex to enable interaction of the

capsid with the nuclear basket protein Nup153, thus promoting nuclear translocation (Fig. 5C).

This model is consistent with the known PF74 resistance and susceptibility of the E45A and

E45A/R132T mutants, respectively, as well as the effects of these mutations on HIV-1 infection

of nondividing cells [29,39]. It may also account for the reduced effect of Nup153 depletion on

infection by the E45A, Q63A/Q67A, E180A and E212A/E213A mutants [1] (Fig. 1E).

Mutants with brittle cores may also exhibit impaired nuclear entry owing to premature

cytoplasmic breakage events. To address this issue, we employed live-cell fluorescence imaging

microscopy to assess the docking efficiency of these cores at the nuclear lamina (Fig. 5D). Our

findings revealed that all mutant cores successfully reach the nucleus, surpassing the threshold for

random interactions. Notably, the docking efficiency of the brittle E45A and Q63A/Q67A mutants

closely mirrored that of their elastic counterparts, E45A/R132T and Q63A/Q67E, respectively.

Furthermore, the docking efficiency of brittle E212A/E213A cores was similar to that of wild-type

(WT) cores. Collectively, our results establish that core elasticity does not correlate with the

changes in docking efficiency. Additionally, our findings emphasize that, although all core mutants

successfully reach the nuclear envelope, the key disparity lies in their ability to penetrate the

nuclear pore, a characteristic exclusively exhibited by elastic cores. Nonetheless, we cannot

exclude the possibility that the brittleness of the mutant cores leads to premature breakage of

mutant capsids during reverse transcription in the cytoplasm, which may impair docking at the

NPC and nuclear entry and result in activation of the cytoplasmic DNA sensor cGAS [40].

Our observation that addition of PF74 and LCV decreases the elasticity of the wild type

capsid further suggests that elasticity is critical for HIV-1 nuclear entry. CA mutations that

decrease the elasticity of the viral core confer resistance to these two inhibitors, likely by rendering

the virus unable to utilize the nuclear pore complex. The elasticity-reducing mutations we have

identified reside at the three major intersubunit interfaces in the capsid lattice, indicating that each

interface is critical for elasticity. The suppressor mutations restored capsid elasticity, infectivity,

nuclear entry, and PF74 and LCV sensitivity. We suggest that the HIV-1 capsid inhibitors PF74

and LCV inhibit nuclear entry by decreasing the elasticity of the viral capsid in addition to

impairing its interactions with nucleoporins.

In conclusion, HIV-1 cores are highly elastic, endowing the core with the ability to undergo

reversible deformation without structural failure. To our knowledge, this is the first reported

example of such a highly elastic viral capsid. Recent structural studies of NPCs within cells have

revealed that the diameter of the nuclear pore is sufficient to permit passage of the intact core [4].

Nonetheless, the pore is normally filled with a high density of FG domains of nucleoporins. Also,

NPCs in cells are dynamic, undergoing dilation and constriction [41]. Docking of the reverse-

transcribing viral core to the NPC may signal both the core and the NPC to dynamically adapt to

one another, thus promoting passage of the core through the pore.

Materials and Methods

HIV-1 core production and core purification

Human embryonic kidney (HEK) 293T cells were used for the preparation of HIV-1 pseudovirus particles as previously described [38]. Briefly, ~10^6 HEK 293T cells were cultured in a 15 cm dishes and transfected with 2.5 µg of Env-defective variants of the full-length HIV-1 construct R9 using 10 µg of polyethyleneimine (PEI; branched; MW ~25000; Sigma-Aldrich). After 20 h, the cell medium (Dulbecco's modified Eagle medium, supplemented with 10% heat-inactivated fetal bovine serum, 1% penicillin-streptomycin, and 1% L-glutamine) was replaced with fresh medium. After 6–7 h, the virus-containing supernatant was harvested, centrifuged at 1,000 rpm for 10 min, and filtered through a 0.45 µm pore size filter. The supernatant was concentrated using ultracentrifugation in an SW-28 rotor (25,000 rpm, 2 h at 4ºC) on a 60% OptiPrep density cushion (Sigma-Aldrich). The pelleted viruses were resuspended in 10 mL TNE buffer (50 mM Tris-HCl, 100 mM NaCl, 0.1 mM EDTA; pH 7.4), and added to 100-kDa molecular mass cutoff Vivaspin 20 centrifugal concentrators (100,000 MWCO; Sartorius AG, Germany). The mixture was centrifuged twice at 2,500 g for 25–30 min at 4ºC until the supernatant level in the concentrator reached a final volume of 300–350 µL.

Virus cores were isolated from the concentrated virus-containing supernatant using a previously described protocol [38]. Purified HIV-1 virus particles were mixed with 1% Triton-X 100 diluted in 100 mM of 3-(N-morpholino) propane sulfonic acid (MOPS) buffer (pH 7.0) and incubated on ice for 2 min. The mixture was centrifuged at 13,800 g for 8 min at 4ºC. After removing the supernatant, the pellet was washed twice using ~80 µL of MOPS buffer and repelleted. The final pellet was resuspended in a 10 µL MOPS buffer. A fresh batch of cores was prepared on the same day as each of the AFM measurements. When specified, IP6, PF74, or

Lenacapavir were added to the buffer at a final concentration of 100 µM, 1.25 µM, or 500 pM,

respectively.

AFM measurements and analysis

Samples for AFM measurements and analysis were prepared as previously described [38].

Briefly, 10 µL of isolated HIV-1 cores were incubated for 45 min at room temperature on

hexamethyldisilazane (HMDS)-coated microscope glass slides (Sigma-Aldrich) in a mildly humid

chamber. AFM measurements were performed on the adhered sample without fixation. Each

experiment was repeated at least three times, each time with independently purified pseudoviruses.

IP6 was received as a gift from the Leo laboratory. PF-3450074 was synthesized by the Vanderbilt

Institute of Chemical Biology Synthesis Core, Vanderbilt University, Nashville, TN. Lenacapavir

(GS-6207, Gilead) was purchased from MedChemExpress (MCE), USA. All measurements were

carried out with a JPK Nanowizard Ultra-Speed atomic force microscope (JPK Instruments,

Berlin, Germany) mounted on an inverted optical microscope (Axio Observer; Carl Zeiss,

Heidelberg, Germany). Silicon nitride probes (mean cantilever spring constant of 0.12 N/m; DNP,

Bruker, Germany) were used. Topographic images were acquired using the quantitative imaging

(QI) mode at a rate of 0.5 lines/s and a loading force of 300 pN and rendered using the WSxM

software (Nanotec Electronica).

Stiffness measurements: Core stiffness was obtained by operating the AFM in nanoindentation

mode as previously described [38,42]. Briefly, to determine the stiffness of each core, 20 force-

distance (FD) curves were obtained from 24 different points on the core surface for a total of 480

FD-curves per core. To confirm that the capsid remained stable during the entire indentation

experiment, each experiment was analyzed by the plotting individual point stiffness as a histogram

and as a function of the measurement count. Samples whose point stiffness values decreased

consistently during measurement, indicative of irreversible deformation, were discarded. The

number of discarded FD-curves was 1 from each sample (corresponding to ~3.5%). Importantly,

this small number of discarded cores was similar for all samples, regardless of their elasticity. The

maximal indentation of the sample surface was 4 nm which corresponds to a maximum loading

force of 0.2–1.5 nN. Each FD-curve was shifted to set the deflection in the non-contact section to

zero (baseline). The set of FD-curves was then averaged. The measured stiffness was

mathematically derived from the slope of the FD-curves. Core stiffness was computed using

Hooke's law, assuming that the experimental system can be modeled as two springs (the core and

the cantilever) arranged in series. The spring constant of the cantilever was determined during

experiments by measuring the thermal fluctuations. The data were analyzed using JPK Data

Processing and MATLAB (The Math Works, Natick, MA) software. To test the mechanical

strength of the core, a single FD curve was acquired from a single central point on the surface of

the core. The maximum loading force was set to either 5 nN, 10 nN or 20 nN.

Volume recovery measurements: The elasticity of the core was assessed in terms of volume

recovery after forcibly compressing the structure. The entire core was imaged using the AFM

operated in the QI mode using a low maximal force of 300 pN. A small rectangular region on the

main body of the core was then selected and scanned. The chosen region was then compressed by

re-scanning it at a maximal loading force of 5 nN. Higher maximal loading forces frequently

resulted in the detachment of cores from the glass substrate as evident by the disappearance of the

core in the consequent images. The force was then lowered back to 300 pN, and the region was

repetitively imaged for a total duration of 5–6 minutes. At the end of the experiment, the entire

core was imaged at a 300 pN loading force. The collected set of images was then processed using

an in-house MatLab script to calculate the volume underneath the selected region. The initial

volume prior to compression was set to 100%, and the rest of the calculated volumes were

normalized accordingly. The obtained volume values were then plotted against time. The initial

and final images of the entire core were used to determine if the core underwent structural breakage

upon compression.

HIV-1 infection assays

Infectivity was quantified by titration of GFP reporter viruses bearing the indicated amino

acid substitutions in CA. Virus stocks were prepared by cotransfection of 3 µg of Env-defective

HIV-GFP proviral plasmid [43] with 0.5 µg pHCMV-G [44], generating VSV-pseudotyped particles.

After culturing for 40-48 h, virus-containing culture supernatants were collected, clarified by

passing through 0.45 µm syringe filters, frozen in aliquots, and stored at -80°C. Virus

concentrations were determined by p24 ELISA [45]. Infections were performed by inoculating

20,000 Hela cells seeded one day previously in 48-well plates with 0.25 ml of virus dilutions

prepared in complete medium containing 20 µg/ml DEAE-dextran. Cultured were incubated for

two days and cells detached by treatment with trypsin, collected, fixed overnight in 0.4%

paraformaldehyde, and analyzed for GFP expression by flow cytometry. Infectivity was

determined as % GFP$^+$ cells in the culture normalized by the concentration of p24 in the inoculum

with wells exhibiting between 1% and 20% GFP$^+$ cells. To assay inhibition of infection by PF-

3450074 (PF74) and Lenacapavir (both purchased from MedChemExpress), the compounds were

added to a dilution of each virus that gave 5-10% infected cells in the absence of inhibitor. Two

days later, the cells were fixed and analyzed for infection by flow cytometry.

In experiments involving cell cycle arrest, Hela cells (40,000) were seeded in medium

containing aphidicolin (1 µg/ml). The next day, medium was removed, and aphidicolin was

included in the virus inocula. The following day, the medium was replaced with fresh medium

lacking aphidicolin. In experiments involving Nup153 depletion, cells were first transfected with

control or Nup153-targeting siRNA duplexes as previously described [32]. Two days later, cells

were replated for infection assays and for immunoblotting to confirm the knockdown efficiency.

To isolate the Q63A/Q67E pseudorevertant, cultures of MT-4 cells (200,000) in 1 ml of

medium were inoculated with R9.Q63A/Q67A mutant virus (25 ng p24) encoding a Vpu

truncation at amino acid 35. The cultures were maintained and assayed for HIV-1 accumulation

by p24 ELISA every four days. After approximately two weeks, p24 was detected in the mutant

virus culture. DNA was extracted and purified using a DNAeasy kit (Qiagen), and the product

used as a template for PCR amplification of the Gag coding region. PCR products were gel

purified, sequenced, and cloned into the wild type R9 and HIV-GFP reporter virus plasmids. The

PCR-amplified regions of the plasmids were sequenced to confirm identity.

For quantifying reverse transcription and nuclear entry, monolayers of 200,000 Hela cells

in 12-well dishes were inoculated with 0.5 ml of virus, corresponding to 20 ng of p24, that had

been previously treated with DNaseI to reduce the level of contaminating plasmid DNA. Infection

was performed in medium containing 20 µg/ml DEAE-dextran. When required, Efavirenz or

Raltegravir was added to diluted viruses to a final concentration of 1 µM. Cultures were incubated

for 16 hours, cells were collected, and DNA was purified on silica columns [46]. DNA was eluted in

100 µl of water, quantified by spectrophotometry, and assayed for late reverse transcripts and 2-

LTR circles by quantitative PCR with standard curves generated from dilutions of appropriate

plasmids.

Simulated AFM nanoindentation experiments

HIV-1 CA conical capsid construction: The coarse grained HIV-1 conical capsid was modeled

using a methodology we developed previously[47]. Details of the coarse graining methodology, as

well as parameterization and model validation, were previously reported [47]. The system includes

models of sodium, chloride, and host-factor inositol hexakisphosphate (IP6), yielding 150 mM

NaCl ionic strength and 253 IP6 molecules bound to central pores of capsomers [48]. The charge-

neutral, ionized and IP6-bound capsid was then equilibrated for 500 ns using NAMD3 [49].

Molecular dynamics simulations: All molecular dynamics (MD) simulations were performed with

the nanoscale molecular dynamics (NAMD) engine [49]. NAMD 2.14 was utilized for energy

minimization and the fully GPU-resident NAMD 3 was utilized for NVT equilibration and

constant velocity steered MD (cv-SMD) simulations. Simulations of wild type capsids utilized a

time step of 48 femtoseconds, whereas mutant E45A and E45A/R132T capsid simulations utilized

a time step of 40 femtoseconds. Long range electrostatic evaluation was accomplished via particle

mesh Ewald (PME) [49,50]. The non-bonded potential excluded bonded neighbors (1-2 exclusion

policy) and utilized a cutoff of 2 nm. Short range non-bonded interactions were computed at every

time step, while long range interactions were computed every other time step. The discrete grid

used in PME for interpolating charges was given a spacing of 2 Å, employing 8th order

interpolation. Temperature was enforced via Langevin dynamics, with a target temperature of 298

K. All beads besides the AFM tip were coupled to the thermal bath with a γ coefficient of 2 ps−1.

A dielectric constant of 80 was employed to mimic charge screening by solvent molecules [51].

In silico AFM simulations: For AFM simulations, constant velocity pulling was applied to a

dummy atom with a velocity of 1.5×10^{-6} Å per time step, or 3.125 nm per microsecond. A

harmonic force constant of 12 kcal/mol was employed to restrain the 13 nm AFM tip's center of

mass to the dummy atom. Additionally, a transverse harmonic force constant of 1 kcal/mol was

employed. Data from the AFM simulations, namely forces acting on the AFM tip throughout

simulation as well as the tip's position data, were output with a frequency of 48 picoseconds, or

every 100 time steps (A detailed description of the AFM simulations is provided in the Extended

Data).

Live-cell imaging of HIV-1 docking at the nuclear envelope and nuclear import

Single HIV-1 infection in live cells was visualized, as previously described [13,52]. In brief, $8 \cdot 10^4$ or

$5 \cdot 10^5$ TZM-bl cells that stably express the emiRFP670-laminB1 nuclear envelope marker was

respectively plated in a 8-well chambered slide (#C8-1.5H-N, CellVis) for single time-point

nuclear import experiments, or in a 35mm 1.5 glass bottom dish (#D35-20-1.5-N, CellVis) for NE-

interactions experiments. Aphidicolin (Sigma Aldrich) 10 µM was added to cells to block cell-

division. About 14h later the cells were infected (MOI 0.2, for NE interactions (or) MOI 1 for

nuclear import) with indicated WT and mutant HIVeGFP particles fluorescently tagged with Vpr-

integrase-mNeongreen (INmNG) to label the viral core. Virus binding to cells was augmented by

spinoculation (1500×g for 30 min, 16°C). Following spinoculation, the cells were washed once,

and virus entry was synchronously initiated by adding pre-warmed complete DMEM medium

containing Aphidicolin (10 µM) to samples mounted on a temperature- and CO_2-controlled

microscope stage. 3D time-lapse live cell imaging was carried out on a Leica SP8 laser scanning

confocal microscopes, using a C-Apo 63x/1.4NA oil-immersion objective. Tile-scanning was

employed to image multiple (4x4) fields of view. Live-cell imaging was performed starting from

0.5 – 16 hpi by acquiring 9-11 Z-stacks spaced by 0.8 μm every 2.5 min. The Adaptive Focus

control (AFC, Leica) was utilized to correct for axial drift. Images were collected at 512 x 512

frame size, at 180 nm pixel sizes (2x digital zoom) with 1.54 μs pixel dwell times and highly

attenuated 488 and 633 nm laser lines that respectively excite the INmNG and emiRFP670-Lamin

B1 fluorescent markers and their respective emission was collected between 502-560 nm

(INmNG) and 645-700 nm (emiRFP670-Lamin) using GaSP-HyD detectors. For assessing HIV-

1 nuclear import in a fixed time-point (8 hpi), stringent imaging conditions were used compared

to those used for long-term time-lapse imaging (2x laser power, with 0.12 μm/pixel in X-Y and

0.3 or 0.5 μm/pixel in Z). 3D-image series were processed off-line using ICY image analysis

software (http://icy.bioimageanalysis.org/) [53].

Image analyses

For assessing HIV-1 nuclear import in fixed time-point images, 3D z-stack images were analyzed

using an in-house script in the ICY protocols module, essentially as described in (PMID:

32665593). Briefly, the nuclear volume in three dimensions was detected using the lamin intensity

by the HK-means and the Connected Components plugins in ICY [53]. The obtained 3D ROI

corresponding to the nuclear volume was shrunk by 0.5 μm in X-Y-Z using an ROI-erosion plugin.

HIV-1 complexes detected within the eroded ROI were considered as nuclear spots.

To track HIV-1 docking at the NE, we extracted >10 cells from each independent extended 0.5 -

16 h imaging experiment. Cells were selected based on their minimal lateral movement during the

imaging time-course. The central z-planes (2x planes, 1.6 um) of the nucleus were projected to a

2D-stack and then analyzed. Additionally, to overcome hurdles associated with single virus

tracking in a crowded environment the time series was cropped between 4-12h. The 4h initial

timepoint was chosen based on prior observations for HIV-1 attrition in the cytoplasm [13], and the

12h later time point was chosen to overcome extensive cellular motion and deformation during

late imaging time points. The nuclear motion was additionally drift corrected using Fast4Dreg in

Fiji [54]. Drift corrected 2D movies (between 4 -12 hpi) was later processed in ICY bio-image

analysis single particle tracking suite. Images were denoised using gaussian filters, and the

laminB1 fluorescence signals was used to mask the pixels occupied by the NE as a region of

interest. The HIV-1 INmNG puncta that were detected within this mask was tracked for each

individual nucleus. Single HIV-1 trajectories coinciding with the NE mask for >2 frames (>5 min)

were considered as NE-interactions. Each track was manually inspected to correspond to true viral

trajectories and segments of each individual track was analyzed for docking behavior. The

segments of tracks when a single HIV-1 particle remained within a 2-pixel (360 nm) radius for 3

or more frames (>7.5 min) was determined by a rolling average of particle displacement, similar

to cage diameter analysis. These segments were considered as interactions representative of HIV-

1 core docking at the NE. The 360nm radius was used taking into consideration the lateral

diffraction limited confocal resolution and the pixel sizes in our time-lapse movies, which is

250nm and 180nm, respectively. These stringent conditions allowed us to robustly define virus

docking and distinguish them from transient interactions within single HIV-1 trajectories along

the NE, that failed to remain at a given location for more than 5 minutes. These stable docking

events, as a fraction of total NE-interactions (total number of HIV-1 tracks, including track

segments that was not localized within the 360nm window) obtained in each individual nuclei over

the 8h time-window was analyzed and plotted. As controls, the interactions between bald

endosome trapped HIV-1 (no-VSV) infections were analyzed and the background in detecting

non-specific overlaps as HIV/NE-docking was estimated to be 4.2%.

Author Contribution

A.D., A.B., J.R.A.M, J.S., J.H., C.T., and C.A. performed research and analyzed data, A.D., A.B., S.H., J.R.A.M., and A.C.F analyzed data, C.A, J.P., A.C.F., and I.R. designed research and wrote the paper.

Acknowledgments

We thank Harrison Bracey and Ashley Fulmer for technical assistance and Alan Engelman for helpful comments. This work was supported in part by the Israel Science Foundation grant 418/21 (Rousso Lab), by NIH R01-AI157843 and U54-AI170791 (Aiken and Perilla Labs), NIH U54 AI170855 and R01-AI148382 (Francis lab). C.T. received support from NIH 5R25AI164610 and a Patrick and Eloise E. Wieland Fellowship. This work used Delta at NCSA and Expanse at SDSC through allocation MCB-170096 from the Advanced Cyberinfrastructure Coordination Ecosystem: Services & Support (ACCESS) program, which is supported by National Science Foundation awards #2138259, #2138286, #2138307, #2137603, and #2138296.

Data availability

The data that support the findings of this study are available in the manuscript and its Supplementary Information and from the corresponding authors on reasonable request.

References

1   Lee, K. *et al.* Flexible use of nuclear import pathways by HIV-1. *Cell Host Microbe* **7**, 221-
233, (2010).

2   Shen, Q., Wu, C., Freniere, C., Tripler, T. N. & Xiong, Y. Nuclear Import of HIV-1.
Viruses **13**, (2021).

3   Li, C., Burdick, R. C., Nagashima, K., Hu, W. S. & Pathak, V. K. HIV-1 cores retain their
integrity until minutes before uncoating in the nucleus. *Proc Natl Acad Sci U S A* **118**,
(2021).

4   Zila, V. *et al.* Cone-shaped HIV-1 capsids are transported through intact nuclear pores.
Cell **184**, 1032-1046 e1018, (2021).

5   Zila, V., Muller, T. G., Muller, B. & Krausslich, H. G. HIV-1 capsid is the key orchestrator
of early viral replication. *PLoS Pathog* **17**, e1010109, (2021).

6   Bejarano, D. A. *et al.* HIV-1 nuclear import in macrophages is regulated by CPSF6-capsid
interactions at the nuclear pore complex. *Elife* **8**, (2019).

7   Dharan, A. & Campbell, E. M. Teaching old dogmas new tricks: recent insights into the
nuclear import of HIV-1. *Curr Opin Virol* **53**, 101203, (2022).

8   Sarkar, S. *et al.* Tough Way In, Tough Way Out: The Complex Interplay of Host and Viral
Factors in Nucleocytoplasmic Trafficking during HIV-1 Infection. *Viruses* **14**, (2022).

9   Achuthan, V. *et al.* Capsid-CPSF6 Interaction Licenses Nuclear HIV-1 Trafficking to Sites
of Viral DNA Integration. *Cell Host Microbe* **24**, 392-404 e398, (2018).

10 Burdick, R. C. *et al.* Dynamics and regulation of nuclear import and nuclear movements
of HIV-1 complexes. *PLoS Pathog* **13**, e1006570, (2017).

11 Di Nunzio, F. *et al.* Human nucleoporins promote HIV-1 docking at the nuclear pore,
nuclear import and integration. *PLoS One* **7**, e46037, (2012).

12 Di Nunzio, F. *et al.* Nup153 and Nup98 bind the HIV-1 core and contribute to the early
steps of HIV-1 replication. *Virology* **440**, 8-18, (2013).

13 Francis, A. C. & Melikyan, G. B. Single HIV-1 Imaging Reveals Progression of Infection
through CA-Dependent Steps of Docking at the Nuclear Pore, Uncoating, and Nuclear
Transport. *Cell Host Microbe* **23**, 536-548 e536, (2018).

14 Matreyek, K. A. & Engelman, A. The requirement for nucleoporin NUP153 during human
immunodeficiency virus type 1 infection is determined by the viral capsid. *J Virol* **85**,
7818-7827, (2011).

15 Muller, T. G. *et al.* HIV-1 uncoating by release of viral cDNA from capsid-like structures
in the nucleus of infected cells. *Elife* **10**, (2021).

16 Schaller, T. *et al.* HIV-1 capsid-cyclophilin interactions determine nuclear import pathway,
integration targeting and replication efficiency. *PLoS Pathog* **7**, e1002439, (2011).

17 Blair, W. S. *et al.* HIV capsid is a tractable target for small molecule therapeutic
intervention. *PLoS Pathog* **6**, e1001220, (2010).

18 Tse, W. *et al.* in *Conference on Retroviruses and Opportunistic Infections.* 13-16 (Seattle
Washington).

19 Bester, S. M. *et al.* Structural and Mechanistic Bases of Viral Resistance to HIV-1 Capsid
Inhibitor Lenacapavir. *mBio* **13**, e0180422, (2022).

20 Bester, S. M. *et al.* Structural and mechanistic bases for a potent HIV-1 capsid inhibitor.
Science **370**, 360-364, (2020).

21   Dvory-Sobol, H., Shaik, N., Callebaut, C. & Rhee, M. S. Lenacapavir: a first-in-class HIV-

1 capsid inhibitor. *Curr Opin HIV AIDS* **17**, 15-21, (2022).

22   Link, J. O. *et al.* Clinical targeting of HIV capsid protein with a long-acting small molecule.

Nature **584**, 614-618, (2020).

23   Peng, K. *et al.* Quantitative microscopy of functional HIV post-entry complexes reveals

association of replication with the viral capsid. *Elife* **3**, e04114, (2014).

24   Price, A. J. *et al.* Host cofactors and pharmacologic ligands share an essential interface in

HIV-1 capsid that is lost upon disassembly. *PLoS Pathog* **10**, e1004459, (2014).

25   Rankovic, S., Ramalho, R., Aiken, C. & Rousso, I. PF74 Reinforces the HIV-1 Capsid To

Impair Reverse Transcription-Induced Uncoating. *J Virol* **92**, (2018).

26   Saito, A. *et al.* Roles of Capsid-Interacting Host Factors in Multimodal Inhibition of HIV-

1 by PF74. *J Virol* **90**, 5808-5823, (2016).

27   Yant, S. R. *et al.* A highly potent long-acting small-molecule HIV-1 capsid inhibitor with

efficacy in a humanized mouse model. *Nat Med* **25**, 1377-1384, (2019).

28   Forshey, B. M., von Schwedler, U., Sundquist, W. I. & Aiken, C. Formation of a human

immunodeficiency virus type 1 core of optimal stability is crucial for viral replication. *J

Virol* **76**, 5667-5677, (2002).

29   Yang, R. *et al.* Second-site suppressors of HIV-1 capsid mutations: restoration of

intracellular activities without correction of intrinsic capsid stability defects. *Retrovirology*

9, 30, (2012).

30   Shi, J., Zhou, J., Shah, V. B., Aiken, C. & Whitby, K. Small-molecule inhibition of human

immunodeficiency virus type 1 infection by virus capsid destabilization. *J Virol* **85**, 542-

549, (2011).

31 Obr, M., Schur, F. K. M. & Dick, R. A. A Structural Perspective of the Role of IP6 in

Immature and Mature Retroviral Assembly. *Viruses* **13**, (2021).

32 Shen, Q. *et al.* The capsid lattice engages a bipartite NUP153 motif to mediate nuclear

entry of HIV-1 cores. *Proc Natl Acad Sci U S A* **120**, e2202815120, (2023).

33 Dismuke, D. J. & Aiken, C. Evidence for a functional link between uncoating of the human

immunodeficiency virus type 1 core and nuclear import of the viral preintegration complex.

J Virol **80**, 3712-3720, (2006).

34 Dominguez-Zotes, S., Fuertes, M. A., Rodriguez-Huete, A., Valbuena, A. & Mateu, M. G.

A Genetically Engineered, Chain Mail-Like Nanostructured Protein Material with

Increased Fatigue Resistance and Enhanced Self-Healing. *Small* **18**, e2105456, (2022).

35 Dominguez-Zotes, S., Valbuena, A. & Mateu, M. G. Antiviral compounds modulate

elasticity, strength and material fatigue of a virus capsid framework. *Biophys J* **121**, 919-

931, (2022).

36 Klug, W. S. *et al.* Failure of viral shells. *Phys Rev Lett* **97**, 228101, (2006).

37 Mateu, M. G. Mechanical properties of viruses analyzed by atomic force microscopy: a

virological perspective. *Virus Res* **168**, 1-22, (2012).

38 Rankovic, S., Varadarajan, J., Ramalho, R., Aiken, C. & Rousso, I. Reverse Transcription

Mechanically Initiates HIV-1 Capsid Disassembly. *J Virol* **91**, (2017).

39 Yamashita, M., Perez, O., Hope, T. J. & Emerman, M. Evidence for direct involvement of

the capsid protein in HIV infection of nondividing cells. *PLoS Pathog* **3**, 1502-1510,

(2007).

40 Siddiqui, M. A. *et al.* A Novel Phenotype Links HIV-1 Capsid Stability to cGAS-Mediated

DNA Sensing. *J Virol* **93**, (2019).

41 Zimmerli, C. E. *et al.* Nuclear pores dilate and constrict in cellulo. *Science* **374**, eabd9776,
(2021).

42 Ramalho, R., Rankovic, S., Zhou, J., Aiken, C. & Rousso, I. Analysis of the mechanical
properties of wild type and hyperstable mutants of the HIV-1 capsid. *Retrovirology* **13**, 17,
(2016).

43 He, J. *et al.* CCR3 and CCR5 are co-receptors for HIV-1 infection of microglia. *Nature*
385, 645-649, (1997).

44 Yee, J. K. *et al.* A general method for the generation of high-titer, pantropic retroviral
vectors: highly efficient infection of primary hepatocytes. *Proc Natl Acad Sci U S A* **91**,
9564-9568, (1994).

45 Wehrly, K. & Chesebro, B. p24 antigen capture assay for quantification of human
immunodeficiency virus using readily available inexpensive reagents. *Methods* **12**, 288-
293, (1997).

46 Liu, X. & Harada, S. DNA isolation from mammalian samples. *Curr Protoc Mol Biol*
Chapter 2, Unit2 14, (2013).

47 Bryer, A. J. & Perilla, J. R. Performance efficient macromolecular mechanics via sub-
nanometer shape based coarse graining. *Nat Commun*, 2014, (2023).

48 Ni, T. *et al.* Structure of native HIV-1 cores and their interactions with IP6 and CypA. *Sci*
Adv **7**, (2021).

49 Phillips, J. C. *et al.* Scalable molecular dynamics on CPU and GPU architectures with
NAMD. *J Chem Phys* **153**, (2020).

50 Essmann, U. *et al.* A Smooth Particle Mesh Ewald Method. *J Chem Phys* **103**, 8577-8593,
(1995).

51   Fernandez, D. P., Mulev, Y., Goodwin, A. R. H. & Sengers, J. M. H. L. A Database for the

550   Static Dielectric-Constant of Water and Steam. *J Phys Chem Ref Data* **24**, 33-69, (1995).

551   52   Francis, A. C. *et al.* HIV-1 replication complexes accumulate in nuclear speckles and

552   integrate into speckle-associated genomic domains. *Nat Commun* **11**, 3505, (2020).

553   53   de Chaumont, F. *et al.* Icy: an open bioimage informatics platform for extended

554   reproducible research. *Nat Methods* **9**, 690-696, (2012).

555   54   Laine, R. F. *et al.* NanoJ: a high-performance open-source super-resolution microscopy

556   toolbox. *J Phys D Appl Phys* **52**, 163001, (2019).

Figure Lengends

Fig. 1: DNA synthesis, nuclear entry, infectivity and averaged measured point stiffness of isolated HIV-1 cores. (A) Relative single-cycle infectivity of wild type and CA mutant viruses. (B) Nuclear entry efficiency of CA mutants relative to WT HIV-1. Value represent the ratio of 2-LTR circle levels to late reverse transcripts. (C) Ratio of infection of control Hela cells vs. aphidicolin-treated cells by WT and mutant viruses. (D) Relative levels of late reverse transcripts in target cells. (E) Fold reduction of infection in Nup153-depleted cells vs. control cells. For A-E, values shown are the mean of three independent experiments, with error bars representing the standard error of the mean. (F) Stiffness measurements of purified HIV-1 cores. Each stiffness value was calculated as the average of ~480 force–distance curves obtained from individual cores. Measurements were conducted in the presence of inositol hexakisphosphate (IP6; 100 µM) except for the first column. The t-test analysis revealed that differences between the stiffness values of the various samples and WT are statistically significant (p values: <0.001 and <0.05 for WT+IP6+PF74 and E45A+IP6, respectively, <0.0001 for the remaining samples). Error bars in all panels represent the standard error of the mean.

Fig. 2: Fluorescence microscopy imaging of HIV-1 nuclear entry. (A) Representative single z-stack images and (B) quantification of HIV-1 INmNG puncta nuclear entry in TZM-bl cells acquired at 8 hours post-infection. The number of fluorescent INmNG puncta (fHIV-1, green) inside 0.5 um of the nuclear membrane (Lamin, blue) was scored from 30 nuclei in individual experiment from each of 3 independent experiments. Scale in (A) is 5 um.

Fig. 3: Experimental and simulated AFM nanoindentation of wild type IP6-treated cores. (A)

Representative force–distance (FD) curves where force loading (black) and unloading (red) curves

are shown. The maximal loading force was set to 5 nN (n=23), 10 nN (n=24), and 20 nN (n=25).

Despite the application of high maximal loading forces, none of the curves exhibited any indication

of structural failure in the core. In one core (of the 72 examined), we observed an abrupt drop in

the loading curve (inset). (B-E) Simulated FD curves for nanoindentation of coarse-grained HIV-

1 capsids, testing the effect of probe location, ultimately capsid curvature, on measured material

properties. The plots show raw data in gray, and a windowed-average in blue. The inset volume

cross-sections show the deformed capsid morphologies and provide a sense of global deformation.

The solid black line shows a linear fit of the FD curve for the first 3 nm of indentation, which was

used to derive the stiffness values given to the right of each plot. Additionally, snapshots from

these nanoindentation simulations are provided on the right, with all components visualized:

spherical AFM probes, capsids, and baseplates. (B) Probe located at the large end of the conical

capsid. Note the global deformation, narrowing, of the cone in the inset graphic. (C) and (D) show

probe locations in the relatively flat, mid-regions of the capsid. (E) Shows nanoindentation of the

narrow, most highly curved region of the capsid.

Fig. 4: Volume recovery of the HIV core following compression. (A) Topographic AFM images

of an isolated IP6-treated WT core before (upper image) and after (lower image) extreme

compression. Both images were acquired using the QI mode at a maximal loading force of 300

pN. Subsequent measurements focused on a selected central region, as marked by a dashed yellow

rectangle. Scale bars are 50 nm. (B) Representative topographic AFM images (left panels) and the

corresponding Matlab plots (right panels) for the selected central region of an isolated IP6-treated

WT core before (top), during (middle), and immediately after (bottom) compression by a 5 nN

loading force. The same region was repeatedly scanned and imaged over time in QI mode at

maximal loading force of 300 pN. (C) Two representative volume recovery trajectories (of the 36

trajectories obtained). The volume beneath the selected region was calculated using MatLab and

plotted as a function of time. The initial volume was set as 100%, and all other measurements were

normalized accordingly. The upper curve shows nearly full volume recovery, whereas the lower

curve shows 80% recovery, which was the minimum volume recovery percentage we detected.

(D) The average volume recovery of the 36 analyzed cores. The average compressed volume was

39% (pressed). The average volume upon reducing the force back to 300 pN was 92% (immediate).

Several minutes (5–7 min) later, the recovery volume remained unchanged at 92% (late). T-test

analysis revealed that the difference between the compressed and immediate or late recovered

volume is significant (p value <0.0001). Error bars represent the standard error of the mean.

Fig. 5: The effect of exposure to a high loading force on the structural integrity of cores. (A)

Topographic AFM images of two representative WT cores before and after compression. Images

were acquired using the QI mode operated at a 300 pN loading force. Typical cone-shaped cores

are observed. A cross-section height profile along the length of the core is displayed. After

compression, most cores (~86%) remained intact. However, those that broke exhibited an opening

at the narrow end of the core, distal from the compressed region. Scale bars are 50 nm. (B) The

percentage of broken wild-type (WT) and mutant cores. Cores were examined in the presence of

IP6 (100 μM). WT cores were also treated with the capsid-inhibiting antiviral drugs PF74 (1.25

μM) or Lenacapavir (LCV; 500 pM). The number of cores analyzed is indicated inside each bar.

Core breakage was determined by AFM imaging of the cores after they were compressed by a

loading force of 5 nN. Error bars represent standard error of the mean. (C) Model depicting the

requirement for core elasticity in HIV-1 nuclear entry. Partially reverse transcribed cores interact

with the external face of the NPC and undergo reversible deformation to insert into the pore.

Capsid interactions with Nup153 promote passage through the pore. (D) Quantification of WT and

mutant HIV-1 docking at the nuclear envelope (NE) in TZM-bl cells. The background level of NE

interactions is shown as a blue dashed line, determined from endosome trapped bald-HIV-1 (no-

VSV) infections that cannot engage NPCs. Data represent mean values acquired from 10 nuclei,

and >600 tracks for each condition. Statistical significance was determined using non-parametric

Mann-Whitney Rank Sum Test. Error bars represent standard error of the mean.

Figures

Figure 1

652   Figure 2

654   Figure 3

656   Figure 4

660   Figure 5