RESEARCH PAPER

MpmiR319 promotes gemma/gemma cup formation in the liverwort *Marchantia*

polymorpha

Authors: Kazutaka Futagami[1], Masayuki Tsuzuki[1,2] [*], Yuichiro Watanabe[1] [*]

[1] Department of Life Sciences, Graduate School of Arts and Sciences, The University of

Tokyo, Tokyo 153-8902, Japan

[2] Faculty of Agriculture and Marine Sciences, Graduate School of Integrated Arts and

Sciences, Kochi University, Nankoku, Kochi 783-8502, Japan

* Corresponding authors

E-mail: ft.kz.m9m@gmail.com (K. Futagami)

mtsuzuki@kochi-u.ac.jp (M. Tsuzuki)

cyuiwat@g.ecc.u-tokyo.ac.jp (Y. Watanabe)

Highlight

This study demonstrates Mp*miR319* expression affects gemma cup and gemma formation

during *Marchantia polymorpha* vegetative propagation by suppressing its target mRNAs.

ORCID

Kazutaka Futagami: https://orcid.org/0000-0002-3321-6508

Masayuki Tsuzuki: https://orcid.org/0000-0002-0642-1143

Yuichiro Watanabe: https://orcid.org/0000-0002-7139-4903

Abstract

After microRNAs (miRNAs) were initially detected as approximately 21-nt sequences in plants, miRNA-mediated regulation of target gene expression was characterized in many cases. The sequences of miR159/319 family miRNAs are highly conserved and their development-related roles have been comprehensively characterized in dicot and monocot plants. However, their effects in basal land plants remain relatively unknown. We used a CRISPR/Cas9 system to edit genome sequences at Mp*MIR319a* and/or Mp*MIR319b* loci. Mutant *mir319b* lines developed relatively few gemma cups, while mutant *mir319a* lines produced relatively few gemmae in each gemma cup. Earlier 5′-RACE or degradome analyses suggested that Mp*miR319* targets Mp*RKD* and Mp*R2R3-MYB21* transcripts and suppresses expression. To determine whether these candidate genes influence the *mir319a/mir319b* phenotype we constructed miR319-resistant Mp*RKD* (mMp*RKD*) and Mp*R2R3-MYB21* (mMp*R2R3-MYB21*) by decreasing the complementarity to miR319. We also introduced the genes into wild-type liverwort and screened for phenotypic changes. We observed that mMp*RKD* resulted in gemma/gemma cup-less liverwort mutants, but mMp*R2R3-MYB21* did not. No major adverse effects were detected during male and female reproductive development induced by far-red illumination. These results indicate that miR319 influences gemma formation mainly by repressing Mp*RKD* expression rather than Mp*R2R3-MYB21* expression.

Keywords: liverwort, *Marchantia polymorpha*, gemma, gemma cup, notch, microRNA, RKD transcription factor, R2R3MYB transcription factor

 Introduction

The complementarity to their target mRNA sequences is generally greater for plant

microRNAs (miRNAs) than for animal miRNAs (Rhoades et al., 2002)(Lin and Bowman,

2018). Each plant miRNA can target a limited number of mRNAs. Accordingly, mRNA(s)

targeted by each miRNA can be predicted by searching genome sequence data for a

sequence complementary to the miRNA sequence of interest. Recent studies revealed that

land plants have 300–500 lineage-specific diverse miRNAs (Axtell et al., 2007)(Lin and

Bowman, 2018)(Pietrykowska et al., 2022). Earlier *in vivo* studies on miRNAs conserved

among land plants showed that their target mRNAs are transcribed from highly conserved

cognate genes across phyla (Axtell et al., 2007)(Tsuzuki et al., 2019)(Alaba et al.,

2015)(Pietrykowska et al., 2022). Such conserved miRNA–target mRNA modules may

form a core set of functionally and biologically important gene regulatory circuits in land

plants. Moreover, as few as seven miRNA families are highly conserved from the

liverwort *Marchantia polymorpha* and the moss *Physcomitrella patens* to angiosperms

(Axtell et al., 2007)(Tsuzuki et al., 2016)(Lin et al., 2016b)(Pietrykowska et al., 2022).

However, most liverwort miRNAs have not been studied sufficiently to determine how

common or specific they are among land plants. Additional experimental data are needed

to substantiate the biological significance of highly conserved miRNAs in basal land

plants. Among conserved miRNAs, miR159/319 family members are unique because

their target mRNA species may have changed during phylogenetic diversification

(Palatnik et al., 2007). In Arabidopsis, miR319 preferentially targets mRNAs of

TEOSINTE BRANCHED1, *CYCLOIDEA*, and *PROLIFERATING CELL FACTOR* genes

(Martín-Trillo and Cubas, 2010)(Kosugi and Ohashi, 1997). These genes encode basic-

helix-loop-helix transcription factors that redundantly affect cotyledon boundary

specification and leaf serration (Palatnik et al., 2003)(Koyama et al., 2017).

Marchantia polymorpha is a liverwort that is a valuable emerging model species used

in studies on land plant diversification, which have generated increasing amounts of

information and resources for comparative evolution research (Bowman et al., 2017). In

M. polymorpha, Mp*miR319* targets mRNAs of *MpR2R3-MYB21* (Arabidopsis MYB33-

like transcription factor belonging to the GAMYB family) (Lin et al., 2016b)(Tsuzuki et

al., 2016) and RWP-RK DOMAIN (RKD)-containing transcription factor genes (Lin et

al., 2016b) according to a 5′-RACE or degradome analysis. Thus, Mp*miR319* may have

some regulatory effects on such candidate target genes, but this possibility remains to be

experimentally verified.

Materials and methods

Plant materials and growth conditions

Takaragaike-1 (Tak-1, male) and Takaragaike-2 (Tak-2, female) were used as wild-type

M. polymorpha samples (Ishizaki et al., 2008). Plants were grown asexually on half-

strength Gamborg's B5 medium (FUJIFILM Wako Pure Chemical Co., Osaka, Japan)

(Gamborg et al., 1968) containing 0.5 g/l 2-morpholinoethanesulfonic acid (MES, Tokyo

Chemical industry Co., Tokyo) and 1% (w/v) agar (STAR Agar L-grade, RIKAKEN,

Tokyo) under continuous white fluorescent light at 22 °C. To induce the sexual

reproductive phase, plants were grown at 22 °C under continuous light conditions

provided by a mixture of white fluorescent light and far-red (FR) light-emitting diodes

(OSXXXXT3C1E, OptoSupply, Hong Kong; peak emission at 730 nm, with a full width

at half-maximum of 20 nm).

Transformation and mutant selection

Regenerating thalli (for *mir319a* or *mir319b* mutants) (Kubota et al., 2013) and a G-

AgarTrap method (to generate gMp*RKD*, mMp*RKD*, gMp*MYB21*, and mMp*MYB21*

lines) (Tsuboyama and Kodama, 2018) were used to transform *M. polymorpha*. Briefly,

transformed plants were selected on half-strength Gamborg's B5 medium containing 0.5

g/l MES, 1% (w/v) agar, 100 µg/ml cefotaxime and 10 µg/ml hygromycin (for

pMpGWB101 and pMpGE010), or 100 µg/ml cefotaxime and 0.5 µM chlorsulfuron (for

pMpGWB301 and pMpGE011). After screening T_1 plants on drug-supplemented media,

we obtained approximately 10 candidate lines for each guide RNA (gRNA) in the male

and female genetic backgrounds.

Genomic DNA extraction

Genomic DNA was extracted from Tak-1 thalli using the DNeasy Plant Mini kit (Qiagen,

Hilden, Germany) according to the manufacturer's recommendations.

Plasmid construction for the CRISPR/Cas9 system

We employed the CasFinder Tool (https://marchantia.info/tools/casfinder/) to design

gRNAs to target Mp*miR319a* and/or Mp*miR319b* loci. The pMpGE_En03 vector

(Sugano et al., 2018) was used to clone oligonucleotide pairs to express gRNAs:

pMpGE_En03-gRNA1,   pMpGE_En03-gRNA3,   and   pMpGE_En03-gRNA4.

pMpGE010 and pMpGE011 were used for Cas9 expression (Sugano et al., 2018).

The *mir319a^{ge}*, *mir319b^{ge}*, and *mir319ab^{ge}* mutants were selected as described

previously (Tsuzuki et al., 2019). Using genomic DNA as the template, *MIR319A* and

MIR319B sequences were amplified by PCR using PrimeSTAR HS DNA polymerase

(TaKaRa Bio, Kusatsu, Japan) and the primer pairs MpMIR319aF1 and MpMIR319a-R1

or MpMIR319bF2 and MpMIR319b-R2. Amplified products were sequenced according

to the Sanger method. Details regarding the PCR and genomic sequencing primers are

listed in Table S1. Mp*MIR319A* and Mp*MIR319B* genomic sequences in *mir319a^{ge}*,

mir319b^{ge}, and *mir319ab^{ge}* mutant plants (T_1 generation) are provided in Figure 1.

Construction of miR319-resistant Mp*RKD* and Mp*R2R3-MYB21* transgenes

To obtain the wild-type *RKD* gene segment, the genomic sequence from the Mp*RKD*

promoter region (3,827 bp upstream of the translation initiation codon) to the 3′

untranslated region (UTR) (568 bp downstream of the translation termination codon) was

amplified by PCR using Tak-1 genomic DNA, PrimeSTAR HS DNA polymerase

(TaKaRa Bio), and primers IF-Hind-pRKD-F and Inf-gMpRKD-Sal-R. To create the

miR319-resistant Mp*RKD* gene segment, two PCR fragments were amplified using Tak-

1 genomic DNA as well as the IF-Hind-pRKD-F and mRWP_RK-R primer set and the

mRWP_RK-F and Inf-gMpRKD-Sal-R primer set to specifically mutate the miR319

target sequence in the Mp*RKD* 3′ UTR sequence. The In-Fusion Cloning kit (TaKaRa

Bio) was used to insert the PCR fragments into the pMpGWB300 vector that had been

cleaved by *Hin*dIII and *Sal*I to generate the pMpGWB300:proMpRKD:gMpRKD and

pMpGWB300:proMpRKD:mMpRKD recombinant plasmids.

To obtain the wild-type *MYB21* gene segment, the Mp*MYB21* promoter region (3008 bp

upstream of the translation initiation codon) to the initiation codon, was first amplified

by PCR using Tak-1 genomic DNA, PrimeSTAR HS DNA polymerase, and primers

Hind-pMpMYB21-3000-F and Sal-pMpMYB21-R, cloned into pMpGWB100 after

both being restricted with *Hin*dIII and *Sal*I to obtain pMpGWB100-pMYB21. Wild-type

MYB21 gene segment (gMYB21) from halfway of promoter region to the termination

codon (4244 bp) were amplified using Tak-1 genomic DNA and primers   pMYB33-

459F and Bam-MpMYB33-R2. To create the miR319-resistant Mp*MYB21* gene segment,

two PCR fragments were amplified using gMYB21 fragment as well as the pMYB33-

459F and mMYB33-R set and the mMYB33-F and Bam-MpMYB33-R2 primer

set to specifically mutate the miR319 target sequence in the Mp*MYB21*

coding sequence. The obtained PCR fragments were mixed and subjected to

the next PCR reaction using primers pMYB33-459F and Bam-MpMYB33-R2

to obtain miR319 resistant MYB21 gene body fragment (mMYB21). The gMYB21 and

mMYB21 fragments were cloned separately into the pMpGWB100 vector, after being

compatibly restricted with *Nru*I and *Bam*HI, to generate the pMpGWB100:pgMp*MYB21*

and pMpGWB100:pmMp*MYB21*.

M. polymorpha was transformed with the recombinant plasmids as described previously

(Tsuboyama et al., 2018a)(Tsuboyama and Kodama, 2018).

Results

Guide RNA-mediated editing of *MIR319* genes

A previous study revealed *M. polymorpha* has a relatively simple genome because of limited gene redundancy (Bowman et al., 2017). *M. polymorpha* has two *MIR319* coding loci, both of which produce an identical mature miRNA sequence (5′-$_p$CUUGGACUGAAGGGAGCUCCC$_{OH}$-3′) (Lin et al., 2016b; Tsuzuki et al., 2016). The two *MIR319* loci have been annotated (*M. polymorpha* genome ver.6.1) as Mp*MIR319A* (Mp3g08965) and Mp*MIR319B* (Mp4g01845) (Fig. 1A).

We focused on whether Mp*MIR319a* and/or Mp*MIR319b* loci are functional. To analyze the possible biological functions of miR319 in *M. polymorpha*, we edited two annotated Mp*miR319* coding sequences to obtain mutants, which were examined for phenotypic changes. Specifically, we designed gRNA#1 for *MIR319A* (Fig. 1B) and gRNA#3 and #4 for *MIR319B* (Fig. 1C) and constructed vectors to introduce indel mutations at the respective *MIR319* loci using a CRISPR/Cas9 genome-editing system (Sugano et al., 2014)(Sugano et al., 2018). We adopted an *Agrobacterium*-mediated transformation protocol to introduce the vectors into Tak-1 thalli to express the gRNAs and Cas9 (Sugano et al., 2014)(Tsuboyama et al., 2018b). After the drug-based selection of G$_1$ plants, we amplified the targeted genomic sequences from selected G$_1$ thalli and analyzed the PCR products via Sanger sequencing to confirm indel mutations were present at the target loci. Multiple lines with indel mutations at the Mp*MIR319a* or Mp*MIR319b* locus were generated. Several representative lines with similar phenotypes were selected (Fig. 1B

and C) for the subsequent analysis. The selected mutants with indel mutations at the

Mp*mir319a* locus were designated as *mir319a*$^{ge-1}$ *a* #5 and *mir319a*$^{ge-1}$ *a* #10, whereas

the selected mutants with indel mutations at the Mp*mir319b* locus were designated as

mir319b^{ge-3} *b* #11, *mir319b*$^{ge-3}$ *b* #15, *mir319b*$^{ge-4}$ *b* #6, and *mir319b*$^{ge-4}$ *b* #9.

***miR319a* and *miR319b* mutant thalli exhibited abnormal gemma cup production**

Although the *mir319a*^{ge} and *mir319b*^{ge} mutant plants grew as fast as the wild-type Tak-1

plants, they produced gemma cups less frequently than Tak-1 plants (Fig. 2). More

specifically, compared with wild-type Tak-1 plants, significantly fewer *mir319a*$^{ge-1}$ (*a* #5

and *a* #10) and *mir319b*$^{ge-3}$ (*b* #11 and *b* #15) plants produced gemma cups

(approximately half as many). Gemma cup production was most inhibited in the

mir319b^{ge-4} (*b* #6 and *b* #9) plants. We speculated that both *mir319a* and *miR319b* alleles

somehow affect gemma cup formation during dorsal organ formation in thalli. In addition,

mir319a^{ge-1} (*a* #5 and *a*#10) mutants produced fewer gemmae in each gemma cup than

the wild-type plants (Fig. 3), whereas the *mir319b*$^{ge-3}$ (*b* #11 and *b* #15) mutants and the

wild-type plants did not differ in terms of the number of gemmae in each gemma cup (Fig.

3). These observations suggest that miR319 contributes to the formation of gemmae and

gemma cups in *M. polymorpha* by regulating target gene expression.

We also created similar *mir319a*^{ge} and *mir319b*^{ge} mutants using female Tak-2 plants. A

decrease in gemma cup formation was also detected in the Tak-2-based mutants (data not

shown). Thus, we focused mainly on lines with a Tak-1 genetic background to study the

miR319 function during vegetative growth stages.

Knockout of both *miR319a* and *miR319b* loci did not result in additive defects

The *mir319a* and *mir319b* knockout lines exhibited similar phenotypic defects in terms of the production of gemma cups/gemmae. We analyzed whether the two loci additively affect liverwort vegetative organ formation by creating and characterizing double knockout mutants. We were unable to obtain a double mutant via a single transformation involving two vectors. Thus, we targeted the *MIR319a* locus of *mir319b*^{ge-3} *b* #15 with pMpGE_En03-gRNA1. After several trials, we obtained a double knockout line that had a 5-base deletion in the *MIR319a* locus identical to *mir319a*^{ge-1} *a* #10 in addition to a 1-base addition in the *MIR319b* locus (Fig. 1D). For brevity, the mutant line was named *miR319ab #10/15* (Fig. 1D). In terms of the number of developed gemma cups and gemmae, the *miR319ab #10/15* mutant phenotype did not differ from the *miR319a* and *mir319b* mutant phenotypes (Fig. 2, *ab* #10/15).

Regulation of Mp*RKD* by miR319 influences gemma cup formation

An earlier degradome analysis indicated that Mp*RKD* (formerly *RWP-RK DOMAIN-CONTAINING 1*) and Mp*MYB33* mRNAs are targeted by Mp*miR319* (Lin et al., 2016a). In the current study, we assessed whether the susceptibility of either or both of these mRNA genes to miR319 affects gemma/gemma cup formation. Mp*RKD* encodes an RWP-RK domain-containing transcription factor that is an evolutionarily conserved regulator of germ cell differentiation (Koi et al., 2016)(Rövekamp et al., 2016).

Additionally, it was reported that Mp*RKD* is expressed on gemma cup rims (Koi et al.,

2016) and is necessary for normal gemma cup growth (Rövekamp et al., 2016).

Accordingly, we examined whether downregulated Mp*RKD* expression due to miR319 is

linked to gemma/gemma cup formation. We constructed a mutated *RKD* coding sequence

that is unaffected by miR319.

The 21-nt sequence complementary to miR319 started from 4 nt downstream of the stop

codon in the *M. polymorpha RKD* (Mp*RKD*) mRNA sequence. We constructed a miR319-

resistant form of the Mp*RKD* gene (designated as mMp*RKD*; Fig. 4A) by introducing a

12-base substitution to decrease the complementarity to Mp*miR319*. The mMp*RKD* gene

was placed downstream of its own 3.8 kb promoter sequence in the pMpGWB300 vector

(Ishizaki et al., 2015). As a control, we cloned wild-type Mp*RKD* (designated as

gMp*RKD*) in the same vector. The mMp*RKD* gene produces *RKD* mRNA that is

insensitive to miR319, whereas gMp*RKD* mRNA is sensitive to miR319. The gMp*RKD*

and mMp*RKD* genes were introduced into separate Tak-1 wild-type plants to obtain

gMp*RKD* and mMp*RKD* lines. Both gMp*RKD* and mMp*RKD* plants had similar thallus

growth rates, but the mMp*RKD* plants failed to produce gemma cups (Fig. 5: mMpRKD).

The gemma cup-less phenotypes were similar to those of the *mir319a^{ge-1}* or *miR319ab*

#10/15 mutants (Fig. 5; mir319ab #10/15). In contrast, the gMp*RKD* lines produced

gemma cups similar to the wild-type Tak-1 control (Fig. 5; gMpRKD). Therefore, the

sequences complementary to Mp*miR319* in Mp*RKD* mRNA are critical for

gemma/gemma cup formation. Moreover, Mp*RKD* translation is likely suppressed by

Mp*miR319*.

As mentioned earlier, the miR319 target sequence is located in the 3′ UTR downstream

of the *RKD* termination codon sequence. Thus, we designed another miR319-insensitive

RKD gene (Mp*RKD*-Δ3′UTR) by removing the complementary sequences from the 3′

UTR without disturbing *RKD* translation (Fig. S1A). When we introduced Mp*RKD*-

Δ3′UTR into liverwort, thalli were able to grow, but only a few gemma cups developed,

similar to the mMp*RKD* liverwort (Fig. S1B).

Gemma cup/gemma formation was unaffected by miR319-resistant Mp*MYB21*

According to a 5′-RACE or degradome analysis, Mp*MYB21* is also a potential target

gene of miR319 (Tsuzuki et al., 2016)(Lin et al., 2016b). To test whether the regulation

of *MYB21* expression by miR319 affects gemma cup formation, we created a miR319-

resistant Mp*MYB21* (mMp*R2R3-MYB21* or m*MYB21*) that was placed under the control

of its own promoter similar to the Mp*RKD* gene (Fig. 4B). mMp*R2R3-MYB21* was

introduced into wild-type Tak-1 liverwort. The obtained m*MYB21* liverwort lines

produced thalli with gemma cups/gemmae, but the thalli had shapes slightly different

from the wild-type thalli (Fig. 5; mMpR2R3-MYB21). These results indicate that

Mp*R2R3-MYB21* affects gemma cup/gemma formation substantially less than Mp*RKD*

(if at all).

mMp*RKD* plants developed deformed antheridiophores/archegoniophores, but

normal sperm and eggs

It was previously reported that Mp*RKD*, which is expressed in developing eggs and

sperm precursors, regulates egg and sperm formation in *M. polymorpha* (Koi et al.,

2016)(Rövekamp et al., 2016). Thus, we analyzed germ cell, gametangia, and

gametangiophore differentiation (Yamaoka et al., 2021)(Shimamura, 2016) in gMp*RKD*

and mMp*RKD* liverworts. In response to FR illumination, the mMp*RKD* line with the

Tak-1 genetic background developed male gametophytes (Fig. 6C) at nearly the same

time as the gMp*RKD* liverwort (Fig. 6A). In addition, mMp*RKD* antheridiophores had

almost normal stalked receptables, except for their irregular asymmetrical shape (Fig. 6C,

D). Antheridia formed and protruded outward, but they were not surrounded by

antheridiophores in mMp*RKD* lines (Fig. 6E, F). Sperm mobility was similar in the

mMp*RKD*, wild-type Tak-1, and gMp*RKD* liverworts.

We also introduced gMp*RKD* and mMp*RKD* genes into wild-type Tak-2 liverwort and

then checked whether archegonia/archegoniophore development was affected. We

generated transgenic lines using two constructs (Fig. 4A). Under FR light, the mMp*RKD*

liverwort in the Tak-2 genetic background developed female gametophytes as early as the

gMp*RKD* liverwort, but its archegoniophores (Fig. 6I) differed from those of gMp*RKD*

(Fig. 6G) or Tak-2 plants in terms of shape. mMp*RKD* archegoniophores had short digits

(Fig. 6I,J), but produced normal archegonia. Furthermore, in terms of their appearance,

the mMp*RKD* archegonia (Fig. 6K) were almost identical to the archegonia of gMp*RKD*

(Fig. 6H) or wild-type Tak-2 liverwort.

Discussion

When we profiled *M. polymorpha* miRNAs by small RNA-seq analysis, we tried to overexpress miR319 precursor RNAs in wild-type liverwort to clarify their biological functions. The ectopic expression of a miR319 precursor in liverwort adversely altered the development of gemma cups (Tsuzuki et al., 2016). We speculated that the balanced expression of miR319/target mRNAs is necessary for normal gemma cup development in *M. polymorpha*. Our RACE assay detected the cleavage of Mp*MYB21* mRNA, which encodes a homolog of Arabidopsis *MYB33*. An earlier RNA degradome analysis revealed Mp*RKD* (formerly *RWP-RK DOMAIN-CONTAINING 1*) mRNA is also cleaved by MpmiR319 (Lin et al., 2016b). Mp*RKD* was previously investigated using *Marchantia* mutants (Koi et al., 2016)(Rövekamp et al., 2016). These studies showed that Mp*RKD* is essential for normal germ cell differentiation in both male and female gametophytes, but it is also involved in the differentiation of non-reproductive cells surrounding the antheridia, developing egg, and sperm precursors (Koi et al., 2016)(Rövekamp et al., 2016). Mp*RKD* also influences gemma cup formation (Rövekamp et al., 2016). This gene is transcribed at the edge of gemma cups and the meristematic regions of apical notches (Koi et al., 2016)(Rövekamp et al., 2016). It remains unknown whether miR319 organizes gene regulatory networks with Mp*RKD* and further research will clarify.

The gemma cup formation has been histologically characterized. Founder gemmae arise directly from above the center of the apical notch of the thallus, which is followed by the formation of the cup rims surrounding gemmae (Shimamura, 2016). Our findings combined with the available data suggest that Mp*RKD* is involved in various cell

proliferation stages, resulting in the formation of a variety of *M. polymorpha* tissues,

including gemmae/gemma cups. Nevertheless, additional research is needed to clarify

how gemma cups are organized by miR319-mediated gene regulatory networks.

There are several reports of mutants with defective pathways mediating gemma cup

formation in thalli, including the RopGEF KARAPPO (Hiwatashi et al., 2019), GCAM1

(Yasui et al., 2019), and MpKAI2–ligand signaling (Komatsu et al., 2023) pathways.

However, the pathway facilitating the formation of gemma cups from the apical notch has

not been fully determined. Solly *et al.* showed that growth rate heterogeneity determines

the flattened shape of *Marchantia* thalli and that the basic shapes are due to growth rate

differences specified by apical notch activities (Solly et al., 2017). Recently, Romani *et*

al. comprehensively revealed the activities of approximately 450 putative transcription

factor gene promoters by analyzing the respective reporter fusion constructs (Romani et

al., 2024). They generated a map of expression domains for each transcription factor gene

related to gemma growth and identified more than 100 promoters that are active

specifically in the apical notch. The mechanism underlying the formation of gemma cups

may be much more complex than expected.

The *mir319a* and *mir319b* loci differentially contribute to gemma cup formation,

possibly because of some differences in cell-specific transcription or processing

efficiency, but this possibility will need to be tested in future studies. In the moss *P. patens*,

miR159/319, which are encoded in the genome, may target mRNAs for cyclin domain

proteins and two MYB transcription factors (Axtell et al., 2007). Moreover, miR159/319

families are highly conserved from bryophytes to angiosperms, but their biological

functions might be more diverse than we initially imagined.

Supplementary Data

Table S1 Primers used in this study

Fig. S1 Introduction of miR319-insensitive *RKD* genes into wild-type liverwort resulted

in the production of relatively few gemma cups.

Acknowledgements

We thank Kohchi, T. and Yamaoka, S. of Kyoto University, Ishizaki, K. of Kobe

University, Shimamura, M. of Hiroshima University, and Nakajima, K. of NAIST for

their advice and discussions. We also thank Bahari M. and Edanz

(https://jp.edanz.com/ac) for editing a draft of this manuscript.

Author contributions

KF performed the experiments. KF and MT designed the experiments. KF, YW, and MT

wrote the manuscript.

Conflict of interest

The authors declare that they have no competing financial interests related to this work.

Funding

This work was supported by MEXT KAKENHI (21K06225 and 24K09499 to Y.W.) and

the Japan Society for the Promotion of Science funding for Bilateral Joint Research

Projects FY2022-23 (JPJSBP 120224601 to Y.W.).

Data Availability

Relevant data can be found within the manuscript and its supplementary data online.

References

Alaba S, Piszczalka P, Pietrykowska H, Pacak AM, Sierocka I, Nuc PW, Singh K, Plewka

P, Sulkowska A, Jarmolowski A, et al (2015) The liverwort Pellia endiviifolia shares

microtranscriptomic traits that are common to green algae and land plants. New Phytol **206**:

352–367

Axtell MJ, Snyder JA, Bartel DP (2007) Common functions for diverse small RNAs of land

plants. Plant Cell **19**: 1750–1769

Bowman JLJL, Kohchi T, Yamato KT, Jenkins J, Shu S, Ishizaki K, Yamaoka S, Nishihama

R, Nakamura Y, Berger F, et al (2017) Insights into Land Plant Evolution Garnered from

the Marchantia polymorpha Genome. Cell **171**: 287-304.e15

Gamborg OL, Miller RA, Ojima K (1968) Nutrient requirements of suspension cultures of

soybean root cells. Exp Cell Res **50**: 151–158

Hiwatashi T, Quan KL, Yasui Y, Takami H, Kajikawa M, Kirita H, Sato M, Wakazaki M,

Yamaguchi K, Shigenobu S, et al (2019) The RopGEF KARAPPO Is Essential for the

Initiation of Vegetative Reproduction in Marchantia polymorpha. Curr Biol **29**: 3525-

3531.e7

Ishizaki K, Chiyoda S, Yamato KT, Kohchi T (2008) Agrobacterium-mediated transformation

of the haploid liverwort Marchantia polymorpha L., an emerging model for plant biology.

Plant Cell Physiol **49**: 1084–1091

Ishizaki K, Nishihama R, Ueda M, Inoue K, Ishida S, Nishimura Y, Shikanai T, Kohchi T

(2015) Development of gateway binary vector series with four different selection markers

for the liverwort marchantia polymorpha. PLoS One **10**: 1–13

Koi S, Hisanaga T, Sato K, Shimamura M, Yamato KT, Ishizaki K, Kohchi T, Nakajima K

(2016) An Evolutionarily Conserved Plant RKD Factor Controls Germ Cell Differentiation.

Curr Biol **26**: 1–7

Komatsu A, Kodama K, Mizuno Y, Fujibayashi M, Naramoto S, Kyozuka J (2023) Control

of vegetative reproduction in Marchantia polymorpha by the KAI2-ligand signaling

pathway. Curr Biol **33**: 1196-1210.e4

Kosugi S, Ohashi Y (1997) PCF1 and PCF2 specifically bind to cis elements in the rice

proliferating cell nuclear antigen gene. Plant Cell **9**: 1607–1619

Koyama T, Sato F, Ohme-Takagi M (2017) Roles of miR319 and TCP transcription factors in

leaf development. Plant Physiol **175**: 874–885

Kubota A, Ishizaki K, Hosaka M, Kohchi T (2013) Efficient Agrobacterium-mediated

transformation of the liverwort Marchantia polymorpha using regenerating thalli. Biosci

Biotechnol Biochem **77**: 167–172

Lin P-CC, Lu C-WW, Shen B-NN, Lee G-ZZ, Bowman JL, Arteaga-Vazquez MA, Liu L-

YD, Hong S-FF, Lo C-FF, Su G-MM, et al (2016a) Identification of miRNAs and their

targets in the liverwort Marchantia polymorpha by integrating RNA-Seq and degradome

analyses. Plant Cell Physiol **57**: 339–358

Lin P, Lu C, Shen B, Lee G, Bowman JL, Arteaga-vazquez MA, Liu LD, Hong S, Lo C, Su

G, et al (2016b) Identification of miRNAs and Their Targets in the Liverwort Marchantia

polymorpha by Integrating RNA-Seq and Degradome Analyses Special Focus Issue –

Regular Paper. **57**: 339–358

Lin SS, Bowman JL (2018) MicroRNAs in Marchantia polymorpha. New Phytol **220**: 409–416

Martín-Trillo M, Cubas P (2010) TCP genes: a family snapshot ten years later. Trends Plant

Sci **15**: 31–39

Palatnik JF, Allen E, Wu X, Schommer C, Schwab R, Carrington JC, Weigel D (2003)

Control of leaf morphogenesis by microRNAs. Nature **425**: 257–263

Palatnik JF, Wollmann H, Schommer C, Schwab R, Boisbouvier J, Rodriguez R,

Warthmann N, Allen E, Dezulian T, Huson D, et al (2007) Sequence and expression

differences underlie functional specialization of Arabidopsis microRNAs miR159 and

miR319. Dev Cell **13**: 115–25

Pietrykowska H, Sierocka I, Zielezinski A, Alisha A, Carrasco-Sanchez JC, Jarmolowski A,

Karlowski WM, Szweykowska-Kulinska Z (2022) Biogenesis, conservation, and function

of miRNA in liverworts. J Exp Bot **73**: 4528–4545

Rhoades MW, Reinhart BJ, Lim LP, Burge CB, Bartel B, Bartel DP (2002) Prediction of

plant microRNA targets. Cell **110**: 513–520

Romani F, Sauret-Güeto S, Rebmann M, Annese D, Bonter I, Tomaselli M, Dierschke T,

Delmans M, Frangedakis E, Silvestri L, et al (2024) The landscape of transcription factor

promoter activity during vegetative development in Marchantia. Plant Cell **36**: 2140–2159

Rövekamp M, Bowman JL, Grossniklaus U (2016) Marchantia MpRKD Regulates the

Gametophyte-Sporophyte Transition by Keeping Egg Cells Quiescent in the Absence of

Fertilization. Curr Biol **26**: 1782–1789

Shimamura M (2016) Marchantia polymorpha: Taxonomy, phylogeny and morphology of a

model system. Plant Cell Physiol **57**: 230–256

Solly JE, Cunniffe NJ, Harrison CJ (2017) Regional Growth Rate Differences Specified by

Apical Notch Activities Regulate Liverwort Thallus Shape. Curr Biol **27**: 16–26

Sugano SS, Nishihama R, Shirakawa M, Takagi J, Matsuda Y, Ishida S, Shimada T, Hara-

Nishimura I, Osakabe K, Kohchi T (2018) Efficient CRISPR/Cas9-based genome editing

and its application to conditional genetic analysis in Marchantia polymorpha. PLoS One **13**:

e0205117

Sugano SS, Shirakawa M, Takagi J, Matsuda Y, Shimada T, Hara-Nishimura I, Kohchi T

(2014) CRISPR/Cas9-mediated targeted mutagenesis in the liverwort Marchantia

polymorpha L. Plant Cell Physiol **55**: 475–481

Tsuboyama K, Tadakuma H, Tomari Y (2018a) Conformational Activation of Argonaute by

Distinct yet Coordinated Actions of the Hsp70 and Hsp90 Chaperone Systems. Mol Cell **70**:

722-729.e4

Tsuboyama S, Kodama Y (2018) Highly efficient G-AgarTrap-mediated transformation of the

Marchantia polymorpha model strains Tak-1 and Tak-2. Plant Biotechnol **35**: 399–403

Tsuboyama S, Nonaka S, Ezura H, Kodama Y (2018b) Improved G-AgarTrap: A highly

efficient transformation method for intact gemmalings of the liverwort Marchantia

polymorpha. Sci Rep **8**: 1–10

Tsuzuki M, Futagami K, Shimamura M, Inoue C, Kunimoto K, Oogami T (2019) An Early

Arising Role of the MicroRNA156 / 529- SPL Module in Reproductive Development

Revealed by the Liverwort Marchantia polymorpha Report An Early Arising Role of the

MicroRNA156 / 529- SPL Module in Reproductive Development Revealed by the

Liverwort. Curr Biol **29**: 1–8

Tsuzuki M, Nishihama R, Ishizaki K, Kurihara Y, Matsui M, Bowman JLJL, Kohchi T,

Hamada T, Watanabe Y (2016) Profiling and characterization of small RNAs in the

liverwort, Marchantia polymorpha, belonging to the first diverged land plants. Plant Cell

Physiol **57**: 359–372

Yamaoka S, Inoue K, Araki T (2021) Regulation of gametangia and gametangiophore initiation

in the liverwort Marchantia polymorpha. Plant Reprod. doi: 10.1007/s00497-021-00419-y

Yasui Y, Tsukamoto S, Sugaya T, Nishihama R, Wang Q, Kato H, Yamato KT, Fukaki H,

Mimura T, Kubo H, et al (2019) GEMMA CUP-ASSOCIATED MYB1, an Ortholog of

Axillary Meristem Regulators, Is Essential in Vegetative Reproduction in Marchantia

polymorpha. Curr Biol **29**: 3987-3995.e5

Figure Legends

Fig. 1 (A) Primary *MIR319a* or *b* RNAs are predicted to be transcribed from two

annotated loci in the *M. polymorpha* genome. (B) The wild-type (WT) *MIR319a* locus

sequence targeted by gRNA#1 is presented. Mutants #5 and #10 were analyzed further.

(C) The WT *MIR319b* locus sequence targeted by gRNA#3 or #4 is presented. Mutants

#11 and #15 from the gRNA#3-treated pool and #6 and #9 from the gRNA#4-treated pool

were analyzed further. In the superscript *ge-x*, *x* indicates the gRNA species used to create

the mutant. (D) Changes in the *MIR319a* and *b* genomic loci in the double knockout line

miR319ab #10/15. The *MIR319a* locus had a 5-bp deletion and the *MIR319b* locus had a

1-bp insertion, similar to *miR319a^{ge-1} #10* and *miR319b^{ge-3}*. For simplicity, we named the

line *miR319ab #10/15*.

Fig. 2 (A) Disruption of the Mp*MIR319a* and/or *b* loci decreased the number of gemma

cups. Five plants were analyzed and the mean number of gemma cups per thallus is

presented. Tukey–Kramer's honestly significant difference test was performed to analyze

the variance. Statistically significant differences are indicated by different letters (a, b, c;

P < 0.05). (B) Appearance of different thalli. Bar = 1 cm. WT: Tak-1 wild-type plant, #5:

miR319a^{ge-1} #5, #10: *miR319a^{ge-1} #10*, #11: *miR319b^{ge-3} #11*, #15: *miR319b^{ge-3} #15*, #6:

miR319b^{ge-4} #6, #10/15: *miR319ab #10/15*.

Fig. 3 Disruption of the Mp*MIR319a* and/or *b* loci inhibited gemma formation. (A)

Number of gemmae formed in each gemma cup. *P < 0.05 (Tukey method). (B)

Appearance of a gemma cup. Bar = 1 cm. WT: Tak-1 wild-type plant, #5: *miR319a^{ge-1} #5*,

#10: *miR319a^{ge-1} #10*, #11: *miR319b^{ge-3} #11*, #15: *miR319b^{ge-3} #15*.

Fig. 4 Exon–intron structures of Mp*RKD* and Mp*R2R3-MYB21* genes in the wild-type

form (or sensitive form, with prefix g) and miR319-resistant form (with prefix m) of (A)

Mp*RKD* and (B) mMp*R2R3-MYB21* genes. Vertical bars indicate conventional base-

pairing, whereas double dots indicate G–U pairs. AUG and UGA are initiation and

termination codons, respectively, of open reading frames. 5′UTR and 3′UTR refer to

untranslated regions.

Fig. 5 Introduction of the miR319-resistant mMp*RKD* gene suppressed gemma cup

formation in wild-type thalli, but the introduction of gMp*RKD* and mMp*MYB21* genes

did not.

Fig. 6 Introduction of the wild-type *RKD* gene (gMp*RKD*) induced normal gametophyte

development in Tak-1 (A, B) and Tak-2 (G, H). Introduction of the miR319-resistant

mutant *RKD* gene (mMp*RKD*) into Tak-1 induced the production of antheridiophores

with stalked receptables with irregular asymmetrical shapes (C, D) and antheridiophores

with exposed antheridia (E, F). Introduction of mMp*RKD* into Tak-2 induced the

production of archegoniophores with short digits (I,J), but normal archegonia (K).

Primers used in this study

primers to construct guide RNA vectors for establishment of *miR319a* or *miR319b* mutants

PAM_MIR319a-174f-F	CTCGACAATCCGTCCTTGGACTGA
PAM_MIR319a-174f-R	AAACTCAGTCCAAGGACGGATTGT
PAM_MIR319b-161r-F	CTCGAGCTCCCTTCAGTCCAAGTC
PAM_MIR319b-161r-R	AAACGACTTGGACTGAAGGGAGCT
PAM-MIR319b-179f-F	CTCGTTGGCCCGACTTGGACTGAA
PAM-MIR319b-179f-R	AAACTTCAGTCCAAGTCGGGCCAA

primers to construct miR319 resistant form Mp*RKD*

IF-Hind-pRKD-F	ACGACGGCCAGTGCCAAGCTTGGACACCGGAGAGAAATCAA
Inf-gMpRKD-Sal-R	CGGGGATCCTCTAGAGTCGACACACAAAGTTTTATAACGCCGATGT
mRWP_RK-F	CTGAATCGCAATCGAGCGGAGTTTCAAATCAAATAGAAGC
mRWP_RK-R	TCCGCTCGATTGCGATTCAGTATGTCATGAGACCAGGGTT

primers to construct miR319 resistant form Mp*R2R3-MYB21*

mMYB33-F	AGAATTGCCAAGTTCTCAGTCTGCAGAGTCTGCTGACAGC
mMYB33-R	ACTGAGAACTTGGCAATTCTAATTTCAGGCTGGAATTCGG
Hind-pMpMYB21-3000-F	GTTAAGCTTcgtgcccttgacctcctcctc
Sal-pMpMYB21-R	GTTGTCGACccaggcaccccagactcttgaac
Bam-MpMYB33-R2	GTTGGATCCGGGATAGGTCAGCAGGTCAGTAGT
pMYB33-459F	TGCCTGCCATGGCATACATCCCATC

Fig.S1 Introduction of the miR319-insensitive iMp*RKD* gene suppressed gemma cup formation in thalli

(A) Construct of wild-type *RKD* (gMp*RKD*) and miR319 insensitive *RKD* (iMp*RKD*) genes. The sequence segment complementary to miR319 sequences located in 3' untranslated region (3' UTR) was deleted to obtain the iMp*RKD* gene. Each construct was placed under the direction of the promoter of Mp*RKD* and introduced into Tak-1 wild type plants to obtain respective transformants. (B) The numbers of gemma cups were counted per thallus for transformants.

Fig. 1 (A) Primary *MIR319a* or *b* RNAs are predicted to be transcribed from two annotated loci in the *M. polymorpha* genome. (B) The wild-type (WT) *MIR319a* locus sequence targeted by gRNA#1 is presented. Mutants *a* #5 and *a* #10 were analyzed further. (C) The WT *MIR319b* locus sequence targeted by gRNA#3 or #4 is presented. Mutants *b* #11 and *b* #15 from the gRNA#3-treated pool and *b* #6 and *b* #9 from the gRNA#4-treated pool were analyzed further. In the superscript ge-x, x indicates the gRNA species used to create the mutant. (D) Changes in the *MIR319a* and *b* genomic loci in the double knockout line *miR319ab #10/15*. The *MIR319a* locus had a 5-bp deletion and the *MIR319b* locus had a 1-bp insertion, as did *miR319a^{ge-1}* *a* #10 and *miR319b^{ge-3}* *b* #15. For simplicity, we named the line *miR319ab #10/15*.

Fig. 2 (A) Disruption of the Mp*MIR319a* and/or *b* loci decreased the number of gemma cups. Five plants were analyzed and the mean number of gemma cups per thallus is presented. Tukey–Kramer's honestly significant difference test was performed to analyze the variance. Statistically significant differences are indicated by different letters (a, b, c; P < 0.05). (B) Appearance of different thalli. Bar = 1 cm. WT: Tak-1 wild-type plant, *a* #5: *miR319a^{ge-1}* #5, *a* #10: *miR319a^{ge-1}* #10, *b* #11: *miR319b^{ge-3}* #11, *b* #15: *miR319b^{ge-3}* #15, *b* #6: *miR319b^{ge-4}* #6, *b* #9: *miR319b^{ge-4}* #9, *ab* #10/15: miR319ab #10/15.

(n = 10)

Fig. 3 Disruption of the Mp*MIR319a* and/or *b* loci inhibited gemma formation. (A) Number of gemmae formed in each gemma cup. *P < 0.05 (Tukey method). (B) Appearance of a gemma cup. Bar = 1 cm. WT: Tak-1 wild-type plant, *a* #5: *miR319a^{ge-1}* #5, *a* #10: *miR319a^{ge-1}* #10, *b* #11: *miR319b^{ge-3}* #11, *b #115*: *miR319b^{ge-3}* #15.

A. Mp*RKD*

gMp*RKD*

···CUCAUGACAUAGGGGCUCCCUUCCGUCCAAAGUUUCAAAUC···

|| : | | | | | | | | | | | | | | | |

3' CCCUCGAGGGAAGUCAGGUUC 5' miR319

mMp*RKD*

: | | | : | | : :

···CUCAUGACAUACUGAAUCGCAAUCGAGCGGAGUUUCAAAUC···

B. Mp*R2R3-MYB21*

gMp*R2R3-MYB21* L E L P S S Q S

5' ···UGGAGCUCCCUUCAAGCCAAU··· 3'

| | | | | | | | | | | | | | | |

3' CCCUCGAGGGAAGUGAGGUUC 5' miR319

mMp*R2R3-MYB21* | | : | | | : : | | :

5 ···UAGAAUUGCCAAGUUCUCAGU··· 3'

Fig. 4 Exon–intron structures of Mp*RKD* and Mp*R2R3-MYB21* genes in the wild-type form (or sensitive form, with prefix g) and miR319-resistant form (with prefix m) of (A) Mp*RKD* and (B) mMp*R2R3-MYB21* genes. Vertical bars indicate conventional base-pairing, whereas double dots indicate G–U pairs. AUG and UGA are initiation and termination codons, respectively, of open reading frames. 5′UTR and 3′UTR refer to untranslated regions.

Fig. 5 Introduction of the miR319-resistant mMp*RKD* gene suppressed gemma cup formation in thalli, but the introduction of gMp*RKD* and mMp*R2R3MYB21* genes did not. Bar = 1 cm.

Fig. 6 Introduction of the wild-type *RKD* gene (gMp*RKD*) induced normal gametophyte development in Tak-1 (A, B) and Tak-2 (G, H). Introduction of the miR319-resistant mutant *RKD* gene (mMp*RKD*) into Tak-1 induced the production of antheridiophores with stalked receptables with irregular asymmetrical shapes (C, D) and antheridiophores with exposed antheridia (E, F). Introduction of mMp*RKD* into Tak-2 induced the production of archegoniophores with short digits (I, J), but normal archegonia (K).