Title:

MpmiR319 promotes gemma/gemma cup formation in the Liverwort *Marchantia*

polymorpha

Authors:

Kazutaka Futagami[1]

Masayuki Tsuzuki[1,2]

Yuichiro Watanabe[1]

[1] Department of Life Sciences, Graduate School of Arts and Sciences, The University of

Tokyo, Tokyo 153-8902, Japan

[2] Faculty of Agriculture and Marine Sciences, Graduate School of Integrated Arts and

Sciences, Kochi University

Correspondence: mtsuzuki@kochi-u.ac.jp (M.Tsuzuki), cyuiwat@g.ecc.u-tokyo.ac.jp

(Y.Watanabe)

ORCID

Kazutaka Futagami: https://orcid.org/0000-0002-3321-6508

Masayuki Tsuzuki: https://orcid.org/0000-0002-0642-1143

Yuichiro Watanabe: https://orcid.org/0000-0002-7139-4903

Abstract

Evolutionally conserved miRNA-target mRNA modules have been identified as a core set of functionally and biologically important gene regulation circuits in land plants. Whether the miR159/319 family could maintain such a common biologically fundamental function remains unknown. In this study, we applied the CRISPR-Cas9 system to knock out two Mp*MIR319* loci, *MIR319a* and/or *MIR319b*, in the liverwort *Marchantia polymorpha* to observe phenotypic changes during development. The *MIR319b* knockout line developed fewer gemma cups, while there was a decrease in the number of gemma produced per gemma cup in the *MIR319a* knockout line. It was predicted that miR319 targets Mp*RKD* and Mp*R2R3-MYB21* mRNA and suppresses their expression. To verify this, we constructed miR319-resistant Mp*RKD* (mMp*RKD*) and Mp*R2R3-MYB2*1 (mMp*R2R3-MYB2*1) genes and introduced these genes into wild-type liverwort plants to monitor phenotypic changes. We found that mMp*RKD* induced gemma/gemma cup-less liverworts but mMp*R2R3-MYB2*1 did not. These results indicate that miR319 mainly targets and represses RKD expression rather than that of R2R3-MYB21. Additionally, negative regulation of RKD expression by miR319 controls the placement and number of gemma/gemma cups produced in thalli.

Keywords: *Marchantia*, miR319, gemma, gemma cup, MpRKD

Introduction

It is generally known that plant miRNAs can form complementary base-pair interactions with their target mRNAs with stricter complementarity than their animal counterparts (Rhoades et al., 2002)(Lin and Bowman, 2018). Each plant miRNA can target quite a limited number of mRNAs. Accordingly, target mRNA(s) of each miRNA could be predicted by searching genome sequence data with the aid of the miRNA sequence of interest. *In vivo* studies of miRNAs conserved among land plants have shown that their target mRNAs are also well-conserved cognate genes across phyla (Axtell et al., 2007)(Tsuzuki et al., 2019)(Alaba et al., 2015)(Pietrykowska et al., 2022). Such conserved miRNA-target mRNA modules could be regarded as a core set of functionally and biologically important gene regulation circuits in land plants. But still more experimental data are expected to show the biological significance of highly conserved miRNAs in basal land plants.

Among conserved miRNAs, the miR159/319 family has a unique feature in that the miRNA may have changed their target mRNA species during phylogenetic diversification (Palatnik et al., 2007). In Arabidopsis, miR319 preferentially targets mRNAs of *TEOSINTE BRANCHED1, CYCLOIDEA,* and *PROLIFERATING CELL FACTOR (TCP)* genes (Martín-Trillo and Cubas, 2010)(Kosugi and Ohashi, 1997). TCP genes encode bHLH motif transcription factors which affect redundantly cotyledon boundary and form leaf serration (Palatnik et al., 2003)(Koyama et al., 2017).

Marchantia polymorpha is a liverwort species that stands out as a valuable model for land plant studies, with it garnering an increasing amount of information and resources

for comparative evolution studies in recent years (Bowman et al., 2017). Recent studies

have revealed that land plants have nearly 300-500 lineage-specific diverse miRNAs

(Axtell et al., 2007)(Lin and Bowman, 2018)(Pietrykowska et al., 2022) and that as few

as seven miRNA families are deeply conserved from the liverwort *Marchantia*

polymorpha, the moss *Physcomitrella patens*, to angiosperms (Axtell et al.,

2007)(Tsuzuki et al., 2016)(Lin et al., 2016b)(Pietrykowska et al., 2022). But most

liverwort miRNAs are yet to be biologically studied in detail enough to obtain a wide

understanding of commonality or specificity among land plants.

In *M. polymorpha*, it was shown that Mp*miR319* targets the mRNA of MpR2R3-MYB21

(an *Arabidopsis* MYB33-like transcription factor, a member of GAMYB families) (Lin

et al., 2016b)(Tsuzuki et al., 2016) and that of the RWP-RK DOMAIN (RKD)-containing

transcription factor (Lin et al., 2016b) based on data from 5'-RACE or degradome

analysis. As such, Mp*miR319* seemingly has some regulatory functions on such target

candidate genes, but this has yet to be experimentally tested.

Materials and methods

Plant materials and growth condition

Takaragaike-1 (Tak-1, male) and Takaragaike-2 (Tak-2, female) were used as wild-type

M. polymorpha (Ishizaki et al., 2008). Plants were grown asexually on half-strength

Gamborg's B5 medium containing 1% agar, under continuous white fluorescent light at

22ºC (Gamborg et al., 1968). For induction of the sexual reproductive phase, plants were

grown under a mixture of white fluorescent light and far-red (FR) power light-emitting

diodes (OSXXXXT3C1E, OptoSupply; peak emission at 730 nm, with a full width at

half-maximum of 20 nm) at 22ºC in continuous light conditions.

Genomic DNA

Genomic DNA was extracted from Tak-1 thallus according to the method described in

(Tsuzuki et al., 2019) or using the DNeasy Plant Mini kit (QIAGEN, Hilden, Germany).

Liverwort transformation and mutant selection

The regenerating thalli method (for *mir319a* or *mir319b* mutants) and G-AgarTrap

method (for establishment of gMp*RKD*, mMp*RKD*, gMp*MYB21* and mMp*MYB21* lines)

were used for transformation of *M. polymorpha* (Kubota et al., 2013) (Tsuboyama and

Kodama, 2018). Briefly, transformed plants were selected on half-strength Gamborg's B5

medium containing 0.5 g/L MES, 1% (w/v) agar and 10 µg/ml hygromycin or 0. 5 µM

chlorsulfuron, or their combination. After screening T1 plants on drug-supplemented

media, we obtained more or less 10 candidate lines for each gRNA in the male and female

backgrounds.

Plasmid construction to apply CRISPR-Cas9 system

The vectors pMpGE010, pMpGE011 and pMpGE_En03 (Sugano et al., 2018) were used

for CRISPR/Cas9-based genome editing of Mp*MIR319A* and Mp*MIR319B* loci. The

pMpGE vectors harboring gRNAs targeting *MIR319A* or *MIR319B* genes were

constructed as described (Sugano et al., 2018) using oligonucleotide pairs of PAM

MIR319a-174f-F and -174f-R, PAM MIR319b-161r-F and -161r-R or PAM-MIR319b-

179f-F and -179f-R, respectively, to make pMpGE_En03-gRNA1, pMpGE_En03-

gRNA3 and pMpGE_En03-gRNA4 constructs.

Selection of *mir319a^{ge}*, *mir319b^{ge}* and *mir319ab^{ge}* mutants was performed as described

previously (Tsuzuki et al., 2019). The DNA segments coding *MIR319A* and *MIR319B* loci

(with genomic DNA as the template) were amplified using PrimeSTAR HS DNA

polymerase (TaKaRa Bio, Kusatsu, Japan) with miR319aF-miR319R or miR319bF-

miR319R primer pairs and were sequenced according to Sanger method. PCR and

sequencing primer details are listed in Table S1.

Introduction of wild-type and miR319 resistant MpRKD/MpR2R3-MYB21 transgenes

The vector pMpGWB300 (Ishizaki et al., 2015) was used to introduce wild-type

(gMp*RKD* and gMp*MYB21*) or miR319-resistant types of *RKD* or *MYB21* genes

(mMp*RKD* and mMp*MYB21*) into wild-type Tak-1 or Tak-2 lines.

To examine the effect of miR319 on repression of MpRKD expression, we generated the

pMpGWB300:proMpRKD:MpRKD   and   pMpGWB300:proMpRKD:mMpRKD

plasmids as follows: for pMpGWB300:proMpRKD:MpRKD, the genomic sequence

from the Mp*RKD* promoter region (3827 bp upstream from start codon) to the 3'UTR

was amplified by PCR with PrimeSTAR HS DNA polymerase (TaKaRa Bio, Kusatsu,

Japan) and the primers IF-Hind-pRKD-F and IF-Sal-gRKD-R, using Tak-1 genomic

DNA as the template. For pMpGWB300:proMpRKD:mMpRKD, two PCR fragments

were amplified from genomic DNA of Tak-1 with the IF-Hind-pRKD-F, Inf-Kpn-gRWP

RK-R and IF-Sal-gRKD-R primer sets to mutate the miR319-target sequence in the

MpRKD transgene. Each one or two PCR fragments were inserted into pMpGWB300

that was cleaved by HindIII and SalI using the In-Fusion Cloning kit (TaKaRa Bio,

Kusatsu,   Japan)   to   generate   pMpGWB300:proMpRKD:MpRKD   and

pMpGWB300:proMpRKD:mMpRKD plasmids.

Results

Targeting (perturbation) of either of the two MIR319 genes by guide-RNAs resulted in

reduced gemma cup production in thalli

Marchantia polymorpha has two *MIR319* coding loci, both of which can produce

miRNAs of an identical mature sequence, 5'-$_p$CUUGGACUGAAGGGAGCUCCC$_{OH}$-3'

(Lin et al., 2016b; Tsuzuki et al., 2016). The two *MIR319* loci have now been annotated

(*M. polymorpha* genome ver.6.1) as Mp*MIR319A* (Mp3g08965) and Mp*MIR319B*

(Mp4g01845), respectively (Fig.1A). To identify the possible biological function(s) of

miR319 in *M. polymorpha*, we decided to observe phenotypic changes after knocking

down its production. To achieve this, we designed a set of guide RNAs (gRNA 1 for

MIR319A (Fig.1B) and gRNAs 3 and 4 for *MIR319B* (Fig.1C) to construct vectors to

introduce in-del mutations at respective *MIR319* loci by applying the CRISPR/Cas9

genome editing system (Sugano et al., 2014)(Sugano et al., 2018).

We adopted the *Agrobacterium*-mediated transformation protocol and introduced the

vectors into the thalli of Tak-1 to express designed gRNAs and Cas9 (Sugano et al.,

2014)(Tsuboyama et al., 2018).   After the corresponding drug selection of T1 plants, we

amplified respectively targeted genomic sequences from a fraction of T1 thalli and

analyzed them by Sanger sequencing. We selected multiple lines that acquired in-del

mutations at *MIR319a* or *MIR319b* loci and characterized several of them (Fig. 1B and

C) in the following analysis. Mutants with in-del mutations in the Mp*mir319a* locus were

characterized as *mir319a^{ge-1}* #5 and *mir319a^{ge-1}*#10; mutants with in-del mutations in the

Mp*mir319b* locus were characterized as *mir319b*^{ge-3} #11, *mir319b*^{ge-3}#15, *mir319b*^{ge-4} #6,

and *mir319b*^{ge-4}#9.

Mir319a^{ge} and *mir319b*^{ge} mutant plants were able to grow as fast as wild-type Tak-1

plants. However, we noticed that the mutants – though able to produce gemma cups –

developed a smaller number of gemma cups than wild-type plants (Fig. 2); *mir319a^{ge-1}*

(#5 and #10) and *mir319b*^{ge-3} plants (#11 and #15) produced fewer gemma cups than wild-

type liverwort, and *mir319b*^{ge-4} plants produced a significantly lower number (#6 and #9).

In addition, we noticed that *mir319a^{ge-1}* (#5 and #10) mutants produced a reduced

number of gemma in each gemma cup (Fig.3A). In contrast, *mir319b*^{ge-3} (#11 and #15)

mutants produced as much gemma as wild-type plants in each gemma cup (Fig.3A,B).

We also created similar *mir319a^{ge}* and *mir319b*^{ge} mutants in female Tak-2 plants.

Reduced formation of gemma cups was observed in Tak-2-based mutants as well. Thus,

we characterized and report mainly Tak-1 background lines in vegetative growth stages.

These observations suggest that miR319 is involved in the formation of gemma and

gemma cups in *M. polymorpha* by regulating target gene expression.

We also tried to obtain double knock-out miR319ab mutants. To achieve this, we

targeted the *MIR319a* locus of *mir319b^{ge-3}* #5 with guideRNA-1.　We obtained one

double knock-out line and found that the new mutant line acquired a five base deletion in

the *MIR319a* locus in addition to a one base addition in the *MIR319b* locus as shown in

Fig.1D. For convenience, the line was named *miR319ab^{ge} #5/15* mutant (Fig.1D). The

severe phenotype of *miR319ab^{ge}#5/15* was no more than those of *miR319a* and *mir319b*

mutants (Fig.2, #5/15) in terms of the number of developed gemma cups/gemma. The

phenotypic severity was expected to be enhanced by disruption of both *miR319a* and *b*

loci.

Regulation of MpRKD by miR319 is involved in gemma cup formation

Earlier degradome analysis studies indicated that Mp*RKD* (RWP-RK DOMAIN-

CONTAINING 1) and Mp*MYB33* mRNAs are targeted by Mp*miR319* (Lin et al., 2016a).

Together with the phenotypes of *mir319a^{ge}* and *mir319b^{ge}* mutants, we attempted to

address whether these genes are targeted by miR319, with the goal of finding whether

susceptibility to miR319 is actually involved in gemma/gemma cup formation in either

or both of these mRNA genes.

First, we focused on Mp*RKD*. Mp*RKD* encodes an RWP-RK domain-containing

transcription factor and was earlier characterized as an evolutionarily conserved regulator

of germ cell differentiation (Koi et al., 2016)(Rövekamp et al., 2016). At that time,

although not clearly stated, it was reported that Mp*RKD* is expressed on gemma cup rims

(Koi et al., 2016) and is necessary for normal growth of cups (Rövekamp et al., 2016).

We first tried to test whether Mp*RKD* downregulation by miR319 is linked to

gemma/gemma cup formation.

We constructed a mutated *RKD* coding sequence which can escape from miR319 action.

The 21-nt sequence stretch complementary to miR319 starts from 4 nt downstream its

stop codon in the *M. polymorpha RKD* (Mp*RKD*) mRNA. We constructed a miR319-

resistant form of the Mp*RKD* gene (designated mMp*RKD*; Fig.4A) by introducing 12

base substitutions to decrease the base complementarity. The mMp*RKD* gene was placed

downstream its own original 3.8 kbp promoter sequences in the pMpGWB300 vector

(Ishizaki et al., 2015). As a control we cloned wild-type Mp*RKD* (designated gMp*RKD*)

in the same vector. gMp*RKD* and mMp*RKD* genes were respectively introduced into Tak-

1 wild-type plants to obtain gMp*RKD* and mMp*RKD* lines.

Both gMp*RKD* and mMp*RKD* line plants showed similar thalli growth rates. But the

mMp*RKD* plants showed a striking phenotype in that they did not develop any gemma

cups (Fig. 5: mMpRKD). The gemma cup-less phenotypes were similar to *mir319a^{ge-1}*,

or *miR319ab #5/15* mutants (Fig.5; ab #5/15) in regards to low frequency of gemma cup

formation. In contrast, the gMp*RKD* lines produced as many gemma cups as wild-type

Tak-1 liverworts (Fig.5; gMpRKD). These results substantiated that the sequences

complementary to Mp*miR319* in Mp*RKD* mRNA are deeply involved in gemma/gemma

cup formation. This is likely to be related to subsequent suppression of MpRKD

translation.

As mentioned before, the miR319 target sequence is located in the 3' untranslated region

downstream the RKD termination codon sequence. Thus, we were able to design another

type of miR319-insensitive *RKD* gene, named Mp*RKD*-Δ3'UTR, by removing the

complementary sequences from the 3'-UTR without disturbing RKD translation (Fig.S1-

A). When we introduced the insensitive form of Mp*RKD*-Δ3'UTR gene into liverwort,

thalli grew but could develop few gemma cups, as was the case with the above mMP*RKD*

liverwort (Fig.S1-B).

miR319-resistant MpMYB21 did not disturb gemma cup/gemma formation

Mp*MYB21* gene is also a target candidate gene of miR319 on the basis of 5'-RACE or

degradome analysis (Tsuzuki et al., 2016)(Lin et al., 2016b). To test whether *MYB21*

regulation by miR319 is involved in gemma cup formation, we created a miR319-

resistant Mp*MYB21* (mMp*R2R3-MYB21*) gene under its own promoter as we did with the

Mp*RKD* gene (Fig.4B). Wild-type, or miR319-sensitive form of Mp*MYB21* (*g*Mp*R2R3-*

MYB21) gene was also constructed in parallel.

gMYB21 and *mMYB21* genes were introduced into Tak-1 wild-type liverwort to obtain

respective transgenic lines. *mMYB21* liverwort lines produced thalli slightly thinner than

gMYB21 lines in width and but could develop gemma cups/gemma without discernible

difference in number (Fig.5; mMpMYB21). These results indicated that the Mp*R2R3-*

MYB21 gene has less contribution, if any, than the Mp*RKD* gene in liverwort gemma

cup/gemma formation.

mMpRKD plants develop deformed antheridiophores/archegoniophores, still with normal

sperm and egg formation

It was previously reported that Mp*RKD* is expressed in developing eggs and sperm

precursors, and regulates egg and sperm formation in *M. polymorpha* (Koi et al.,

2016)(Rövekamp et al., 2016). Thus we next checked possible phenotypic changes in

germ cell, gametangia and gametangiophore differentiation (Yamaoka et al.,

2021)(Shimamura, 2016) in gMp*RKD* and mMp*RKD* liverworts.

mMp*RKD* liverworts of Tak-1 background developed male gametophytes under FR

illumination (Fig.6C) as early as gMp*RKD* liverworts (Fig.6A). mMp*RKD*

antheridiophores had normally-shaped stalked receptables, but with an irregular shape

(Fig. 6C,D). Antheridia were formed as if they were protruding outwards but notably not

surrounded by antheridiophores in mMp*RKD* lines (Fig.6E,F). We could observe similar

sperm mobility in mMp*RKD* liverworts as was found in wild type Tak-1 and gMp*RKD*

liverworts.

We next checked whether introduction of gMp*RKD* and mMp*RKD* genes into Tak-2

wild-type liverworts affects archegonia/archegoniophore development. We established

respective transgenic lines using the two constructs described in Fig.4A. mMp*RKD*

liverworts of Tak-2 background developed female gametophytes under FR illumination

as early as gMp*RKD* liverworts, but they produced archegoniophores with a different

shape (Fig.6I) from those of gMp*RKD* (Fig.6G) or Tak-2 plants. mMp*RKD*

archegoniophores had short digit shapes but grew normal archegonia (Fig.6I). The

archegonia had almost identical appearance (Fig.6J) with those of gMp*RKD* (Fig.6H) or

wild-type Tak-2 liverworts.

Discussion

While we were profiling *Marchantia* miRNAs after RNA-seq analysis, we tried to

induce overexpression of miR319 precursor RNAs to speculate what biological function

miR319 might have. Ectopic expression of liverwort miR319 precursor induced

abnormal development of gemma cups (Tsuzuki et al., 2016). This suggested that

balanced expression of miRNA/target mRNAs is necessary for normal gemma cup

development in *M. polymorpha*.

The possible function of Mp*RKD* have previously been investigated via mutant

analyses in *Marchantia* (Koi et al., 2016)(Rövekamp et al., 2016). It was revealed that

Mp*RKD* is essential for normal germ cell differentiation in both male and female

gametophytes. Notably, both studies also reported that Mp*RKD* is involved in non-

reproductive cells surrounding the antheridia and developing egg and sperm precursors

and acts as an essential regulator of germ cell differentiation in *M. polymorpha* (Koi et

al., 2016)(Rövekamp et al., 2016). However, it remained uncertain whether biologically

significant gene regulatory networks (GRNs) are organized by miR319 and Mp*RKD* and

functional at the vegetative stage.

It has been described that the founder gemma arise directly from above the center of the

apical notch of the thallus, followed by the formation of the cup rims surrounding the

gemmae (Shimamura, 2016). Together it was presumed that MpRKD is involved in

various stages of cell proliferation to form a variety of tissues in *M. polymorpha*,

including gemmae/gemma cups. Our preliminary data showed that the Mp*miR319a*,

Mp*miR319b* and Mp*RKD* gene promoters are active in midribs, apical notches and

gemma cup areas, which is consistent with recognized phenotypic changes. Still, further

investigation is needed to understand how gemma cups are organized by miR319-

mediated GRNs.

The phenotypes of *mir319b*$^{ge-4}$ #6 and *mir319b*$^{ge-4}$ #9 were more severe than those of

mir319b^{ge-3} #11 and *mir319b*$^{ge-3}$#15. This difference in severity may have been caused by

the position of base changes that occurred inside the miR319 sequence. The *mir319b*$^{ge-4}$

plants would be able to produce miRNA, but with differing sequences in the middle

portion. Such RNA would not anneal to the target mRNAs effectively and would show

more distinct phenotypic changes as indicated in Fig.2.

In the moss *Physcomitrella patens*, it was found that miR159/319 are encoded in the

genome and possibly target cyclin domain proteins and two MYB transcription factors

(Axtell et al., 2007). miR159/319 families could be considered as highly conserved

miRNAs from bryophytes to angiosperms.

Acknowledgements

We thank Kohchi, T., Yamaoka, S. of Kyoto University, Ishizaki, K. of Kobe University,

Shimamura, M. of Hiroshima University, Nakajima, K. of NAIST for their advice and

discussions and Bahari M. for English editing.

Author contributions

KF performed the experiments. KF and MT designed the experiments. KF, YW and MT

wrote the manuscript.

Conflict of interest

The authors declare no conflict of interest.

Funding

This work was supported by MEXT KAKENHI (21K06225 to Y,W.) and JSPS (Japan

Society for the Promotion of Science) funding for Bilateral Joint Research Projects

FY2022-23 (JPJSBP 120224601 to Y.W.)

References

Alaba S, Piszczalka P, Pietrykowska H, Pacak AM, Sierocka I, Nuc PW, Singh K,

Plewka P, Sulkowska A, Jarmolowski A, et al (2015) The liverwort Pellia

endiviifolia shares microtranscriptomic traits that are common to green algae and

land plants. New Phytol **206**: 352–367

Axtell MJ, Snyder JA, Bartel DP (2007) Common functions for diverse small RNAs

of land plants. Plant Cell **19**: 1750–1769

Bowman JLJL, Kohchi T, Yamato KT, Jenkins J, Shu S, Ishizaki K, Yamaoka S,

Nishihama R, Nakamura Y, Berger F, et al (2017) Insights into Land Plant

Evolution Garnered from the Marchantia polymorpha Genome. Cell **171**: 287-

304.e15

Gamborg OL, Miller RA, Ojima K (1968) Nutrient requirements of suspension

cultures of soybean root cells. Exp Cell Res **50**: 151–158

Ishizaki K, Chiyoda S, Yamato KT, Kohchi T (2008) Agrobacterium-mediated

transformation of the haploid liverwort Marchantia polymorpha L., an emerging

model for plant biology. Plant Cell Physiol **49**: 1084–1091

Ishizaki K, Nishihama R, Ueda M, Inoue K, Ishida S, Nishimura Y, Shikanai T,

Kohchi T (2015) Development of gateway binary vector series with four different

selection markers for the liverwort marchantia polymorpha. PLoS One **10**: 1–13

Koi S, Hisanaga T, Sato K, Shimamura M, Yamato KT, Ishizaki K, Kohchi T,

Nakajima K (2016) An Evolutionarily Conserved Plant RKD Factor Controls

Germ Cell Differentiation. Curr Biol **26**: 1–7

Kosugi S, Ohashi Y (1997) PCF1 and PCF2 specifically bind to cis elements in the rice

proliferating cell nuclear antigen gene. Plant Cell **9**: 1607–1619

Koyama T, Sato F, Ohme-Takagi M (2017) Roles of miR319 and TCP transcription

factors in leaf development. Plant Physiol **175**: 874–885

Kubota A, Ishizaki K, Hosaka M, Kohchi T (2013) Efficient Agrobacterium-

mediated transformation of the liverwort Marchantia polymorpha using

regenerating thalli. Biosci Biotechnol Biochem **77**: 167–172

Lin P-CC, Lu C-WW, Shen B-NN, Lee G-ZZ, Bowman JL, Arteaga-Vazquez MA,

Liu L-YD, Hong S-FF, Lo C-FF, Su G-MM, et al (2016a) Identification of

miRNAs and their targets in the liverwort Marchantia polymorpha by integrating

RNA-Seq and degradome analyses. Plant Cell Physiol **57**: 339–358

Lin P, Lu C, Shen B, Lee G, Bowman JL, Arteaga-vazquez MA, Liu LD, Hong S,

Lo C, Su G, et al (2016b) Identification of miRNAs and Their Targets in the

Liverwort Marchantia polymorpha by Integrating RNA-Seq and Degradome

Analyses Special Focus Issue – Regular Paper. **57**: 339–358

Lin SS, Bowman JL (2018) MicroRNAs in Marchantia polymorpha. New Phytol **220**:

409–416

Martín-Trillo M, Cubas P (2010) TCP genes: a family snapshot ten years later. Trends

Plant Sci **15**: 31–39

Palatnik JF, Allen E, Wu X, Schommer C, Schwab R, Carrington JC, Weigel D

(2003) Control of leaf morphogenesis by microRNAs. Nature **425**: 257–263

Palatnik JF, Wollmann H, Schommer C, Schwab R, Boisbouvier J, Rodriguez R,

Warthmann N, Allen E, Dezulian T, Huson D, et al (2007) Sequence and

expression differences underlie functional specialization of Arabidopsis

microRNAs miR159 and miR319. Dev Cell **13**: 115–25

Pietrykowska H, Sierocka I, Zielezinski A, Alisha A, Carrasco-Sanchez JC,

Jarmolowski A, Karlowski WM, Szweykowska-Kulinska Z (2022) Biogenesis,

conservation, and function of miRNA in liverworts. J Exp Bot **73**: 4528–4545

Rhoades MW, Reinhart BJ, Lim LP, Burge CB, Bartel B, Bartel DP (2002)

Prediction of plant microRNA targets. Cell **110**: 513–520

Rövekamp M, Bowman JL, Grossniklaus U (2016) Marchantia MpRKD Regulates

the Gametophyte-Sporophyte Transition by Keeping Egg Cells Quiescent in the

Absence of Fertilization. Curr Biol **26**: 1782–1789

Shimamura M (2016) Marchantia polymorpha: Taxonomy, phylogeny and

morphology of a model system. Plant Cell Physiol **57**: 230–256

Sugano SS, Nishihama R, Shirakawa M, Takagi J, Matsuda Y, Ishida S, Shimada

T, Hara-Nishimura I, Osakabe K, Kohchi T (2018) Efficient CRISPR/Cas9-

based genome editing and its application to conditional genetic analysis in

Marchantia polymorpha. PLoS One **13**: e0205117

Sugano SS, Shirakawa M, Takagi J, Matsuda Y, Shimada T, Hara-Nishimura I,

Kohchi T (2014) CRISPR/Cas9-mediated targeted mutagenesis in the liverwort

Marchantia polymorpha L. Plant Cell Physiol **55**: 475–481

Tsuboyama S, Kodama Y (2018) Highly efficient G-AgarTrap-mediated

transformation of the Marchantia polymorpha model strains Tak-1 and Tak-2.

Plant Biotechnol **35**: 399–403

Tsuboyama S, Nonaka S, Ezura H, Kodama Y (2018) Improved G-AgarTrap: A

highly efficient transformation method for intact gemmalings of the liverwort

Marchantia polymorpha. Sci Rep **8**: 1–10

Tsuzuki M, Futagami K, Shimamura M, Inoue C, Kunimoto K, Oogami T (2019)

An Early Arising Role of the MicroRNA156 / 529- SPL Module in Reproductive

Development Revealed by the Liverwort Marchantia polymorpha Report An Early

Arising Role of the MicroRNA156 / 529- SPL Module in Reproductive

Development Revealed by the Liverwort. Curr Biol **29**: 1–8

Tsuzuki M, Nishihama R, Ishizaki K, Kurihara Y, Matsui M, Bowman JLJL,

Kohchi T, Hamada T, Watanabe Y (2016) Profiling and characterization of

small RNAs in the liverwort, Marchantia polymorpha, belonging to the first

diverged land plants. Plant Cell Physiol **57**: 359–372

Yamaoka S, Inoue K, Araki T (2021) Regulation of gametangia and gametangiophore

initiation in the liverwort Marchantia polymorpha. Plant Reprod. doi:

10.1007/s00497-021-00419-y

Figure Legends

Fig.1 (A) Primary *MIR319a or b* RNAs are predicted to be transcribed from two

annotated loci in the *M. polymorpha* genome. (B) The row wt shows the wild-type

MIR319a locus sequence and is targeted by gRNA#1. Mutants #5 and #10 were selected

for further analysis among obtained in-del mutants. (C) The row wt shows the wild-type

MIR319b locus sequence and is targeted by gRNA#3 or #4. Mutants #11 and #15 from

gRNA#3 treated-pool and mutants #6 and #9 from gRNA#4 treated-pool were selected

for further analysis. X in the superscript *ge-x* indicates guide RNA species with which the

respective mutant was created. (D) The genomic change of the double knock-out line for

MIR319a and *b* loci. *MIR319a* locus has a 5-bp deletion and *MIR319b* ha a 1-bp insertion

as depicted.

Fig. 2   (A) Disruption of Mp*MIR319a* and/or *b* genes reduced the numbers of gemma

cups. Five plants were analyzed and the means of the numbers of gemma cups per each

thallus were shown. For statistical analysis, the analysis of variance followed by Tukey-

Kramer's honestly significant difference test was performed. Statistically significant

differences are indicated by different lowercase letters ($P < 0.05$).   (B) Appearances of

respective thalli.   Bar = 1 cm.   wt: Tak-1 wild-type plant, #5: *miR319a^{ge-1}* #5, #10:

miR319a^{ge-1} #10, #11: *miR319b^{ge-3}*#11, #15: *miR319b^{ge-3}*#15, #6: *miR319b^{ge-4}*#6, #5/15:

miR319ab #5/15.

Fig. 3   Disruption of Mp*MIR319a* and/or *b* genes reduced gemma formation. (A) The

number of gemma formed in each gemma cup. wt: Tak-1 wild-type plant, #5: *miR319a^{ge-}*

^{1} #5, #10: *miR319a^{ge-1}* #10, #11: *miR319b^{ge-3}*#11, #15: *miR319b^{ge-3}*#15. *p < 0.05 (Tukey

method).   (B) Images of single gemma cups. Bar = 1 cm

Fig. 4   Exon-intron structures of Mp*RKD* and Mp*R2R3-MYB21* genes and target

sequence modifications to construct miR319-resistant mMp*RKD* (A) and mMp*R2R3-*

MYB21 (B) genes. gMpRKD (A) and gMpR2R3-MYB21 genes of miR319-sensitive type

were also constructed. Vertical bars indicate conventional base-pairing, double dots

indicate G-U pairs. AUG and UGA are the initiation and termination codons of the

respective open reading frames. 5'UTR and 3'UTR are respective untranslated regions.

Fig.5   Introduction of miR319-resistant mMpRKD gene but not mMpMYB21 gene

resulted in thalli with fewer numbers of gemma cups.

Fig.6   Introduction of wild-type RKD gene (gMpRKD) induced normal development of

gametophytes respectively in Tak-1 (A,B) and Tak-2 (G, H). Introduction of miR319-

resistant type mutant RKD gene (mMpRKD) into Tak-1 induced antheridiophores with

normally-shaped stalked receptables but with irregular asymmetric shapes and antheridia

exposed out of antheridiophores (C,D,E,F). Introduction of mMpRKD into Tak-2 induced

archegoniophores with short digit shapes (I), however they retained normal development

of archegonia (J).

Fig. S1 Introduction of wild type RKD and miR319 resistant RKD genes.

(A)Wild-type *RKD* (w) and miR319 resistant *RKD* (R) genes were placed under the

direction of the promoter of Mp*RKD* gene （pRKD). The sequence segment

complementary to miR319 sequences downstream RKD termination codon was deleted

to obtain the miR319 resistant *RKD* gene.

(B) Phenotypes of growing thalli of transformants to which wild-type (w#1 and w#2)

and resistant (r#1 and #2) transgenes were introduced.

(C) The numbers of gemma cups were counted per thallus.

Fig.1 (A) Primary *MIR319a or b* RNAs are predicted to be transcribed from two annotated loci in the *M. polymorpha* genome. (B) The row wt shows the wild-type *MIR319a* locus sequence and is targeted by gRNA#1. Mutants #5 and #10 were selected for further analysis among obtained in-del mutants. (C) The row wt shows the wild-type *MIR319b* locus sequence and is targeted by gRNA#3 or #4. Mutants #11 and #15 from gRNA#3 treated-pool and mutants #6 and #9 from gRNA#4 treated-pool were selected for further analysis. X in the superscript *ge-x* indicates guide RNA species with which the respective mutant was created. (D) The genomic change of the double knock-out line for *MIR319a* and *b* loci. *MIR319a* locus has a 5-bp deletion and *MIR319b* ha a 1-bp insertion as depicted.

A.

B.

Fig. 2 (A) Disruption of Mp*MIR319a* and/or *b* genes reduced the numbers of gemma cups. Five plants were analyzed and the means of the numbers of gemma cups per each thallus were shown. For statistical analysis, the analysis of variance followed by Tukey-Kramer's honestly significant difference test was performed. Statistically significant differences are indicated by different lowercase letters (*P* < 0.05). (B) Appearances of respective thalli. Bar = 1 cm. wt: Tak-1 wild-type plant, #5: *miR319a^{ge-1}* #5, #10: *miR319a^{ge-1}* #10, #11: *miR319b^{ge-3}*#11, #15: *miR319b^{ge-3}*#15, #6: *miR319b^{ge-4}*#6, #5/15: *miR319ab* #5/15.

Fig. 3 Disruption of Mp*MIR319a* and/or *b* genes reduced gemma formation. (A) The number of gemma formed in each gemma cup. wt: Tak-1 wild-type plant, #5: *miR319a^{ge-1}* #5, #10: *miR319a^{ge-1}* #10, #11: *miR319b^{ge-3}*#11, #15: *miR319b^{ge-3}*#15. *p < 0.05 (Tukey method). (B) Images of single gemma cups. Bar = 1 cm

A. Mp*RKD*

AUG

UGA

5' UTR
3' UTR

gMp*RKD*

···CUCAUGACAUAGGGGCUCCCUUCCGUCCAAAGUUUCAAAUC···

‖ : ‖‖‖‖‖‖‖‖‖ ‖‖‖‖‖‖

3' CCCUCGAGGGAAGUCAGGUUC 5' miR319

mMp*RKD*

: ‖ ‖ : ‖ ‖::

···CUCAUGACAUACUGAAUCGCAAUCGAGCGGAGUUUCAAAUC···

B. Mp*R2R3-MYB21*

AUG

UGA

5' UTR
3' UTR

gMp*R2R3-MYB21*

···UGGAGCUCCCUUCAAGCCAAU···

‖‖‖‖‖‖‖‖‖‖‖‖ ‖‖‖‖

3' CCCUCGAGGGAAGUGAGGUUC 5' miR319

mMp*R2R3-MYB21*

‖‖ :‖ ‖‖ : :‖‖:

···UAGAAUUGCCAAGUUCUCAGU···

Fig. 4 Exon-intron structures of Mp*RKD* and Mp*R2R3-MYB21* genes and target sequence modifications to construct miR319-resistant mMp*RKD* (A) and mMp*R2R3-MYB21* (B) genes. gMpRKD (A) and gMpR2R3-MYB21 genes of miR319-sensitive type were also constructed. Vertical bars indicate conventional base-pairing, double dots indicate G-U pairs. AUG and UGA are the initiation and termination codons of the respective open reading frames. 5'UTR and 3'UTR are respective untranslated regions.

Fig.5 Introduction of miR319-resistant mMpRKD gene but not mMpMYB21 gene resulted in thalli with fewer numbers of gemma cups.

Fig.6 Introduction of wild-type RKD gene (gMp*RKD*) induced normal development of gametophytes respectively in Tak-1 (A,B) and Tak-2 (G, H). Introduction of miR319-resistant type mutant *RKD* gene (mMp*RKD*) into Tak-1 induced antheridiophores with normally-shaped stalked receptables but with irregular asymmetric shapes and antheridia exposed out of antheridiophores (C,D,E,F). Introduction of mMp*RKD* into Tak-2 induced archegoniophores with short digit shapes (I), however they retained normal development of archegonia (J).

Fig. S1 Introduction of wild type RKD and miR319 resistant RKD genes.

(A)Wild-type *RKD* (w) and miR319 resistant *RKD* (R) genes were placed under the direction of the promoter of Mp*RKD* gene （pRKD). The sequence segment complementary to miR319 sequences downstream RKD termination codon was deleted to obtain the miR319 resistant *RKD* gene.
 (B) Phenotypes of growing thalli of transformants to which wild-type (w#1 and w#2) and resistant (r#1 and #2) transgenes were introduced.
(C) The numbers of gemma cups were counted per thallus.

miR319a F	GTTCTGCAGGGACTGTTTTCAGCATCCC	*mir319a* mutant screening
miR319a R	GTTGGTACCTCGACAAGAATTGAGCGAGA	
miR319b F	GTTGTCGACGTGAACTCAGCGCTCATG	*mir319b* mutant screening
miR319b R	GTTGGTACCTACAATCTTTCAAATCAGTACGGAT	
PAM_MIR319a-174f-F	CTCGACAATCCGTCCTTGGACTGA	miR319a gRNA-1
PAM_MIR319a-174f-R	AAACTCAGTCCAAGGACGGATTGT	
PAM_MIR319b-161r-F	CTCGAGCTCCCTTCAGTCCAAGTC	miR319b gRNA-3
PAM_MIR319b-161r-R	AAACGACTTGGACTGAAGGGAGCT	
PAM_MIR319a-166f-F	CTCGCGATCTGTACAATCCGTCCT	miR319a gRNA-2
PAM_MIR319a-166f-R	AAACAGGACGGATTGTACAGATCG	
PAM-MIR319b-179f-F	CTCGTTGGCCCGACTTGGACTGAA	miR319b gRNA-4
PAM-MIR319b-179f-R	AAACTTCAGTCCAAGTCGGGCCAA	
IF-Hind-pRKD-F	ACGACGGCCAGTGCCAAGCTTGGACACCGGAGAGAAATCAA	Mp*RKD* gene construction
IF-Sal-gRKD-R	CGGGGATCCTCTAGAGTCGACTCATGAGACCAGGGTTGAGACAC	
Inf-Kpn-gRWP_RK-R	GGAAATTCGAGCTCGGTACCCGGAAGGTTATGATATCAAGTAGCA	
pMpMIR319a-83F	AGCGGAGAGCATTTCTCAGACC	sequencing of MIR319a promoter
pMpMIR319a-140F	GTCCCCAGACGCATGCAGGTCG	
pMpMIR319a-192F	GGCCGAGCGAGATCCCAGAGAT	
pMpMIR319a-254F	GCCTCGAAACGTAGTGTAGTGT	
pMpMIR319a-315F	CCATGCGCTGTAAGACATGGAC	
pMpMIR319a-375F	TGCGTGACTCTTCCGCATATTT	
pMpMIR319a-423F	TTCAGTCGAGAATGCGATCCCT	
pMpMIR319b-23F	CTGCAAGGCGATTAAGTTGGGT	sequencing of MIR319b promoter
pMpMIR319b-83F	CATTCTCGTACTTTGAACCGTC	
pMpMIR319b-135F	CCAGGTCTCCTTCGCGGAAAAG	
pMpMIR319b-195F	TGGATGCATAGCTTTCCATAAAG	
pMpMIR319b-255F	AGTGGAGATGAGGCGGAGTCAG	
pMpMIR319b-312F	CAAGTCTTCAAGGTGCACTACA	
pMpMIR319b-372F	TAATTGCGCTTGGAAACTACAC	
pMpMIR319b-413F	GTCCATGAGCGCTCATACGTCA	
pRWP_RK-53F	AAATGATACGAAGGAGGTGGACATC	sequencing of MpRKD construct
pRWP_RK-109F	GTGGCAATTTGTCACATTCTTCCAG	
pRWP_RK-162F	GGACGATTGAGCTTACAGAATCTGG	
pRWP_RK-227F	TCGATGGGTTGTGAGAGAACTTGAG	
pRWP_RK-287F	AGACACTAAGCGTGAATTTAGAGGA	
pRWP_RK-336F	AGCCAAAACTACGGACATCTTATTC	
pRWP_RK-395F	AGTGGTTTTCGTTCGGTACAGTCAC	
gRWP_RK-61F	GAATGTGTGGGTGTCACTGGTGGCT	
gRWP_RK-117F	GAATAAGTTTACGACCATGGCTGTA	
gRWP_RK-170F	GTTAGGAAATGTATGTTTCATCCTC	
gRWP_RK-230F	TTGCCGTCTTTAGTCATTGATTCTG	
gRWP_RK-288F	TTATAATTTGGAGATGCGGCTTAGC	
gRWP_RK-350F	TTGTAGCCACGATCTGTCTCCACAA	
gRWP_RK-405F	AGGATTCACGACTAATGGAGAATTC	

Table S1   primers and oligonucleotides used in this study