Title: Cell type-specific gene expression dynamics during human brain maturation

Christina Steyn[1,2], Ruvimbo Mishi[1,2], Stephanie Fillmore[1,2], Matthijs B. Verhoog[1,2], Jessica
More[1,2], Ursula K. Rohlwink[2,3], Roger Melvill[3], James Butler[2,4], Johannes M. N. Enslin[2,3],
Muazzam Jacobs[2,5,6,7], Sadi Quiñones[8,9], Chris G. Dulla[8], Joseph V. Raimondo[1,2,5], Anthony
Figaji[2,3] and Dorit Hockman[1,2,*]

Affiliations:
[1]Division of Cell Biology, Department of Human Biology, University of Cape Town, Cape Town,
South Africa
[2]Neuroscience Institute, University of Cape Town, Cape Town, South Africa
[3]Division of Neurosurgery, Department of Surgery, University of Cape Town, Cape Town,
South Africa
[4]Division of Neurology, Department of Medicine, University of Cape Town, Cape Town, South
Africa
[5]Institute of Infectious Disease and Molecular Medicine, University of Cape Town, Cape Town,
South Africa
[6]Division of Immunology, Department of Pathology University of Cape Town
[7]National Health Laboratory Service, South Africa
[8]Department of Neuroscience, Graduate School of Biomedical Sciences, Tufts University
School of Medicine, Boston, MA, USA
[9]Graduate School of Biomedical Science, Tufts University School of Medicine, Boston, MA,
USA
*Corresponding author

Abstract

The human brain undergoes protracted post-natal maturation, guided by dynamic changes in
gene expression. To date, studies exploring these processes have used bulk tissue analyses,
which mask cell type-specific gene expression dynamics. Here, using single nucleus (sn)RNA-
seq on temporal lobe tissue, including samples of African ancestry, we build a joint paediatric
and adult atlas of 54 cell subtypes, which we verify with spatial transcriptomics. We explore
the differences in cell states between paediatric and adult cell types, revealing the genes and
pathways that change during brain maturation. Our results highlight excitatory neuron
subtypes, including the LTK and FREM subtypes, that show elevated expression of genes
associated with cognition and synaptic plasticity in paediatric tissue. The new resources we
present here improve our understanding of the brain during a critical period of its
development and contribute to global efforts to build an inclusive cell map of the brain.

Introduction

The adult human brain is a complex assembly of diverse cell types, which has now been defined with unprecedented accuracy using single cell transcriptomics[1-4]. This adult transcriptomic signature is set up over a protracted period of development, which begins in the embryo and continues after birth as the brain matures. However, while the single cell diversity of the embryonic human brain has been explored[5,6], little is known about how these cell type-specific gene expression profiles change during childhood. Existing studies of gene expression dynamics during human brain development and maturation have used bulk transcriptomic approaches, which revealed a dramatic period of global gene expression change during the late foetal/early infancy transition, that stabilises during childhood and adolescence[6-10]. Bulk transcriptomics, however, cannot reveal the more subtle, cell type-specific changes in gene expression that drive brain maturation from childhood, through adolescence to adulthood.

Childhood and adolescence are periods of important changes in brain structure, during which neuronal connections are refined and strengthened. While synaptogenesis peaks in the early postnatal period, synaptic pruning activity begins during late childhood, peaks during adolescence, and then gradually decreases[11-13]. Together, synaptogenesis and synaptic pruning influence brain plasticity, a feature which allows the brain to adapt in response to experience[14]. Considering synaptic changes are most prominent in infancy, childhood and adolescence, these stages represent periods of enhanced susceptibility to environmental influence, as well as being periods of increased neuropsychiatric risk[15]. Describing the typical cell type-specific gene expression trajectories of the maturing brain will serve as an important reference to assess the effects of genetic perturbations and early adverse experiences on brain maturation. Furthermore, investigating the driving forces behind cell type-specific maturational processes may help in developing targeted therapies for neurological disease[16].

To this end, the Paediatric Cell Atlas (PCA)[17], a branch of the Human Cell Atlas (HCA) efforts to map the cellular diversity of the human body[3], aims to ensure that the benefits of single cell transcriptomics, which will likely include vast improvements in precision medicine, are available to children as well as adults from diverse populations[3,17]. Considering that Africa has the most genetically diverse[18] and youngest population[19] worldwide and that, by 2050, 37% of the world's children will grow up in Africa[20], the inclusion of the African paediatric population in the PCA's efforts has never been more pertinent. A reference paediatric cell atlas for the brain that includes data from African donors will contribute to developing the best treatment for locally prevalent conditions, such as tuberculosis meningitis and HIV infection[21,22]. In addition, studying the differences in gene expression dynamics between adult and paediatric brains may explain why the manifestation of neurological conditions and responses to therapies differ across the lifespan[17].

To contribute to these endeavours, we present a joint paediatric and adult temporal cortex
cell atlas, including samples from eight South African donors. To facilitate a direct comparison
of our snRNA-seq datasets with the existing adult temporal cortex cell atlas, we annotate our
data using the Allen Brain Map middle temporal gyrus (MTG) cell taxonomy[1]. We validate this
annotation approach using spatial transcriptomics analysis. In addition, we use *de novo*
marker gene analysis with machine learning tools to compare our paediatric and adult
datasets to the existing MTG cell taxonomy and highlight markers that define paediatric
versus adult cell states. Using differential gene expression analysis, we highlight six cell
subtypes, including two layer 2/3 excitatory neuron subtypes, that show differential
expression of genes involved in synaptic plasticity and cognition. Overall, we highlight the
subtle cell type-specific differences between the paediatric and adult brain and expand the
representation of diverse paediatric populations in the HCA.

92
93   **Results**
94
95   **A joint paediatric and adult temporal cortex cell atlas**
96
We generated snRNA-seq libraries from five paediatric and three adult donor temporal cortex
tissue samples using the 10x Genomics Chromium platform. These new libraries were
analysed alongside similar published datasets from one paediatric and three adult donors[23],
resulting in a total of 23 snRNA-seq datasets (including technical replicates) from 12
individuals (six paediatric and six adult) (Fig. 1a, Extended Data Table 1). The samples were
sequenced to a median depth of 19,853 reads per nucleus, with 176,012 nuclei remaining
after filtering low quality barcodes (Methods) (Extended data Fig. 1, Extended Data Table 2).
104
Using data integration and clustering (Methods), we aligned similar cell types across the 23
datasets (Fig. 1a, Extended data Fig. 1h), yielding 40 clusters (Extended data Fig. 1i). To
facilitate a direct comparison between our datasets and the current draft human brain cell
atlas, we used *Seurat*'s label transfer method[24] to classify each nucleus according to the Allen
Brain Map MTG atlas[1] (Fig. 1b; Extended Data Table 3). Of the 75 reference cell types, 21
neuronal subtypes were absent from our datasets (Extended Data Table 3). The majority of
the absent neuronal cell types are rare (<2% of all cells; see Extended Data Table 3), and
therefore their absence is likely due to the lower proportion of neurons in our datasets
compared to the reference atlas, which was made using cell-sorting to enrich for neurons[1].
114
As an initial validation of the label transfer, we confirmed that the majority of annotated cell
types expressed the expected cell type-specific marker genes[1] (Fig. 1c). Additionally, we
performed a correlation analysis to compare the transcriptomic similarity of the annotated
cell types to the reference MTG cell types[1] (Fig. 1d). The non-neuronal cell types showed high
correlation with the corresponding reference cell types, as well as high specificity (Fig. 1d).

Similarly, most neuronal subtypes showed high correlation with the corresponding subtype
in the reference datasets, however, there was also correlation with other subtypes within
their class. Exceptions were Exc_L4-6_FEZF2_IL26, which expressed the excitatory neuron
marker gene *FEZF2* but correlated more strongly with microglia, and Exc_L5-
6_THEMIS_FGF10, which expressed the inhibitory marker *LHX6* and correlated more strongly
with the inhibitory neuron subtypes (Fig. 1c-d). The overall cell composition of the paediatric
and adult samples was very similar. Oligodendrocytes and oligodendrocyte precursor cells
(OPCs) were the most common non-neuronal cell types and Exc_L2_LAMP5_LTK was the most
common neuronal cell subtype (Fig. 1e; Extended data Figure 2a-c).
129
Neuronal clusters had a greater number of expressed genes and unique molecular identifiers
(UMIs) compared to non-neuronal cells (Extended data Figure 3a). Similar to previous snRNA-
seq analyses of the adult brain[1,2], excitatory neurons had a greater number of genes detected
per nucleus than inhibitory neurons, with Exc_L2-3_LINC00507_FREM3 and Exc_L4-
5_RORB_FOLH1B among the cell types with the highest median gene detection. When
comparing the paediatric to adult samples, only two cell subtypes showed significant
differences in the number of genes (Exc_L2_LAMP5_LTK and Exc_L3-5_RORB_FILIP1L) and
UMIs (Exc_L2_LAMP5_LTK and Exc_L2-3_LINC00507_FREM3) between the age categories,
with the paediatric samples having a significantly larger value in each case (Extended data
Figure 3b-c). This result points towards higher transcriptional diversity in these neuronal
subtypes during childhood.
141
142
143 **Spatial mapping of cell types reveals similar tissue cytoarchitecture in adult and paediatric**
144 **temporal cortex**
145
Next, we used spatial transcriptomics to verify the positions of our annotated cell types within
the layered structure of the temporal cortex. We generated 10x Genomics Visium Spatial
Gene Expression datasets for frozen tissue sections from adult (31-year-old) and paediatric
(15-year-old) temporal cortex samples (two sections each; Extended Data Table 1; Extended
data Fig. 4). The four Visium libraries were sequenced to a median depth of 87,178 reads per
spot (median of 5,878 UMIs and 2,745 genes per spot) (Extended Data Table 4).
152
Using *cell2location*[25] (Methods), we calculated cell type abundance estimates for each Visium
spot, with our annotated snRNA-seq dataset as a reference. As seen in our snRNA-seq
datasets, oligodendrocytes were the most common cell type, while Exc_L2_LAMP5_LTK was
the most abundant neuronal cell type (Extended data Fig. 5a). Spatial plots of estimated cell
abundance for a selection of cell types revealed that the annotated cell types mapped to their
expected cortical layer locations across all tissue sections (Fig. 2a-b; Extended data Fig. 5b-c).
The two cell types that did not correlate strongly with the expected MTG reference atlas cell
types in our snRNA-seq analysis, Exc_L4-6_FEZF2_IL26 and Exc_L5-6_THEMIS_FGF10, were

most strongly distributed in layers 4-6, providing support for their annotation (Extended data
Fig. 5d). The layered pattern of the annotated cell types coincided with the spatial expression
of known cortical layer marker genes[1,26,27] in the Visium datasets (Fig. 2c). These layered
expression patterns were verified for a subset of layer-specific marker genes using *in situ*
hybridisation analysis on frozen temporal cortex tissue sections from the same 31-year-old
and 15-year-old donors (Extended data Fig. 6).

To confirm the co-location of cell types within the layered structure of the temporal cortex,
non-negative matrix factorization (NMF) was performed using the cell abundance estimates
from *cell2location*, resulting in 12 cellular compartments (Fig. 2d). The NMF weights for the
identified cellular compartments were visualised across the Visium samples to assess their
spatial distribution (Fig. 2e). In both the paediatric and adult datasets, there was clear co-
location of the expected neuronal cell types within overlapping compartments across the
cortical layers, including layers 1-3 (factor_9), layers 3-5 (factor_3) and layers 4-6 (factor_7).
Several excitatory neuron subtypes formed discreet cellular compartments, including Exc_L2-
3_LINC00507_FREM (factor_4), Exc_L5-6_THEMIS_CRABP1 (factor_11) and Exc_L5-
6_FEZF2_ABO (factor_6). The two astrocyte subtypes were confirmed to have distinct
distributions profiles, with Astro_L1-2_FGFR3_GFAP (factor_5) located primarily in layer 1
and the white matter, and Astro_L1-6_FGFR3_SLC14A1 (factor_8) more widely distributed.
The remaining non-neuronal cell types were largely associated with factors located in layer 1
and the white matter.

Overall, our spatial transcriptomic analyses provide support for our annotation approach,
showing the expected spatial distribution of annotated cell types, and revealing a similar
tissue cytoarchitecture in adult and paediatric temporal cortex tissue.

A machine learning approach identifies gene pathways that distinguish paediatric and adult
cell states

To establish a standardized and scalable approach for defining cell types, it has been proposed
to use the minimum combination of gene markers that can classify a cell type and distinguish
it from other cell types[28,29]. Towards achieving this, Aevermann et al. (2021)[28] developed the
machine learning tool, *NS-Forest V2.0*, which they applied to the Allen Brain Map MTG
dataset. Ideally, theses MTG minimal markers would be conserved in similar datasets to
facilitate accurate comparisons across different studies[30]. We found that the MTG cell atlas
minimal markers[28] are indeed highly expressed in the expected cell types (Extended data Fig.
7).

Next, we applied the *NS-Forest V2.0*[28] algorithm to our datasets, firstly, to assess if the
identified minimal markers overlap with the published MTG minimal markers and, secondly,

to identify combinations of marker genes that distinguish the paediatric and adults states of
each cell type (Methods). 151 paediatric and 149 adult minimal marker genes were identified
across 53 cell types (Fig. 3; Extended data Table 5). There was little overlap with the MTG
atlas, with only 11 paediatric (7.3%) and 4 adult (2.7%) minimal markers showing overlap with
existing datasets[1,28] (Fig. 3; Extended data Table 5). On the other hand, there was a greater
overlap in minimal markers between the paediatric and adult datasets, with 35 markers
(~23%) present in both lists.

Our minimal marker analysis revealed improved markers for several cell types when
compared to the reference MTG cell atlas. *DDR2* shows high specificity for Inh_L1-
2_PAX6_CDH12 (Fig. 3; Extended data Fig. 8a), while the existing minimal marker for this cell
type, *TGFBR2*, is more highly expressed in the microglia and endothelial cells (Extended data
Fig. 7; Extended data Fig. 8b). Similarly, *SEMA3E* is very specific to Exc_L5-6_FEZF2_ABO (Fig.
3; Extended data Fig. 8c), while an existing minimal marker for this cell type, *SULF1*, is also
expressed at appreciable levels in other neurons (Extended data Fig. 7; Extended data Fig.
8d). Additionally, UMAP analysis of the annotated datasets using our minimal marker gene
list for each age group, in comparison to an equivalent number of random genes, resulted in
better grouping of the cell subtypes into clusters (Fig. 4 a-b). While the cell clusters are not as
clearly separated as the original UMAP plot, (see Fig. 1a), this analysis reveals that our short
list of ~150 marker genes captures much of the underlying transcriptomic diversity in our
datasets.

To explore the cellular functions of the paediatric and adult minimal marker genes
respectively, we used Gene ontology (GO) analysis (Extended Data Table 6), which revealed
significant enrichment of GO terms related to neuronal development and cell signalling for
both datasets. Interestingly, only the paediatric dataset was enriched for cellular migration
terms (Extended Data Table 6). Genes included in these sets were *RELN*, *CXCL14* and *SEMA3A*,
which play roles in neuronal migration during brain development[31-33]. On the other hand, only
the adult datasets were enriched for extracellular matrix and cell death terms (Extended Data
Table 6). These broad analyses of minimal marker gene function indicate that genes involved
in neuronal development pathways remain key to neuronal identity, alongside functional
signalling molecules, as the brain matures and in adult life. Our results also suggest that genes
involved in cellular migration processes may continue to define cell states during childhood
and adolescence.

To further assess the difference in cell type-specific markers between our paediatric and adult
datasets, we expanded our analysis to include the top 50 genes identified as cell type
classification features for each cell type. For most cell types, the majority of these top markers
(>20 genes) were shared between our paediatric and adult datasets (Fig. 4c; Extended Data
Tables 7-8). The non-neuronal cell types showed the highest number of shared marker genes
(≥40). GO analysis of these shared marker genes showed significant enrichment of terms

related to cell type-specific functions, such as "leukocyte proliferation" for microglia, and
"myelin sheath" for oligodendrocytes (Extended Data Table 6). These results suggest that
paediatric and adult non-neuronal cell states are relatively similar, although it is likely that
more diversity in the marker gene profiles could be revealed with subdivision of the cells into
further subtypes.
245
The cell types with the fewest shared markers between our paediatric and adult datasets
(≤10) were mostly represented by fewer than 20 nuclei and therefore the lack of marker
overlap between our datasets is possibly due to sampling bias (Extended Data Table 6). An
exception was Exc_L4-6_FEZF2_IL26, represented by 2,579 nuclei, which had no shared
genes, two adult-specific genes (*SNAP25* and *CALM1*) and four paediatric-specific genes
(*MEF2A*, *MEF2C*, *DOCK4* and *PLXDC2*) (Fig. 4c; Extended Data Tables 7-8). Interestingly, the
paediatric-specific markers for this cell type are implicated in neuronal development
processes, including synaptic plasticity[34], dendritic branching[35] and neuronal proliferation[36].
The cell type with the greatest number of paediatric-specific marker genes was Exc_L4-
5_RORB_DAPK2 (29 paediatric vs 3 adult markers) (Fig. 4c; Extended Data Tables 7-8). GO
analysis of the 29 paediatric-specific marker genes for this cell type revealed enriched terms
related to synapse development, while similar analysis of the 19 shared genes together with
the three adult-specific genes highlighted terms involved in synaptic function (Extended Data
Table 6).
260
261 Overall, our expanded marker gene analysis suggests that neuronal cell types show greater
262 dissimilarity between their paediatric and adult states than non-neuronal cells, and reveals
263 several paediatric-specific markers that reflect the less mature state of paediatric neuron
264 subtypes.
265
266
267 **Differential gene expression analysis highlights six cell subtypes and enriched expression of**
268 **genes associated with synaptogenesis and cognition in paediatric samples.**
269
While our minimal marker gene analysis provides insight into the genetic signatures that
distinguish cell types from one another over the course of brain maturation, we were also
interested in investigating how the general transcriptomic profile of each cell type differs in
children when compared to adults. In particular, we sought to identify genes that were
upregulated in the paediatric cell populations and thus might be involved in childhood brain
development and function. To this end, we conducted cell type-specific differential gene
expression analysis with *DESeq2*[37] (Methods).
277
In total, we detected 166 unique significantly differentially expressed genes (DEGs) across 12
out of the 54 annotated cell types, with some DEGs associated with multiple cell types (Fig.
5a; Extended data Table 9). When the magnitude of fold change of the DEGs was considered,

143 genes across six cell types changed by at least 10%, with a positive fold change indicating
higher expression in our paediatric samples (Fig. 5b-c). The six cell types included four
excitatory neuron subtypes, one inhibitory subtype and one astrocyte subtype (Fig. 5b-c;
Extended data Table 10). For the majority of DEGs, the change in expression was accompanied
by a corresponding change in the percentage of nuclei expressing the gene (Extended data
Table 10). We assessed the expression patterns of a subset of DEGs in our Visium datasets
using *BayesSpace*[38] (Methods), confirming that the genes were expressed at higher levels and
in a greater number of spots in the 15-year-old compared to the 31-year-old (Extended data
Fig. 9).
290
The layer 2/3 excitatory neuron subtypes, Exc_L2_LAMP5_LTK and Exc_L2-
3_LINC00507_FREM3, shared several upregulated DEGs that are developmentally regulated
in the mammalian brain (Fig. 5c-d). *FNBP1L* (*TOCA-1*) regulates neurite outgrowth[39] and is
associated with intelligence[40]. In line with our findings, its expression declines over the course
of brain maturation in the rat[39]. Similarly, both *KCNG1,* a voltage gated-potassium channel
(Kv6.1), and *MYO16* (*MYR8*), an unconventional myosin protein, decrease in expression with
age in the mouse[41] and rat[42] brain, respectively. These findings indicate that previously
reported expression dynamics for these genes in mammalian models are conserved in the
human temporal cortex, with higher expression in children. Importantly, our analysis reveals
that these patterns are specific to two layer 2/3 excitatory neuron subtypes.
301
Two significantly upregulated genes in the layer 2/3 excitatory neurons, *LUZP2* and *RERGL*
(Fig. 5c-d), are associated with neuropsychiatric disorders, as well as brain cancers. *LUZP2*, a
leucine zipper protein, is associated with Alzheimer's Disease and schizophrenia, as well as
cognitive performance in the normal elderly population[43]. It is differentially expressed in
several cancers, including low grade gliomas[44]. Similarly, *RERGL* is differentially expressed in
schizophrenia[45], while its expression is significantly higher in meningiomas compared to
healthy tissue[46]. Our results suggest that these genes likely play a role in the maturation of
layer 2/3 excitatory neurons.
310
The layer 3-5 excitatory neuron subtypes, Exc_L3-5_RORB_ESR1 and Exc_L4-5_RORB_DAPK2,
shared 3 upregulated DEGs, which have known roles in the development and reorganisation
of neuronal connections (Fig. 5c,e). *XKR4* is a member of a plasma membrane protein family
involved in signalling cellular compartments for engulfment, such as during synaptic
pruning[47]. These proteins decrease in expression with age in the mouse brain[48]. *TENM1*, is a
member of the teneurin transmembrane protein family that regulate cytoskeletal
organisation and neurite outgrowth, as well as shaping synaptic connections[49-51]. *AGBL1*
(*CCP4*) is a glutamate decarboxylase that mediates deglutamylation of tubulin[52]. This process
is essential for the maintenance of neuronal tubulin, and when disrupted, contributes to
neurodegeneration[52,53]. Our results point towards a role for these genes specifically in Exc_L3-

5_RORB_ESR1 and Exc_L4-5_RORB_DAPK2 neurons, where they possibly contribute to the
re-shaping of neuronal connections during brain maturation.
323
The majority of the DEGs in the four neuronal subtypes were not shared across the cell types
and represent promising candidates for future explorations into molecular mechanisms
guiding cell type-specific brain maturation. For example, *FGF13* (*FHF2*) was significantly
upregulated in Exc_L3-5_RORB_ESR1 (Fig. 5c). *FGF13* decreases in expression with age in the
mouse brain, where it regulates post-natal neurogenesis[54] and axonal formation[55]. Similarly,
PLPPR1 (*PRG1*) was significantly upregulated in Exc_L2-3_LINC00507_FREM3 (Fig. 5c). *PLPPR1*
is higher in the postnatal mouse hippocampus than in the adult[56] and is also known to
regulate axon growth by modulating cytoskeletal dynamics[57].
332
In Inh_L2-6_VIP_QPCT, a single lncRNA, *LINC00276*, was upregulated (Fig. 5b). While this non-
coding RNA has been shown to be expressed in the brain[58], nothing is known of its function
there. Similarly, in Astro_L1-6_FGFR3_SLC14A1, a single non-coding gene, *AC109439.2* (*CTB-
1I21*), was significantly upregulated (Fig. 5c, f). This lncRNA has recently been identified as a
protective factor in oesophageal cancer[59] and glioma[60]. One of the downregulated genes in
this cell type, *ADAM28*, a metalloproteinase, has been found to be upregulated in breast and
lung cancers[61]. These results provide new molecular candidates to expand our understanding
of molecular mechanisms of astrocyte maturation.
341
Genes associated with intelligence quotient (IQ) and educational attainment (EA) have
recently been shown to be enriched in adult temporal lobe cortical neurons, especially the
Exc_L2-3_LINC00507_FREM3 subtype[62]. Since childhood is a key period of cognitive
development[63], we explored whether the same genes were found amongst our DEGs. Of the
137 DEGs found in at least one neuronal cell type, 15 (11%) are known to be significantly
associated with EA[64] and 4 (3%) with IQ[65]. These included several genes enriched in paediatric
samples, such as *MYO16*, *KCNG1* and *LUZP2* (Extended data Table 10).
349
Overall, our differential expression analysis highlights six cell subtypes that show significant
changes in gene expression between children and adults. Several of the genes that are
upregulated in in children have known roles in brain development and have been associated
with cognitive ability. Our analysis builds on this knowledge by implicating specific paediatric
cell subtypes and provides new candidate genes that likely contribute to cell type-specific
maturation processes.
356
357 **Gene pathways involved in synaptic development and functioning are enriched in**
358 **paediatric cell types**
359
360 We next used gene set enrichment analysis (GSEA) to conduct a broad analysis of the gene
361 pathways that are differentially regulated across all brain cell types during brain maturation

362   (Methods). GSEA aggregates the information from many genes to identify enriched functional
363   pathways, allowing us to interrogate the gene signature changes across all cell types,
364   including those that did not show any significant DEGs[66].
365
In total, 2,003 GOBP terms where enriched in the paediatric samples compared to the adults,
while 866 were depleted (p<0.01 and q<0.1) (Extended data Table 11). When focusing on the
25 most frequently enriched terms, the majority (11 terms) were associated with cellular
respiration pathways (Fig. 6; Extended data Table 11). Six of the most commonly enriched
terms were linked to synaptic development and functioning. A similar trend was observed
when focussing on the cell types from our differential expression analysis, with gene
expression, cellular respiration and synapse development pathways dominating the top
enriched terms (Fig. 5g).
374
Focusing on pathways that are depleted in the paediatric brain, four of the top ten depleted
terms, including the top term, were associated with neuronal ensheathment (Fig. 6).
Interestingly, none of these terms were significantly enriched in oligodendrocytes or OPCs,
while they were associated with neuronal sub types, astrocytes and microglia. The remainder
of the top ten depleted terms included neuronal morphogenesis, cell adhesion and gene
expression pathways.
381
382   Overall, our GSEA analysis points towards putative genetic pathways that may drive the
383   differences in synaptic plasticity between the paediatric and adult brain. Pathways related to
384   reorganising and strengthening synapses may be enhanced across multiple cell types during
385   childhood, while those required to limit synaptic growth, such as axonal ensheathment
386   pathways, may need to be suppressed.
387
388
389   **Discussion**
390
The brain is the most complex organ in the human body, which continuously changes as we
mature and age. Existing studies exploring the transcriptomic changes across the full span of
brain maturation have used bulk transcriptomic techniques[6-10], which drown out the subtle
molecular events taking place within specific cell types. Here, we unmask these processes,
using single cell transcriptomics to compare similar cell types between paediatric and adult
datasets.
397
To facilitate accurate comparisons of cell types across age groups, we used the existing Allen
Brain Map MTG cell atlas[1] to annotate our datasets. This demonstrated that the reference
atlas, generated from eight adult snRNA-seq datasets, is indeed generalisable[30], and can be
used to classify cell types in similar datasets from samples of different ages. This
generalisability is essential for healthy human reference atlases to serve as a baseline to

improve our understanding of human development and disease[3]. Our machine-learning
marker gene analysis also shows that while the cell type classifications, which are based on
the expression of thousands of genes, can be transferred onto new datasets, the minimal
markers that define the cell types do vary across datasets. Very few of our *NS-Forest* minimal
markers overlap with the existing MTG cell atlas minimal markers[1,28] and some provide better
discrimination between cell type than the existing markers. These results highlight that the
minimal markers that currently define the MTG cell taxonomy will likely need to be revised
as more samples are made available to ensure that the cell type classification is as widely
applicable as possible.

Similar to previous analyses of aging in the mouse[66], our integrated analysis of paediatric and
adult datasets showed there is little change in cell type composition within the temporal
cortex during human brain maturation. However, both our minimal marker analysis and
differential gene expression analysis highlight differences in cell states between paediatric
and adult cell types. GO analysis of our minimal marker genes revealed a signal for migratory
pathways amongst the paediatric minimal markers, including genes that code for
chemoattractants, such as *RELN* and *CXCL14*. While these genes are expressed in the
mammalian adult brain[26,67,68], our analysis suggests that they play a greater role in defining
cell identity in childhood, where these signals possibly function to mediate processes such as
dendritic outgrowth and the rearrangement of neuronal synaptic connections.

In recent years, the excitatory pyramidal neurons in the supragranular layers of the MTG have
been shown to have high transcriptional diversity[1,69] and to possess unique features including
exceptionally large arborisations[70] and electrophysiological properties that impact signal
integration and encoding[71-74] in ways that may enhance neuronal computational abilities,
contributing to human cognition. Since cognitive ability is a key feature that is established
during childhood[68], our differential expression analysis offers an extraordinary opportunity to
explore how cell type-specific gene expression dynamics contribute to cognitive
development. We highlight six cell subtypes that show significant gene expression changes
between childhood and adulthood. Interestingly, two of these cell types were the layer 2/3
excitatory neurons, Exc_L2_LAMP5_LTK and Exc_L2-3_LINC00507_FREM3, that have recently
been associated with human cognition[62]. In line with these findings, several of the DEGS
shared by these cell types, including *FNBP1L*[40] and *LUZP2*[43], have been implicated in cognitive
ability and intelligence. Overall, our data points towards genes and pathways that likely play
key roles in cognitive development specifically within these layer 2/3 excitatory neurons.

The relatively low number of genes and cell types implicated in our differential expression
analysis in comparison to similar studies in mouse[66] suggests that the difference between the
paediatric and adult brain are subtle. However, the inherent high variability in human gene
expression data may potentially mask some of the differential gene expression in our limited
sample. As the HCA database for the human temporal cortex expands, it will be important to

build on these analyses with more samples for each developmental stage to provide more
support for our findings.
446
We have provided the first single cell gene expression datasets for the brain that includes
data from black South African donors, thus increasing the diversity of the HCA database. Our
paediatric datasets will form important baseline references for future studies aiming to
explore how locally important challenges to child brain health, including infectious diseases
and traumatic brain injury, impact on the normal gene expression profiles. Importantly, these
investigations will contribute to the development of effective treatments, that are tailored to
specific needs of paediatric patients.
454
455

456   **Methods**
457

Human samples
Ethical approval was granted for the collection and use of paediatric and adult human brain
tissue by the University of Cape Town Human Research Ethics Committee (UCT HREC REF
016/2018; sub-studies 146/2022 and 147/2022). The human brain tissue samples used to
generate new 10x Genomics snRNA-seq and Visium datasets were obtained by informed
consent for studies during temporal lobe surgical resections to treat epilepsy and/or cancer
performed at the Red Cross War Memorial Children's Hospital and Constantiaberg Mediclinic
in Cape Town, South Africa. The samples used in this study were of temporal cortex origin and
represent radiologically and macroscopically normal neocortex within the pathological
context (details in Extended Data Table 1). Upon resection, samples were placed in
carbogenated ice-cold artificial cerebral spinal fluid (aCSF) containing in (mM): 110 choline
chloride, 26 NaHCO$_3$, 10 D-glucose, 11.6 sodium ascorbate, 7 MgCl$_2$, 3.1 sodium pyruvate, 2.5
KCl, 1.25 NaH$_2$PO$_4$, and 0.5 CaCl$_2$ (300 mOsm) and immediately transported to the laboratory
(~20 minutes). Tissue blocks containing the full span from pia to white matter were prepared
and either flash frozen in liquid nitrogen or embedded in optimal cutting temperature
compound (OCT) and stored at -80°C. The OCT-embedded samples were flash frozen in a
10×10 mm^2 cryomold which was either frozen directly in liquid nitrogen or placed in a
container of isopentane (Merck) which was in turn placed in liquid nitrogen at the same level
as the isopentane. The publicly available snRNA-seq datasets[23], generated from samples
obtained during elective surgeries performed at Universitair Ziekenhuis Leuven, Belgium,
were downloaded from the Sequence Read Archive database.
479

Nuclei isolation for snRNA-seq
Nuclei were isolated according to a protocol adapted from Habib et al. (2017)[75] and the 10X
Genomics nuclei isolation protocol (CG000124, User Guide Rev E). Frozen brain tissue was
homogenised in a dounce-homogeniser containing 2 ml ice-cold lysis solution (Nuclei EZ Lysis
Buffer [Sigma-Aldrich, NUC101] or Nuclei PURE Lysis buffer [Sigma-Aldrich, NUC201] with 1

mM dithiothreitol [DTT, Promega, P1171, US] and 0.1% Triton X-100 [Sigma-Aldrich, NUC201-
1KT, US]). Homogenisation was done 20 times with the loose pestle A followed by 20 times
with the tight pestle B. An additional 2 ml lysis solution was added, and the sample was
incubated for 5 minutes on ice. The sample was centrifuged at 500 x g for 5 minutes at 4°C
after which the supernatant was discarded and the nuclei resuspended in 3 ml ice cold nuclei
suspension buffer (1x phosphate-buffered saline [PBS, Sigma-Aldrich, P4417-50TAB, US]),
0.01% bovine serum albumin [BSA, Sigma-Aldrich, A2153-10G, US], and 0.2 U/µl RNAsin Plus
RNase inhibitor [Promega, N2615, US]). Resuspended nuclei were passed through a 40 µm
filter and centrifuged at 900 x g for 10 minutes at 4°C. The supernatant was discarded and
pelleted nuclei were resuspended in 3 ml blocking buffer (1xPBS [Sigma-Aldrich, P4417-
50TAB, US], 1% BSA [Sigma-Aldrich, A2153-10G, US], 0.2 U/µl RNAsin Plus RNase inhibitor
[Promega, N2615, US]).
497
To remove myelin debris, 30 µl of myelin removal beads [Miltenyi Biotec. 130-096-733, US]
was added to the solution which was mixed by gently pipetting 5 times. The sample was
incubated for 15 minutes at 4°C after which it was mixed with 3 ml blocking buffer and
centrifuged at 300 x g for 5 minutes at 4°C. The supernatant was removed and the nuclei were
resuspended in 2 ml clean blocking buffer. The sample was transferred to a 2 ml tube and
placed on a Dynamag magnet for 15 minutes at 4°C. The supernatant was transferred to a
new tube and stored on ice. An aliquot of trypan blue stained nuclei was counted using a
haemocytometer to determine the nuclei concentration and the volume to use in snRNA-seq
library preparation.
507

10X Genomics snRNA-seq library preparation
snRNA-seq library preparation was carried out using the 10x Genomics Chromium Next Gen
Single Cell 3' Reagent Kit (v3.1) according to manufacturer's protocols (CG000204, User Guide
Rev D), targeting 10,000 nuclei per sample. At step 2.2d and 3.5e, the libraries were amplified
using 11 cycles and 13 cycles, respectively. Library quality and concentration was assessed
using either the TapeStation or Bioanalyser (Agilent) and Qubit (Invitrogen) at the Central
Analytical Facility (CAF, University of Stellenbosch). cDNA libraries were sequenced by
Novogene (Singapore) on either the Illumina HiSeq or NovaSeq system using the Illumina High
Output kits (150 cycles).
517

snRNA-seq read alignment and gene expression quantification
Fastq files were aligned to the human reference transcriptome (GRCh38) and quantified using
the count function from the 10X Genomics Cell Ranger v6.1.1 software (Cell Ranger, RRID
SCR_017344) (Code availability: script 1). The inclusion of introns was specified in the count
function. An automatic filtering process was performed to remove barcodes corresponding
to background noise which have very low UMI counts.
524

snRNA-seq quality control

snRNA-seq quality control

The resulting count matrices were processed using a pipeline adapted from the Harvard Chan

Bioinformatics Core (https://hbctraining.github.io/scRNA-seq_online/). The filtered gene

barcode matrix for each sample was imported into R (V.4.2.0) using the Read10X function

from the Seurat (v.2.0)[24]. Nuclei-level filtering was performed to remove poor quality nuclei

according to their number of UMIs (nUMIs) detected, number of genes detected (nGene),

number of genes detected per UMI (log10GenesPerUMI), and the fraction of mitochondrial

read counts to total read counts (mitoRatio) (Code availability: script 2). Nuclei that met the

following criteria were retained: nUMI > 500, nGene > 250, log10GenesPerUMI > 0.8 and

mitoRatio < 0.2. Gene-level filtering was performed to remove genes that had zero counts in

all nuclei, remove genes expressed in fewer than 10 nuclei, and remove mitochondrial genes

from the gene by cell counts matrix. Three doublet removal tools namely

DoubletFinder[76](Code availability: script 3), DoubletDecon[77] (Code availability: script 4), and

Scrublet[77] (Code availability: script 5,6) were used to identify doublets for each dataset

individually. The sample-specific parameters of each of the tools were adjusted according to

the specified guidelines. To achieve a balance between the false positive and false negative

rate of the different doublet detection tools, all doublets identified by DoubletFinder as well

as the intersection of the doublets identified by DoubletDecon and Scrublet, were removed[77].

snRNA-seq data normalization, integration and clustering

Principal component analysis was performed to evaluate known sources of within-sample

variation between nuclei, namely the mitoRatio and cell cycle phase (Code availability: script

7). The UMI counts of the 3000 most variable features were normalised and scaled on a per

sample basis by applying Seurat's SCTransform function with mitoRatio regressed out. A

Uniform Manifold Approximation and Projection (UMAP) analysis was performed on the

merged object to assess whether integration was necessary. The datasets were subsequently

integrated using Seurat's SelectIntegrationFeatures, PrepSCTIntegration,

FindIntegrationAnchors, and IntegrateData functions (Code availability: script 7). To cluster

the datasets following integration, dimensionality reduction was first performed using UMAP

embedding, specifying 40 dimensions (Code availability: script 8). The Seurat FindClusters

function was then applied at a resolution of 0.8.

snRNA-seq cluster annotation

Label transfer was performed using Seurat's TransferData function with Allen Brain Map MTG

atlas[1] as a reference dataset (Code availability: scripts 9-10). This resulted in each barcode in

the query dataset receiving a predicted annotation based on a similarity score to an

annotated cell type in the reference. To validate the annotation, the expression of known

marker genes was assessed. Additionally, cosine similarity scores were computed to compare

the transcriptomic similarity of each of the 54 annotated query cell types to the 75 reference

middle temporal gyrus cell types using the SCP package (https://github.com/zhanghao-

[njmu/SCP)](#) (Code availability: script 11). This was achieved by computing cosine similarity
scores for each pair of query and reference cell types using the expression of the top 2000
shared highly variable features between the query and reference datasets. The log
normalised expression counts were used for this purpose (RNA assay, data slot).
569

NS-Forest machine learning marker analysis of snRNA-seq datasets

The NS-Forest tool (v2.0)[28,29] was used to identify combinations of marker genes uniquely
defining each annotated cell type (Code availability: script 12-14) in the paediatric and adult
datasets separately. The number of nuclei per sample was randomly down-sampled to that
of the sample with the fewest nuclei (n=4865). A random-forest model was used to select a
maximum of 50 marker genes per cell type based on them being both highly expressed as
well as uniquely expressed within a cell type compared to other cell types (i.e., the top Gini
Index ranked features with positive expression values). The number of trees chosen for this
model was 35,000, the cluster median expression threshold was set to the default value of
zero, the number of genes used to rank permutations of genes by their F-beta-score was 6,
and the beta weight of the F score was set to 0.5 allowing the outputs to be directly compared
to the Allen Brain Map MTG atlas minimal markers[1]. To assess the relevance of these markers
in terms of their capacity to distinguish different cell types in a UMAP analysis, the SCT and
integration methods were repeated using either a random set of genes or the NS-Forest
markers as anchors[28] (Code availability: script 15).
585

DESeq2 age-dependent differential gene expression analysis of snRNA-seq datasets

DESeq2[37] was used to identify genes that were differentially expressed with age (Code
availability: script 16). The unnormalized counts were aggregated across all nuclei for each
cluster and sample to generate a 'pseudobulk' counts matrix with the counts from technical
replicates collapsed to the level of biological replicates. Genes were filtered to only include
those expressed in more than 10% of nuclei for a given cell type. Principal component analysis
was performed on each cell type separately in order to assess the variation between samples
and determine which variables were contributing most to inter-sample variation from a set
of possible variables. The collapsed counts served as input into DESeq2's
DESeqDataSetFromMatrix function in which the design formula ~single_cell_chemistry +
age_group was specified to treat the age_group (paediatric vs adult) as the variable of interest
while the effect of single_cell_chemistry (version2 vs version3 chemistry) was regressed out.
A hypothesis test was performed using the Wald test. The null hypothesis for each gene was
that there is no difference in gene expression between the sample groups (i.e Log2 Fold
Change = 0). A Wald test statistic was determined for each gene together with the associated
p-value after which the p-values were adjusted for multiple testing using the Benjamini and
Hochberg method. Positive log2 Fold Changes represent genes which are upregulated in
paediatric samples compared to adult samples (p_{adj} <0.05).
604

Pathway enrichment analysis of snRNA-seq datasets

GO analysis of NS-Forest marker genes was performed on the gProfiler web server[78] using default settings ($p_{adj} < 0.05$) with "highlight diver terms in GO" selected.

Neuronal DEGs identified by DESeq2 (see Extended data Table 10) that were associated with EA and IQ were determined by comparing the list of neuronal DEGS to the EA and IQ gene lists used by Driessens et al. (2023)[62], which were subsets of the lists from Lee et al. (2018)[64] and Savage et al. (2018)[65] respectively.

GSEA on the DESeq2 output for all genes was performed using the Broad Institute's GSEA software (https://www.gsea-msigdb.org/gsea/msigdb) (Code availability: script 17). The gene lists for each cell type were queried against the C5 GO Biological Processes collection comprising of gene sets derived from the GO Biological Process ontology. The input lists of genes were ranked according to the $-\log(\text{p-value}) * \log_2\text{FoldChange}$ for each gene. The parameters specified to the GSEA function included number of permutations (nperm)=1000, minimum gene set size (set_min=15), maximum gene set size (set_max=200), excludes genes that have no gene symbols (collapse)= No_Collapse, value to use for the single identifier that will represent all identifiers for the gene (mode)=Max_probe, normalised enrichment score method (norm)= meandiv, weighted scoring scheme (scoring_scheme) = classic. Positive Normalised Enrichment Scores (NES) represent genes that were upregulated in the paediatric population compared to the adult population ($p < 0.01$ and $q < 0.1$). To visualise the output of universally enriched pathways across multiple cell types, the top 25 most frequently appearing positively and negatively associated terms were plotted. Additionally, for five cell types of interest [which had DEGs meeting the threshold of $p < 0.05$ and $abs(\log_2\text{FC}) > 0.1$], the top 5 positively associated terms were plotted.

snRNA-seq data plots

Plots were produced with Seurat[24], ggplot2[79], ShinyCell[80] and Microsoft Excel.

10x Genomics Visium library preparation

Frozen OCT embedded brain tissue samples were scored using a pre-chilled razor blade to fit in the Spatial Gene Expression slide capture areas. 10 µm-thick sections were cut using a cryostat (Leica CM1860/CM1950) and collected onto the Spatial Gene Expression slide capture areas. Two replicate sections of the 15-year-old (10 µm apart) and two replicate sections of 31-year-old (40 µm apart) were collected. The spatial Gene Expression slides with tissue sections were stored in a sealed container at -80°C. Captured sections were Haematoxylin and Eosin (H&E) stained according to the 10x Genomics Demonstrated Protocol Guide (CG000160, Rev B). Brightfield images of the stained sections were captured using an EVOS M5000 microscope (Thermo Fisher Scientific) at 20x magnification without coverslipping. Overlapping images of the sections including the fiducial frame were stitched together using Image Composite Editor-2.0.3 (Microsoft). Visium libraries were prepared

from the stained tissue sections following the Visium Spatial Gene Expression Reagents Kit
User Guide (CG000239, Rev D). At Step 1.1 the tissue was permeabilised for 12 minutes as
determined using the Visium Spatial Gene Expression Tissue Optimisation User Guide
(CG000238, Rev D). At Step 3.2, cDNA was amplified using 20 cycles. Library quality and
concentration was assessed using TapeStation (Agilent) and Qubit (Invitrogen) at the Central
Analytical Facility (CAF, University of Stellenbosch). Libraries were sequenced by Novogene
(Singapore) on the Illumina NovaSeq system using the Illumina High Output kits (150 cycles).
654

655 **Visium read alignment and gene expression quantification**
656 The H&E images were processed using the 10X Genomics Loupe Browser V4.0 Visium Manual
657 Alignment Wizard. 10X Genomics Space Ranger *count* (10X Space Ranger V1.3.0) was used to
658 perform alignment of FASTQ files to the human reference transcriptome (GRCh38), tissue
659 detection, fiducial detection and barcode/UMI counting.
660

cell2location analysis of Visium datasets
The average number of nuclei per Visium spot was determined using Vistoseg[81] (Code
availability: script 18). Cell2location (version 0.7a0)[25] was used to spatially map the brain cell
types by integrating the Visium data count matrices (Space Ranger output) with the
annotated snRNAseq datasets (Code availability: script 19). To avoid mapping artifacts,
mitochondrial genes were removed from the Visium datasets prior to spatial mapping.
Reference signatures of the 54 annotated cell populations were derived using a negative
binomial regression model using the default values. Unnormalized and untransformed
snRNA-seq mRNA counts were used as input in the regression model for estimating the
reference signatures (Code availability: script 20). The snRNA-seq mRNA counts were filtered
to 13,870 genes and 176,012 cells. The cell2location model for estimating the spatial
abundance of cell populations was filtered to 13,858 genes and 14,324 cells that were shared
in both the snRNA-seq and Visium data. The following cell2location parameters were used:
training iterations = 30,000 cell per location, N^ = 10 (estimated using Vistoseg segmentation
results), Normalization (ys) alpha prior = 20 (Code availability: script 21). To visualise the cell
abundance in spatial coordinates 5 % quantile of the posterior distribution was used, which
represents the value of cell abundance that the model has high confidence in (Code
availability: script 22). Cell2location's Non-negative Matrix Factorization (NMF) was used to
identify cellular compartments and cell types that co-locate from the cell type abundance
estimates, using n_fact=12 (Code availability: script 23)
681

BayesSpace analysis of Visium datasets
The raw gene expression counts from Space Ranger were normalized, log transformed and
principal component analysis was performed on the top 2000 highly variable genes. To obtain
high-resolution gene expression, the principal component values were mapped back to their
original log-transformed gene expression space (spot level) using the default BayeSpace[38]
regression (Code availability: script 24). To do this the principal components from the original

data were used as predictors in training the model for each gene, in which the results were the measured gene expression at the spot level. The trained model was then used to predict the gene expression at sub spot level using high resolution PCs. The high-resolution model was trained using default values except for the following parameters: 7 PCs, Number of clusters = 8, nrep = 100,000, burn-in = 10,000.

***In situ* Hybridisation Chain Reaction (HCR) on frozen human tissue sections**

10 μm thick frozen sections were collected on Histobond+ slides (Marienfeld) and stored at -20°C. The *In situ* HCR protocol was carried out on tissue sections as detailed in Choi et al. (2016)[82] using reagents, probes and hairpins purchased from Molecular Instruments. Probes were ordered for the following genes: *RELN* (NM_005045.4), *FABP7* (CR457057.1), *AQP4* (NM_001650.5), *RORB* (NM_006914.4), *CLSTN2* (NM_022131.3) and *TSHZ2* (NM_173485.6). When necessary to quench lipofuscin autofluorescence, sections were rinsed after HCR in 1x PBS and treated with 200 μl TrueBlack (Biotium) for 30 sec. Slides were rinsed in PBS, stained with Hoescht (Thermofisher) and mounted using SlowFade Gold Antifade Reagent (Invitrogen). Sections were imaged using the LSM 880 Airyscan confocal microscope (Carl Zeiss, ZEN SP 2 software) using the 40X or 60X objective.

References

Hodge, R. D. *et al.* Conserved cell types with divergent features in human versus mouse cortex. *Nature* **573**, 61-68 (2019).
Bakken, T. E. *et al.* Comparative cellular analysis of motor cortex in human, marmoset and mouse. *Nature* **598**, 111-119, doi:10.1038/s41586-021-03465-8 (2021).
Regev, A. *et al.* The Human Cell Atlas. *Elife* **6**, doi:10.7554/eLife.27041 (2017).
Network, B. I. C. C. A multimodal cell census and atlas of the mammalian primary motor cortex. *Nature* **598**, 86-102, doi:10.1038/s41586-021-03950-0 (2021).
Darmanis, S. *et al.* A survey of human brain transcriptome diversity at the single cell level. *Proc Natl Acad Sci U S A* **112**, 7285-7290, doi:10.1073/pnas.1507125112 (2015).
Li, M. *et al.* Integrative functional genomic analysis of human brain development and neuropsychiatric risks. *Science* **362**, doi:10.1126/science.aat7615 (2018).
Werling, D. M. *et al.* Whole-Genome and RNA Sequencing Reveal Variation and Transcriptomic Coordination in the Developing Human Prefrontal Cortex. *Cell Rep* **31**, 107489, doi:10.1016/j.celrep.2020.03.053 (2020).
Kang, H. J. *et al.* Spatio-temporal transcriptome of the human brain. *Nature* **478**, 483-489, doi:10.1038/nature10523 (2011).
Colantuoni, C. *et al.* Temporal dynamics and genetic control of transcription in the human prefrontal cortex. *Nature* **478**, 519-523, doi:10.1038/nature10524 (2011).
Donertas, H. M. *et al.* Gene expression reversal toward pre-adult levels in the aging human brain and age-related loss of cellular identity. *Sci Rep* **7**, 5894, doi:10.1038/s41598-017-05927-4 (2017).

11   Bourgeois, J. P. & Rakic, P. Changes of synaptic density in the primary visual cortex of
the macaque monkey from fetal to adult stage. *J. Neurosci.* **13**, 2801-2820,
doi:10.1523/JNEUROSCI.13-07-02801.1993 (1993).
12   Huttenlocher, P. R. & Dabholkar, A. S. Regional differences in synaptogenesis in human
cerebral cortex. *J. Comp. Neurol.* **387**, 167-178, doi:10.1002/(sici)1096-
9861(19971020)387:2<167::aid-cne1>3.0.co;2-z (1997).
13   Petanjek, Z. *et al.* Extraordinary neoteny of synaptic spines in the human prefrontal
cortex. *Proc Natl Acad Sci U S A* **108**, 13281-13286, doi:10.1073/pnas.1105108108
(2011).
14   Fu, M. & Zuo, Y. Experience-dependent structural plasticity in the cortex. *Trends*
Neurosci. **34**, 177-187, doi:10.1016/j.tins.2011.02.001 (2011).
15   Paus, T., Keshavan, M. & Giedd, J. N. Why do many psychiatric disorders emerge
during adolescence? *Nature Reviews Neuroscience* **9**, 947-957, doi:10.1038/nrn2513
(2008).
16   Jaffe, A. E. *et al.* Developmental and genetic regulation of the human cortex
transcriptome illuminate schizophrenia pathogenesis. *Nat. Neurosci.* **21**, 1117-1125,
doi:10.1038/s41593-018-0197-y (2018).
17   Taylor, D. M. *et al.* The pediatric cell atlas: defining the growth phase of human
development at single-cell resolution. *Dev. Cell* **49**, 10-29 (2019).
18   Tishkoff, S. A. *et al.* The genetic structure and history of Africans and African
Americans. *Science* **324**, 1035-1044, doi:10.1126/science.1172257 (2009).
19   PRB. *World Population Datasheet 2022*, <https://www.prb.org/wp-
content/uploads/2022/09/2022-World-Population-Data-Sheet-Booklet.pdf> (2022).
20   O'Malley, J., Wardlaw, T., You, D., Hug, L. & Anthony, D. Africa's child demographics
and the world's future. *Lancet* **384**, 730-732, doi:10.1016/S0140-6736(14)61331-3
(2014).
21   Schutte, C. M. Analysis of HIV-related mortality data in a tertiary South African
neurology unit, 2006-2012. *Southern African Journal of HIV Medicine* **14**, 121-124
(2013).
22   Rohlwink, U. K. *et al.* Clinical characteristics and neurodevelopmental outcomes of
children with tuberculous meningitis and hydrocephalus. *Dev Med Child Neurol* **58**,
461-468, doi:10.1111/dmcn.13054 (2016).
23   Thrupp, N. *et al.* Single-nucleus RNA-Seq is not suitable for detection of microglial
activation genes in humans. *Cell reports* **32**, 108189 (2020).
24   Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-
1902.e1821, doi:https://doi.org/10.1016/j.cell.2019.05.031 (2019).
25   Kleshchevnikov, V. *et al.* Cell2location maps fine-grained cell types in spatial
transcriptomics. *Nat. Biotechnol.* **40**, 661-671, doi:10.1038/s41587-021-01139-4
(2022).
26   Maynard, K. R. *et al.* Transcriptome-scale spatial gene expression in the human
dorsolateral prefrontal cortex. *Nat. Neurosci.* **24**, 425-436 (2021).
27   Hintsch, G. *et al.* The calsyntenins—a family of postsynaptic membrane proteins with
distinct neuronal expression patterns. *Mol. Cell. Neurosci.* **21**, 393-409 (2002).
28   Aevermann, B. *et al.* A machine learning method for the discovery of minimum marker
gene combinations for cell type identification from single-cell RNA sequencing.
Genome Res. **31**, 1767-1780 (2021).

29 Aevermann, B. D. *et al.* Cell type discovery using single-cell transcriptomics:
implications for ontological representation. *Hum. Mol. Genet.* **27**, R40-R47,
doi:10.1093/hmg/ddy100 (2018).

30 Tan, S. Z. K. *et al.* Brain Data Standards - A method for building data-driven cell-type
ontologies. *Sci Data* **10**, 50, doi:10.1038/s41597-022-01886-2 (2023).

31 Tissir, F. & Goffinet, A. M. Reelin and brain development. *Nat. Rev. Neurosci.* **4**, 496-
505, doi:10.1038/nrn1113 (2003).

32 Park, C. R. *et al.* Spatiotemporal expression and functional implication of CXCL14 in
the developing mice cerebellum. *Mol Cells* **34**, 289-293, doi:10.1007/s10059-012-
0116-0 (2012).

33 Tamamaki, N., Fujimori, K., Nojyo, Y., Kaneko, T. & Takauji, R. Evidence that Sema3A
and Sema3F regulate the migration of GABAergic neurons in the developing
neocortex. *J. Comp. Neurol.* **455**, 238-248 (2003).

34 Akhtar, M. W. *et al.* In vivo analysis of MEF2 transcription factors in synapse regulation
and neuronal survival. *PLoS One* **7**, e34863, doi:10.1371/journal.pone.0034863
(2012).

35 Ueda, S., Fujimoto, S., Hiramoto, K., Negishi, M. & Katoh, H. Dock4 regulates dendritic
development in hippocampal neurons. *J. Neurosci. Res.* **86**, 3052-3061,
doi:10.1002/jnr.21763 (2008).

36 Miller-Delaney, S. F., Lieberam, I., Murphy, P. & Mitchell, K. J. Plxdc2 is a mitogen for
neural progenitors. *PloS one* **6**, e14565 (2011).

37 Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome biology* **15**, 1-21 (2014).

38 Zhao, E. *et al.* Spatial transcriptomics at subspot resolution with BayesSpace. *Nat.*
Biotechnol. **39**, 1375-1384 (2021).

39 Kakimoto, T., Katoh, H. & Negishi, M. Regulation of neuronal morphology by Toca-1,
an F-BAR/EFC protein that induces plasma membrane invagination. *J. Biol. Chem.* **281**,
29042-29053 (2006).

40 Benyamin, B. *et al.* Childhood intelligence is heritable, highly polygenic and associated
with FNBP1L. *Mol. Psychiatry* **19**, 253-258 (2014).

41 Khatir, I. *et al.* Decoupling of mRNA and Protein Expression in Aging Brains Reveals the
Age-Dependent Adaptation of Specific Gene Subsets. *Cells* **12**,
doi:10.3390/cells12040615 (2023).

42 Patel, K. G., Liu, C., Cameron, P. L. & Cameron, R. S. Myr 8, a novel unconventional
myosin expressed during brain development associates with the protein phosphatase
catalytic subunits 1alpha and 1gamma1. *J. Neurosci.* **21**, 7954-7968,
doi:10.1523/JNEUROSCI.21-20-07954.2001 (2001).

43 Stepanov, V. *et al.* Analysis of Association of Genetic Markers in the LUZP2 and FBXO40
Genes with the Normal Variability in Cognitive Performance in the Elderly. *Int J*
Alzheimers Dis **2018**, 2686045, doi:10.1155/2018/2686045 (2018).

44 Feng, D. *et al.* A pan-cancer analysis of the oncogenic role of leucine zipper protein 2
in human cancer. *Exp Hematol Oncol* **11**, 55, doi:10.1186/s40164-022-00313-x (2022).

45 Perez-Santiago, J. *et al.* A combined analysis of microarray gene expression studies of
the human prefrontal cortex identifies genes implicated in schizophrenia. *J Psychiatr*
Res **46**, 1464-1474, doi:10.1016/j.jpsychires.2012.08.005 (2012).

46 Chen, J. *et al.* Identification of the Key Immune Cells and Genes for the Diagnostics
and Therapeutics of Meningioma. *World Neurosurg* **176**, e501-e514,
doi:10.1016/j.wneu.2023.05.090 (2023).
47 Maruoka, M. & Suzuki, J. Regulation of phospholipid dynamics in brain. *Neurosci Res*
167, 30-37, doi:10.1016/j.neures.2021.01.003 (2021).
48 Neniskyte, U. *et al.* Phospholipid scramblase Xkr8 is required for developmental axon
pruning via phosphatidylserine exposure. *The EMBO Journal*, e111790 (2023).
49 Zhang, X., Lin, P.-Y., Liakath-Ali, K. & Südhof, T. C. Teneurins assemble into presynaptic
nanoclusters that promote synapse formation via postsynaptic non-teneurin ligands.
Nature Communications **13**, 2297 (2022).
50 Cheung, A. *et al.* Teneurin paralogues are able to localise synaptic sites driven by the
intracellular domain and have the potential to form cis-heterodimers. *Front Neurosci*
16, 915149, doi:10.3389/fnins.2022.915149 (2022).
51 Beckmann, J., Schubert, R., Chiquet-Ehrismann, R. & Müller, D. J. Deciphering teneurin
domains that facilitate cellular recognition, cell–cell adhesion, and neurite outgrowth
using atomic force microscopy-based single-cell force spectroscopy. *Nano Lett.* **13**,
2937-2946 (2013).
52 Rogowski, K. *et al.* A family of protein-deglutamylating enzymes associated with
neurodegeneration. *Cell* **143**, 564-578, doi:10.1016/j.cell.2010.10.014 (2010).
53 Bodakuntla, S., Janke, C. & Magiera, M. M. Tubulin polyglutamylation, a regulator of
microtubule functions, can cause neurodegeneration. *Neurosci. Lett.* **746**, 135656,
doi:10.1016/j.neulet.2021.135656 (2021).
54 Yang, Q. Q. *et al.* Nuclear isoform of FGF13 regulates post-natal neurogenesis in the
hippocampus through an epigenomic mechanism. *Cell Rep* **35**, 109127,
doi:10.1016/j.celrep.2021.109127 (2021).
55 Wu, Q.-F. *et al.* Fibroblast growth factor 13 is a microtubule-stabilizing protein
regulating neuronal polarization and migration. *Cell* **149**, 1549-1564 (2012).
56 Bräuer, A. U. *et al.* A new phospholipid phosphatase, PRG-1, is involved in axon growth
and regenerative sprouting. *Nat. Neurosci.* **6**, 572-578 (2003).
57 Agbaegbu Iweka, C., Hussein, R. K., Yu, P., Katagiri, Y. & Geller, H. M. The lipid
phosphatase-like protein PLPPR1 associates with RhoGDI1 to modulate RhoA
activation in response to axon growth inhibitory molecules. *J. Neurochem.* **157**, 494-
507, doi:10.1111/jnc.15271 (2021).
58 Fagerberg, L. *et al.* Analysis of the human tissue-specific expression by genome-wide
integration of transcriptomics and antibody-based proteomics. *Mol. Cell. Proteomics*
13, 397-406, doi:10.1074/mcp.M113.035600 (2014).
59 Zheng, Z.-J. *et al.* Construction of the Six-lncRNA prognosis signature as a novel
biomarker in esophageal squamous cell carcinoma. *Frontiers in Genetics* **13**, 839589
(2022).
60 Xing, Z. *et al.* Clinical Significance and Immune Landscape of a Pyroptosis-Derived
LncRNA Signature for Glioblastoma. *Front Cell Dev Biol* **10**, 805291,
doi:10.3389/fcell.2022.805291 (2022).
61 Mochizuki, S. *et al.* Effect of ADAM28 on carcinoma cell metastasis by cleavage of von
Willebrand factor. *J. Natl. Cancer Inst.* **104**, 906-922, doi:10.1093/jnci/djs232 (2012).
62 Driessens, S. L. *et al.* Genes associated with cognitive ability and HAR show overlapping
expression patterns in human cortical neuron types. *Nature communications* **14**, 4188
(2023).

63 Gauvain, M. *Cognitive development in infancy and childhood*. (Cambridge University
Press, 2022).

64 Lee, J. J. *et al.* Gene discovery and polygenic prediction from a genome-wide
association study of educational attainment in 1.1 million individuals. *Nat. Genet.* **50**,
1112-1121, doi:10.1038/s41588-018-0147-3 (2018).

65 Savage, J. E. *et al.* Genome-wide association meta-analysis in 269,867 individuals
identifies new genetic and functional links to intelligence. *Nat. Genet.* **50**, 912-919,
doi:10.1038/s41588-018-0152-6 (2018).

66 Ximerakis, M. *et al.* Single-cell transcriptomic profiling of the aging mouse brain. *Nat.*
Neurosci. **22**, 1696-1708, doi:10.1038/s41593-019-0491-3 (2019).

67 Wu, J., Zhao, Z., Shi, Y. & He, M. Cortical VIP(+) Interneurons in the Upper and Deeper
Layers Are Transcriptionally Distinct. *J. Mol. Neurosci.* **72**, 1779-1795,
doi:10.1007/s12031-022-02040-8 (2022).

68 Tasic, B. *et al.* Shared and distinct transcriptomic cell types across neocortical areas.
Nature **563**, 72-78, doi:10.1038/s41586-018-0654-5 (2018).

69 Berg, J. *et al.* Human neocortical expansion involves glutamatergic neuron
diversification. *Nature* **598**, 151-158, doi:10.1038/s41586-021-03813-8 (2021).

70 Mohan, H. *et al.* Dendritic and Axonal Architecture of Individual Pyramidal Neurons
across Layers of Adult Human Neocortex. *Cereb. Cortex* **25**, 4839-4853,
doi:10.1093/cercor/bhv188 (2015).

71 Gidon, A. *et al.* Dendritic action potentials and computation in human layer 2/3
cortical neurons. *Science* **367**, 83-87, doi:10.1126/science.aax6239 (2020).

72 Eyal, G. *et al.* Human Cortical Pyramidal Neurons: From Spines to Spikes via Models.
Front Cell Neurosci **12**, 181, doi:10.3389/fncel.2018.00181 (2018).

73 Testa-Silva, G. *et al.* High bandwidth synaptic communication and frequency tracking
in human neocortex. *PLoS Biol.* **12**, e1002007, doi:10.1371/journal.pbio.1002007
(2014).

74 Beaulieu-Laroche, L. *et al.* Enhanced Dendritic Compartmentalization in Human
Cortical Neurons. *Cell* **175**, 643-651 e614, doi:10.1016/j.cell.2018.08.045 (2018).

75 Habib, N. *et al.* Massively parallel single-nucleus RNA-seq with DroNc-seq. *Nat.*
Methods **14**, 955-958, doi:10.1038/nmeth.4407 (2017).

76 McGinnis, C. S., Murrow, L. M. & Gartner, Z. J. DoubletFinder: Doublet Detection in
Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst* **8**, 329-
337 e324, doi:10.1016/j.cels.2019.03.003 (2019).

77 DePasquale, E. A. K. *et al.* DoubletDecon: Deconvoluting Doublets from Single-Cell
RNA-Sequencing Data. *Cell Rep* **29**, 1718-1727 e1718,
doi:10.1016/j.celrep.2019.09.082 (2019).

78 Raudvere, U. *et al.* g:Profiler: a web server for functional enrichment analysis and
conversions of gene lists (2019 update). *Nucleic Acids Res.* **47**, W191-W198,
doi:10.1093/nar/gkz369 (2019).

79 Wickham, H. *ggplot2: elegant graphics for data analysis*. (Springer, 2016).

80 Ouyang, J. F., Kamaraj, U. S., Cao, E. Y. & Rackham, O. J. L. ShinyCell: simple and
sharable visualization of single-cell gene expression data. *Bioinformatics* **37**, 3374-
3376, doi:10.1093/bioinformatics/btab209 (2021).

81 Tippani, M. *et al.* Vistoseg: a matlab pipeline to process, analyze and visualize high
resolution histology images for visium spatial transcriptomics data. *bioRxiv* (2021).

82   Choi, H. M. *et al.* Mapping a multiplexed zoo of mRNA expression. *Development* **143**,
3632-3637, doi:10.1242/dev.140137 (2016).

916
917

Data Availability

All scripts used to analyse the data are indicated in the methods section and are available in the supplementary material. A description of the raw and analysed data files will be made available on the University of Cape Town's ZivaHub data sharing platform on publication. As the data is from living donors, access to the data will be mediated through contact with the corresponding author. A ShinyApp will be made publicly available on publication for exploration of the annotated snRNA-seq data.

Acknowledgments

This research was support by a Royal Society/Global Challenges Research Fund/African Academy of Sciences Future Leaders African Independent Researcher Fellowship to DH (FLR\R1\191008), a National Research Foundation (NRF) Research Development Grant for Y-rated Researchers Award to DH (CSRP210415595025), a University of Cape Town (UCT) Building Research Active Academic Staff Grant award to DH and a National Institutes of Health (NIH) R21 Exploratory/Development Grant to DH, JVR, CGD and MJ (TW011225). CS was supported by a Harry Crossley Research Scholarship, an Oppenheimer Memorial Trust scholarship, an NRF scholarship and a UCT Vice Chancellor's Research Scholarship. SF was supported by a DAAD-NRF joint In-country Scholarship and a UCT Vice Chancellor's Research Scholarship. MBV was supported by an EMBO long-term fellowship (ALTF 415-2018) and a Claude Leon Foundation research fellowship. AF was supported by the NRF SARChI Chair of Clinical Neurosciences. Computations were performed using facilities provided by the University of Cape Town's ICTS High Performance Computing team: hpc.uct.ac.za. This publication is part of the Human Cell Atlas: www.humancellatlas.org/publications/

Fig. 1: Annotation of nuclei by label transfer identifies 54 cortical subtypes across the 23 datasets. a, Data
integration shows alignment of nuclei across the technical (T) and biological (B) replicates from donors ranging
in age from 4 to 50 years. **b,** Annotated UMAP plot for the 23 merged datasets identifies 34 inhibitory, 14
excitatory, and 6 non-neuronal populations using the Allen Brain Map MTG dataset as a reference. Each cell type
is annotated with 1) a major cell class (e.g. Exc for excitatory neuron), 2) the cortical layer the cell is associated
with (e.g. L2 for layer 2), 3) a subclass marker gene and 4) a cluster-specific marker gene. **c,** Validation of the
high-resolution cell type annotations shows a high degree of correspondence in the expression of known cell
type-specific markers (x axis) with their expected cell type (y axis) (left). Number of nuclei per cell type (right). **d,**
Correlation plot showing the cosine similarity scores assessing similarity between the annotated cell types in our
dataset (y axis) and the MTG reference dataset (x axis) based on the log normalized expression counts of the top
2000 shared highly variable features between query and reference datasets. **e,** Stacked barplot showing the
proportion of nuclei per cell type (y axis) for each age category (x axis) out of the total number of nuclei for each
group. The colour scheme for the cell types is in accordance with the MTG cell taxonomy.

Fig. 2: Fig. 2: Visium spatial transcriptomics in the adult and paediatric temporal cortex validates snRNA-seq annotation. a,b, Estimated cell type abundances (colour intensity) in the 31-year-old and 15-year-old temporal cortex tissue sections for a selection of cell types including non-neuronal cell types (a,b), excitatory neurons (a) and inhibitory neurons (b). c, Visium gene expression profiles (colour intensity) for a selection of known cortical layer marker genes in the 31-year-old and 15-year-old temporal cortex tissue sections including *AQP4* (layer 1), *LAMP5* (layer 2), *CUX2* (layer 2-3), RORB (layer 4) and *CLSTN2* (layer 5-6). Dashed white lines and numbers indicate estimated cortical layer boundaries. d,e, Identification of co-locating cell types using NMF. The dot plot (d) shows the NMF weights of the cell types (rows) across each of the NMF factors (columns), which correspond to tissue compartments. Block boxes indicate cell types that co-locate within the indicated tissue compartments. Spatial plots show (e) show the NMF weights for each NMF factor/tissue compartment across the 31-year-old and 15-year-old temporal cortex tissue sections. Panels are displayed in the same order as the dotplot in (d), with the dominant cell types for each factor indicated in brackets. WM: white matter. See also Extended Data Figs 4-6.

991

992 **Fig. 3: NS-Forest identifies minimal marker genes distinguishing the cell types in the paediatric and adult**
993 **temporal cortex snRNA-seq datasets. a,b,** Heatmap showing the scaled average normalised expression counts
994 of the NS-Forest minimal marker genes (y-axis) identified for 53 cortical cell types (x-axis) across the six adult (**a**)
995 and six paediatric (**b**) datasets. As input into NS-Forest, the nuclei of each sample were randomly down-sampled
996 to the size of the sample with the fewest nuclei. Heatmaps show gene expression values for the down-sampled
997 datasets. Inh_L2–4_VIP_SPAG17 was excluded from the down-sampled datasets due to a low number of nuclei.
998 The minimal marker genes are annotated (colour codes on the y-axes) according to whether they are unique to
999 a given cell type, whether they are coding/non-coding genes, whether they are unique to the indicated age
1000 group, whether they overlap with existing MTG minimal marker gene sets for the same cell type, and according
1001 to the cell type they define.

1002

1003

1004

1005

1006

1007

1008

1009

Fig. 4: Validation of NS-Forest minimal markers and assessment of the top 50 NS-forest markers. a,b, Annotated UMAP plots following data integration using either the minimal marker genes (left) or the equivalent number of a random set of genes (right) as anchors for the adult (**a**) and paediatric (**b**) datasets. The colour scheme for the cell types is in accordance with the MTG cell taxonomy. **c,** Overlap of the top 50 paediatric and top 50 adult NS-Forest markers per cell type. The bar plot shows the number of shared markers between paediatric and adult datasets (blue), the number of markers unique to the paediatric datasets (orange), and the number of markers unique to the adult datasets (grey) for each cell type.

1022

Fig. 5: Differential expression analysis reveals genes guiding temporal cortex maturation. a, The number of differentially expressed genes (DEGs) identified per cell type. **b**, Strip chart showing the log$_2$FoldChange for all DESeq2-tested genes (dots) across 54 cell types (x axis). Genes that were significantly upregulated (110) or downregulated (52) in paediatric samples [padj<0.05 and abs(log$_2$FoldChange)>10%)] are coloured according to the MTG cell taxonomy. Non-significant genes [padj>0.05 or abs(log$_2$FoldChange)<10%] are coloured in grey. Cell types with significant DEGS are coloured red. **c**, Volcano plots showing log$_2$FoldChange (x axis) and -log$_{10}$padj values (y axis) for all DESeq2-tested genes in Exc_L2_LAMP5_LTK, Exc L2-3_LINC00507_FREM3, Exc_L4-5_RORB_DAPK2, Exc_L3-5_RORB_ESR1 and Astro_L1-6_FGFR3_SLC14A1. Genes that were significantly upregulated or downregulated in paediatric samples (padj<0.05 & abs(log$_2$FoldChange)>10%) are coloured in red and selected genes are labelled. Non-significant genes (padj>0.05 or abs(log$_2$FoldChange)<10%) are coloured in blue. Red labels indicate DEGs shared between similar cell types. **d,e**, Dot plots showing the scaled average normalised expression across samples for DEGS shared between either Exc_L2_LAMP5_LTK and Exc L2-3_LINC00507_FREM3 (**d**) or Exc_L4-5_RORB_DAPK2 and Exc_L3-5_RORB_ESR1 (**e**). **f**, Dot plot showing the scaled average normalised expression across samples of DEGs in Astro_L1-6_FGFR3_SLC14A1. **g**, Enrichment plot showing the top 5 GSEA terms (y axis) that were enriched in the paediatric datasets and their associated NES values (x axis) for the five indicated cell types.

1039

1040 **Fig. 6: Pathways that are enriched or depleted across multiple cell types in paediatric cell types.** GSEA heatmap
1041 showing the top 25 most frequently enriched (top 25 rows) or depleted (bottom 25 rows) terms appearing across
1042 all cell types. Only significantly (p < 0.01 and q < 0.1) terms are shown. NES value represents the normalized
1043 enrichment scores. Grey indicates that the term was not significantly enriched or depleted in the indicted cell
1044 type. See also Extended Data Table 11.

1045

1046

1047

1048

1049

1050

1051

Extended Data Fig. 1: Nuclei filtering and clustering. a, Number of doublets identified across all 23 datasets by DoubletDecon, DoubletFinder, and Scrublet. Red outline indicates the subset of barcodes called as doublets that were removed. **b**, Total number of nuclei per dataset before (yellow) and after filtering (green). **c**, Mean number of reads per nucleus (y axis) by dataset prior to filtering split by age group (x axis). p value determined by two-tailed Welch's t-test. **d**, Number of nuclei (y axis) by sample after filtering split by age group (x axis). p value determined by Brunnermunzel permutation test. **e**, Violin plots showing the number of unique molecular identifiers (UMIs) (top) and the number of genes detected (bottom) per nucleus per sample after filtering. Black dots indicate the median value. Error bars show 95% confidence intervals. **f,g**, Median number of UMIs (2,263 paediatric and 2,011 adult) (**f**) and the median number of genes (1,372 paediatric and 1,226 adult) (**g**) detected per nucleus (y axes) by sample after filtering split by age group (x axis). p values determined by two-tailed Brunnermunzel permutation test. **h**, UMAP plot for the 23 datasets prior to integration. **i**, UMAP plot showing the resulting clusters determined by the shared nearest neighbour algorithm. Data in all box plots represent mean ± sem for six paediatric and six adult samples. No significant differences were detected between paediatric and adult samples. B, biological replicate; NS, not significant; T, technical replicate. See also Extended Data Table 2.

1067

Extended Data Fig. 2: Assessment of cell composition across datasets. a, b, Stacked barplots showing the
proportion of nuclei per cell type (y axis) for each technical replicate (**a**) or biological replicate (**b**) (x axis) out of
the total number of nuclei for each group. The colour scheme for the cell types is in accordance with the MTG
cell taxonomy. Samples with technical replicates showed high degrees of similarity in cell composition between
their replicates (**a**). Technical replicates from each donor were merged to allow comparisons between the 12
samples (**b**). **c,** Boxplots showing the proportion of nuclei (y axis) per cell type per sample (x axis) split by age
group (red: paediatric, grey: adult). Data represents mean ± sem for six paediatric and six adult samples. No
significant differences were detected between paediatric and adult samples; See also Extended Data Table 3 for
details of statistical tests performed.

 Extended Data Fig. 3: Assessment of the sequencing metrics for the annotated cell types. a, Violin plots showing
the distribution of the number of genes (left) and transcripts (right) detected per nucleus per cell type across all
datasets. Black dots indicate the median value. Error bars show 95% confidence intervals. **b,c**, Boxplots showing
the number of genes (**b**) and the number of UMIs (**c**) (y axis) detected per cell type per sample (x axis) split by
age group (red: paediatric, grey: adult). Data in all box plots represent mean ± sem for six paediatric and six adult
samples for each cell type. p values are given for cell types that showed a significant difference between
paediatric and adult samples. See also Extended Data Table 3 for details of statistical tests performed.

1085

1086

Extended Data Fig. 4: Visium Spatial Gene Expression samples. a,b, 31-year-old (**a**) and 15-year-old (**b**) temporal cortex tissue blocks embedded in OCT. Black dashed boxes outline the regions collected onto the Visium Spatial Gene Expression slide. **c-f**, H&E stained technical replicate tissue sections used to generate Visium Spatial Gene Expression libraries for the 31-year-old (**c,e**) and 15-year-old (**d,f**) tissue samples. T, technical replicate.

1098

Extended Data Fig. 5: Spatial mapping of cell types in the human temporal cortex. a, Estimated cell abundance of 52 cell types across all Visium samples. Shown is a heatmap with the colour indicating the relative cell abundance of cell types (rows) across the different samples (columns). **b,** Estimated cell type abundances (colour intensity) in the technical replicate 31-year-old and 15-year-old temporal cortex tissue sections for a selection of cell types including non-neuronal cell types (**b,c**), excitatory neurons (**b**) and inhibitory neurons (**c**). **d,** Estimated cell type abundances (colour intensity) for Exc_L4-6 _FEZF2_IL26 and Exc_L5-6_THEMIS_FGF10 in the 31-year-old and 15-year-old temporal cortex tissue sections. T, technical replicate.

1106

1107

1108

1109

1110

1111

1112

1113

1114

1115

Extended Data Fig. 6: *In situ* **HCR analysis of selected cortical layer marker genes. a,** Expression of layer 1 markers *AQP4*, *FABP7* and *RELN* in 31-year-old and 15-year-old temporal cortex tissue sections, with high magnification views of layer 1 indicating *AQP4/RELN*-positive cells (yellow arrowheads) and *FABP7* positive cells (green arrowhead). **b,** Expression of layer 4-6 markers *RORB*, *CLSTN2* and *TSHZ2* in 31-year-old and 15-year-old temporal cortex tissue sections. In high magnification views of layer 4 in the 31-year-old tissue section *RORB/CLSTN2*-positive (white arrowhead) and *RORB/TSHZ2*-positive cells (green arrowhead) are indicated. In high magnification views of layer 4 in the 15-year-old tissue section *RORB/CLSTN2/TSHZ2*-positive cells (white arrowheads) are indicated. Dashed white lines indicate layer boundaries. Solid white line indicates tissue edge. Scale bars are 100 µm in low magnification views (tile scan at 40x) and 20 µm in high magnification views (63x).

1125

1126

1127

Extended Data Fig. 7: Expression of the reference MTG atlas minimal markers. Heatmap showing the scaled average normalised expression counts of the NS-Forest minimal marker genes identified for the reference MTG cell atlas dataset (y-axis) in each of the 54 query cortical cell types identified in the combined adult and paediatric snRNA-seq datasets (x-axis). The minimal marker genes are annotated (colour codes on the y-axes) according to the cell type they define.

1133

1134

1135

1136

1137

Extended Data Fig. 8: Evaluation of NS-Forest minimal marker gene expression across cell types in comparison to MTG cell taxonomy markers. a-d, Boxplots showing the normalised expression counts for *DDR2* (**a**)*, TGFBR2* (**b**)*, SEMA3E* (**c**) and *SULF1* (**d**) in paediatric (top) and adult (bottom) datasets. The cell types expressing the markers at high levels are indicated in bold.

Extended Data Fig. 9: BayesSpace analysis of differentially expressed genes. High resolution Visium spatial gene expression profiles for selected DEGs using BayesSpace analysis to compare sub-spot level expression intensities between 31-year-old and 15-year-old temporal cortex tissue sections.

1142

1143

1144

1145

1146

1147

1148

1149

1150

1151

1152

1153

1154

1155

1156

Extended data Tables:

Extended Data Table 1: Summary of snRNA-seq and Visium sample metadata. Samples are ordered by age. The eight "P00" datasets were generated in the Hockman laboratory while the four "Nuc" datasets were generated by Thrupp et al. (2020).

Extended Data Table 2: Summary of average quality control metrics for snRNA-seq datasets across nuclei for each sample before and after filtering. Several measures for quality control were evaluated on a per sample basis including the sequencing saturation, the mean number of reads per nucleus, the number of barcodes, the median number of genes detected per nucleus, the median number of UMIs detected per nucleus, and the number of doublets removed.

Extended Data Table 3: Label transfer annotation of snRNA-seq datasets using the Allen Brain Map MTG atlas as a reference. Sheet 1, Number of nuclei per MTG cell type per sample. The number of barcodes corresponding to each MTG cell type and sample is shown. Additionally, the total, minimum, and maximum number of nuclei per cell type and sample was computed. The number of cell types represented per sample was also determined. **Sheet 2,** Cell types present in reference MTG dataset which are absent from the query datasets. Label transfer resulted in 21 cell populations that were not annotated in the query datasets but were present in the reference MTG dataset. The cluster size and percentage of the total cell count for each cell type in the reference MTG dataset is also shown. **Sheet 3-5,** p values and the test performed for each cell type when comparing the proportion of nuclei (sheet 3; see Extended Data Fig. 2c), number of genes (sheet 4; see Extended Data Fig. 3b) and number of UMIs (sheet 5; see Extended Data Fig. 3c) for each cell type between paediatric and adult samples shown.

Extended Data Table 4: Summary of average quality control metrics for Visium datasets. Several measures for quality control were evaluated on a per sample basis including the sequencing saturation, the percentage of read mapped to the transcriptome, the number of spots under the tissue, the average number of nuclei per spot determined by Vistoseg analysis, the mean reads detected per spot, the median genes detected per spot, the total number of genes detected, the median UMI Counts per Spot and the total number of nuclei.

Extended Data Table 5: NS-Forest minimal marker analysis. Sheet 1, Statistics from NS-Forest analysis including the mean F-beta score (the measure of the discriminative power of a given combination of marker genes) and the mean binary expression score per cell type (a measure of an individual gene's classification power). **Sheets 2-3,** Metadata for each feature identified by NS-Forest marker in the down-sampled paediatric (**sheet 2**) and down-sampled adult (**sheet 3**) datasets describing the cell type, overlap with Aevermann et al. (2021) and Hodge et al. (2019), uniqueness to the age group of interest, coding status, and uniqueness to the associated cell type as shown in Fig. 3. As input to NS-Forest, all datasets (six paediatric and six adult) were randomly down-sampled such that the total number of nuclei per sample was equal to the sample with the fewest number of nuclei.

Extended Data Table 6: gProfiler analysis of NS-forest markers. (**Sheet1-2**) Significantly enriched GO terms associated with the paediatric (**sheet1**) and adult (**sheet2**) minimal marker genes identified by NS-forest. (**Sheet3-7**) Significantly enriched GO terms associated with shared (i.e associated with both adult and paediatric sample) or paediatric-specific marker genes from the list of top 50 positive NS-Forest markers for selected cell types. Terms for which "highlighted" is true are driver terms.

Extended Data Table 7: Summary of metadata for the top 50 positive NS-Forest markers per cell type. The top 50 positive NS-Forest markers (or total number of positive markers if < 50) per cell type for the down-sampled paediatric and down-sampled adult datasets. The number of intersecting markers, the number of markers unique to paediatric samples, and the number of markers unique to adults is shown for each cell type. The number of nuclei per cell type is shown for the combined paediatric and adult down-sampled datasets, the down-sampled paediatric datasets, and down-sampled adult datasets.

Extended Data Table 8: Overlap of the top 50 positive NS-Forest markers per cell type between the paediatric
and adult datasets. The top 50 positive NS-Forest markers (or total number of positive markers if < 50) per cell
type were extracted for the down-sampled paediatric and down-sampled adult datasets. Each sheet represents
1 of 53 cortical cell types (Inh_L2-4_VIP_SPAG17 was excluded due to too few nuclei) and the NS-Forest features
which were shared between the paediatric and adult datasets, unique to paediatric datasets, or unique to adult
datasets are shown (sheet 1-53).

1215

Extended Data Table 9: DESeq2 output of all genes tested for differential expression between paediatric and
adult brains per cell type. (**Sheet 1-53**) Differential expression analysis was performed using DESeq2's Wald Test
for each cell type separately (Inh L2-4 VIP SPAG17 was excluded due to too few nuclei). Genes were filtered prior
to testing to only include those expressed in > 10% of nuclei for that cell type across all paediatric and adult
datasets. The associated log2FoldChanges, p-adjusted values (padj), and description of each feature are shown.
Positive $\log_2$FoldChanges represent genes upregulated in paediatrics versus adults. See DESeq2 documentation
for explanation of NA values
(https://bioconductor.org/packages/release/bioc/vignettes/DESeq2/inst/doc/DESeq2.html#why-are-some-p-
values-set-to-na).

1225

Extended Data Table 10: DESeq2 output of significant DEGs between paediatric and adult brains in a subset of
cell types. (**sheet 1-5**) Significant DEGs (padj < 0.05) for Exc_L2_LAMP5_LTK, Exc_L2-3_LINC00507 _FREM3,
Exc_L3-5_RORB_ESR1, Exc_L4-5_RORB_DAPK2 and Astro_L1-6_FGFR3_SLC14A1. The associated
$\log_2$FoldChanges, p-adjusted values (padj), description, percentage of paediatric nuclei expressing the gene,
percentage of adult nuclei expressing the gene, average normalised expression across paediatric nuclei, and
average normalised expression across adult nuclei are shown. Positive $\log_2$FoldChanges represent genes
upregulated in paediatric versus adults datasets. See DESeq2 documentation for explanation of NA values
(https://bioconductor.org/packages/release/bioc/vignettes/DESeq2/inst/doc/DESeq2.html#why-are-some-p-
values-set-to-na). (**sheet 6-7**) EA (sheet 6) and IQ (sheet7) associated DEGs and their associated cell types.

1235

Extended Data Table 11: **GSEA terms associated with each cell type showing enriched or depleted pathways in**
paediatric versus adult samples. GSEA was performed using DESeq2's output gene lists for each cell type ranked
according to the log2FoldChange*-log$_2$(padj) for each gene. All DESeq2-tested genes served as input into GSEA
(genes were expressed in > 10% of nuclei for the cell type of interest). Matrix shows the corresponding positive
(**sheet 1**) and negative (**sheet 2**) NES values for each GSEA term (y axis) and cell type (x axis) based on the analysis
using the ranked list of genes for each cell type. Terms were filtered to only include significantly associated terms
(p<0.01, q<0.1). Positive NES values indicate pathways that are enriched in paediatric versus adult samples;
negative NES values indicate pathways that are depleted in paediatric versus adult samples. The total number of
terms per cell type and the total number of cell types associated with a given term are shown. 1,290 terms were
enriched in at least two cell types, while 250 were depleted in at least two cell types.

1246

1247

1248

1249

1250

1251

1252

1253

1254

1255

1256