RAG1 and RAG2 Non-core Regions Are Implicated in Leukemogenesis and Off-target V(D)J Recombination in BCR-ABL1-driven B-cell Lineage Lym-phoblastic Leuke-mia

Xiaozhuo Yu[1*], Wen Zhou[1*], Xiaodong Chen[1], Shunyu He[1], Mengting Qin[1], Meng Yuan[1], Yang Wang[1], Woodvine otieno Odhiambo[1], Yinsha Miao[2 **] and Yanhong Ji[1,2 **]

[1]Department of Pathogenic Biology and Immunology. School of Basic Medical Sciences, Xi'an Jiaotong University Health Science Center.

[2] Department of Clinical laboratory, Xi'an No. 3 Hospital, the Affiliated Hospital of Northwest University.

*Xiaozhuo Yu and Wen Zhou contributed equally to this work as co-first anthor

**Correspondences:

Yanhong Ji (jiyanhong@xjtu.edu.cn), Yinsha Miao (miaoyinsha@med.nwu.edu.cn),

Running title: RAG's Non-core Regions Suppress Leukemogenesis and Off-target V(D)J Recombination

Key words: RAG, Non-core regions, Off-target V(D)J recombination, *BCR-ABL1*[+] B-ALL, Genomic stability

Abstract

The evolutionary conservation of non-core RAG regions suggests significant roles that might involve quantitative or qualitative alterations in RAG activity. Off-target V(D)J recombination contributes to lymphomagenesis and is exacerbated by RAG2' C-terminus absence in Tp53$^{-/-}$ mice thymic lymphomas. However, the genomic stability effects of non-core regions from both cRAG1 and cRAG2 in *BCR-ABL1*$^+$ B-lymphoblastic leukemia (*BCR-ABL1*$^+$ B-ALL), the characteristics, and mechanisms of non-core regions in suppressing off-target V(D)J recombination remains unclear. Here, we established three mouse models of *BCR-ABL1*$^+$ B-ALL in mice expressing full-length RAG (fRAG), core RAG1 (cRAG1), and core RAG2 (cRAG2). The cRAG (cRAG1 and cRAG2) leukemia cells exhibited greater malignant tumor characteristics compared to fRAG cells. Additionally, cRAG cells showed higher frequency of off-target V(D)J recombination and oncogenic mutations than fRAG. We also revealed decreased RAG cleavage accuracy in cRAG cells and a smaller recombinant size in cRAG1 cells, which could potentially exacerbate off-target V(D)J recombination in cRAG cells. In conclusion, these findings indicate that the non-core RAG regions, particularly the non-core region of RAG1, play a significant role in preserving V(D)J recombination precision and genomic stability in *BCR-ABL1*$^+$ B-ALL.

Introduction

V(D)J recombination serves as the central process for early lymphocyte development and generates diversity in antigen receptors. This process involves the double-strand DNA breaks of gene segments by the V(D)J recombinase, including RAG1 and RAG2. RAG recognizes conserved recombination signal sequences (RSSs) positioned adjacent to V, D, and J gene segments. A bona fide RSS contains a conserved palindromic heptamer (consensus 5′-CACAGTG) and A-rich nonamer (consensus 5′-ACAAAAACC) separated by a degenerate spacer of either 12 or 23 base pairs (*Hirokawa, et al.*,*2020*; *Schatz and Ji*,*2011*). The process of efficient recombination is contingent upon the presence of recombination signal sequences (RSSs) with differing spacer lengths, as dictated by the "12/23 rule" (*Banerjee and Schatz*,*2014*; *Eastman, et al.*,*1996*; *Shi, et al.*,*2020*). Following cleavage, the DNA ends are joined via non-homologous end joining (NHEJ), resulting in the precise alignment of the two coding ends and the signal ends (*Rooney, et al.*,*2004*). V(D)J recombination promotes B cell development, but aberrant V(D)J recombination can lead to precursor B-cell malignancies through RAG mediated off-target effects (*Mendes, et al.*,*2014*; *Onozawa and Aplan*,*2012*; *Thomson, et al.*,*2020*).

The regulation of RAG expression and activity is multifactorial, serving to ensure V(D)J recombination and B cell development (*Gan, et al.*,*2021*; *Kumari, et al.*,*2021*). The RAGs consist of core and non-core region. Although non-core regions of RAG1/2 are not strictly required for V(D)J recombination, the evolutionarily conserved non-core RAG regions indicate their potential significance in vivo that may involve quantitative or qualitative modifications in RAG activity and expression (*Braams, et al.*,*2023*; *Curry and Schlissel*,*2008*; *Liu, et al.*,*2022*; *Liu, et al.*,*2022*; *Sekiguchi, et al.*,*2001*). Specifically, the non-core RAG2 region (amino acids 384–527 of 527 residues) contains a plant homeodomain (PHD) that can recognize histone H3K4 trimethylation, as well as a T490 locus that mediates a cell cycle-regulated protein degradation signal in proliferated pre-B cells stage (*Liu, et al.*,*2007*;

Matthews, et al.,2007). Failure to degrade RAG2 during the S stage poses a threat

to the genome (*Zhang, et al.,2011*). Moreover, the off-target V(D)J recombination

frequency is significantly higher when RAG2 is C-terminally truncated, thereby estab-

lishing a mechanistic connection between the PHD domain, H3K4me3-modified

chromatin, and the suppression of off-target V(D)J recombination (*Lu, et al.,2015*;

Mijušković, et al.,2015). The RAG1' non-core region (amino acids1–383 of 1040

residues) has been identified as a RAG1 regulator. While the core RAG1 maintains

its catalytic activity, it's in vivo recombination efficiency and fidelity are reduced in

comparison to the full-length RAG1 (fRAG1). In addition, the RAG1 binding to the

genome is more indiscriminate (*Beilinson, et al.,2021*; *Sadofsky, et al.,1993*). The

N-terminal domain (NTD), which is evolutionarily conserved, is predicted to contain

multiple zinc-binding motifs, including a Really Interesting New Gene (RING) domain

(aa 287 to 351) that can ubiquitylate various targets, including RAG1 itself (*Deng, et*

al.,2015).. Furthermore, NTD contains a specific region (amino acids 1 to 215) that

facilitates interaction with DCAF1, leading to the degradation of RAG1 in a CRL4-

dependent manner (*Schabla, et al.,2018*). Additionally, the NTD plays a role in

chromatin binding and the genomic targeting of the RAG complex (*Schatz and*

Swanson,2011). Despite increased evidence emphasizing the significance of non-

core RAG regions, particularly RAG1's non-core region, the function of non-core

RAG regions in off-target V(D)J recombination and the underlying mechanistic basis

have not been fully clarified.

Typically, genomic DNA is safeguarded against inappropriate RAG cleavage by

the inaccessibility of cryptic RSSs (cRSSs), which are estimated to occur once per

600 base pairs (*Teng, et al.,2015*). However, recent research has demonstrated that

epigenetic reprogramming in cancer can result in heritable alterations in gene ex-

pression, including the accessibility of cRSSs (*Becker, et al.,2020*; *Fatma, et*

al.,2022; *Goel, et al.,2022*; *Khoshchehreh, et al.,2019*). We selected the *BCR-*

ABL1[+] B-ALL model, which is characterized by ongoing V(D)J recombinase activity

and *BCR-ABL1* gene rearrangement in pre-B leukemic cells (*Schjerven, et al.,2017*;

Wong and Witte,*2004*). The genome structural variations (SVs) analysis was con-

ducted on leukemic cells from *fRAG, cRAG1*, and *cRAG2, BCR-ABL1*[+] B-ALL mice to

examine the involvement of non-core RAG regions in off-target V(D)J recombination

events. The non-core domain deletion in both *RAG1* and *RAG2* led to accelerated

leukemia onset and progression, as well as an increased off-target V(D)J recombina-

tion. Our analysis showed a reduction in RAG cleavage accuracy in *cRAG* cells and

a decrease in recombinant size in *cRAG1* cells, which may be responsible for the in-

creased off-target V(D)J recombination in *cRAG* leukemia cells. In conclusion, our

results highlight the potential importance of the non-core RAG region, particularly

RAG1's non-core region, in maintaining accuracy of V(D)J recombination and ge-

nomic stability in *BCR-ABL1*[+] B-ALL.

Results

cRAG give more aggressive leukemia in a mouse model of *BCR-ABL1*[+] B-ALL

In order to assess the impact of RAG activity on the clonal evolution of *BCR-ABL1*[+]

B-ALL through a genetic experiment, we utilized bone marrow transplantation (BMT)

to compare disease progression in fRAG, cRAG1, and cRAG2 *BCR-ABL1*[+] B-ALL

(*Schjerven, et al.*,*2017*; *Wong and Witte*,*2004*). Bone marrow cells transduced with

a BCR-ABL1/GFP retrovirus were administered into syngeneic lethally irradiated

mice, and CD19[+] B cell leukemia developed within 30-80 days (Figure 1A, Supple-

mentary Figure 1). Western blotting results confirmed equivalent transduction effi-

ciencies of the retroviral *BCR-ABL1* in all three cohorts (Supplementary Figure 2A).

To investigate potential variances in leukemia outcome across different genomic

backgrounds, we employed Mantel-Cox analysis to evaluate survival rates in fRAG,

cRAG1, or cRAG2 mice transplanted with *BCR-ABL1*-transformed bone marrow cells.

Our results show that, during the primary transplant phase, *BCR-ABL1*[+] B-ALL mice

expressing cRAG1 or cRAG2 demonstrated lower survival rates compared to their

counterparts with fRAG (median 74.5 days versus 39 or 57 days, $P < 0.0425$, Figure

1A). This survival rates discrepancy was also observed during the secondary trans-

plant phase, wherein leukemic cells were extracted from the spleens of primary recip-

ients and subsequently purified via GFP$^+$ cell sorting. A total of 10^5,10^4 and 10^3 GFP$^+$

leukemic cells that originated from fRAG, cRAG1, or cRAG2 leukemic mice were

transplanted into corresponding non-irradiated immunocompetent syngenetic recipi-

ents (survival days fRAG,11-26 days, cRAG1,10-16 days, cRAG2,11-21 days, Sup-

plementary Figure 2B). Additionally, the cRAG mice exhibited significantly higher leu-

kemia burdens in the peripheral blood, bone marrow, and spleen compared to the

fRAG mice (Figure 1B-D). To elucidate the cellular mechanisms driving the acceler-

ated proliferation observed in cRAG *BCR-ABL1*$^+$ B-ALL, flow cytometry analyses

were conducted to evaluate cell cycle dynamics and apoptotic activity. Results re-

vealed a higher fraction of cRAG *BCR-ABL1*$^+$ B-ALL cells residing in the S/G2-M

phase of the cell cycle compared to their fRAG counterparts (Figure 1E). Additionally,

the enhanced proliferation in cRAG leukemic cells was attributed to a reduction in

apoptosis rates (Supplementary Figure 2C). RNA-seq analysis demonstrated the

changes of cell differentiation and proliferation/apoptotic pathways (Supplementary

Figure 3) These findings indicate that the absence of non-core RAG regions acceler-

ates malignant transformation and leukemic proliferation, leading to a more aggres-

sive disease phenotype in the cRAG *BCR-ABL1*$^+$ B-ALL mouse model.

147   The loss of non-core RAG regions corresponds to a less ma-
ture cell surface phenotype but does not impede IgH VDJ re-
combination

To delineate the developmental stages of B cells from which the leukemic cells origi-

nated, we performed flow cytometry on single cells stained with B cell-specific sur-

face markers. Analysis revealed that 91%-98% of GFP$^+$ cells in cRAG mice were

CD19$^+$BP-1$^+$B220$^+$CD43$^+$, indicating that most leukemic cells were at the large pre-B

cell stage (Figure 2A) (**Hardy and Hayakawa**,*2001*). Conversely, in fRAG leukemic

mice, the distribution was 65% large pre-B cells (GFP$^+$CD19$^+$BP-1$^+$B220$^+$CD43$^+$) and

35% small pre-B cells (GFP$^+$CD19$^+$BP-1$^+$B220$^+$CD43$^-$) (Figure 2A). Moreover, ap-

proximately 5% of leukemic cells in fRAG mice expressed μHC, in con-trast to mini-

mal expression in cRAG leukemic cells. This suggests that fRAG leukemic cells may

differentiate further, associated with an immune phenotype (Figure 2B). *IgH* rear-

rangement initiates with D_H-J_H joining in pro-B cells, followed by V_H-DJ_H joining in pro-

B cells, and ultimately, V_L-J_L rearrangements occur at the *IgL* loci in small pre-B cells

(**Schatz and Ji**,*2011*). Genomic PCR analysis of DNA from GFP$^+$CD19$^+$ cells was

utilized to assess $V_H DJ_H$ rearrangement. The results showed a pronounced

oligoclonality in cRAG leukemic cells, with tumors consistently demonstrating rear-

rangements involving a restricted set of V_H family members. In contrast, fRAG

leukemias displayed significant polyclonality, evidenced by the widespread rear-

rangement of various V_H family members to all potential J_H1-3 segments, indicative of

a broader clonal diversity (Supplementary Figure 4). This observation aligns with the

more aggressive leukemia phenotype seen in cRAG *BCR-ABL1*$^+$ B-ALL mice. Such

oligoclonality in cRAG leukemic cells suggests a selection process driven by BCR-

ABL1-induced leukemia, favoring the emergence of a limited number of dominant

leukemic clones. The absence of non-core RAG regions appears to restrict the diver-

sity of leukemic clones, leading to the formation of oligoclonal tumors.

174　The loss of non-core RAG regions highlights genomic DNA
175　damage

The findings indicate that leukemic cells from three types of mice exhibited variable

arrests at the large pre-B cell stage, deviating from normal B cell developmental tra-

jectory. Typically, at this juncture, B cells initiate the degradation of RAG2 via the cy-

clin-dependent kinase cyclinA/Cdk2, leading to a downregulation of RAG activity. It is

therefore crucial to explore the impact of deletions in non-core regions on the ex-

pression and functionality of RAG in these leukemic cells. Analysis showed that both

RAG1 (cRAG1) and RAG2 (cRAG2) were present in GFP$^+$CD19$^+$ splenic leukemic

cells from *BCR-ABL1*$^+$ B-ALL mice across different genetic backgrounds (Figure 3A).

Notably, we observed an upregulation of cRAG1 and cRAG2 in leukemic cells from

cRAG1 or cRAG2 mice compared to those from fRAG mice (Figure 3A, Supplemen-

tary Figure 5A). The in vitro V(D)J recombination assay confirmed that different forms

of RAG exhibited cleavage activity in *BCR-ABL1*$^+$B-ALL (Figure 3B and Supplemen-

tary Figure 5B).

To examine the potential correlation between aberrant RAG activity and in-

creased DNA double-strand breaks (DSBs), we assessed levels of phosphorylated

H2AX (□-H2AX), a marker of DSB response, in leukemic cells from fRAG, cRAG1,

and cRAG2 mice (gated on GFP$^+$). This evaluation aimed to gauge DNA DSBs and

overall genomic instability. Flow cytometry analysis revealed elevated □-H2AX levels

in cRAG1 and cRAG2 leukemic cells compared to those from fRAG, indicating a

more pronounced role of cRAG in mediating somatic structural variants in *BCR-*

ABL1$^+$ B cells. These findings suggest enhanced expression of cRAG1 endonucle-

ases in cRAG1 leukemic cells and increased DNA damage in cells lacking core RAG

regions.

Off-target recombination mediated by RAG in *BCR-ABL1*$^+$ B cells

Genome-wide sequencing and analysis were performed to compare somatic struc-

tural variants (SVs) in *BCR-ABL1*$^+$ B cells derived from fRAG*,* cRAG1 and cRAG2

mice. The leukemic cells were sequenced with an average coverage of 25✗ (Sup-

plementary Table 1). The SVs generated by RAG were screened based on two crite-

ria: the presence of a CAC to the right (or GTG to the left) of both breakpoints, and its

occurrence within 21 bp from the breakpoint (***Mijušković, et al.***,***2015***). Further elabo-

ration on these criteria can be found in Supplementary Figure 6. Consequently, aber-

rant *V*-to-*V* junctions and *V* to intergenic regions were encompassed in five validated

abnormal rearrangements at *Ig* loci in cRAG leukemic mice (Supplementary Table 2).

Additionally, seven samples had 24 somatic structural variations, with an average of

3.4 coding region mutations per sample (range of 0-9), which is consistent with the

limited number of acquired somatic mutations observed in hematological cancers

(Figure 4 and Supplementary Table 3). The results of the study demonstrate that

fRAG cells had low SVs (0-1 per sample), cRAG1 cells exhibited higher SVs (6-9 per

sample) while cRAG2 cells had moderate SVs incidence (1-4 per sample) (Figure 4,

Supplementary Table 3). These findings suggest that cRAG may lead to an elevated

off-target recombination, eventually posing a threat to the *BCR-ABL1*[+] B cells ge-

nome.

Off-target V(D)J recombination characteristics in *BCR-ABL1*[+] B cells

We further examined the characteristics of the identified structural variants (SVs).

Specifically, we analyzed the exon-intron distribution profiles of 41 breakpoints from

24 SVs through genome analysis. The results indicated that 57% of the breakpoints

were located within the gene body, while 43% were enriched in the flanking se-

quences, the majority of which were identified as transcriptional regulatory sequenc-

es (Figure 5A). P and N nucleotides are recognized as distinctive characteristics of

V(D)J recombination (***Repasky, et al.,2004***). RSS-to-RSS and cRSS-to-cRSS re-

combination have P nucleotides lengths of 7 and 9, respectively, and N lengths of 5,

so nucleotide lengths are basically the same during RSS-to-RSS and cRSS-to-cRSS

recombination (Figure 5B). However, the frequency of P and N sequences in RSS-to-

RSS recombination was 50%/50% (P/N), compared to 4%/8% (P/N) in cRSS-to-

cRSS recombination (Figure 5B). This significant reduction in the frequency of P and

N sequences suggests that DNA repair at off-target sites in *BCR-ABL1*[+] B cells di-

verges from the classical V(D)J re-combination repair process.

The hybrid joints were notably prevalent in cRAG1 and cRAG2 leukemic cells
(93% and 100%, respectively), suggesting that the non-core regions of RAG may
play a role in inhibiting harmful transposition events (Figure 5C). To evaluate the ef-
fect of deleting non-core RAG regions on the emergence of oncogenic mutations, we
performed a comparative analysis of cancer-related genes across three types of leu-
kemic cells. We found that cRAG1 leukemic cells harbored a significantly higher
number of cancer genes compared to the other groups. This finding corresponds with
the most aggressive leukemia phenotype observed in cRAG1 *BCR-ABL1*[+] B-ALL
mice and associated changes in their transcription profiles.

The non-core regions have effects on RAG cleavage and off-target
244
recombination size in *BCR-ABL1*[+] B cells

Sequence logos were employed to visually contrast RSS and cRSSs within *Ig* and
non-*Ig* loci, respectively. Notably, the RSS elements in *Ig* loci displayed a higher simi-
larity to the canonical RSS, especially at critical functional sites. The initial four nu-
cleotides (highlighted) of the canonical heptamer sequence CACAGTG were recog-
nized as the cleavage site for fRAG. Conversely, in leukemic cells, the cleavage site
for cRAG was pinpointed to the first three nucleotides, the CAC trinucleotide, of the
heptamer sequence (Figure 6A). While both motifs (CACA and CAC) align with the
highly conserved segment of the RSS heptamer sequence, differences in the cRSS
sequences across off-target genes in both fRAG and cRAG mice suggest that dele-
tion of RAG's non-core regions broadens the spectrum of off-target substrates in
BCR-ABL1[+] B cells.
Antigen receptor genes are assembled by large-scale deletions and inversions
(*Teng, et al.,2015*). The off-target recombination size was determined as the DNA
fragment size spanning the two breakpoints. Our analysis demonstrated that both
fRAG and cRAG2 leukemic cells produced off-target recombinations with 100% and
92% of events, respectively, spanning over 10,000 bp in length. In contrast, cRAG1

leukemic cells showed only 6% of off-target recombinations exceeding 10,000 bp,

with 48% under 1,000 bp and 46% ranging between 1,000 to 10,000 bp (Figure 6BC).

These findings suggest that the cRAG1 variant primarily facilitates smaller-scale off-

target recombinations in *BCR-ABL1*[+] B cells, highlighting the role of the non-core

RAG1 region in influencing the extent of off-target recombination. The deletion of the

non-core RAG1 region appears to constrict the size of off-target recombination, po-

tentially contributing to the elevated frequency of off-target V(D)J recombination ob-

served in cRAG1 leukemic cells (Figure 6D).

270 **Discussion**

In this study, we have demonstrated that non-core region deletion of both RAG1 and

RAG2 leads to accelerated development of leukemia and increased off-target V(D)J

recombination in mice models of *BCR-ABL1*[+] B cells. Furthermore, we report re-

duced cRAG cleavage accuracy and off-target recombination size in cRAG leukemia

cells, which might contribute to exacerbated off-target V(D)J recombination of cRAG

BCR-ABL1[+] B cells. These findings suggest that the non-core regions, particularly the

non-core region of RAG1, play a crucial role in maintaining accuracy of V(D)J recom-

bination and genomic stability in *BCR-ABL1*[+] B cells.

Our findings suggest that leukemic cells with cRAG regions exhibit increased produc-

tion of hybrid joints, implying that non-core RAG regions might suppress the for-

mation of these hybrid joints in vivo. Post-cleavage synaptic complexes (PSCs),

comprising RAG proteins, coding ends, and RSS ends, are believed to have evolved

to form with optimal conformation and/or stability for conventional coding and RSS

end-joining(**Fugmann, et al.**,**2000**; **Libri, et al.**,**2021**). In contrast, cRAG PSCs could

promote RAG-mediated hybrid joints by facilitating closer proximity of coding and

RSS ends or by increasing PSC stability. It is also conceivable that fRAGs recruit

disassembly/remodeling factors to PSCs, a process that could allow non-homologous

end joining (NHEJ) factors to complete the normal reaction(**Fugmann, et al.**,**2000**).

Conversely, cRAGs may have a diminished recruitment capacity due to changes in

overall conformation or the absence of specific motifs, leading to more unstable

PSCs and a heightened risk of accumulating incomplete hybrid joints (**Raghavan, et**

al.,**2006**; **Talukder, et al.**,**2004**). Our data reveal that over 90% of junctions were hy-

brid joints in cRAG leukemic cells, a frequency exceeding that reported in previous

studies. This suggests that deficiencies in the NHEJ pathway could contribute to

chromosomal instability and lymphomagenesis(**Gaymes, et al.**,**2002**; **Rassool**,**2003**;

Scully, et al.,**2019**; **Wiegmans, et al.**,**2021**). Significantly, our analysis uncovered

variations in the NHEJ repair pathway among leukemic cells from different genetic

backgrounds, suggesting a potential aberrant expression of DNA repair pathways in

cRAG leukemic cells (Supplementary Figure 3B). These findings highlight the poten-

tial of cRAG to foster increased hybrid joint formation, especially when a normal

pathway for efficient coding and RSS joining is compromised in an NHEJ-aberrant

context.

Our data demonstrates that the deletion of the RAG1 non-core region results in

more severe off-target V(D)J recombination compared to the deletion of the RAG2

non-core region. This observation is supported by the fact that the RAG1 terminus

contains multiple zinc-binding motifs and ubiquitin ligase activity, which are known to

enhance the efficiency of the rearrangement reaction (**Beilinson, et al.**,**2021**; **Burn,**

et al.,**2022**). Furthermore, our research reveals that RAG1 expression persists in

BCR-ABL1$^+$ progenitor B cells, and deletion of the non-core region of RAG1 results in

elevated expression of RAG in comparison to fRAG. Consequently, as demonstrated

in this study, cRAG1 from *BCR-ABL1*$^+$ B leukemic mice is prone to generating off-

target V(D)J recombination. The distinct function of RAG1's non-core region in thymic

lymphomas of Tp53-/- mice and *BCR-ABL1*$^+$ B leukemic mice leads to dissimilar off-

target activity of cRAG1 (**Mijušković, et al.**,**2015**). Therefore, it would be intriguing to

replicate these analyses across various subtypes of ALL to further investigate this

phenomenon.

In human *ETV6-RUNX1* ALL, the *ETV6-RUNX1* fusion gene is believed to initi-

ate prenatally, yet the disease remains clinically latent until critical secondary events

occur, leading to leukemic transformation-"pre-leukemia to leukemia" (*Mijušković, et*

al.,2015). Genomic rearrangement, mediated by aberrant RAG recombinase activity,

is a frequent driver of these secondary events in *ETV6-RUNX1* ALL (*Chen, et*

al.,2021; *Papaemmanuil, et al.,2014*). In contrast, RAG mediated off-target V(D)J

recombination is also observed in *BCR-ABL1*[+] B-ALL. These oncogenic structural

variations can also be considered as secondary events that promote the transition -

"leukemia to aggressive leukemia". The enhancement of *BCR-ABL1*[+] B-ALL deterio-

ration and progression by cRAG in mouse model was consistent with our previous

study that RAG enhances BCR-ABL1 positive leukemic cell growth through its endo-

nuclease activity (*Yuan, et al.,2021*). Additionally, we showed that non-core RAG1

region deletion leads to increased cRAG1 expression and high RAG expression re-

lated to low survival in pediatric acute lymphoid leukemia (Figure 3A and Supplemen-

tary Figure 7). Therefore, more attention should be paid to the non-core RAG region

mutation in *BCR-ABL1*[+] B-ALL for the role of non-core region in leukemia suppres-

sion and off-target V(D)J recombination.

Methods

Mice

The C57BL/6 mice were purchased from the Experimental Animal Center of Xi'an

Jiaotong University, while cRAG1 (amino acids 384-1040) and cRAG2 (amino acids

1-383) were obtained from Dr. David G. Schatz (Yale University, New Haven, Con-

necticut, USA). The mice were bred and maintained in a specific pathogen-free (SPF)

environment at the Experimental Animal Center of Xi'an Jiaotong University. All ani-

mal-related procedures were in accordance with the guidelines approved by the Xi'an

Jiaotong University Ethics Committee for Animal Experiments.

Generation of Retrovirus Stocks

The pMSCV-BCR-BAL1-IRES-GFP vector is capable of co-expressing the human BCR-ABL1 fusion protein and green fluorescence protein (GFP), while the pMSCV-GFP vector serves as a negative control by solely expressing GFP. To produce viral particles, 293T cells were transfected with either the MSCV-BCR-BAL1-IRES-GFP or MSCV-GFP vector, along with the packaging vector PKAT2, utilizing the X-tremeGENE HP DNA Transfection Reagent from Roche (Basel, Switzerland). After 48 hours, the viral supernatants were collected, filtered, and stored at -80°C.

Bone Marrow Transduction and Transplantation

Experiments were conducted using mice aged between 6 to 10 weeks. *BCR-ABL1$^+$* B-ALL murine model was induced by utilizing marrow from donor mice who had not undergone 5-FU treatment. The donor mice were euthanized through CO_2 asphyxiation, and the bone marrow was harvested by flushing the femur and tibia with a syringe and 26-gauge needle. Erythrocytes were not removed, and 1×10^6 cells per well were plated in six-well plates. A single round of co-sedimentation with retroviral stock was performed in medium containing 5% WEHI-3B-conditioned medium and 10 ng/mL IL-7 (Peprotech, USA). After transduction, cells were either transplanted into syngeneic female recipient mice (1×10^6 cells each) that had been lethally irradiated (2×450 cGy), or cultured in RPMI-1640 (Hyclone, Logan, UT) medium supplemented with 10% fetal calf serum (Hyclone), 200 mmol/L L-glutamine, 50 mmol/L 2-mercaptoethanol (Sigma, St Louis, MO), and 1.0 mg/mL penicillin/streptomycin (Hyclone). Subsequently, recipient mice were monitored daily for indications of morbidity, weight loss, failure to thrive, and splenomegaly. Weekly assessment of peripheral blood GFP percentage was done using FACS analysis of tail vein blood. Hematopoietic tissues and cells were utilized for histopathology, in vitro culture, FACS analysis, secondary transplantation, genomic DNA preparation, protein lysate preparation, or lineage analysis, contingent upon the unique characteristics of mice under

370   study.

Secondary Transplants

372   Thawed BM cells were sorted using a BD FACS Aria II (Becton Dickinson, San Jose

373   California, USA). GFP positive leukemic cells (1×10^6, 1×10^5, 1×10^4, and 1×10^3) were

374   then resuspended in 0.4 mL Hank's Balanced Salt Solution (HBSS) and intravenous-

375   ly administered to unirradiated syngeneic mice.

Flow cytometry analysis and sorting

Bone marrow, spleen cells, and peripheral blood were harvested from leukemic mice.

Red blood cells were eliminated using NH4Cl RBC lysis buffer, and the remaining

nucleated cells were washed with cold PBS. In order to conduct in vitro cell surface

receptor staining, 1×10^6 cells were subjected to antibody staining for 20 minutes at

4°C in 1×PBS containing 3% BSA. Cells were then washed with 1×PBS and ana-

lyzed using a CytoFLEX Flow Cytometer (Beckman Coulter, Miami, FL) or sorted on

a BD FACS Aria II. Apoptosis was analysed by resuspending the cells in Binding

Buffer (BD Biosciences, Baltimore, MD, USA), and subsequent labeling with anti-

annexin V-AF647 antibody (BD Biosciences) and propidium iodide (BD Biosciences)

for 15 minutes at room temperature. The lineage analysis was performed using the

following antibodies, which were purchased from BD Biosciences: anti-BP-1-PerCP-

Cy7, anti-CD19-PerCP-CyTM$^{5.5}$, anti-CD43-PE, anti-B220-APC, and anti-µHC-APC.

BrdU incorporation and analysis

390   Cells obtained from primary leukemic mice were cultured in six-well plates containing

391   RPMI-1640 medium supplemented with 10% FBS and 50 µg/mL BrdU. After incuba-

392   tion at 37°C for 30 minutes, the cells were harvested and intranuclearly stained with

393   anti-BrdU and 7-AAD antibodies, following the manufacturer's instructions.

The in vitro V(D)J recombination assay

The retroviral recombination substrate pINV-12/23 was introduced into primary leuke-mic cells utilizing X-treme GENE HP DNA Transfection Reagent (Roche). Recombina-tion efficiency of pINV-12/23 was evaluated through flow cytometry anal-ysis for mouse CD90 (mCD90) and hCD4 expression (**Yuan, et al.,2021**).

Western blotting analysis

Over 1×10^6 leukemic cells were centrifuged and washed with ice-cold PBS. The cells were then treated with ice-cold RIPA buffer, consisting of 50 mM Tris-HCl (pH 7.4), 0.15 M NaCl, 1% Triton X-100, 0.5% NaDoc, 0.1% sodium dodecyl sulphide (SDS), 1 mM ethylene diamine tetraacetic acid (EDTA), 1 mM phenylmethane sulphony fluoride (PMSF) (Amresco), and fresh protease inhibitor cocktail Pepstain A (Sigma). After sonication using a Bioruptor TMUCD-200 (Diagenode, Seraing, Bel-gium), the suspension was spined at 14,000 g for 3 minutes at 4°C. The total cell ly-sate was either utilized immediately or stored at -80°C. Protein concentrations were determined using DC Protein Assay (Bio–Rad Laboratories, Hercules, California, USA). Subsequently, the protein samples (20 μg) were incubated with α-RAG1 (mAb 23) and α-RAG2 (mAb 39) antibodies (**Teng, et al.,2015**), with GAPDH serving as the loading control. The signal was further detected using secondary antibody of goat anti-rabbit IgG conjugated with horseradish peroxidase (Thermo Scientific, Waltham, MA). The band signal was developed with Immobilon™ Western Chemiluminescent HRP substrate (Millipore, Billerica, MA). The band development was analyzed using GEL-PRO ANALYZER software (Media Cybernetics, Bethesda, MD).

Genomic PCR

Genomic PCR was performed in a 20μl reaction containing 50 ng of genomic DNA, 0.2 μm of forward and reverse primer, and 10 μl Premix Ex Taq (TaKaRa, Shiga, Ja-pan). Amplification conditions were as follows: 94°C for 5 minutes; 35 cycles of 30

seconds at 94°C, 30 seconds at 60°C and 1 minutes at 72°C; 72°C for 5 minutes

(BioRad, Hercules, CA). Genomic PCR was performed using the following primers:

D_HL-5′-GGAATTCGMTTTTTGTSAAGGGATCTACTACTGTG-3′;J_H3-5′-

GTCTAGATTCTCAC

-AAGAGTCCGATAGACCCTGG-3′; V_HQ52-5′-CGGTACCAGACTGARCATCASCAAG

-GACAAYTCC-3′; V_H558-5′-CGAGCTCTCCARCACAGCCTWCATGCARCTCARC-3′;

V_H7183-5′-CGGTACCAAGAASAMCCTGTWCCTGCAAATGASC-3′.(***Schlissel, et***

al.,1991):

RNA-seq library preparation and sequencing

GFP^+CD19^+ cells were sorted from the spleen of cRAG1 (n=3, $1×10^6$ cells /sample),

cRAG2 (n=3, $1×10^6$ cells /sample), and fRAG (n=3, $1×10^6$ cells /sample) *BCR-ABL1*[+]

B-ALL mice. Total RNA was extracted using Trizol reagent (Invitrogen, CA, USA) foll-

owing the manufacturer's guidelines. RNA quantity and purity analysis was done us-

ing Bioanalyzer 2100 and RNA 6000 Nano LabChip Kit (Agilent, CA, USA) with RIN

number >7.0. RNA-seq libraries were prepared by using 200 ng total RNA with

TruSeq RNA sample prep kit (Illumina). Oligo(dT)-enriched mRNAs were fragmented

randomly with fragmentation buffer, followed by first- and second-strand cDNA syn-

thesis. After a series of terminal repair, the double-stranded cDNA library was ob-

tained through PCR enrichment and size selection. cDNA libraries were sequenced

with the Illumina Hiseq 2000 sequencer (Illumina HiSeq 2000 v4 Single-Read 50 bp)

after pooling according to its expected data volume and effective concentration. Two

biological replicates were performed in the RNA-seq analysis. Raw reads were then

aligned to the mouse genome (GRCm38) using Tophat2 RNA-seq alignment soft-

ware, and unique reads were retained to quantify gene expression counts from

Tophat2 alignment files. The differentially expressed mRNAs and genes were select-

ed with log2 (fold change) >1 or log2 (fold change) <-1 and with statistical signifi-

cance (p value < 0.05) by R package. Bioinformatic analysis was performed using

the OmicStudio tools athttps://www.omicstudio.cn/tool.

Preparation of tumor DNA samples

GFP$^+$CD19$^+$ splenic cells, tail and kidney tissue were obtained from *cRAG1, cRAG2*

and *fRAG BCR-ABL1*$^+$ B-ALL mice, and genomic DNA was extracted using a

TIANamp Genomic DNA Kit (TIANGEN-DP304). Subsequently, paired-end libraries

were constructed from 1 µg of the initial genomic material using the TruSeq DNA v2

Sample Prep Kit (Illumina, #FC-121-2001) as per the manufacturer's instructions.

The size distribution of the libraries was assessed using an Agilent 2100 Bioanalyzer

(Agilent Technologies, #5067-4626), and the DNA concentration was quantified using

a Qubit dsDNA HS Assay Kit (Life Technologies, #Q32851). The Illumina HiSeq 4000

was utilized to sequence the samples, with two to four lanes allocated for sequencing

the tumor and one lane for the control DNA library of the kidney or liver, each with

150 bp paired end reads.

Read alignment and structural variant calling

Fastq files were generated using Casava 1.8 (Illumina), and BWA 37 was employed

to align the reads to mm9. PCR duplicates were eliminated using Picard's Mark Du-

plicates   tool   *(source-forge.net/apps/mediawiki/picard)*.   Our   custom   scripts

(http://sourceforge.net/projects/svdetection)   were   utilized   to   eliminate   BWA-

designated concordant and read pairs with low BWA mapping quality scores.

Intrachromosomal and inter-chromosomal rearrangements were identified using SV

Detect from discordant, quality prefiltered read pairs. The mean insertion size and

standard deviation for this analysis were obtained through Picard's InsertSizeMetrics

tool (sourceforge.net/apps/mediawiki/picard). Tumor-specific structural variants (SVs)

were   identified   using   the   manta   software

(https://github.com/Illumina/manta/blob/mater/docs/userGuide/README.md#introduc

tion).

Validation of high confidence off-target candidates

The elimination of non-specific structural mutations from the kidney or tail was necessary for tumor-specific structural variants identification. Subsequently, the method involving 21-bp CAC-to-breakpoint was employed to filter RAG-mediated off-target gene. The validation of high confidence off-target candidates was carried out through PCR. Oligonucleotide primers were designed to hybridize within the "linking" regions of SV Detect, in the appropriate orientation. The PCR product was subjected to Sanger sequencing and aligned to the mouse mm9 reference genome using BLAST *(https://blast.ncbi.nlm.nih.gov/Blast.cgi)*.

Statistics.

Statistical analysis was conducted using SPSS 20.0 (IBM Corp.) and GraphPad Prism 6.0 (GraphPad Software). Descriptive statistics were reported as means ± standard deviation for continuous variables. Statistical analyses were applied to biologically independent mice or technical replicates for each experiment which was independently repeated at least three times. The equality of variances was assessed using Levene's test. Two-group comparisons, multiple group comparisons, and survival comparisons were performed using independent-samples t-test, one-way analyses of variance (ANOVA) with post hoc Fisher's LSD test, and log-rank Mantel-Cox analysis, respectively. Kaplan-Meier survival curves were utilized to depict the changes in survival rate over time. Statistical significance was set at $P<0.05$.

Disclosure of Potential Conflicts of Interest

The authors declare no potential conflicts of interest.

Authors' Contributions

Yanghong Ji: Conceptualization, resources, data curation, funding acquisition, validation, writing-review, and editing. Xiaozhuo Yu and Wen Zhou: Conceptualization, validation, visualization, methodology, writing-original draft, writing-review, and editing. Xiaodong Chen: validation, writing-review, and editing. Shunyu He: methodology, writing-review, and editing. Mengting Qin: writing-review, and editing. Meng Yuan: validation, writing-review, and editing. Yang Wang: validation, writing-review and editing. Woodvine otieno Odhiambo: writing-review and editing. YinSha Miao: funding, validation, writing-review, and editing.

Acknowledgments

This study was supported by grants (no. 31170821, no. 31370874 and no. 81670157) from the National Natural Scientific Foundation of China and by a grant (no. 2016JZ0 -30) from the Natural Scientific Foundation of Shaanxi. The authors would like to thank Professor Shaoguang Li from the Division of Hematology/Oncology, University of Massachusetts Medical School, for providing the MSCV-BCR-BAL1-IRES-GFP construct. The authors would also like to thank Mr. Xiaofei Wang (Xi'an Jiaotong University Health Science Centre) for providing expert technical assistance with cell sorting.

References:

Banerjee JK, Schatz DG. 2014. Synapsis alters RAG-mediated nicking at Tcrb recombination signal sequences: implications for the "beyond 12/23" rule. *MOL CELL BIOL* **34**:2566-2580. doi:10.1128/MCB.00411-14

Bateman CM, Colman SM, Chaplin T, Young BD, Eden TO, Bhakta M, Gratias EJ, van Wering ER, Cazzaniga G, Harrison CJ, Hain R, Ancliff P, Ford AM, Kearney L, Greaves M. 2010. Acquisition of genome-wide copy number alterations in monozygotic twins with acute lymphoblastic leukemia. *BLOOD* **115**:3553-3558. doi:10.1182/blood-2009-10-251413

Becker LM, O'Connell JT, Vo AP, Cain MP, Tampe D, Bizarro L, Sugimoto H, McGow AK, Asara JM, Lovisa S, McAndrews KM, Zielinski R, Lorenzi PL, Zeisberg M, Raza S, LeBleu VS, Kalluri R.

2020. Epigenetic Reprogramming of Cancer-Associated Fibroblasts Deregulates Glucose Metabolism and Facilitates Progression of Breast Cancer. *CELL REP* **31**:107701. doi:10.1016/j.celrep.2020.107701

Beilinson HA, Glynn RA, Yadavalli AD, Xiao J, Corbett E, Saribasak H, Arya R, Miot C, Bhattacharyya A, Jones JM, Pongubala J, Bassing CH, Schatz DG. 2021. The RAG1 N-terminal region regulates the efficiency and pathways of synapsis for V(D)J recombination. *J EXP MED* **218**doi:10.1084/jem.20210250

Beilinson HA, Glynn RA, Yadavalli AD, Xiao J, Corbett E, Saribasak H, Arya R, Miot C, Bhattacharyya A, Jones JM, Pongubala J, Bassing CH, Schatz DG. 2021. The RAG1 N-terminal region regulates the efficiency and pathways of synapsis for V(D)J recombination. *J EXP MED* **218**doi:10.1084/jem.20210250

Bhojwani D, Pei D, Sandlund JT, Jeha S, Ribeiro RC, Rubnitz JE, Raimondi SC, Shurtleff S, Onciu M, Cheng C, Coustan-Smith E, Bowman WP, Howard SC, Metzger ML, Inaba H, Leung W, Evans WE, Campana D, Relling MV, Pui CH. 2012. ETV6-RUNX1-positive childhood acute lymphoblastic leukemia: improved outcome with contemporary therapy. *LEUKEMIA* **26**:265-270. doi:10.1038/leu.2011.227

Braams M, Pike-Overzet K, Staal F. 2023. The recombinase activating genes: architects of immune diversity during lymphocyte development. *FRONT IMMUNOL* **14**:1210818. doi:10.3389/fimmu.2023.1210818

Burn TN, Miot C, Gordon SM, Culberson EJ, Diamond T, Kreiger PA, Hayer KE, Bhattacharyya A, Jones JM, Bassing CH, Behrens EM. 2022. The RAG1 Ubiquitin Ligase Domain Stimulates Recombination of TCRβ and TCRα Genes and Influences Development of αβ T Cell Lineages. *J IMMUNOL* **209**:938-949. doi:10.4049/jimmunol.2001441

Chen D, Camponeschi A, Nordlund J, Marincevic-Zuniga Y, Abrahamsson J, Lönnerholm G, Fogelstrand L, Mårtensson IL. 2021. RAG1 co-expression signature identifies ETV6-RUNX1-like B-cell precursor acute lymphoblastic leukemia in children. *Cancer Med* **10**:3997-4003. doi:10.1002/cam4.3928

Curry JD, Schlissel MS. 2008. RAG2's non-core domain contributes to the ordered regulation of V(D)J recombination. *NUCLEIC ACIDS RES* **36**:5750-5762. doi:10.1093/nar/gkn553

Deng Z, Liu H, Liu X. 2015. RAG1-mediated ubiquitylation of histone H3 is required for chromosomal V(D)J recombination. *CELL RES* **25**:181-192. doi:10.1038/cr.2015.1

Eastman QM, Leu TM, Schatz DG. 1996. Initiation of V(D)J recombination in vitro obeying the 12/23 rule. *NATURE* **380**:85-88. doi:10.1038/380085a0

Fatma H, Maurya SK, Siddique HR. 2022. Epigenetic modifications of c-MYC: Role in cancer cell reprogramming, progression and chemoresistance. *SEMIN CANCER BIOL* **83**:166-176. doi:10.1016/j.semcancer.2020.11.008

Fugmann SD, Lee AI, Shockett PE, Villey IJ, Schatz DG. 2000. The RAG proteins and V(D)J recombination: complexes, ends, and transposition. *ANNU REV IMMUNOL* **18**:495-527. doi:10.1146/annurev.immunol.18.1.495

Gan T, Wang Y, Liu Y, Schatz DG, Hu J. 2021. RAG2 abolishes RAG1 aggregation to facilitate V(D)J recombination. *CELL REP* **37**:109824. doi:10.1016/j.celrep.2021.109824

Gaymes TJ, North PS, Brady N, Hickson ID, Mufti GJ, Rassool FV. 2002. Increased error-prone non homologous DNA end-joining--a proposed mechanism of chromosomal instability in Bloom's syndrome. *ONCOGENE* **21**:2525-2533. doi:10.1038/sj.onc.1205331

Goel S, Bhatia V, Biswas T, Ateeq B. 2022. Epigenetic reprogramming during prostate cancer progres-
sion: A perspective from development. *SEMIN CANCER BIOL* **83**:136-151.
doi:10.1016/j.semcancer.2021.01.009

Hardy RR, Hayakawa K. 2001. B cell development pathways. *ANNU REV IMMUNOL* **19**:595-621.
doi:10.1146/annurev.immunol.19.1.595

Hirokawa S, Chure G, Belliveau NM, Lovely GA, Anaya M, Schatz DG, Baltimore D, Phillips R. 2020.
Sequence-dependent dynamics of synthetic and endogenous RSSs in V(D)J recombination. *NUCLEIC*
ACIDS RES **48**:6726-6739. doi:10.1093/nar/gkaa418

Khoshchehreh R, Totonchi M, Carlos RJ, Torres R, Baharvand H, Aicher A, Ebrahimi M, Heeschen C.
2019. Epigenetic reprogramming of primary pancreatic cancer cells counteracts their in vivo
tumourigenicity. *ONCOGENE* **38**:6226-6239. doi:10.1038/s41388-019-0871-x

Kumari R, Roy U, Desai S, Nilavar NM, Van Nieuwenhuijze A, Paranjape A, Radha G, Bawa P,
Srivastava M, Nambiar M, Balaji KN, Liston A, Choudhary B, Raghavan SC. 2021. MicroRNA miR-
29c regulates RAG1 expression and modulates V(D)J recombination during B cell development. *CELL*
REP **36**:109390. doi:10.1016/j.celrep.2021.109390

Lewis SM, Agard E, Suh S, Czyzyk L. 1997. Cryptic signals and the fidelity of V(D)J joining. *MOL*
CELL BIOL **17**:3125-3136. doi:10.1128/MCB.17.6.3125

Libri A, Marton T, Deriano L. 2021. The (Lack of) DNA Double-Strand Break Repair Pathway Choice
During V(D)J Recombination. *FRONT GENET* **12**:823943. doi:10.3389/fgene.2021.823943

Liu C, Zhang Y, Liu CC, Schatz DG. 2022. Structural insights into the evolution of the RAG
recombinase. *NAT REV IMMUNOL* **22**:353-370. doi:10.1038/s41577-021-00628-6

Liu C, Zhang Y, Liu CC, Schatz DG. 2022. Structural insights into the evolution of the RAG
recombinase. *NAT REV IMMUNOL* **22**:353-370. doi:10.1038/s41577-021-00628-6

Liu Y, Subrahmanyam R, Chakraborty T, Sen R, Desiderio S. 2007. A plant homeodomain in RAG-2
that binds Hypermethylated lysine 4 of histone H3 is necessary for efficient antigen-receptor-gene rear-
rangement. *IMMUNITY* **27**:561-571. doi:10.1016/j.immuni.2007.09.005

Lu C, Ward A, Bettridge J, Liu Y, Desiderio S. 2015. An autoregulatory mechanism imposes allosteric
control on the V(D)J recombinase by histone H3 methylation. *CELL REP* **10**:29-38.
doi:10.1016/j.celrep.2014.12.001

Matthews AG, Kuo AJ, Ramón-Maiques S, Han S, Champagne KS, Ivanov D, Gallardo M, Carney D,
Cheung P, Ciccone DN, Walter KL, Utz PJ, Shi Y, Kutateladze TG, Yang W, Gozani O, Oettinger MA.
2007. RAG2 PHD finger couples histone H3 lysine 4 trimethylation with V(D)J recombination. *NA-*
TURE **450**:1106-1110. doi:10.1038/nature06431

Mendes RD, Sarmento LM, Canté-Barrett K, Zuurbier L, Buijs-Gladdines JG, Póvoa V, Smits WK,
Abecasis M, Yunes JA, Sonneveld E, Horstmann MA, Pieters R, Barata JT, Meijerink JP. 2014. PTEN
microdeletions in T-cell acute lymphoblastic leukemia are caused by illegitimate RAG-mediated re-
combination events. *BLOOD* **124**:567-578. doi:10.1182/blood-2014-03-562751

Mijušković M, Chou YF, Gigi V, Lindsay CR, Shestova O, Lewis SM, Roth DB. 2015. Off-Target
V(D)J Recombination Drives Lymphomagenesis and Is Escalated by Loss of the Rag2 C Terminus.
CELL REP **12**:1842-1852. doi:10.1016/j.celrep.2015.08.034

Mori H, Colman SM, Xiao Z, Ford AM, Healy LE, Donaldson C, Hows JM, Navarrete C, Greaves M.
2002. Chromosome translocations and covert leukemic clones are generated during normal fetal devel-
opment. *Proc Natl Acad Sci U S A* **99**:8242-8247. doi:10.1073/pnas.112218799

Onozawa M, Aplan PD. 2012. Illegitimate V(D)J recombination involving nonantigen receptor loci in

lymphoid malignancy. *Genes Chromosomes Cancer* **51**:525-535. doi:10.1002/gcc.21942

Papaemmanuil E, Rapado I, Li Y, Potter NE, Wedge DC, Tubio J, Alexandrov LB, Van Loo P, Cooke
SL, Marshall J, Martincorena I, Hinton J, Gundem G, van Delft FW, Nik-Zainal S, Jones DR, Rama-
krishna M, Titley I, Stebbings L, Leroy C, Menzies A, Gamble J, Robinson B, Mudie L, Raine K,
O'Meara S, Teague JW, Butler AP, Cazzaniga G, Biondi A, Zuna J, Kempski H, Muschen M, Ford AM,
Stratton MR, Greaves M, Campbell PJ. 2014. RAG-mediated recombination is the predominant driver
of oncogenic rearrangement in ETV6-RUNX1 acute lymphoblastic leukemia. *NAT GENET* **46**:116-125.
doi:10.1038/ng.2874

Raghavan SC, Tong J, Lieber MR. 2006. Hybrid joint formation in human V(D)J recombination re-
quires   nonhomologous   DNA   end   joining.   *DNA   Repair   (Amst)*   **5**:278-285.
doi:10.1016/j.dnarep.2005.09.008

Rassool FV. 2003. DNA double strand breaks (DSB) and non-homologous end joining (NHEJ) path-
ways in human leukemia. *CANCER LETT* **193**:1-9. doi:10.1016/s0304-3835(02)00692-4

Repasky JA, Corbett E, Boboila C, Schatz DG. 2004. Mutational analysis of terminal
deoxynucleotidyltransferase-mediated N-nucleotide addition in V(D)J recombination. *J IMMUNOL*
172:5478-5488. doi:10.4049/jimmunol.172.9.5478

Rooney S, Chaudhuri J, Alt FW. 2004. The role of the non-homologous end-joining pathway in lym-
phocyte development. *IMMUNOL REV* **200**:115-131. doi:10.1111/j.0105-2896.2004.00165.x

Sadofsky MJ, Hesse JE, McBlane JF, Gellert M. 1993. Expression and V(D)J recombination activity of
mutated RAG-1 proteins. *NUCLEIC ACIDS RES* **21**:5644-5650. doi:10.1093/nar/21.24.5644

Schabla NM, Perry GA, Palmer VL, Swanson PC. 2018. VprBP (DCAF1) Regulates RAG1 Expression
Independently of Dicer by Mediating RAG1 Degradation. *J IMMUNOL* **201**:930-939.
doi:10.4049/jimmunol.1800054

Schatz DG, Ji Y. 2011. Recombination centres and the orchestration of V(D)J recombination. *NAT*
REV IMMUNOL **11**:251-263. doi:10.1038/nri2941

Schatz DG, Swanson PC. 2011. V(D)J recombination: mechanisms of initiation. *ANNU REV GENET*
45:167-202. doi:10.1146/annurev-genet-110410-132552

Schjerven H, Ayongaba EF, Aghajanirefah A, McLaughlin J, Cheng D, Geng H, Boyd JR, Eggesbø
LM, Lindeman I, Heath JL, Park E, Witte ON, Smale ST, Frietze S, Müschen M. 2017. Genetic analy-
sis of Ikaros target genes and tumor suppressor function in BCR-ABL1(+) pre-B ALL. *J EXP MED*
214:793-814. doi:10.1084/jem.20160049

Schlissel MS, Corcoran LM, Baltimore D. 1991. Virus-transformed pre-B cells show ordered activa-
tion but not inactivation of immunoglobulin gene rearrangement and transcription. *J EXP MED*
173:711-720. doi:10.1084/jem.173.3.711

Scully R, Panday A, Elango R, Willis NA. 2019. DNA double-strand break repair-pathway choice in
somatic mammalian cells. *Nat Rev Mol Cell Biol* **20**:698-714. doi:10.1038/s41580-019-0152-0

Sekiguchi JA, Whitlow S, Alt FW. 2001. Increased accumulation of hybrid V(D)J joins in cells ex-
pressing   truncated   versus   full-length   RAGs.   *MOL   CELL*   **8**:1383-1390.   doi:10.1016/s1097-
2765(01)00423-3

Shi B, Dong X, Ma Q, Sun S, Ma L, Yu J, Wang X, Pan J, He X, Su D, Yao X. 2020. The Usage of
Human IGHJ Genes Follows a Particular Non-random Selection: The Recombination Signal Sequence
May   Affect   the   Usage   of   Human   IGHJ   Genes.   *FRONT   GENET*   **11**:524413.
doi:10.3389/fgene.2020.524413

Silver DP, Spanopoulou E, Mulligan RC, Baltimore D. 1993. Dispensable sequence motifs in the

RAG-1 and RAG-2 genes for plasmid V(D)J recombination. *Proc Natl Acad Sci U S A* **90**:6100-6104.

doi:10.1073/pnas.90.13.6100

Talukder SR, Dudley DD, Alt FW, Takahama Y, Akamatsu Y. 2004. Increased frequency of aberrant

V(D)J recombination products in core RAG-expressing mice. *NUCLEIC ACIDS RES* **32**:4539-4549.

doi:10.1093/nar/gkh778

Teng G, Maman Y, Resch W, Kim M, Yamane A, Qian J, Kieffer-Kwon KR, Mandal M, Ji Y, Meffre

E, Clark MR, Cowell LG, Casellas R, Schatz DG. 2015. RAG Represents a Widespread Threat to the

Lymphocyte Genome. *CELL* **162**:751-765. doi:10.1016/j.cell.2015.07.009

Thomson DW, Shahrin NH, Wang P, Wadham C, Shanmuganathan N, Scott HS, Dinger ME, Hughes

TP, Schreiber AW, Branford S. 2020. Aberrant RAG-mediated recombination contributes to multiple

structural rearrangements in lymphoid blast crisis of chronic myeloid leukemia. *LEUKEMIA* **34**:2051-

2063. doi:10.1038/s41375-020-0751-y

Wiegmans AP, Ward A, Ivanova E, Duijf P, Adams MN, Najib IM, Van Oosterhout R, Sadowski MC,

Kelly G, Morrical SW, O'Byrne K, Lee JS, Richard DJ. 2021. Genome instability and pressure on non-

homologous end joining drives chemotherapy resistance via a DNA repair crisis switch in triple nega-

tive breast cancer. *NAR Cancer* **3**:b22. doi:10.1093/narcan/zcab022

Wong S, Witte ON. 2004. The BCR-ABL story: bench to bedside and back. *ANNU REV IMMUNOL*

22:247-306. doi:10.1146/annurev.immunol.22.012703.104753

Yu X, Zhang H, Yuan M, Zhang P, Wang Y, Zheng M, Lv Z, Odhiambo WO, Li C, Liu C, Ma Y, Ji Y.

2019. Identification and characterization of a murine model of BCR‑ABL1+ acute B‑lymphoblastic

leukemia with central nervous system metastasis. *ONCOL REP* **42**:521-532. doi:10.3892/or.2019.7184

Yuan M, Wang Y, Qin M, Zhao X, Chen X, Li D, Miao Y, Otieno OW, Liu H, Ma Y, Ji Y. 2021. RAG

enhances BCR-ABL1-positive leukemic cell growth through its endonuclease activity in vitro and in

vivo. *CANCER SCI* **112**:2679-2691. doi:10.1111/cas.14939

Zhang L, Reynolds TL, Shan X, Desiderio S. 2011. Coupling of V(D)J recombination to the cell cycle

suppresses   genomic   instability   and   lymphoid   tumorigenesis.   *IMMUNITY*   **34**:163-174.

doi:10.1016/j.immuni.2011.02.003

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6