Title: Organoid-based *in vitro* system and reporter for the study of *Cryptosporidium parvum* sexual

reproduction

Running Title: Tools for studying *Cryptosporidium* sexual reproduction

Bethany R. Korwin-Mihavics[1,2,3], Emmett A. Dews[1,2], Peter Miller[1,2,] Alexandra Cameron[1,2], Bruno

Martorelli di Genova[2,3], Christopher D. Huston[1,2,3*]

[1]Department of Medicine, University of Vermont, Burlington, Vermont, USA

[2]Microbiology and Molecular Genetics, University of Vermont, Burlington, Vermont, USA

[3]Cellular, Molecular, and Biomedical Sciences, University of Vermont, Burlington, Vermont, USA

14   *Corresponding Author: Christopher Huston

15   95 Carrigan Drive, Stafford 202B, Burlington, VT 05401

16   (802) 656-9115

17   christopher.huston@med.uvm.edu

Abstract

Many advances have been made recently in our understanding of *Cryptosporidium*'s asexual cycle and sexual differentiation. However, the process of fertilization, which is required for transmission of infectious oocysts, is not well understood. Typical cancer cell-based culture only allows robust exploration of asexual cycle and sexual differentiation of *Cryptosporidium*. To facilitate exploration of sexual reproduction in *C. parvum* we developed an organoid-based culture system that supports *Cryptosporidium's* full life cycle and a novel fertilization reporter. Organoid derived monolayers (ODMs) supported fertilization and oocyst production and maintained the infection for up to 3 weeks. ODM derived oocysts were infectious *in vivo*. Fertilization was confirmed by successfully mating two strains of *C. parvum* and with a novel fertilization switch reporter. The fertilization switch reporter utilizes a DiCre system in which cre fragments are expressed under the control of sexual stage promoters resulting in a rapamycin-inducible switch in fluorescent protein expression from mCherry to mNeonGreen after fertilization that is spatially and temporally controlled. This results in mCherry positive parasites in the first generation and offspring that express mNeonGreen. *In vivo* validation of the fertilization switch reporter demonstrated the precision and efficiency of the fertilization switch reporter and confirmed excision of the mCherry gene sequence only after rapamycin treatment. The start of a second generation of parasites was also shown in the ODMs and rarely in HCT8s. Use of this reporter in ODMs can help investigate the *Cryptosporidium* lifecycle post sexual differentiation in a physiologically relevant *in vitro* system.

Importance

Organoid derived monolayers provide an opportunity to elucidate previously inaccessible aspects of *Cryptosporidium*'s biology. This system overcomes the disadvantages of previous

organoid-based methods for *Cryptosporidium* culture. It is faster and simpler than previously

described systems, uses defined media to increase reproducibility and consistency, enables real-

time observation, supports parasite fertilization and oocyst production, and provides a

physiologically relevant tissue culture system to facilitate studies of *Cryptosporidium* cell

biology. The ODM system could facilitate the study of host-pathogen interactions,

Cryptosporidium-host specificity, or innate or cellular immune responses to *Cryptosporidium*

infection stimulated in the intestinal epithelium. The fertilization switch reporter could be used to

test factors or drugs that may have potential to interfere with *Cryptosporidium's* sexual

reproduction. Organoid-based cell cultures in combination with the fertilization switch reporter

could increase our understanding of sexual reproduction in *Cryptosporidium,* leading to vital

information for the development of sexual reproduction inhibitors or vaccines that could shorten

disease duration and prevent transmission.

Introduction

Cryptosporidium is an intestinal protozoan parasite that is a major cause of diarrheal disease

worldwide and is among the 5 most common infectious causes of moderate to severe diarrhea in

children under 2 years[1–4]. Although not officially designated a neglected tropical disease, the

morbidity and mortality of *Cryptosporidium* outpaces that of many officially recognized

neglected tropical diseases[5]. Half of all waterborne outbreaks in the United States are caused by

Cryptosporidium, with *C. hominis* and *C. parvum* being the main species that cause outbreaks[6].

Cryptosporidium is found on every continent but Antarctica, but the heaviest burden of disease

falls on low-income nations and low-resource communities, especially in sub-Saharan Africa[1,3,5].

In immunocompetent adults, cryptosporidiosis is self-limited and usually resolves within 2-3

weeks, but immunocompromised individuals and young children, particularly malnourished

young children, are the most prone to developing severe disease that can become chronic and/or

life-threatening[1,2,5]. Currently, there is no vaccine and nitazoxanide, the only drug approved for

cryptosporidiosis, is poorly effective in the most vulnerable populations prone to severe

cryptosporidiosis. Chronic or repeated infection with *Cryptosporidium* can cause intestinal

damage resulting in loss of absorptive surface area and contribute to future malabsorption and

malnutrition, resulting in a vicious cycle of malnutrition and infection that can result in long-

term growth stunting and cognitive deficits in young children[7].

Cryptosporidium exhibits monoxenous development, whereby the entire lifecycle is completed

within a single host. Upon ingestion, oocysts pass through the acidic stomach environment and

into the small intestine where bile acids and digestive enzymes promote excystation to release

motile sporozoites[8,9]. Sporozoites invade the host cell, establish an epicellular niche in which the

parasite is intracellular but extracytoplasmic, and develop into trophozoites that mature into

meronts containing 8 merozoites which egress and reinvade new host cells[9–11]. After three rounds

of merogony, merozoites differentiate into sexual stages, the microgamont (male stage) and the

macrogamont (female stage)[10,12]. Microgamonts release microgametes which find and fertilize a

macrogamont forming a zygote which sporulates to form a new oocyst. Oocysts may excyst

within the gut lumen and begin a new round of infection or be excreted into the environment

where they wait to be ingested by a new host[10–12].

Due to its monoxenous development, a system of continuous culture should be simple, but such a

system has been elusive for *Cryptosporidium*. Approaches that have supported oocyst production

have not been widely adopted, suggesting excessive complexity, incompatibility with study

needs, or an inability to reproduce the results. Conventional HCT8 culture supports asexual

reproduction and sexual differentiation, but robust production of new oocysts is not supported,

suggesting a block to fertilization and zygote formation[7,12]. Culturing HCT8s in a candle jar has

been reported to support the production of thin-walled oocysts[13]. Culture in a bioreactor with

hollow nanofibers lined with HCT8s produced infectious oocysts for up to 20 weeks and

maintained oocyst production for 2 years, but this has not been widely adopted and is not

compatible with microscopy methods[14,15].

Organoid-based infection models offer a more promising and physiologically relevant approach

due to the presence of differentiated intestinal epithelial stem cells. One study using three

dimensional organoids derived from human intestinal epithelial cells showed new oocyst

production but required tedious microinjection of the enteroids for the infection[16]. Another

intestinal stem cell derived system utilized targeted laser ablation to create a Matrigel™ channel

that mimicked intestinal crypt and villi structures to serve as a scaffold for intestinal stem cell

attachment and differentiation and reported oocyst production for 4 weeks[17]. A stem cell-derived

air-liquid interface culture system also reported new oocyst production but required the

maintenance of two cell lines and the initial cell differentiation took up to 11 days[18]. The method

described here has been adapted from a number of protocols to support the full lifecycle of

Cryptosporidium in vitro[19–22]. Importantly, our method uses a defined media with readily

available reagents and equipment to support complete *Cryptosporidium* lifecycle development

physiologically relevant intestinal environment.

CRISPR/Cas9 systems enabling genetic modification[23,24], RNA-Seq and single cell RNA

sequencing data sets refining our knowledge of sexual differentiation[10,12,25,26] and new culture

systems have opened the possibility of studying fertilization *in vitro*[16,18]. The programmed nature

of *Cryptosporidium*'s life cycle in which sexual differentiation occurs after completion of three

rounds of asexual reproduction means that sexual reproduction is required for continued

expansion within the host, which makes treatments or vaccines targeting fertilization

mechanisms extremely attractive as they could shorten the disease and also help prevent

transmission[10].

Past approaches to measuring fertilization have relied on mating of different strains of *C.*

parvum[12,18]. While excellent approaches, they only capture interstrain crossing and we sought to

develop a reporter that would detect all fertilization events, including fertilization within the

same parasite strain. Here we present a novel fertilization reporter for use in studying

Cryptosporidium fertilization.

Results

Organoid derived monolayers exhibit the characteristics of differentiated intestinal

epithelium and support *C. parvum* infection. Confluent mouse intestinal stem cells were

grown on Matrigel-coated permeable supports with 50% stem cell maintenance media (SCMM)

for the first 24 hours, 5% SCMM for the second 24 hours, and were then allowed to differentiate

in differentiation media (DM) for 2 to 3 days before *C. parvum infection* (Fig. 1A).

Characteristics of functional intestinal epithelial tissue, e.g., microvilli (Fig. 1J and 1K) and tight

junctions (Fig. 1B, F, and K), and presence of differentiated cell types were confirmed by

immunofluorescence microscopy and gene expression. Goblet, enteroendocrine, and Paneth cells

were identified by staining for Muc2, Chromogranin A, and Lysozyme, respectively (Fig 1 G-I).

ZO-1 gene expression, indicating tight junction formation, was unchanged between spheroids

and ODMs (Fig. 1B). Stem cell marker Lgr5 gene expression was significantly reduced in

differentiated ODMs compared to spheroids. *Muc2* gene expression was significantly increased

in ODMs compared to spheroid culture, and chromogranin A (*ChgA*) expression was elevated in

ODMs, although not significantly (Fig. 1C-D). Altogether, these data confirmed differentiation

of the ODMs after 2-3 days in DM. Turnover of the ODMs was also observed. During media

changes, dead cells were observed in spent media, but the monolayers remained intact, indicating

that dead cells that had sloughed off were likely replaced with new cells (Supplemental Fig. 1D).

The cell turnover rate appeared comparable to that of intestinal epithelium *in vivo* (2-3 days in

mice and 3-5 days in humans)[27].

Differentiated monolayers infected with 10^6, 10^5, or 10^4 mNeonGreen expressing *C. parvum*

oocysts were tracked over the course of 3 weeks with live widefield epifluorescence microscopy

(representative images in Fig. 1L). The parasite vacuole (PV) load decreased slowly over the

course of 3 weeks (Fig. 1M). By 21 dpi, PVs were still present, but they were no longer present

in each field of view. The biggest drops in vacuole load were concurrent with supernatant

collections (Fig. 1M), and after an initial drop, the population often increased and leveled out

until the next supernatant collection was performed. These trends occurred in multiple similar

experiments (Supplementary Fig. 1). Over the course of the infection, an average of about 4.1

$x10^5$, 6.6 $x10^4$, and 2.3 x 10^4 PVs were estimated to have formed per monolayer based on

calculating the area under the curve of PVs for the total infection course for 10^6, 10^5, or 10^4

infective doses, respectively, with the estimated total PVs for the 10^6 dose being significantly

higher than the others (Fig. 1N). Oocysts collected from the supernatant of ODMs were

quantified by flow cytometry (Fig. 1O). Even though the 10^6 infective dose resulted in a

significantly higher number of estimated PVs (Fig. 1M-N), this did not correlate with the number

of oocysts produced (Fig. 1O-P) compared to the smaller infective doses. In fact, oocyst output

for each condition was comparable (Fig. 1P). Oocyst counts did not directly increase with the

drop in number of vacuoles observed after supernatant collection, likely because many of the

host cells that sloughed off the monolayer were also infected (Fig. 1D). Overall, organoid

derived monolayers support robust infection with *C. parvum* and production of new oocysts.

ODM derived oocysts are infectious *in vivo*. To verify the infectiousness of oocysts derived

from ODMs, HCT8 and ODM cultures were infected with tdTomato-nLuc *C. parvum*. The

parasites were excysted and filtered to exclude any unexcysted oocysts and ensure only free

sporozoites were used to infect the cultures and any oocysts collected from the cultures were

newly generated. Approximately equivalent surface areas were used for each set of cultures.

Supernatants were collected for 2 weeks, then concentrated by centrifugation and used to gavage

5 IFNγ-/- mice for each condition (Fig. 2A). Mouse fecal collections were performed over the

course of 3 weeks. Nanoluciferase assay analysis of the fecal samples indicated that all 5 mice

gavaged with ODM derived supernatants were infected, while none of the mice gavaged with

HCT8 derived supernatants were ever positive for nanoluciferase in the feces (Fig. 2B). This

indicates that ODMs support the *in vitro* development of new infectious oocysts in great enough

numbers to support subsequent *in vivo* infections.

Organoid derived monolayers support *C. parvum* mating. Organoid derived monolayers were

coinfected with mNeonGreen (mNG) and tdTomato (tdT) reporter *C. parvum* strains and

infections were observed and recorded with live imaging over the course of 3 weeks. Double

positive vacuoles indicative of mating were present (Fig. 3A), but rare compared to single

positive vacuoles (Fig. 4A-C). Oocyst output from ODMs can serve as a proxy for fertilization

and total oocyst output from the 3 experiments was not statistically significantly different (Fig.

4G-H). New oocysts with single positive, double positive, and double negative phenotypes were

also produced from these infections (Fig. 4D-F). Double positive oocysts were again rare events

and were not observed in all supernatant collections (Fig. 4D-F, Fig 4B). While most of the

oocyst population collected in supernatant from the ODMs was composed of double negative

oocysts, we cannot confirm that they resulted from outcrossing because a high proportion of

double negative oocysts were detected in the washes performed immediately after infection (Sup

Fig. 3B), suggesting that they were present in the input. Upon retrospective examination of the

tdT and mNG stocks used for these experiments, the mNG stock was confirmed to be a mixed

population of mNG and non-fluorescent oocysts. This means that in truth the results reflect the

mixing of 3 phenotypes rather than 2 and that vacuole counts from epifluorescence microscopy

lacked data for the double negative population and underrepresented the total parasite load.

Regardless of the limitations, the presence of double positive vacuoles within the ODM and

double positive oocysts in the supernatant combined with the total oocyst output and infection of

mice with ODM-derived oocysts confirm that fertilization and oocyst production is supported in

organoid derived monolayers.

A novel fertilization reporter: Fertilization Switch Reporter. Previous studies have focused

on mating 2 strains of parasites to measure fertilization and explore recombination[12,28]. These

studies successfully captured events resulting from interstrain crosses, but did not capture

intrastrain mating. Similarly, we initially crossed tdTomato and mNeonGreen reporter parasite

strains to visualize fertilization (Figures 3-4), which confirmed fertilization but quantification

was confounded by presence of non-fluorescent parasites and inability to identify intrastrain

mating. To be able to measure all fertilization events including "selfing", i.e. fertilization within

the same parasite strain, we developed a reporter that exhibits a color switch from red to green

after fertilization occurs (Figure 5). To this end, we utilized the DiCre system that has been used

by others in the parasitology field and has recently been demonstrated to be tractable in

Cryptosporidium[26,29–31]. The fertilization switch reporter is inducible and spatially and

temporally controlled. Cre subunits fused to FKB12 and FRB subunits dimerize in the presence

of rapamycin, bringing the Cre subunits together to make a functional CRE recombinase. To gain

spatial control of dimerization and precisely measure fertilization, one Cre subunit was expressed

in the microgametocyte under control of the HAP2 promoter (cgd8_2220) and the other

expressed only in the macrogamont under control of the COWP1 promoter (cgd6_2090). In this

way, the Cre subunits are only expressed in the sexual stages and are only in the same physical

space after fertilization and during zygote development. In the presence of rapamycin, functional

Cre recombinase will assemble in zygotes and excise a floxed mCherry-STOP sequence under

control of a constitutive aldolase promoter (cgd1_3020), leaving behind an mNeonGreen

sequence. A switch from red to green parasites in the second generation should be observed.

We first sought to validate the functionality of the fertilization switch reporter *in vivo*.

Mice infected with the fertilization switch reporter strain were either left untreated or gavaged

with 10 mg/kg rapamycin for 4 days (three doses/day for a total of 10 mg/kg) (Fig. 6A). Seven

days post treatment, fecal samples were collected, purified, and analyzed. Oocysts in mouse fecal

samples were analyzed by epifluorescence microscopy (Fig. 6B) and flow cytometric analysis

(Fig. 6C-D). Microscopy confirmed mCherry fluorescence and mNeonGreen fluorescence of

oocysts that were overwhelmingly mNeonGreen positive and negative for mCherry following

rapamycin treatment (Fig. 6B). An mCherry positive oocyst was observed only once under the

microscope in oocyst preparations made from the rapamycin treated group, while oocysts from

untreated mouse fecal samples were exclusively mCherry positive. When quantified by flow

cytometry, the oocyst population in feces from mice that received rapamycin treatment shifted

from mCherry positive to over 95% mNeonGreen positive, 11.5% of which were positive for

both mCherry and mNeonGreen (Fig. 6C-D). PCR of the region containing the loxP-

mCherrySTOP-loxP-mNeonGreen sequence from purified fecal oocysts showed a band shift

consistent with successful excision of mCherry in the rapamycin treated group (Fig. 6E). No

unexcised band was detected in the rapamycin treated group. Collectively, the data indicated that

the color switch post rapamycin treatment was a highly specific and efficient marker of

fertilization.

Fertilization switch reporter functions in ODMs. Next, we sought to test the fertilization

switch reporter in ODM culture. We had previously shown interstrain mating in ODMs and

wanted to determine if we could capture intrastrain mating in ODMs as well. Unfortunately, the

fluorescence of the fertilization switch reporter was not bright enough to readily observe the

fertilization reporter live in culture as it was intended. However, by staining with anti-RFP and

anti-mNeonGreen antibodies the fluorescent proteins could be detected. After validating the

function of the fertilization switch reporter *in vivo*, we wanted to use it to confirm its function in

the ODMs.

ODMs were infected with the fertilization switch reporter, wells were washed 16 hpi, and

(apical) media was replaced with differentiation media with or without 100 nM rapamycin. A

week after infection, the ODMs were fixed, stained, and imaged. In untreated transwells,

mNeonGreen positive vacuoles were not observed while between 25-40% of vacuoles in the

rapamycin treated ODMs were mNeonGreen positive (Fig 7A-B). Interestingly, mononucleated

and multinucleated mNeonGreen positive vacuoles corresponding to the sizes of trophozoites

and meronts were also observed, indicative of autoinfection and a second generation of

Cryptosporidium in the ODMs (Figure 8).

Fertilization in HCT8 cell monolayers: The fertilization switch reporter was also used to infect

HCT8 cells, expecting to confirm a complete block to fertilization. No mNeonGreen clusters

were observed in the absence of rapamycin treatment (Fig. 9A). To our surprise, clusters of

mNeonGreen positive vacuoles were observed repeatedly following rapamycin treatment,

although they were rare compared to mCherry positive vacuoles (Fig. 9B-C; 88 hpi). Based on

the number, size, and distribution of vacuoles, these clusters likely represented sites of

fertilization followed by a single round of merogony. Interestingly, mNeonGreen positive

parasite clusters were not detected at 48 hpi but at 90 hpi were detected more frequently when

infecting HCT8 cell monolayers grown in fibronectin-coated MatTek plates (Fig 9D). This could

be related to coating the MatTek plates with fibronectin before seeding HCT8s, which may have

promoted stronger polarization of the cell monolayer.

Discussion

This study shows that organoid-based cultures generated using defined culture media contain differentiated cell types representative of intestinal epithelial tissue and support *Cryptosporidium* infection and sexual reproduction. Organoid derived monolayers (ODMs) support *C. parvum* infection and new infectious oocyst production for at least 21 days and enable studies of fertilization. Important advances specific to this study include use of a defined culture medium that will enable simpler reproduction and use of the system by the community of *Cryptosporidium* researchers, and a precise tool in the fertilization switch reporter parasite with which to quantify fertilization that should enable new insights into the mechanisms underlying the largely unstudied *Cryptosporidium* sexual replication cycle.

Organoid-based models represent the major cell types found in the small intestine and provide an environment that better mimics the *in vivo* environment. We confirmed the presence of enteroendocrine cells, goblet cells, Paneth cells, enterocytes, and stem cells by gene and protein expression, which aligns with reported data[16,22,32–34]. The presence of tuft cells and M cells was not confirmed here, but other groups have demonstrated them after adding media supplements to promote their differentiation[35,36]. We have observed cell turnover in both formats resembling cell turnover found in the gut *in vivo*, which occurs about every 3 days in the mouse intestine (3-5 in human intestine)[7,27,37–40]. Host cells sloughed off the monolayer and were replaced by new cells in both uninfected and infected conditions. We did not investigate whether turnover rate is affected by infection. Many cells that sloughed off ODMs or enteroids were infected by various stages of *C. parvum* (Sup Fig. 1D). Like many other protozoans, *Cryptosporidium* inhibits host cell apoptosis through NFκB[41–46]. Based on previous studies in biliary epithelium, it may be that uninfected cells in proximity to infected cells are undergoing

apoptosis despite prolonged survival of infected cells[47]. A recent study determined that infection

did increase cell turnover, but that infected cells were less likely to be apoptotic than neighboring

uninfected cells[48]. Mechanisms mediating cell turnover and/or inhibition of apoptosis during

Cryptosporidium infection are not fully understood, but this system could be helpful in exploring

them.

ODMs are ready for infection within 5 days, which more closely resembles the maturation,

differentiation, and turnover of cells in the intestine. This is 5-6 days faster than monolayers

using the stem cell derived air-liquid interface (ALI) system used in a previously reported

Cryptosporidium infection model[18]. The defined media used here does not rely on serum

supplementation and should not be subject to the same lot-to-lot variation inherent in fetal bovine

serum-based media used for the ALI method, and thus be more reproducible. Past studies have

demonstrated that serum supplementation may not be sufficient alone to provide the necessary

components to support robust oocyst production[14] or that serum may actually inhibit the

infection in some cases[49].

The main goal of this project was to model successful sexual reproduction leading to

production of infectious oocysts in an *in vitro* system, since the programmed nature of

*Cryptosporidium'*s lifecycle requires sex for continued infection[10]. Our mating assay confirmed

that ODMs support *C. parvum* fertilization *in vitro*. Double positive vacuoles indicated fusion of

macrogamonts and microgametes and confirmed that fertilization does indeed occur in ODMs.

Double positive vacuoles were likely indicative of zygotes rather than recombined double

positive offspring due to the proximity of the mNG and tdT genes on chromosome 5 lowering

the probability of recombination resulting in double positive parasites. We would have expected

an expansion of the double positive population if successful recombination between those two

genes had occurred. This does not mean no recombination occurred, only that it did not occur

frequently between the mNG and tdT genes. However, this mating assay method greatly

underestimates the number of fertilization events because it fails to reflect the intrastrain

fertilization events observed as single positive.

Our fertilization switch reporter is a sensitive tool to measure fertilization that overcomes

these shortcomings. Validation of the fertilization switch reporter in mice demonstrated that the

color switch is tightly controlled and that rapamycin administration results in a robust shift from

tdT to mNG expressing parasites after fertilization occurs. In ODMs, the fertilization switch

reporter demonstrated production of mNeonGreen vacuoles of various sizes that were both

mononucleated and multinucleated, suggesting they were trophozoites or meronts. Many

mCherry positive vacuoles also remained; some appeared to be meronts (multinucleated) and

some appeared to be, presumably, unfertilized macrogamonts. If unfertilized macrogamonts can

survive and remain in the intestine until another round of sexual differentiation occurs, they

could have another chance to be fertilized. The presence of apparent mNeonGreen positive

trophozoites and meronts indicates that new oocysts were produced, excysted, and sporozoites

infected new host cells. Oocyst excystation has been shown to have a number of triggers, but

sometimes only one trigger is required to induce excystation[8]. Changes in temperature, pH,

exposure to digestive enzymes and bile acids, and even binding to oocyst wall receptors can

trigger excystation[8,9]. It is possible that during media changes oocysts present in culture

experienced a temperature change significant enough to trigger excystation. Alternatively, oocyst

wall receptors may have bound to host ligands that could have triggered excystation.

Surprisingly, our data using the fertilization switch reporter indicate that the block to

fertilization in HCT8 cell culture is not absolute. This result runs counter to a recent mating

experiment that detected fertilization *in vivo* but not in HCT8 cell culture using a method in

which a Cre recombinase bearing *C. parvum* strain was crossed with a *C. parvum* strain with a

floxed termination sequence upstream of a fluorescent reporter gene[12]. In contrast, our data

strongly suggest that fertilization and oocyst production, followed by autoinfection occurs in

HCT8 cell culture, albeit rarely. The accuracy of this conclusion is supported by the apparent

specificity of the fertilization switch reporter *in vivo* and in tissue culture, and by observation of

clustered NeonGreen positive vacuoles of various sizes corresponding to different life cycle

stages. Interestingly, more mNG clusters were observed after infecting HCT8 monolayers

cultured on fibronectin-coated coverslips. As a member of the extracellular matrix, fibronectin

could have contributed to enhanced monolayer polarization, but this was not explored[50,51]. The

discrepancy with prior work is likely due to enhanced sensitivity of the fertilization switch

reporter which theoretically detects all fertilization events compared to previously used methods

unable to detect intrastrain mating. Our data are consistent with earlier *Cryptosporidium* research

in which production of small numbers of new, usually thin-walled, oocysts in various culture

systems was reported, including in HCT8 cell culture[13,14,52]. Nonetheless and consistent with

numerous prior reports, the rate of fertilization and oocyst production from HCT8 cell culture

was inadequate to support infection of mice, which was only possible with material derived from

the new organoid-based culture method. However, our experiments show that HCT8 culture did

not support the development of sufficient oocysts to infect mice, which is an inefficient process

that would not be detected when coupled with low numbers of oocysts produced in HCT8s. It is

also possible that HCT8-derived oocysts are defective. The necessary fertilization enabling

factors may be present in HCT8 cell culture but at low levels leading to inefficient and infrequent

fertilization. If HCT8s are only capable of supporting production of thin-walled oocysts, those

oocysts may not survive passage through stomach acid in the mouse. Explorations into why

fertilization and autoinfection occur rarely in HCT8s and more frequently in ODMs and *in vivo*

could provide opportunities to develop still simpler continuous culture methods and/or to identify

drug or vaccine targets to shorten the duration of infection and prevent transmission.

The programmed nature of the *Cryptosporidium* lifecycle suggests that persistent

infection and the disease course are dependent on an appreciable rate of autoinfection. The

parasite must need to balance the rate of autoinfection and the rate of release of mature oocysts

in the environment in order to find new hosts. In immunocompromised individuals where the

infection can become chronic, there may be factors at play that increase the rate of autoinfection.

This could be an interesting avenue of research that could benefit the populations most prone to

severe cryptosporidiosis.

Proximity of the different strains may be a factor for detecting fertilization via mating

experiments using different colored reporters. As has been proposed previously, male and female

gamonts develop in close proximity which can facilitate fertilization within the same strain

allowing the lifecycle to restart [10]. Our ability to detect the rare fertilization events in HCT8s

using the more sensitive fertilization switch reporter may hinge on this principle.

Whether or to what extent microgametes are motile is still unknown, thus proximity to

female gametes may be extremely important for facilitating fertilization. Crossing separate

strains may not have detected fertilization due to spacing of initial infection clusters of the

different strains and perhaps the sexual stages were not in close enough proximity to facilitate a

cross. It may be that *in vivo* more than one complete turn of the life cycle is required before cross

fertilization of different strains can occur. Alternatively, the intestinal environment *in vivo* likely

provides help in the way of moving microgametes around via peristalsis and flow of luminal

contents. Additionally, the tube shape of the intestinal lumen also suggests that parasites may

happen upon a new host cell or sexual partner no matter which direction they go, whereas in 2D

culture they are limited to a single plane.

Organoid-based methods supporting the complete lifecycle of *Cryptosporidium* have the

potential to aid in making big strides toward understanding *Cryptosporidium*'s biology. This

system can capitalize on the vast library of transgenic mice to generate organoids with the

desired genotype and be used in tandem with genetic manipulation of *C. parvum* to interrogate

both sides of the host-parasite interactions[24]. Transfection of the organoids has also been

demonstrated by others if the desired host genotype does not exist[53]. Additionally, ODM co-

culture with various immune cell types could facilitate *in vitro* studies of immune cell responses

to infection[54,55]. Other studies have co-cultured various organoids or ODMs with T cells,

intraepithelial lymphocytes, or macrophages[56–60]. Because this method is based on a protocol that

makes shifting between host species simple with minor adjustments, it could facilitate the study

of the basis of host-parasite specificity[22]. Combining the ODMs with the Fertilization Switch

reporter can facilitate studies to identify and verify factors that enable fertilization and to test

sexual reproduction inhibitors. Interfering with the parasite's ability to complete sexual

reproduction has the potential to significantly reduce the duration of the infection and reduce

associated pathology.

Materials and Methods

Media Preparations

L-WRN conditioned media was prepared according to methods previously described[53].

Advanced DMEM/F12 supplemented with 20% FBS was conditioned with stem cell

maintenance ligands by culturing L-WRN cells for 12 days. L-WRN cells were engineered to

overexpress Wnt3a, R-spondin, and noggin which help maintain and grow intestinal epithelial

stem cells. Conditioned media was collected every 4 days and stored at -20°C.

To prepare stem cell maintenance media (SCMM), L-WRN conditioned media was diluted to

50% with Advanced DMEM/F12 and supplemented with 10 mM nicotinamide, 1 mM N-

acetylcysteine, 50 ng/mL mouse epidermal growth factor, 1x Pen/strep, 2 mM GlutaMax, 1 mM

HEPES, 1X N2 and 1X B27 supplements as described previously[22]. Differentiation media (DM)

consisted of Advanced DMEM/F12 supplemented with the same components as SCMM[22]. Five

percent SCMM was prepared by diluting 50% SCMM with DM.

Mouse Intestinal Epithelial Stem Cell Isolation, Culture, and Maintenance: Stem cells were

isolated for the small intestinal crypts of an 8 week old female C57Bl6/NJ mouse as described by

Miyoshi et al[53]. Briefly, the mouse was euthanized and the small intestine was harvested, flushed

with sterile PBS, opened lengthwise, scraped with a sterile glass slide to remove the villi, then

minced and submerged in a collagenase solution (2 mg/mL collagenase type 1 and 50 ug/mL

gentamicin in DMEM/F12). The tissue fragments were incubated at 37°C with vigorous pipette

mixing every 10 minutes for 1 hour until the crypts had separated from larger tissue fragments.

Isolated crypts were washed, resuspended in Matrigel™, and dropped on a cell culture plate to

form domes. The plate was inverted during polymerization to prevent cell clusters from

accumulating at the bottom of the Matrigel™ dome. Once polymerized, SCMM with 10 μM

Y27632 inhibitor and 10 μM SB431542 was overlaid. SCMM with 500 nM A83-01 and 1 μM

SB202190 added was changed every 2-3 days. For passage, spent media was aspirated, the

Matrigel™ domes were washed with PBS, and then overlaid with Trypsin/EDTA and

mechanically dissociated with a pipette. Spheroids were incubated with Trypsin/EDTA at 37°C

for 2-5 minutes and then washed with washing medium (DMEM/F12, 10% FBS, 2 mM L-

glutamine, 15 mM HEPES, and 1x pen/strep). Cells (usually a 1:5 passage) were centrifuged at

200 x g for 5 minutes. The supernatant was removed, and spheroids were resuspended in 2:1

Matrigel™ to SCMM and dotted onto plates to form domes, inverted, and incubated at 37°C

until the Matrigel™ polymerized. SCMM was then overlaid on the Matrigel™ domes. Stem cell

spheroids were passaged every 3-7 days depending on the size and density of the spheroids.

Organoid derived monolayer preparation: ODM preparation was adapted from a previously

described protocol[22]**.** Spheroids were expanded appropriately to set up desired experiments.

Permeable supports were coated with Matrigel™ diluted 1:10 in SCMM and incubated overnight

at 4°C. Prior to seeding, diluted Matrigel™ suspension was removed from wells and permeable

supports were allowed to dry at 37°C for 10 minutes. To prepare mISCs for seeding, spheroids

were mechanically dissociated in Trypsin-EDTA with vigorous pipetting and incubated for 3-5

minutes at 37°C. Cells were then washed in washing medium, filtered through 40 µm mesh cell

strainer, and washed once with PBS. Each permeable support was seeded with $3x10^5$ cells

resuspended in 100 µL SCMM supplemented with 10 µM Y27632 inhibitor per well ($3x10^5$

cells/permeable support). After 24 hours, media was changed to 5% SCMM with 10 µM Y27632

inhibitor for 24 hours and then switched to differentiation medium without Y27632 inhibitor

thereafter. After differentiating for 48 hours ODMs were ready for infection. ODMs were

maintained with 5% CO_2 atmosphere at 37°C. Apical and basal media was changed every 2-3

days. Frequency of media change for apical chambers may have varied due to supernatant

collections. New media was added to apical chambers after supernatant collections.

HCT8 Cell Culture: HCT8s were maintained in RPMI supplemented with 10% FBS, and 1x

Pen/Strep. HCT8s were seeded into a 96 well plate or fibronectin coated MatTek plates and

infected once confluent. Media was changed every other day. Cells were maintained with 5%

CO_2 atmosphere at 37°C.

Excystation and Infection: Appropriate numbers of oocysts were disinfected with bleach (10%)

for 10 minutes on ice, washed twice with PBS, incubated with 10 mM HCl at 37°C for 10

minutes, centrifuged for 5 minutes at 14000xg, resuspended in 2 mM sodium taurocholate and

incubated at 37°C for 10 minutes before infection of ODMs or HCT8s. For transfection, oocysts

were allowed to excyst for 45 minutes-1 hour in 2 mM sodium taurocholate at 37°C before

transfection. For filtered infections, oocysts were allowed to excyst for 1 hour in 2 mM sodium

taurocholate at 37°C before being filtered through a 1 μm syringe filter to remove unexcysted

oocysts and retain free sporozoites. Free sporozoites were resuspended in 100 μL/well of DM

(ODMs) or RPMI (HCT8s) and added to apical chambers of cell culture supports or to HCT8

cultures. For unfiltered infections, excysted oocysts and sporozoites in sodium taurocholoate

were resuspended in warm DM and dispensed into wells in 100 μL volumes. Sporozoites were

allowed to infect for 2-3 hours in the incubator at 37°C before washing 3 times with PBS to

remove unattached parasites.

Mouse infection with ODM or HCT8 derived supernatants: HCT8s were maintained in

RPMI supplemented with 10% FBS, and Pen/Strep with 5% CO_2 atmosphere at 37°C. HCT8s

were seeded into 24 wells of a 96 well plate and were infected once confluent. This format was

chosen to keep the culture surface area comparable between ODM (33.6 mm^2) and HCT8 culture

(33 mm^2/well). HCT8s and differentiated ODMs were infected with filtered tdTomato *C. parvum*

sporozoites (as described above). Supernatants were collected 3 times/week for 2 weeks and

stored at 4°C. Media volumes in HCT8s and ODM apical chambers were kept at 100 μL/well

throughout the experiment. After 2 weeks, supernatants were spun down at 16000xg for 5

minutes to concentrate any oocysts present and then resuspended in 500 µL of PBS.

Concentrated supernatants were gavaged and then used to gavage 5 mice/condition (IFNg-/-, 3-4

weeks old Male). Infection was monitored via the nanoluciferase assay (Promega).

Acknowledgements:

Funding for this study was provided by the Vermont Immunobiology and Infectious Disease Training

Grant 2T32AI055402-16A1 and RO1AI143951 to CDH.

We would like to thank the Harry Hood Bassett Flow Cytometry and Cell Sorting Facility

(RRID:SCR_022147) at the University of Vermont Larner College of Medicine in the generation of our

flow cytometry data.

Transmission electron microscopy sample preparation and imaging was performed at the Microscopy

Imaging Center at the University of Vermont (RRID# SCR_018821).

The qPCR and qRT-PCR were performed using equipment in Vermont Integrative Genomics Resource

DNA Facility and was supported by University of Vermont Cancer Center, Lake Champlain Cancer

Research Organization, and the UVM Larner College of Medicine.

We would like to thank Dr. Christian Klotz for consultations while refining this protocol for

Cryptosporidium infection, Dr. Boris Striepen for the generous gift of the tdTomato and mNeonGreen

strains of *C. parvum*, and Camilla Strother and Ethan Mattice for their help during the initial stem cell

isolations.

Figure 1A, 2A, and 3A were created with BioRender.com.

References

1. Kotloff, K. L. *et al.* The Global Enteric Multicenter Study (GEMS) of Diarrheal Disease in Infants and Young Children in Developing Countries: Epidemiologic and Clinical Methods of the Case/Control Study. *Clin. Infect. Dis. Off. Publ. Infect. Dis. Soc. Am.* **55**, S232–S245 (2012).

2. Checkley, W. *et al.* A review of the global burden, novel diagnostics, therapeutics, and vaccine targets for cryptosporidium. *Lancet Infect. Dis.* **15**, 85–94 (2015).

3. Kotloff, K. L. *et al.* Burden and aetiology of diarrhoeal disease in infants and young children in developing countries (the Global Enteric Multicenter Study, GEMS): a prospective, case-control study. *The Lancet* **382**, 209–222 (2013).

4. Troeger, C. *et al.* Estimates of global, regional, and national morbidity, mortality, and aetiologies of diarrhoeal diseases: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet Infect. Dis.* **17**, 909–948 (2017).

5. Gilbert, I. H. *et al.* Safe and effective treatments are needed for cryptosporidiosis, a truly neglected tropical disease. *BMJ Glob. Health* **8**, e012540 (2023).

6. Hlavsa, M. C. *et al.* Outbreaks Associated with Treated Recreational Water - United States, 2000-2014. *MMWR Morb. Mortal. Wkly. Rep.* **67**, 547–551 (2018).

7. Liu, J., Bolick, D. T., Kolling, G. L., Fu, Z. & Guerrant, R. L. Protein Malnutrition Impairs Intestinal Epithelial Cell Turnover, a Potential Mechanism of Increased Cryptosporidiosis in a Murine Model. *Infect. Immun.* **84**, 3542–3549 (2016).

8. Smith, H. V., Nichols, R. A. B. & Grimason, A. M. Cryptosporidium excystation and invasion: getting to the guts of the matter. *Trends Parasitol.* **21**, 133–142 (2005).

9. Lendner, M. & Daugschies, A. Cryptosporidium infections: molecular advances. *Parasitology* **141**, 1511–32 (2014).

10. English, E. D., Guérin, A., Tandel, J. & Striepen, B. Live imaging of the Cryptosporidium parvum life cycle reveals direct development of male and female gametes from type I meronts. *PLOS Biol.* **20**, e3001604 (2022).

11. *Cryptosporidium and Cryptosporidiosis*. (CRC Press Taylor & Francis Group, 2008).

12. Tandel, J. *et al.* Life cycle progression and sexual development of the apicomplexan parasite

Cryptosporidium parvum. *Nat. Microbiol.* **4**, 2226–2236 (2019).

13. Hijjawi, N. S., Meloni, B. P., Morgan, U. M. & Thompson, R. C. Complete development and long-

term maintenance of Cryptosporidium parvum human and cattle genotypes in cell culture. *Int. J.*

Parasitol. **31**, 1048–1055 (2001).

14. Morada, M. *et al.* Continuous culture of Cryptosporidium parvum using hollow fiber technology. *Int.*

J. Parasitol. **46**, 21–29 (2016).

15. Yarlett, N., Morada, M., Gobin, M., Van Voorhis, W. & Arnold, S. In Vitro Culture of

Cryptosporidium parvum Using Hollow Fiber Bioreactor: Applications for Simultaneous

Pharmacokinetic and Pharmacodynamic Evaluation of Test Compounds. *Methods Mol. Biol. Clifton*

NJ **2052**, 335–350 (2020).

16. Heo, I. *et al.* Modeling Cryptosporidium infection in human small intestinal and lung organoids. *Nat.*

Microbiol. **3**, 814–823 (2018).

17. Nikolaev, M. *et al.* Homeostatic mini-intestines through scaffold-guided organoid morphogenesis.

Nature **585**, 574–578 (2020).

18. Wilke, G. *et al.* A Stem-Cell-Derived Platform Enables Complete Cryptosporidium Development

In Vitro and Genetic Tractability. *Cell Host Microbe* **26**, 123-134.e8 (2019).

19. Co, J. Y., Margalef-Català, M., Monack, D. M. & Amieva, M. R. Controlling the polarity of human

gastrointestinal organoids to investigate epithelial biology and infectious diseases. *Nat. Protoc.* **16**,

5171–5192 (2021).

20. Nash, T. J., Morris, K. M., Mabbott, N. A. & Vervelde, L. Inside-out chicken enteroids with

leukocyte component as a model to study host–pathogen interactions. *Commun. Biol.* **4**, 1–15 (2021).

21. Stroulios, G. *et al.* Culture Methods to Study Apical-Specific Interactions using Intestinal Organoid

Models. *J. Vis. Exp. JoVE* (2021) doi:10.3791/62330.

22. Holthaus, D., Delgado-Betancourt, E., Aebischer, T., Seeber, F. & Klotz, C. Harmonization of

Protocols for Multi-Species Organoid Platforms to Study the Intestinal Biology of Toxoplasma

gondii and Other Protozoan Infections. *Front. Cell. Infect. Microbiol.* **10**, (2021).

23. Vinayak, S. *et al.* Genetic modification of the diarrhoeal pathogen Cryptosporidium parvum. *Nature*

523, 477–480 (2015).

24. Vinayak, S. Recent advances in genetic manipulation of Cryptosporidium. *Curr. Opin. Microbiol.* **58**,

146–152 (2020).

25. Walzer, K. A. *et al.* Transcriptional control of the Cryptosporidium life cycle. *Nature* 1–7 (2024)

doi:10.1038/s41586-024-07466-1.

26. Tandel, J. *et al.* Genetic Ablation of a Female-Specific Apetala 2 Transcription Factor Blocks Oocyst

Shedding in Cryptosporidium parvum. *mBio* **14**, e03261-22 (2023).

27. Barker, N. Adult intestinal stem cells: critical drivers of epithelial homeostasis and regeneration. *Nat.*

Rev. Mol. Cell Biol. **15**, 19–33 (2014).

28. Kimball, A. *et al.* Mendelian segregation and high recombination rates facilitate genetic analyses in

Cryptosporidium parvum. *PLOS Genet.* **20**, e1011162 (2024).

29. Knuepfer, E., Napiorkowska, M., van Ooij, C. & Holder, A. A. Generating conditional gene

knockouts in Plasmodium – a toolkit to produce stable DiCre recombinase-expressing parasite lines

using CRISPR/Cas9. *Sci. Rep.* **7**, 3881 (2017).

30. Collins, C. R. *et al.* Robust inducible Cre recombinase activity in the human malaria parasite

Plasmodium falciparum enables efficient gene deletion within a single asexual erythrocytic growth

cycle. *Mol. Microbiol.* **88**, 687–701 (2013).

31. Jullien, N., Sampieri, F., Enjalbert, A. & Herman, J. Regulation of Cre recombinase by ligand-

induced complementation of inactive fragments. *Nucleic Acids Res.* **31**, e131 (2003).

32. Luu, L. *et al.* An Open-Format Enteroid Culture System for Interrogation of Interactions Between

Toxoplasma gondii and the Intestinal Epithelium. *Front. Cell. Infect. Microbiol.* **9**, (2019).

33. Dutta, D., Heo, I. & O'Connor, R. Studying Cryptosporidium Infection in 3D Tissue-derived Human

Organoid Culture Systems by Microinjection. *J. Vis. Exp. JoVE* (2019) doi:10.3791/59610.

34. Hares, M. F. *et al.* Stem cell-derived enteroid cultures as a tool for dissecting host-parasite

interactions in the small intestinal epithelium. *Parasite Immunol.* **43**, e12765 (2021).

35. Fasciano, A. C., Blutt, S. E., Estes, M. K. & Mecsas, J. Induced Differentiation of M Cell-like Cells

in Human Stem Cell-derived Ileal Enteroid Monolayers. *JoVE J. Vis. Exp.* e59894 (2019)

doi:10.3791/59894.

36. Jelinsky, S. A. *et al.* Molecular and Functional Characterization of Human Intestinal Organoids and

Monolayers for Modeling Epithelial Barrier. *Inflamm. Bowel Dis.* **29**, 195–206 (2023).

37. Krndija, D. *et al.* Active cell migration is critical for steady-state epithelial turnover in the gut.

Science **365**, 705–710 (2019).

38. Park, J. *et al.* Promotion of Intestinal Epithelial Cell Turnover by Commensal Bacteria: Role of

Short-Chain Fatty Acids. *PLOS ONE* **11**, e0156334 (2016).

39. Cheng, H. & Leblond, C. P. Origin, differentiation and renewal of the four main epithelial cell types

in the mouse small intestine. V. Unitarian Theory of the origin of the four epithelial cell types. *Am. J.*

Anat. **141**, 537–561 (1974).

40. Darwich, A. S., Aslam, U., Ashcroft, D. M. & Rostami-Hodjegan, A. Meta-Analysis of the Turnover

of Intestinal Epithelia in Preclinical Animal Species and Humans. *Drug Metab. Dispos.* **42**, 2016–

2022 (2014).

41. Chen, X. *et al.* Cryptosporidium parvum activates nuclear factor κB in biliary epithelia preventing

epithelial cell apoptosis. *Gastroenterology* **120**, 1774–1783 (2001).

42. Carmen, J. C. & Sinai, A. P. Suicide prevention: disruption of apoptotic pathways by protozoan

parasites. *Mol. Microbiol.* **64**, 904–916 (2007).

43. McCole, D. F., Eckmann, L., Laurent, F. & Kagnoff, M. F. Intestinal Epithelial Cell Apoptosis

following Cryptosporidium parvum Infection. *Infect. Immun.* **68**, 1710–1713 (2000).

44. Heussler, V. T., Küenzi, P. & Rottenberg, S. Inhibition of apoptosis by intracellular protozoan

parasites. *Int. J. Parasitol.* **31**, 1166–1176 (2001).

45. Nishikawa, Y., Makala, L., Otsuka, H., Mikami, T. & Nagasawa, H. Mechanisms of apoptosis in

murine fibroblasts by two intracellular protozoan parasites, Toxoplasma gondii and Neospora

caninum. *Parasite Immunol.* **24**, 347–354 (2002).

46. Mele, R., Gomez Morales, M. A., Tosini, F. & Pozio, E. Cryptosporidium parvum at Different

Developmental Stages Modulates Host Cell Apoptosis In Vitro. *Infect. Immun.* **72**, 6061–6067

(2004).

47. Chen, X.-M., Gores, G. J., Paya, C. V. & LaRusso, N. F. Cryptosporidium parvum induces apoptosis

in biliary epithelia by a Fas/Fas ligand-dependent mechanism. *Am. J. Physiol.-Gastrointest. Liver*

Physiol. **277**, G599–G608 (1999).

48. Wallbank, B. A., Pardy, R. D., Brodsky, I. E., Hunter, C. A. & Striepen, B. Cryptosporidium impacts

epithelial turnover and is resistant to induced death of the host cell. *mBio* **0**, e01720-24 (2024).

49. Woods, K. M. & Upton, S. J. In vitro development of Cryptosporidium parvum in serum-free media.

Lett. Appl. Microbiol. **44**, 520–523 (2007).

50. Trinh, L. A. & Stainier, D. Y. R. Fibronectin Regulates Epithelial Organization during Myocardial

Migration in Zebrafish. *Dev. Cell* **6**, 371–382 (2004).

51. Increased basal fibronectin is sufficient to promote excess endothelial cell matrix assembly causing

localized barrier dysfunction | Molecular Biology of the Cell.

https://www.molbiolcell.org/doi/10.1091/mbc.E24-02-0090.

52. Arrowood, M. J. In Vitro Cultivation of Cryptosporidium Species. *Clin. Microbiol. Rev.* **15**, 390–400

(2002).

53. Miyoshi, H. & Stappenbeck, T. S. In vitro expansion and genetic modification of gastrointestinal

stem cells in spheroid culture. *Nat. Protoc.* **8**, 2471–2482 (2013).

54. Schreurs, R. R. C. E., Baumdick, M. E., Drewniak, A. & Bunders, M. J. In vitro co-culture of human

intestinal organoids and lamina propria-derived CD4+ T cells. *STAR Protoc.* **2**, 100519 (2021).

55. Read, E., Jowett, G. M., Coman, D. & Neves, J. F. Co-Culture of Murine Small Intestine Epithelial

Organoids with Innate Lymphoid Cells. *J. Vis. Exp. JoVE* (2022) doi:10.3791/63554.

56. Wagar, L. E. *et al.* Modeling human adaptive immune responses with tonsil organoids. *Nat. Med.* **27**,

125–135 (2021).

57. Cook, L. *et al.* Suppressive and Gut-Reparative Functions of Human Type 1 T Regulatory Cells.

Gastroenterology **157**, 1584–1598 (2019).

58. Rogoz, A., Reis, B. S., Karssemeijer, R. A. & Mucida, D. A 3-D enteroid-based model to study T-cell

and epithelial cell interaction. *J. Immunol. Methods* **421**, 89–95 (2015).

59. Noel, G. *et al.* A primary human macrophage-enteroid co-culture model to investigate mucosal gut

physiology and host-pathogen interactions. *Sci. Rep.* **7**, 45270 (2017).

60. Nozaki, K. *et al.* Co-culture with intestinal epithelial organoids allows efficient expansion and

motility analysis of intraepithelial lymphocytes. *J. Gastroenterol.* **51**, 206–213 (2016).

61. Pawlowic, M. C., Vinayak, S., Sateriale, A., Brooks, C. & Striepen, B. Generating and maintaining

transgenic Cryptosporidium parvum parasites. *Curr. Protoc. Microbiol.* **46**, 20B.2.1-20B.2.32 (2017).

 Figure Legends:

Figure 1: Organoid derived monolayers exhibit intestinal epithelium characteristics and support *C. parvum*

infection. (**A**) Schematic for organoid derived monolayer (ODM) differentiation and infection. Mouse intestinal stem cells

(mISC) were embedded and grown in Matrigel domes and maintained as spheroids in SCMM until expanded enough to seed

permeable supports. Prior to seeding, permeable supports were coated with a 1:10 Matrigel in SCMM and incubated overnight at

4°C. Matrigel dilution was removed and permeable supports were incubated at 37°C for at least 15 minutes prior to seeding with

mISC. Permeable supports were seeded with $3x10^5$ cells/well. Once confluent, the SCMM concentration was reduced to 5% for

24 hours, and then differentiation media (DM) which lacked L-WRN conditioned media entirely. ODMs were maintained in DM

for 2-3 days and then infected with *C. parvum*. (**B-E**) qRT-PCR of mRNA expression of ZO-1 for tight junctions (**B**), Muc2 for

goblet cells (**C**), ChgA for enteroendocrine cells (**D**), and lgr5 for stem cells (**E**). A Student's t-test was used to assess statistical

significance with a cut-off of 0.05. P-values are displayed on each graph. (**F-I**) Immunofluorescent staining of ZO-1 (**F**), Muc2

(**G**), Chromogranin A (**H**), and Lysozyme (**I**) in green and nuclei in blue confirming the presence of tight junctions, goblet cells,

enteroendocrine cells, and Paneth cells respectively in differentiated ODMs. Scale bars are 10 μm. (**J-K**) TEM confirmation of

tight junction (**J**) and microvilli (**K**) on ODMs. Scale bars are 200 nm. (**L**) Representative live epifluorescence images of ODM

infection with mNeonGreen parasites 3, 6, 9, 12, 15, and 20 dpi. Scale bar is 50 μm. (**M**) Vacuole counts from image J macro

analysis of fluorescent parasite vacuoles adjusted for total cell culture area. Lines represent infection with different infectious

doses (10^6 oocysts (blue), 10^5 oocysts (yellow) and 10^4 oocysts (magenta). Vertical grey lines show days of supernatant

collection (3, 8, 11, 15, 17, 22, and 25 dpi); images were taken after supernatants were collected. Standard deviation is shown as

shaded area surrounding each line. (**N**) Area under the curve (AUC) of experiment in M. (**O**) Flow cytometry quantification of

COWP1 + oocysts collected from ODM supernatant. (**P**) AUC of oocyst counts in O. One-way ANOVA (Tukey post-hoc

multiple comparisons) was used to assess statistical significance of AUC in N and P with a p-value cut-off of 0.05. P-values are

displayed on graphs. Error bars show standard deviation. n=2 for each condition.

Figure 2: Organoid derived monolayers support *C. parvum* production of infectious oocysts. (**A**) Graphical

experimental representation. tdTomato-nLuc-Neo oocysts were excysted and free sporozoites were purified by filtration through

a 1.2 μm pore syringe filter. Purified sporozoites were used to infect 24 wells (~33 mm² each) of typical HCT8 culture or ODMs

($1.5 x 10^5$ oocysts/well). Supernatants were collected every other day for 2 weeks, pooled and concentrated by centrifugation, and

stored at 4°C. IFNγ -/- mice were gavaged with supernatants from either HCT8 culture or ODM culture. Fecal samples were

collected on days 1-14, 19 and 24 post infection and fecal nanoluciferase was assessed. (**B**) Nanoluciferase assay of fecal samples

collected from mice gavaged with infected HCT8 or ODM supernatants. Thick lines represent the mean Relative Luminescence

Units (RLU) for each group and thin lines represent the range of RLU for each condition. Results are representative of 1

experiment, n=5 mice per group.

Figure 3: Interstrain mating confirmed by double positive zygote and oocyst observation. (**A**) Live image of

ODM infected with mNeonGreen (mNG) (left) and tdTomato (tdT) (middle) expressing *C. parvum* strains 8 dpi. Merge on the

right shows double positive vacuoles in white. Scale bar is 50 μm. Image was processed for 2D deconvolution using AutoQuant

X. (**B**) Oocyst collected from supernatant expressing both mNG and tdT. Scale bar is 5 μm.

Figure 4: Successful mating of two transgenic *C. parvum* strains in organoid derived monolayers. (**A-C**)

Quantification of parasitophorous vacuoles (PVs) from Image J macro adjusted for total cell culture area. Each graph shows

results from a different experiment. (**D-F**) Flow cytometry quantification of total oocysts collected from supernatant. Each graph

shows oocyst results from individual experiments in A-C respectively. Error bars show standard deviation. (**G**) Total COWP1+

oocysts from supernatant over the course of the experiment. Each line represents a different experiment. Standard deviation is

shown as shaded area surrounding each line. (**H**) AUC of COWP1+ oocysts for each experiment shown. One-way ANOVA

(Tukey post-hoc multiple comparisons) used to assess statistical significance. P-values are displayed on graph. P-values ≤ 0.05

were considered significant. Error bars show standard deviation. n=2 for each experiment. Data is representative of 2

experiments.

Figure 5: Fertilization Switch reporter. Sexual stage specific promoters drive the expression of Cre recombinase

fragments fused to FRB and FKB12 which will dimerize in the presence of rapamycin. Because they will be spatially separated

due to expression in different sexual stages, they cannot dimerize in the presence of rapamycin unless they are in the same cell

which occurs only after fertilization. LoxP sites flanking an mCherry sequence immediately upstream of an mNeonGreen

sequence will be excised in the presence of rapamycin and when the Cre fragments are in the same physical space in a zygote.

Figure 6: In vivo validation of the Fertilization Switch reporter demonstrates its inducibility and efficiency.

(**A**) Oocysts were purified from fecal samples collected from Fertilization Switch reporter infected NSG mice that were left

untreated or were treated with 10 mg/kg rapamycin daily for 4 days. Purified oocysts were used for microscopic analysis (**B**),

flow cytometry (**C-D**), and PCR analysis (**E**). (**B**) Top row: no rapamycin treatment; bottom row: with rapamycin treatment. Far

right column: zoomed in view of oocysts of merged images. mCherry in magenta, mNeonGreen in cyan. Scale bars are 5 μm. (**C**)

Flow cytometry plots of purified oocysts collected from mice with or without rapamycin treatment. (**D**) Percentages of COWP

positive oocysts expressing mCherry or mNeonGreen. (COWP+/mCherry or mNeonGreen+) (**E**) DNA gel of DNA collected

from oocysts purified from fecal samples from mice with (right lane) or without (left lane) rapamycin treatment.

Figure 7: Fertilization switch reporter infection of ODMs. (**A**) Representative images of fertilization switch reporter

infected ODMs with (bottom row) or without (top row) rapamycin treatment. (**B**) Percentage of vacuoles of large tiled images

counted as mCherry or mNeonGreen positive. Error bars represent SD. (**C**) Higher magnification of infected area showing a

mixed population of mCherry and mNeonGreen vacuoles. In merged images, mNeonGreen fluorescence is shown in cyan and

mCherry fluorescence is magenta. Data are representative of 2 experiments. Scale bars are 5 µm.

Figure 8: Evidence of autoinfection in ODMs. (**A**) Infected area with a mixed population of mNeonGreen and mCherry

positive vacuoles. Scale bar is 5 µm. (**B**) Z-stack slice showing trophozoites. (**C**) Larger mCherry positive vacuole with only 1

nucleus, likely an unfertilized macrogamont. (**D-E**) Z-stack slice showing multiple nuclei in a larger mNeonGreen positive

vacuole, likely a meront. Scale bars for B-E are 1 µm.

Figure 9: Fertilization switch reporter infection in HCT8 cells. Fertilization switch infected HCT8 cells in

the absence (**A**) or presence (**B**) of rapamycin treatment at 88 hpi. (**B**) Representative mNeonGreen

positive cluster of parasite vacuoles. Scale bars are 5 µm. (**C**) Number of clusters of mNeonGreen

positive parasite vacuoles counted at 88 hpi. Each data point indicates total mNG+ vacuole clusters

present on a 33 mm^2 culture well with data accrued in two independent experiments.

A

| Hoescht | mCherry | mNeonGreen | Merge |

-RAPA

+RAPA

B

% PV by color

-RAPA   +RAPA

■ mCherry   ■ mNG

C

| mCherry | mNeonGreen | Merge |

-RAPA

+RAPA

A

B

C

D

E

mNG   mCh   Hoescht   Merge