Deep learning reveals the complex genetic architecture of a highly polymorphic sexual trait

Wouter van der Bijl[1*], Jacelyn J. Shu[1], Versara S. Goberdhan[1], Linley M. Sherin[1], Changfu Jia[1], María Cortázar-Chinarro[1,2], Alberto Corral-López[1,2] & Judith E. Mank[1]

1. Department of Zoology & Biodiversity Research Centre, University of British Columbia, Vancouver, Canada

2. Division of Ecology and Genetics, Uppsala University, Uppsala, Sweden

* Corresponding author

Abstract

The extraordinary variation in male guppy coloration has proven a powerful model for studying the interplay of natural and sexual selection. However, this variation has hampered the high-resolution characterization and determination of the genetic architecture underlying male guppy color, as well as clouded our understanding of how this exceptional level of diversity is maintained. Here we identify the heritability and genetic basis of male color variation using convolutional neural networks for high-resolution phenotyping coupled with selection experiments, controlled pedigrees and whole-genome resequencing for a Genome Wide Association Study (GWAS) of color. Our phenotypic and genomic results converge to show that color patterning in guppies is a combination of many heritable features, each with a largely independent genetic architecture spanning the entire genome. Autosomally-inherited ornaments are polygenic, with significant contributions from loci involved in neural crest cell migration. Unusually, our GWAS results suggest that gene duplicates from the autosomes to the Y chromosome are responsible for much of the sex-linked variation in color in guppies, providing a potential mechanism for the maintenance of variation of this classic model trait.

Introduction

The study of color polymorphisms has been pivotal to our understanding of how variation is maintained

in populations. Perhaps none are as polymorphic as the color patterns of male guppies, *Poecilia*

reticulata, which have been extensively studied in the context of fitness[1–4] and female preference[5,6]. The

widespread variation in the extent and patterning of male color across wild populations[4,7,8] and the rapid

response to natural and artificial selection[9,10], coupled with high heritability estimates[11,12] all suggest

that substantial genetic variation for this trait is maintained in guppy populations. These colors are not

only unusually variable for sexual ornaments but have some of the highest coefficients of genetic

variation ever reported for morphological traits[11,13]. Although female mate preference for rare male

patterns likely acts to preserve some variation from the eroding forces of genetic drift and

selection[5,13,14], the genetic architecture underlying extraordinary diversity in male guppy color remains

mysterious, as does the maintenance of this exceptional diversity.

Part of the difficulty in determining the genetic basis of male guppy coloration lies in the multiple forms

of variation. Yellow-orange xanthophores and black melanophores form spots that vary among males in

their location, as well as in the extent and saturation of these colors. Color pattern variation arises from

embryonic neural crest cells that form the precursors of xanthophores and melanophores[15]. As the

administration of testosterone can induce color traits in females, coloration appears to be under

hormonal control[8,16]. Therefore, the genetic basis of color variation is expected to include loci involved

in the migration of neural crest cells, as well as genes involved in the synthesis and deposition of

pigments, and possibly the production of and response to testosterone. Importantly, orange carotenoids

are not synthesized *de novo* in vertebrates, but are collected from food sources[17] and modified prior to

deposition, and so the genetic basis of orange coloration could also include loci involved in the

bioavailability of these molecules.

It has often been suggested that, despite the striking diversity of guppy color patterns, their genetic

basis might be surprisingly simple. However, the excessive complexity of guppy pattern phenotypes has

significantly complicated the identification of the underlying genetic loci. Early studies, which assessed

pattern inheritance by visual inspection, found a mix of Y-linked, X-linked, and autosomal inheritance for

specific color traits[8,12,18–21]. More recently, a study suggested that the complete color patterns in one

population are inherited as a single Y-linked locus with twenty-seven alleles[5]. These findings indicate

that at least some ornaments are monogenic and co-inherited due to linkage, but they leave unresolved

how ornaments with more complex inheritance patterns are controlled. More quantitative and

objective summary pattern measures, such as the total orange or black area, suggest a mix of

inheritance patterns[11,12,16,22]. Here we overcome these longstanding obstacles by developing a pipeline

of convolutional neural networks for high-resolution phenotyping of 3,229 fish coupled with selection

experiments and controlled pedigrees to objectively determine the heritability of the full complexity of

color patterns. We then used our phenotype data for a Genome Wide Association Study (GWAS) of color

to identify the genetic architecture of a suite of color traits.

Our phenotypic and genomic results converge to show that color patterning in guppies is a combination

of many heritable features, each with largely independent genetic architectures. We delineate fifteen

separate ornaments with heterogeneous genetic architectures, with some ornaments inherited as a

single sex-linked locus, while others have a polygenic autosomal basis. Finally, our results suggest that

gene duplications from the autosomes to the sex chromosomes are responsible for much of the sex-

linked variation in color in guppies, providing a potential mechanism for the maintenance of variation of

this classic model trait.

Artificial selection on orange coloration

Starting from an outbred laboratory population with large variation in color patterning, we created three replicate pairs of selection lines for reduced and increased amounts of orange coloration as a percent of total body area. Each male in the experiment (n = 3,229) had multiple photos taken from each side to track measurement error (n = 14,100). Using a series of neural networks, we performed image segmentation to separate the fish from the background, and to identify the orange and black coloration on the fish (Fig 1A). In addition to orange, xanthophores also create the less frequent yellow color patches, which we include in our orange measurements. We then warped images to a common reference shape based on placed landmarks to enable fine-scale color mapping[12]. Combined with a full pedigree, this dataset allowed us to trace the inheritance of color patterns across four generations (Fig 1B). While some body areas are almost always pigmented (Fig 1C), color is rare in most places on the male body plan, giving each male a distinct pattern.

Figure 1: Quantification of pattern variation. A) Overview of the phenotyping pipeline. From each photo, the fish is extracted, the body is aligned to a common reference shape using landmarks, and the position of orange and black color is extracted. B) Randomly selected examples of color inheritance along patrilines. Images on the same row belong to the same patriline, and columns are successive generations of the selection experiment (both selection directions). C) The incidence of color across the guppy body among all males. Body positions where the incidence of color is less than 1% are colored grey.

The percent of orange color responded rapidly and strongly to selection (Fig 2A). After just three

generations, up-selected males had more than double the amount of orange color compared to their

down-selected counterparts (% up – % down in F_3 [95% credible interval]: 6.05 [5.67, 6.48]). We

modeled the inheritance of black and orange color along the pedigree[23], separating autosomal, X-linked

and Y-linked genetic effects, as well as influences due to the maternal and tank environments. The

strong response in orange color to selection is consistent with its substantial heritability (narrow-sense

h^2: 0.59, [0.50, 0.68], Fig S1). While orange and black ornaments share the same body plan, selection on

orange nonetheless had a negligible effect on the amount of black coloration (up – down in F_3: -0.09 [-

0.17, 0.01], Fig S2A). Although the amount of black color is also substantially heritable (h^2: 0.67 [0.53,

0.80], Fig S3), the amount of the two colors is inherited largely independently as evidenced by their low

genetic correlations ($r_{autosomal}$: 0.07 [-0.15, 0.26], Fig S4). Although previous work on other

populations[11,24–26] has identified strong Y-linkage, our estimates of Y effects are low ($h^2_{Y-linked}$ orange:

0.03 [0.00, 0.09], black: 0.04 [0.00, 0.08]), and instead the genetic variation for the proportion of orange

and black color on the body has an autosomal and, to a lesser extent, X-linked origin (Figs S1&S2).

Our selection regime affected the incidence of orange color at nearly all positions on the body, with

effect sizes increasing with each successive generation of selection (Fig 2B). The strength and direction

of the effects were spatially heterogeneous. In the F_3 generation, up-selected males were approximately

5x more likely to have orange color at any position in the tail or caudal fin (posterior to the

gonopodium). The strongest effects were located close to the pectoral fin and gills, where up-selected

males were more than 100x as likely to have orange color on the ventral side, while surprisingly, they

were more than 100x *less* likely to be orange dorsally (although this dorsal color was much less

prevalent overall, Fig 1C). The response against the direction of selection at some locations, and

correlated local changes in black color (Fig S2B), illustrate that some elements of the color pattern are

genetically linked.

To investigate whether other traits were co-selected in our selection lines, we looked at changes in

behavior and life history, since selection on embryonic neural crest cells has resulted in changes in these

traits in other vertebrate systems[27,28]. Guppy males flaunt their coloration during characteristic sigmoid

display behavior, where they curve their body into an S-shape, quiver, and swim laterally in front of

females[29,30]. Using a subset of 126 male offspring from the F3 generation (63 from each selection

direction, evenly split across the replicate lines), we evaluated the effects of our selection regime on

male sexual behavior. In a standard no-choice test with a non-receptive female, males from the up-

selected lines performed both more (short displays: ratio up/down = 1.74, z = 2.15, p = 0.031, long

displays: up/down = 1.84, z = 2.28, p = 0.022) and longer (up/down = 1.9, z = 2.44, p = 0.015) displays to

Figure 2: Orange color responds rapidly to selection. A) Density plots show the distribution of the percent of orange
coloration per generation and selection regime. Thin lines between generations connect fathers to their sons. Thick lines
inside the densities show the median value of color area. B) Heatmaps illustrating the effect of selection across the body.
Each heatmap cell is colored by the log odds ratio (as estimated by a generalized linear mixed model), illustrating the
relative odds that a male has orange color at that location. Body positions where the incidence of orange color is less than
1% are colored grey. C) Selection moves populations through orange patternspace. Axes show orange patternspace (after
UMAP dimension reduction), with points indicating individuals, colored by selection direction, and facets showing the four
generations.

the female (Fig S5). In addition, they spent more time following the female (up/down = 1.9, z = 2.44, p =

0.015, Fig S6), and performed more attempts at coercive sneak copulations (up/down = 1.99, z = 2. 09, p

= 0.037), suggesting the more colorful males were 1.7-2x more sexually active overall (Fig S6). Males did

not show changes in body size or tail size (standard length in mm, up – down in F_3: -0.36 [-0.84, 0.12],

tail length: 0.10 [-0.21, 0.43], tail area mm^2: 0.019 [-2.5, 2.2], Fig S7), nor did selection alter the fecundity

of females (F_2 offspring per brood, down-selected: 7.74, up-selected: 7.79, ratio up/down: 0.99 [0.87,

1.12]) or the time between pairing and the first brood (F_2, up - down in days: -1.23 [-3.4, 0.97], Fig S8).

Localized heritability, ornaments, and patternspace

While the quantity of orange and black coloration are important to female choice[7,31], male patterning is

largely inherited as discrete spots and stripes at fixed body positions[12,24–26]. To understand the spatial

structure of color inheritance, we modeled the heritability of the presence of color separately for each

body location (Fig 3), confirming the fine-scale heritability of color patterning. In many places the

Figure 3: The localized heritability of color is high, but the architecture is heterogeneous. The narrow-sense heritability of the presence of orange (A) and black (B) color on the guppy body. For each heatmap cell, a Bayesian animal model was used to estimate the contribution of additive genetic and environmental factors. Heritabilities are expressed as a proportion of the between-individual phenotypic variance. The autosomal, X-linked, and Y-linked heritability add up to the total heritability. Body positions where the incidence of color is less than 1% are colored grey.

localized heritability is high (> 0.8), much higher than the heritability of the total amount of color.

Decomposing the genetic variance into autosomal and sex-linked components shows that the genetic

architecture is highly spatially heterogeneous. For example, orange color in the top of the tail fin and

black color ventrally of the dorsal fin were strongly X-linked, while both colors near the pectoral fin

showed a large Y-linked component. This overlapping patchwork of genetic architecture generates the

complexity of the color pattern.

We defined seven orange (Fig 4) and eight black ornaments (Fig S9), based on the incidence of color (Fig

1C) and the localized genetic architecture (Fig 3). As expected, the incidence of all ornaments is

heritable. In addition, their size is also heritable for all but one ornament (O3). All orange ornaments

responded to selection on the total amount of orange, either in their incidence, their size or both (Fig 4C

& 4D). The large effect sizes for ornament incidence are the result of high heritability and strong

selection and are consistent with the expectation arising from the breeder's equation (Fig S10).

Concordant with the localized analysis of inheritance, we find that the ornaments each have their own

characteristic combination of autosomal, sex-linked and environmental causes. This can also be seen

from the low correlations between many different ornaments with autosomal or X-linked architectures

(Figs S11 & S12). In contrast, we observed strong Y-linked genetic correlations, consistent with alleles

across multiple loci linked together on a non-recombining Y chromosome. Within each ornament,

however, the architecture of incidence and size were typically similar, which may indicate that additive

genetic factors for ornament size also control its presence as a threshold trait.

In principle, the seven orange and eight black ornaments can combine into 32,768 (2^{15}) unique pattern

combinations. We observed 691 of these unique patterns among the 3,229 males in our dataset, with

just 130 males (4%) sharing the most common pattern (O2, O4, O5, O6, & B7, Fig S13). Such pattern

diversity is consistent with previous qualitative descriptions[8], although much greater than the only

previously reported count of twenty-seven patterns in a different population[5]. We observed fewer

Figure 4: Orange ornaments are strongly heritable and their incidence and size responded to selection. A) Pictograms of seven orange ornaments. B) Heritabilities of the incidence and size (when present) of each ornament. Dots and lines reflect point estimates and 95% credible intervals, and the gradient bars show the cumulative posterior density. C&D) Effect of selection on the incidence and size of orange ornaments. Bars and points represent means, with error bars showing 95% bootstrapped confidence intervals. X-axes show consecutive generations.

patterns than expected from the prevalence of each ornament (1,010), suggesting that linkage

disequilibrium between ornaments caused by physical linkage and correlated selection reduced pattern

diversity by 32% (confidence interval: [29%, 34%], Fig S14).

These localized approaches will underestimate the true pattern diversity in the population if separate

color elements are co-located. For example, some males exhibit large patches of color in the tail that do

not neatly fit the three ornaments defined there (Fig S15). Similarly, they do not capture potential

heritable differences in spot shape or color characteristics such as hue and saturation. To describe the

orange and black patterns free from assumptions about developmental mechanism, we developed a

novel high-dimensional phenomics approach using a deep neural network embedder which places the

patterns within five-dimensional patternspace. Briefly, using triplet learning[32], these convolutional

neural nets were trained to encode the heritable pattern variation of orange and black patterns (see

Supplemental Methods). The networks are only given color patterns but are rewarded when they place

patterns from related males closer together than those of unrelated males. As a result, patternspace is a

holistic summary of the complex pattern variation among males (Fig S16). Patterns which are close in

patternspace share heritable features, as can be observed when we map the presence and size of the

ornaments onto this space (Fig S17).

Orange patternspace is divided into two large clusters, differentiated by the presence of the highly

heritable and Y-linked O6 ornament (Fig 5 & S17). However, because the Y does not recombine,

complete Y-linkage of all color patterns would result in discrete patterns associated with each patriline,

which would be expected to form additional clusters, which we did not observe. Instead, the many

heritable features combine into a continuous space of pattern variation consistent with high levels of

autosomal and X-linked variation across the recombining portion of the genome. Our selection regime

gradually separated the populations in patternspace over the generations (Fig 2C). Nonetheless, as there

are many routes to increase or reduce the amount of orange, pattern variation is largely maintained

within the selection lines among succeeding generations.

Mapping color pattern to the genome

We selected 300 males from the F_3 generation for short-read whole genome resequencing, sampling

across all families and across patternspace to maximize the variety of patterns captured. The guppy Y-

chromosome is highly diverse, both within[33] and among populations[34], and using a male reference

genome with a single Y haplotype creates major potential reference bias for all other Y haplotypes (see

Supplemental Methods). However, the high degree of similarity between the X and Y for homologous

regions[33,35,36] ensures that the vast majority of Y-linked reads will map to the X assembly. We therefore

aligned our reads to the guppy female reference genome[37], which avoids reference bias to any particular

Y haplotype, but confirmed our conclusions using a male reference[38] (Fig S18). After variant calling and

filtering, we performed a GWAS for the presence/absence of common orange and black ornaments, and

multivariate GWAS for orange and black patternspace. As some of our phenotypes showed sex-linked

inheritance, we controlled our GWAS for relatedness across the autosomes as well as the sex-

chromosomes (see *Methods*).

The GWAS for ornaments with substantial autosomal inheritance revealed genetic architectures

consistent with complex traits (Figs 5, S19 & S20). We recovered multiple peaks on autosomes for each

of these ornaments (Fig S21), each explaining a modest proportion of variation in the phenotype (logistic

regression, Nagelkerke's R^2: <16%), suggesting an oligogenic or polygenic architecture, consistent with

threshold trait. To investigate the specificity of these loci, we performed PheWAS and related the

genotype at the loci to pigmentation across the body. The loci showed specific local effects for its

associated ornament, supporting the conclusion that these ornaments have an independent genetic

basis (Fig S21).

Figure 5: **The genetic architecture of color pattern is a mix of autosomal loci and sex-linked loci**. A) Manhattan plot showing the association of genomic variants with ornament O2. Points represent genomic variants, with their genomic location on the x-axis, and the p-value of the association on the y-axis. Dotted line denotes 5% FDR. Sex-linked loci are placed to the side, with both the sex-chromosome (LG12) and sex-linked loci cross-mapping to the autosomes shown, colored by their putative origin (red = X, blue = Y). Peaks are labelled with gene names, but unnamed and uncharacterized genes are shown. Names with asterisk (*) indicate that more significant cross-mapping SNPs were present in the same peak. B C, D) As in A, but for ornament B1, orange patternspace, and black patternspace respectively. Results for other ornaments in Figs S19 & S20.

Genes with significant variants include those related to cell-to-cell adhesion, which plays a crucial role in

the migration of neural crest cells[39] (NCC), which are the precursors of xanthophores and

melanophores[15]. Significant variants include cadherin genes *fat1a, cdh4* and *cdh12* which encode

adhesion molecules. Expression of *cdh12* has been implicated in variation in human skin pigmentation[40],

and other cadherins have been associated with population-level differences in guppy coloration[41]. In

addition, a variant close to *sema6a* explained 16% of the variation in the tail ornament B1. This gene is a

target of the melanocyte master regulator *mitf* in humans[42], is upregulated during the growth of

pigmented tail fins in betta fish[43], and semaphorins have been broadly implicated in guiding NCC

migration in zebrafish[44]. Similarly, *sema4f* is associated with ornament B6. Chromatin remodeler *bptf* is

necessary for *mitf* function[45], and is also implicated in ornament B1. We also find signal upstream of

mitfa itself for ornament O2, supporting a recent study showing *mitf* also affecting the fate of

xanthophores in medaka and zebrafish[46]. A variant associated with B6 falls in *lin28a*, a key regulator of

position-dependent NCC differentiation[47]. *Cygb1* affects NCC development and melanophore patterning

in zebrafish[48], and *fscn1a* is involved in directional NCC migration[49]. Together, these results suggest that

many of the ornaments are ultimately under the control of loci modifying the trajectory of NCCs, as they

migrate towards the skin and develop into chromatophores.

Next, we investigated the genetic basis of color traits with X- and Y-linked patterns of inheritance. For

orange and black patternspace, highly significantly-associated variants span the entirety of linkage group

(LG) 12, the sex chromosomes, largely due to strong linkage across this region. Given the low

differentiation between the X and Y for homologous regions[33] (male:female coverage at LG12: 21-26 Mb

= 0.986), most reads from the Y chromosome map to the X chromosome in the female reference

genome, and variants on LG12 represent a combination of pseudo-autosomal, X-linked and Y-linked

variants[35]. Ornament level GWAS (Figs S19 & S20) confirmed that ornaments with strongly sex-linked

inheritance are associated most with LG12.

Surprisingly, we observed additional peaks of associated variants on nearly all autosomes for ornaments

that our heritability analysis showed to be sex-linked. Strikingly, many of these regions showed excess

heterozygosity, elevated read depth, and an increased read depth from heterozygotes compared to

homozygotes, suggesting reads from copy number variants (CNVs, Fig S22). Given recent work that

identified CNVs spanning autosomes and the Y chromosome in natural guppy populations[50] and the role

of sex-linked variation in this system, we hypothesized that reads from the Y chromosome regions that

are not homologous to the X could map to these autosomal regions, representing CNVs spanning

multiple genomic locations. We tested this conjecture by comparing the genotypes at each variant with

expectations from autosomal, X- or Y-linked segregation. Briefly, we modelled the dose of the variant as

following the pedigree-based covariance between individuals in each of the three scenarios. We then

called the putative source of these variants as X- or Y-linked if their relative likelihood (AIC weight)

exceeded 80% (Fig6). Using this pedigree-informed approach, we identified that 1% of called variants

were potentially the result of cross-mapping reads between the autosome and sex-chromosome.

Variants implicated in the patternspace GWAS were highly enriched for these cross-mapping variants

(orange pattern: 19 X-linked & 154 Y-linked out of 510 called variants; black pattern: 46 & 252 out of

563; Figs S23 & S24), also suggesting the involvement of CNVs with sex-linked copies.

One possible explanation for the abundance of Y-linked variants in our GWAS is that they are all strongly

linked to one or a few causal loci of large effect which control the Y-linked components of the color

traits. To test this, we performed principal component analysis on the variants associated with orange

pattern and a Y-linked inheritance pattern. If these loci have significant associations with orange pattern

due to strong linkage to the same few alleles, we expect these variants to cluster into clear haplotypes

that are strongly associated to color. Instead, we identify four putative Y-haplogroups with substantial

variation within each haplogroup (Fig S27), and which only explain a small amount of color variation

compared to individual variants (Figs 6C & S28). The fully non-recombining region of the Y chromosome

Figure 6: **Large effect sex-linked CNVs underlie sex-linked variation in color pattern**. A) GWAS peaks on autosomes with significant loci with sex-linked inheritance, near *texim* and in *prex1*. B) Relative coverage (1x = genome wide sample average) across the region for 297 males (this study) and 3 females, calculated in moving windows of 1kb, 100bp apart. C) Phenotypic effects of the top variant (diamond in A) on ornaments O6 (orange pattern) and B5 & B7 (black pattern). The proportion of variation in the phenotype that is explained by the top SNP is displayed as Nagelkerke's R^2 from a logistic regression. D) Median estimated allele counts for the top SNP, based on read depth, for homozygous reference and heterozygous individuals. No heterozygous alternate individuals were found. E) PheWAS heatmaps displaying Z-scores for the effect of the top variant on orange and black color across the body, controlling for autosomal and sex-linked relatedness.

is relatively small[36], and the majority of the chromosome recombines with the X exceedingly rarely[51].

Although these rare X-Y recombination events might explain the variation without our four Y-

haplogroups, this would also lead to expected X and Y segregation of SNPs. We therefore suggest that

our data are more consistent with a substantial amount of Y diversity in our population.

The locus most strongly associated with orange pattern mapped to LG14 (an autosome), with many

significant variants in a 35kb region that are inherited in a Y-linked manner (Fig 6A, left column),

suggesting these variants represent fixed differences between the autosomal region and its copies on

the Y-chromosome. Consistent with this scenario, we observed highly elevated coverage consistent with

multiple Y-linked copies in males, and reduced coverage in females (Fig 6B).

The gene located in this region (LOC103476393) is uncharacterized in the annotation but has 72%

sequence identity with *teximYa* from the related Southern platyfish (*Xiphophorus maculatus*)[52]. Y-

specific duplication of *texim* has been documented in multiple guppy populations[38], and copies of *texim*

have also been found on the Y-chromosome of the closely related *Poecilia parae*[53], suggesting

duplication of *texim* to the Y-chromosome is common in this clade, and most likely mediated by

helitrons[38,52]. In fact, the most strongly associated variant in this region is located downstream of this

texim copy, and falls in the *texim*-associated helitron. This top SNP predicts 92% of the phenotypic

variation of the Y-linked ornament O6 (Fig 6C & 6E). Coverage for most of this region is elevated five to

ten-fold and heterozygous individuals have increased coverage compared to individuals that are

homozygous reference (Fig 6D), so that coverage in this region is strongly predictive of the presence of

ornament O6 (Fig S29). Using long-read RNA sequencing, we confirmed the existence of at least five

texim copies whose presence is polymorphic on the Y-chromosome. We also confirmed that the helitron

is transcribed together with the *texim* gene. One such copy is expressed in adult tail tissue, but only in

males, again consistent with Y-linkage (male vs. female: 2.14 vs 0.00 CPM; t_4 = 3.8, *p* = 0.019; Fig S30).

Genotyping the males in our GWAS for the presence of this copy revealed that it is predictive of the

presence O6 (R^2 = 0.30, Fig S29), but can only partially explain the observed association of the top GWAS

SNP (R^2 = 0.92).

Similarly, the locus most strongly associated with black pattern mapped to LG7, also an autosome (Fig 6,

right column). We observed many highly associated sex-linked variants in a 3.5kb region, again paired

with an increased read-depth for males, but not females. This locus all shows non-Mendelian

inheritance with increased read depth for heterozygotes and a lack of alternate homozygous genotypes,

all supporting the notion that these variants are the result of cross-mapping from the sex-chromosomes.

The inheritance pattern of the significant SNPs indicate that the presence or absence of an X-linked copy

is most strongly associated with the phenotype, although increased coverage in males compared to

females suggests the presence of additional Y-linked copies. All significant variants fall into the gene

PIP$_3$-dependent Rac-exchanger 1 (*prex1*). This gene has previously been found to be important to

melanoblast migration in mice, where disruption of this gene leads to white paws and belly, the most

distal endpoints of NCC migration[54]. This locus alone explains 46% and 50% of the phenotypic variation

in ornaments B5 and B7, but in opposite directions. Since these ornaments are in close vicinity, this may

suggest that genetic variation in *prex1* modifies NCC trajectories to terminate in one or the other

location. Overall, the sex-linked ornaments appear to have a monogenic or possibly oligogenic

architecture, with evidence suggesting a large role of copy number variants that span the autosomes

and sex chromosomes.

Discussion

The remarkable variation in male guppy coloration has made them a model for studies of female

preference[5,6] and the interaction of sexual and natural selection[4,55]. This variation has long been

suggested to have a simple genetic basis, but the large variation in phenotypes has precluded analysis of

the full complexity of the pattern and identification of responsible genomic loci. Here we used a

combination of high-resolution phenotyping via neural nets, controlled pedigrees, and GWAS to dissect

the detailed genetic architecture for male color variation.

Our population has shown a rapid response to artificial selection for other complex traits[56–58], suggesting

significant underlying genetic variation. Consistent with this, we observe substantial variation in male

color (Fig 1) and a rapid response to artificial selection for increased and decreased orange coloration in

all our replicate selection lines (Figs 2, 4). Male color has previously been shown to respond to artificial

selection in other populations[10,59,60], and consistent with this, we observe high heritability (Figs 3 and 4)

for both total color area and specific color ornaments. Nonetheless, we see that color is not entirely

genetically determined, and there is a significant role of the environment for the size of ornaments and

locations in the tail fin. The role of color as an honest signal of a male's ability to e.g. forage and fend off

parasites[1,2] may either be mediated through these less-heritable components of the color phenotype, or

color loci may be genetically linked to loci conferring such fitness benefits.

We confirmed that sex-linked variation plays a substantial role in the genetic architecture of color in this

population, although for only a subset of ornaments (Fig 3). Previous reports on other populations based

on the visual assessment of select co-inherited ornaments have often emphasized their Y-linkage,

although there are also many which have been observed segregating on the X or autosomes[5,21,24–26,61]. It

is therefore unclear how much of the total color variation these classic patterns capture. Quantitative

analyses based on color area and spot number have also shown a larger contribution of the sire

genotype, but dam heritability nonetheless plays a significant role[11,62]. Finally, testosterone-treated

females will generally express some ornaments, demonstrating that complete Y-linkage is the

exception[16,26]. As both the observable ornaments and their sex-linkage varies considerably among

populations, we look forward to population comparisons using the comprehensive analyses now

available.

One might expect that if black is epistatic to orange, selection to increase the area of the fish with

orange coloration would reduce the total black area. That is not the case, as we observe little change in

black area throughout the selection experiment but do see modest localized shifts in black color (Fig S3),

more consistent with the repulsion of developing xanthophores and melanophores during color

development[63].

We then used whole genome sequencing for 300 males from the F_3 generation, sampled to maximize

their distribution within color-space to perform GWAS for multiple color traits (Figs 5 and 6).

Interestingly, genes previously identified as key to melanophore formation in guppies, *colony-*

stimulating factor 1 receptor a (also called *fms*), *kit ligand a* (*kita*)[27,64] and *adenylate cyclase 5* (*adcy5*)[65],

were not implicated in our GWAS (Fig S31). This suggests that although these loci are fundamental and

necessary in the development of color cells, they do not contribute substantially to the variation in color

pattern we observe in our population. Instead, we find significant associations with genes involved in

cell-cell adhesion and neural crest cell migration, suggesting that the migration routes of

chromatophores and their pre-cursors during embryonic development, rather than the production of

pigments, determines the black pattern. Genetic variation in one such gene, *prex1*, explains half the

phenotypic variation in two black ornaments.

In contrast to melanin, carotenoids are not synthesized in vertebrates *de novo*, but rather collected from

dietary intake and modified prior to deposition[66]. This mode of color development is hypothesized to

make carotenoid-based coloration a key mediator of honest sexual signaling of male quality[1,67,68].

Consistent with this, males from selection lines with greater orange area exhibited greater sexual vigor

(Fig S4 & S5), though we did not observe any correlated changes in female life history traits. The

behavioral response to selection on color may be the result of pleiotropic changes to neural crest cell

development, which plays a central role in both pigmentation and neurological development[27,28].

We might expect genes associated with carotenoid metabolism in other vertebrates to play a role in

orange coloration in guppies. A key gene identified in red coloration of birds, cytochrome P450 locus

CYP2J19[69,70], lacks a reciprocal ortholog in teleost fish. Aldehyde dehydrogenase family 1 member A1,

associated with red carotenoid coloration in lizards[71], is also not implicated in our GWAS. Instead, the

region most strongly associated with orange color is a helitron that is closely linked to and co-expressed

with *texim*. Although the functional role of *texim* remains unclear, it is thought to encode an

esterase/lipase[52], important in carotenoid modification[72,73] and therefore may play a role in orange

coloration.

Similarly, we do not find support for a large haplotype on LG1, which has been previously been indirectly

implicated in guppy color variation in another population[74]. Instead, our results suggest a far more

complicated genetic architecture for color pattern. For both black and orange color, we observe specific

ornaments encoded through largely independent suites of loci (Figs S23 & S24). Most ornaments have a

polygenic and largely autosomal architecture. Moreover, the sex-linked architecture is heavily

influenced by CNVs, although it is challenging to differentiate between gene dose effects, with more

copies of a gene associated with greater color, and allelic variation, with some copies associated with

specific patterns. CNV dose has also been implicated in rapidly evolving color phenotypes during

domestication. In rock pigeons, at least seven alleles that vary in the copy number of the Z-linked *st*

locus produce different depigmented phenotypes[75], while additional copies of a CNV near the melanin-

associated *kitlg* gene causes darker coats both within and across dog breeds[76]. This aspect of the genetic

architecture of guppy color is particularly interesting in that it might be expected to help maintain male

phenotypic color variation more effectively than allelic variation in single-copy genes. It has been a long-

standing mystery how the remarkable variation in male color is maintained against the eroding forces of

genetic drift, selection and sweeps. Although female mate preferences for rare male patterns can be

expected to preserve some male patterning variation[5,13,14], it is difficult to envisage how this would be

sufficient to maintain the extensive variation observed in guppy populations, many of which formed

with significant bottlenecks and founder effects. Moreover, the evolution of female preference for rare

patterns has been proposed to be beneficial to females through the indirect benefit of sexy sons[5,77].

However, such benefits rely on a strong genetic covariance between the pattern of fathers and their

sons. It is unlikely that populations like the one studied here, with its many autosomal and X-linked

inherited pattern elements, would confer such benefits to females.

The fact that we observe CNVs associated with color spanning autosomes and the sex chromosome may

help to explain recent observations of variation in the extent of Y-linkage for color in wild

populations[16,22]. Although the size of the fully non-recombining region of the Y chromosome is relatively

small[34–36], there is evidence that it can accumulate substantial recent duplications of genes from the

remainder of the genome[50]. If this process involves coloration loci and varies across populations, it is

easy to envisage how the association between the Y chromosome and color variation could differ, even

among closely related populations. It may also suggest that the comparatively low-levels of Y-linkage we

observe in our population, originally collected from the high predation region of the Quare River,

Trinidad, may be observed in other natural populations that either have lower rates of duplication to the

Y, or lower levels of Y-duplicate retention. Finally, our data suggests that the Y-linked regions that do

contribute to color in our population are surprisingly diverse.

Taken together, our results present a critical link between the extraordinary variation in male guppy

coloration and the genome that encodes it. The fine-scale genetic architecture of color that we reveal

presents an exciting opportunity for future molecular analyses of sexual selection dynamics and a critical

point of comparison for future studies of the genetic architecture of highly polymorphic traits.

Acknowledgements

We thank Lengxob Yong for sharing insights on guppy photography. We thank Ben Sandkam, Ben

Furman, David Metzger, Lydia Fong, Yuying Lin, Christina Hodson, and James Lewis for help with animal

husbandry and valuable discussions, and Tom Booker for advice and comments on an earlier draft.

Yuying Lin contributed the female coverage data. Funding was provided by a Canada 150 Research Chair,

NSERC and an ERC grant (680951) to JEM.

Author contributions

Conceptualization: WvdB, AC-L & JEM, Methodology: WvdB, Formal analysis: WvdB & AC-L,

Investigation: WvdB, JJS, VSG, LMS, CJ, MC-C, AC-L & JEM, Writing – Original Draft: WvdB & JEM, Writing

– Review & Editing: WvdB, JJS, VSG, LMS, CJ, MC-C, AC-L & JEM, Visualization: WvdB, Supervision: JEM,

Project administration: JJS, Funding acquisition: JEM.

Data and code availability

Code is available at https://github.com/Ax3man/vdBijl_etal_2024_GuppyColorPatterns. Sequence data

will be deposited upon publication.

Materials and methods

Study population and animal husbandry

Our study population are descendants of guppies that were captured in Trinidad in 1998 and

subsequently maintained in large, outbreeding stock populations to maintain genetic diversity, and

males show large diversity in their ornamentation. 240 females for the parental generation (P) were

randomly sampled as juveniles from our three stock populations and maintained as virgins until sexual

maturity. 300 males for the parental generation were sampled as adults from the stock populations.

Fish were kept in 1.4 l tanks in three flow-through systems, with each experimental replicate in a

separate system. Water was maintained at 25 °C, and lights were on a 12:12h cycle. Fish were fed daily

with a mix of live brine shrimp and tropical fish micro pellets (Hikari, Japan), with immature fish fed an

additional meal on weekdays of frozen daphnia and bloodworms.

Experimental procedure and selection

Each replicate was started using 100 males, evenly sampled across the three stock populations. We

quantified the amount of orange coloration for each male (see *Color pattern extraction and*

quantification, below), and selected the 30 males with the largest and 30 males with the smallest total

orange area to form paired up- and down-selected lines. For the P generation, these males were then

randomly paired with females. To increase our statistical power to detect sex-linked variation, we paired

10 males of each line with two females each to generate half siblings, and 20 males with a single female

each, creating a partial half-sib breeding design. For subsequent generations, we selected the 30 males

with greatest orange area in up-selected lines, and the 30 males with least orange for down-selected

lines. Additionally, we increased the strength of selection by selecting on females. Since females do not

express orange coloration, we used an animal model to estimate breeding values (see *Quantitative*

genetics, below). We then selected the females with greatest and least breeding value (40 for each

category), incorporating uncertainty by ranking across draws from the posterior distribution. To avoid a

rapid loss of genetic variation, and to limit the selective advantage of males paired with two females, we

further limited our selection to include a maximum of three sons from each sire, and three daughters

from each dam.

For the males matched with two females, females were kept in separate tanks, and the male was moved

between females each week. We collected broods from the up- and down-selected lines until we had 90

broods per generation per replicate. 93.3% of paired males and 89.3% of paired females successfully

sired offspring, resulting in an average of 1.46 broods per sire, and 1.10 broods per dam. Having >1

brood per pair allowed us to estimate the magnitude of brood effects.

Broods were collected within a few days of birth to avoid infanticide and moved to their own tank. To

maintain similar developmental rates, broods larger than ten individuals were randomly reduced to ten.

Broods were checked three times per week for signs of male maturation (first signs of gonopodium

growth), and males were moved to a separate tank to avoid sibling mating. The tanks with male brood

members were then queued for phenotyping after visual confirmation of sexual maturity (fully

developed gonopodium and color pattern), and a minimum age of 60 days. After phenotyping (see *Color*

pattern extraction and quantification, below), males were housed in individual tanks to maintain their

identities.

The three replicates were set up staggered in time, so that one replicate could be phenotyped while the

others were breeding and maturing. Fish were only marked with their individual IDs, blinding the

experimenters as to which line individuals belonged to throughout experimental procedures and

phenotyping (except when setting up the next generation of breeding pairs).

Animal photography and calibration

We (WvdB & JJS) captured photos of male color patterns using a Canon EOS Rebel T7i (Canon, Japan)

with Canon EF 100mm f/2.8 Macro USM lens (Canon, Japan). All photos were captured using identical

camera settings, at an aperture of f/11, an exposure time of 1/320s, an ISO of 800 and an approximate

distance of 35-40 cm, and we captured a color standard *in situ* before each photography session

(ColorChecker Passport Photo 2, X-rite, USA). Photos were captured in RAW format, then color

corrected in Adobe Lightroom (Adobe, USA), see *Supplemental methods*.

The photarium (photo tank) was placed on a pedestal covered with matte black foam, inside a 30 x 60 x

30 cm photo box. The walls and top of the box were constructed of opaque white plastic, to minimize

the influence of external lights. In the top, directly above the pedestal, we mounted 2 Neewer NL660

LED panels with white diffusers, with a color rendering index of >= 96 (Neewer, China).

We photographed both sides of every fish, and for 99.9% of fish we had at least three photos, with the

replicate photos allowing us to parse measurement error from asymmetry. We analyzed a total of

14,100 photos, for an average of 4.37 photos per fish.

Color pattern extraction and quantification

After color correction, we used a custom pipeline to analyze the images. The pipeline consists of deep

neural networks for image segmentation and landmark placement, and geometric morphometrics for

image alignment. To be able to phenotype fish during the selection procedure, when we still had little

data available to train the models, we adopted a semi-autonomous strategy. We used an interactive

script to phenotype fish, which prompted the experimenter (WvdB) to visually check the output of each

analysis step, and correct manually when the model prediction was incorrect. We then intermittently

updated model weights with the growing number of manually annotated examples to improve model

performance, gradually reducing the need for manual intervention throughout the experiment. The

pipeline consisted of five steps (SI). By comparing the number of pixels in the extracted fish mask from

Step 1, and the number of pixels in the extracted carotenoid and melanin masks from Step 2 (i.e. before

warping), we calculated the amount of coloration as a percentage of the body area. For use in our

selection paradigm, we calculated the mean across multiple photos of each side of each fish, and then

the mean across the sides.

Quantifying effects of selection

We quantified the effect of selection on the amount of orange and black coloration using Bayesian linear

mixed models (LMMs) using $brms$[78,79]. We modeled the amount of color as a percentage of the body

area as the dependent variable, with fixed effects of generation, selection direction and their

interaction, and a random intercept per generation nested within experimental replicate. We then

calculated the contrast in marginal means between up- and down-selected fish of the F_3 generation

using $emmeans$[80]. We used the same modelling approach for the effect of selection on life history

parameters, except that we modelled fecundity using the negative binomial distribution. In addition, we

estimated the effect of selection on orange and black color per body location (Fig 2B & S3) using

independent binomial generalized linear mixed models with $lme4$[81], using the same model structure,

and the presence/absence of orange as the dependent variable.

Male sexual behavior

We assessed the sexual behavior of 126 male offspring from the F3 generation (63 from each selection

direction, evenly split across the replicate lines). We isolated males upon early indications of sexual

maturity (first signs of gonopodial development), then at around three months of age tested their sexual

behaviors using a no choice test with nonreceptive females, a standard for studying male sexual

behavior in poecilids[30]. Behavioral trials took place in a 47 cm diameter arena, with fish transfer

followed by a 5-minute acclimation to minimize stress and a 15-minute observation using a webcam at

1080p and 30 fps. We analyzed the recorded videos to quantify male sexual behaviors. We utilized

idTracker[82] for positional data during each trial. Trials where females exhibited receptivity or where fish

showed signs of stress were excluded, leading to a final dataset of 95 individuals. Statistical analysis on

behavioral differences between up- and down-selected males used generalized linear mixed models

implemented in *glmmTMB*[83], with behavioral metrics as dependent variables, selection direction, male

body size, and female body size as fixed effects, and replicate line as random effect. Behavioral counts

were modelled using the negative binomial distribution, while for time spent following we used the

Gaussian distribution with log link function.

Ornaments

We defined areas of the fish as ornaments manually, based on the visible spatial structure in the

incidence of color and the localized genetic architecture (Figs 1C & 3). We then overlayed these areas on

the extracted color patterns from each photo and counted the number of pigmented pixels within each

area. We scored an ornament as "absent" (ornament size = 0) if < 10% of the ornament area was

pigmented. This limit was chosen based on the bimodal distribution of the number of pixels, as in some

cases a few pixels from a neighboring ornament spilled over into other areas.

Quantitative genetics

Estimating of quantitative genetic (QG) parameters for the amount of orange and black color was

performed by MCMC sampling of Gaussian animal models implemented in *Stan*[84] with *brms*[78,79]. Additive

genetic variance-covariance matrices were calculated from the pedigree using *nadiv*[85], with the *S* matrix

for X-linked effects assuming no global dosage compensation[36]. Y-linked genetic variances were

estimated by including patriline as a group level effect. Similarly, we included dam identity to estimate

maternal effects, included brood identity to capture shared environmental effects between siblings, and

individual identity to estimate the remaining environmental influence, all as group level effects. We

included the side of the fish, nested in individual identity, as a group level effect to estimate the variance

due to asymmetry, and included the residual variance to estimate measurement error. For all variances,

including the residual variances, we estimated their correlation among the two colors. We used default

non-informative priors for all parameters.

For the QG heatmaps (Fig 3), we estimated separate animal models for each cell, estimating orange and

black colors separately. We used the same model structure as the model above, now with a Bernoulli

error structure. We also removed the maternal effect, as the clutch and maternal effects are correlated

and their magnitudes are small. Since estimating these parameters independently thousands of times

using MCMC is prohibitively computationally expensive, we used an approach that makes use of the fact

that the QG estimations will be similar between neighboring cells. To limit the number of models, we

reduced the dimensions of the color pattern images by a factor of two (250 x 70 cells or pixels), and only

ran models for cells with an incidence of at least 1%. We then performed model fitting in two rounds.

First, we fit animal models for the cells with x- and y-coordinates divisible by five. For these models, we

used four chains, with 500 warmup iterations and 1500 sampling iterations per chain, and uninformative

StudentT(3, 0, 5) priors on the intercept and variance components. In a second step, we then used the

output of these models to fill in the cells between this five-by-five grid. These models only used two

chains of 500 warmup iterations and 500 sampling iterations, but were seeded with initial parameter

values and step sizes that were the average of the closest nine models from Step 1, weighted by their

inverse distance. Similarly, we set normal priors for the intercept and variance components with the

mean (also weighted by inverse distance) and standard deviation of those parameters of the closest

nine models.

For ornament heritability, we used similar models to the amount of orange and black color, but used

hurdle-lognormal models instead to jointly estimate the variance components for the incidence and size

(when present) of the ornaments. Given the very small maternal effects on total color, we did not

include that parameter here. We specified an uninformative logistic(0, 5) prior on the hurdle parameter,

and a StudentT(3, 0, 5) parameter on the standard deviations of the group level effects, with otherwise

default priors.

For the pairwise correlations across ornaments, we used the Gaussian approximation and fitted the

multivariate animal model on individual-level means using the restricted maximum likelihood method

from *sommer*[86]. Given the small magnitude of brood effects from the full Bayesian animal models for

each ornament, we did not include that effect here.

Learning of patternspace

We devised a deep learning approach to map images of guppies onto a "patternspace". Our objective

was to capture the heritable components of color patterns by learning a space where images of related

guppies are placed closer in this patternspace compared to those of unrelated males. Our method is

predicated on the "triplet loss learning" paradigm[32]. In this paradigm, a model maps images into a space,

one at a time, using a learned metric. Because the model uses the same metric for each image, images

that are placed close together are similar according to that metric. As our goal was to obtain a space

describing the heritable variation in color patterns, we trained the model to use a metric that captured

this heritable variation. This training used triplets of images, which comprise an anchor image (a random

male), a positive image (of a related male), and a negative image (of an unrelated male). By comparing

the distances between these three images, we can evaluate whether the metric is capturing heritable

variation. During training, each image is passed through the model and the triplet loss function is

applied, optimizing the metric to reduce the distances between anchors and positives, while increasing

the distances between anchors and negatives. Unrelated males had a coefficient of relatedness r of 0,

while for related males r was at least 1/8.

The deep learning architecture was designed to process images of dimensions 250x70x3, with the last

dimension being color channels. Our model, a Convolutional Neural Network (CNN), had a series of

layers that successively compress the images to lower dimensions. Specifically, we used separable

convolutional layers with a kernel size of 3x3 and *relu* activation, followed by a max pooling layer,

repeating this four times. Each convolutional layer had double the filters of the previous, starting at 8

filters. After the fourth pooling layer, we flattened the output to a dense layer of size 8, and then a

second dense layer of the embedding dimension. Lastly, a batch normalization layer normalized

activations to unit variance. Embeddings were obtained at this normalization layer.

We trained for 50 epochs using a batch size of 128 images. At each epoch, we randomly generated 5,000

triplets. We randomly allocated 20% of individuals to the validation set, from which we generated 500

triplets per epoch to track model performance. Moreover, learning rate schedules were implemented to

adjust the learning rate across epochs, with specified rates of 10^{-4}, 10^{-5}, and 10^{-6} at epochs 0, 30, and 40

respectively. This scheduling ensured a faster convergence initially, with fine-tuning during the later

stages of training.

We evaluated the number of embedding dimensions by training the model at one to eight dimensions,

then chose five dimensions, since the model accuracy for the validation sample stopped increasing

above five. After training, we used the CNN to embed all 14,100 images. All analyses use the five

embedding dimensions while for visualization we performed dimension reduction using UMAP[87], which

aims to preserve the local neighborhood structure of the data, as implemented in R package *uwot*[88].

Genetic sampling and whole genome sequencing

To maximize power for the genetic mapping of color traits, we sampled 50 fish from each selection

line (N = 300) for whole genome sequencing (WGS). To maximize genetic and phenotypic variation in our

sample, and therefore power, we sampled males from each family with offspring in generation F_3, and

sampled males with the most/least carotenoid coloration in up/down-selected lines respectively. This

meant we sampled one son from each family, with some families contributing a second son until we

reached the desired 50 males. Additional methods are in the SI.

DNA was extracted from sampled males using Qiagen Dneasy blood & tissue kits (Qiagen, USA). Samples

were then sequenced on four lanes of Illumina NovaSeq 6000 S4 PE150 to an average depth of 10x per

individual. One individual was not included because the library failed. We checked the identity of

samples by visually comparing the genome-wide relatedness (GRM) with the pedigree-based

relatedness (A) and excluded two samples which did not match, leaving 297 samples for analysis.

Genome-wide association analysis

We first trimmed the adapters using *Trimmomatic* and then mapped the reads to the reference using

bwa mem using default mapping parameters. We marked PCR duplicates using GATK's *MarkDuplicates*.

We then called variants using *bcftools mpileup* where we downgraded the mapping quality of reads

containing mismatches with *–adjust-MQ 60*, and set a minimal mapping quality using *–min-MQ 30*,

followed by variant calling using the consensus caller *bcftools call*. We then performed variant filtering,

and dropped sites with a QUAL score < 30, MQ < 40, FS > 60, |BQBZ| > 10, |MQSBZ| > 10 or |RPBZ| >

10, and sites where more than 20% of individuals had a depth < 5, or > 20% of individuals had a GQ

score < 10. Additionally, we filtered out sites with a major allele frequency < 10% or a missingness > 5%,

as well as indels.

We performed GWAS using the two-step "population parameters previously determined" (P3D)

approach[89] implemented in *sommer*[86]. We controlled for relatedness and population structure by

including random effect terms for the autosomal, X-linked and Y-linked portions of the genome, as well

as a fixed effect for selection direction. We determined the population level parameters using the full

set of phenotyped individuals. For the autosomal relatedness, we combined the pedigree-based matrix

A with the genomic relatedness matrix (GRM) estimated using GEMMA[90] for the sequenced samples, to

obtain the single-step matrix H[91]. We used the *S* matrix for X-linked relatedness, and patriline identity

for Y-linked relatedness. We then performed marker-wise effect size estimation for the subset of

sequenced individuals. We performed multivariate GWAS by performing this procedure on each of the 5

axes of patternspace. We then combined the Z-scores to obtain multivariate p-values using CPASSOC[92].

For univariate ornament GWAS, we used the same approach, using the normal approximation, and only

including ornaments with an incidence between 5% and 95%.

The phenotypic effects of individual variants were evaluated with PheWAS using *sommer*, where we

controlled for autosomal, X- and Y-linked inheritance, fixing those variance parameters at the posterior

medians of the per-pixel quantitative genetic analyses described above.

Coverage statistics were calculated using *samtools depth*, with a window size of 1000bp and a window

distance of 100bp. We additionally calculated coverage for sequencing data from 3 females from the

same population taken from[93] (NCBI BioProject accession PRJNA858015, sequenced at ~30x coverage),

which were re-analyzed using the identical pipeline used for the males from the current study. We

controlled the false discovery rate (FDR) using the Benjamini & Hochberg (BH) procedure[94] separately

for the orange and black patternspace GWAS. For ornament GWAS, we performed the BH procedure

jointly across the ornaments, but separately per color. FDR is controlled to 5% within each of these four

sets of tests. We compared the overlap in associated variants between traits using the *ComplexUpset*

package[95].

We performed principal component analysis to detect Y-haplogroups using *irlba*[96]. We assigned the

haplogroups using hierarchical clustering on the Euclidian distance matrix of the first 20 principal

components using *hclust*.

Inference of variant inheritance pattern

The inheritance pattern of all called variants was estimated using animal models. We encoded the

genotypes as doses of the alternative allele, and then fitted three models using *sommer*[86] with dose as

the dependent variable. The three models included a random effect for the autosomal, X-linked or Y-

linked covariance, respectively, as described in *Quantitative genetics*. For each model we calculated

Akaike's Information Criterion (AIC) and their AIC weight (i.e. the relative likelihood) to compare

between the three hypotheses. We called the inheritance of a variant as sex-linked if the relative

likelihood of X- or Y-linkage exceeded 80%. We quantified the performance of this approach by calling

the inheritance for variants called on the autosomes (autosomal: 98.8%, X-linked: 0.8%, Y-linked: 0. 3%),

confirming that a sex-linked result is highly informative for placement on LG12. In addition, we

visualized the inheritance of 100,000 variants along LG12 showing patterns of sex-linked inheritance

outside the pseudo-autosomal region (Fig S32).

Expression analysis

Following the end of selection, we sampled three replicate male and female pools of seven individuals

each from each resulting post-selection population. RNA was extracted from whole tail tissue (posterior

to the anal fin) using Qiagen RNeasy Mini kits (Qiagen, USA). Long-read sequencing was conducted using

PacBio Revio (Pacific Biosciences, USA) and reads were subsequently processed with the PacBio IsoSeq

pipeline. Reads were mapped to the female guppy reference using minimap2[97] after clustering. We used

FLAIR[98] to further collapse redundant reads and quantify expression using counts per million (CPM). Five

copies of the *texim* gene on the Y chromosome were identified by mapping Y-linked genotypes (as

determined above) to collapsed full-length RNA reads, of which one (copy "1889") was expressed at >1

CPM. Finally, we genotyped the 297 sequenced individuals for the presence of copy 1889 if they

possessed the Y-linked genotype for > 50% of the SNPs in this copy.

Software

Unless stated otherwise above, all analyses were performed in the R statistical environment[99].

References

1. Stephenson, J. F., Stevens, M., Troscianko, J. & Jokela, J. The Size, Symmetry, and Color Saturation of a Male
Guppy's Ornaments Forecast His Resistance to Parasites. *Am. Nat.* **196**, 597–608 (2020).
2. Karino, K., Shinjo, S. & Sato, A. Algal-searching ability in laboratory experiments reflects orange spot coloration
of the male guppy in the wild. *Behaviour* **144**, 101–113 (2007).
3. Evans, J. P., Kelley, J. L., Bisazza, A., Finazzo, E. & Pilastro, A. Sire attractiveness influences offspring
performance in guppies. *Proc. R. Soc. Lond. B Biol. Sci.* **271**, 2035–2042 (2004).
4. Endler, J. A. Natural selection on color patterns in *Poecilia reticulata*. *Evolution* **34**, 76–91 (1980).
5. Potter, T. *et al.* Female preference for rare males is maintained by indirect selection in Trinidadian guppies.
Science **380**, 309–312 (2023).
6. Houde, A. E. Mate choice based upon naturally occurring color-pattern variation in a guppy population.
Evolution **41**, 1–10 (1987).
7. Houde, A. E. & Endler, J. A. Correlated evolution of female mating preferences and male color patterns in the
guppy *Poecilia reticulata*. *Science* **248**, 1405–1408 (1990).
8. Haskins, C. P., Haskins, E. F., McLaughlin, J. J. A. & Hewitt, R. E. Polymorphism and population structure in
Lebistes reticulatus, an ecological study. in *Vertebrate speciation* (1961).
9. Kemp, D. J., Reznick, D. N., Grether, G. F. & Endler, J. A. Predicting the direction of ornament evolution in
Trinidadian guppies (*Poecilia reticulata*). *Proc. R. Soc. B Biol. Sci.* **276**, 4335–4343 (2009).
10. Herdegen-Radwan, M., Cattelan, S., Buda, J., Raubic, J. & Radwan, J. What do orange spots reveal about male
(and female) guppies? A test using correlated responses to selection. *Evolution* **75**, 3037–3055 (2021).
11. Brooks, R. & Endler, J. A. Direct and indirect sexual selection and quantitative genetics of male traits in guppies
(*Poecilia reticulata*). *Evolution* **55**, 1002–1015 (2001).
12. Morris, J., Darolti, I., van der Bijl, W. & Mank, J. E. High-resolution characterization of male ornamentation and
re-evaluation of sex linkage in guppies. *Proc. R. Soc. B Biol. Sci.* **287**, 20201677 (2020).
13. Hughes, K. A., Houde, A. E., Price, A. C. & Rodd, F. H. Mating advantage for rare males in wild guppy
populations. *Nature* **503**, 108–110 (2013).
14. Farr, J. A. Male Rarity or Novelty, Female Choice Behavior, and Sexual Selection in the Guppy, Poecilia
reticulata Peters (Pisces: Poeciliidae). *Evolution* **31**, 162–168 (1977).
15. Parichy, D. M. & Spiewak, J. E. Origins of adult pigmentation: diversity in pigment stem cell lineages and
implications for pattern evolution. *Pigment Cell Melanoma Res.* **28**, 31–50 (2015).
16. Gordon, S. P., López-Sepulcre, A. & Reznick, D. N. Predation-associated differences in sex linkage of wild guppy
coloration. *Evolution* **66**, 912–918 (2012).
17. Toomey, M. B. *et al.* A mechanism for red coloration in vertebrates. *Curr. Biol.* **32**, 4201-4214.e12 (2022).
18. Lindholm, A. K. & Breden, F. Sex Chromosomes and Sexual Selection in Poeciliid Fishes. *Am. Nat.* **160**, (2002).
19. Tripathi, N. *et al.* Genetic linkage map of the guppy, Poecilia reticulata, and quantitative trait loci analysis of
male size and colour variation. *Proc. R. Soc. B Biol. Sci.* **276**, 2195–2208 (2009).
20. Nanda, I. *et al.* Sex chromosome polymorphism in guppies. *Chromosoma* **123**, 373–383 (2014).
21. Kirpichnikov, V. S. *Genetic Bases of Fish Selection*. (Berlin : Springer, 1981).

22. Gordon, S. P., López-Sepulcre, A., Rumbo, D. & Reznick, D. N. Rapid changes in the sex linkage of male
coloration in introduced guppy populations. *Am. Nat.* **189**, 196–200 (2017).
23. Kruuk, L. E. B. Estimating genetic parameters in natural populations using the 'animal model'. *Philos. Trans. R.*
Soc. Lond. B. Biol. Sci. **359**, 873–890 (2004).
24. Winge, Ø. One-sided masculine and sex-linked inheritance inLebistes reticulatus. *J. Genet.* **12**, 145–162 (1922).
25. Winge, Ø. & Ditlevsen, E. Colour inheritance and sex determination in Lebistes. *Heredity* **1**, 65–83 (1947).
26. Haskins, C. P., Young, P., Hewitt, R. E. & Haskins, E. F. Stabilised heterozygosis of supergenes mediating certain
Y-linked colour patterns in populations of Lebistes Reticulatus. *Heredity* **25**, 575–589 (1970).
27. Parichy, D. M., Ransom, D. G., Paw, B., Zon, L. I. & Johnson, S. L. An orthologue of the kit-related gene fms is
required for development of neural crest-derived xanthophores and a subpopulation of adult melanocytes in
the zebrafish, Danio rerio. *Development* **127**, 3031–3044 (2000).
28. Wilkins, A. S., Wrangham, R. W. & Fitch, W. T. The "Domestication Syndrome" in Mammals: A Unified
Explanation Based on Neural Crest Cell Behavior and Genetics. *Genetics* **197**, 795–808 (2014).
29. Liley, N. R. Ethological Isolating Mechanisms in Four Sympatric Species of Poeciliid Fishes. *Behav. Suppl.* **1**, 1–
197 (1966).
30. Houde, A. E. *Sex, Color, and Mate Choice in Guppies*. (Princeton University Press, 1997).
31. Brooks, R. & Endler, J. A. Female Guppies Agree to Differ: Phenotypic and Genetic Variation in Mate-Choice
Behavior and the Consequences for Sexual Selection. *Evolution* **55**, 1644–1655 (2001).
32. Schroff, F., Kalenichenko, D. & Philbin, J. FaceNet: A unified embedding for face recognition and clustering. in
2015 IEEE Conference on Computer Vision and Pattern Recognition (CVPR) 815–823 (2015).
doi:10.1109/CVPR.2015.7298682.
33. Almeida, P. *et al.* Divergence and Remarkable Diversity of the Y Chromosome in Guppies. *Mol. Biol. Evol.* **38**,
619–633 (2021).
34. Du, K. *et al.* Identification of the male-specific region on the guppy Y Chromosome from a haplotype-resolved
assembly. *Genome Res.* (2025) doi:10.1101/gr.279582.124.
35. Wright, A. E. *et al.* Convergent recombination suppression suggests role of sexual selection in guppy sex
chromosome formation. *Nat. Commun.* **8**, 14251 (2017).
36. Darolti, I. *et al.* Extreme heterogeneity in sex chromosome differentiation and dosage compensation in
livebearers. *Proc. Natl. Acad. Sci.* **116**, 19031–19036 (2019).
37. Künstner, A. *et al.* The Genome of the Trinidadian Guppy, Poecilia reticulata, and Variation in the Guanapo
Population. *PLOS ONE* **11**, e0169087 (2016).
38. Fraser, B. A. *et al.* Improved Reference Genome Uncovers Novel Sex-Linked Regions in the Guppy (Poecilia
reticulata). *Genome Biol. Evol.* **12**, 1789–1805 (2020).
39. Barriga, E. H. & Mayor, R. Chapter Nine - Embryonic Cell–Cell Adhesion: A Key Player in Collective Neural Crest
Migration. in *Current Topics in Developmental Biology* (ed. Yap, A. S.) vol. 112 301–323 (Academic Press,
2015).
40. Yin, L. *et al.* Epidermal gene expression and ethnic pigmentation variations among individuals of Asian,
European and African ancestry. *Exp. Dermatol.* **23**, 731–735 (2014).
41. Yong, L., Croft, D. P., Troscianko, J., Ramnarine, I. W. & Wilson, A. J. Sensory-based quantification of male
colour patterns in Trinidadian guppies reveals no support for parallel phenotypic evolution in multivariate trait
space. *Mol. Ecol.* **31**, 1337–1357 (2022).
42. Hoek, K. S. *et al.* Novel MITF targets identified using a two-step DNA microarray strategy. *Pigment Cell*
Melanoma Res. **21**, 665–676 (2008).
43. Zhang, Y. *et al.* Transcriptome analyses of betta fish (*Betta splendens*) provide novel insights into fin
regeneration and color-related genes. *Gene* **876**, 147508 (2023).
44. Yu, H.-H. & Moens, C. B. Semaphorin signaling guides cranial neural crest cell migration in zebrafish. *Dev. Biol.*
280, 373–385 (2005).
45. Koludrovic, D. *et al.* Chromatin-Remodelling Complex NURF Is Essential for Differentiation of Adult
Melanocyte Stem Cells. *PLOS Genet.* **11**, e1005555 (2015).
46. Miyadai, M. *et al.* Pax3 and Pax7 function in combination with Mitf to generate melanophores and
xanthophores in medaka and zebrafish. 2023.06.22.546052 Preprint at
https://doi.org/10.1101/2023.06.22.546052 (2023).

47. Bhattacharya, D., Rothstein, M., Azambuja, A. P. & Simoes-Costa, M. Control of neural crest multipotency by
Wnt signaling and the Lin28/let-7 axis. *eLife* **7**, e40556 (2018).
48. Takahashi, K. *et al.* A globin-family protein, Cytoglobin 1, is involved in the development of neural crest-
derived tissues and organs in zebrafish. *Dev. Biol.* **472**, 1–17 (2021).
49. Boer, E. F., Howell, E. D., Schilling, T. F., Jette, C. A. & Stewart, R. A. Fascin1-Dependent Filopodia are Required
for Directional Migration of a Subset of Neural Crest Cells. *PLOS Genet.* **11**, e1004946 (2015).
50. Lin, Y. *et al.* Gene duplication to the Y chromosome in Trinidadian guppies. *Mol. Ecol.* **31**, 1853–1863 (2022).
51. Bergero, R., Gardner, J., Bader, B., Yong, L. & Charlesworth, D. Exaggerated heterochiasmy in a fish with sex-
linked male coloration polymorphisms. *Proc. Natl. Acad. Sci.* **116**, 6924–6931 (2019).
52. Tomaszkiewicz, M., Chalopin, D., Schartl, M., Galiana, D. & Volff, J.-N. A multicopy Y-chromosomal SGNH
hydrolase gene expressed in the testis of the platyfish has been captured and mobilized by a Helitron
transposon. *BMC Genet.* **15**, 44 (2014).
53. Sandkam, B. A. *et al.* Extreme Y chromosome polymorphism corresponds to five male reproductive morphs of
a freshwater fish. *Nat. Ecol. Evol.* **5**, 939–948 (2021).
54. Lindsay, C. R. *et al.* P-Rex1 is required for efficient melanoblast migration and melanoma metastasis. *Nat.*
Commun. **2**, 555 (2011).
55. Endler, J. A. Natural and sexual selection on color patterns in poeciliid fishes. in *Evolutionary ecology of*
neotropical freswater fishes 95–111 (Springer, The Netherlands, 1984).
56. Kotrschal, A. *et al.* Artificial Selection on Relative Brain Size in the Guppy Reveals Costs and Benefits of
Evolving a Larger Brain. *Curr. Biol.* **23**, 168–171 (2013).
57. Fong, S. *et al.* Rapid mosaic brain evolution under artificial selection for relative telencephalon size in the
guppy (Poecilia reticulata). *Sci. Adv.* **7**, eabj4314 (2021).
58. Kotrschal, A. *et al.* Rapid evolution of coordinated and collective movement in response to artificial selection.
Sci. Adv. **6**, eaba3148 (2020).
59. Houde, A. E. Effect of artificial selection on male colour patterns on mating preference of female guppies.
Proc. R. Soc. Lond. B Biol. Sci. **256**, 125–130 (1994).
60. Cole, G. L. & Endler, J. A. Change in male coloration associated with artificial selection on foraging colour
preference. *J. Evol. Biol.* **31**, 1227–1238 (2018).
61. Yamamoto, T. The Medaka, Oryzias latipes, and the Guppy, Lebistes reticularis. in *Handbook of Genetics:*
Volume 4 Vertebrates of Genetic Interest (ed. King, R. C.) 133–149 (Springer US, Boston, MA, 1975).
doi:10.1007/978-1-4613-4470-4_7.
62. Houde, A. E. Sex-linked heritability of a sexually selected character in a natural population of *Poecilia*
reticulata (Pisces: Poeciliidae) (guppies). *Heredity* **69**, 229–235 (1992).
63. Inaba, M., Yamanaka, H. & Kondo, S. Pigment Pattern Formation by Contact-Dependent Depolarization.
Science **335**, 677–677 (2012).
64. Kottler, V. A., Fadeev, A., Weigel, D. & Dreyer, C. Pigment Pattern Formation in the Guppy, Poecilia reticulata,
Involves the Kita and Csf1ra Receptor Tyrosine Kinases. *Genetics* **194**, 631–646 (2013).
65. Kottler, V. A. *et al.* Adenylate cyclase 5 is required for melanophore and male pattern development in the
guppy (Poecilia reticulata). *Pigment Cell Melanoma Res.* **28**, 545–558 (2015).
66. Brush, A. H. Metabolism of carotenoid pigments in birds. *FASEB J.* **4**, 2969–2977 (1990).
67. Hill, G. E. & Johnson, J. D. The Vitamin A–Redox Hypothesis: A Biochemical Basis for Honest Signaling via
Carotenoid Pigmentation. *Am. Nat.* **180**, E127–E150 (2012).
68. Grether, G. F., Hudon, J. & Millie, D. F. Carotenoid limitation of sexual coloration along an environmental
gradient in guppies. *Proc. R. Soc. Lond. B Biol. Sci.* **266**, 1317–1322 (1999).
69. Lopes, R. J. *et al.* Genetic Basis for Red Coloration in Birds. *Curr. Biol.* **26**, 1427–1434 (2016).
70. Mundy, N. I. *et al.* Red Carotenoid Coloration in the Zebra Finch Is Controlled by a Cytochrome P450 Gene
Cluster. *Curr. Biol.* **26**, 1435–1440 (2016).
71. McLean, C. A. *et al.* Red carotenoids and associated gene expression explain colour variation in frillneck
lizards. *Proc. R. Soc. B Biol. Sci.* **286**, 20191172 (2019).
72. Pérez-Gálvez, A. & Mínguez-Mosquera, M. I. Esterification of xanthophylls and its effect on chemical behavior
and bioavailability of carotenoids in the human. *Nutr. Res.* **25**, 631–640 (2005).
73. Breithaupt, D. E., Bamedi, A. & Wirt, U. Carotenol fatty acid esters: easy substrates for digestive enzymes?
Comp. Biochem. Physiol. B Biochem. Mol. Biol. **132**, 721–728 (2002).

74. Paris, J. R. *et al.* A large and diverse autosomal haplotype is associated with sex-linked colour polymorphism in
the guppy. *Nat. Commun.* **13**, 1233 (2022).
75. Bruders, R. *et al.* A copy number variant is associated with a spectrum of pigmentation patterns in the rock
pigeon (Columba livia). *PLOS Genet.* **16**, e1008274 (2020).
76. Weich, K. *et al.* Pigment Intensity in Dogs is Associated with a Copy Number Variant Upstream of KITLG. *Genes*
11, 75 (2020).
77. Kokko, H., Jennions, M. D. & Houde, A. Evolution of frequency-dependent mate choice: keeping up with
fashion trends. *Proc. R. Soc. B Biol. Sci.* **274**, 1317–1324 (2007).
78. Bürkner, P.-C. Advanced Bayesian Multilevel Modeling with the R Package brms. *R J.* **10**, 395–411 (2018).
79. Bürkner, P.-C. brms: An R Package for Bayesian Multilevel Models Using Stan. *J. Stat. Softw.* **80**, 1–28 (2017).
80. Lenth, R. V. *Emmeans: Estimated Marginal Means, Aka Least-Squares Means*. (2023).
81. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models Using lme4. *J. Stat. Softw.*
67, 1–48 (2015).
82. Pérez-Escudero, A., Vicente-Page, J., Hinz, R. C., Arganda, S. & de Polavieja, G. G. idTracker: tracking
individuals in a group by automatic identification of unmarked animals. *Nat. Methods* **11**, 743–748 (2014).
83. Brooks, M. E. *et al.* glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized
Linear Mixed Modeling. *R J.* **9**, 378–400 (2017).
84. Carpenter, B. *et al.* Stan: A Probabilistic Programming Language. *J. Stat. Softw.* **76**, (2017).
85. Wolak, M. E. nadiv: an R package to create relatedness matrices for estimating non-additive genetic variances
in animal models. *Methods Ecol. Evol.* **3**, 792–796 (2012).
86. Giovanny, C.-P. Genome assisted prediction of quantitative traits using the R package sommer. *PLoS ONE* **11**,
1–15 (2016).
87. McInnes, L., Healy, J. & Melville, J. UMAP: Uniform Manifold Approximation and Projection for Dimension
Reduction. Preprint at https://doi.org/10.48550/arXiv.1802.03426 (2020).
88. Melville, J. *Uwot: The Uniform Manifold Approximation and Projection (UMAP) Method for Dimensionality*
Reduction. (2023).
89. Zhang, Z. *et al.* Mixed linear model approach adapted for genome-wide association studies. *Nat. Genet.* **42**,
355–360 (2010).
90. Zhou, X. & Stephens, M. Genome-wide efficient mixed-model analysis for association studies. *Nat. Genet.* **44**,
821–824 (2012).
91. Legarra, A., Aguilar, I. & Misztal, I. A relationship matrix including full pedigree and genomic information. *J.*
Dairy Sci. **92**, 4656–4663 (2009).
92. Li, X. & Zhu, X. Cross-Phenotype Association Analysis Using Summary Statistics from GWAS. in *Statistical*
Human Genetics: Methods and Protocols (ed. Elston, R. C.) 455–467 (Springer, New York, NY, 2017).
doi:10.1007/978-1-4939-7274-6_22.
93. Lin, Y., Darolti, I., Bijl, W. van der, Morris, J. & Mank, J. E. Extensive variation in germline de novo mutations in
Poecilia reticulata. *Genome Res.* **33**, 1317–1324 (2023).
94. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to
Multiple Testing. *J. R. Stat. Soc. Ser. B Methodol.* **57**, 289–300 (1995).
95. Krassowski, M. ComplexUpset. (2020) doi:10.5281/zenodo.3700590.
96. Baglama, J., Reichel, L. & Lewis, B. W. *Irlba: Fast Truncated Singular Value Decomposition and Principal*
Components Analysis for Large Dense and Sparse Matrices. (2022).
97. Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–3100 (2018).
98. Tang, A. D. *et al.* Full-length transcript characterization of SF3B1 mutation in chronic lymphocytic leukemia
reveals downregulation of retained introns. *Nat. Commun.* **11**, 1438 (2020).
99. R Core Team. *R: A Language and Environment for Statistical Computing*. (R Foundation for Statistical
Computing, Vienna, Austria, 2023).
835