1   Resident Synovial Macrophages in Synovial Fluid:
2   Implications for Immunoregulation in Infectious and Inflammatory Arthritis
3

Karen I. Cyndari, MD, PhD[1,2], Breanna M. Scorza, PhD[2,3], Zeb R. Zacharias, PhD[4,5],
Leela Strand[6], Kurayi Mahachi, MPH, PhD[7], Juan Marcos Oviedo, PhD[8], Lisa Gibbs,
BS[8], Danielle Pessoa-Pereira, PhD[2,3], Graham Ausdal, MPH[2,3], Dylan Hendricks, MS[2,3],
Rika Yahashiri[9], Jacob M. Elkins, MD, PhD[10], Trevor Gulbrandsen, MD[10], Andrew R.
Peterson, MSPH, MD[10], Michael C. Willey, MD[10], Keke C. Fairfax, PhD[8,] and Christine
A. Petersen, DVM, PhD[2,3]
10
[1]Department of Emergency Medicine, University of Iowa, Iowa City, IA
[2]Center for Emerging Infectious Diseases
[3]Department of Epidemiology, University of Iowa, Iowa City, IA
[4]Human Immunology Core, University of Iowa, Iowa City, IA
[5]Holden Comprehensive Cancer Center, Iowa City, IA
[6]North Liberty High School, North Liberty, IA
[7]Research and Analytics, Enterprise Analytics, Sentara Health
[8]Department of Pathology, University of Utah, Salt Lake City, UT
[9]Williams College, Williamstown, MA
[10]Department of Orthopedics, University of Iowa, Iowa City, IA

21
22

23   Sources of funding:
24   Cyndari: This project was supported by funding from the Herbert N. Hultgren Award
25   (Wilderness Medicine Society), the University of Iowa Physician-Scientist Training
26   Program, and the University of Iowa Department of Emergency Medicine.

27
28

29   Corresponding Author:
30   Karen Cyndari, MD, PhD
31   Tel: 585-797-3464
32   karen-cyndari@uiowa.edu
33
34   University of Iowa Hospitals and Clinics
35   200 Hawkins Drive
36   GH-SE203
37   Iowa City, IA 52242

38
39

40   Keywords: resident synovial macrophage, Type A synoviocyte, joint space
41   immunoregulation, septic arthritis, inflammatory arthritis

42
43

44
45 **Abstract:**
46
47 **Objectives**: Resident synovial macrophages (RSM) provide immune sequestration of
48 the joint space and are likely involved in initiation and perpetuation of the joint-specific
49 immune response. We sought to identify RSM in synovial fluid (SF) and demonstrate
50 migratory ability, in additional to functional changes that may perpetuate a chronic
51 inflammatory response within joint spaces.
52
Methods: We recruited human patients presenting with undifferentiated arthritis in
multiple clinical settings. We used flow cytometry to identify mononuclear cells in
peripheral blood and SF. We used a novel transwell migration assay with human *ex-*
vivo synovium obtained intra-operatively to validate flow cytometry findings. We used
single cell RNA-sequencing (scRNA-seq) to further identify macrophage/monocyte
subsets. ELISA was used to evaluate the bone-resorption potential of SF.
59
Results: We were able to identify a rare population of CD14^{dim}, OPG$^+$, ZO-1$^+$ cells
consistent with RSM in SF via flow cytometry. These cells were relatively enriched in
the SF during infectious processes, but absolutely decreased compared to healthy
controls. Similar putative RSM were identified using *ex vivo* migration assays when
MCP-1 and LPS were used as migratory stimulus. scRNA-seq revealed a population
consistent with RSM transcriptionally related to CD56$^+$ cytotoxic dendritic cells and IDO$^+$
M2 macrophages.
67
68 **Conclusion**: We identified a rare cell population consistent with RSM, indicating these
69 cells are likely migratory and able to initiate or coordinate both acute (septic) or chronic
70 (autoimmune or inflammatory) arthritis. RSM analysis via scRNA-seq indicated these
71 cells are M2 skewed, capable of antigen presentation, and have consistent functions in
72 both septic and inflammatory arthritis.

Introduction: Damage to the articular surface of joints resulting in arthritis may

be secondary to infection, inflammation, and chronic or acute trauma. Different

etiologies of arthritis result in unique local immune environments within the joint

space. While healthy synovium delineates an immune-privileged space to which few

circulating cells gain entry (1), there are significant numbers of immune cells in the

synovial fluid (SF) of pathologic joints (2; 3; 4). This indicates the localized synovial

immune response is coordinated by the synovium itself, which limits entry to the SF by

actively sequestering inflammatory damage (5) versus allowing circulating immune cells

to enter SF.

Synovium is primarily composed of two types of cells: Fibroblast-like

Synoviocytes (FLS) and Resident Synovial Macrophages (RSMs), also known as Type

A cells. FLS express MHC Class II and produce lubricating joint fluid, including

hyaluronan (6). In pathologic settings, FLS are involved in joint inflammation and,

ultimately, cartilage destruction (6). Conversely, RSMs compose approximately 10% of

the synovium and have only been identified using tissue histology to date. RSMs are

described as constitutively anti-inflammatory, as opposed to circulating monocytes

which may either assist with sequestration of pathology or provide further momentum

toward significant cellular collateral damage (5).

It was recently discovered that RSMs are derived from embryonic precursor cells

and perpetuate through self-proliferation within synovium (7). Certain cellular surface

receptors, chemokines, or structural proteins such as CD68, osteoprotegerin

(OPG/TNFRSF11B), CX3CR1, ZO-1/TJP1, F11R/JAM-A/JAM-1/CD321, and Triggering

Receptor Expressed on Myeloid cells 2 (TREM2) (8; 7) have all been proposed to

identify these cells, yet it remains difficult to differentiate RSMs from macrophages

recruited from circulation. RSMs have not been further evaluated in SF as it is unclear if

they can migrate out of tissue. As inflammation dysregulates the tight junctions

connecting these epithelial-like RSMs (7), it is possible that these cells could leave the

synovium and participate in joint space immune responses.

Here, we used flow cytometry to describe a subset of CD14^{dim}OPG+ZO-1+ M2

macrophages enriched in SF of pathologic joints consistent with previously published

histological descriptions of RSMs. *Ex vivo* migration experiments validated migration

from tissue. Single cell RNA sequencing (scRNA-seq) reveals these putative RSMs had

dysregulated complement in settings of inflammatory arthritis, and a unique reactome

signature involving threonine, niacin, and thiamine metabolism. This work is important in

understanding how damage to the joint space is initiated and perpetuated both during

infectious and inflammatory arthritis.

Materials and Methods:

Patient Recruitment: These studies were approved by the Institutional Review Board

(IRB) at the University of Iowa Hospitals and Clinics. For SF studies, we recruited

patients under evaluation for septic arthritis. Exclusion criteria included significant joint

trauma, joint surgery, or immunomodulatory medications. All patients provided a blood

sample. Not all patients required or had successful arthrocentesis. SF samples were

classified as Normal, Non-Inflammatory, Inflammatory, or Septic based on established

guidelines (9). For synovium procurement, 6 patients >18 years of age receiving a

scheduled, non-emergent, total joint replacement or resection arthroplasty secondary to

infection were recruited. Synovium (knee) or pulvinar (hip) was sterilely obtained during

normal operating protocol.

PBMC and SF Cell Sample Preparation: Peripheral blood mononuclear cells (PBMCs)

were isolated from whole blood over Ficoll-Paque PLUS (Fisherbrand). For SF samples,

a 200 µl aliquot was centrifuged at 1000 RCF for 10 minutes and stored at -80°C for

ELISA. The remainder of the SF was treated with bovine testes hyaluronidase (Sigma-

Adritch) according to manufacturer's SF clarification protocol. SF was then filtered

through a 70µm nylon mesh strainer (Fisher Scientific), diluted to 10 mL with PBS, and

centrifuged at 400 RCF for 10 minutes at room temperature. Supernatant was

discarded. SF cells (SFCs) and PBMCs were counted and then cryopreserved in 90%

FBS with 10% DMSO.

Transwell Migration Assay: Synovium (knee) or pulvinar (hip) acquired in the

operating room was transferred to the laboratory in PBS on ice. Whole synovium was

washed twice in PBS, then sterilely dissected into 4x4 mm segments, and washed

again. Segments were placed into 24-well transwell inserts with 5 µm pores (Corning), a

size that should allow monocyte and macrophage migration, but prevent fibroblast

migration (10). Bottom wells were treated with LPS (1, 10, or 100 µg/mL, Sigma-Aldrich)

or MCP-1 (25 ng or 250 ng/mL, Fischer Scientific) in DMEM supplemented with 10%

FBS and 1% penicillin/streptomycin. The inserts were then placed in the well, and

enough media to cover the tissue was placed in the insert (approximately 200-400 µl).

The transwell plates were incubated for 24 hours at 37°C and 5% CO_2. At day 1, 2, 3, 5,

and 7, changes to the synovium were compared to *in vivo* controls also obtained intra-

operatively (**Suppl. Figure 2**). H&E staining was performed to evaluate changes to the

synovial intimal and sub-intimal lining with MCP-1 and LPS treatment (**Suppl. Figure 3**)

Validation for tissue survival was performed out to 7 days with normoxic (21% oxygen)

and hyperoxic (50% oxygen) conditions and Caspace-3 immunohistochemistry (**Suppl.**

Figure 4). After incubation, the cells in the bottom well media were counted, and a

crystal violet assay was performed on cells adhered to the bottom of the well and the

underside of the transwell per standard protocol. Migratory cells in solution were

analyzed using flow cytometry.

Flow Cytometry Analysis: Samples were plated in a 96-well round-bottom plate for

single stain, unstained, patient test samples, and/or Fluorescence Minus One (FMO)

controls. Controls were plated at 2×10^5 cells per well, and patient test samples at 1-

2×10^6 cells per well. Antibodies are listed in **Suppl. Table 1**. For intracellular staining

(OPG, CD68, F11R, TREM2, ZO-1 and RANKL), cells were fixed and permeabilized

with Fixation Buffer and Intracellular Staining Perm Wash Buffer (BioLegend) or

Cytofix/Cytoperm Fixation/Permeabilization Solution Kit (BD Bioscences). Final flow

cytometry gating strategy for SFCs and PBMCs shown in **Suppl. Figure 1**. Flow

cytometry was performed on a Cytek Aurora cytometer (Bethesda, MD). Analysis was

performed using FlowJo (Ashland, OR) software.

Enzyme-linked Immunosorbent Assays (ELISA): Human TIMP-1 (RAB0466-1KT),

TGF-BETA (RAB0460-1KT), TRACP (RAB1755-1KT), OPG/TNFRSF11B (RAB0484-

1KT), IFN GAMMA (RAB0222-1KT), TNF-ALPHA (RAB0476-1KT), MMP-9 (RAB0372-

1KT) and BMP2 (RAB0028-1KT) ELISA kits were obtained from Millipore Sigma, and

sRANKL (MBS262624) kit from MyBioSource. For TIMP-1 and MMP-9, SF was diluted

1:500, for sRANKL dilution was 1:100, and all others were diluted 1:20. TGF-β1 was

activated and then neutralized per manufacturer's protocol. Plates were read using a

VERSAmax plate reader (Molecular Devices) and analyzed using MyAssays.com,

Microsoft Excel, and GraphPad Prism 9.4.1.

Cell Sorting: To prepare the highest quality sample for single cell RNA sequencing

(scRNA-seq), patients who had the highest percentage of viable, non-neutrophil SFCs

were selected, with 3 patients having septic arthritis, and 3 having inflammatory arthritis,

regardless of crystal status. As SF in non-pathologic states lacks sufficient cellularity for

scRNA-seq analysis, healthy patients were not included. SFCs were thawed in a 37ºC

water bath and diluted in 4 mL Fluorescence Activated Cell Sorting (FACS) buffer and

centrifuged at 4ºC at 1400 rpm. Supernatant was gently decanted, and cells were

resuspended in 50 µL of ice cold FACS buffer before staining 1:1000 with DAPI and 5

µl/reaction of: CD244/APC (Clone C1.7), CD11b/BV695 (Clone ICRF44), CD66b/FITC

(Clone G10F5) and CD56/PE (Clone 5.1H11) (BioLegend). Cells were incubated in the

dark at 4ºC for 30 minutes, washed once in ice cold FACS buffer, and resuspended in

50 µL of FACS buffer. Samples were then sorted on a Sony MA900 (San Jose, CA) with

100 µm sorting chip, sorting out CD66b$^+$ and DAPI positive cells, then sorting on

CD11b$^+$, CD56$^+$, or CD244$^+$ positive cells into tubes containing cold PBS with 1% BSA.

Cells were then counted for viability using trypan blue on a hemocytometer and

concentrated according to 10X Chromium 3' kit guidelines.

Single Cell RNA sequencing: Cells were delivered to the Sequencing Core where

RNA library generation was performed on a 10X Chromium Controller according to

manufacturer's guidelines. RNA libraries were then sent to NovoGene (Sacramento,

CA) for sequencing. Analysis was performed in RStudio with R v4.2.2 using Seurat (11),

ReactomeGSA (12) and EnhancedVolcano (13). MT-DNA percentage was limited to

15% during quality control analysis (14). The number of unique genes was set as 200 to

5000. A minimum of 3 cells were required to express each gene. Data was normalized

using a global-scaling normalization method per Seurat with a scale factor of 10,000

with log transformation, then the data was integrated into a single database. A

resolution of 0.4 was used to define clusters with dimensions set 1:30. Clusters were

annotated by identifying top 10 gene expression in addition to expression of markers

such as *CD56, CD206, F11R*, and *CD68*. Log2FC threshold was set to 0.5 for gene

expression. Analysis was performed comparing septic arthritis to inflammatory arthritis.

Adjusted p-values were used to determine significance of Differentially Expressed

Genes (DEGs), with Log2FC threshold of 1.5 given the homogenous sample. For

Conserved Markers, included genes had min.diff.pct set to 0.7, and min.pct to 0.25.

Statistics: Descriptive statistics were used to compare patient demographics and

underlying diagnoses. Statistical analysis was performed using GraphPad Prism v9.4.1.

Flow Cytometry data was tabulated in FlowJo, and 2-way ANOVA with Tukey's *post-hoc*

correction was used to compare %parent or %total cells of control, inflammatory, and

septic arthritis populations. Mann-Whitney U-tests were used to compare ELISA results

between the 3 arthritis groups (control, inflammatory, and septic). DEG and Conserved

gene analysis was performed in Seurat per individual cell cluster using adjusted p-

values.

Results:

Patient Demographics: To identify RSM in SF, 52 patients were recruited into the

initial study. Of these, 32% of patients were female and 87% were Caucasian (**Suppl.**

Table 2). SF was obtained from 36 of 52 patients. Eighteen patients were found to have

septic arthritis based on SF analysis and final culture results, and 8 had inflammatory

arthritis. Of the remainder, 10 were designated non-inflammatory (<2000 WBCs) or

normal (<200 WBCs) (**Suppl. Table 3**).

Identification of SF Macrophages Consistent with RSM: We identified a cell subset

expressing myeloid marker CD11b, pan-macrophage marker CD68, anti-inflammatory

macrophage marker TREM2 (15), Major Histocompatibility Complex marker Class II

HLA-DR (16), CX3CR1 (7), OPG (16), and hematopoietic/macrophage adhesion marker

CD45 (17). These cells were also negative for Receptor Activator of NF-kB

Ligand/RANKL (16) and for the dendritic cell (DC) marker CD11c. There were two

distinct subpopulations: a CD14^{hi} and a CD14^{dim} (**Figure 1A**). The CD45$^+$CD14^{dim} RSM-

like population was found to have an absolute decrease in frequency in pathologic

states compared to control SFCs (**Figure 1B**). However, it was also relatively enriched

in SF compared to the PBMC fraction (**Figure 1C**). The Alive/CD45$^+$CD14^{dim} was then

back-gated to better describe this population, first to evaluate the frequency of

macrophages by CD68 and TREM2, then if macrophages also co-expressed RSM

markers OPG and ZO-1 (**Figure 1D**). For patients with inflammatory arthritis, 1.96% of

the CD14^{dim} cells were macrophages, and of those macrophages, 34.35% were RSM-

like by OPG and ZO-1 expression. For patients with septic arthritis, 0.92% of CD14^{dim}

cells were macrophages, and of those 9.97% were RSM-like.

Identification of RSM-like Cells Migrating from Intact Synovium: We obtained

synovium from patients undergoing total joint replacement or resection arthroplasty and

placed tissue into transwells (**Figure 2A**). This model was validated out 24 hours to

have a dose-dependent loss of intimal lining cells with MCP-1 and LPS stimulation

(**Suppl. Figure 3**), and negligible apoptosis by Caspace-3 IHC at day 3 in standard

culture conditions (**Suppl. Figure 4**). At 24 hours, the migratory synovial myeloid

population (Live/CD56$^-$/CD3$^-$/CD11c$^-$/CD20$^-$/CD14$^+$/CD11b$^+$) in a representative patient

sample contained OPG+, TREM-2+, ZO-1+, CX3CR1+, and F11R+ populations in the

lower transwell media (**Figure 2B**). Sixty-five percent of CD11b+CD14^{dim} cells migrating

out of the synovium tissue were double positive for OPG and CX3CR1, and of those

cells, 93.8% were double positive for tight junction markers F11R and ZO-1. However,

we noted the CD14^{hi} macrophages displayed the greatest increase in the proposed

RSM markers (**Figure 2B**), and the relevance of CD14 dim versus high in the acute *ex*

vivo setting requires further clarification. To evaluate co-expression of these RSM-

specific markers, t-distributed Stochastic Neighbor Embedding (tSNE) plots were

created. Of the synovial myeloid population described above, there was weak or low

expression of all RSM-specific markers, but ZO-1 may be most specific to identifying

migratory RSM in SF (**Figure 2C, top right**). Cells with markers of circulating immune

subsets, including neutrophils (CD66b), NK cells (CD56), and T cells (CD3) were also

present in the lower transwell chambers (**Suppl. Figure 2**). Though there were no

significant differences found between treatments in this experiment, in all cases the

stimuli resulted in decreased RSM-like cell migration compared to control, indicating

these RSMs may remain active in the tissue, and migratory cells present in the control

may be in response to tissue trauma, an effect countered by the stimuli. Cells found

adhered to the underside of the transwell or in the bottom of the lower well were below

the limit of detection by Crystal Violet assay (data not shown). Therefore, we conclude

that RSMs can migrate out of the synovium, and ZO-1 is likely the most specific RSM

marker expressed by myeloid cells, but tight junction markers in conjunction with M2

markers and OPG are necessary for identification.

Evaluation of Pro and Anti-Resorptive Potential of SF: SF supernatant was

analyzed for cytokines implicated in bone and/or cartilage destruction. OPG is the decoy

receptor for RANKL, which in turn is required for the formation of osteoclasts; the ratio

between these two cytokines is an important method to evaluate bone maintenance

versus destruction (**Figure 3A**) (18). The collagenase Tartrate Resistant Acid

Phosphatase (TRAP), which is secreted by osteoclasts, had no observed change.

Transforming Growth Factor Beta (TGF-β) not only inhibits osteoclastogenesis (19), but

also stimulates osteoblasts (20) and further, it is stored in the latent phase within the

extracellular matrix to be released during bone turnover (21; 22). Likewise, there was no

significant difference between pathologic SF and control TGF-β levels (**Figure 3B**). We

also tested Bone Morphogenic Protein 2 (BMP2), TNF-α and IFN-γ, however all were

below the limit of detection (data not shown). Finally, Tissue Inhibitor of Matrix

Metalloproteinases 1 (TIMP1) and Matrix Metallopeptidase 9 (MMP9) were evaluated

(**Figure 3C**). TIMP1 is an inhibitor of MMPs. MMP9 is expressed by osteoclasts and is

an important enzyme for bone remodeling (23). There was an increase in MMP9 in

patients with inflammatory and septic arthritis, and a significantly decreased ratio of

TIMP1:MMP9 in these patients as well. Therefore, there is increased potential for bone

and cartilage damage in patients with inflammatory and septic arthritis due mainly to

increased MMP9, which is possibly due to increased osteoclastogenesis secondary to

increased RANKL, or increased synovial fibroblast production. As RSM-like cells were

decreased in SF of septic arthritis patients that also had highest RANKL and MMP9, it is

possible RSM provide a protective mechanism against bone and joint destruction. The

specific cytokine production profile of RSM specifically will require further evaluation.

Identification of Rare Cell Subsets Using scRNA-seq: We analyzed sorted myeloid

cells from patients with infectious and inflammatory arthritis for highly variable features,

demonstrating a relevant focus of M1 and M2 functions (**Suppl. Figure 6A**). Cell

subpopulations were clustered into 11 groups (**Figure 4A**) with manual annotation of

the clusters based on top ten gene expression (**Figure 4B**). There were insufficient cell

events to separately cluster NK and NKT cells, therefore they are represented in a

single, though spatially separate, cluster.

After assigning known and widely accepted monocyte/macrophage designations

based on gene profiles, clusters 9 and 10 remained. Cluster 10 expressed NK-marker

CD56, low *CD68* (**Suppl. Figure 6B**), while also having high *HLA* expression (**Figure

4B**). However, Cluster 10 did not express any granzymes. Therefore, we putatively

classified this cluster as cytotoxic DCs (24). Cluster 9 expressed *F11R* (7), *CD68,* and

M2-marker *IDO1* (25) (**Figure 4B, Suppl. Figure 6B**). It was also the only cluster with

OPG expression, though this was not significant. This is putatively consistent with the

RSM phenotype. *ZO-1* was not expressed in any cluster. We also evaluated expression

of resident macrophage transcription factor *GATA6* (26), which was minimally

expressed, but only in Cluster 3. This may indicate that RSMs are split between multiple

clusters, as Cluster 3 was designated as DCs based on high expression of HLA—which

is also consistent with sub-intimal RSM, but not intimal RSM (27). Based on PCA

analysis (**Suppl. Figure 6C**), cytotoxic DCs were more closely related to NK/NKT cells

than to RSM, and the cytotoxic DC and RSM populations may represent phases of

differentiation of the same cell of origin, as both were distinct from the circulating

monocyte/macrophage population. The plasticity of tissue macrophages, and ability to

survive in new compartments has been advocated and challenged; the exact lineage

remains unknown.

Differentially Expressed Genes (DEGs) were explored using a volcano plot

(**Suppl. Figure 6D**) and a Log2FC threshold of 1.5. DEGs were statistically significant in

Clusters 0-8 (**Figure 5A**). Transcripts highly upregulated in septic arthritis (and therefore

down-regulated in inflammatory arthritis) among multiple clusters included *HLA-DRB5,*

C15orf48, IL1B, AC025580.2, and *SOD2*. In inflammatory arthritis, common gene

upregulation included *PPARG, FABP5, CD36*, *NLRP3, SPP1, and MITF*. M1

macrophages from the two different arthritis etiologies showed a distinctly different gene

expression patterns with an upregulation of *GBP1, GBP5*, and *TIMP1.*

Conserved genes that remained highly expressed in both inflammatory and

infectious arthritis were also examined, as these could represent novel targets in the

treatment, or prevent the conversion of infectious to inflammatory arthritis. Conserved

genes were only significant in DC, NK/NKT, RSM, and Cytotoxic DC clusters (**Figure**

5B, Suppl. Table 4). Nearly all genes were associated with cytolytic function, antigen

presentation, M1/M2 polarity, or lysosomes.

Gene Set Analysis (GSA) was then assessed to determine the overall, broad

picture, of cell function and metabolism (**Figure 6**) with the 15 most upregulated or

downregulated pathways, which identified complement, bone-derived FGF23 signaling,

and COX signaling. Pathways specifically immune relevant, related to phagocytic

potential, complement associated and adhesion related (**Suppl. Figure 7**) were also

examined. In all cases except adhesion, Cytotoxic DC, NK/NKT, and RSMs populations

shared a similar pattern of activity, indicating functional overlap. GSA was further

performed specifically on the RSM cluster to identify the maximum changes in septic

and inflammatory arthritis (**Suppl. Figure 8**), which identified threonine, pyridoxine, and

thiamine metabolism.

Discussion:

RSMs are regulatory immune cells of the joint space that are historically

identified using tissue histology. We identified a rare subset of SF macrophages

identified by flow cytometry consistent with previously described RSMs, indicating

migratory capacity especially during pathologies that dysregulate tight junctions. The

role of these cells in SF has yet to be established.

Though the chemokine receptor CX3CR1 was found to be specific for murine

RSM (1), we were unable to distinguish SFCs from peripheral monocytes using this

marker. Likewise, HLA-DR expression was downregulated in the intimal-lining RSMs,

but upregulated in sub-lining RSMs (8), and it is unclear if HLA-DR expression would

assist in identification of RSM-like cells in the SF. We first identified putative RSMs as

CD14^{dim}OPG$^+$ M2-macrophages, and while CD14^{dim} monocytes have been described

previously as non-classical and poorly phagocytic (28; 29), data is limited on CD14^{dim}

macrophages. CD14^{dim} gingival macrophages were M2 and likely osteo-protective by

high expression of IL-10 and TGF-β in the setting of gingivitis (30). To fully elucidate the

utility of CX3CR1, HLA-DR, and CD14 in human SF macrophage subsets requires

further work.

To determine whether our identified cells came from synovium or from

circulation, we piloted a novel transwell migration assay with human *ex vivo* synovium

that validated our flow cytometry findings. Therefore, we believe the

CD68/TREM2/OPG/ZO-1/F11R myeloid cell fraction is the most representative of

putative RSMs. This explant model could be used widely to study other synovial

pathology.

We then utilized scRNA-seq to identify rare macrophage subpopulations.

Markers identified using flow cytometry were not always identified in gene transcripts,

including *ZO-1*. Instead we identified M2/Tumor Associated Macrophage markers such

as *IRF4* (31) and *IDO1* (32) (**Suppl. Figure 6B**). While the putative RSM cluster

resembled M2 macrophages, these cells also had a similar transcriptional signature to

inflammatory NK/NKT and Cytotoxic DCs, indicating that RSMs may be capable of

taking on an inflammatory and/or joint destructive phenotype in settings of chronic

inflammation. This was demonstrated by the pro-inflammatory expression of *CCR7*

(Log2FC 2.25) and *CD86* (Log2FC 0.92) in infectious settings.

To provide context to our findings, we compared our findings to scRNA-seq

performed on synovium by other groups. Human MERTK+CD206+ RSMs were anti-

inflammatory in patients in remission from rheumatoid arthritis (RA) (27). We found

MERTK expressed highest in Cluster 5/M2 Monocytes (Log2FC 0.94), but it was not

expressed in 9/RSM. Interestingly, MERTK-CD206- RSM indicated active RA (27). As all

patients who received arthrocentesis and participated in the scRNA-seq were acutely

symptomatic, a MERTK+CD206+ RSM profile was likely physiologically improbable. For

CD206, this was expressed highest in Cluster 4/M2 Macrophages, which additionally

expressed *TREM2, FOLR2*, and *LYVE1* (27), though only *TREM2* had Log2FC >0.5.

Once RSMs exit synovial tissue to enter the SF, previously established profiles may no

longer apply, and RSMs may be distributed amongst clusters rather than a discrete

cluster.

Others found that RSMs arose from *CSF1R+* interstitial macrophages (7). We

found *CSF1R* expressed in clusters 3-6, and 8-10. It was also found that interstitial RSM

expressed *RETNLA, STMN1*, and *AQP1* (7). *RETNLA* and *AQP1* transcripts were not

identified in this study, while *STMN1* was only expressed by Cluster 3/DCs, but this was

not significant. Likewise, markers of tight junctions and cell polarity previously found

included *F11R, CLDN5, FAT4*, and *VANGL2*. Of these, only *F11R* was identified, and

expressed mainly in RSM and Cytotoxic DC clusters (**Suppl. Figure 6B**).

We identified *LAMP3/CD208/DC-LAMP* as the top gene expressed by Cluster

9/RSM, and while the exact function of LAMP3 has yet to be elucidated, it is likely to be

involved with MHC Class II peptide presentation (34). *LAMP3* is also traditionally

considered a marker of mature DCs. DCs expressing LAMP3 are regulatory in nature

and more enriched in draining lymph nodes rather than tumors (35). However, *LAMP3*

is upregulated by THP-1 macrophages with *in vitro* LPS stimulation (36), and is

constitutively expressed in primary macrophages in multiple species (37).

Further analysis of Cluster 9/RSM gene expression revealed high expression of *ENOX1*

(**Figure 4**), involved with reduction of oxygen to superoxide (38). Likewise, *CERS6*

contributes to mitochondrial dysfunction by promoting reactive oxygen species

production in hepatocytes (39). *IDO1* expression in macrophages has been associated

with increased tumor immune cell infiltration (40) and tryptophan metabolism, the

metabolites of which inhibit oxidative cell death (41). Together this indicates that these

cells are likely to have potent generation of superoxide with inhibition of apoptosis,

which may indicate perpetuation of chronic inflammation. Until these putative RSM can

be compared to similar cells from healthy SF, the baseline role and function are unclear.

Given we believe these cells are capable of migratory function, markers of

migration and extravasation were also examined. *CADM1* was highly and

conservatively expressed (**Figure 5**) and is strongly associated with TREM2+ tumor

associated macrophages (42). *ALCAM* (Log2FC 0.73) is expressed by endothelial cells

of the blood-brain barrier and migrating monocytes (43), which may be consistent with

the relative immune privilege of the joint space. As ALCAM stabilizes tight junctions

(44), this integrin may be important in the homeostasis of the joint space as maintained

by RSM. *PECAM1* (Log2FC -1.03) assists with leukocyte migration through tight

junctions (45). Our data seems to suggest a differential regulation of tight junctions by

macrophages depending on pathology.

As there were no DEGs identified in Cluster 9/RSM, GSA was performed to

identify unique pathways and discovered threonine, pyridoxine, and thiamine

metabolism (**Suppl. Figure 8**). The role of threonine catabolism is unclear in

macrophages but is critically necessary for murine stem cell viability (46), which may be

similar function to locally renewing macrophage populations. Pyridoxine suppresses IL-

1β release through NLRP3 inhibition (47), while thiamine precursors have been found to

both increase cellular glutathione stores and inhibit NF-kB translocation to the nucleus

in microglial cells (48), the resident macrophages of the brain. Putative RSMs are

upregulating anti-inflammatory pathways, but the question remains if the concurrent

upregulation of complement and superoxide transcripts may supersede the protective

mechanisms in place through B vitamins.

We also evaluated the protein levels of multiple bone-relevant cytokines and

related this to cell populations in the joint space. The DEG *TIMP1* was identified in

Cluster 7/M1 Macrophage as highly upregulated in septic arthritis, with concurrent

downregulation in inflammatory arthritis. As a 1:1 inhibitor of MMP9 (49), TIMP1 is likely

protective in the setting of joint inflammation by preserving the extracellular matrix of

cartilage and bone from enzymatic degradation. We observed a decrease in the

stoichiometric ratio of TIMP1:MMP9 protein concentration in both infectious and

inflammatory arthritis SF, which is concerning for MMP9 as a major cause of bone

destruction in these pathologic states, and a potential therapeutic target. *MMP9* was not

differentially expressed, and MMP9 found by ELISA is likely from other cells, specifically

fibroblast like synoviocytes and/or neutrophils (50), but possibly also osteoclasts (23).

TIMP1 was expressed in Cluster 9/RSM (Log2FC -2.67) indicating RSM would not be

the primary source of this in SF.

In conclusion, the profile of a subset of M2, likely osteoprotective macrophages in

SF that can be stimulated to migrate out of synovial tissue *ex vivo* suggests these are

equivalent to tissue resident macrophages. However, these putative RSM expressed

transcripts heavily involved with antigen presentation (*LAMP3*), oxidative stress

(*ENOX1, CERS6, IDO1*), and cell migration (*ALCAM, PECAM*). This suggests that in

settings of infectious or inflammatory arthritis, these cells may perpetuate rather than

attenuate inflammation, and could be involved in the transition to chronic symptomology

once out of the normal synovial tissue niche. Further work will focus on the cytokine

expression of these putative migratory RSM, and the interactions of RSM with other

cells in the joint space, including T cells, B cells, and NK or NKT cells.

Acknowledgements:

The authors would like to thank Catherine Fairfield BSN, EM Research Coordinator, and

Alex Peebles, Cameron Williams, Shannon Landers, Klaudia Golebiewski, Allison Herr,

Noble Briggs, Vanko Bicar, Malea Pinckney, Heath Gibbs, Nicole Grossmann, Scott

Tibbetts, Ike Appleton, Jay Miller, and Jacob Hampton with the Emergency Department

Research Enroller Program (ED-REP) for their essential role in enrolling patients and

collecting data for this project.

The authors would also like to thank Dr. Daniel Livorsi and the Division of Infectious

Diseases for assistance with patient recruitment.

The data presented herein were obtained at the Flow Cytometry Facility, which is a

Carver College of Medicine / Holden Comprehensive Cancer Center core research

facility at the University of Iowa. The facility is funded through user fees and the

generous financial support of the Carver College of Medicine, Holden Comprehensive

Cancer Center, and Iowa City Veteran's Administration Medical Center. Research

reported in this publication was supported by: the National Center for Research

Resources of the National Institutes of Health under Award Number 1 S10 OD034193-

01; and the National Cancer Institute of the National Institutes of Health under Award

Number P30CA086862

References

1. *Origin and function of synovial macrophage subsets during inflammatory joint disease.* **Culemann S,**
Grüneboom A, Krönke G. 2019, Adv Immunol, Vol. 143, pp. 75-98.

2. *Immune cell profiles in synovial fluid after anterior cruciate ligament and meniscus injuries.* **Kim-**
Wang, S.Y., Holt, A.G., McGowan, A.M. et al. 280, 2021, Arthritis Res Ther, Vol. 23.

3. *Synovial fluid monocyte/macrophage subsets and their correlation to patient-reported outcomes in*
osteoarthritic patients: a cohort study. **Gómez-Aristizábal, A., Gandhi, R., Mahomed, N.N. et al.** 26,
2019, Arthritis Res Ther, Vol. 21.

4. *High percentages and activity of synovial fluid NK cells present in patients with advanced stage active*
Rheumatoid Arthritis. **Yamin R, Berhani O, Peleg H, Aamar S, Stein N, Gamliel M, Hindi I, Scheiman-**
Elazary A, Gur C. 1, 2019, Sci Rep, Vol. 9, p. 1351.

5. *Resident Macrophages Cloak Tissue Microlesions to Prevent Neutrophil-Driven Inflammatory Damage.*
Stefan Uderhardt, Andrew J. Martins, John S. Tsang, Tim Lämmermann, Ronald N. Germain,. 3, 2019,
Cell, Vol. 177, pp. 541-55.

6. *Fibroblast-like synoviocytes: key effector cells in rheumatoid arthritis.* **Bartok B, Firestein GS.** 1, 2010,
Immunol Rev, Vol. 233, pp. 233-55.

7. *Locally renewing resident synovial macrophages provide a protective barrier for the joint.* **S,**
Culemann, et al. 7771, 2019, Nature, Vol. 572, pp. 670-675.

8. *Critical Role of Synovial Tissue-Resident Macrophage and Fibroblast Subsets in the Persistence of Joint*
Inflammation. **Kemble S, Croft AP.** 2021, Front Immunol.

9. *Synovial Fluid Tests: What Should Be Ordered?* **Shmerling RH, Delbanco TL, Tosteson ANA, Trentham**
DE. 8, 1990, JAMA, Vol. 264, pp. 1009-1014.

10. **Corning Incorporated.** Transwell Permeable Supports Selection and Use Guide. 2013.

11. *Comprehensive Integration of Single-Cell Data.* **Stuart T, Butler A, Hoffman P, Hafemeister C,**
Papalexi E, III WMM, Hao Y, Stoeckius M, Smibert P, Satija R. 2019, Cell, Vol. 177, pp. 1888-1902.

12. *ReactomeGSA - Efficient Multi-Omics Comparative Pathway Analysis.* **Griss J, Viteri G, Sidiropoulos**
K, Nguyen V, Fabregat A, Hermjakob H. 2020, bioRxiv.

13. **Blighe K, Rana S, Lewis M.** EnhancedVolcano: Publication-ready volcano plots with enhanced
colouring and labeling. [Online] https://github.com/kevinblighe/EnhancedVolcano.

14. *Systematic determination of the mitochondrial proportion in human and mice tissues for single-cell*
RNA-sequencing data quality control. **Osorio D, Cai JJ.** 7, 2021, Bioinformatics, Vol. 37, pp. 963-967.

15. *Trem-2 Promotes Emergence of Restorative Macrophages and Endothelial Cells During Recovery*
From Hepatic Tissue Damage. **Coelho Inês, Duarte Nádia, Barros André, Macedo Maria Paula, Penha-**
Gonçalves Carlos. 2021, Frontiers in Immunology , Vol. 11.

16. *Synovial tissue macrophages: friend or foe?* **Kurowska-Stolarska M, Alivernini S.** 2017, RMD Open,
Vol. 3, p. e000527.

17. *CD45 regulates Src family member kinase activity associated with macrophage integrin-mediated*
adhesion. **Tamara Roach, Suzanne Salter, Michael Koval et al.** 6, 1997, Current Biology, Vol. 7, pp. 408-
17.

18. *Biology of RANK, RANKL, and osteoprotegerin.* **Boyce BF, Xing L.** Suppl 1, 2007, Arthritis Res Ther,
Vol. 9.

19. *Interferon-Gamma-Mediated Osteoimmunology.* **Tang M, Tian L, Luo G, Yu X.** 9, 2018, Front
Immunol, Vol. 29, p. 1508.

20. *Role of transforming growth factor-beta in bone remodeling.* **Bonewald LF, Mundy GR.** 250, 1990,
Clin Orthop Relat Res, pp. 261-76.

21. *The extracellular matrix and transforming growth factor-β1: Tale of a strained relationship.* **Hinz,**
Boris. 2015, Matrix Biology, Vol. 47, pp. 54-65.

22. *TGF-β and BMP signaling in osteoblast, skeletal development, and bone formation, homeostasis and*
disease. **Wu M, Chen G, Li YP.** 4, 2016, Bone Res, Vol. 26, p. 16009.

23. *Matrix metalloproteinases in bone development and pathology: current knowledge and potential*
clinical utility. **Liang HPH, Xu J, Xue M, Jackson CJ.** 2016, Metalloproteinases In Medicine, Vol. 3, pp. 93-
102.

24. *CD56 marks human dendritic cell subsets with cytotoxic potential.* **Roothans D, Smits E, Lion E, Tel J,**
Anguille S. 2, 2013, Oncoimmunology, Vol. 2, p. e23037.

25. *Critical role of synovial tissue-resident macrophage niche in joint homeostasis and suppression of*
chronic inflammation. **Huang QQ, Doyle R, Chen SY, Sheng Q, Misharin AV, Mao Q, Winter DR, Pope**
RM. 2, 2021, Sci Adv, Vol. 7, p. eabd0515.

26. *GATA6+ Peritoneal Resident Macrophage: The Immune Custodian in the Peritoneal Cavity.*
Jayakumar P, Laganson A, Deng M. 866993, 2022, Front Pharmacol, Vol. 13.

27. *Distinct synovial tissue macrophage subsets regulate inflammation and remission in rheumatoid*
arthritis. . **Alivernini, S., MacDonald, L., Elmesmari, A. et al.** 2020, Nat Med, Vol. 26, pp. 1295-1306.

28. *Human Monocyte Subsets and Phenotypes in Major Chronic Inflammatory Diseases.* **Kapellos**
Theodore S., Bonaguro Lorenzo, Gemünd Ioanna, Reusch Nico, Saglam Adem, Hinkley Emily R.,
Schultze Joachim L. 2019, Frontiers in Immunology , Vol. 10.

29. *Human CD14dim monocytes patrol and sense nucleic acids and viruses via TLR7 and TLR8 receptors.*
Cros J, Cagnard N, Woollard K, Patey N, Zhang SY, Senechal B, Puel A, Biswas SK, Moshous D, Picard C,
Jais JP, D'Cruz D, Casanova JL, Trouillet C, Geissmann F. 3, 2010, Immunity, Vol. 33, pp. 375-86.

30. *Macrophage subset sensitivity to endotoxin tolerisation by Porphyromonas gingivalis.* **Foey AD,**
Crean S. 7, 2013, PLoS One, Vol. 8, p. e67955.

31. *Interferon-regulatory factors determine macrophage phenotype polarization.* **Günthner R, Anders**
HJ. 2013, Mediators Inflamm, p. 731023.

32. *The role of indoleamine 2,3-dioxygenase (IDO) in immune tolerance: focus on macrophage*
polarization of THP-1 cells. **Wang XF, Wang HS, Wang H, Zhang F, Wang KF, Guo Q, Zhang G, Cai SH, Du**
J. 1-2, 2014, Cell Immunol, Vol. 289, pp. 42-8.

33. *TGF-β induces M2-like macrophage polarization via SNAIL-mediated suppression of a pro-*
inflammatory phenotype. **Zhang F, Wang H, Wang X, Jiang G, Liu H, Zhang G, Wang H, Fang R, Bu X, Cai**
S, Du J. 32, 2016, Oncotarget, Vol. 7, pp. 52294-52306.

34. *Vitamin D3 regulates LAMP3 expression in monocyte derived dendritic cells.* **L. Malaguarnera, A.**
Marsullo, K. Zorena, G. Musumeci, M. Di Rosa. 2017, Cellular Immunology, Vol. 311, pp. 13-21.

35. *Mature dendritic cells enriched in immunoregulatory molecules (mregDCs): A novel population in the*
tumour microenvironment and immunotherapy target. **Li J, Zhou J, Huang H, Jiang J, Zhang T, Ni C.** 2,
2023, Clin Transl Med, Vol. 13, p. e1199.

36. *LAMP-3 (Lysosome-Associated Membrane Protein 3) Promotes the Intracellular Proliferation of*
Salmonella typhimurium. **Lee EJ, Park KS, Jeon IS, Choi JW, Lee SJ, Choy HE, Song KD, Lee HK, Choi JK.** 7,
2016, Mol Cells, Vol. 39, pp. 566-72.

37. *Characterisation and expression analysis of the rainbow trout (Oncorhynchus mykiss) homologue of*
the human dendritic cell marker CD208/lysosomal associated membrane protein 3,. **Petronella**
Johansson, Yolanda Corripio-Miyar, Tiehui Wang, Bertrand Collet, Chris J. Secombes, Jun Zou. 3-4,
2012, Developmental & Comparative Immunology, Vol. 37, pp. 402-413.

38. *Increased Expression of Ecto-NOX Disulfide-thiol Exchanger 1 (ENOX1) in Diabetic Mice Retina and its*
Involvement in Diabetic Retinopathy Development. **YU-CHUEN HUANG, SHIH-PING LIU, SHIH-YIN CHEN,**

JANE-MING LIN, HUI-JU LIN, YU-JIE LEI, YEH-HAN WANG, WAN-TING HUANG, WEN-LING LIAO, FUU-JEN TSAI. 6, 2019, In Vivo, Vol. 33, pp. 1801-6.

39. *CerS6 triggered by high glucose activating the TLR4/IKKβ pathway regulates ferroptosis of LO2 cells through mitochondrial oxidative stress.* **Li D, Tian L, Nan P, Zhang J, Zheng Y, Jia X, Gong Y, Wu Z.** 2023, Mol Cell Endocrinol, Vol. 572, p. 111969.

40. *Evaluating the role of IDO1 macrophages in immunotherapy using scRNA-seq and bulk-seq in colorectal cancer.* **Liu Xingwu, Yan Guanyu, Xu Boyang, Yu Han, An Yue, Sun Mingjun.** 2022, Frontiers in Immunology , Vol. 13.

41. *IL4i1 and IDO1: Oxidases that control a tryptophan metabolic nexus in cancer.* **Zeitler L, Murray PJ.** 6, 2023, J Biol Chem, Vol. 299, p. 104827.

42. *Tissue-resident FOLR2+ macrophages associate with CD8+ T cell infiltration in human breast cancer.* **Ramos, RN, Yoann Missolo-Koussou, Yohan Gerber-Ferder, et al.** 7, 2022, Cell, Vol. 185, pp. 1189-1207.

43. *ALCAM (CD166) is involved in extravasation of monocytes rather than T cells across the blood-brain barrier.* **Lyck R, Lécuyer MA, Abadier M, Wyss CB, Matti C, Rosito M, Enzmann G, Zeis T, Michel L, García Martín AB, Sallusto F, Gosselet F, Deutsch U, Weiner JA, Schaeren-Wiemers N, Prat A, Engelhardt B.** 8, 2017, J Cereb Blood Flow Metab, Vol. 37, pp. 2894-2909.

44. *Dual role of ALCAM in neuroinflammation and blood-brain barrier homeostasis.* **Lécuyer MA, Saint-Laurent O, Bourbonnière L, Larouche S, Larochelle C, Michel L, Charabati M, Abadier M, Zandee S, Haghayegh Jahromi N, Gowing E, Pittet C, Lyck R, Engelhardt B, Prat A.** 4, 2017, Proc Natl Acad Sci U S A, Vol. 114, pp. E524-E533.

45. *PECAM-1: a multi-functional molecule in inflammation and vascular biology.* **Woodfin A, Voisin MB, Nourshargh S.** 12, 2007, Arterioscler Thromb Vasc Biol, Vol. 27, pp. 2514-23.

46. *Dependence of Mouse Embryonic Stem Cells on Threonine Catabolism.* **al, Jian Wang et.** 2009, Science, Vol. 325, pp. 435-439.

47. *Vitamin B6 Prevents IL-1β Protein Production by Inhibiting NLRP3 Inflammasome Activation.* **Zhang P, Tsuchiya K, Kinoshita T, Kushiyama H, Suidasari S, Hatakeyama M, Imura H, Kato N, Suda T.** 47, 2016, J Biol Chem, Vol. 291, pp. 24517-24527.

48. *Benfotiamine Upregulates Antioxidative System in Activated BV-2 Microglia Cells.* **Bozic I., Savic D., Stevanovic I., Pekovic S., Nedeljkovic N., Lavrnja I.** 9, 2015, Front. Cell. Neurosci, p. 351.

49. *Structural characterisation of inhibitory and non-inhibitory MMP-9–TIMP-1 complexes and implications for regulatory mechanisms of MMP-9.* **Charzewski, Ł., Krzyśko, K.A. & Lesyng, B.** 2021, Sci Rep, Vol. 11, p. 13376.

50. *Matrix metalloproteinase-9: Many shades of function in cardiovascular disease.* **Yabluchanskiy A, Ma Y, Iyer RP, Hall ME, Lindsey ML.** 6, 2013, Physiology (Bethesda), Vol. 28, pp. 391-403.

610   **Figures:**

611
612

Figure 1: Resident Synovial Macrophage-like cells. Live, CD56$^-$CD3$^-$CD20$^-$CD11c$^-$
TREM2$^+$OPG$^+$CD68$^+$CD11b$^+$HLA-DR$^+$CX3CR1$^+$ cells were identified with an enrichment
of CD14^{dim} cells in the SF compared to PBMCs (**A**, n=6 patients for preliminary
evaluation for RSM cells). These CD14^{dim} macrophages were decreased in absolute
frequency in inflammatory and septic arthritis compared to controls (**B**, n=22 patients, 5-
12 patients per group) but were relatively enriched in the SF of pathologic joints when
SFCs were compared to PBMC, shown by ratio >1 (**C**, n=22 patients, 5-12 patients per
group). Back-gating on the Alive/CD14^{dim} population to identify M2 macrophages (D,
inset), 78.6% were double positive for OPG and ZO-1 (**D**, representative patient with
inflammatory arthritis). Two-way ANOVA with Tukey's *post-hoc* correction.
623

Figure 2: Transwell synovial cell migration assay. Schematic of the experimental set up (**A**). Resident synovial macrophage markers TREM-2+, OPG+, CX3CR1+, ZO-1+ and F11R+ were evaluated within the migratory monocyte/macrophage population (Alive/CD56-/CD3-/CD11c-/CD20-/CD14+/CD11b+) in a representative patient treated with 250 ng/mL of MCP-1 (**B**). tSNE plots of the same patient's migratory myeloid cells demonstrating ZO-1 has the most delineation from other markers (**C**).

634
635
636 **Figure 3**: ELISA results of SF supernatant protein concentration. OPG, sRANKL, and
637 ratio of OPG:sRANKL (A). TRAP and TGF-β1, (B). Measures of TIMP1, MMP9, or ratio
638 of TIMP1:MMP9. Normal (n=9), inflammatory (n=6), and septic (n=11). Multiple Mann-
639 Whitney tests with FDR rate <0.05.
640

Figure 4: Unsupervised Cluster Analysis showing composite of 3 patients with inflammatory arthritis (A). Clusters were manually identified based on top ten expressed genes (B) in addition to classical markers. Analysis performed in R with resolution of 0.4 and dimensions 1:30. N = 3 patients per inflammatory and septic arthritis.

646
647
Figure 5: Differentially Expressed Genes (DEGs) (**A**). Conserved genes where Log2FC
> 1.5 and the change in percent expression in both septic arthritis and inflammatory
arthritis was > 0.7 in each cluster compared to all other clusters. Gene names in •**bold**
represent genes that were also in the top 10 expressed genes in **Figure 4**.
652

653
654 **Figure 6:** Top 15 most up or downregulated pathways using DEGs and ReactomeGSA.
655

Supplementary Table 1: Flow cytometry panel, initial (top) and modified (bottom). Mac = macrophage; Mono = monocyte; OB = osteoblast; RSM = Resident Synovial Macrophage; APC = Antigen Presenting Cell

Molecular Target	Fluorochrome	Clone	Vendor	Cell of Interest
CD11b	BV605	ICRF44	BioLegend	Myeloid
CD11c	BV785	3.9	BioLegend	Dendritic Cell
CD14	PE-Cy7	63D3	BioLegend	Mac/Mono
CD20	BV785	2H7	BioLegend	B cell
CD3	BV650	UCHT1	BioLegend	T Cell
CD45	AF700	2D1	BioLegend	Leukocytes
CD56	PE-Cy5	5.1H11	BioLegend	NK cell
CD68	PerCP-Cy5.5	Y1/82A	BioLegend	Macrophage
CX3CR1	BV421	2A9-1	BioLegend	RSM/Mac/Mono
HLA-DR	PE	Tu36	BioLegend	Activated APC
OPG/TNFRSF11B	AF488	155321	R&D Systems	RSM
RANKL/TNFSF11	AF350	685857	R&D Systems	OB/activated T cells
TREM2	APC	237920	R&D Systems	Macrophage
Viability	Live/Dead Aqua		Thermofisher	

Molecular Target	Fluorochrome	Clone	Vendor	Cell of Interest
CD11b	BV605	ICRF44	BioLegend	Myeloid
CD11c	BV785	3.9	BD Biosciences	Dendritic Cell
CD14	Spark Blue 550	63D3	BioLegend	Mac/Mono
CD20	BV785	2H7	BD Biosciences	B cell
CD3	BV510	SK7	BioLegend	T Cell
CD45	AF700	HI30	BioLegend	Leukocytes
CD56	PE-Cy5	5.1H11	BioLegend	NK cell
CD66b	PE	MIH24	BioLegend	Neutrophil
CD68	R718	Y1/82A	BD Biosciences	Macrophage
CX3CR1	BV711	2A9-1	BioLegend	RSM/Mac/Mono
F11R/JAM-1/JAM-A	BV421	M.Ab.F11	BD Biosciences	RSM
OPG/TNFRSF11B	AF488	155321	R&D Systems	RSM
TREM2	APC	237920	R&D Systems	Macrophage
ZO-1/TJP1	Coralite 594	polyclonal	ThermoFisher	RSM
Viability	Live/Dead Blue		ThermoFisher	

673 **Supplemental Table 2: Patient Demography.** Arthritis Type was determined by
674 synovial fluid WBC count per mm^3, %PMNs, and/or culture results. * = 3 patients had
675 Lyme Arthritis, 1 of which had both a *S. aureus* and positive IgG with confirmatory
676 Western Blot.

Demographics	n=52	
Age	(years)	
	Mean (STD)	51.7 (21)
	Median	58
	Range	3-77
Gender	n, (%)	
	Male	35 (68)
	Female	17 (32)
Race and Ethnicity	n, (%); Hispanic n, (%)	
	White	45 (87); 1 (2)
	Black	3 (6); 0
	Asian	1 (2); 0
	Multiracial	1 (2); 0
	Unknown	1 (2); 0
Arthritis Type	n (%)	
	Normal/Non-I	9 (25)
	Inflammatory	8 (22)
	Septic*	18 (50)
	Hemorrhagic	1 (3)

681 **Supplementary Table 3**: Patient Diagnosis. SF analysis was performed on patients
682 when it was clinically indicated. SF diagnosis was based on WBC count (# nucleated
683 cells) and % PMN based on neutrophil count. * = not performed. ** = Lyme Arthritis in
684 addition to other bacterial arthritis.

Patient	Color	Clarity	Crystals	# Nucleated Cells	# Neutrophils	# Lymphocytes	Arthritis Type
2	Red	Turbid	CCP, MSU	29783	22635	101	Inflammatory
3	Yellow	Turbid	CPP	4716	2453	20	Inflammatory
5	Red	Turbid	none	822	674	107	Non-Inflammatory
6	Yellow	Turbid	none	29497	28022	0	Septic
7	Yellow	Hazy	none	714	29	64	Non-Inflammatory
9	Yellow	Hazy	none	63726	60539	1275	Septic
10	Yellow	Slightly Hazy	none	464	28	70	Non-Inflammatory
11	Red	Turbid	none	671	658	0	Non-Inflammatory
13	Orange	Hazy	none	831	632	58	Non-Inflammatory
14	None	Clear	none	29	0	7	Normal
15	Orange	Turbid	none	63184	60025	632	Inflammatory
16	Pink	Hazy	none	166627	128303	9998	Septic
18	Amber	Turbid	none	134280	119509	2686	Septic
19	Red	Turbid	none	11522	8987	1959	Inflammatory
21	N/A	N/A	none	N/A	N/A	N/A	Septic
22	Yellow	Turbid	CPP	34482	27586	2	Inflammatory
23	Yellow	Clear	none	348	28	52	Non-Inflammatory
24	Red	Turbid	none	1819	1564	14	Hemorrhagic
26	Yellow	Clear	none	26	5	12	Normal
27	Orange	Turbid	MSU	33805	26030	0	Inflammatory
28	Brown	Turbid	none	66945	*	*	Septic
30	Yellow	Slightly Hazy	none	26536	25209	531	Septic
31	Red	Hazy	none	48363	46428	484	Septic
32	Orange	Hazy	none	43355	42054	434	Septic
33	Red	Turbid	none	304	76	85	Septic
34	Red	Hazy	none	854	384	154	Non-Inflammatory
38	Red	Turbid	none	114000	1000320	2	Septic
39	Red	Turbid	MSU	76890	76121	0	Inflammatory
40	Pale Yellow	Turbid	CPP	67368	64673	0	Inflammatory
41	Pink	Turbid	None	79118	75162	0	Septic**
43	Orange	Turbid	None	104595	92044	2	Septic
46	Pink	Turbid	None	80562	74117	2417	Septic
47	Red	Turbid	None	9884	8303	692	Septic
48	Orange	Hazy	None	130	83	26	Normal
49	Yellow	Hazy	MSU	20566	18715	0	Inflammatory

Supplementary Table 4: List of all significant conserved genes in RSM, Cytotoxic DC, and NK/NKT clusters.

Gene	Log2FC	Δ % Septic	Δ % Inflam	Cluster	Gene Function
PKIB	2.10657	0.71	0.726	DC	Also known as AKT, causes M2 formation
SKAP1	3.16933	0.89	0.954	NK/NKT	TCR adaptor protein
CD96	2.690931	0.77	0.874	NK/NKT	Inhibitory, marker of exhaustion
CD2	2.571958	0.78	0.817	NK/NKT	Regulates lytic activity
CD247	2.473159	0.79	0.869	NK/NKT	Subunit of T cell antigen receptor complex
ETS1	2.430819	0.85	0.88	NK/NKT	Required for NK differentiation and cytotoxic function
SLC38A1	2.222173	0.82	0.77	NK/NKT	Glutamine transporter
IDO1	3.679276	0.93	0.764	Cytotoxic DC	Immunosuppressive, expressed by DCs, macs, and epithelial cells
C1orf54	3.555745	0.94	0.794	Cytotoxic DC	cDC1 gene
CCSER1	3.110627	0.78	0.818	Cytotoxic DC	upregulated in anti-inflammatory conditions
CLEC9A	3.009585	1.00	0.888	Cytotoxic DC	DC receptor for necrotic cell death
CPNE3	2.908024	0.82	0.76	Cytotoxic DC	cDC1 gene
AUTS2	2.836438	0.78	0.74	Cytotoxic DC	associated with autism
CADM1	2.766723	0.99	0.936	Cytotoxic DC	expressed by pre-cDC1, cell adhesion
NEGR1	2.54191	0.97	0.798	Cytotoxic DC	immunoglobulin superfamily cell adhesion molecule subgroup IgLON, has been implicated in neuronal growth and connectivity
CLNK	2.441584	0.99	0.885	Cytotoxic DC	upregulated in response to IL-2 and IL-3, associated with SLP-76
DNASE1L3	2.326661	0.89	0.998	Cytotoxic DC	Cytokine secretion following inflammasome activation in response to DNA in circulating apoptotic bodies
ARL4C	2.315802	0.71	0.709	Cytotoxic DC	oncogene, tubulogenesis, wnt-beta catenin signaling
NAAA	2.166039	0.71	0.78	Cytotoxic DC	inflammatory, lysosomal
KIF16B	1.944567	0.79	0.778	Cytotoxic DC	microtubule formation associated with endosomes and cross presentation
DST	1.91132	0.77	0.764	Cytotoxic DC	junctional adhesion protein
PLPP1	1.872874	0.74	0.736	Cytotoxic DC	phospholipid lipase associated with mac inflammation, cDC1 marker
MCOLN2	1.841604	0.73	0.824	Cytotoxic DC	MHCII presentation, innate immune cell activation, enhances infectivity of certain viruses, interferon stimulating gene
CALCRL	1.840268	0.76	0.951	Cytotoxic DC	regulates DC function through NF-kB

Gene	Value	A	B	Type	Description
TACSTD2	1.765963	0.78	0.989	Cytotoxic DC	Marker for TGF-B1 dependent DCs
ZEB1	1.672937	0.74	0.941	Cytotoxic DC	cDC1 cells
C1orf21	1.647088	0.84	0.9	Cytotoxic DC	enriched in cytotoxic CD8 T cells
FLT3	1.645133	0.96	0.895	Cytotoxic DC	The growth factor Flt3 ligand (Flt3L) is central to dendritic cell (DC) homeostasis and development, controlling survival and expansion by binding to Flt3 receptor tyrosine kinase on the surface of DCs.
ENPP1	1.548458	1.00	0.777	Cytotoxic DC	suppresses innate immune response,
NUBPL	1.540362	0.83	0.911	Cytotoxic DC	mitochondrial gene, iron and sulfur
XCR1	1.53506	0.78	0.995	Cytotoxic DC	DC, induces CD8 responses
DAPP1	1.52997	0.72	0.717	Cytotoxic DC	immune related mucosal tissues, hyper-reactive airway disease, IDO1/CCR7 signaling, associated with DCs
LAMP3	3.181144	0.88	0.967	RSM	lysosome
CST7	2.710421	0.93	0.748	RSM	Cistatin F, lysosomal cathepsin inhibitor
DAPP1	2.225168	0.72	0.781	RSM	immune related mucosal tissues, hyper-reactive airway disease, IDO1/CCR7 signaling, associated with DCs
GPR157	2.163834	0.75	0.897	RSM	G protein coupled receptor 157
TBC1D4	2.161417	0.79	0.941	RSM	Rab GTPase Activating Protein
RFTN1	2.085313	0.77	0.736	RSM	glucose transport
CCR7	2.083676	0.92	0.837	RSM	M1 polarization
LAD1	1.96187	0.77	0.713	RSM	epithelialization
SLC41A2	1.415962	0.76	0.78	RSM	magnesium transporter
REPIN1	1.325367	0.76	0.72	RSM	increasing cell size and tumor metastasis
SPIB	1.005641	0.75	0.726	RSM	recruits TAMs via CCL4 signaling

Supplemental Figure 1: Final gating strategy to identify TREM2$^+$CX3CR1$^+$OPG$^+$F11R$^+$ZO-1$^+$ macrophages consistent with RSM. FMO and single stain (SS) shown where used to define gates.

Supplemental Figure 2: *In vivo* negative (A, C, same patient, 10X and 40X respectively) and positive (B, D, second patient, 10X and 40X respectively) controls. Negative control patient displays fibrosis, but positive control (chronic bacterial infection) patient has evident dysfunction of synovium with cartilage formation, evident migration of inflammatory cells into the sublining, and thinning/loss of cellularity of the initial lining (B, D).

Supplementary Figure 3: disruption of the synovial intimal lining and migration of inflammatory cells at 24 hours is dependent upon stimulus dose. A-C are from same patient, D-F are from a second patient, all at 24 hours.

Supplemental Figure 4: Caspace-3 IHC and quantification for explant culture out to 7 days in normoxic (B, D, F) and hyperoxic (C, E, G) conditions. Arrowheads point out positive IHC for Caspace-3. Apoptosis became evident in normoxic conditions on day 3 of culture, compared to day 1 for hyperoxic conditions. Two-way ANOVA with Tukey's *post-hoc* correction (H).

Supplementary Figure 5: Evaluation of migratory cell populations in the bottom transwell. Neutrophils (CD66b), NK (CD56), NKT (CD3 and CD56), T cells (CD3), B cells or Dendritic Cells (CD20 and CD11c, same fluorophore), and CD14 high or dim myeloid (CD11b) cells (**A**). Sequential analysis of the CD14^{dim} myeloid population demonstrating nearly all CD14^{dim} macrophages are positive for OPG, CX3CR1, ZO-1, and F11R (**B**). LPS in µg/mL, MCP-1 in ng/mL.

Supplemental Figure 6: Evaluation of scRNA-seq data. Variance analysis with top 20 genes labeled (A). Ridge plots of NK, M2, and RSM associated genes (B). Principle component analysis of all 10 clusters not including dead cells (C). Volcano plot of most up- and down-regulated genes, where minimum Log2FC was 1.5 for significance given homogeneity of sample (D).

Supplemental Figure 7: GSA for top 10 immune-relevant pathways (A), phagocytosis (B), complement (C), and adhesion-related (D).

Supplemental Figure 8: GSA for RSMs: top ten genes upregulated in inflammatory arthritis (top) and septic arthritis (bottom).