TgLaforin, a glucan phosphatase, reveals the dynamic role of storage polysaccharides

in *Toxoplasma gondii* tachyzoites and bradyzoites

Robert D. Murphy[1,2,a], Cortni A. Troublefield[1b#], Joy S. Miracle[1c#], Lyndsay E.A. Young[2],

Aashutosh Tripathi[1d], Corey O. Brizzee[2e], Animesh Dhara[1], Abhijit Patwardhan[3], Ramon

C. Sun[2fg], Craig W. Vander Kooi[2fg], Matthew S. Gentry[2fg], Anthony P. Sinai[1*]

Affiliations at time research was conducted:

1. Department of Microbiology, Immunology, and Molecular Genetics, College of

Medicine, University of Kentucky, Lexington, KY 40536, USA

2. Department of Molecular and Cellular Biochemistry, College of Medicine,

University of Kentucky, Lexington, KY 40536, USA

3. F. Joseph Halcomb III, MD. Department of Biomedical Engineering, College of

Engineering, University of Kentucky, Lexington KY 40506, USA

#These authors contributed equally to this work.

*For correspondence: Anthony P. Sinai, sinai@uky.edu

Current affiliations:

a. Maryland Department of Health, Laboratories Administration, Baltimore, MD

21205

b. Good Samaritan Hospital and Albert B. Chandler Medical Center Emergency

Departments, University of Kentucky, Lexington KY 40536

c. Department of Forestry and Natural Resources, Martin-Gatton College of

Agriculture, Food and Environment, University of Kentucky, Lexington, KY 40546

d. Department of Biochemistry, Stanford University School of Medicine, Palo Alto

CA 94305

e. Demeetra AgBio, 2277 Thunderstick Dr., Lexington, KY, USA

f. Department of Biochemistry and Molecular Biology, College of Medicine,

University of Florida, Gainesville, FL, USA

g. Center for Advanced Spatial Biomolecule Research, University of Florida,

Gainesville, FL, USA

Running title: TgLaforin regulation of amylopectin granules in Toxoplasma

Keywords: *Toxoplasma gondii*, amylopectin granule, glucan, reversible

phosphorylation, laforin, phosphatase, starch, glycogen, steady-state metabolomics,

glutamine

ABSTRACT

The asexual stages of *Toxoplasma gondii* are defined by the rapidly growing tachyzoite

during the acute infection and by the slow growing bradyzoite housed within tissue cysts

during the chronic infection. These stages represent unique physiological states, each

with distinct glucans reflecting differing metabolic needs. A defining feature of *T. gondii*

bradyzoites is the presence of insoluble storage glucans known as amylopectin

granules (AGs), the function of which remains largely unexplored during the chronic

infection. The presence of storage glucans has more recently been established in

tachyzoites, a finding corroborated by specific labeling with the anti-glycogen antibody

IV58B6. The *T. gondii* genome encodes activities needed for glucan turnover inlcuding:

a glucan phosphatase (TgLaforin; TGME49_205290) and a glucan kinase (TgGWD;

TGME49_214260) that catalyze a cycle of reversible glucan phosphorylation required

for glucan degradation by amylases. Disruption of TgLaforin in tachyzoites had no

impact on growth under nutrient-replete conditions. Growth of TgLaforin-KO tachyzoites

was however severely stunted when starved of glutamine despite being glucose replete.

Loss of TgLaforin attenuated acute virulence in mice and was accompanied by a lower

tissue cyst burden, without a direct impact on tissue cyst size. Quantification of relative

AG levels using AmyloQuant, an imaging based application, revealed the starch-excess

phenotype associated with the loss of TgLaforin is heterogeneous and linked to an

emerging AG cycle in bradyzoites.   Excessive AG accumulation TgLaforin-KO

bradyzoites promoted intra-cyst bradyzoite death implicating reversible glucan

phosphorylation as a legitimate target for the development of new drugs against chronic

T. gondii infections.

Importance

Storage of glucose is associated with a projected need for future metabolic potential.

Accumulation of glucose in insoluble amylopectin granules (AG) is associated with

encysted forms of *Toxoplasma gondii.* AG which are not observed in rapidly growing

tachyzoites do appear to possess glycogen, a soluble storage glucan. Here we address

the role of reversible glucan phosphorylation by targeting TgLaforin, a glucan

phosphatase and key component of reversible glucan phosphorylation controlling AG

and glycogen turnover. Loss of TgLaforin fundamentally alters tachyzoite metabolism

making them dependent on glutamine. These changes directly impact acute virulence

resulting in lowering tissue cyst yields. The effects of the loss of TgLaforin on AG levels

in encysted bradyzoites is heterogenous, manifesting non-uniformly with the

progression of the chronic infection. With the loss of TgLaforin culminating with the

death of encysted bradyzoites, AG metabolism presents a potential target for

therapeutic intervention, the need for which is acute.

INTRODUCTION

Toxoplasma gondii is an opportunistic protozoan parasite of all warm-blooded animals that infects one-third of humans worldwide (1, 2). Humans are primarily infected through the consumption of an encysted form of the parasite: either the oocysts shed in cat feces or tissue cysts found in undercooked meat from a chronically infected animal (3). Encysted parasites convert into tachyzoites that rapidly divide and disseminate throughout the body of the host, defining the acute phase of infection (4). Under host immune pressure, tachyzoites convert into slow-growing bradyzoites that populate tissue cysts which are found predominantly in the central nervous system and muscle, defining the chronic phase of infection (5, 6). Tissue cysts are believed to persist for the lifetime of the host and possess the ability to reactivate into tachyzoites in the context of immunosuppression. Reactivation can result in the life-threatening symptoms of toxoplasmosis, with toxoplasmic encephalitis being the primary condition leading to mortality (7, 8). The current lack of insights into bradyzoite physiology *in vivo* precludes the basic understanding needed for the development of drugs that either can clear tissue cysts or target encysted bradyzoites so as to prevent reactivation (9).

Until recently, bradyzoites within tissue cysts were considered to be dormant, metabolically inert entities. This view was challenged by our demonstration that encysted bradyzoites replicate (10, 11). Moreover, bradyzoite physiology is both diverse and complex as viewed through the lens of mitochondrial activity (12, 13), replication status (11), and, importantly, amylopectin granule (AG) accumulation (14). Although the function of AGs in bradyzoites has not been confirmed, an understanding of the roles of

polysaccharides elsewhere suggests that AGs are a source of energy and biosynthetic

potential needed for persistence, replication, reactivation, and transmission (15). These

assumptions remain to be tested, and thus much like bradyzoites themselves, the role

of AGs in the *T. gondii* lifecycle is poorly understood. Our first insights into the potential

relationships connecting AG to intermediate metabolism via mitochondrial and

replication activities have been exposed using imaging-based approaches (14).

AGs are large glucans found in the cytoplasm of bradyzoites that have classically

served as a morphological feature distinguishing them from tachyzoites (16-19).

Toxoplasma AGs are much like plant starch in that they are water-insoluble storage

polysaccharides composed of branched chains of glucose (18). Unlike plant starch,

however, AGs contain no detectable amylose (unbranched chains of glucose) (18).

More recently, the presence of small, punctate, cytoplasmic glucans in tachyzoites that

are only visible by periodic acid-Schiff (PAS) staining have been recognized (20-22),

and the presence of the glucan is dependent on the *T. gondii* starch synthase (TgSS;

TGME49_222800) (23). Like animal glycogen, this tachyzoite storage polysaccharide is

rapidly turned over (20), as has been observed in other protozoa (24-26), and provides

glucose for glycolysis (23, 27). The observation that large, insoluble glucans do not

accumulate within tachyzoites as they do within bradyzoites suggests that the

tachyzoite glucan could be a distinct and labile form of stored glucose, likely glycogen-

like, although its exact chemical and structural identity remains unknown.

Glucose release from starch in plants requires a cycle of direct, reversible glucan

phosphorylation to solubilize the starch surface, allowing access to degradation

enzymes such as amylases, branching enzymes, and a phosphorylase (28-30). The

cycle begins with the addition of phosphate directly to glucose by the glucan, water-

dikinase (GWD) and phospho-glucan, water dikinase (PWD) that results in the

unwinding of glucose chains within starch, solubilizing the starch surface (31, 32).

Glucose-releasing enzymes (amylases) then degrade starch until the glucan-bound

phosphate becomes a steric hindrance, at which point a glucan phosphatase is needed

to remove the phosphate and reset the cycle (33-35). *T. gondii* encodes all the activities

needed for glucan degradation and reversible glucan phosphorylation including the

glucan phosphatase, TgLaforin (TGME49_205290) (36), and glucan dikinase, *T. gondii*

GWD (TgGWD; TGME49_214260) (27).   The central role of reversible glucan

phosphorylation in plants is seen in *Arabidopsis thaliana* where loss of the plant glucan

phosphatase, starch-excess 4 (SEX4), results in excess starch accumulation, aberrant

starch morphology, and severely stunted plant growth (37, Zeeman, 1998 #1387).

Additionally, loss of the glucan phosphatase, laforin, in humans, results in

hyperphosphorylated glycogen that aggregates in neurons and astrocytes causing a

fatal neurodegenerative childhood dementia and epilepsy (38-40).   In *T. gondii*,

perturbations of several genes related to glucan metabolism also result in a variety of

similar defects including aberrant glucan accumulation, rewiring of central carbon

metabolism, and virulence defects in mice, highlighting the central metabolic role of

glucan metabolism in *T. gondii* (20-23, 27, 41-44)

In this study, we build on our understanding of reversible glucan phosphorylation and its

relevance to parasite metabolism in *T. gondii*. We have recently demonstrated that

TgLaforin is the glucan phosphatase in *T. gondii*, and that TgLaforin represents a

unique and viable drug target (36, 45, 46). Here, we investigate the role of TgLaforin

throughout the asexual stages by knocking out TgLaforin in Type II ME49 parasites.

While we expected to observe effects related to the loss of TgLaforin exclusively in

bradyzoites where AGs are typically observed, these effects appeared to be connected

to an emerging AG temporal cycle (14). More surprisingly, the loss of TgLaforin also

resulted in phenotypic effects in tachyzoites, also in a context-specific manner. We thus

established a role for TgLaforin, and by extension reversible glucan phosphorylation,

across both tachyzoite and bradyzoite life stages. These findings build upon previous

studies that increasingly demonstrate a central role for glucan metabolism throughout

the parasite's asexual life cycle.

RESULTS

T. gondii tachyzoites contain a cytoplasmic glucan with a punctate distribution.

Previous studies have presented biochemical evidence for rapid glucan turnover in *T.*

gondii Type I RH tachyzoites (20). Moreover, small granules that stain with periodic acid

Schiff reagent (PAS) have also been noted in the cytoplasm of tachyzoites (20, 21, 23).

Under acid-stress conditions, these tachyzoite glucans have been biochemically

characterized as pure amylopectin, and resemble AGs seen in bradyzoites (18). To

further characterize the nature of this tachyzoite glucan, we used multiple methods to

visualize them under unstressed, normal growth conditions (**Figure 1A**). PAS staining

confirmed that Type II ME49 tachyzoites contain small punctate granules distributed

throughout the cytoplasm. To determine if these PAS-stained granules in unstressed

tachyzoites were more glycogen- or starch-like, they were stained with IV58B6. IV58B6

is an anti-glycogen IgM monoclonal antibody that has previously been demonstrated to

be specific to glycogen (47-49) by recognizing the highly frequent branch-points found

in glycogen (48). Moreover, IV58B6 does not detect other glucans such as amylopectin

or amylose (the primary constituents of plant starch) (50). Tachyzoites stained with

IV58B6 in a similar pattern to PAS-stained parasites, containing small punctate granules

distributed throughout the cytoplasm (**Figure 1A**), suggesting that the glucan found in

tachyzoites is more glycogen-like than starch-like. Finally, as is well-known, *T. gondii*

tachyzoites contain almost no visible glucan within their cytoplasm when visualized by

transmission electron microscopy (TEM) (**Figure 1A**), suggesting that the glucan

detected by both PAS staining and IV58B6 is either water-soluble or too small to be

visualized, consistent with this glucan being glycogen-like.

In contrast, bradyzoites have been extensively characterized as containing starch-like

AGs (17-19, 51). After *in vitro* bradyzoite conversion, much of the cytoplasm stained

heavily with PAS (**Figure 1A**). Interestingly, IV58B6 staining intensity appeared to

correlate negatively with *Dolichos biflorus* agglutinin (DBA) staining intensity that

defines the cyst wall, implying that IV58B6 does not stain the PAS-stained glucan in

bradyzoites, further reinforcing the observation that structurally distinct polysaccharides

exist in tachyzoites and bradyzoites (**Figure 1A**). Finally, unlike in tachyzoites, AGs

were readily identified as electron-lucent structures throughout the cytoplasm of in vitro

generated bradyzoites by TEM (**Figure 1A**).

To verify the specificity of PAS and IV58B6 for glucose polymers, tachyzoites and

bradyzoites were treated with acid-α-amyloglucosidase (GAA) after parasite fixation and

before staining. GAA cleaves both α-1,4- and α-1,6-glycosidic bonds and can therefore

completely digest glucans into glucose monomers. Indeed, GAA treatment resulted in

the disappearance of staining within both tachyzoites and in vitro bradyzoites (**Figure**

1B) demonstrating their specificity for glucose polymers.

TgLaforin colocalizes with the tachyzoite glucan

Because *T. gondii* encodes TgLaforin, a glucan phosphatase that is more animal-like

than plant-like (45, 46), we reasoned that TgLaforin could be involved in the metabolism

of the glycogen-like glucan found in tachyzoites. To determine if TgLaforin co-localizes

with the tachyzoite glucan, endogenous TgLaforin was epitope-tagged with

hemagglutinin (HA) in *T. gondii* Type II ME49ΔHXGPRT parasites (52) with a

CRISPR/Cas9 mediated strategy (**Figure 2A**) (53). Successful tagging of TgLaforin was

confirmed by western blotting (**Figure 2B**). Immunofluorescence analysis (IFA) of *T.*

gondii tachyzoites indicated that TgLaforin is present in small puncta throughout the

cytoplasm, similar to the distribution of the tachyzoite glucan (**Figure 2C**). Surprisingly,

TgLaforin was not detected in *in vitro* bradyzoites by IFA, 6 days post conversion

(**Figure 2C**).   To verify that TgLaforin levels decrease during the tachyzoite to

bradyzoite transition, we converted *T. gondii* tachyzoites to bradyzoites in cell culture

using alkaline stress for 6 days and then probed the converted parasites using western

blot analysis. As observed using IFA, TgLaforin-HA expression decreased dramatically

over the course of bradyzoite differentiation (**Figure 2D**). Transcriptomic data from a

previous study obtained from ToxoDB.org indicates that the transcript levels for

TgLaforin do not substantially change over the course of differentiation, suggesting the

possibility that levels of TgLaforin protein are regulated by post-translational

mechanisms (54). To determine if TgLaforin colocalizes with the glucan present in

tachyzoites, we co-stained TgLaforin-HA tachyzoites with either PAS or IV58B6 along

with an anti-HA antibody. In tachyzoites, TgLaforin colocalized with both PAS (**Figure

2E**) and with IV58B6 (**Figure 2F**), suggesting its involvement in the metabolism of the

tachyzoite glucan.

Initial characterization of TgLaforin-KO tachyzoites

To dissect the role of TgLaforin in *T. gondii* glucan metabolism, TgLaforin was knocked

out using CRISPR/Cas9 to disrupt the gene with a pyrimethamine-resistant form of the

dihydrofolate reductase (DHFR-TS*) under a *Neospora caninum* GRA7 (NcGRA7)

promoter (55) (56) (**Figure 3A**). In agreement with a genome-wide CRISPR KO screen

(57), TgLaforin is a non-essential gene under standard cell culture conditions, as

multiple TgLaforin-KO clones were successfully recovered. Integration of the DHFR-TS*

construct into the TgLaforin locus was verified using inside/out PCR at the chimeric

locus and by verifying the loss of TgLaforin transcription (**Figures 3B,C**). The TgLaforin-

KO line further used in this study (designated "ΔTgLaf") was complemented by the

introduction of an epitope tagged (HA) gene driven by the TgLaforin promoter. The

complementation construct was introduced at an ectopic site in the genome that lacks

known coding sequences or regulatory elements on chromosome VI (**Figure S1A**),

while leaving the ΔTgLaf/DHFR-TS* KO lesion intact for true complementation (58).

This complemented strain, henceforth designated "COMP," was successfully isolated

and confirmed by PCR (**Figure S1B**). Expression levels and localization were similar to

those seen in the TgLaforin-HA line as confirmed by western blotting and IFA (**Figures**

S1C-D),

To evaluate effects of a TgLaforin-KO, glucan levels in WT and ΔTgLaf tachyzoites

were first compared using our suite of glucan detection techniques (**Figure 3D**).

Surprisingly, the size and number of PAS-stained granules were not significantly

changed in ΔTgLaf tachyzoites relative to WT parasites. Levels of IV58B6 also

remained unaltered after the loss of TgLaforin, and no aberrant glucan accumulation

was observed by TEM as has been previously reported when genes related to AG or

central carbon metabolism were knocked out in *T. gondii* tachyzoites (20-23, 41, 43, 44)

(**Figure 3D**)

Loss of glucan phosphatases in plants and animals results in aberrant glucan

accumulation, and such a phenotype was not observed within tachyzoites under

standard growth conditions.

Loss of TgLaforin results in upregulation of glutaminolysis and glutamine dependence in tachyzoites

Glucan catabolism is significantly affected by the presence of covalently bound phosphate, and, therefore, loss of glucan phosphatases has profound downstream metabolic impacts in other systems (59, 60). We thus speculated that loss of TgLaforin would result in the reduced efficiency of glucan utilization in tachyzoites and also affect downstream central carbon metabolism. To test this hypothesis, we used gas chromatography/mass spectrometry (GC/MS) steady-state metabolomic analysis of 3 μm filter-purified, syringed-passaged intracellular tachyzoites employing a previously developed sample preparation technique (61).

Previously, it was demonstrated that *T. gondii* tachyzoites primarily utilize glucose and glutamine to drive central carbon metabolism, synthesize macromolecules, and proceed normally through the lytic cycle (62). Glucose primarily fuels glycolysis, and glutamine undergoes glutaminolysis to drive the tricarboxylic acid (TCA) cycle. In the absence of glucose, *T. gondii* can upregulate both glutaminolysis and gluconeogenesis to make up for the loss of glucose (62, 63).

While ΔTgLaf metabolite levels remained unaltered relative to WT tachyzoites across much of the TCA cycle, steady-state levels of metabolites immediately downstream of glutamine were consistently more abundant in ΔTgLaf parasites compared to their WT counterparts (**Figure 4A**), supporting our hypothesis that ΔTgLaf parasites were

deficient in glucan/glucose utilization. An increase in metabolites downstream of

glutamine in ∆TgLaf parasites demonstrates that ∆TgLaf parasites are possibly

compensating for deficiencies in glucose metabolism, supporting a role for the

tachyzoite glucan in intermediate *T. gondii* glucose metabolism.

To determine if loss of TgLaforin resulted in increased dependence on glutamine due to

impaired access to glucose, we performed plaque assays in the presence and absence

of glutamine (**Figure 4B**). In replete media, ∆TgLaf parasites established a similar

number of plaques (data not shown), indicating no defect in infectivity. ∆TgLaf plaques

were slightly larger than both the WT and COMP lines after 10 days of growth (**Figure**

4C). To test the effects of glutamine starvation on ∆TgLaf parasites, glutamine was

removed from plaque assays after parasite invasion to evaluate the effects of glutamine

removal on parasite growth independent of the initial invasion event. In the absence of

glutamine, ∆TgLaf parasites were unable to form visible plaques, whereas both the WT

and COMP parasites formed plaques comparable to those formed under glutamine-

replete conditions (**Figures 4B, C**).

TgLaforin is required for repeated rounds of progression through the lytic cycle

The absence of differences in plaque number suggest that there is no specific defect in

infectivity. To determine which aspects of the *T. gondii* lytic cycle were impaired in the

absence of glutamine, the effects of glutamine starvation on initial parasite replication

and egress (stimulated with both A23187 (64) and zaprinast (65)) were evaluated. In

both assays, intracellular parasites were pre-starved of glutamine for at least 72 h

before assay initiation. Surprisingly, glutamine starvation had no effect on stimulated

egress or initial parasite replication across the three lines (**Figures S2A-C**). These data

demonstrate that the absence of plaques under glutamine deficient conditions cannot

be pinpointed to a single aspect of the ΔTgLaf lytic cycle, and that the reason for the

apparent absence of plaques manifested later in the infection cycle.

Plaques develop due to repeated cycles of localized infection and cell lysis resulting in the

clearance of infected cells over time. The absence of visible clearance prompted us to

examine infected host-cell monolayers for clusters of infected cells using a higher

magnification than is typically used in a traditional plaque assay. Low numbers of parasites

were seeded onto glass coverslips and fixed at 3- and 6-days post-infection, allowing for

visualization of developing plaques at a high magnification. In these experiments, glutamine-

depleted host cells were pre-starved of glutamine prior to infection with parasites to allow for

potential for invasion defects. Importantly, ΔTgLaf parasites demonstrated similar infectivity to

WT parasites under both glutamine-replate and depleted conditions, indicting no gross initial

invasion defect. After 3 days of growth, no statistical differences of nascent plaque sizes were

noted between glutamine-replete and starved conditions in both the WT and COMP lines.

However, ΔTgLaf parasites in glutamine starved conditions were already 1.5x smaller in area

than their counterparts in replete conditions (**Figure 4D**). By day 6 of growth, this difference

had widened to a >3x difference between glutamine replete/depleted ΔTgLaf parasites (**Figure

4E**). Such a difference was not detected between the two conditions in WT/COMP parasite

lines. By measuring the internal clearing area relative to the total plaque perimeter, it was also

noted that ΔTgLaf parasites were much less capable of forming clearings than the WT/COMP

lines (**Figures S2D, E**), rather they formed clusters of infected cells akin to "turbid plaques"

(66, 67) due to their presumed inability to compete with host cell growth, as the infection

progressed. This observation explains the apparent absence of plaques seen at the lower

magnification used in traditional plaque assays (**Figure 4B**). The modified plaque assay

therefore confirmed that the loss of TgLaforin penalized the summation of repeated rounds of

the energy-demanding lytic cycle rather than one particular aspect of the lytic cycle.

Representative images from this assay can be found in **Figure S2D**.

Loss of TgLaforin results in aberrant bradyzoite AGs in vitro

To determine if loss of TgLaforin resulted in bradyzoite conversion defects, or aberrant

AG accumulation, parasites were converted to bradyzoites *in vitro* using alkaline stress.

During differentiation, the parasitophorous vacuole membrane (PVM), delimiting the

replicative niche established by tachyzoites, converts into the cyst wall that surrounds

bradyzoites within their host cell (68, 69). The cyst wall is heavily glycosylated and

contains N-acetylgalactosamine (Gal-NAc) that is detectible with Dolichos biflorus

agglutinin (DBA) (68). Using DBA-FITC intensity as a marker for differentiation, no

penalty was imposed by the loss of TgLaforin on cyst wall formation over the course of

six days (**Figure 5A**). Somewhat surprisingly, ΔTgLaf mutant parasites tended to exhibit

stronger labeling with DBA at day 6. We additionally assessed the levels of

accumulated glucans using PAS staining (**Figure 5B**). Semi-quantitative analysis of

PAS intensity within vacuoles during stage conversion showed an expected increase

over time, but no significant difference between the WT and ΔTgLaf parasites was

detected over the time course examined.

Because PAS is not specific to glucans and can stain other glucose-containing

molecules such as glycosylated protein and provides no resolution on glucan

morphology, we utilized TEM to gain higher resolution on AG formation during

bradyzoite differentiation. After 6 days of conversion, WT parasites produced AGs that

were circular/ovoid and white (**Figure 5C**). In contrast, ΔTgLaf parasites contained

irregular AGs that were morphologically distinct from AGs that were observed in WT

parasites (**Figure 5C**). AGs in ΔTgLaf parasites appeared amorphous and grayer, while

appearing to occupy more area of the parasite cytoplasm compared to WT parasites. To

quantify this phenotype, the area of AGs was calculated relative to total parasite area to

determine the percentage of the parasite body occupied by AGs in both WT and ΔTgLaf

strains (**Figure 5D**). Strikingly, AGs occupied approximately 4x more relative area in

ΔTgLaf parasites when compared to WT, indicating that PAS staining may lack the

sensitivity to capture this difference, in tissue culture generated bradyzoites. When

analyzed on an 8-bit gray scale, AGs in ΔTgLaf parasites were significantly grayer

(grayscale 0-255 is black-white) than those found in WT parasites, highlighting potential

chemical differences (such as predicted hyperphosphorylation) resulted in differential

interactions of ΔTgLaf AGs with the TEM contrast reagents, likely the heavy metals

used in processing (**Figure 5E**).

Examination and quantification of AGs in the COMP line revealed that complementation

of TgLaforin restored most of the circular/ovoid AG cross sections while they also

occupied less space in the cytoplasm and were overall more like those found in WT

parasites (**Figures 5C-E**). Thus, cell culture experiments demonstrate that the loss of

TgLaforin presents itself in both a context and life cycle stage-specific manner.

Loss of TgLaforin results in attenuated virulence and cyst formation in vivo

We hypothesized that loss of TgLaforin may impose a steep penalty under the stresses

and potential nutrient scarcities encountered *in vivo* as it does when nutrients (such as

glutamine) are scarce *in vitro*. To test this hypothesis, equal numbers of male and

female CBA/J mice were infected with 100 tachyzoites intraperitoneally (i.p.) and

monitored daily using a previously developed five-stage body index score to track the

severity of symptoms associated with a tachyzoite infection over the course of 28 days

(52).

Mice infected with WT parasites began demonstrating symptoms of infection ten days

after infection with tachyzoites (**Figure 6A**). However, mice infected with ΔTgLaf

parasites did not begin to exhibit symptoms until 15 days after infection. Moreover, mice

that became symptomatic from WT parasite infections often proceeded through all

stages of symptomology, and only a minor proportion of mice that became sick were

able to recover from infection (>70% of mice became moribund or died). Infection from

ΔTgLaf parasites, however, resulted in the majority of mice only developing mild

symptoms (Stage 2 or less) with many of these mice recovering (**Figure 6A**). The

attenuated capacity of the ΔTgLaf parasites to cause symptoms in mice was reflected in

the mortality rates of the infected mice: infection with WT parasites resulted in 73%

mortality rate after 28 days whereas ΔTgLaf parasites only caused 17% mortality

(**Figure 6B**). Complementation of TgLaforin partially rescued this defect in virulence as

COMP parasites resulted in an earlier onset of symptomatic infection at Day 11, and the

majority (53%) of mice succumbed to infection during the first 28 days (**Figures 6A, B**).

Because the acute phase of infection was significantly attenuated by the loss of

TgLaforin, we hypothesized that cyst numbers would be significantly lowered. To

determine the number of cysts formed after 28 days of acute infection (Week 4), we

used a previously established protocol for harvesting and counting tissue cysts from

infected mouse brains, following purification on Percoll gradients (11, 70). Consistent

with the ability of ΔTgLaf parasites to stage convert in culture, mutant parasites were

able to establish tissue cysts *in vivo*. However, the number of cysts recovered from

ΔTgLaf infected animals was lower than those obtained from WT infected animals

(**Figure 6C**). Restoration of TgLaforin in the COMP line effectively restored tissue cyst

yields.

ΔTgLaf tissue cysts can reestablish infections in naïve mice

To determine if the loss of TgLaforin impacted the overall viability/infectivity of *in vivo*

tissue cysts, we examined the disease progression in WT, ΔTgLaf, and COMP infected

animals following injection of 20 tissue cysts i.p. Consistent with prior data (52),

infection with tissue cysts results in markedly lower pathology and consequent mortality

during the acute phase for WT as well as both the ΔTgLaf and COMP lines (**Figure 7A,**

B). Mortality from cyst infections did not differ statistically among the three lines (**Figure**

7A, B). Twenty-eight days post-infection, cyst burdens were again enumerated for each

line. ΔTgLaf parasites were once again much less competent at forming cysts *in vivo*

(**Figure 7C**). However, unlike the tachyzoite infection, the COMP line was unable to

rescue this defect in cyst formation (**Figure 7C**), suggesting that physiological and

metabolic changes associated with the loss of TgLaforin manifest differently based on

the life cycle stage, impacting their capacity to be complemented.

Loss of TgLaforin results in a delayed starch-excess phenotype within encysted

bradyzoites

We leveraged the inherent fluorescence of PAS coupled with an optimized labeling

protocol to determine the mean intensity of WT, ΔTgLaf and complemented (COMP)

tissue cyst using Image J (**Figure. 8A**). Similar to the pattern observed with in vitro

generated bradyzoites (**Figure. 5**), *in vivo* derived tissue cysts harvested at 28 days

(Week 4) post infection failed to demonstrate a starch-excess phenotype in the ΔTgLaf

mutant (**Figure 8A**). Notable here, is the fact that at week 4 post infection, bradyzoites

are in a state of low AG accumulation (14), and presumably, low turnover, potentially

minimizing the impact of the loss of TgLaf. Our prior data indicate that tissue cysts

harvested at week 6 post infection are in markedly more dynamic state noted by

increased accumulation of AG, higher levels of active mitochondria and increased

replicative activity (14). Quantification of mean PAS intensity exposed a dramatic

increase in AG levels comparing WT and ΔTgLaf tissue cysts, with overall levels

between WT and COMP cysts remaining not significantly different (**Figure 8A**). This

points to the effect of the loss of TgLaf demonstrating a delayed starch excess

phenotype, indicating the penetrance of the mutation is potentially connected to

achieving a threshold level of AG as a part of an emerging AG cycle (14).

The development of AmyloQuant, an imaging-based application, permits the direct

measurement of relative AG levels within tissue cysts (14).   With is application,

individual PAS-labeled pixels are classified into 4 bins defining background pixels

(black: 0-10 grayscale), low intensity (blue: 10-25), intermediate (green: 25-50) and high

(red: > 50 grayscale). Thirty randomly acquired tissue cysts, (the mean intensity of

which were reported in **Figure 8A**) from WT, ΔTgLaf and COMP mice infected for 4

weeks (**Figure 8B, top row**) and 6 weeks (**Figure 8B, bottom row**) are arrayed based

on their overall mean intensity. The percentage of pixels within each of the 4 bins is

presented as a stacked plot, with the AmyloQuant generated thumbnails representing

spatial heatmaps of AG intensity at 5 cyst intervals as presented below each stacked

plot. This analysis reveals that despite the mean PAS intensities not being statistically

significant (**Figure 8A**), several ΔTgLaf cysts present with varying levels of high

intensity (red) PAS labeling. Notable here is that with this cohort of cysts, effective

complementation of AG levels is not observed.

As expected for wild type Week 6 tissue cysts (14), markedly increased AG

accumulation is evident compared to Week 4 cysts (**Figure 8B**). This phenotype is

grossly exaggerated in the case of ΔTgLaf tissue cysts (**Figure 8B**), displaying the

expected starch excess phenotype. Even so, not all ΔTgLaf cysts present uniformly

high levels AG based on PAS intensity (**Figure 8B**). Notable here, the levels and

distribution of AG levels within the COMP line appear to be intermediate to that

observed for the WT and ΔTgLaf, pointing to the plasticity of the phenotype.

Several recent studies targeting activities connected with AG metabolism report

differences in cyst yield as well as cyst size (21-23, 41). We therefore examined if the

loss of TgLaf resulted in any significant impact of cyst size. Contrary to other reports

however no differences in cyst diameter were noted for either week 4 or week 6

harvested tissue cysts (**Figure S3A**). Furthermore, we found no strong correlation

between cyst size and mean AG levels as defined by PAS labeling intensity (**Figure

S3B**).

Aberrant AG accumulation in ΔTgLaf tissue cysts promotes bradyzoite death.

The inherent harshness of PAS labeling causes differential labeling intensity related

issues with regard to the labeling of bradyzoite nuclei (14) as well as labeling for

TgIMC3 a marker of recent replication (14). As a result, the relationship between AG

levels and replication associated outputs (packing density and recency of replication)

could not be reliably implemented in the context of the TgLaf mutant. We therefore

resorted to the direct examination of purified tissue cysts by transmission electron

microscopy.

We adapted a protocol designed to capture and image low abundance cells by TEM by

making it compatible with our tissue cyst purification protocol (see Methods) (71, 72).

TEM imaging revealed that while WT parasites formed largely normal/canonical AGs *in*

vivo as seen *in vitro* (**Figure 9** [compare with **Figure 5A**]), ΔTgLaf parasites contained

almost exclusively aberrant AGs that mirrored the same morphological defects seen *in*

vitro (**Figure 9** and **Figure S4**). ΔTgLaf AGs were irregularly sharpened with a flat,

multi-lobed appearance. Importantly, COMP parasites neither over-accumulated nor

formed aberrant AGs, demonstrating that this defect is specific to loss of TgLaforin

(**Figure 9**).

In addition to containing aberrant AGs, the internal morphology of ΔTgLaf parasites

appeared to be altered by the presence of the AGs, as significant organelle

displacement was also noted (**Figure 9, S4**). Many of the ΔTgLaf parasites also

appeared hollow or "ghost-like" in appearance resulting from their unstained/absent

cytoplasm, which included the apparent loss of the nucleus in several bradyzoites

(**Figure 9, S4**). These data suggest that a significant number of the ΔTgLaf bradyzoites

were inviable within the cyst (**Figure 9, S4**). Together, these direct and selective

impacts of TgLaforin's loss on both bradyzoite viability and growth *in vivo* establish

TgLaforin as a potentially druggable target.

DISCUSSION

The asexual life cycle of *Toxoplasma gondii* is defined by two fundamentally distinct

forms: the rapidly replicating tachyzoite and the slowly growing bradyzoite. These forms

represent distinct physiological states that can be further subdivided, particularly within

encysted bradyzoites (73). Insights into these physiological states, inferred from

transcriptomic analyses, are reinforced in emerging metabolomic studies (74, 75). In

these studies, glucose and glutamine, which are both linked to energetics, biosynthesis,

and intermediary metabolism, appear as critical metabolites. Importantly, glucose can

be stored in polymers like glycogen and amylopectin during times of low energy need.

Stored glucose can be present in an accessible and labile form for rapid mobilization

such as glycogen, or in a less soluble forms as AGs from which it can be accessed

more slowly (76, 77). Reversible glucan phosphorylation facilitates the breakdown of

such glucan polymers by disrupting the crystalline helices on the glucan surface (78). *T.*

gondii encodes the capacity for reversible glucan phosphorylation (27, 36, 45, 46). The

contribution of this process to tachyzoite and bradyzoite biology was evaluated through

targeted manipulation of the glucan phosphatase, TgLaforin (36).

The accumulation of AGs within bradyzoites and their apparent absence in tachyzoites

has been used as discriminator between these life cycle stages (16, 79). Detailed

examination, however, presents a considerably more nuanced picture, alongside

emerging evidence that points to rapid glucan turnover within tachyzoites (20). In the

current study, using an IgM monoclonal antibody (IV58B6) that specifically recognizes

glycogen-like glucose polymers (47, 48), we demonstrate that the stored glucan within

tachyzoites is structurally closer to animal glycogen (47, 48) than the insoluble plant-like

AG granules found in bradyzoites (17, 19). This duality between life stages may be

additionally reflected in the observation that TgLaforin, the glucan phosphatase, and

TgGWD (TGME49_214260), the partner kinase that is predicted to phosphorylate *T.*

gondii glucan polymers, trace their structural lineages to animals and plants,

respectively (45, 46). The glycogen-like glucan polymer appears to be specific to

tachyzoites as its levels decrease upon *in vitro* differentiation while the overall PAS

intensity increases (**Figure 1A**). This suggests that the tachyzoite glucan and bradyzoite

AG are architecturally distinct polymers with respect to both branching frequency,

solubility, and phosphorylation status (**Figure 2D**). The glucose-based nature of both

particles is supported by the elimination of both IV58B6 and PAS staining with α-

amyloglucosidase treatment (**Figure 1B**).

To address the contribution of stored glucans in both tachyzoites and bradyzoites, we

disrupted the glucan phosphatase TgLaforin. This enzyme preferentially removes

phosphate groups from the C3 carbon on glucose facilitating access to enzymes that

release glucose (36). The loss of other glucan phosphatases such as SEX4 in *A.*

thaliana and laforin in mammals is accompanied by excessive accumulation of aberrant

starch and hyperphosphorylated glycogen in plants and animals (37, 39, 40, 80, 81).

Surprisingly, given these penalties in other systems, ΔTgLaf tachyzoites exhibited no

gross morphological changes in glucan content, consistent with a recent study in which

TgGWD was knocked out (27). These observations contrasts with other KO studies of

glucan pathway associated proteins such as CDPK2 (20), glycogen phosphorylase (41),

the PP2A holoenzyme (44), and α-amylase (22)which all reported exaggerated glucan

accumulation in tachyzoites.

Despite the absence of glucan accumulation in ΔTgLaf tachyzoites, the loss of this gene

exerts an effect on tachyzoite glucan metabolism. Consistent with the metabolic defects

associated with the loss of laforin in humans (60), loss of TgLaforin in *T. gondii* resulted

in altered central carbon metabolism that manifested as ΔTgLaf parasites' dependence

on glutamine (**Figure 4, S2D,E**). ΔTgLaf tachyzoite dependence on glutamine supports

recent studies demonstrating that tachyzoites utilize storage glucans for glucose

allocation (23) because the presumed loss of efficient glucan degradation results in

tachyzoite dependence on glutamine (**Figures 4B,C**). As *T. gondii* tachyzoites primarily

utilize glucose and glutamine to support their rapid growth, this dependence on

glutamine reinforces previous observations that glutamine can substitute for glucose in

this altered metabolic landscape (62, 82). These results are consistent with many

previous studies that disrupt glucose and/or glucan metabolism, but contrast with

others. Disruption of TgGT1 (TGME49_214320), the only plasma-membrane glucose

transporter in *T. gondii* (62, 83), or TgHK (TGME49_265450), the *T. gondii* hexokinase,

resulted in upregulation of gluconeogenesis, and parasite growth was highly attenuated

with glutamine depletion (82). Most strikingly, parasites lacking starch synthase (TgSS;

TGME49_222800) displayed no dependence on glutamine and, in fact, grew faster than

WT parasites when both glucose and glutamine were removed from the culture media

(23). Interestingly, however loss of TgSS did result in lower glucose flux through

glycolysis (23), consistent with our findings that demonstrate a role for the tachyzoite

glucan in glucose allocation. This finding may suggest that while the absence of AG in

the ΔTgSS parasites may not be detrimental, overaccumulation as observed in ΔTgLaf

cysts can contribute to toxicity (**Figure 9**). Perturbations of other glycolytic enzymes

also demonstrated varied effects related to the presence of glutamine: loss of the

glycolytic enzyme glyceraldehyde-3-phospahte dehydrogenase (GAPDH1) could be

rescued with high levels of glutamine (84) (84), but glutamine could not rescue pyruvate

kinase (TgPYK1) knockdown parasites (85). Our data indicate that loss of access to key

nutrients such as glucose and glutamine has a profound impact on the repeated rounds

of the lytic cycle without being attributable to one specific process within the cycle,

suggesting that the penetrance of the phenotypic defect manifests cumulatively over

time, rather than being hard wired in each infection cycle (**Figures 4D,E, S2D,E**).

Despite glucan metabolism being historically viewed as being important in the chronic

infection, TgLaforin protein expression decreased during the tachyzoite to bradyzoite

conversion *in vitro* even though its transcript levels do not change (**Figures 2C,D**). This

could be a transient observation as the downregulation of glucan catabolism during

conversion would facilitate accumulation of AGs for the chronic infection. We therefore

examined how the loss of TgLaforin affected the capacity of ΔTgLaf parasites to

differentiate *in vitro.* The ΔTgLaf parasites exhibited no defect in AG-accumulation

kinetics, detected by PAS staining, or in cyst wall formation, detected with DBA lectin

over the course of the in vitro conversion assay (**Figure 5A, B**). The lack of difference in

PAS labeling between both the WT and the ΔTgLaf lines, however, did not reveal the

differences noted by TEM (**Figure 5C-E**). As initially hypothesized would be the case in

both tachyzoites and bradyzoites, loss of TgLaforin resulted in aberrant AG

accumulation within *in vitro* bradyzoites that is marked by changes in both level and

morphology (**Figures 5C-E**), as seen in plants and vertebrates (37-40, 81). AGs in the

ΔTgLaf parasites were not only present at higher levels but were potentially chemically

distinct considering their differential binding to TEM contrast metals (**Figure 5E**). Given

that TgLaforin is a confirmed glucan phosphatase (36), we speculate that AG

hyperphosphorylation may account for both altered morphology and appearance by

TEM.

These context-specific phenotypes suggested that the ΔTgLaf mutant would manifest

phenotypic differences in both the acute and chronic phases on infection *in vivo*.

Indeed, the loss of TgLaforin was associated with a markedly reduced symptomology

and associated mortality compared to both the parental and complemented parasites

during acute tachyzoite-initiated infection (**Figures 6A,B**). Not only was there a delay in

symptomatic disease, but also a reduction in disease severity and overall cyst burden.

Symptomology in the acute infection is driven by an increasing parasite burden driving

an overexuberant host inflammatory response (4, 86). The delayed symptom onset

suggests growth inhibition by the stringent *in vivo* environment that more effectively

controls ΔTgLaf parasite infection with less robust inflammation. Notably, the delayed

and milder course of the tachyzoite infection resulted in a lower overall cyst burden in

surviving animals compared to infection with both WT and COMP parasites (**Figure

6C**).

Infection with ΔTgLaf tachyzoites resulted in fewer tissue cysts being generated relative

to WT and COMP parasite (**Figure 6C**) The basis for this is not clear, but suggestive of

a regulatory imbalance that is masked in tachyzoites but evident in tissue cyst-initiated

infections which have the additional burden of converting from bradyzoites to

tachyzoites, surviving the chronic infection before forming new tissue cysts. This could

potentially parallel the observations with plaque formation in vitro (**Figure 4, S2**).

Our ability to quantify and map relative AG levels within tissue cysts using

AmyloQuant (14), confirm that the loss of TgLaf does not fundamentally alter the

initiation and early progression of an AG cycle as evidenced by the effect of the

mutation not being evident in Week 4 tissue cyst in the form of the expected starch

excess phenotype (**Figure 8**). The effect of the ΔTgLaf mutation becomes evident in

week 6 tissue cysts where a massive increase in AG accumulation (**Figure 8**) is likely

due to an imbalance caused by the predicted defect in AG turnover. As with other

phenotypes associated with this mutation, the effect continues to exhibit phenotypic

variation noted by roughly a third of the imaged cysts lacking a significant proportion of

high intensity pixels (red), while others are completely oversaturated (**Figure 8**). These

distinct populations are not reflected in any way based on the size of the cyst (**Figure

S3**). In fact, contrary to other reports regarding AG metabolism associated genes, the

TgLaf mutation has no significant effect on tissue cyst size (**Figure S3**).

While we were not able to accurately quantify either the nuclear number or TgIMC3

intensity distributions as markers of packing density and replicative activity in PAS

stained tissue cysts (14), TEM analysis exposed the true consequence of the loss of

TgLaf within tissue cysts in vivo (**Figure 9, S4**). Wild type tissue cysts presented with

clusters of small amylopectin granules as well as evidence of active endodyogeny

(**Figure 9**). In contrast ΔTgLaf cysts presented bradyzoites laden with AG to the point

where other organellar structures were obscured (**Figure 9, S4**). Several bradyzoites

lacked nuclei while others lacked the cytoplasm (**Figure 9, S4**). Together, these

features are incompatible with viability. The presence of such features does not apply to

all bradyzoites within the cyst accounting for the fact that these cysts are able to initiate

a new infection (**Figure 7**.), albeit with lower virulence and cystogenic potential. These

finding suggest that AG metabolism is under tight control as dysregulated accumulation

can result in cumulative defects resulting in toxicity and death. The high frequency of

these abnormal parasites suggests that reversible glucan phosphorylation and

TgLaforin specifically represent legitimate bradyzoite specific drug targets. We recently

described a small molecule that inhibits recombinant TgLaforin (36) which serves as a

potential starting point in the development of a new class of anti-*Toxoplasma*

therapeutic agents. Particularly exciting in this context is the fact that a class of drugs

exhibiting efficacy with tissue cyst clearance (atovaquone (87, 88) endochin-like

quinolones (89-91) and JAG21 (92)), all target mitochondrial respiration. When glucose

is limiting, mitochondrial respiration can be driven by glutamine. This provides an

opportunity for combination therapy to promote the clearance of toxoplasma tissue cysts

as a means of mitigating the risk of reactivation.

METHODS

Fibroblast and parasite culture and maintenance

All parasite lines were maintained in human foreskin fibroblasts (HFFs; ATCC) in

Minimal Essential Media-α (MEM-α; Gibco) supplemented with 7% heat-inactivated fetal

bovine serum (FBS; Gemini Bio), 100 U/mL penicillin, 100 μg/mL streptomycin, and an

additional 2 mM L-glutamine (Gibco; 4 mM total L-glutamine). Cells and parasites were

incubated at 37^{o}C and 5% CO_2 in a humidified incubator. Genetically modified parasites

were maintained in MEM-α containing 7% dialyzed FBS (Gemini Bio) and either

pyrimethamine (1 μM), mycophenolic acid/xanthine (MPA: 25 μg/mL, xanthine: 50

μg/mL), or 6-thioxanthine (6-Tx: 80 μg/mL).

Assays analyzing the effects of glutamine deprivation used Dulbecco's Modified Eagle

Medium (DMEM). Both glutamine-replete (Gibco, 11966025) and depleted (Gibco,

11054020) DMEM were supplemented with 7% dialyzed FBS. Glutamine-replete media

from the supplier lacked other key nutrients and was modified to contain 5 mM glucose,

1 mM sodium pyruvate, and 4 mM L-glutamine.

Generation of T. gondii mutant lines

Type II ME49ΔHXGPRT ("WT"—the parental line utilized to generate all other lines in

this study): This line was generated in a previous study using CRISPR/Cas9 targeting of

TgHXGPRT and selection with 6-Thioxanthine (52).

TgLaforin-3xHA-HXGPRT: TgLaforin was epitope tagged with HA at the C-terminus

using CRISPR-Cas9 to disrupt the TgLaforin 3'UTR immediately downstream of the

endogenous stop codon as has been previously described (53). Briefly, a sgRNA

immediately downstream of the TgLaforin stop codon was designed using the

EuPaGDT design tool (http://grna.ctegd.uga.edu). The top hit was selected (**Table S1**)

and used to replace the sgRNA sequence in pSAG1::CAS9-U6::sgUPRT, a plasmid

containing both Cas9-green fluorescent protein (GFP) and an interchangeable sgRNA

scaffold (56); (**Table S2**). Replacement of the interchangeable sgRNA was

accomplished using a Q5 site-directed mutagenesis kit (**Table S3**) (New England

BioLabs). The TgLaforin-HA tagging construct was generated by amplifying the 3' end

of the TgLaforin-HA construct generated for complementation (see generation of COMP

line below and **Tables S2 and S3**) along with the connected HXGPRT selectable

marker. Both the TgLaforin-HA PCR-amplicon and the CRISPR-Cas9-GFP were

transfected into $1.4x10^7$ *T. gondii* ME49ΔHXGPRT parasites (2:1 insert:plasmid molar

ratio; 30 μg DNA total) by electroporation with a time constant between 0.16 and 0.20

msec (BioRad Gene Pulser II). After 24 h, surviving parasites were syringe-passaged

from infected HFFs with a 27 G needle to lyse host cells, and gravity-filtered through a

10 μm filter to remove host-cell debris. Successful transformants were then enriched by

use of fluorescence-activated cell sorting (FACS; Sony SY3200, installed in a biosafety

level II cabinet) to select parasites expressing Cas9-GFP from the transfected plasmid

by isolating GFP+ parasites. HFFs were infected with GFP+ parasites, and then placed

in media containing MPA/xanthine 24 h later to select for restoration of HXGPRT.

MPA/xanthine-resistant parasites were cloned by limiting dilution into a 96 well plate.

Wells containing single plaques were picked 7 days later and expanded. Genomic DNA

was extracted from clones using a Proteinase K treatment detailed elsewhere (93).

Successful tagging of TgLaforin was verified using sequencing, immunoblotting, and

IFA.

ME49ΔHXΔTgLaforin ("ΔTgLaf"): TgLaforin was disrupted using a CRISPR-Cas9

mediated strategy as detailed above, with several differences. Briefly, a single sgRNA

was designed to target the first exon of TgLaforin with the top hit from EuPaGDT (**Table**

S1). To disrupt TgLaforin with a selectable drug marker, DHFR-TS*, a pyrimethamine-

resistant mutant of the DHFR gene, containing a 5'-NcGra7 promotor and DHFR 3'UTR

was amplified from pJET-NcGra7_DHFR (**Table S2**). Amplification utilized primers

containing 40 nt extensions homologous to the 5'- and 3'-UTR of TgLaforin to

encourage homologous recombination-mediated whole-gene replacement with the drug

cassette (**Table S3**). Both the PCR-amplified DHFR* homology cassette and the

CRISPR-Cas-GFP plasmid were transfected and FACS-sorted as described above.

GFP+ parasites underwent drug selection in pyrimethamine. Parasites were then cloned

and expanded as detailed above. Successful integration of the DHFR* cassette into the

TgLaforin locus was verified using PCR with inside/out primer pairs to the chimeric,

interrupted gene (**Table S3**). Loss of TgLaforin transcription was verified by purifying

RNA from TgLaforin clones on RNeasy spin columns (Qiagen). Using the Promega

Reverse Transcriptase System, cDNA was synthesized from RNA extracts. Primers

designed for full-length TgLaforin amplification were then used to verify loss of

TgLaforin cDNA in knockout lines.

ME49ΔHXΔTgLaforin+ChrVI-TgLaforin ("COMP"): Complementation of TgLaforin was

also executed using a CRISPR-mediated strategy. A sgRNA to a neutral locus on

chromosome VI identified previously (58) was generated using the same mutagenesis

strategy as above (**Table S1 and S3**). A full length TgLaforin cDNA containing its

endogenous 5'UTR (2000 bp upstream from gDNA) was synthesized by GenScript and

inserted into a pHA3x-LIC vector (**Table S2**) containing a C-terminal HA tag and a

DHFR 3'UTR, linked to the HXGPRT selectable marker (named "TgLaforin-HA3x-LIC";

also used above for endogenous tagging to create the TgLaforin-HA line). The entire

construct (5'UTR:TgLaforin-cDNA:DHFR-3'UTR:HXGPRT) was amplified from the

vector and co-transfected into ΔTgLaf parasites with the CRISPR-Cas9 plasmid as done

above. Successful transformants that received the HXGPRT marker were selected with

MPA/xanthine. Successful insertion of TgLaforin along with its promoter was verified

using PCR (**Table S3**), immunoblotting, and IFA with an anti-HA antibody (Abcam).

Immunofluorescence (IF) staining

HFFs were grown on glass coverslips until confluent and subsequently infected.

Infected HFFs were fixed with either methanol (MeOH) (100%, -20^{o}C) or methanol-free

paraformaldehyde (PFA) (4% in phosphate-buffered saline (PBS); Electron Microscopy

Sciences) as indicated below for each antibody. Infected HFFs fixed with PFA were

permeabilized in 0.1% TritonX-100 in PBS++ (PBS containing 0.5 mM CaCl$_2$ and 0.5

mM MgCl$_2$) for 10 min at room temperature (RT). Primary and secondary antibodies

were diluted in 3% (w/v) bovine serum albumin (BSA; Fisher) in PBS++. Samples were

first incubated with the primary antibody (αHA-1:1,000; αSAG-1:10,000; αGAP45-

1:5,000; αGRA3-1:1500; IV58B6-1:50) at RT for 45 min, washed 3x with PBS++, and

then incubated with fluorescent secondary antibodies (1:2,000) and 4',6-diamidino-2-

phenylindole (DAPI; 300 nM) for 45 min. Secondary antibodies (Invitrogen) were

conjugated to either Oregon Green or Texas Red fluorophores and specific to the

species and class of primary antibody used. Samples were then washed 3x with PBS++

before mounting the coverslip on a glass slide using MOWIOL mounting media.

Immunofluorescence staining was visualized using a Zeiss AxioVision upright

microscope with a 100X 1.4 numerical-aperture oil immersion objective, and images

were acquired using a grayscale Zeiss AxioCam MRM digital camera. Grayscale

images were pseudo-colored in ImageJ using magenta (Texas Red), yellow (Oregon

Green), and cyan (DAPI), and further alterations to brightness and contrast were also

made in ImageJ when deemed appropriate. For all assays in which staining intensity

was compared across treatments and parasite lines, concentrations of antibodies,

exposure times, and alterations to brightness/contrast were identical. Colocalization of

fluorescent antibodies/reagents was quantified using Pearson's coefficient calculated

with the JACoP plugin on ImageJ (94).

***PAS staining*-tachyzoite and in vitro bradyzoite**

PAS staining of tachyzoites and in vitro bradyzoites was done on infected HFFs fixed in

4% PFA and permeabilized as above. Coverslips were then washed 3x in tap water

before the addition of 1% periodic acid (Sigma-Aldrich) for 5 min. Coverslips were then

washed with three changes of tap water. Schiff's reagent (diluted 1:4 in tap water) was

added for 15 min. Coverslips were subsequently washed 10x with tap water to develop

stain before being incubated with DAPI for 10 min and then mounted as above. PAS-

stained samples were visualized using fluorescence microscopy (excitation: 545 nm,

emission: 605 nm). When PAS was co-stained with antibodies, primary antibodies were

incubated with PAS-stained slides overnight in BSA at 4 ^{o}C before standard secondary

staining.

PAS labeling of methanol fixed tissue cysts was performed as described elsewhere,

using Schiff reagent diluted 1:10 in tap water as described elsewhere ().

Samples treated with acid-α-amylogucosidase (GAA) (from *Aspergillus niger*, >260

U/mL, Sigma) were incubated with GAA after permeabilization. GAA was diluted 1:50 in

50 mM sodium phthalate buffer, pH 5.5, and samples were treated for 24 h at room

temperature. Untreated controls were incubated in phthalate buffer without GAA.

Samples were then stained with PAS or IV58B6 as described in the IF-staining workflow

above.

In vitro bradyzoite conversion assay

Tachyzoites were converted to bradyzoites *in vitro* using alkaline stress as has been

done previously with several modifications (95). HFFs grown were infected with

tachyzoites in standard cell culture media. 4 h later, media was replaced with RPMI

1640 (Gibco 31800022) supplemented with 50 mM HEPES and adjusted to pH 8.2 with

NaOH. Parasites were then cultured for 2-6 days at 37°C, ambient CO_2, and sealed in

Parafilm. Media was replaced every other day to maintain the basic pH. Parasites were

fixed in PFA and stained with fluorescein conjugated *Dolichos biflorus* agglutinin (DBA;

1:1000, Vector Laboratories) and PAS. Images were obtained in grayscale on a Zeiss

AxioVision upright microscope as described above. To determine the degree of labeling

with DBA or PAS, the Fiji/ImageJ (96) was used to create a binary mask outlining cysts

that was applied to the PAS-stained image to measure the greyscale intensity of each

ROI (i.e. each individual vacuole/*in vitro* cyst).

Transmission electron microscopy of in vitro tachyzoites and bradyzoites

Transmission-electron microscopy (TEM) was performed as done previously (97).

Blocks were stained at the University of Kentucky's Imaging Center in the College of

Arts and Sciences. Blocks were trimmed and sectioned on an ultramicrotome with a

diamond knife. Sections were placed on copper grids and then contrast stained with

lead citrate. Micrographs were collected at the University of Kentucky's Electron

Microscopy Center on a Talos F200X TEM (Thermo) operated at 200 kV accelerating

voltage with a 50 μm objective aperture inserted to enhance contrast using a 16M pixel

4k x 4k CMOS camera (Ceta, Thermo Scientific). AG size and grayscale values were

measured in ImageJ.

Immunoblotting

Parasites were syringe lysed from host cells, pelleted, and $2x10^6$ parasites were

resuspended in SDS-PAGE sample buffer and boiled for 10 min before being run on a

single lane of a 10% polyacrylamide gel. The gel was then transferred to a 0.2 μm

PVDF membrane (BioRad) using a Turbotransfer System (BioRad) for 7 min at 25 V.

The PVDF membrane was blocked in 5% (w/v) non-fat milk in Tris-buffered saline plus

Tween-20 detergent (TBST; 0.1% Tween-20) for 20 min before being probed with a

primary antibody (αHA-1:1,000; αGAP45-1:5,000; αSRS9-1:1,000; αSAG1-1:10,000) in

non-fat milk overnight at 4^oC (Cell Signaling C29F4). The blot was washed 3x with

TBST before probing with either HRP-conjugated α-rabbit or α-mouse-IgG (Jackson

Laboratories). Blot was washed and developed for 5 min using SuperSignal^{TM} West

Pico PLUS (Thermo Scientific) and visualized on a GelDoc station (BioRad).

Steady state polar metabolite analysis

Parasites were prepared as previously described (61). Confluent HFFs were infected

with parasites at a multiplicity of infection (MOI) of 2 to achieve a high density of

parasites after 48 h of growth (>80% cells containing >32 parasites each). Plates

containing infected HFFs were placed on ice, media removed, and the monolayer was

washed 2X with ice-cold PBS. Parasites were harvested on ice in a 4 ^{o}C cold-room.

Cells were scraped from plate surface, resuspended in PBS (8 plates/50 mL PBS), and

centrifuged at 1000g for 10 min at 4 ^{o}C. PBS was removed, the cell pellet was

resuspended in 2 mL PBS, and syringe passaged successively in 23 G and 27 G

needles. The soluble host cell lysate was removed by centrifugation (1000g). The pellet

was resuspended in 5 mL PBS and host-cell debris was removed by syringe-filtering the

suspension through a 3 μm filter (Whatman). Filtered parasites were then pelleted,

resuspended in 1 mL PBS, and counted on a hemacytometer. Parasites were pelleted a

final time at 14,000g for 30 s at 4 ^{o}C, supernatant was removed, and pelleted parasites

were flash frozen in liquid nitrogen and stored at -80 ^{o}C until metabolite extraction.

Polar metabolite extraction: Polar metabolites were extracted in 0.5 mL -20 ^{o}C 50%

methanol (MeOH) containing 20 μM L-norvaline (procedural control) for 30 min on ice.

During the 30 min incubation, samples were regularly vortexed. Samples were then

centrifuged at 14000g for 10 min to pellet insoluble material (protein, DNA, RNA, and

glycans). Supernatant containing polar metabolites and pellet were dried separately on

a SpeedVac (Thermo) at 10^{-3} mBar until methanol (MeOH) was completely sublimated

and only dried pellet remained.

Pellet hydrolysis and extraction: Dried fraction containing protein was hydrolyzed by

resuspending the pellet in 2 N HCl (final concentration) at 95^{o}C for 2 h. Hydrolysis was

quenched, and hydrolyzed amino acids were extracted by the addition of an equal

volume of 100% MeOH with 40 μM L-norvaline such that the final concentration was

50% and 20 μM, respectively. Extraction and drying then proceeded as described

above.

Sample derivatization: Dried samples (both polar metabolites and hydrolyzed protein)

were derivatized in 70 μL 20 mg/mL methoxyamine hydrochloride in pyridine for 90 min

at 30 ^{o}C. Samples were then centrifuged at 14000g for 10 min to remove any

particulate, and 50 μL of the methoxyamine supernatant was mixed with 80 μL *N-*

methyl-N-trimethylsilyl trifluoroacetamide (MSTFA) and incubated for 30 min at 37^{o}C.

Samples were then transferred to amber glass chromatography vials and analyzed by

GC/MS.

GC/MS analysis: Metabolites were analyzed on an Agilent 7800B GC coupled to a

5977B MS detector using a previously established protocol (98). Automated Mass

Spectral Deconvolution and Identification System (AMDIS) was used to analyze

metabolites by matching metabolites to the FiehnLib metabolomics library via retention

time and fragmentation pattern. Quantification of metabolite levels was performed in

Mnova. Sample abundance was normalized to L-norvaline (procedural control) and

protein from the protein pellet (experimental control). Steady state metabolites are

presented as the mean of three independent replicates.

Conventional plaque assays

HFFs were grown in 12-well plates until confluent. HFFs were subsequently infected

with 200 parasites/well under standard cell culture conditions. Wells were washed with

PBS to remove residual invasion media, and media was changed to glutamine replete

or depleted media 4 h post-infection to allow for invasion. Plates remained undisturbed

for 10 days before the infected HFFs were fixed with 100%, -20^{o}C MeOH for 20 min,

stained with 1% crystal violet solution for 20 min, and then the plaques were de-stained

with repeated tap water washes. Zones of lysis (white clearings) could be visualized

against intact cells (purple). Images of plaques were obtained by scanning plates on an

Epson Perfection V600 photo scanner at a resolution of 600 dpi. The plaques were

measured by pixel area using ImageJ. Plaque assays were conducted on three

independent replicates, and plaque size from these experiments were aggregated to

highlight variability in plaque sizes.

Modified plaque assays

A modified plaque assay was developed to visualize foci of infection at higher

magnification where zones of clearing were not readily evident. Confluent HFF

monolayers on glass coverslips in 24 well plates were infected with WT, ΔTgLaf, and

COMP parasites in replete media for 4 hours to allow for invasion. The monolayers

were washed gently 3 times with PBS and either fresh replete media or glutamine-

depleted media were added for 3 or 6 days to appropriate wells. Infected monolayers

were washed and fixed with MeOH (-20 $^{\circ}$C) and subjected to IF using DAPI and GRA3,

an antibody that detects the PVM. Individual plaques were imaged on coded blinded

slides using a 10X objective, and their perimeters and encompassed areas were

measured using Image J. Host cell clearance within plaques was similarly measured,

and the extent of clearance was represented as a percentage for each individual

plaque. A total of three independent replicates were performed.

Egress assays

HFFs were grown to confluency in 35 mm glass bottom dishes (MatTek, P35G-0-14-C).

Two days before infecting HFFs on glass bottom dishes, both HFFs and parasites were

independently pre-treated in either glutamine-replete or -depleted media (see above for

media formulations). After 48 h pre-treatment, 10^5 parasites of each line (WT, ΔTgLaf,

and COMP) in each pre-treatment (gln+/-) were added to fresh dishes and allowed to

grow for 48 h so that most vacuoles contained >32 parasites each. Several hours before

egress, media in each infected plate was adjusted to 1.5 mL and allowed to equilibrate

at 37^{o}C in 5% CO_2. The calcium ionophore A23187 (Cayman Chemical Company) was

prepared as a 2 mM stock in DMSO and diluted in (+/-) gln media to make a 4X

concentration of 12 μM and maintained at 37^{o}C throughout the assay. Zaprinast was

likewise prepared as a 100 mM stock in DMSO and diluted into media at a 4X

concentration of 2 mM. Egress was triggered by the addition of 0.5 mL 4X A23187 to

infected HFFs (3.0 μM final concentration) or 0.5 mL 4X-zaprinast (500 μM final

concentration). Egress was monitored on a Nikon Eclipse Ti2 inverted microscope with

a 40X phase air objective modified with a 1.5X optivar. Several fields containing

vacuoles were selected from each plate, and an image was obtained 10 s after

triggering egress from each field once every 5 s for 5 mins (61 images/field) on a Nikon

DS-Ri2 color camera. Videos of each field were assembled on NIS Elements software.

Egress was monitored using standard deviation of pixel intensity and determined by

inflection point of change in standard deviation of pixel intensity. Inflection point was

calculated by fitting a gaussian curve to the first derivative of the standard deviation in

pixel intensity and calculating the mean of the curve. Technical replicates (fields on

each plate) were averaged for each biological replicate (average of fields from each

plate).

Replication assays

HFFs were grown on glass coverslips in a 24-well plate until confluent. Two days before

infecting HFFs on coverslips, both HFFs and parasites were independently pre-treated

in either glutamine-replete or -depleted media. After pre-treatment, 10^4 parasites of

each line (WT, ΔTgLaf, and COMP) in each condition (gln+/-) were added to 3

coverslips each. 24 hours later, infected HFFs were fixed in MeOH and stained with Rb-

α-SAG1 (1:10,000) and DAPI for ease of visualization. Counting of parasites/vacuole

was performed for each line/condition on coded blinded slides with the identity of

samples revealed upon completion of the counting. A total of three independent

replicates were performed.

Mouse infection studies

4- to 6-week old CBA/J mice of both sexes (Jackson Laboratories, Bar Harbor, ME)

were injected intraperitoneally (i.p.) with either 100 WT, ΔTgLaf, or COMP tachyzoites,

or with 20 tissue-cysts from brain homogenates derived from previously infected mice.

In either case, parasites/cysts were suspended in a final volume of 0.2 mL serum-free,

Opti-MEM media (Gibco). Mice were then monitored and assigned a body index score

daily. Monitoring frequency increased to twice a day once symptomatic throughout the

course of infection as previously described (52). When symptomatic, mice were

administered a gel diet and wet chow on the cage floor and given 0.25-0.5 mL saline

solution subcutaneously as needed. Moribund mice were humanely euthanized.

Euthanasia of both moribund mice and mice sacrificed at the time of tissue cyst harvest

was performed by CO_2 asphyxiation, followed by cervical dislocation. The number of

mice used for each experiment is indicated within relevant figures. All protocols were

carried out under the approval of the University of Kentucky's Institutional Animal Care

and Use Committee (IACUC).

Tissue cyst purification

Tissue cysts were purified as previously described using discontinuous Percoll

gradients (11, 70). Processing of two sex/infection-matched brains was performed on

each gradient. Cysts were collected in 1 mL fractions from the bottom of the centrifuged

Percoll gradient using a peristaltic pump adjusted to a flow rate of 2 mL/min. To quantify

tissue cysts, 10-20 μL of each fraction was placed into 100 μL PBS in the well of a 96-

well plate, pelleted and directly enumerated at 20X magnification in each well. Total

cysts per mouse were calculated by summing the total number of cysts in each fraction

and dividing the total by two to adjust for brain homogenization in pairs. Each pair of

mice was presented as a single averaged data point. Tissue cysts were pelleted onto

slides using a Cytospin centrifuge and fixed and stored in 100% MeOH (-20^{o}C) until

staining.

Quantification of AG levels in tissue cysts based on PAS labeling intensity.

Tissue cysts deposited on glass slides were fixed and stored in methanol at -20^{o}C.

Slides were equilibrated to room temperature and stained with PAS using conditions

optimized for staining tissue cysts (14). Cysts were additionally stained with DBA lectin

to demarcate the cyst boundary. Thirty randomly acquired cysts were imaged using a

fixed exposure as a z-stack with a 0.24μm step, with the center slice used for

quantification without deconvolution using AmyloQuant, a purpose developed

quantitative imaging based application (14). The distribution of pixel intensities was

defined in 4 bins representing the background (black: 0-10 grayscale), low (blue: 10-25),

intermediate (green: 25-50) and high (red: >50) proportion of pixels (14). The relative

distribution of PAS intensities across the 4 bins is represented using a stacked plot with

the distribution patterns arrayed from lowest to highest intensity for each cohort.

AmyloQuant additionally presents a spatial heat map to allow for the distribution of PAS

intensity (AG) to be revealed within the imaged cyst. The mean pixel intensity of each

cyst was additionally determined following the definition of the ROI defined by the

boundary of the cyst wall using Image J. Measurements of cyst diameters in microns

was achieved using the Zeiss Zen imaging software.

Preparation of in vivo tissue cysts for TEM imaging

To prepare _T. gondii_ tissue cysts generated _in vivo_ for TEM, cysts were isolated from

the brains of infected mice as detailed above through the counting step. The Percoll

fraction containing mouse red blood cells (RBCs) was also recovered. After combining

Percoll fractions containing tissue cysts and diluting with PBS to a volume of 15 mL

(maximum of two 1-mL fractions were combined before dilution), cysts were pelleted for

15 min at 1000g at 4 ^{o}C. To maximize cyst recovery, 10 mL supernatant was removed,

and the remaining 5 mL was divided into 1 mL fractions for the top 4 mL, and the bottom

1 mL directly above the pellet was sub-fractionated into 100 μL volumes. Typically, the

majority of cysts were localized to within 300 μL of the Percoll pellet, rather than in the

pellet itself. Sub-fractions containing cysts were once again combined and diluted

(typically 200-300 μL diluted with 1-mL PBS) in a 1.5 mL Eppendorf tube, and then

pelleted in a swinging-bucket rotor for 10 min at 1000g and 4 ^{o}C. Leaving the pellet

undisturbed, all but 50 μL of the supernatant was removed. A small volume (~5-10 μL)

of the reserved RBC fraction was added to the remaining volume for ease of visualizing

the pellet throughout the remaining processing steps.

To ensure the detection of the relatively rare cyst population, a previously described

protocol (71, 72) was adapted to concentrate the cysts into a small agarose block. A

1.33X fixative solution of glutaraldhehyde (GA) in cacodylate buffer was prepared

containing 4% GA and 133 mM sodium cacodylate. 150 μL fixative solution was then

added to the 50 μL sample, bringing the total volume to 200 μL such that the final

concentration of GA was 3% and sodium cacodylate was 100 mM. Cysts were then

incubated at room temperature for 1 h in fixative. While cysts were in fixative, 4% low-

melt agarose (BioRad) was prepared in 100 mM sodium cacodylate buffer and kept

liquid at 70 $^{\circ}$C until needed. After fixation, cysts were pelleted again at 1000g for 10 min

at room temperature in a table-top centrifuge. All but 50 μL supernatant was once again

removed, and 200 μL warm low-melt agarose was slowly added on top of fixed, pelleted

cysts (3.2% agarose, final concentration). Suspension was then centrifuged again at

1000g for 10 min at 30 $^{\circ}$C to keep the agarose semi-liquid, and then placed on ice for

20 min to solidify agarose. After solidification, entire agarose plug was removed from

tube with a small a wooden dowel that had been whittled into a thin scoop. This agarose

plug was placed in a Petri dish, and the pellet was carefully cut out of the plug with a

razor blade to create a 1 mm^3 block. The agarose block was then stored in 1X

GA/cacodylate buffer overnight at 4°C. Processing of the block from post-stain onward

was then identical to TEM processing described above. During sectioning, thick

sections cut on a glass knife were stained with toluidine blue and examined for cysts

using a light microscope prior to ultra-thin sectioning once the cyst containing later was

identified.

Data analyses

All data analyses, including graph preparation and statistics, were performed using

GraphPad Prism 9 or 10. Details on statistical tests applied are presented in the specific

figure legends.

1020

Figure Legends.

Figure 1. Glucan dynamics in *T. gondii* ME49 tachyzoites and in vitro bradyzoites.

A, Microscopy-based glucan evaluation of *T. gondii* tachyzoites and bradyzoites using

PAS (left), IV58B6 (middle; α-glycogen IgM mAb), and TEM (right). **B,** GAA digest of

AGs in tachyzoites and bradyzoites confirms specificity of IV58B6 antibody and PAS

staining. All scale bars = 5 μm.

1027

Figure 2. Endogenous epitope tagging and localization of TgLaforin.

A, Schematic depicting the TgLaforin 3xHA-epitope tagging strategy. X = stop codon. **B,**

Successful tagging of TgLaforin (62 kDa) was verified using immunoblot analysis with

an α-HA antibody with SAG1 used as a loading control. **C,** IFA of Wild type (WT) and

TgLaforin-3XHA (TgLaf-HA) tagged parasites with α-HA antibody under tachyzoite and

in vitro bradyzoite conditions. Stage conversion is confirmed by Dolichos lectin (DBA)

staining. **D,** Western blot analysis of TgLaforin expression levels in tachyzoites (T) and

in vitro bradyzoites (B). Expression of TgLaf-HA is absent under in vitro bradyzoite

conditions. GAP45 is the loading control. Decrease in SAG1 alongside increase in

SRS9 confirms tachyzoite to bradyzoite conversion. **E**, IFA of TgLaforin-HA

colocalization with PAS. Pearson's coefficient: 0.765. **F**, IFA of TgLaforin-HA

colocalization with IV58B6. Pearson's coefficient: 0.737. All scale bars = 5 μm.

1040

Figure 3. Loss of TgLaforin results in no gross glucan abnormalities under glucose

replete conditions. **A,** Schematic of TgLaforin KO strategy: the pyrimethamine-

resistance DHFR* gene containing 40-nt homologous arms (dark gray boxes on either

end of gene) was inserted into the TgLaforin locus via homologous recombination. A

double stranded break was induced using CRISPR/Cas9-GFP with a PAM site in the

first exon. **B,** Inside/out PCR verification of DHFR integration into the TgLaforin locus.

Amplicons (PCR1-4) are illustrated in (A). **C,** Loss of TgLaforin mRNA was confirmed by

amplifying full-length TgLaforin cDNA generated from both WT and ΔTgLaf strains.

Actin cDNA amplification serves both as a loading control and as a control to verify the

absence of gDNA. **D,** Analysis of glucan levels in ΔTgLaf tachyzoites using three

different approaches: PAS and IV58B6 immunofluorescence staining, and TEM.

1052

Figure 4. ΔTgLaf parasites are dependent on glutamine for normal plaque formation.

A, Steady-state metabolomics suggests upregulation of glutaminolysis in ΔTgLaf

parasites. Metabolite levels of intracellular tachyzoites were analyzed after 48 hours of

growth in HFFs by GC/MS analysis. Data were collected from 3 independent replicates.

Statistical comparisons were done using unpaired two-tailed t-tests. Statistical

significance is as follows: **p<0.01, ns=p>0.05, nd=not detected.

B, Representative plaque assays following staining of infected HFF monolayers with

crystal violet under nutrient replete (+gln) and glutamine free (gln-) conditions. WT,

ΔTgLaf (KO) and complemented (COMP) lines were evaluated.

C, Quantification of visible plaques for WT, ΔTgLaf and COMP lines under nutrient

replete and glutamine free conditions represented as the plaque area (pixels) was

measured across three independent replicates 6 days following infection.

D, Pixel area of nascent plaques/vacuoles after 3 days of growth as monitored by IFA

microscopy following staining with the PVM maker GRA3 under both glutamine replete

(+) and depleted conditions (-). **E**, Pixel area of plaques monitored as in (D) after 6 days

of growth. Statistical comparisons for C-E were done using an ordinary one-way

ANOVA using Tukey's post-hoc test to correct for multiple comparisons. Statistical

significance is indicated as follows: $*p<0.05$, $****p<0.0001$, ns=$p>0.05$.

1071

Figure 5. Loss of TgLaforin results in aberrant glucan morphology and accumulation in

in vitro bradyzoites only visible by TEM. **A,** *In vitro* tachyzoite to bradyzoite conversion

efficiency of ΔTgLaf vs WT parasites as measured by DBA intensity confirms no defect

in induced stage conversion in ΔTgLaf parasites. **B,** Change in AG levels detected

using the intensity of PAS labeling during bradyzoite conversion reveals no statistically

different levels in AG accumulation based on PAS intensity. **C,** Representative TEM

images of bradyzoites from each indicated parasite line. At 4300x magnification, scale

bar = 2 μm; at 8600x magnification, scale bar = 1 μm. Arrowhead = canonical AG

(white, round/ovoid); Arrow = aberrant AG (grey, flattened, multi-lobed). **D,**

Quantification of relative parasite AG content and **E,** AG grayness across parasite lines

using 8-bit grayscale (0=black, 255=white). Statistical comparisons were done using an

ordinary one-way ANOVA with Tukey's post-hoc test to correct for multiple

comparisons. Statistical significance is indicated as follows: *p<0.05, ***p<0.001,

****p<0.0001, ns=p>0.05.

1086

Figure 6. Loss of TgLaforin attenuates tachyzoite virulence and cyst burden in mice.

Equal numbers of male and female CBA/J mice were infected i.p. with 100 tachyzoites

from each group and monitored for both symptom progression and mortality.

A, Symptomology throughout the acute (< day 20) and early chronic phases of infection

(day 21-28). Mice were monitored 1-2x/day and assigned a body score index ranging

from asymptomatic (Stage 0) to moribund/deceased. Staging of disease progression is

discussed in (52). **B,** Kaplan-Meier curve of mouse survival throughout acute and early

chronic infection following tachyzoite infection. **C,** A cohort of mice that survived 4-

weeks were euthanized, and the cyst burden determined as done previously described

(70). Error bars depict SD from the mean. Statistical comparison for Kaplan Meier

curves is indicated on plot, and statistical comparison of cyst burden was done using

unpaired two-tailed t-tests. Statistical significance: *p<0.05, ***p<0.0002.

1099

Figure 7. Mouse infection with ∆TgLaf tissue cysts results in milder illness and lower

cyst burden. CBA/J mice were infected i.p. with 20 cysts taken from previously infected

mouse brains and monitored for symptomology and death. Tissue cysts were harvested

from mice infected for 4 or 6 weeks respectively. **A,** Symptomology throughout the

acute and early chronic phase of infection as described in Figure 6B. **B,** Kaplan-Meier

curve of mouse survival during the acute and early chronic infection reveal low overall

mortality associated with cyst initiated infections for all parasite lines. **C,** Tissue cyst

burden following a 4-week bradyzoite initiated infection. Cyst numbers were determined

as described in Figure 6C. Statistical comparison of cyst burden was done using

unpaired two-tailed t-tests. Statistical significance: $*p<0.05$, ns=$p>0.05$.

1110

Figure 8. Quantification of amylopectin levels in tissue cysts based on PAS intensity.

A. The mean intensity of PAS labeled tissue cysts measured using Image J for WT,

ΔTgLaf and COMP tissue cysts harvested at 4 and 6 weeks post infection reveal no

difference in AG levels across these parasite lines at week 4. Highly significant

differences are evident between WT/COMP and DTgLaf tissue cysts at week 6 post

infection indicating a temporal component for the emergence of the starch excess

phenotype. n=30 for each parasite line at each time point of harvest. Statistical analysis

was done using Tukey's multiple comparison test. P values: $*$ $p <0.05$, $****$ $p< 0.0001$

B. Analysis of WT, DTgLaf and COMP tissue cysts harvested at weeks 4 and 6 using

AmyloQuant. The 30 tissue cysts per sample presented in A. were analyzed using

AmyloQuant. While acquired at random, the data are arrayed from low to high PAS

intensity presented as the percentage of pixels in each classification bin. The

classification bins defined as: background: Black: 0-10 grayscale, low: Blue: 10-25

grayscale, intermediate: Green 25-50 grayscale and high intensity pixels: Red >50

grayscale. The higher overall sensitivity of AmyloQuant reveals increased accumulation

of AG at week 4 in the DTgLAf and COMP lines. This phenotype is greatly exaggerated

at week 6 post infection consistent with a temporal component for the phenotypic

manifestation of the starch excess phenotype. The pattern in the COMP parasites is

intermediate between the WT and ΔTgLaf cysts. The distribution of PAS intensity levels

within the imaged tissue cysts generated by AmyloQuant for cysts in 5 cyst intervals is

presented under each set of stacked plots.

1132

Figure 9. Transmission electron microscopy of purified tissue cyst from infected mouse

brains confirms the presence of aberrant AG accumulation within ΔTgLaf cysts. The

accumulation of amylopectin granules within the cytoplasm of WT bradyzoites with

parasites exhibiting different levels. Additionally, evidence of active endodyogeny is

present. ΔTgLaf tissue cysts contain a mix of bradyzoites with expected cytoplasmic

and organellar contrast as well as others with grossly exaggerated AG levels that and

the apparent loss of both the cytoplasmic and organellar contents. The COMP line

exhibits AG accumulation levels similar to that observed in WT parasites with the tissue

cyst itself appearing to be very tightly packed, with high levels of granular material

between individual bradyzoites. Upper panels: scale bar = 5 μm; lower panels (zoom of

boxed region from upper panel): scale bar = 2 μm.

1144

Supplemental Files

Supplemental File 1.

Figure S1. Schematic of TgLaforin complementation strategy and confirmation of

successful expression of TgLaforin. **A.** Schematic of TgLaforin complementation into

ΔTgLaf parasites in which a PAM site was chosen at a neutral locus previously

identified in chromosome VI (99) to insert TgLaforin cDNA under its endogenous

promoter. The TgLaforin construct was connected to the HXGPRT selectable drug

marker and inserted using NHEJ. **B.** PCR confirmation of integration of TgLaforin

construct into chromosome VI. Primer sets are indicated above amplicons. WT primers

amplify the same locus as in Figure 3 ("PCR 1"), also present in the COMP line. KO

primers amplify the chimeric locus depicted in Figure 3 ("PCR 2"). Presence of the KO

amplicon confirms that KO locus remains intact in COMP line. COMP primers amplify

the chimeric locus generated upon insertion of the complementation construct. VI

primers amplify the native chromosome VI locus, which is lost only in the COMP line. **C,**

Western blot confirms expression of TgLaforin-HA in complemented parasites. Tagged

LAF-HA parasites serve as a comparison to confirm the correct MW (62 kDa) and

expression level. **D,** IFA demonstrates restoration of cytoplasmic, punctate localization

of TgLaforin. Scale bar = 5 μm.

1163

Supplemental File 2

Figure S2. Loss of TgLaforin results in cumulative defects that cannot be pinpointed to

a single aspect of lytic cycle. **A.** Calcium ionophore-stimulated egress assay in which

parasites were pre-starved of glutamine for 48 hours, seeded onto HFFs and allowed to

grow for 48 hours to produce vacuoles containing >16 parasites, and stimulated with 3

μM A23187. Egress was monitored by video microscopy, and time to egress was

monitored as described in Materials and Methods. Data is the average of 3 biological

replicates that each consist of 4-5 technical replicates. COMP experiments measured 2

biological replicates. **B.** Zaprinast stimulated egress assay performed as described for

ionophore, however 500-μM zaprinast was used to stimulate egress. Data is the

average of 3 biological replicates that each consist of 2-5 technical replicates. COMP

experiments measured 2 biological replicates. **C.** Replication assay in which parasites

were pre-starved of glutamine for 48 hours, re-seeded into HFFs, and counted after 24

hours of growth. Numbers (2, 4, or 8) indicate the number of tachyzoites counted per

vacuole. Data is the average of 3 biological replicates with at least 70 vacuoles counted

per replicate. **D.** Representative images of plaque formation at days 3 and 6, +/-

glutamine. Images were taken at 10X magnification using a SAG1 antibody to visualize

vacuoles and developing plaque size. The boundaries of vacuoles/ plaques were traced

manually to define their area. Regions of cell clearance was additionally scored. **E,**

Percent of plaques cleared was measured at both days 3 and 6 by dividing the area of

the clearing by total plaque size. Statistical comparisons were done using an ordinary

one-way ANOVA using Tukey's post-hoc test to correct for multiple comparisons. Error

bars depict SD from the mean. Statistical significance is indicated as follows:

ns=p>0.05.

1188

Supplemental File 3

Figure S3. The loss of TgLaforin does not have any impact on distribution of tissue

cyst sizes. **A**. Tissue cyst diameters of 30 cysts each per line (WT, ΔTgLaf, COMP) per

time point (Week 4 and Week 6) were measured using DBA labeled cyst wall as the

delimiter using the Zeiss Zen software functionality. While all lines exhibit considerable

variability, no statistically significant differences in cyst size were noted in the DTgLaf

mutant relative to WT and COMP cysts. **B**. The relationship between tissue cyst size

and AG accumulation defined by mean PAS intensity was not found to have any

significant correlation.

1198

Supplemental File 4

Figure S4. Additional images of *in* vivo tissue cysts. **A.** A WT tissue cyst with two

zoomed in areas (orange and yellow dashed line boxes) exhibit variability in the levels

of AG and present with well defined cytoplasmic and organellar contents. **B.** A ΔTgLaf

tissue cyst presents with a mix of bradyzoites with both cytoplasmic and organelles as

well as bradyzoites lacking clearly defined organelles and/or cytoplasm. In addition

enucleated bradyzoites are evident (asterisks) as well as the displacement of rhoptries

by elevated levels of AG in the cytoplasm. Left column scale bar = 5 μm; right column

zoom scale bars = 2 μm.

1208

1209 **Supplemental File 5**

1210 **Supplemental Table 1**. sgRNA sequences used in this study.

1211

1212 **Supplemental File 6**

1213 **Supplemental Table 2.** Plasmid constructs used in this study.

1214

Supplemental File 7

Supplemental Table 3. DNA primer sequences and their application in this study

----

Acknowledgements: The authors wish to acknowledge Dr. Elizabeth Watts who

conducted early exploratory studies that encouraged the pursuit of this project, Jim

Begley (Imaging Center, University of Kentucky) for preparation of EM samples and

willingness to explore new methodologies for isolating rare *in vivo* cysts, Jillian Cramer

(Electron Microscopy Center, University of Kentucky) for acquisition of TEM images,

and Jennifer Strange (Flow Cytometry and Immune Monitoring Core Facility, University

of Kentucky) for assistance with isolation of GFP+ *T. gondii* mutants using flow

cytometry.

We additionally thank Dr's David Sibley (Washington University) and Peter Bradley

(UCLA) for their gifts of DNA constructs and antibodies.

1228

Author contributions:

Conceptualization (R.D.M., A.P.S., M.S.G.), Methodology (R.D.M., A.D., C.O.B., A.T.,

A.P.S., M.S.G.), Formal analysis (R.D.M., C.A.T., J.S.M., L.E.A.Y., R.C.S., A.T, A.P.,

C.W.V.K., A.P.S.), Investigation (R.D.M., C.A.T., J.S.M., L.E.A.Y., A.P.S.), Resources

(M.S.G., A.P.S.), Writing—original draft (R.D.M., A.P.S.), Writing—review and editing

(all authors), Visualization (R.D.M., C.A.T., J.S.M., A.P.S.), Supervision (M.S.G.,

A.P.S.), Project administration (R.D.M., M.S.G., A.P.S.), Funding acquisition (R.D.M.,

C.W.V.K., M.S.G., A.P.S.).

1237

1238 **Funding:** This work was supported by: GRFP 1247392 to R.D.M., NIH grants R21

1239 AI150631 to A.P.S and R01 AI145335 to A.P.S. and A.P, and NIH grants R35

1242

Conflicts of Interest: none

1244

Abbreviations: AG, amylopectin granule; CDPK2, calcium dependent protein kinase2; COMP, TgLaforin-complemented parasite line; CRISPR, clustered regularly interspersed short palindromic repeats; DBA, *Dolichos biflorus* agglutinin lectin; ΔTgLaf, TgLaforin-KO parasite line; DHFR, dihydrofolate reductase; FACS, fluorescence activated cell sorting; GAA, acid-α-amyloglucosidase; Gal-NAc, N-acetylgalactosamine; GAPDH1, glyceraldehyde-3-phosphate dehydrogenase; GC/MS, gas chromatography/mass spectrometry; GFP, green fluorescent protein; GT1, glucose transporter1; GWD, glucan, water di-kinase; HA, hemagglutinin; HFF, human foreskin fibroblasts; HK, hexokinase; HR, homologous recombination; HXGPRT, hypoxanthine-xanthine-guanine phosphoribosyl transferase; IF, immunofluorescence; i.p., intraperitoneally; KO, knockout; MPA, mycophenolic acid; PAM, protospacer adjacent motif; PAS, periodic acid-Schiff; PV, parasitophorous vacuole; PWD, phospho-glucan, water di-kinase; PYK1, pyruvate kinase1; RBC, red blood cell; RT, room temperature; SEX4, starch-excess4; SS, starch/glycogen synthase; TCA, tricarboxylic acid; TEM, transmission electron microscopy; UTR, untranslated region; WT, parental ME49ΔHXGPRT parasite line used in this study.

1261

Literature Cited

1. Flegr J, Prandota J, Sovickova M, Israili ZH. 2014. Toxoplasmosis--a global threat.

Correlation of latent toxoplasmosis with specific disease burden in a set of 88

countries. PLoS One 9:e90203.

2. Tenter AM, Heckeroth AR, Weiss LM. 2000. *Toxoplasma gondii*: from animals to

humans. Int J Parasitol 30:1217-58.

3. Jones JL, Dubey JP. 2012. Foodborne toxoplasmosis. Clin Infect Dis 55:845-51.

4. Derouin F, Garin YJ. 1991. *Toxoplasma gondii*: blood and tissue kinetics during

acute and chronic infections in mice. Exp Parasitol 73:460-8.

5. Pittman KJ, Knoll LJ. 2015. Long-Term Relationships: the Complicated Interplay

between the Host and the Developmental Stages of *Toxoplasma gondii* during

Acute and Chronic Infections. Microbiol Mol Biol Rev 79:387-401.

6. Zhao XY, Ewald SE. 2020. The molecular biology and immune control of chronic

Toxoplasma gondii infection. J Clin Invest 130:3370-3380.

7. Luft BJ, Hafner R, Korzun AH, Leport C, Antoniskis D, Bosler EM, Bourland DD, 3rd,

Uttamchandani R, Fuhrer J, Jacobson J, et al. 1993. Toxoplasmic encephalitis in

patients with the acquired immunodeficiency syndrome. Members of the ACTG

077p/ANRS 009 Study Team. N Engl J Med 329:995-1000.

8. Luft BJ, Remington JS. 1992. Toxoplasmic encephalitis in AIDS. Clin Infect Dis

15:211-22.

9. Cerutti A, Blanchard N, Besteiro S. 2020. The Bradyzoite: A Key Developmental

Stage for the Persistence and Pathogenesis of Toxoplasmosis. Pathogens 9.

10. Sinai AP, Watts EA, Dhara A, Murphy RD, Gentry MS, Patwardhan A. 2016.

Reexamining Chronic *Toxoplasma gondii* Infection: Surprising Activity for a

"Dormant" Parasite. Curr Clin Microbiol Rep 3:175-185.

11. Watts E, Zhao Y, Dhara A, Eller B, Patwardhan A, Sinai AP. 2015. Novel

Approaches Reveal that *Toxoplasma gondii* Bradyzoites within Tissue Cysts Are

Dynamic and Replicating Entities In Vivo. MBio 6:e01155-15.

12. Place BC, Troublefield C, Murphy RD, Sinai AP, Patwardhan A. 2021. Computer

Aided Image Processing to Facilitate Determination of Congruence in Manual

Classification of Mitochondrial Morphologies in *Toxoplasma gondii* Tissue Cysts.

Annu Int Conf IEEE Eng Med Biol Soc 2021:3509-3513.

13. Place BC, Troublefield CA, Murphy RD, Sinai AP, Patwardhan AR. 2023. Machine

learning based classification of mitochondrial morphologies from fluorescence

microscopy images of *Toxoplasma gondii* cysts. PLoS One 18:e0280746.

14. Tripathi A, Donkin RW, Miracle JS, Murphy RD, Patwardhan A, Sinai AP. 2024.

Dynamics of amylopectin granule accumulation during the course of the chronic

Toxoplasma infection is linked to intra-cyst bradyzoite replication. bioRxiv

doi:10.1101/2024.09.02.610794:2024.09.02.610794.

15. MacNeill GJ, Mehrpouyan S, Minow MAA, Patterson JA, Tetlow IJ, Emes MJ. 2017.

Starch as a source, starch as a sink: the bifunctional role of starch in carbon

allocation. J Exp Bot 68:4433-4453.

16. Dubey JP, Lindsay DS, Speer CA. 1998. Structures of *Toxoplasma gondii*

tachyzoites, bradyzoites, and sporozoites and biology and development of tissue

cysts. Clin Microbiol Rev 11:267-99.

17. Coppin A, Dzierszinski F, Legrand S, Mortuaire M, Ferguson D, Tomavo S. 2003.

1308   Developmentally regulated biosynthesis of carbohydrate and storage

1309   polysaccharide during differentiation and tissue cyst formation in *Toxoplasma*

1310   *gondii*. Biochimie 85:353-61.

1311   18. Guerardel Y, Leleu D, Coppin A, Lienard L, Slomianny C, Strecker G, Ball S,

1312   Tomavo S. 2005. Amylopectin biogenesis and characterization in the protozoan

1313   parasite *Toxoplasma gondii*, the intracellular development of which is restricted in

1314   the HepG2 cell line. Microbes Infect 7:41-8.

1315   19. Coppin A, Varre JS, Lienard L, Dauvillee D, Guerardel Y, Soyer-Gobillard MO,

1316   Buleon A, Ball S, Tomavo S. 2005. Evolution of plant-like crystalline storage

1317   polysaccharide in the protozoan parasite *Toxoplasma gondii* argues for a red

1318   alga ancestry. J Mol Evol 60:257-67.

1319   20. Uboldi AD, McCoy JM, Blume M, Gerlic M, Ferguson DJ, Dagley LF, Beahan CT,

1320   Stapleton DI, Gooley PR, Bacic A, Masters SL, Webb AI, McConville MJ, Tonkin

1321   CJ. 2015. Regulation of Starch Stores by a Ca(2+)-Dependent Protein Kinase Is

1322   Essential for Viable Cyst Development in *Toxoplasma gondii*. Cell Host Microbe

1323   18:670-81.

1324   21. Yang J, He Z, Chen C, Zhao J, Fang R. 2022. Starch Branching Enzyme 1 Is

1325   Important for Amylopectin Synthesis and Cyst Reactivation in *Toxoplasma*

1326   *gondii*. Microbiol Spectr 10:e0189121.

1327   22. Yang J, Yang C, Qian J, Li F, Zhao J, Fang R. 2020. *Toxoplasma gondii* alpha-

1328   amylase deletion mutant is a promising vaccine against acute and chronic

1329   toxoplasmosis. Microb Biotechnol 13:2057-2069.

23. Lyu C, Yang X, Yang J, Hou L, Zhou Y, Zhao J, Shen B. 2021. Role of amylopectin

synthesis in *Toxoplasma gondii* and its implication in vaccine development

against toxoplasmosis. Open Biol 11:200384.

24. Blume M, Nitzsche R, Sternberg U, Gerlic M, Masters SL, Gupta N, McConville MJ.

2015. A *Toxoplasma gondii* Gluconeogenic Enzyme Contributes to Robust

Central Carbon Metabolism and Is Essential for Replication and Virulence. Cell

Host Microbe 18:210-20.

25. Ralton JE, Sernee MF, McConville MJ. 2021. Evolution and function of carbohydrate

reserve biosynthesis in parasitic protists. Trends Parasitol 37:988-1001.

26. Sernee MF, Ralton JE, Nero TL, Sobala LF, Kloehn J, Vieira-Lara MA, Cobbold SA,

Stanton L, Pires DEV, Hanssen E, Males A, Ward T, Bastidas LM, van der Peet

PL, Parker MW, Ascher DB, Williams SJ, Davies GJ, McConville MJ. 2019. A

Family of Dual-Activity Glycosyltransferase-Phosphorylases Mediates Mannogen

Turnover and Virulence in Leishmania Parasites. Cell Host Microbe 26:385-399

e9.

27. Chen P, Lyu CC, Wang YD, Pan M, Lin XY, Shen B. 2023. Key roles of amylopectin

synthesis and degradation enzymes in the establishment and reactivation of

chronic toxoplasmosis. Animal Diseases 3.

28. Stitt M, Zeeman SC. 2012. Starch turnover: pathways, regulation and role in growth.

Curr Opin Plant Biol 15:282-92.

29. Kotting O, Kossmann J, Zeeman SC, Lloyd JR. 2010. Regulation of starch

metabolism: the age of enlightenment? Curr Opin Plant Biol 13:321-9.

30. Silver DM, Kotting O, Moorhead GB. 2014. Phosphoglucan phosphatase function

sheds light on starch degradation. Trends Plant Sci 19:471-8.

31. Ritte G, Lloyd JR, Eckermann N, Rottmann A, Kossmann J, Steup M. 2002. The

starch-related R1 protein is an alpha -glucan, water dikinase. Proc Natl Acad Sci

U S A 99:7166-71.

32. Kotting O, Pusch K, Tiessen A, Geigenberger P, Steup M, Ritte G. 2005.

Identification of a novel enzyme required for starch metabolism in Arabidopsis

leaves. The phosphoglucan, water dikinase. Plant Physiol 137:242-52.

33. Edner C, Li J, Albrecht T, Mahlow S, Hejazi M, Hussain H, Kaplan F, Guy C, Smith

SM, Steup M, Ritte G. 2007. Glucan, water dikinase activity stimulates

breakdown of starch granules by plastidial beta-amylases. Plant Physiol 145:17-

28.

34. Hejazi M, Fettke J, Haebel S, Edner C, Paris O, Frohberg C, Steup M, Ritte G. 2008.

Glucan, water dikinase phosphorylates crystalline maltodextrins and thereby

initiates solubilization. Plant J 55:323-34.

35. Hansen PI, Spraul M, Dvortsak P, Larsen FH, Blennow A, Motawia MS, Engelsen

SB. 2009. Starch phosphorylation--maltosidic restrains upon 3'- and 6'-

phosphorylation investigated by chemical synthesis, molecular dynamics and

NMR spectroscopy. Biopolymers 91:179-93.

36. Murphy RD, Chen T, Lin J, He R, Wu L, Pearson CR, Sharma S, Vander Kooi CD,

Sinai AP, Zhang ZY, Vander Kooi CW, Gentry MS. 2022. The Toxoplasma

glucan phosphatase TgLaforin utilizes a distinct functional mechanism that can

be exploited by therapeutic inhibitors. J Biol Chem 298:102089.

37. Kotting O, Santelia D, Edner C, Eicke S, Marthaler T, Gentry MS, Comparot-Moss S,

Chen J, Smith AM, Steup M, Ritte G, Zeeman SC. 2009. STARCH-EXCESS4 Is

a Laforin-Like Phosphoglucan Phosphatase Required for Starch Degradation in

Arabidopsis thaliana. Plant Cell 21:334-46.

38. Minassian BA, Lee JR, Herbrick JA, Huizenga J, Soder S, Mungall AJ, Dunham I,

Gardner R, Fong CY, Carpenter S, Jardim L, Satishchandra P, Andermann E,

Snead OC, 3rd, Lopes-Cendes I, Tsui LC, Delgado-Escueta AV, Rouleau GA,

Scherer SW. 1998. Mutations in a gene encoding a novel protein tyrosine

phosphatase cause progressive myoclonus epilepsy. Nat Genet 20:171-4.

39. Gentry MS, Guinovart JJ, Minassian BA, Roach PJ, Serratosa JM. 2018. Lafora

disease offers a unique window into neuronal glycogen metabolism. J Biol Chem

293:7117-7125.

40. Duran J, Hervera A, Markussen KH, Varea O, Lopez-Soldado I, Sun RC, Del Rio JA,

Gentry MS, Guinovart JJ. 2021. Astrocytic glycogen accumulation drives the

pathophysiology of neurodegeneration in Lafora disease. Brain 144:2349-2360.

41. Sugi T, Tu V, Ma Y, Tomita T, Weiss LM. 2017. *Toxoplasma gondii* Requires

Glycogen Phosphorylase for Balancing Amylopectin Storage and for Efficient

Production of Brain Cysts. MBio 8.

42. Cao XZ, Wang JL, Elsheikha HM, Li TT, Sun LX, Liang QL, Zhang ZW, Lin RQ.

2019. Characterization of the Role of Amylo-Alpha-1,6-Glucosidase Protein in the

Infectivity of *Toxoplasma gondii*. Front Cell Infect Microbiol 9:418.

43. Wang JL, Li TT, Elsheikha HM, Liang QL, Zhang ZW, Wang M, Sibley LD, Zhu XQ.

2022. The protein phosphatase 2A holoenzyme is a key regulator of starch

metabolism and bradyzoite differentiation in *Toxoplasma gondii*. Nat Commun

13:7560.

44. Zhao M, Yang Y, Shi Y, Chen X, Yang Y, Pan L, Du Z, Sun H, Yao C, Ma G, Du A.

2023. PP2Acalpha-B'/PR61 Holoenzyme of *Toxoplasma gondii* Is Required for

the Amylopectin Metabolism and Proliferation of Tachyzoites. Microbiol Spectr

11:e0010423.

45. Gentry MS, Dowen RH, 3rd, Worby CA, Mattoo S, Ecker JR, Dixon JE. 2007. The

phosphatase laforin crosses evolutionary boundaries and links carbohydrate

metabolism to neuronal disease. J Cell Biol 178:477-88.

46. Gentry MS, Pace RM. 2009. Conservation of the glucan phosphatase laforin is

linked to rates of molecular evolution and the glycogen metabolism of the

organism. BMC Evol Biol 9:138.

47. Baba O. 1993. [Production of monoclonal antibody that recognizes glycogen and its

application for immunohistochemistry]. Kokubyo Gakkai Zasshi 60:264-87.

48. van de Weerd R, Berbis MA, Sparrius M, Maaskant JJ, Boot M, Paauw NJ, de Vries

N, Boon L, Baba O, Canada FJ, Geurtsen J, Jimenez-Barbero J, Appelmelk BJ.

2015. A murine monoclonal antibody to glycogen: characterization of epitope-fine

specificity by saturation transfer difference (STD) NMR spectroscopy and its use

in mycobacterial capsular alpha-glucan research. Chembiochem 16:977-89.

49. Oe Y, Baba O, Ashida H, Nakamura KC, Hirase H. 2016. Glycogen distribution in

the microwave-fixed mouse brain reveals heterogeneous astrocytic patterns. Glia

64:1532-45.

50. Skurat AV, Segvich DM, DePaoli-Roach AA, Roach PJ. 2017. Novel method for

detection of glycogen in cells. Glycobiology 27:416-424.

51. Guimaraes EV, de Carvalho L, Barbosa HS. 2003. An alternative technique to reveal

polysaccharides in *Toxoplasma gondii* tissue cysts. Mem Inst Oswaldo Cruz

98:915-7.

52. Troublefield CA, Miracle JS, Murphy RD, Donkin RW, Sinai AP. 2023. Factors

Influencing Tissue Cyst Yield in a Murine Model of Chronic Toxoplasmosis. Infect

Immun 91:e0056622.

53. Sidik SM, Hackett CG, Tran F, Westwood NJ, Lourido S. 2014. Efficient genome

engineering of *Toxoplasma gondii* using CRISPR/Cas9. PLoS One 9:e100450.

54. Fritz HM, Buchholz KR, Chen X, Durbin-Johnson B, Rocke DM, Conrad PA,

Boothroyd JC. 2012. Transcriptomic analysis of toxoplasma development reveals

many novel functions and structures specific to sporozoites and oocysts. PLoS

One 7:e29998.

55. Choi CP, Moon AS, Back PS, Jami-Alahmadi Y, Vashisht AA, Wohlschlegel JA,

Bradley PJ. 2019. A photoactivatable crosslinking system reveals protein

interactions in the *Toxoplasma gondii* inner membrane complex. PLoS Biol

17:e3000475.

56. Shen B, Brown KM, Lee TD, Sibley LD. 2014. Efficient gene disruption in diverse

strains of *Toxoplasma gondii* using CRISPR/CAS9. mBio 5:e01114-14.

57. Sidik SM, Huet D, Ganesan SM, Huynh MH, Wang T, Nasamu AS, Thiru P, Saeij

JP, Carruthers VB, Niles JC, Lourido S. 2016. A Genome-wide CRISPR Screen

in Toxoplasma Identifies Essential Apicomplexan Genes. Cell 166:1423-1435

e12.

58. Markus BM, Bell GW, Lorenzi HA, Lourido S. 2019. Optimizing Systems for Cas9

Expression in *Toxoplasma gondii*. mSphere 4.

59. Sun RC, Young LEA, Bruntz RC, Markussen KH, Zhou Z, Conroy LR, Hawkinson

TR, Clarke HA, Stanback AE, Macedo JKA, Emanuelle S, Brewer MK, Rondon

AL, Mestas A, Sanders WC, Mahalingan KK, Tang B, Chikwana VM, Segvich

DM, Contreras CJ, Allenger EJ, Brainson CF, Johnson LA, Taylor RE, Armstrong

DD, Shaffer R, Waechter CJ, Vander Kooi CW, DePaoli-Roach AA, Roach PJ,

Hurley TD, Drake RR, Gentry MS. 2021. Brain glycogen serves as a critical

glucosamine cache required for protein glycosylation. Cell Metab 33:1404-1417

e9.

60. Brewer MK, Uittenbogaard A, Austin GL, Segvich DM, DePaoli-Roach A, Roach PJ,

McCarthy JJ, Simmons ZR, Brandon JA, Zhou Z, Zeller J, Young LEA, Sun RC,

Pauly JR, Aziz NM, Hodges BL, McKnight TR, Armstrong DD, Gentry MS. 2019.

Targeting Pathogenic Lafora Bodies in Lafora Disease Using an Antibody-

Enzyme Fusion. Cell Metab 30:689-705 e6.

61. King EFB, Cobbold SA, Uboldi AD, Tonkin CJ, McConville MJ. 2020. Metabolomic

Analysis of *Toxoplasma gondii* Tachyzoites. Methods Mol Biol 2071:435-452.

62. Nitzsche R, Zagoriy V, Lucius R, Gupta N. 2016. Metabolic Cooperation of Glucose

and Glutamine Is Essential for the Lytic Cycle of Obligate Intracellular Parasite

Toxoplasma gondii. J Biol Chem 291:126-41.

63. Oppenheim RD, Creek DJ, Macrae JI, Modrzynska KK, Pino P, Limenitakis J,

Polonais V, Seeber F, Barrett MP, Billker O, McConville MJ, Soldati-Favre D.

2014. BCKDH: the missing link in apicomplexan mitochondrial metabolism is

required for full virulence of *Toxoplasma gondii* and Plasmodium berghei. PLoS

Pathog 10:e1004263.

64. Endo T, Sethi KK, Piekarski G. 1982. *Toxoplasma gondii*: calcium ionophore

A23187-mediated exit of trophozoites from infected murine macrophages. Exp

Parasitol 53:179-88.

65. Lourido S, Tang K, Sibley LD. 2012. Distinct signalling pathways control Toxoplasma

egress and host-cell invasion. EMBO J 31:4524-34.

66. Hotchin J, Kinch W, Benson L. 1971. Lytic and turbid plaque-type mutants of

lymphocytic choriomeningitis virus as a cause of neurological disease or

persistent infection. Infect Immun 4:281-6.

67. Fiala M. 1968. Plaque formation by 55 rhinovirus serotypes. Appl Microbiol 16:1445-

50.

68. Ferguson DJ. 2004. Use of molecular and ultrastructural markers to evaluate stage

conversion of *Toxoplasma gondii* in both the intermediate and definitive host. Int

J Parasitol 34:347-60.

69. Sinai AP. 2008. Biogenesis of and activities at the *Toxoplasma gondii*

parasitophorous vacuole membrane. Subcell Biochem 47:155-64.

70. Watts EA, Dhara A, Sinai AP. 2017. Purification *Toxoplasma gondii* Tissue Cysts

Using Percoll Gradients. Curr Protoc Microbiol 45:20C 2 1-20C 2 19.

71. Kumar S, Ciraolo G, Hinge A, Filippi MD. 2014. An efficient and reproducible

process for transmission electron microscopy (TEM) of rare cell populations. J

Immunol Methods 404:87-90.

72. Kumar S, Filippi MD. 2016. An Alternative Approach for Sample Preparation with

Low Cell Number for TEM Analysis. J Vis Exp doi:10.3791/54724.

73. Sinai AP, Suvorova ES. 2020. The RESTRICTION checkpoint: a window of

opportunity governing developmental transitions in *Toxoplasma gondii*. Curr Opin

Microbiol 58:99-105.

74. Xue Y, Theisen TC, Rastogi S, Ferrel A, Quake SR, Boothroyd JC. 2020. A single-

parasite transcriptional atlas of *Toxoplasma gondii* reveals novel control of

antigen expression. Elife 9.

75. Olson WJ, Martorelli Di Genova B, Gallego-Lopez G, Dawson AR, Stevenson D,

Amador-Noguez D, Knoll LJ. 2020. Dual metabolomic profiling uncovers

Toxoplasma manipulation of the host metabolome and the discovery of a novel

parasite metabolic capability. PLoS Pathog 16:e1008432.

76. Zeeman SC, Delatte T, Messerli G, Umhang M, Stettler M, Mettler T, Streb S,

Reinhold H, Kotting O. 2007. Starch breakdown: recent discoveries suggest

distinct pathways and novel mechanisms. Funct Plant Biol 34:465-473.

77. Zeeman SC, Smith SM, Smith AM. 2007. The diurnal metabolism of leaf starch.

Biochem J 401:13-28.

78. Mahlow S, Orzechowski S, Fettke J. 2016. Starch phosphorylation: insights and

perspectives. Cell Mol Life Sci 73:2753-64.

79. Ferguson DJ, Hutchison WM. 1987. An ultrastructural study of the early
development and tissue cyst formation of *Toxoplasma gondii* in the brains of
mice. Parasitol Res 73:483-91.

80. Niittyla T, Comparot-Moss S, Lue WL, Messerli G, Trevisan M, Seymour MD,
Gatehouse JA, Villadsen D, Smith SM, Chen J, Zeeman SC, Smith AM. 2006.
Similar protein phosphatases control starch metabolism in plants and glycogen
metabolism in mammals. J Biol Chem 281:11815-8.

81. Zeeman SC, Northrop F, Smith AM, Rees T. 1998. A starch-accumulating mutant of
Arabidopsis thaliana deficient in a chloroplastic starch-hydrolysing enzyme. Plant
J 15:357-65.

82. Shukla A, Olszewski KL, Llinas M, Rommereim LM, Fox BA, Bzik DJ, Xia D,
Wastling J, Beiting D, Roos DS, Shanmugam D. 2018. Glycolysis is important for
optimal asexual growth and formation of mature tissue cysts by *Toxoplasma*
gondii. Int J Parasitol 48:955-968.

83. Blume M, Rodriguez-Contreras D, Landfear S, Fleige T, Soldati-Favre D, Lucius R,
Gupta N. 2009. Host-derived glucose and its transporter in the obligate
intracellular pathogen *Toxoplasma gondii* are dispensable by glutaminolysis.
Proc Natl Acad Sci U S A 106:12998-3003.

84. Dubey R, Staker BL, Foe IT, Bogyo M, Myler PJ, Ngo HM, Gubbels MJ. 2017.
Membrane skeletal association and post-translational allosteric regulation of
Toxoplasma gondii GAPDH1. Mol Microbiol 103:618-634.

85. Xia N, Ye S, Liang X, Chen P, Zhou Y, Fang R, Zhao J, Gupta N, Yang S, Yuan J,

Shen B. 2019. Pyruvate Homeostasis as a Determinant of Parasite Growth and

Metabolic Plasticity in *Toxoplasma gondii*. mBio 10.

86. Suresh K, Mak JW, Yong HS. 1991. Immune response in acute Toxoplasma

infection of Balb/C, ICR and CBA/J mice. Southeast Asian J Trop Med Public

Health 22:452-4.

87. Araujo FG, Huskinson-Mark J, Gutteridge WE, Remington JS. 1992. In vitro and in

vivo activities of the hydroxynaphthoquinone 566C80 against the cyst form of

Toxoplasma gondii. Antimicrob Agents Chemother 36:326-30.

88. Araujo FG, Lin T, Remington JS. 1993. The activity of atovaquone (566C80) in

murine toxoplasmosis is markedly augmented when used in combination with

pyrimethamine or sulfadiazine. J Infect Dis 167:494-7.

89. Doggett JS, Nilsen A, Forquer I, Wegmann KW, Jones-Brando L, Yolken RH,

Bordon C, Charman SA, Katneni K, Schultz T, Burrows JN, Hinrichs DJ, Meunier

B, Carruthers VB, Riscoe MK. 2012. Endochin-like quinolones are highly

efficacious against acute and latent experimental toxoplasmosis. Proc Natl Acad

Sci U S A 109:15936-41.

90. Doggett JS, Schultz T, Miller AJ, Bruzual I, Pou S, Winter R, Dodean R, Zakharov

LN, Nilsen A, Riscoe MK, Carruthers VB. 2020. Orally Bioavailable Endochin-

Like Quinolone Carbonate Ester Prodrug Reduces *Toxoplasma gondii* Brain

Cysts. Antimicrob Agents Chemother 64.

91. Vidadala RS, Rivas KL, Ojo KK, Hulverson MA, Zambriski JA, Bruzual I, Schultz TL,

Huang W, Zhang Z, Scheele S, DeRocher AE, Choi R, Barrett LK, Siddaramaiah

LK, Hol WG, Fan E, Merritt EA, Parsons M, Freiberg G, Marsh K, Kempf DJ,

Carruthers VB, Isoherranen N, Doggett JS, Van Voorhis WC, Maly DJ. 2016.

Development of an Orally Available and Central Nervous System (CNS)

Penetrant *Toxoplasma gondii* Calcium-Dependent Protein Kinase 1 (TgCDPK1)

Inhibitor with Minimal Human Ether-a-go-go-Related Gene (hERG) Activity for

the Treatment of Toxoplasmosis. J Med Chem 59:6531-46.

92. McPhillie MJ, Zhou Y, Hickman MR, Gordon JA, Weber CR, Li Q, Lee PJ,

Amporndanai K, Johnson RM, Darby H, Woods S, Li ZH, Priestley RS, Ristroph

KD, Biering SB, El Bissati K, Hwang S, Hakim FE, Dovgin SM, Lykins JD,

Roberts L, Hargrave K, Cong H, Sinai AP, Muench SP, Dubey JP, Prud'homme

RK, Lorenzi HA, Biagini GA, Moreno SN, Roberts CW, Antonyuk SV, Fishwick

CWG, McLeod R. 2020. Potent Tetrahydroquinolone Eliminates Apicomplexan

Parasites. Front Cell Infect Microbiol 10:203.

93. Brown KM, Long S, Sibley LD. 2018. Conditional Knockdown of Proteins Using

Auxin-inducible Degron (AID) Fusions in *Toxoplasma gondii.* Bio Protoc 8.

94. Bolte S, Cordelieres FP. 2006. A guided tour into subcellular colocalization analysis

in light microscopy. J Microsc 224:213-32.

95. Mayoral J, Di Cristina M, Carruthers VB, Weiss LM. 2020. *Toxoplasma gondii*:

Bradyzoite Differentiation In Vitro and In Vivo. Methods Mol Biol 2071:269-282.

96. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T,

Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein

V, Eliceiri K, Tomancak P, Cardona A. 2012. Fiji: an open-source platform for

biological-image analysis. Nat Methods 9:676-82.

97. Sinai AP, Webster P, Joiner KA. 1997. Association of host cell endoplasmic

reticulum and mitochondria with the *Toxoplasma gondii* parasitophorous vacuole

membrane: a high affinity interaction. J Cell Sci 110 (Pt 17):2117-28.

98. Andres DA, Young LEA, Veeranki S, Hawkinson TR, Levitan BM, He D, Wang C,

Satin J, Sun RC. 2020. Improved workflow for mass spectrometry-based

metabolomics analysis of the heart. J Biol Chem 295:2676-2686.

99. Markus BM, Bell GW, Lorenzi HA, Lourido S. 2019. Optimizing systems for Cas9

expression in *Toxoplasma gondii*. MSphere 4:e00386-19.

1583

Murphy et al. 2024 Figure 1

Murphy et al 2024 Figure 2

A.

WT

PCR 1

5'UTR | | | | | 3'UTR

PAM

pyrimethamine
resistance

DHFR*

ΔTgLaf

PCR 2

5'UTR | | DHFR* | 3'UTR

PCR 4

B.

Diagnostic PCRs

| | PCR 1 | | PCR 2 | | PCR 3 | | PCR 4 | |
|bp| WT | KO | WT | KO | WT | KO | WT | KO |

C.

| | PAS | | IV58B6 | | TEM |
| | PAS DAPI | Merge | IV58B6 GAP45 DAPI | Phase | |

WT

ΔTgLaf

5-μm

2-μm

Murphy et al. 2024 Figure 3

Murphy et al 2024 Figure 4

Murphy et al. 2024 Figure 5

Murphy et al. 2024 Figure 6

Murphy et al. 2024 Figure 7

Murphy et al. 2024 Figure 8

WT **ΔTgLaf** **COMP**

Murphy et al. 2024 Figure 9