**TITLE:** TgLaforin, a glucan phosphatase, reveals the dynamic role of storage polysaccharides

in *Toxoplasma gondii* tachyzoites and bradyzoites

**AUTHORS:** Robert D. Murphy[1,2], Cortni A. Troublefield[1a#], Joy S. Miracle[1#], Lyndsay E.A.

Young[2], Corey O. Brizzee[2b], Animesh Dhara[1], Ramon C. Sun[2cd], Craig W. Vander Kooi[2cd],

Matthew S. Gentry[2cd], Anthony P. Sinai[2*]

1. Department of Microbiology, Immunology, and Molecular Genetics, College of

Medicine, University of Kentucky

2. Department of Molecular and Cellular Biochemistry, College of Medicine, University of

Kentucky

Current affiliations:

a. Good Samaritan Hospital and Albert B. Chandler Medical Center Emergency

Departments, University of Kentucky

b. Demeetra AgBio, 2277 Thunderstick Dr., Lexington, KY, USA

c. Department of Biochemistry and Molecular Biology, College of Medicine, University of

Florida, Gainesville, FL, USA

d. Center for Advanced Spatial Biomolecule Research, University of Florida, Gainesville,

FL, USA

#These authors contributed equally to this work.

*For correspondence: Anthony P. Sinai, sinai@uky.edu

***Running title:*** TgLaforin regulation of amylopectin granules in Toxoplasma

***Keywords:*** *Toxoplasma gondii* , amylopectin granule, glucan, reversible phosphorylation,

laforin, phosphatase, starch, glycogen, steady-state metabolomics, glutamine

**ABSTRACT**

The asexual stages of *Toxoplasma gondii* are defined by the rapidly growing tachyzoite during

the acute infection and by the slow growing bradyzoite housed within tissue cysts during the

chronic infection. These stages represent unique physiological states, each with distinct

glucans reflecting differing metabolic needs. A defining feature of *T. gondii* bradyzoites is the

presence of insoluble storage glucans known as amylopectin granules (AGs) that are believed

to play a role in reactivation, but their functions during the chronic infection remain largely

unexplored. More recently, the presence of storage glucans has been recognized in

tachyzoites where their precise function and architecture have yet to be fully defined.

Importantly, the *T. gondii* genome encodes activities needed for glucan turnover: a glucan

phosphatase (TgLaforin; TGME49_205290) and a glucan kinase (TgGWD; TGME49_214260)

that catalyze a cycle of reversible glucan phosphorylation required for glucan degradation by

amylases. The expression of these enzymes in tachyzoites supports the existence of a storage

glucan, evidence that is corroborated by specific labeling with the anti-glycogen antibody

IV58B6. Disruption of reversible glucan phosphorylation via a CRISPR/Cas9 knockout (KO) of

TgLaforin revealed no growth defects under nutrient-replete conditions in tachyzoites.

However, the growth of TgLaforin-KO tachyzoites was severely stunted when starved of

glutamine, even under glucose replete conditions. The loss of TgLaforin also resulted in the

attenuation of acute virulence in mice accompanied by a lower cyst burden. Defective cyst

formation due to profound changes in AG morphology was also observed in TgLaforin-KO

parasites, both *in vitro* and *in vivo*. Together, these data demonstrate the importance of glucan

turnover across the *T. gondii* asexual cycle. These findings, alongside our previously identified

class of small molecules that inhibit TgLaforin, implicate reversible glucan phosphorylation as

a legitimate target for the development of new drugs against chronic *T. gondii* infections.

**INTRODUCTION**

*Toxoplasma gondii* is an opportunistic protozoan parasite of all warm-blooded animals that infects one-third of humans worldwide (1). Humans are primarily infected through the consumption of an encysted form of the parasite: either the oocysts shed in cat feces or tissue cysts found in undercooked meat from a chronically infected animal (2). Encysted parasites convert into tachyzoites that rapidly divide and disseminate throughout the body of the host, defining the acute phase of infection (3). Under host immune pressure, tachyzoites convert into slow-growing bradyzoites that populate tissue cysts which are found predominantly in the central nervous system and muscle, defining the chronic phase of infection (4). Tissue cysts are believed to persist for the lifetime of the host and possess the ability to reactivate into tachyzoites in the context of immunosuppression. Reactivation can result in the life-threatening symptoms of toxoplasmosis, with toxoplasmic encephalitis being the primary condition leading to mortality (5). The current lack of insights into bradyzoite physiology *in vivo* precludes the basic understanding needed for the development of drugs that either can clear tissue cysts or prevent reactivation (6).

Until recently, bradyzoites within tissue cysts were considered to be dormant, metabolically inert entities. This view was challenged by our demonstration that encysted bradyzoites replicate (7). Moreover, bradyzoite physiology is both diverse and complex as viewed through the lens of mitochondrial activity, replication status, and, importantly, amylopectin granule (AG) accumulation (8). Although the function of AGs in bradyzoites has not been confirmed, an understanding of the roles of polysaccharides elsewhere suggests that AGs are a source of energy and biosynthetic potential needed for persistence, replication, reactivation, and

transmission (9). These assumptions remain to be tested, and thus much like bradyzoites

themselves, the role of AGs in the *T. gondii* lifecycle is poorly understood.

AGs are large glucans found in the cytoplasm of bradyzoites that have classically served as a

morphological feature distinguishing them from tachyzoites (10, 11). Early studies

characterized AGs as pure amylopectin (12), so AGs are much like plant starch in that they

are water-insoluble storage polysaccharides composed of branched chains of glucose. Unlike

plant starch, however, AGs contain no detectable amylose (unbranched chains of glucose)

(12, 13). More recently, the presence of small, punctate, cytoplasmic glucans in tachyzoites

that are only visible by periodic acid-Schiff (PAS) staining have been recognized (14-16), and

the presence of the glucan is dependent on the *T. gondii* starch synthase (TgSS;

TGME49_222800) (17). Like animal glycogen, this tachyzoite storage polysaccharide is rapidly

turned over (14), as has been observed in other protozoa (18-20), and provides glucose for

glycolysis (17). The observation that large, insoluble glucans do not accumulate within

tachyzoites as they do within bradyzoites suggests that the tachyzoite glucan could be a

distinct and labile form of stored glucose, likely glycogen-like, although its exact chemical and

structural identity remain unknown.

Glucose release from starch in plants requires a cycle of direct, reversible glucan

phosphorylation to solubilize the starch surface, allowing access to degradation enzymes such

as amylases, branching enzymes, and a phosphorylase (21-23). The cycle begins with the

addition of phosphate directly to glucose by the glucan, water-dikinase (GWD) and phospho-

glucan, water dikinase (PWD) that results in the unwinding of glucose chains within starch,

solubilizing the starch surface (24, 25). Glucose-releasing enzymes (amylases) then degrade

starch until the glucan-bound phosphate becomes a steric hindrance, at which point a glucan

phosphatase is needed to remove the phosphate and reset the cycle (26-29). *T. gondii*

encodes all the activities needed for glucan degradation and reversible glucan phosphorylation

including the glucan phosphatase, TgLaforin (TGME49_205290) (30), and glucan dikinase, *T.*

*gondii* GWD (TgGWD; TGME49_214260) (31, 32) .

The central role of reversible glucan phosphorylation in plants is seen in *Arabidopsis thaliana*

where loss of the plant glucan phosphatase, starch-excess 4 (SEX4), results in excess starch

accumulation, aberrant starch morphology, and severely stunted plant growth (33, 34).

Additionally, loss of the glucan phosphatase, laforin, in humans results in hyperphosphorylated

glycogen that aggregates in neurons and astrocytes causing a fatal, neurodegenerative

childhood dementia and epilepsy (35-38). In *T. gondii* , perturbations of several genes related

to glucan metabolism also result in a variety of similar defects including aberrant glucan

accumulation, rewiring of central carbon metabolism, and virulence defects in mice,

highlighting the central metabolic role of glucan metabolism in *T. gondii* (14-17, 39-42).

In this study, we build on our understanding of reversible glucan phosphorylation and its

relevance to parasite metabolism in *T. gondii* . We have recently demonstrated that TgLaforin

is the glucan phosphatase in *T. gondii* , and that TgLaforin represents a unique and viable drug

target (30, 43, 44). Here, we investigate the role of TgLaforin throughout the asexual stages

by knocking out TgLaforin in Type II ME49 parasites. While we expected to observe effects

related to the loss of TgLaforin exclusively in bradyzoites where AGs are typically observed,

the loss of TgLaforin also resulted in striking phenotypic effects in tachyzoites in a context-

specific manner. We thus established a role for TgLaforin, and by extension reversible glucan

phosphorylation, across both tachyzoite and bradyzoite life stages. These findings build upon

previous studies that increasingly demonstrate a central role for glucan metabolism throughout

the parasite's asexual life cycle.

**RESULTS**

***T. gondii  tachyzoites contain a punctate, cytoplasmic glucan***

Previous studies have presented biochemical evidence for rapid glucan turnover in *T. gondii*

Type I RH tachyzoites (14). Moreover, small granules that stain with PAS have also been noted

in the cytoplasm of tachyzoites (14, 16, 17). Under acid-stress conditions, these tachyzoite

glucans have been biochemically characterized as pure amylopectin, and resemble AGs seen

in bradyzoites (13). To further characterize the nature of this tachyzoite glucan, we used

multiple methods to visualize them under unstressed, normal growth conditions (**Figure 1A**).

PAS staining confirmed that Type II ME49 tachyzoites contain small punctate granules

distributed throughout the cytoplasm. To determine if these PAS-stained granules in

unstressed tachyzoites were more glycogen- or starch-like, they were stained with IV58B6.

IV58B6 is an anti-glycogen IgM monoclonal antibody that has previously been demonstrated

to be specific to glycogen (45, 46) by recognizing the highly frequent branch-points found in

glycogen (47). Moreover, IV58B6 does not detect other glucans such as amylopectin or

amylose (the primary constituents of plant starch) (48). Tachyzoites stained with IV58B6 in a

similar pattern to PAS-stained parasites, containing small punctate granules distributed

throughout the cytoplasm (**Figure 1A**), suggesting that the glucan found in tachyzoites is more

glycogen-like than starch-like. Finally, as is well-known, *T. gondii*  tachyzoites contain almost

no visible glucan within their cytoplasm when visualized by transmission electron microscopy

(TEM) (**Figure 1A**), suggesting that the glucan detected by both PAS staining and IV58B6 is

either water-soluble or too small to be visualized, consistent with this glucan being glycogen-

like.

In contrast, bradyzoites have been extensively characterized as containing starch-like AGs

(11-13, 49). After *in vitro* bradyzoite conversion, much of the cytoplasm stained heavily with

PAS (**Figure 1A**). Interestingly, IV58B6 staining intensity appeared to correlate negatively with

*Dolichos biflorus* agglutinin (DBA) staining intensity that defines the cyst wall, implying that

IV58B6 does not stain the PAS-stained glucan in mature bradyzoites, further reinforcing the

observation that structurally distinct polysaccharides exist in tachyzoites and bradyzoites

(**Figure 1A**). Finally, unlike in tachyzoites, AGs were readily identified as electron-lucent

structures throughout the bradyzoite cytoplasm by TEM (**Figure 1A**).

To verify the specificity of PAS and IV58B6 for glucose polymers, tachyzoites and bradyzoites

were treated with acid-$\alpha$-amyloglucosidase (GAA) after parasite fixation and before staining.

GAA cleaves both $\alpha$-1,4- and $\alpha$-1,6-glycosidic bonds and can therefore completely digest

glucans into glucose monomers. Indeed, GAA treatment resulted in the disappearance of

staining with both reagents (**Figure 1B**) demonstrating their specificity for glucose polymers.

***TgLaforin colocalizes with the tachyzoite glucan***

Because *T. gondii* encodes TgLaforin, a glucan phosphatase that is more animal-like than

plant-like (43, 44), we reasoned that TgLaforin could be involved in the metabolism of the

glycogen-like glucan found in tachyzoites. To determine if TgLaforin co-localizes with the

tachyzoite glucan, endogenous TgLaforin was epitope-tagged with hemagglutinin (HA) in *T.*

*gondii* Type II ME49$\Delta$HXGPRT parasites with a CRISPR/Cas9 mediated strategy (**Figure 2A**)

(50). Successful tagging of TgLaforin was confirmed by western blotting (**Figure 2B**).

Immunofluorescence analysis (IFA) of *T. gondii* tachyzoites indicated that TgLaforin is present

in small puncta throughout the cytoplasm, similar to the distribution of the tachyzoite glucan

(**Figure 2C**). Surprisingly, TgLaforin was not detected in *in vitro* bradyzoites by IFA, 6 days

post conversion.

To verify that TgLaforin levels decrease during the tachyzoite to bradyzoite transition, we

converted *T. gondii* tachyzoites to bradyzoites in cell culture using alkaline stress for 6 days

and then probed the converted parasites using western blot analysis. As observed using IFA,

TgLaforin-HA expression decreased dramatically over the course of bradyzoite differentiation

(**Figure 2D**). Transcriptomic data from a previous study obtained from ToxoDB.org indicates

that the transcript levels for TgLaforin do not substantially change over the course of

differentiation, suggesting the possibility that levels of TgLaforin protein are regulated by post-

translational mechanisms (51).

To determine if TgLaforin colocalizes with the glucan present in tachyzoites, we co-stained

TgLaforin-HA tachyzoites with either PAS or IV58B6 along with an anti-HA antibody. In

tachyzoites, TgLaforin colocalized with both PAS (**Figure 2E**) and with IV58B6 (**Figure 2F**),

suggesting its involvement in the metabolism of the tachyzoite glucan.

***Initial characterization of TgLaforin-KO tachyzoites***

To dissect the role of TgLaforin in *T. gondii* glucan metabolism, TgLaforin was knocked out

using CRISPR/Cas9 to disrupt the gene with a pyrimethamine-resistant form of the

dihydrofolate reductase (DHFR-TS*) under a *Neospora caninum* GRA7 (NcGRA7) promoter

(52, 53) (**Figure 3A**). In agreement with a genome-wide CRISPR KO screen (54), TgLaforin is

a non-essential gene under standard cell culture conditions, as multiple TgLaforin-KO clones

were successfully recovered. Integration of the DHFR-TS* construct into the TgLaforin locus

was verified using inside/out PCR at the chimeric locus and by verifying the loss of TgLaforin

transcription (**Figures 3B, C**). The TgLaforin-KO line further used in this study (designated

"ΔTgLaf") was complemented by the introduction of an epitope tagged gene driven by the

TgLaforin promoter. The complementation construct was introduced at an ectopic site in the

genome that lacks known coding sequences or regulatory elements on chromosome VI

(**Figure S1A**), while leaving the ΔTgLaf/DHFR-TS* KO lesion intact for true complementation

(55). This complemented strain, henceforth designated "COMP," was successfully isolated and

confirmed by PCR (**Figure S1B**). Expression levels and localization were similar to those seen

in the TgLaforin-HA line as confirmed by western blotting and IFA (**Figures S1C-D**),

To evaluate effects of a TgLaforin-KO, glucan levels in WT and ΔTgLaf tachyzoites were first

compared using our suite of glucan detection techniques (**Figure 3D**). Surprisingly, the size

and number of PAS-stained granules were not significantly changed in ΔTgLaf tachyzoites

relative to WT parasites. Levels of IV58B6 also remained unaltered after the loss of TgLaforin,

and no aberrant glucan accumulation was observed by TEM as has been previously reported

when genes related to AG or central carbon metabolism were knocked out in *T. gondii*

tachyzoites (**Figure 3D**) (14, 39, 41, 42, 56, 57). Loss of glucan phosphatases in plants and

animals results in aberrant glucan accumulation, and such a phenotype was not observed

here.

***Loss of TgLaforin results in upregulation of glutaminolysis and glutamine dependence***

***in tachyzoites***

Glucan catabolism is significantly affected by the presence of covalently bound phosphate,

and, therefore, loss of glucan phosphatases has profound downstream metabolic impacts in

other systems (58, 59). We thus speculated that loss of TgLaforin would result in the reduced

efficiency of glucan utilization in tachyzoites and also affect downstream central carbon

metabolism. To test this hypothesis, we used gas chromatography/mass spectrometry

(GC/MS) steady-state metabolomic analysis of 3 $\mu$m filter-purified, syringed-passaged

intracellular tachyzoites employing a previously developed sample preparation technique (60).

Previously, it was demonstrated that *T. gondii* tachyzoites primarily utilize glucose and

glutamine to drive central carbon metabolism, synthesize macromolecules, and proceed

normally through the lytic cycle (61). Glucose primarily fuels glycolysis, and glutamine

undergoes glutaminolysis to drive the tricarboxylic acid (TCA) cycle. In the absence of glucose,

*T. gondii* can upregulate both glutaminolysis and gluconeogenesis to make up for the loss of

glucose (61, 62).

While ΔTgLaf metabolite levels remained unaltered relative to WT tachyzoites across much of

the TCA cycle, steady-state levels of metabolites immediately downstream of glutamine were

consistently more abundant in ΔTgLaf parasites compared to their WT counterparts (**Figure**

**4A**), supporting our hypothesis that ΔTgLaf parasites were deficient in glucan/glucose

utilization. An increase in metabolites downstream of glutamine in ΔTgLaf parasites

demonstrates that ΔTgLaf parasites are possibly compensating for deficiencies in glucose

metabolism, supporting a role for the tachyzoite glucan in intermediate *T. gondii* glucose

metabolism.

To determine if loss of TgLaforin resulted in increased dependence on glutamine due to

impaired access to glucose, we performed plaque assays in the presence and absence of

glutamine (**Figure 4B**). In replete media, ΔTgLaf parasites established a similar number of

plaques, indicating no defect in infectivity, that were slightly larger than both the WT and COMP

lines after 10 days of growth (**Figure 4C**). To test the effects of glutamine starvation on ΔTgLaf

parasites, glutamine was removed from plaque assays after parasite invasion to evaluate the

effects of glutamine removal on parasite growth independent of the initial invasion event. In

the absence of glutamine, ΔTgLaf parasites were unable to form visible plaques, whereas both

the WT and COMP parasites formed plaques comparable to those formed under glutamine-

replete conditions (**Figures 4B, C**).

***TgLaforin is required for repeated rounds of progression through the lytic cycle***

To determine which aspects of the *T. gondii* lytic cycle were impaired in the absence of

glutamine, preventing visible plaque formation in ΔTgLaf parasites, the effects of glutamine

starvation on initial parasite replication and egress (stimulated with both A23187 and zaprinast)

were evaluated (63, 64). In both assays, intracellular parasites were pre-starved of glutamine

for at least 72 h before assay initiation. Surprisingly, glutamine starvation had no effect on

stimulated egress or initial parasite replication across the three lines (**Figures S2A-C**). These

data demonstrate that the absence of plaques under glutamine deficient conditions cannot be

pinpointed to a single aspect of the ΔTgLaf lytic cycle, and that the reason for the apparent

absence of plaques manifested later in the infection cycle.

Plaques develop due to repeated cycles of lysis resulting in the clearance of infected cells over

time. The absence of visible clearance prompted us to examine infected host-cell monolayers

for clusters of infected cells using a higher magnification than is typically used in a traditional

plaque assay. Low numbers of parasites were seeded onto glass coverslips and fixed at 3-

and 6-days post-infection, allowing for visualization of developing plaques at a high

magnification. In these experiments, glutamine-depleted host cells were pre-starved of

glutamine prior to infection with parasites to allow for potential for invasion defects. Importantly,

ΔTgLaf parasites demonstrated similar infectivity to WT parasites under both glutamine-replate

and depleted conditions, indicting no gross initial invasion defect.

After 3 days of growth, no statistical differences of nascent plaque sizes were noted between

glutamine-replete and starved conditions in both the WT and COMP lines. However, ΔTgLaf

parasites in glutamine starved conditions were already 1.5x smaller in area than their

counterparts in replete conditions (**Figure 4D**). By day 6 of growth, this difference had widened

to a >3x difference between glutamine replete/depleted ∆TgLaf parasites (**Figure 4E**). Such a

difference was not detected between the two conditions in WT/COMP parasite lines. By

measuring the internal clearing area relative to the total plaque perimeter, it was also noted

that ∆TgLaf parasites were much less capable of forming clearings than the WT/COMP lines

(**Figures S2D, E**), rather they formed clusters of infected cells akin to "turbid plaques" due to

their presumed inability to compete with host cell growth, as the infection progressed. This

observation explains the apparent absence of plaques seen at the lower magnification used in

traditional plaque assays (**Figure 4B**). The modified plaque assay therefore confirmed that the

loss of TgLaforin penalized the summation of repeated rounds of the energy-demanding lytic

cycle rather than one particular aspect of the lytic cycle. Representative images from this assay

can be found in **Figure S2D**.

***Loss of TgLaforin results in aberrant bradyzoite AGs in vitro***

To determine if loss of TgLaforin resulted in bradyzoite conversion defects or aberrant AG

accumulation, parasites were converted to bradyzoites *in vitro* using alkaline stress. During

differentiation, the parasitophorous vacuole membrane (PVM), delimiting the replicative niche

established by tachyzoites, converts into the cyst wall that surrounds bradyzoites within their

host cell (65, 66). The cyst wall is heavily glycosylated and contains N-acetylgalactosamine

(Gal-NAc) that is detectible with DBA (67). Using DBA intensity as a marker for differentiation

progress, no penalty was imposed by the loss of TgLaforin on cyst wall formation over the

course of six days, indicating that ∆TgLaf parasites are not defective in this aspect of

bradyzoite differentiation (**Figure 5A**). Somewhat surprisingly, ∆TgLaf mutant parasites tended

to exhibit stronger labeling with DBA at day 6. We additionally assessed the levels of

accumulated glucans using PAS staining (**Figure 5B**). Semi-quantitative analysis of PAS

intensity within vacuoles during stage conversion showed an expected increase over time, but

no significant difference between the WT and ΔTgLaf parasites was detected over the time

course examined.

Because PAS is not specific to glucans and can stain other glucose-containing molecules such

as glycosylated protein and provides no resolution on glucan morphology, we utilized TEM to

gain higher resolution on AG formation during bradyzoite differentiation. After 6 days of

conversion, WT parasites produced AGs that were circular/ovoid and white (**Figure 5C**). In

contrast, ΔTgLaf parasites contained irregular AGs that were morphologically distinct from AGs

that were observed in WT parasites.  AGs in ΔTgLaf parasites appeared amorphous and

grayer, while appearing to occupy more area of the parasite cytoplasm compared to WT

parasites. To quantify this phenotype, the area of AGs was calculated relative to total parasite

area to determine the percentage of the parasite body occupied by AGs in both WT and ΔTgLaf

strains (**Figure 5D**). Strikingly, AGs occupied approximately 4x more relative area in ΔTgLaf

parasites when compared to WT, indicating that PAS staining may lack the specificity to

capture this difference. When analyzed on an 8-bit gray scale, AGs in ΔTgLaf parasites were

significantly grayer than those found in WT parasites, highlighting potential chemical

differences (such as hyperphosphorylation) resulted in differential interactions of ΔTgLaf AGs

with the TEM contrast reagents, likely the heavy metals used in processing (**Figure 5E**).

Examination and quantification of AGs in the COMP line revealed that complementation of

TgLaforin restored most of the circular/ovoid AGs while they also occupied less space in the

cytoplasm and were overall more like those found in WT parasites (**Figures 5C-E**). Thus, cell

culture experiments demonstrate that the loss of TgLaforin presents itself in both a context and

stage-specific manner.

***Loss of TgLaforin results in attenuated virulence and cyst formation in vivo***

We hypothesized that loss of TgLaforin may impose a steep penalty under the stresses and

potential nutrient scarcities encountered *in vivo* as it does when nutrients (such as glutamine)

are scarce *in vitro*. To test this hypothesis, equal numbers of male and female CBA/J mice

were infected with 100 tachyzoites intraperitoneally (i.p.) and monitored daily using a

previously developed five-stage body index score to track the severity of symptoms associated

with a tachyzoite infection over the course of 28 days (68).

Mice infected with WT parasites began demonstrating symptoms of infection ten days after

infection with tachyzoites (**Figure 6A**). However, mice infected with ΔTgLaf parasites did not

begin to exhibit symptoms until 15 days after infection. Moreover, mice that became

symptomatic from WT parasite infections often proceeded through all stages of symptomology,

and only a minor proportion of mice that became sick were able to recover from infection (>70%

of mice became moribund or died). Infection from ΔTgLaf parasites, however, resulted in the

majority of mice only developing mild symptoms (Stage 2 or less) with many of these mice

recovering (**Figure 6A**). The attenuated capacity of the ΔTgLaf parasites to cause symptoms

in mice was reflected in the mortality rates of the infected mice: infection with WT parasites

resulted in 73% mortality rate after 28 days whereas ΔTgLaf parasites only caused 17%

mortality (**Figure 6B**). Complementation of TgLaforin partially rescued this defect in virulence

as COMP parasites resulted in an earlier onset of symptomatic infection at Day 11, and the

majority (53%) of mice succumbed to infection during the first 28 days (**Figures 6A, B**).

Because the acute phase of infection was significantly attenuated by the loss of TgLaforin, we

hypothesized that cyst numbers would be significantly lowered. To determine the number of

cysts formed after 28 days of acute infection, we used a previously established protocol for

harvesting and counting tissue cysts from infected mouse brains, following purification on

Percoll gradients (7, 69). Consistent with the ability of ΔTgLaf parasites to stage convert in

culture, mutant parasites were able to establish tissue cysts *in vivo*.  However, the number of

cysts recovered from ΔTgLaf infected animals was lower than those obtained from WT infected

animals (**Figure 6C**). Restoration of TgLaforin in the COMP line effectively restored tissue cyst

yields.

*ΔTgLaf tissue cysts can reestablish infections in naïve mice*

To determine if the loss of TgLaforin impacted the overall viability/infectivity of *in vivo* tissue

cysts, we examined the disease progression in WT, ΔTgLaf, and COMP infected animals

following injection of 20 tissue cysts i.p. Consistent with prior data, infection with tissue cysts

results in markedly lower pathology and consequent mortality during the acute phase for WT

as well as both the ΔTgLaf and COMP lines (**Figure 7A, B**) (68). The death rate from cyst

infection did not differ statistically among the three lines. Twenty-eight days post-infection, cyst

burdens were again enumerated for each line. ΔTgLaf parasites were once again much less

competent at forming cysts *in vivo*. However, unlike the tachyzoite infection, the COMP line

was unable to rescue this defect in cyst formation (**Figure 7C**), suggesting that physiological

and metabolic changes associated with the loss of TgLaforin manifest differently based on the

life cycle stage, impacting their capacity to be complemented.

**Proper cyst formation relies on the presence of TgLaforin**

Recent work has also demonstrated that loss of phosphoglucomutase 1, an enzyme related to

AG metabolism, results in smaller cysts (70). Our prior work established that bradyzoite

replication within tissue cysts occurs in a cyclical pattern that is reflected in the packing density

of bradyzoites within the cyst (7).  Notably, expansion of tissue cyst diameter precedes

replication.  We therefore analyzed the packing densities and diameters of cysts from WT,

∆TgLaf, and COMP lines to investigate the relationship between the number of bradyzoites

within a cyst to the overall volume of the cyst.

Packing density is calculated by dividing the number of bradyzoite nuclei within an optical

section of a cyst by the volume of the cylinder representing that optical section (7). ∆TgLaf

parasites had significantly lower packing density than WT or COMP parasites at both 4 and 6

weeks post-infection (**Figure 8A,C** and **Figure S3A,B**). This indicates that loss of TgLaforin

resulted either in lower replication within the cyst, increased parasite death, or both. These

data provide the first evidence of a mutation resulting in lower bradyzoite numbers within tissue

cysts as opposed to merely assessing overall cyst numbers.

To determine the relationship of packing density to cyst diameter, we analyzed the cyst

diameter alone as a surrogate for cyst expansion at both time points (**Figure 8B,D** and **Figure**

**S3C,D)**. At 4 weeks post-infection, ∆TgLaf tissue cysts were smaller in diameter than WT and

COMP cysts. At week 6, however, there was no difference between average diameter of

∆TgLaf tissue cysts and that of the WT and COMP lines. Together, these data indicate that the

decreased packing density seen in ∆TgLaf parasites is a result of continued cyst expansion

accompanied by reduced bradyzoite replication, increased bradyzoite death, or a combination

of these factors.

To establish whether the changes in AG levels and morphology observed in ∆TgLaf

bradyzoites following *in vitro* conversion were observed *in vivo* by TEM, we adapted a protocol

designed to capture and image low abundance cells by TEM by making it compatible with our

tissue cyst purification protocol (see Methods) (71, 72). TEM imaging revealed that while WT

parasites formed largely normal/canonical AGs *in vivo* as seen *in vitro* (**Figure 9** [compare with

**Figure 5A**])*,* ∆TgLaf parasites contained almost exclusively aberrant AGs that mirrored the

same morphological defects seen *in vitro* (**Figure 9** and **Figure S4**). ΔTgLaf AGs were

irregularly sharpened with a flat, multi-lobed appearance. Importantly, COMP parasites neither

over-accumulated nor formed aberrant AGs, demonstrating that this defect is specific to loss

of TgLaforin (**Figure 9**).

In addition to containing aberrant AGs, the internal morphology of ΔTgLaf parasites appeared

to be altered by the presence of the AGs, as significant organelle displacement was also noted.

Many of the ΔTgLaf parasites also appeared hollow or "ghost-like" in appearance resulting

from their unstained/absent cytoplasm, which included the apparent loss of the nucleus in

several parasite forms. These data suggest that a significant number of the ΔTgLaf bradyzoites

were inviable within the cyst (additional images of parasites from both WT and ΔTgLaf

parasites are presented in **Figure S4**) and that increased death is a significant contributing

factor to the reduced packing density (**Figure 8**). Together, these direct and selective impacts

of TgLaforin's loss on both bradyzoite viability and growth *in vivo* establish TgLaforin as a

potentially druggable target.

**DISCUSSION**

The asexual life cycle of *Toxoplasma gondii* is defined by two fundamentally distinct forms: the

rapidly replicating tachyzoite and the slowly growing bradyzoite. These forms represent distinct

physiological states that can be further subdivided, particularly within encysted bradyzoites

(73). Insights into these physiological states, inferred from transcriptomic analyses, are

reinforced in emerging metabolomic studies (74, 75). In these studies, glucose and glutamine,

which are both linked to energetics, biosynthesis, and intermediary metabolism, appear as

critical metabolites. Importantly, glucose can be stored in polymers like glycogen and

amylopectin during times of low energy need. Stored glucose can be present in an accessible

and labile form for rapid mobilization such as glycogen, or in a less soluble forms as AGs from

which it can be accessed more slowly (76). Reversible glucan phosphorylation facilitates the

breakdown of such glucan polymers by disrupting the crystalline helices on the glucan surface

(77). *T. gondii* encodes the capacity for reversible glucan phosphorylation (30, 32, 43, 44).

The contribution of this process to tachyzoite and bradyzoite biology was evaluated through

targeted manipulation of the glucan phosphatase, TgLaforin.

The accumulation of AGs within bradyzoites and their apparent absence in tachyzoites has

been used as discriminator between these life cycle stages. Detailed examination, however,

presents a considerably more nuanced picture, alongside emerging evidence that points to

rapid glucan turnover within tachyzoites (14).  Here, using an IgM monoclonal antibody

(IV58B6) that specifically recognizes glycogen-like glucose polymers (45), we demonstrate

that the stored glucan within tachyzoites is structurally closer to animal glycogen than the

insoluble plant-like AG granules found in bradyzoites (12). This duality between life stages may

be additionally reflected in the observation that TgLaforin, the glucan phosphatase, and

TgGWD (TGME49_214260), the accompanying kinase that is predicted to phosphorylate *T.*

*gondii* glucan polymers, trace their structural lineages to animals and plants, respectively (43,

44). The glycogen-like glucan polymer appears to be specific to tachyzoites as its levels

decrease upon *in vitro* differentiation while the overall PAS intensity increases (**Figure 1A**).

This suggests that the tachyzoite glucan and bradyzoite AG are architecturally distinct

polymers with respect to both branching frequency, solubility, and phosphorylation status

(**Figure 2D**). The glucose-based nature of both particles is supported by the elimination of both

IV58B6 and PAS staining with $\alpha$-amyloglucosidase treatment (**Figure 1B**).

To address the contribution of stored glucans in both tachyzoites and bradyzoites, we disrupted

the glucan phosphatase TgLaforin. This enzyme preferentially removes phosphate groups

from the C3 carbon on glucose facilitating access to enzymes that release glucose (30).  The

loss of other glucan phosphatases such as SEX4 in *A. thaliana* and laforin in mammals is

accompanied by excessive accumulation of aberrant starch and hyperphosphorylated

glycogen in plants and animals (33, 34, 37, 38, 78). Surprisingly, given these penalties in other

systems, ΔTgLaf tachyzoites exhibited no gross morphological changes in glucan content,

consistent with a recent study in which TgGWD was knocked out (32). These observations

contrasts with other KO studies of glucan pathway associated proteins such as CDPK2,

glycogen phosphorylase, the PP2A holoenzyme, and α-amylase which all reported

exaggerated glucan accumulation in tachyzoites (14, 39, 41, 42, 56).

Despite the absence of glucan accumulation in ΔTgLaf tachyzoites, the loss of this gene exerts

an effect on tachyzoite glucan metabolism. Consistent with the metabolic defects associated

with the loss of laforin in humans (58), loss of TgLaforin in *T. gondii* resulted in altered central

carbon metabolism that manifested as ΔTgLaf parasites' dependence on glutamine. ΔTgLaf

tachyzoites' dependence on glutamine supports recent studies demonstrating that tachyzoites

utilize storage glucans for glucose allocation (14, 17) because the presumed loss of efficient

glucan degradation results in tachyzoite dependence on glutamine (**Figures 4B, C**).  As *T.*

*gondii* tachyzoites primarily utilize glucose and glutamine to support their rapid growth, this

dependence on glutamine reinforces previous observations that glutamine can substitute for

glucose in this altered metabolic landscape (61, 79).

These results are consistent with many previous studies that disrupt glucose and/or glucan

metabolism, but contrast with others. Disruption of TgGT1 (TGME49_214320), the only

plasma-membrane glucose transporter in *T. gondii* (61, 80), or TgHK (TGME49_265450), the

*T. gondii* hexokinase, resulted in upregulation of gluconeogenesis, and parasite growth was

highly attenuated with glutamine depletion (79). Most strikingly, parasites lacking starch

synthase (TgSS; TGME49_222800) displayed no dependence on glutamine and, in fact, grew

faster than WT parasites when both glucose and glutamine were removed from the culture

media (17). Interestingly, however loss of TgSS did result in lower glucose flux through

glycolysis, consistent with our findings that demonstrate a role for the tachyzoite glucan in

glucose allocation. This finding may suggest that while the absence of AG in the ΔTgSS

parasites may not be detrimental, overaccumulation as observed in ΔTgLaf cysts can

contribute to toxicity (**Figure 9**). Perturbations of other glycolytic enzymes also demonstrated

varied effects related to the presence of glutamine: loss of the glycolytic enzyme

glyceraldehyde-3-phospahte dehydrogenase (GAPDH1) could be rescued with high levels of

glutamine (81), but glutamine could not rescue pyruvate kinase (TgPYK1) knockdown

parasites (57). Our data indicate that loss of access to key nutrients such as glucose and

glutamine has a profound impact on the repeated rounds of the lytic cycle without being

attributable to one specific process within the cycle, suggesting that the penetrance of the

phenotypic defect manifests cumulatively over time, rather than being hard wired in each

infection cycle (**Figures 4D, E**).

Despite glucan metabolism being historically viewed as being important in the chronic infection,

TgLaforin protein expression decreased during the tachyzoite to bradyzoite conversion *in vitro*

even though its transcript levels do not change (**Figures 2C, D**). This could be a transient

observation as the downregulation of glucan catabolism during conversion would facilitate

accumulation of AGs for the chronic infection. We therefore examined how the loss of

TgLaforin affected the capacity of ΔTgLaf parasites to differentiate *in vitro*. The ΔTgLaf

parasites exhibited no defect in AG-accumulation kinetics, detected by PAS staining, or in cyst

wall formation, detected with DBA lectin (**Figure 5A, B**). The lack of difference in PAS labeling

between both the WT and the ΔTgLaf lines, however, did not reveal the differences noted by

TEM. As initially hypothesized would be the case in both tachyzoites and bradyzoites, loss of

TgLaforin resulted in aberrant AG accumulation within *in vitro* bradyzoites that is marked by

changes in both level and morphology (**Figures 5C-E**), as seen in plants and vertebrates (33,

34, 36-38). AGs in the ΔTgLaf parasites were not only present at higher levels but were

potentially chemically distinct considering their differential binding to TEM contrast metals.

Given that TgLaforin is a confirmed glucan phosphatase (30), we speculate that AG

hyperphosphorylation may account for both altered morphology and appearance by TEM.

These context-specific phenotypes suggested that the ΔTgLaf mutant would manifest

phenotypic differences in both the acute and chronic phases on infection *in vivo*. Indeed, the

loss of TgLaforin was associated with a markedly reduced symptomology and associated

mortality compared to both the parental and complemented parasites during acute tachyzoite-

initiated infection (**Figures 6A, B**). Not only was there a delay in symptomatic disease, but also

a reduction in disease severity and overall cyst burden. Symptomology in the acute infection

is driven by an increasing parasite burden driving an overexuberant host inflammatory

response. The delayed symptom onset suggests growth inhibition by the stringent *in vivo*

environment that more effectively controls ΔTgLaf parasite infection with less robust

inflammation.  Notably, the delayed and milder course of the tachyzoite infection resulted in a

lower overall cyst burden in surviving animals compared to infection with both WT and COMP

parasites (**Figure 6C**).

Bradyzoite replication within cysts is preceded by the expansion of the cyst to accommodate

newly formed parasites (7).  Bradyzoite occupancy can be established by quantifying the

number of nuclear cross sections in imaged tissue cysts to define the packing density. The

finding that ΔTgLaf cysts were smaller, and less densely packed than WT or complemented

cysts, indicates that TgLaforin, and by extension reversible glucan phosphorylation, may have

a significant impact on the progression of the chronic phase of infection, manifesting at the

level of encysted bradyzoites.

The capacity of ΔTgLaf cysts to initiate new infections was assessed *in vivo*. Consistent with

other studies (7, 68), infection with tissue cysts is associated with limited symptomology and

low overall mortality. Similar to what was observed with tachyzoite-initiated infections, the

ΔTgLaf parasites consistently presented with lower cyst burdens relative to WT cyst infections.

One puzzling outcome, however, is the fact that the complemented line failed to rescue the

overall cyst burden for tissue cyst-initiated infections, despite completely rescuing infections

initiated with tachyzoites. This recurring theme for the penetrance or lack thereof in a context

specific manner appears to apply selectively to the rescue of the phenotype as well.

As was observed *in vitro*, TEM revealed a phenotype of aberrant AG accumulation which was

accompanied by apparent extensive parasite death **(Figure 9)**. Together, these phenotypes

manifest as AGs occupying much of the cytoplasm, the enucleation of parasites, and the

detection of "ghost parasites" that appear to be empty shells surrounded by the cytoskeleton.

The presence of "ghost parasites" adjacent to seemingly normal parasites highlights the

variable penetrance of the phenotype in what is a clonal population, given that each tissue cyst

originates from a single parasite. These finding suggest that AG metabolism is under tight

control as dysregulated accumulation can result in cumulative defects resulting in toxicity and

death. The high frequency of these abnormal parasites suggests that reversible glucan

phosphorylation and TgLaforin specifically represent legitimate bradyzoite specific drug

targets. We recently described a small molecule that inhibits recombinant TgLaforin (30) which

serves as a potential starting point in the development of a new class of anti-*Toxoplasma*

therapeutic agents. Particularly exciting in this context is the fact that a class of drugs exhibiting

efficacy with tissue cyst clearance (atovaquone (82), endochin-like quinolones (83), and

JAG21 (84)) all target mitochondrial respiration. When glucose is limiting, mitochondrial

respiration can be driven by glutamine. This provides an opportunity for combination therapy

to promote the clearance of toxoplasma tissue cysts as a means of mitigating the risk of

reactivation.

**METHODS**

***Fibroblast and parasite culture and maintenance***

All parasite lines were maintained in human foreskin fibroblasts (HFFs; ATCC) in Minimal

Essential Media-$\alpha$ (MEM-$\alpha$; Gibco) supplemented with 7% heat-inactivated fetal bovine serum

(FBS; Gemini Bio), 100 U/mL penicillin, 100 $\mu$g/mL streptomycin, and an additional 2 mM L-

glutamine (Gibco; 4 mM total L-glutamine). Cells and parasites were incubated at 37$^{\circ}$C and

5% $CO_2$ in a humidified incubator. Genetically modified parasites were maintained in MEM-$\alpha$

containing 7% dialyzed FBS (Gemini Bio) and either pyrimethamine (1 $\mu$M), mycophenolic

acid/xanthine (MPA: 25 $\mu$g/mL, xanthine: 50 $\mu$g/mL), or 6-thioxanthine (6-Tx: 80 $\mu$g/mL).

Assays analyzing the effects of glutamine deprivation used Dulbecco's Modified Eagle Medium

(DMEM). Both glutamine-replete (Gibco, 11966025) and depleted (Gibco, 11054020) DMEM

were supplemented with 7% dialyzed FBS. Glutamine-replete media from the supplier lacked

other key nutrients and was modified to contain 5 mM glucose, 1 mM sodium pyruvate, and 4

mM L-glutamine.

***Generation of T. gondii mutant lines***

*Type II ME49ΔHXGPRT* ("WT"—the parental line utilized to generate all other lines in this

study): This line was generated in a previous study using CRISPR/Cas9 targeting of

TgHXGPRT and selection with 6-Tx (68).

*TgLaforin-3xHA-HXGPRT*: TgLaforin was epitope tagged with HA at the C-terminus using

CRISPR-Cas9 to disrupt the TgLaforin 3'UTR immediately downstream of the endogenous

stop codon as has been previously described (50). Briefly, a sgRNA immediately downstream

of the TgLaforin stop codon was designed using the EuPaGDT design tool

(http://grna.ctegd.uga.edu). The top hit was selected (**Table S1**) and used to replace the

sgRNA sequence in pSAG1::CAS9-U6::sgUPRT, a plasmid containing both Cas9-green

fluorescent protein (GFP) and an interchangeable sgRNA scaffold; (**Table S2**) (53).

Replacement of the interchangeable sgRNA was accomplished using a Q5 site-directed

mutagenesis kit (**Table S3**) (New England BioLabs). The TgLaforin-HA tagging construct was

generated by amplifying the 3' end of the TgLaforin-HA construct generated for

complementation (see generation of COMP line below and **Tables S2 and S3**) along with the

connected HXGPRT selectable marker. Both the TgLaforin-HA PCR-amplicon and the

CRISPR-Cas9-GFP were transfected into $1.4 \times 10^7$ *T. gondii* ME49ΔHXGPRT parasites (2:1

insert:plasmid molar ratio; 30 μg DNA total) by electroporation with a time constant between

0.16 and 0.20 msec (BioRad Gene Pulser II). After 24 h, surviving parasites were syringe-

passaged from infected HFFs with a 27 G needle to lyse host cells, and gravity-filtered through

a 10 μm filter to remove host-cell debris. Successful transformants were then enriched by use

of fluorescence-activated cell sorting (FACS; Sony SY3200, installed in a biosafety level II

cabinet) to select parasites expressing Cas9-GFP from the transfected plasmid by isolating

GFP+ parasites. HFFs were infected with GFP+ parasites, and then placed in media containing

MPA/xanthine 24 h later to select for restoration of HXGPRT. MPA/xanthine-resistant parasites

were cloned by limiting dilution into a 96 well plate. Wells containing single plaques were picked

7 days later and expanded. Genomic DNA was extracted from clones using a Proteinase K

treatment detailed elsewhere (85). Successful tagging of TgLaforin was verified using

sequencing, immunoblotting, and IFA.

*ME49ΔHXΔTgLaforin* ("ΔTgLaf"): TgLaforin was disrupted using a CRISPR-Cas9 mediated

strategy as detailed above, with several differences. Briefly, a single sgRNA was designed to

target the first exon of TgLaforin with the top hit from EuPaGDT (**Table S1**). To disrupt

TgLaforin with a selectable drug marker, DHFR-TS*, a pyrimethamine-resistant mutant of the

DHFR gene, containing a 5'-NcGra7 promotor and DHFR 3'UTR was amplified from pJET-

NcGra7_DHFR (**Table S2**). Amplification utilized primers containing 40 nt extensions

homologous to the 5'- and 3'-UTR of TgLaforin to encourage homologous recombination-

mediated whole-gene replacement with the drug cassette (**Table S3**). Both the PCR-amplified

DHFR* homology cassette and the CRISPR-Cas-GFP plasmid were transfected and FACS-

sorted as described above. GFP+ parasites underwent drug selection in pyrimethamine.

Parasites were then cloned and expanded as detailed above. Successful integration of the

DHFR* cassette into the TgLaforin locus was verified using PCR with inside/out primer pairs

to the chimeric, interrupted gene (**Table S3**). Loss of TgLaforin transcription was verified by

purifying RNA from TgLaforin clones on RNeasy spin columns (Qiagen). Using the Promega

Reverse Transcriptase System, cDNA was synthesized from RNA extracts. Primers designed

for full-length TgLaforin amplification were then used to verify loss of TgLaforin cDNA in

knockout lines.

*ME49ΔHXΔTgLaforin+ChrVI-TgLaforin* ("COMP"): Complementation of TgLaforin was also

executed using a CRISPR-mediated strategy. A sgRNA to a neutral locus on chromosome VI

identified previously (55) was generated using the same mutagenesis strategy as above (**Table**

**S1 and S3**). A full length TgLaforin cDNA containing its endogenous 5'UTR (2000 bp upstream

from gDNA) was synthesized by GenScript and inserted into a pHA3x-LIC vector (**Table S2**)

containing a C-terminal HA tag and a DHFR 3'UTR, linked to the HXGPRT selectable marker

(named "TgLaforin-HA3x-LIC"; also used above for endogenous tagging to create the

TgLaforin-HA line). The entire construct (5'UTR:TgLaforin-cDNA:DHFR-3'UTR:HXGPRT) was

amplified from the vector and co-transfected into ΔTgLaf parasites with the CRISPR-Cas9

plasmid as done above. Successful transformants that received the HXGPRT marker were

selected with MPA/xanthine. Successful insertion of TgLaforin along with its promoter was

verified using PCR (**Table S3**), immunoblotting, and IFA with an anti-HA antibody (Abcam).

***Immunofluorescence (IF) staining***

HFFs were grown on glass coverslips until confluent and subsequently infected. Infected HFFs

were fixed with either methanol (MeOH) (100%, -20°C) or methanol-free paraformaldehyde

(PFA) (4% in phosphate-buffered saline (PBS); Electron Microscopy Sciences) as indicated

below for each antibody. Infected HFFs fixed with PFA were permeabilized in 0.1% TritonX-

100 in PBS++ (PBS containing 0.5 mM $CaCl_2$ and 0.5 mM $MgCl_2$) for 10 min at room

temperature (RT). Primary and secondary antibodies were diluted in 3% (w/v) bovine serum

albumin (BSA; Fisher) in PBS++. Samples were first incubated with the primary antibody (αHA-

1:1,000; αSAG-1:10,000; αGAP45-1:5,000; IV58B6-1:50) at RT for 45 min, washed 3x with

PBS++, and then incubated with fluorescent secondary antibodies (1:2,000) and 4',6-

diamidino-2-phenylindole (DAPI; 300 nM) for 45 min. Secondary antibodies (Invitrogen) were

conjugated to either Oregon Green or Texas Red fluorophores and specific to the species and

class of primary antibody used. Samples were then washed 3x with PBS++ before mounting

the coverslip on a glass slide using MOWIOL mounting media.

IF staining was visualized using a Zeiss AxioVision upright microscope with a 100X 1.4

numerical-aperture oil immersion objective, and images were acquired using a grayscale Zeiss

AxioCam MRM digital camera. Grayscale images were pseudo-colored in ImageJ using

magenta (Texas Red), yellow (Oregon Green), and cyan (DAPI), and further alterations to

brightness and contrast were also made in ImageJ when deemed appropriate. For all assays

in which staining intensity was compared across treatments and parasite lines, concentrations

of antibodies, exposure times, and alterations to brightness/contrast were identical.

Colocalization of fluorescent antibodies/reagents was quantified using Pearson's coefficient

calculated with the JACoP plugin on ImageJ (86).

***PAS staining***

PAS staining was done on infected HFFs fixed in 4% PFA and permeabilized as above.

Coverslips were then washed 3x in tap water before the addition of 1% periodic acid (Sigma-

Aldrich) for 5 min. Coverslips were then washed with three changes of tap water. Schiff's

reagent (diluted 1:4 in tap water) was added for 15 min. Coverslips were subsequently washed

10x with tap water to develop stain before being incubated with DAPI for 10 min and then

mounted as above. PAS-stained samples were visualized using fluorescence microscopy

(excitation: 545 nm, emission: 605 nm). When PAS was co-stained with antibodies, primary

antibodies were incubated with PAS-stained slides overnight in BSA at 4 °C before standard

secondary staining.

Samples treated with acid-$\alpha$-amyloglucosidase (GAA) (from *Aspergillus niger*, >260 U/mL,

Sigma) were incubated with GAA after permeabilization. GAA was diluted 1:50 in 50 mM

sodium phthalate buffer, pH 5.5, and samples were treated for 24 h at room temperature.

Untreated controls were incubated in phthalate buffer without GAA. Samples were then stained

with PAS or IV58B6 as described in the IF-staining workflow above.

***In vitro bradyzoite conversion assay***

Tachyzoites were converted to bradyzoites *in vitro* using alkaline stress as has been done

previously with several modifications (87). HFFs grown were infected with tachyzoites in

standard cell culture media. 4 h later, media was replaced with RPMI 1640 (Gibco 31800022)

supplemented with 50 mM HEPES and adjusted to pH 8.2 with NaOH. Parasites were then

cultured for 2-6 days at 37$^{o}$C, ambient $CO_2$, and sealed in Parafilm. Media was replaced every

other day to maintain the basic pH. Parasites were fixed in PFA and stained with fluorescein

conjugated *Dolichos biflorus* agglutinin (DBA; 1:1000, Vector Laboratories) and PAS. Images

were obtained in grayscale on a Zeiss AxioVision upright microscope as described above.

To determine the degree of labeling with DBA or PAS, the Fiji/ImageJ (88) was used to create

a binary mask outlining cysts that was applied to the PAS-stained image to measure the

greyscale intensity of each ROI (i.e. each individual vacuole/*in vitro* cyst).

***Transmission electron microscopy of in vitro tachyzoites and bradyzoites***

Transmission-electron microscopy (TEM) was performed as done previously (89). Blocks were

stained at the University of Kentucky's Imaging Center in the College of Arts and Sciences.

Blocks were trimmed and sectioned on an ultramicrotome with a diamond knife. Sections were

placed on copper grids and then contrast stained with lead citrate. Micrographs were collected

at the University of Kentucky's Electron Microscopy Center on a Talos F200X TEM (Thermo)

operated at 200 kV accelerating voltage with a 50 $\mu$m objective aperture inserted to enhance

contrast using a 16M pixel 4k x 4k CMOS camera (Ceta, Thermo Scientific). AG size and

grayscale values were measured in ImageJ.

***Immunoblotting***

Parasites were syringe lysed from host cells, pelleted, and $2x10^6$ parasites were resuspended

in SDS-PAGE sample buffer and boiled for 10 min before being run on a single lane of a 10%

polyacrylamide gel. The gel was then transferred to a 0.2 $\mu$m PVDF membrane (BioRad) using

a Turbotransfer System (BioRad) for 7 min at 25 V. The PVDF membrane was blocked in 5%

(w/v) non-fat milk in Tris-buffered saline plus Tween-20 detergent (TBST; 0.1% Tween-20) for

20 min before being probed with a primary antibody ($\alpha$HA-1:1,000; $\alpha$GAP45-1:5,000; $\alpha$SRS9-

1:1,000; $\alpha$SAG1-1:10,000) in non-fat milk overnight at 4$^o$C (Cell Signaling C29F4). The blot

was washed 3x with TBST before probing with either HRP-conjugated $\alpha$-rabbit or $\alpha$-mouse-

IgG (Jackson Laboratories). Blot was washed and developed for 5 min using SuperSignal$^{TM}$

West Pico PLUS (Thermo Scientific) and visualized on a GelDoc station (BioRad).

***Steady state polar metabolite analysis***

Parasites were prepared as previously described (60). Confluent HFFs were infected with

parasites at a multiplicity of infection (MOI) of 2 to achieve a high density of parasites after 48

h of growth (>80% cells containing >32 parasites each).  Plates containing infected HFFs were

placed on ice, media removed, and the monolayer was washed 2X with ice-cold PBS.

Parasites were harvested on ice in a 4 $^o$C cold-room. Cells were scraped from plate surface,

resuspended in PBS (8 plates/50 mL PBS), and centrifuged at 1000$g$ for 10 min at 4 $^o$C. PBS

was removed, the cell pellet was resuspended in 2 mL PBS, and syringe passaged

successively in 23 G and 27 G needles. The soluble host cell lysate was removed by

centrifugation (1000*g*). The pellet was resuspended in 5 mL PBS and host-cell debris was

removed by syringe-filtering the suspension through a 3 $\mu$m filter (Whatman). Filtered parasites

were then pelleted, resuspended in 1 mL PBS, and counted on a hemacytometer. Parasites

were pelleted a final time at 14,000*g* for 30 s at 4 °C, supernatant was removed, and pelleted

parasites were flash frozen in liquid nitrogen and stored at -80 °C until metabolite extraction.

*Polar metabolite extraction:* Polar metabolites were extracted in 0.5 mL -20 °C 50% methanol

(MeOH) containing 20 $\mu$M L-norvaline (procedural control) for 30 min on ice. During the 30 min

incubation, samples were regularly vortexed. Samples were then centrifuged at 14000*g* for 10

min to pellet insoluble material (protein, DNA, RNA, and glycans). Supernatant containing polar

metabolites and pellet were dried separately on a SpeedVac (Thermo) at $10^{-3}$ mBar until

methanol (MeOH) was completely sublimated and only dried pellet remained.

*Pellet hydrolysis and extraction:* Dried fraction containing protein was hydrolyzed by

resuspending the pellet in 2 N HCl (final concentration) at 95°C for 2 h. Hydrolysis was

quenched, and hydrolyzed amino acids were extracted by the addition of an equal volume of

100% MeOH with 40 $\mu$M L-norvaline such that the final concentration was 50% and 20 $\mu$M,

respectively. Extraction and drying then proceeded as described above.

*Sample derivatization*: Dried samples (both polar metabolites and hydrolyzed protein) were

derivatized in 70 $\mu$L 20 mg/mL methoxyamine hydrochloride in pyridine for 90 min at 30 °C.

Samples were then centrifuged at 14000*g* for 10 min to remove any particulate, and 50 $\mu$L of

the methoxyamine supernatant was mixed with 80 $\mu$L *N*-methyl-*N*-trimethylsilyl

trifluoroacetamide (MSTFA) and incubated for 30 min at 37°C. Samples were then transferred

to amber glass chromatography vials and analyzed by GC/MS.

*GC/MS analysis*: Metabolites were analyzed on an Agilent 7800B GC coupled to a 5977B MS

detector using a previously established protocol (90). Automated Mass Spectral Deconvolution

and Identification System (AMDIS) was used to analyze metabolites by matching metabolites

to the FiehnLib metabolomics library via retention time and fragmentation pattern.

Quantification of metabolite levels was performed in Mnova. Sample abundance was

normalized to L-norvaline (procedural control) and protein from the protein pellet (experimental

control). Steady state metabolites are presented as the mean of three independent replicates.

**Conventional plaque assays**

HFFs were grown in 12-well plates until confluent. HFFs were subsequently infected with 200

parasites/well under standard cell culture conditions. Wells were washed with PBS to remove

residual invasion media, and media was changed to glutamine replete or depleted media 4 h

post-infection to allow for invasion. Plates remained undisturbed for 10 days before the infected

HFFs were fixed with 100%, -20°C MeOH for 20 min, stained with 1% crystal violet solution for

20 min, and then the plaques were de-stained with repeated tap water washes. Zones of lysis

(white clearings) could be visualized against intact cells (purple). Images of plaques were

obtained by scanning plates on an Epson Perfection V600 photo scanner at a resolution of 600

dpi. The plaques were measured by pixel area using ImageJ. Plaque assays were conducted

on three independent replicates, and plaque size from these experiments were aggregated to

highlight variability in plaque sizes.

**Egress assays**

HFFs were grown to confluency in 35 mm glass bottom dishes (MatTek, P35G-0-14-C). Two

days before infecting HFFs on glass bottom dishes, both HFFs and parasites were

independently pre-treated in either glutamine-replete or -depleted media (see above for media

formulations). After 48 h pre-treatment, $10^5$ parasites of each line (WT, $\Delta$TgLaf, and COMP) in

each pre-treatment (gln+/-) were added to fresh dishes and allowed to grow for 48 h so that

most vacuoles contained >32 parasites each. Several hours before egress, media in each

infected plate was adjusted to 1.5 mL and allowed to equilibrate at 37°C in 5% $CO_2$. The

calcium ionophore A23187 (Cayman Chemical Company) was prepared as a 2 mM stock in

DMSO and diluted in (+/-) gln media to make a 4X concentration of 12 $\mu$M and maintained at

37°C throughout the assay. Zaprinast was likewise prepared as a 100 mM stock in DMSO and

diluted into media at a 4X concentration of 2 mM. Egress was triggered by the addition of 0.5

mL 4X A23187 to infected HFFs (3.0 $\mu$M final concentration) or 0.5 mL 4X-zaprinast (500 $\mu$M

final concentration). Egress was monitored on a Nikon Eclipse Ti2 inverted microscope with a

40X phase air objective modified with a 1.5X optivar. Several fields containing vacuoles were

selected from each plate, and an image was obtained 10 s after triggering egress from each

field once every 5 s for 5 mins (61 images/field) on a Nikon DS-Ri2 color camera. Videos of

each field were assembled on NIS Elements software. Egress was monitored using standard

deviation of pixel intensity and determined by inflection point of change in standard deviation

of pixel intensity. Inflection point was calculated by fitting a gaussian curve to the first derivative

of the standard deviation in pixel intensity and calculating the mean of the curve. Technical

replicates (fields on each plate) were averaged for each biological replicate (average of fields

from each plate).

***Replication assays***

HFFs were grown on glass coverslips in a 24-well plate until confluent. Two days before

infecting HFFs on coverslips, both HFFs and parasites were independently pre-treated in either

glutamine-replete or -depleted media. After pre-treatment, $10^4$ parasites of each line (WT,

ΔTgLaf, and COMP) in each condition (gln+/-) were added to 3 coverslips each. 24 hours later,

infected HFFs were fixed in MeOH and stained with Rb-α-SAG1 (1:10,000) and DAPI for ease

of visualization. Counting of parasites/vacuole was performed for each line/condition on coded

blinded slides with the identity of samples revealed upon completion of the counting. A total of

three independent replicates were performed.

***Modified plaque assays***

A modified plaque assay was developed to visualize foci of infection at higher magnification

where zones of clearing were not readily evident. Confluent HFF monolayers on glass

coverslips in 24 well plates were infected with WT, ΔTgLaf, and COMP parasites in replete

media for 4 hours to allow for invasion. The monolayers were washed gently 3 times with PBS

and either fresh replete media or glutamine-depleted media were added for 3 or 6 days to

appropriate wells. Infected monolayers were washed and fixed with MeOH (-20 °C) and

subjected to IF using DAPI and GRA3, an antibody that detects the PVM. Individual plaques

were imaged on coded blinded slides using a 10X objective, and their perimeters and

encompassed areas were measured using Image J.  Host cell clearance within plaques was

similarly measured, and the extent of clearance was represented as a percentage for each

individual plaque. A total of three independent replicates were performed.

***Mouse infection***

4- to 6-week old CBA/J mice of both sexes (Jackson Laboratories, Bar Harbor, ME) were

injected intraperitoneally (i.p.) with either 100 WT, ΔTgLaf, or COMP tachyzoites, or with 20

tissue-cysts from brain homogenates derived from previously infected mice. In either case,

parasites/cysts were suspended in a final volume of 0.2 mL serum-free, Opti-MEM media

(Gibco). Mice were then monitored and assigned a body index score daily. Monitoring

frequency increased to twice a day once symptomatic throughout the course of infection as

previously described (68). When symptomatic, mice were administered a gel diet and wet chow

on the cage floor and given 0.25-0.5 mL saline solution subcutaneously as needed. Moribund

mice were humanely euthanized. Euthanasia of both moribund mice and mice sacrificed at the

time of tissue cyst harvest was performed by $CO_2$ asphyxiation, followed by cervical

dislocation. The number of mice used for each experiment is indicated within relevant figures.

All protocols were carried out under the approval of the University of Kentucky's Institutional

Animal Care and Use Committee (IACUC).

***Tissue cyst purification***

Tissue cysts were purified as previously described using discontinuous Percoll gradients (7,

91). Processing of two sex/infection-matched brains was performed on each gradient. Cysts

were collected in 1 mL fractions from the bottom of the centrifuged Percoll gradient using a

peristaltic pump adjusted to a flow rate of 2 mL/min. To quantify tissue cysts, 10-20 $\mu$L of each

fraction was placed into 100 $\mu$L PBS in the well of a 96-well plate, pelleted and directly

enumerated at 20X magnification in each well. Total cysts per mouse were calculated by

summing the total number of cysts in each fraction and dividing the total by two to adjust for

brain homogenization in pairs. Each pair of mice was presented as a single averaged data

point. Tissue cysts were pelleted onto slides using a Cytospin centrifuge and fixed and stored

in 100% MeOH (-20$^\circ$C) until staining.

***Determination of cyst diameter and packing density***

Tissue cysts fixed on slides were co-stained with DAPI and DBA to obtain nuclear profiles for

bradyzoites and an outline of the cyst for diameter measurement. The diameter was

determined in Fiji (ImageJ) by drawing a circle around the cyst and computing the diameter.

Packing density was determined as described previously (7). Briefly, DAPI images of cysts

were opened in BradyCount1.0, and threshold levels were adjusted as needed to capture all

nuclear profiles. After BradyCount1.0 provided the number of bradyzoite nuclei, the packing

density was determined by dividing the number of bradyzoites by the volume of the section.

The formula describing this calculation is: $PD = N / (\pi r^2 \times h)$ where PD=packing density, N = #

of bradyzoites, r = radius calculated from cyst diameter, and h = height of section (0.238 $\mu$m).

At least 10 cysts from each line at each timepoint were measured.

***Preparation of in vivo tissue cysts for TEM imaging***

To prepare *T. gondii* tissue cysts generated *in vivo* for TEM, cysts were isolated from the

brains of infected mice as detailed above through the counting step. The Percoll fraction

containing mouse red blood cells (RBCs) was also recovered. After combining Percoll

fractions containing tissue cysts and diluting with PBS to a volume of 15 mL (maximum of two

1-mL fractions were combined before dilution), cysts were pelleted for 15 min at 1000*g* at 4

°C. To maximize cyst recovery, 10 mL supernatant was removed, and the remaining 5 mL was

divided into 1 mL fractions for the top 4 mL, and the bottom 1 mL directly above the pellet was

sub-fractionated into 100 $\mu$L volumes. Typically, the majority of cysts were localized to within

300 $\mu$L of the Percoll pellet, rather than in the pellet itself. Sub-fractions containing cysts were

once again combined and diluted (typically 200-300 $\mu$L diluted with 1-mL PBS) in a 1.5 mL

Eppendorf tube, and then pelleted in a swinging-bucket rotor for 10 min at 1000*g* and 4 °C.

Leaving the pellet undisturbed, all but 50 $\mu$L of the supernatant was removed. A small volume

(~5-10 $\mu$L) of the reserved RBC fraction was added to the remaining volume for ease of

visualizing the pellet throughout the remaining processing steps.

To ensure the detection of the relatively rare cyst population, a previously described protocol

(71, 72) was adapted to concentrate the cysts into a small agarose block. A 1.33X fixative

solution of glutaraldhehyde (GA) in cacodylate buffer was prepared containing 4% GA and 133

mM sodium cacodylate. 150 $\mu$L fixative solution was then added to the 50 $\mu$L sample, bringing

the total volume to 200 $\mu$L such that the final concentration of GA was 3% and sodium

cacodylate was 100 mM. Cysts were then incubated at room temperature for 1 h in fixative.

While cysts were in fixative, 4% low-melt agarose (BioRad) was prepared in 100 mM sodium

cacodylate buffer and kept liquid at 70 $^o$C until needed. After fixation, cysts were pelleted again

at 1000$g$ for 10 min at room temperature in a table-top centrifuge. All but 50 $\mu$L supernatant

was once again removed, and 200 $\mu$L warm low-melt agarose was slowly added on top of

fixed, pelleted cysts (3.2% agarose, final concentration). Suspension was then centrifuged

again at 1000$g$ for 10 min at 30 $^o$C to keep the agarose semi-liquid, and then placed on ice for

20 min to solidify agarose. After solidification, entire agarose plug was removed from tube with

a small a wooden dowel that had been whittled into a thin scoop. This agarose plug was placed

in a Petri dish, and the pellet was carefully cut out of the plug with a razor blade to create a 1

mm$^3$ block. The agarose block was then stored in 1X GA/cacodylate buffer overnight at 4$^o$C.

Processing of the block from post-stain onward was then identical to TEM processing

described above. During sectioning, thick sections cut on a glass knife were stained with

toluidine blue and examined for cysts using a light microscope prior to ultra-thin sectioning

once the cyst containing later was identified.

***Data analyses***

All data analyses, including graph preparation and statistics, were performed using GraphPad

Prism 9.

----------------------------------------------------------------------------------------------------------------

***Acknowledgements:*** The authors wish to acknowledge Dr. Elizabeth Watts who conducted

early exploratory studies that encouraged the pursuit of this project, Jim Begley (Imaging

Center, University of Kentucky) for preparation of EM samples and willingness to explore new

methodologies for isolating rare *in vivo* cysts, Jillian Cramer (Electron Microscopy Center,

University of Kentucky) for acquisition of TEM images, and Jennifer Strange (Flow Cytometry

and Immune Monitoring Core Facility, University of Kentucky) for assistance with isolation of

GFP+ *T. gondii* mutants using flow cytometry.

***Author contributions:***

Conceptualization (R.D.M., A.P.S., M.S.G.), Methodology (R.D.M., A.D., C.O.B., A.P.S.,

M.S.G.), Formal analysis (R.D.M., C.A.T., J.S.M., L.E.A.Y., R.C.S., C.W.V.K., A.P.S.),

Investigation (R.D.M., C.A.T., J.S.M., L.E.A.Y., A.P.S.), Resources (M.S.G., A.P.S.), Writing—

original draft (R.D.M.), Writing—review and editing (all authors), Visualization (R.D.M., C.A.T.,

J.S.M., A.P.S.), Supervision (M.S.G., A.P.S.), Project administration (R.D.M., M.S.G., A.P.S.),

Funding acquisition (R.D.M., C.W.V.K., M.S.G., A.P.S.).

***Funding:*** This work was supported by: GRFP 1247392 to R.D.M., NIH grants R21 AI150631

and R01 AI145335 to A.P.S., and NIH grants R35 NS116824, P01 NS097197, and NSF CHE

1808304 to C.W.V.K. and MCB 2308488 to M.S.G.

***Conflicts of Interest:*** none

***Abbreviations:*** AG, amylopectin granule; CDPK2, calcium dependent protein kinase2;

COMP, TgLaforin-complemented parasite line; CRISPR, clustered regularly interspersed short

palindromic repeats; DBA, *Dolichos biflorus* agglutinin lectin; $\Delta$TgLaf, TgLaforin-KO parasite

line; DHFR, dihydrofolate reductase; FACS, fluorescence activated cell sorting; GAA, acid-$\alpha$-

amyloglucosidase; Gal-NAc, N-acetylgalactosamine; GAPDH1, glyceraldehyde-3-phosphate

dehydrogenase; GC/MS, gas chromatography/mass spectrometry; GFP, green fluorescent

protein; GT1, glucose transporter1; GWD, glucan, water di-kinase; HA, hemagglutinin; HFF,

human foreskin fibroblasts; HK, hexokinase; HR, homologous recombination; HXGPRT,

hypoxanthine-xanthine-guanine phosphoribosyl transferase; IF, immunofluorescence; i.p.,

intraperitoneally; KO, knockout; MPA, mycophenolic acid; PAM, protospacer adjacent motif;

PAS, periodic acid-Schiff; PV, parasitophorous vacuole; PWD, phospho-glucan, water di-

kinase; PYK1, pyruvate kinase1; RBC, red blood cell; RT, room temperature; SEX4, starch-

excess4; SS, starch/glycogen synthase; TCA, tricarboxylic acid; TEM, transmission electron

microscopy; UTR, untranslated region; WT, parental ME49$\Delta$HXGPRT parasite line used in this

study.

**REFERENCES**

1. Flegr J, Prandota J, Sovičková M, Israili ZH. Toxoplasmosis--a global threat. Correlation of latent toxoplasmosis with specific disease burden in a set of 88 countries. PLoS One. 2014;9(3):e90203.

2. Jones JL, Dubey JP. Foodborne toxoplasmosis. Clinical infectious diseases : an official publication of the Infectious Diseases Society of America. 2012;55(6):845-51.

3. Derouin F, Garin YJF. *Toxoplasma gondii* : Blood and tissue kinetics during acute and chronic infections in mice. Experimental Parasitology. 1991;73(4):460-8.

4. Jeffers V, Tampaki Z, Kim K, Sullivan WJ, Jr. A latent ability to persist: differentiation in *Toxoplasma gondii* . Cell Mol Life Sci. 2018;75(13):2355-73.

5. Luft BJ, Remington JS. Toxoplasmic Encephalitis in AIDS. Clinical Infectious Diseases. 1992;15(2):211-22.

6. Cerutti A, Blanchard N, Besteiro S. The Bradyzoite: A Key Developmental Stage for the Persistence and Pathogenesis of Toxoplasmosis. Pathogens. 2020;9(3):234.

7. Watts E, Zhao Y, Dhara A, Eller B, Patwardhan A, Sinai AP. Novel approaches reveal that *Toxoplasma gondii* bradyzoites within tissue cysts are dynamic and replicating entities in vivo. MBio. 2015;6(5):e01155-15.

8. Place BC, Troublefield C, Murphy RD, Sinai AP, Patwardhan A. Computer Aided Image Processing to Facilitate Determination of Congruence in Manual Classification of Mitochondrial Morphologies in *Toxoplasma gondii* Tissue Cysts. Annu Int Conf IEEE Eng Med Biol Soc. 2021;2021:3509-13.

9.  MacNeill GJ, Mehrpouyan S, Minow MA, Patterson JA, Tetlow IJ, Emes MJ, et al. Starch

as a source, starch as a sink: the bifunctional role of starch in carbon allocation. Journal of

Experimental Botany. 2017;68(16):4433-53.

10. Dubey JP, Frenkel JK. Feline toxoplasmosis from acutely infected mice and the

development of Toxoplasma cysts. J Protozool. 1976;23(4):537-46.

11. Coppin A, Dzierszinski F, Legrand S, Mortuaire M, Ferguson D, Tomavo S.

Developmentally regulated biosynthesis of carbohydrate and storage polysaccharide during

differentiation and tissue cyst formation in *Toxoplasma gondii* . Biochimie. 2003;85(3-4):353-

61.

12. Coppin A, Varre J-S, Lienard L, Dauvillee D, Guerardel Y, Soyer-Gobillard M-O, et al.

Evolution of plant-like crystalline storage polysaccharide in the protozoan parasite *Toxoplasma*

*gondii* argues for a red alga ancestry. Journal of Molecular Evolution. 2005;60(2):257-67.

13. Guérardel Y, Leleu D, Coppin A, Liénard L, Slomianny C, Strecker G, et al. Amylopectin

biogenesis and characterization in the protozoan parasite *Toxoplasma gondii* , the intracellular

development of which is restricted in the HepG2 cell line. Microbes and infection. 2005;7(1):41-

8.

14. Uboldi AD, McCoy JM, Blume M, Gerlic M, Ferguson DJ, Dagley LF, et al. Regulation of

starch stores by a Ca2+-dependent protein kinase is essential for viable cyst development in

*Toxoplasma gondii* . Cell Host & Microbe. 2015;18(6):670-81.

15. Yang J, Yang C, Qian J, Li F, Zhao J, Fang R. *Toxoplasma gondii* α-amylase deletion

mutant is a promising vaccine against acute and chronic toxoplasmosis. Microb Biotechnol.

2020.

16. Yang J, He Z, Chen C, Zhao J, Fang R. Starch branching enzyme 1 is important for

amylopectin synthesis and cyst reactivation in *Toxoplasma gondii* . Microbiology Spectrum.

2022;10(3):e01891-21.

17. Lyu C, Yang X, Yang J, Hou L, Zhou Y, Zhao J, et al. Role of amylopectin synthesis in

*Toxoplasma gondii*  and its implication in vaccine development against toxoplasmosis. Open

Biol. 2021;11(6):200384.

18. Blume M, Nitzsche R, Sternberg U, Gerlic M, Masters SL, Gupta N, et al. A *Toxoplasma*

*gondii* gluconeogenic enzyme contributes to robust central carbon metabolism and is essential

for replication and virulence. Cell host & microbe. 2015;18(2):210-20.

19. Ralton JE, Sernee MF, McConville MJ. Evolution and function of carbohydrate reserve

biosynthesis in parasitic protists. Trends Parasitol. 2021.

20. Sernee MF, Ralton JE, Nero TL, Sobala LF, Kloehn J, Vieira-Lara MA, et al. A family of

dual-activity glycosyltransferase-phosphorylases mediates mannogen turnover and virulence

in Leishmania parasites. Cell host & microbe. 2019;26(3):385-99. e9.

21. Kötting O, Kossmann J, Zeeman SC, Lloyd JR. Regulation of starch metabolism: the age

of enlightenment? Curr Opin Plant Biol. 2010;13(3):321-9.

22. Silver DM, Kötting O, Moorhead GB. Phosphoglucan phosphatase function sheds light on

starch degradation. Trends in plant science. 2014;19(7):471-8.

23. Stitt M, Zeeman SC. Starch turnover: pathways, regulation and role in growth. Curr Opin

Plant Biol. 2012;15(3):282-92.

24. Ritte G, Lloyd JR, Eckermann N, Rottmann A, Kossmann J, Steup M. The starch-related

R1 protein is an alpha -glucan, water dikinase. Proc Natl Acad Sci U S A. 2002;99(10):7166-

71.

25. Kötting O, Pusch K, Tiessen A, Geigenberger P, Steup M, Ritte G. Identification of a novel

enzyme required for starch metabolism in Arabidopsis leaves. The phosphoglucan, water

dikinase. Plant Physiol. 2005;137(1):242-52.

26. Takeda Y, Hizukuri S. Re-examination of the action of sweet-potato beta-amylase on

phosphorylated (1→4)-χ-d-glucan. Carbohydrate Research. 1981;89(1):174-8.

27. Edner C, Li J, Albrecht T, Mahlow S, Hejazi M, Hussain H, et al. Glucan, water dikinase

activity stimulates breakdown of starch granules by plastidial beta-amylases. Plant Physiol.

2007;145(1):17-28.

28. Hejazi M, Fettke J, Haebel S, Edner C, Paris O, Frohberg C, et al. Glucan, water dikinase

phosphorylates crystalline maltodextrins and thereby initiates solubilization. Plant J.

2008;55(2):323-34.

29. Hansen PI, Spraul M, Dvortsak P, Larsen FH, Blennow A, Motawia MS, et al. Starch

phosphorylation--maltosidic restrains upon 3'- and 6'-phosphorylation investigated by chemical

synthesis, molecular dynamics and NMR spectroscopy. Biopolymers. 2009;91(3):179-93.

30. Murphy RD, Chen T, Lin J, He R, Wu L, Pearson CR, et al. The Toxoplasma glucan

phosphatase TgLaforin utilizes a distinct functional mechanism that can be exploited by

therapeutic inhibitors. Journal of Biological Chemistry. 2022:102089.

31. Kloehn J, Krishnan A, Tonkin CJ, McConville MJ, Soldati-Favre D. Chapter 10 - Metabolic

networks and metabolomics. In: Weiss LM, Kim K, editors. *Toxoplasma gondii* (Third Edition):

Academic Press; 2020. p. 451-97.

32. Chen P, Lyu C, Wang Y, Pan M, Lin X, Shen B. Key roles of amylopectin synthesis and

degradation enzymes in the establishment and reactivation of chronic toxoplasmosis. Animal

Diseases. 2023;3(1):1-11.

33. Zeeman SC, Northrop F, Smith AM, Rees T. A starch-accumulating mutant of Arabidopsis

thaliana deficient in a chloroplastic starch-hydrolysing enzyme. Plant J. 1998;15(3):357-65.

34. Kötting O, Santelia D, Edner C, Eicke S, Marthaler T, Gentry MS, et al. STARCH-EXCESS4

is a laforin-like Phosphoglucan phosphatase required for starch degradation in Arabidopsis

thaliana. Plant Cell. 2009;21(1):334-46.

35. Lafora GR. Band 205. In: Johannes O, editor. XXI Über das Vorkommen amyloider

Körperchen im Innern der Ganglienzellen; zugleich Ein Beitrag zum Studium der amyloiden

Substanz im Nervensystem: De Gruyter; 2021. p. 295-303.

36. Minassian BA, Lee JR, Herbrick JA, Huizenga J, Soder S, Mungall AJ, et al. Mutations in a

gene encoding a novel protein tyrosine phosphatase cause progressive myoclonus epilepsy.

Nat Genet. 1998;20(2):171-4.

37. Gentry MS, Guinovart JJ, Minassian BA, Roach PJ, Serratosa JM. Lafora disease offers a

unique window into neuronal glycogen metabolism. Journal of Biological Chemistry.

2018;293(19):7117-25.

38. Duran J, Hervera A, Markussen KH, Varea O, López-Soldado I, Sun RC, et al. Astrocytic

glycogen accumulation drives the pathophysiology of neurodegeneration in Lafora disease.

Brain. 2021;144(8):2349-60.

39. Sugi T, Tu V, Ma Y, Tomita T, Weiss LM. *Toxoplasma gondii* requires glycogen

phosphorylase for balancing amylopectin storage and for efficient production of brain cysts.

MBio. 2017;8(4):e01289-17.

40. Cao XZ, Wang JL, Elsheikha HM, Li TT, Sun LX, Liang QL, et al. Characterization of the

role of Amylo-alpha-1, 6-glucosidase protein in the infectivity of *Toxoplasma gondii*. Frontiers

in Cellular and Infection Microbiology. 2019;9:418.

41. Wang J-L, Li T-T, Elsheikha HM, Liang Q-L, Zhang Z-W, Wang M, et al. The protein

phosphatase 2A holoenzyme is a key regulator of starch metabolism and bradyzoite

differentiation in *Toxoplasma gondii* . Nature communications. 2022;13(1):7560.

42. Zhao M, Yang Y, Shi Y, Chen X, Yang Y, Pan L, et al. PP2Acα-B′/PR61 Holoenzyme of

*Toxoplasma gondii* Is Required for the Amylopectin Metabolism and Proliferation of

Tachyzoites. Microbiology Spectrum. 2023;11(3):e00104-23.

43. Gentry MS, Dowen RH, Worby CA, Mattoo S, Ecker JR, Dixon JE. The phosphatase laforin

crosses evolutionary boundaries and links carbohydrate metabolism to neuronal disease. The

Journal of cell biology. 2007;178(3):477-88.

44. Gentry MS, Pace RM. Conservation of the glucan phosphatase laforin is linked to rates of

molecular evolution and the glucan metabolism of the organism. BMC evolutionary biology.

2009;9(1):138.

45. Baba O. Production of monoclonal antibody that recognizes glycogen and its application

for immunohistochemistry. Kokubyo Gakkai zasshi The Journal of the Stomatological Society,

Japan. 1993;60(2):264-87.

46. Oe Y, Baba O, Ashida H, Nakamura KC, Hirase H. Glycogen distribution in the microwave-

fixed mouse brain reveals heterogeneous astrocytic patterns. Glia. 2016;64(9):1532-45.

47. van de Weerd R, Berbís MA, Sparrius M, Maaskant JJ, Boot M, Paauw NJ, et al. A Murine

Monoclonal Antibody to Glycogen: Characterization of Epitope-Fine Specificity by Saturation

Transfer Difference (STD) NMR Spectroscopy and Its Use in Mycobacterial Capsular α-Glucan

Research. Chembiochem. 2015;16(6):977-89.

48. Skurat AV, Segvich DM, DePaoli-Roach AA, Roach PJ. Novel method for detection of

glycogen in cells. Glycobiology. 2017;27(5):416-24.

49. Guimarães EV, de Carvalho L, Barbosa HS. An alternative technique to reveal

polysaccharides in *Toxoplasma gondii* tissue cysts. Mem Inst Oswaldo Cruz. 2003;98(7):915-

7.

50. Sidik SM, Hackett CG, Tran F, Westwood NJ, Lourido S. Efficient genome engineering of

*Toxoplasma gondii* using CRISPR/Cas9. PloS one. 2014;9(6):e100450.

51. Fritz HM, Buchholz KR, Chen X, Durbin-Johnson B, Rocke DM, Conrad PA, et al.

Transcriptomic analysis of toxoplasma development reveals many novel functions and

structures specific to sporozoites and oocysts. PloS one. 2012;7(2):e29998.

52. Choi CP, Moon AS, Back PS, Jami-Alahmadi Y, Vashisht AA, Wohlschlegel JA, et al. A

photoactivatable crosslinking system reveals protein interactions in the *Toxoplasma gondii*

inner membrane complex. PLoS biology. 2019;17(10):e3000475.

53. Shen B, Brown KM, Lee TD, Sibley LD. Efficient gene disruption in diverse strains of

*Toxoplasma gondii* using CRISPR/CAS9. mBio. 2014;5(3):e01114-14.

54. Sidik SM, Huet D, Ganesan SM, Huynh MH, Wang T, Nasamu AS, et al. A Genome-wide

CRISPR Screen in Toxoplasma Identifies Essential Apicomplexan Genes. Cell.

2016;166(6):1423-35.e12.

55. Markus BM, Bell GW, Lorenzi HA, Lourido S. Optimizing systems for Cas9 expression in

*Toxoplasma gondii* . MSphere. 2019;4(3):e00386-19.

56. Yang J, Yang C, Qian J, Li F, Zhao J, Fang R. *Toxoplasma gondii* α-amylase deletion

mutant is a promising vaccine against acute and chronic toxoplasmosis. Microb Biotechnol.

2020;13(6):2057-69.

57. Xia N, Ye S, Liang X, Chen P, Zhou Y, Fang R, et al. Pyruvate homeostasis as a

determinant of parasite growth and metabolic plasticity in *Toxoplasma gondii* . MBio.

2019;10(3):e00898-19.

58. Brewer MK, Uittenbogaard A, Austin GL, Segvich DM, DePaoli-Roach A, Roach PJ, et al.

Targeting pathogenic Lafora bodies in Lafora disease using an antibody-enzyme fusion. Cell

metabolism. 2019;30(4):689-705. e6.

59. Sun RC, Young LEA, Bruntz RC, Markussen KH, Zhou Z, Conroy LR, et al. Brain glycogen

serves as a critical glucosamine cache required for protein glycosylation. Cell Metab.

2021;33(7):1404-17 e9.

60. King EF, Cobbold SA, Uboldi AD, Tonkin CJ, McConville MJ. Metabolomic Analysis of

*Toxoplasma gondii* Tachyzoites. *Toxoplasma gondii* : Springer; 2020. p. 435-52.

61. Nitzsche R, Zagoriy V, Lucius R, Gupta N. Metabolic cooperation of glucose and glutamine

is essential for the lytic cycle of obligate intracellular parasite *Toxoplasma gondii* . Journal of

Biological Chemistry. 2016;291(1):126-41.

62. Oppenheim RD, Creek DJ, Macrae JI, Modrzynska KK, Pino P, Limenitakis J, et al.

BCKDH: the missing link in apicomplexan mitochondrial metabolism is required for full

virulence of *Toxoplasma gondii* and Plasmodium berghei. PLoS pathogens. 2014;10(7).

63. Endo T, Sethi K, Piekarski G. *Toxoplasma gondii* : Calcium Ionophore A23187-mediated

exit of trophozoites from infected murine macrophages. Experimental parasitology.

1982;53(2):179-88.

64. Lourido S, Tang K, Sibley LD. Distinct signalling pathways control Toxoplasma egress and

host-cell invasion. The EMBO journal. 2012;31(24):4524-34.

65. Ferguson DJ. Use of molecular and ultrastructural markers to evaluate stage conversion of

*Toxoplasma gondii* in both the intermediate and definitive host. International journal for

parasitology. 2004;34(3):347-60.

66. Sinai AP. Chapter 11 - The *Toxoplasma gondii* Parasitophorous Vacuole Membrane: A

Multifunctional Organelle in the Infected Cell. In: Weiss LM, Kim K, editors. *Toxoplasma gondii*

(Second Edition). Boston: Academic Press; 2014. p. 375-87.

67. Boothroyd JC, Black M, Bonnefoy S, Hehl A, Knoll LJ, Manger ID, et al. Genetic and

biochemical analysis of development in *Toxoplasma gondii*. Philos Trans R Soc Lond B Biol

Sci. 1997;352(1359):1347-54.

68. Troublefield CA, Miracle JS, Murphy RD, Donkin RW, Sinai AP. Factors influencing tissue

cyst yield in a murine model of chronic toxoplasmosis. Infection and Immunity. 2023:e00566-

22.

69. Watts EA, Dhara A, Sinai AP. Purification *Toxoplasma gondii* Tissue Cysts Using Percoll

Gradients. Curr Protoc Microbiol. 2017;45:20C 2 1-C 2 19.

70. Quach EV, Cao B, Babacarkhial E, Ho D, Sharma J, Guiton PS. Phosphoglucomutase 1

contributes to optimal cyst development in *Toxoplasma gondii*. BMC Research Notes.

2022;15(1):1-8.

71. Kumar S, Ciraolo G, Hinge A, Filippi MD. An efficient and reproducible process for

transmission electron microscopy (TEM) of rare cell populations. J Immunol Methods.

2014;404:87-90.

72. Kumar S, Filippi MD. An Alternative Approach for Sample Preparation with Low Cell

Number for TEM Analysis. J Vis Exp. 2016(116).

73. Sinai AP, Suvorova ES. The RESTRICTION checkpoint: a window of opportunity governing

developmental transitions in *Toxoplasma gondii*. Current opinion in microbiology. 2020;58:99-

105.

74. Xue Y, Theisen TC, Rastogi S, Ferrel A, Quake SR, Boothroyd JC. A single-parasite

transcriptional atlas of *Toxoplasma gondii* reveals novel control of antigen expression. Elife.

2020;9:e54129.

75. Olson WJ, Martorelli Di Genova B, Gallego-Lopez G, Dawson AR, Stevenson D, Amador-

Noguez D, et al. Dual metabolomic profiling uncovers Toxoplasma manipulation of the host

metabolome and the discovery of a novel parasite metabolic capability. PLoS pathogens.

2020;16(4):e1008432.

76. Zeeman SC, Smith SM, Smith AM. The diurnal metabolism of leaf starch. Biochemical

Journal. 2007;401(1):13-28.

77. Mahlow S, Orzechowski S, Fettke J. Starch phosphorylation: insights and perspectives.

Cellular and molecular life sciences. 2016;73(14):2753-64.

78. Niittyla T, Comparot-Moss S, Lue WL, Messerli G, Trevisan M, Seymour MD, et al. Similar

protein phosphatases control starch metabolism in plants and glycogen metabolism in

mammals. The Journal of biological chemistry. 2006;281(17):11815-8.

79. Shukla A, Olszewski KL, Llinás M, Rommereim LM, Fox BA, Bzik DJ, et al. Glycolysis is

important for optimal asexual growth and formation of mature tissue cysts by *Toxoplasma*

*gondii* . International journal for parasitology. 2018;48(12):955-68.

80. Blume M, Rodriguez-Contreras D, Landfear S, Fleige T, Soldati-Favre D, Lucius R, et al.

Host-derived glucose and its transporter in the obligate intracellular pathogen *Toxoplasma*

*gondii* are dispensable by glutaminolysis. Proceedings of the National Academy of Sciences.

2009;106(31):12998-3003.

81. Dubey R, Staker BL, Foe IT, Bogyo M, Myler PJ, Ngô HM, et al. Membrane skeletal

association and post-translational allosteric regulation of *Toxoplasma gondii* GAPDH1. Mol

Microbiol. 2017;103(4):618-34.

82. Araujo FG, Huskinson-Mark J, Gutteridge WE, Remington JS. In vitro and in vivo activities

of the hydroxynaphthoquinone 566C80 against the cyst form of *Toxoplasma gondii* .

Antimicrobial agents and chemotherapy. 1992;36(2):326-30.

83. Doggett JS, Nilsen A, Forquer I, Wegmann KW, Jones-Brando L, Yolken RH, et al.

Endochin-like quinolones are highly efficacious against acute and latent experimental

toxoplasmosis. Proc Natl Acad Sci U S A. 2012;109(39):15936-41.

84. McPhillie MJ, Zhou Y, Hickman MR, Gordon JA, Weber CR, Li Q, et al. Potent

Tetrahydroquinolone Eliminates Apicomplexan Parasites. Front Cell Infect Microbiol.

2020;10:203.

85. Brown KM, Long S, Sibley LD. Conditional Knockdown of Proteins Using Auxin-inducible

Degron (AID) Fusions in *Toxoplasma gondii* . Bio Protoc. 2018;8(4).

86. Bolte S, Cordelières FP. A guided tour into subcellular colocalization analysis in light

microscopy. J Microsc. 2006;224(Pt 3):213-32.

87. Mayoral J, Di Cristina M, Carruthers VB, Weiss LM. *Toxoplasma gondii* : Bradyzoite

Differentiation In Vitro and In Vivo. Methods in molecular biology. 2020;2071:269-82.

88. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an

open-source platform for biological-image analysis. Nat Methods. 2012;9(7):676-82.

89. Sinai AP, Webster P, Joiner KA. Association of host cell endoplasmic reticulum and

mitochondria with the *Toxoplasma gondii* parasitophorous vacuole membrane: a high affinity

interaction. J Cell Sci. 1997;110 ( Pt 17):2117-28.

90. Andres DA, Young LEA, Veeranki S, Hawkinson TR, Levitan BM, He D, et al. Improved

workflow for mass spectrometry–based metabolomics analysis of the heart. Journal of

Biological Chemistry. 2020;295(9):2676-86.

91. Watts EA, Dhara A, Sinai AP. Purification *Toxoplasma gondii* Tissue Cysts Using Percoll

Gradients. Curr Protoc Microbiol. 2017;45:20c.2.1-c.2.19.

1157

1158

Figure 1. Glucan dynamics in *T. gondii* ME49 tachyzoites and bradyzoites. **A,** Microscopy-based glucan evaluation of *T. gondii* tachyzoites and bradyzoites using PAS (left), IV58B6 (middle; α-glycogen IgM mAb), and TEM (right). **B,** GAA digest of AGs in tachyzoites and bradyzoites confirms specificity of IV58B6 antibody and PAS staining. All scale bars = 5 μm.

**Figure 2.** Endogenous tagging and localization of TgLaforin. **A,** Schematic depicting the
TgLaforin 3xHA-epitope tagging strategy. X = stop codon. **B,** Successful tagging of TgLaforin
(62 kDa) was verified using immunoblot analysis with an α-HA antibody with SAG1 used as a
loading control. **C,** IFA in WT/tagged parasites with α-HA antibody. **D,** Western blot analysis
of TgLaforin expression levels in tachyzoites (T) and bradyzoites (B). GAP45 is the loading
control. Decrease in SAG1 alongside increase in SRS9 confirms tachyzoite to bradyzoite
conversion. **E**, IFA of TgLaforin colocalization with PAS. Pearson's coefficient: 0.765. **F**, IFA of
TgLaforin colocalization with IV58B6. Pearson's coefficient: 0.737. All scale bars = 5 μm.

1178
1179
**Figure 3.** Loss of TgLaforin results in no gross glucan abnormalities under glucose replete
conditions. **A,** Schematic of TgLaforin KO strategy: the pyrimethamine-resistant DHFR* gene
containing 40-nt homologous arms (dark gray boxes on either end of gene) was inserted into
the TgLaforin locus via homologous recombination. A double stranded break was induced
using CRISPR/Cas9-GFP with a PAM site in the first exon. **B,** Inside/out PCR verification of
DHFR integration into the TgLaforin locus. Amplicons (PCR1-4) are illustrated in (A). **C,** Loss
of TgLaforin mRNA was confirmed by amplifying full-length TgLaforin cDNA generated from
both WT and ΔTgLaf strains. Actin cDNA amplification serves both as a loading control and as
a control to verify the absence of gDNA. **D,** Analysis of glucan levels in ΔTgLaf tachyzoites
using three different approaches: PAS and IV58B6 immunofluorescence staining, and TEM.
1190

1191
1192

**Figure 4.** ΔTgLaf parasites are dependent on glutamine for normal plaque formation. **A**, Steady-state metabolomics suggests upregulation of glutaminolysis in ΔTgLaf parasites. Metabolite levels of intracellular tachyzoites were analyzed after 48 hours of growth in HFFs by GC/MS analysis. Data were collected from 3 independent replicates. Statistical comparisons were done using unpaired two-tailed t-tests. Statistical significance is as follows: **p<0.01, ns=p>0.05, nd=not detected. **B**, Representative plaque assays are presented for each line and condition. **C**, Total plaque area (pixels) was measured across three independent replicates. **D**, Pixel area of nascent plaques after 3 days of growth as monitored by IF microscopy under both glutamine replete (+) and depleted conditions (-). **E**, Pixel area of plaques monitored as in (D) after 6 days of growth. Statistical comparisons for C-E were done using an ordinary one-way ANOVA using Tukey's post-hoc test to correct for multiple comparisons. Statistical significance is indicated as follows: *p<0.05, ****p<0.0001, ns=p>0.05.

1205
1206

1207
1208
**Figure 5**. Loss of TgLaforin results in aberrant glucan morphology and accumulation in *in vitro*
bradyzoites only visible by TEM. **A,** *In vitro* tachyzoite to bradyzoite conversion efficiency of
ΔTgLaf vs WT parasites as measured by DBA intensity. **B,** Change in PAS levels during
bradyzoite conversion by PAS levels. **C,** Representative TEM images of bradyzoites from each
indicated parasite line. At 4300x magnification, scale bar = 2 μm; at 8600x magnification, scale
bar = 1 μm. Arrowhead = canonical AG (white, round/ovoid); Arrow = aberrant AG (grey,
flattened, multi-lobed). **D,** Quantification of relative parasite AG content and **E,** AG grayness

across parasite lines using 8-bit grayscale (0=black, 255=white). Statistical comparisons were done using an ordinary one-way ANOVA with Tukey's post-hoc test to correct for multiple comparisons. Statistical significance is indicated as follows: $*p<0.05$, $***p<0.001$, $****p<0.0001$, ns=$p>0.05$.

Figure 6. Loss of TgLaforin attenuates tachyzoite virulence and cyst burden in mice. Equal numbers of male and female CBA/J mice were infected i.p. with 100 tachyzoites from each group for monitoring symptoms and death: **A,** Symptomology throughout the acute phase of infection. Mice were monitored 1-2x/day and assigned a body score index ranging from asymptomatic (Stage 0) to moribund/deceased. **B,** Kaplan-Meier curve of mouse survival throughout acute tachyzoite infection. **C,** Mice that survived 4-weeks were euthanized, and cysts were counted as done previously (1, 2). Error bars depict SD from the mean. Statistical comparison for Kaplan Meier curves is indicated on plot, and statistical comparison of cyst burden was done using unpaired two-tailed t-tests. Statistical significance: *p<0.05, ***p<0.0002.

Figure 7. Mouse infection with ΔTgLaf tissue cysts results in milder illness and lower cyst burden. CBA/J mice were infected i.p. with 20 cysts taken from previously infected mouse brains and monitored for symptomology and death. **A,** Symptomology throughout the acute phase of infection as described in Figure 6B. **B,** Kaplan-Meier curve of mouse survival during acute tachyzoite infection. **C,** Cysts/brain after 4-week bradyzoite infection. Cyst numbers were determined as described in Figure 6C. Statistical comparison of cyst burden was done using unpaired two-tailed t-tests. Statistical significance: *p<0.05, ns=p>0.05.

**Figure 8.** Cysts formed by ΔTgLaf parasites are smaller and less densely packed with bradyzoites. **A,** Packing density comparison among WT, ΔTgLaf, and COMP lines at week 4. Calculations were done using BradyCount1.0 to identify and quantify nuclear profiles from DAPI-stained tissue cysts (1). **B,** Cyst diameter comparison among WT, ΔTgLaf, and COMP lines as measured in ImageJ at week 4. **C,** Packing density comparison at week 6, as described in (A). **D,** Cyst diameter comparison at week 6, as described in (B). Statistical comparisons were done using an ordinary one-way ANOVA using Tukey's post-hoc test to correct for multiple comparisons. Error bars depict SD from the mean. Statistical significance is indicated as follows: *p<0.05, **p<0.01, ns=p>0.05.

**Figure 9.** *T. gondii* bradyzoites accumulate excess AGs in the absence of TgLaforin. Upper panels: scale bar = 5 μm; lower panels (zoom of boxed region from upper panel): scale bar = 2 μm.