TgLaforin, a glucan phosphatase, reveals the dynamic role of storage polysaccharides in

Toxoplasma gondii tachyzoites and bradyzoites

Robert D. Murphy[1,2,a], Cortni A. Troublefield[1b#], Joy S. Miracle[1c#], Lyndsay E.A. Young[2],

Aashutosh Tripathi[1d], Corey O. Brizzee[2e], Animesh Dhara[1], Abhijit Patwardhan[3], Ramon

C. Sun[2fg], Craig W. Vander Kooi[2fg], Matthew S. Gentry[2fg], Anthony P. Sinai[1*]

Affiliations at time research was conducted:

1. Department of Microbiology, Immunology, and Molecular Genetics, College of

Medicine, University of Kentucky, Lexington, KY 40536, USA

2. Department of Molecular and Cellular Biochemistry, College of Medicine,

University of Kentucky, Lexington, KY 40536, USA

3. F. Joseph Halcomb III, MD. Department of Biomedical Engineering, College of

Engineering, University of Kentucky, Lexington KY 40506, USA

#These authors contributed equally to this work.

*For correspondence: Anthony P. Sinai, sinai@uky.edu

Current affiliations:

a. Maryland Department of Health, Laboratories Administration, Baltimore, MD

21205

b. Good Samaritan Hospital and Albert B. Chandler Medical Center Emergency

Departments, University of Kentucky, Lexington KY 40536

c. Department of Forestry and Natural Resources, Martin-Gatton College of

Agriculture, Food and Environment, University of Kentucky, Lexington, KY 40546

d. Department of Biochemistry, Stanford University School of Medicine, Palo Alto CA

94305

e. Demeetra AgBio, 2277 Thunderstick Dr., Lexington, KY, USA

f. Department of Biochemistry and Molecular Biology, College of Medicine,

University of Florida, Gainesville, FL, USA

g. Center for Advanced Spatial Biomolecule Research, University of Florida,

Gainesville, FL, USA

Running title: TgLaforin regulation of amylopectin granules in Toxoplasma

Keywords: *Toxoplasma gondii*, amylopectin granule, glucan, reversible phosphorylation,

laforin, phosphatase, starch, glycogen, steady-state metabolomics, glutamine

ABSTRACT

The asexual stages of *Toxoplasma gondii* are defined by the rapidly growing tachyzoite

during the acute infection and by the slow growing bradyzoite housed within tissue cysts

during the chronic infection. These stages represent unique physiological states, each

with distinct glucans reflecting differing metabolic needs. A defining feature of *T. gondii*

bradyzoites is the presence of insoluble storage glucans known as amylopectin granules

(AGs) that are believed to play a role in reactivation, but their functions during the chronic

infection remain largely unexplored. More recently, the presence of storage glucans has

been recognized in tachyzoites where their precise function and architecture have yet to

be fully defined. Importantly, the *T. gondii* genome encodes activities needed for glucan

turnover: a glucan phosphatase (TgLaforin; TGME49_205290) and a glucan kinase

(TgGWD; TGME49_214260) that catalyze a cycle of reversible glucan phosphorylation

required for glucan degradation by amylases. The expression of these enzymes in

tachyzoites supports the existence of a storage glucan, evidence that is corroborated by

specific labeling with the anti-glycogen antibody IV58B6. Disruption of reversible glucan

phosphorylation via a CRISPR/Cas9 knockout (KO) of TgLaforin revealed no growth

defects under nutrient-replete conditions in tachyzoites. However, the growth of

TgLaforin-KO tachyzoites was severely stunted when starved of glutamine, even under

glucose replete conditions. The loss of TgLaforin also resulted in the attenuation of acute

virulence in mice accompanied by a lower cyst burden. While the absence of TgLaforin

did not impact the size distribution of tissue cysts, bradyzoites within both in vitro

generated and in vivo tissue cysts exhibited profound changes in AG levels and

morphology. Quantification of relative AG levels using AmyloQuant, an imaging-based

application, revealed the starch-excess phenotype associated with the loss of TgLaforin

is heterogeneous across tissue cysts and exhibits a temporal profile linked to an emerging

AG cycle. Together, these data demonstrate the importance of glucan turnover across

the *T. gondii* asexual cycle. These findings, alongside our previously identified class of

small molecules that inhibit TgLaforin, implicate reversible glucan phosphorylation as a

legitimate target for the development of new drugs against chronic *T. gondii* infections.

INTRODUCTION

Toxoplasma gondii is an opportunistic protozoan parasite of all warm-blooded animals

that infects one-third of humans worldwide (1, 2). Humans are primarily infected through

the consumption of an encysted form of the parasite: either the oocysts shed in cat feces

or tissue cysts found in undercooked meat from a chronically infected animal (3).

Encysted parasites convert into tachyzoites that rapidly divide and disseminate

throughout the body of the host, defining the acute phase of infection (4). Under host

immune pressure, tachyzoites convert into slow-growing bradyzoites that populate tissue

cysts which are found predominantly in the central nervous system and muscle, defining

the chronic phase of infection (5, 6). Tissue cysts are believed to persist for the lifetime

of the host and possess the ability to reactivate into tachyzoites in the context of

immunosuppression. Reactivation can result in the life-threatening symptoms of

toxoplasmosis, with toxoplasmic encephalitis being the primary condition leading to

mortality (7, 8). The current lack of insights into bradyzoite physiology *in vivo* precludes

the basic understanding needed for the development of drugs that either can clear tissue

cysts or target encysted bradyzoites so as to prevent reactivation (9).

Until recently, bradyzoites within tissue cysts were considered to be dormant,

metabolically inert entities. This view was challenged by our demonstration that encysted

bradyzoites replicate (10, 11). Moreover, bradyzoite physiology is both diverse and

complex as viewed through the lens of mitochondrial activity (12, 13), replication status

(11), and, importantly, amylopectin granule (AG) accumulation (14). Although the function

of AGs in bradyzoites has not been confirmed, an understanding of the roles of

polysaccharides elsewhere suggests that AGs are a source of energy and biosynthetic

potential needed for persistence, replication, reactivation, and transmission (15). These

assumptions remain to be tested, and thus much like bradyzoites themselves, the role of

AGs in the *T. gondii* lifecycle is poorly understood. Our first insights into the potential

relationships connecting AG to intermediate metabolism via mitochondrial and replication

activities have been exposed using imaging-based approaches (14).

AGs are large glucans found in the cytoplasm of bradyzoites that have classically served

as a morphological feature distinguishing them from tachyzoites (16-19). Toxoplasma

AGs are much like plant starch in that they are water-insoluble storage polysaccharides

composed of branched chains of glucose (18). Unlike plant starch, however, AGs contain

no detectable amylose (unbranched chains of glucose) (18). More recently, the presence

of small, punctate, cytoplasmic glucans in tachyzoites that are only visible by periodic

acid-Schiff (PAS) staining have been recognized (20-22), and the presence of the glucan

is dependent on the *T. gondii* starch synthase (TgSS; TGME49_222800) (23). Like animal

glycogen, this tachyzoite storage polysaccharide is rapidly turned over (20), as has been

observed in other protozoa (24-26), and provides glucose for glycolysis (23, 27). The

observation that large, insoluble glucans do not accumulate within tachyzoites as they do

within bradyzoites suggests that the tachyzoite glucan could be a distinct and labile form

of stored glucose, likely glycogen-like, although its exact chemical and structural identity

remains unknown.

Glucose release from starch in plants requires a cycle of direct, reversible glucan

phosphorylation to solubilize the starch surface, allowing access to degradation enzymes

such as amylases, branching enzymes, and a phosphorylase (28-30). The cycle begins

with the addition of phosphate directly to glucose by the glucan, water-dikinase (GWD)

and phospho-glucan, water dikinase (PWD) that results in the unwinding of glucose

chains within starch, solubilizing the starch surface (31, 32). Glucose-releasing enzymes

(amylases) then degrade starch until the glucan-bound phosphate becomes a steric

hindrance, at which point a glucan phosphatase is needed to remove the phosphate and

reset the cycle (33-35). *T. gondii* encodes all the activities needed for glucan degradation

and reversible glucan phosphorylation including the glucan phosphatase, TgLaforin

(TGME49_205290) (36), and glucan dikinase, *T. gondii* GWD (TgGWD;

TGME49_214260) (27). The central role of reversible glucan phosphorylation in plants

is seen in *Arabidopsis thaliana* where loss of the plant glucan phosphatase, starch-excess

4 (SEX4), results in excess starch accumulation, aberrant starch morphology, and

severely stunted plant growth (37, Zeeman, 1998 #1387). Additionally, loss of the glucan

phosphatase, laforin, in humans, results in hyperphosphorylated glycogen that

aggregates in neurons and astrocytes causing a fatal neurodegenerative childhood

dementia and epilepsy (38-40). In *T. gondii*, perturbations of several genes related to

glucan metabolism also result in a variety of similar defects including aberrant glucan

accumulation, rewiring of central carbon metabolism, and virulence defects in mice,

highlighting the central metabolic role of glucan metabolism in *T. gondii* (20-23, 27, 41-

44)

In this study, we build on our understanding of reversible glucan phosphorylation and its

relevance to parasite metabolism in *T. gondii*. We have recently demonstrated that

TgLaforin is the glucan phosphatase in *T. gondii*, and that TgLaforin represents a unique

and viable drug target (36, 45, 46). Here, we investigate the role of TgLaforin throughout

the asexual stages by knocking out TgLaforin in Type II ME49 parasites. While we

expected to observe effects related to the loss of TgLaforin exclusively in bradyzoites

where AGs are typically observed, these effects appeared to be connected to an

emerging AG temporal cycle (14). More surprisingly, the loss of TgLaforin also resulted

in phenotypic effects in tachyzoites, also in a context-specific manner. We thus

established a role for TgLaforin, and by extension reversible glucan phosphorylation,

across both tachyzoite and bradyzoite life stages. These findings build upon previous

studies that increasingly demonstrate a central role for glucan metabolism throughout the

parasite's asexual life cycle.

RESULTS

***T. gondii* tachyzoites contain a cytoplasmic glucan with a punctate distribution.**

Previous studies have presented biochemical evidence for rapid glucan turnover in *T.*

gondii Type I RH tachyzoites (20). Moreover, small granules that stain with periodic acid

Schiff reagent (PAS) have also been noted in the cytoplasm of tachyzoites (20, 21, 23).

Under acid-stress conditions, these tachyzoite glucans have been biochemically

characterized as pure amylopectin, and resemble AGs seen in bradyzoites (18). To

further characterize the nature of this tachyzoite glucan, we used multiple methods to

visualize them under unstressed, normal growth conditions (**Figure 1A**). PAS staining

confirmed that Type II ME49 tachyzoites contain small punctate granules distributed

throughout the cytoplasm. To determine if these PAS-stained granules in unstressed

tachyzoites were more glycogen- or starch-like, they were stained with IV58B6. IV58B6

is an anti-glycogen IgM monoclonal antibody that has previously been demonstrated to

be specific to glycogen (47-49) by recognizing the highly frequent branch-points found in

glycogen (48). Moreover, IV58B6 does not detect other glucans such as amylopectin or

amylose (the primary constituents of plant starch) (50). Tachyzoites stained with IV58B6

in a similar pattern to PAS-stained parasites, containing small punctate granules

distributed throughout the cytoplasm (**Figure 1A**), suggesting that the glucan found in

tachyzoites is more glycogen-like than starch-like. Finally, as is well-known, *T. gondii*

tachyzoites contain almost no visible glucan within their cytoplasm when visualized by

transmission electron microscopy (TEM) (**Figure 1A**), suggesting that the glucan

detected by both PAS staining and IV58B6 is either water-soluble or too small to be

visualized, consistent with this glucan being glycogen-like.

In contrast, bradyzoites have been extensively characterized as containing starch-like

AGs (17-19, 51). After *in vitro* bradyzoite conversion, much of the cytoplasm stained

heavily with PAS (**Figure 1A**). Interestingly, IV58B6 staining intensity appeared to

correlate negatively with *Dolichos biflorus* agglutinin (DBA) staining intensity that defines

the cyst wall, implying that IV58B6 does not stain the PAS-stained glucan in bradyzoites,

further reinforcing the observation that structurally distinct polysaccharides exist in

tachyzoites and bradyzoites (**Figure 1A**). Finally, unlike in tachyzoites, AGs were readily

identified as electron-lucent structures throughout the cytoplasm of in vitro generated

bradyzoites by TEM (**Figure 1A**).

To verify the specificity of PAS and IV58B6 for glucose polymers, tachyzoites and

bradyzoites were treated with acid-α-amyloglucosidase (GAA) after parasite fixation and

before staining. GAA cleaves both α-1,4- and α-1,6-glycosidic bonds and can therefore

completely digest glucans into glucose monomers. Indeed, GAA treatment resulted in the

disappearance of staining within both tachyzoites and in vitro bradyzoites (**Figure 1B**)

demonstrating their specificity for glucose polymers.

TgLaforin colocalizes with the tachyzoite glucan

Because *T. gondii* encodes TgLaforin, a glucan phosphatase that is more animal-like than

plant-like (45, 46), we reasoned that TgLaforin could be involved in the metabolism of the

glycogen-like glucan found in tachyzoites. To determine if TgLaforin co-localizes with the

tachyzoite glucan, endogenous TgLaforin was epitope-tagged with hemagglutinin (HA) in

T. gondii Type II ME49ΔHXGPRT parasites (52) with a CRISPR/Cas9 mediated strategy

(**Figure 2A**) (53). Successful tagging of TgLaforin was confirmed by western blotting

(**Figure 2B**). Immunofluorescence analysis (IFA) of *T. gondii* tachyzoites indicated that

TgLaforin is present in small puncta throughout the cytoplasm, similar to the distribution

of the tachyzoite glucan (**Figure 2C**). Surprisingly, TgLaforin was not detected in *in vitro*

bradyzoites by IFA, 6 days post conversion (**Figure 2C**). To verify that TgLaforin levels

decrease during the tachyzoite to bradyzoite transition, we converted *T. gondii*

tachyzoites to bradyzoites in cell culture using alkaline stress for 6 days and then probed

the converted parasites using western blot analysis. As observed using IFA, TgLaforin-

HA expression decreased dramatically over the course of bradyzoite differentiation

(**Figure 2D**). Transcriptomic data from a previous study obtained from ToxoDB.org

indicates that the transcript levels for TgLaforin do not substantially change over the

course of differentiation, suggesting the possibility that levels of TgLaforin protein are

regulated by post-translational mechanisms (54). To determine if TgLaforin colocalizes

with the glucan present in tachyzoites, we co-stained TgLaforin-HA tachyzoites with either

PAS or IV58B6 along with an anti-HA antibody. In tachyzoites, TgLaforin colocalized with

both PAS (**Figure 2E**) and with IV58B6 (**Figure 2F**), suggesting its involvement in the

metabolism of the tachyzoite glucan.

Initial characterization of TgLaforin-KO tachyzoites

To dissect the role of TgLaforin in *T. gondii* glucan metabolism, TgLaforin was knocked

out using CRISPR/Cas9 to disrupt the gene with a pyrimethamine-resistant form of the

dihydrofolate reductase (DHFR-TS*) under a *Neospora caninum* GRA7 (NcGRA7)

promoter (55) (56) (**Figure 3A**). In agreement with a genome-wide CRISPR KO screen

(57), TgLaforin is a non-essential gene under standard cell culture conditions, as multiple

TgLaforin-KO clones were successfully recovered. Integration of the DHFR-TS* construct

into the TgLaforin locus was verified using inside/out PCR at the chimeric locus and by

verifying the loss of TgLaforin transcription (**Figures 3B,C**). The TgLaforin-KO line further

used in this study (designated "ΔTgLaf") was complemented by the introduction of an

epitope tagged (HA) gene driven by the TgLaforin promoter. The complementation

construct was introduced at an ectopic site in the genome that lacks known coding

sequences or regulatory elements on chromosome VI (**Figure S1A**), while leaving the

ΔTgLaf/DHFR-TS* KO lesion intact for true complementation (58). This complemented

strain, henceforth designated "COMP," was successfully isolated and confirmed by PCR

(**Figure S1B**). Expression levels and localization were similar to those seen in the

TgLaforin-HA line as confirmed by western blotting and IFA (**Figures S1C-D**),

To evaluate effects of a TgLaforin-KO, glucan levels in WT and ΔTgLaf tachyzoites were

first compared using our suite of glucan detection techniques (**Figure 3D**). Surprisingly,

the size and number of PAS-stained granules were not significantly changed in ΔTgLaf

tachyzoites relative to WT parasites. Levels of IV58B6 also remained unaltered after the

loss of TgLaforin, and no aberrant glucan accumulation was observed by TEM as has

been previously reported when genes related to AG or central carbon metabolism were

knocked out in *T. gondii* tachyzoites (20-23, 41, 43, 44) (**Figure 3D**)

Loss of glucan phosphatases in plants and animals results in aberrant glucan

accumulation, and such a phenotype was not observed within tachyzoites under standard

growth conditions.

Loss of TgLaforin results in upregulation of glutaminolysis and glutamine

dependence in tachyzoites

Glucan catabolism is significantly affected by the presence of covalently bound phosphate, and, therefore, loss of glucan phosphatases has profound downstream metabolic impacts in other systems (59, 60). We thus speculated that loss of TgLaforin would result in the reduced efficiency of glucan utilization in tachyzoites and also affect downstream central carbon metabolism. To test this hypothesis, we used gas chromatography/mass spectrometry (GC/MS) steady-state metabolomic analysis of 3 μm filter-purified, syringed-passaged intracellular tachyzoites employing a previously developed sample preparation technique (61).

Previously, it was demonstrated that *T. gondii* tachyzoites primarily utilize glucose and glutamine to drive central carbon metabolism, synthesize macromolecules, and proceed normally through the lytic cycle (62). Glucose primarily fuels glycolysis, and glutamine undergoes glutaminolysis to drive the tricarboxylic acid (TCA) cycle. In the absence of glucose, *T. gondii* can upregulate both glutaminolysis and gluconeogenesis to make up for the loss of glucose (62, 63).

While ΔTgLaf metabolite levels remained unaltered relative to WT tachyzoites across much of the TCA cycle, steady-state levels of metabolites immediately downstream of glutamine were consistently more abundant in ΔTgLaf parasites compared to their WT counterparts (**Figure 4A**), supporting our hypothesis that ΔTgLaf parasites were deficient

in glucan/glucose utilization. An increase in metabolites downstream of glutamine in

ΔTgLaf parasites demonstrates that ΔTgLaf parasites are possibly compensating for

deficiencies in glucose metabolism, supporting a role for the tachyzoite glucan in

intermediate *T. gondii* glucose metabolism.

To determine if loss of TgLaforin resulted in increased dependence on glutamine due to

impaired access to glucose, we performed plaque assays in the presence and absence

of glutamine (**Figure 4B**). In replete media, ΔTgLaf parasites established a similar number

of plaques (data not shown), indicating no defect in infectivity. ΔTgLaf plaques were

slightly larger than both the WT and COMP lines after 10 days of growth (**Figure 4C**). To

test the effects of glutamine starvation on ΔTgLaf parasites, glutamine was removed from

plaque assays after parasite invasion to evaluate the effects of glutamine removal on

parasite growth independent of the initial invasion event. In the absence of glutamine,

ΔTgLaf parasites were unable to form visible plaques, whereas both the WT and COMP

parasites formed plaques comparable to those formed under glutamine-replete conditions

(**Figures 4B, C**).

TgLaforin is required for repeated rounds of progression through the lytic cycle

The absence of differences in plaque number suggest that there is no specific defect in

infectivity. To determine which aspects of the *T. gondii* lytic cycle were impaired in the

absence of glutamine, the effects of glutamine starvation on initial parasite replication and

egress (stimulated with both A23187 (64) and zaprinast (65)) were evaluated. In both

assays, intracellular parasites were pre-starved of glutamine for at least 72 h before assay

initiation. Surprisingly, glutamine starvation had no effect on stimulated egress or initial

parasite replication across the three lines (**Figures S2A-C**). These data demonstrate that

the absence of plaques under glutamine deficient conditions cannot be pinpointed to a

single aspect of the ΔTgLaf lytic cycle, and that the reason for the apparent absence of

plaques manifested later in the infection cycle.

Plaques develop due to repeated cycles of localized infection and cell lysis resulting in the

clearance of infected cells over time. The absence of visible clearance prompted us to

examine infected host-cell monolayers for clusters of infected cells using a higher

magnification than is typically used in a traditional plaque assay. Low numbers of parasites

were seeded onto glass coverslips and fixed at 3- and 6-days post-infection, allowing for

visualization of developing plaques at a high magnification. In these experiments, glutamine-

depleted host cells were pre-starved of glutamine prior to infection with parasites to allow for

potential for invasion defects. Importantly, ΔTgLaf parasites demonstrated similar infectivity to

WT parasites under both glutamine-replate and depleted conditions, indicting no gross initial

invasion defect. After 3 days of growth, no statistical differences of nascent plaque sizes were

noted between glutamine-replete and starved conditions in both the WT and COMP lines.

However, ΔTgLaf parasites in glutamine starved conditions were already 1.5x smaller in area

than their counterparts in replete conditions (**Figure 4D**). By day 6 of growth, this difference

had widened to a >3x difference between glutamine replete/depleted ΔTgLaf parasites (**Figure**

4E). Such a difference was not detected between the two conditions in WT/COMP parasite

lines. By measuring the internal clearing area relative to the total plaque perimeter, it was also

noted that ΔTgLaf parasites were much less capable of forming clearings than the WT/COMP

lines (**Figures S2D, E**), rather they formed clusters of infected cells akin to "turbid plaques"

(66, 67) due to their presumed inability to compete with host cell growth, as the infection

progressed. This observation explains the apparent absence of plaques seen at the lower

magnification used in traditional plaque assays (**Figure 4B**). The modified plaque assay

therefore confirmed that the loss of TgLaforin penalized the summation of repeated rounds of

the energy-demanding lytic cycle rather than one particular aspect of the lytic cycle.

Representative images from this assay can be found in **Figure S2D**.

Loss of TgLaforin results in aberrant bradyzoite AGs in vitro

To determine if loss of TgLaforin resulted in bradyzoite conversion defects, or aberrant

AG accumulation, parasites were converted to bradyzoites *in vitro* using alkaline stress.

During differentiation, the parasitophorous vacuole membrane (PVM), delimiting the

replicative niche established by tachyzoites, converts into the cyst wall that surrounds

bradyzoites within their host cell (68, 69). The cyst wall is heavily glycosylated and

contains N-acetylgalactosamine (Gal-NAc) that is detectible with Dolichos biflorus

agglutinin (DBA) (68). Using DBA-FITC intensity as a marker for differentiation, no penalty

was imposed by the loss of TgLaforin on cyst wall formation over the course of six days

(**Figure 5A**). Somewhat surprisingly, ΔTgLaf mutant parasites tended to exhibit stronger

labeling with DBA at day 6. We additionally assessed the levels of accumulated glucans

using PAS staining (**Figure 5B**). Semi-quantitative analysis of PAS intensity within

vacuoles during stage conversion showed an expected increase over time, but no

significant difference between the WT and ΔTgLaf parasites was detected over the time

course examined.

Because PAS is not specific to glucans and can stain other glucose-containing molecules

such as glycosylated protein and provides no resolution on glucan morphology, we

utilized TEM to gain higher resolution on AG formation during bradyzoite differentiation.

After 6 days of conversion, WT parasites produced AGs that were circular/ovoid and white

(**Figure 5C**). In contrast, ΔTgLaf parasites contained irregular AGs that were

morphologically distinct from AGs that were observed in WT parasites (**Figure 5C**). AGs

in ΔTgLaf parasites appeared amorphous and grayer, while appearing to occupy more

area of the parasite cytoplasm compared to WT parasites. To quantify this phenotype,

the area of AGs was calculated relative to total parasite area to determine the percentage

of the parasite body occupied by AGs in both WT and ΔTgLaf strains (**Figure 5D**).

Strikingly, AGs occupied approximately 4x more relative area in ΔTgLaf parasites when

compared to WT, indicating that PAS staining may lack the sensitivity to capture this

difference, in tissue culture generated bradyzoites. When analyzed on an 8-bit gray scale,

AGs in ΔTgLaf parasites were significantly grayer (grayscale 0-255 is black-white) than

those found in WT parasites, highlighting potential chemical differences (such as

predicted hyperphosphorylation) resulted in differential interactions of ΔTgLaf AGs with

the TEM contrast reagents, likely the heavy metals used in processing (**Figure 5E**).

Examination and quantification of AGs in the COMP line revealed that complementation

of TgLaforin restored most of the circular/ovoid AG cross sections while they also

occupied less space in the cytoplasm and were overall more like those found in WT

parasites (**Figures 5C-E**). Thus, cell culture experiments demonstrate that the loss of

TgLaforin presents itself in both a context and life cycle stage-specific manner.

Loss of TgLaforin results in attenuated virulence and cyst formation in vivo

We hypothesized that loss of TgLaforin may impose a steep penalty under the stresses

and potential nutrient scarcities encountered *in vivo* as it does when nutrients (such as

glutamine) are scarce *in vitro*. To test this hypothesis, equal numbers of male and female

CBA/J mice were infected with 100 tachyzoites intraperitoneally (i.p.) and monitored daily

using a previously developed five-stage body index score to track the severity of

symptoms associated with a tachyzoite infection over the course of 28 days (52).

Mice infected with WT parasites began demonstrating symptoms of infection ten days

after infection with tachyzoites (**Figure 6A**). However, mice infected with ΔTgLaf parasites

did not begin to exhibit symptoms until 15 days after infection. Moreover, mice that

became symptomatic from WT parasite infections often proceeded through all stages of

symptomology, and only a minor proportion of mice that became sick were able to recover

from infection (>70% of mice became moribund or died). Infection from ΔTgLaf parasites,

however, resulted in the majority of mice only developing mild symptoms (Stage 2 or less)

with many of these mice recovering (**Figure 6A**). The attenuated capacity of the ΔTgLaf

parasites to cause symptoms in mice was reflected in the mortality rates of the infected

mice: infection with WT parasites resulted in 73% mortality rate after 28 days whereas

ΔTgLaf parasites only caused 17% mortality (**Figure 6B**). Complementation of TgLaforin

partially rescued this defect in virulence as COMP parasites resulted in an earlier onset

of symptomatic infection at Day 11, and the majority (53%) of mice succumbed to infection

during the first 28 days (**Figures 6A, B**).

Because the acute phase of infection was significantly attenuated by the loss of TgLaforin,

we hypothesized that cyst numbers would be significantly lowered. To determine the

number of cysts formed after 28 days of acute infection (Week 4), we used a previously

established protocol for harvesting and counting tissue cysts from infected mouse brains,

following purification on Percoll gradients (11, 70). Consistent with the ability of ΔTgLaf

parasites to stage convert in culture, mutant parasites were able to establish tissue cysts

in vivo. However, the number of cysts recovered from ΔTgLaf infected animals was lower

than those obtained from WT infected animals (**Figure 6C**). Restoration of TgLaforin in

the COMP line effectively restored tissue cyst yields.

ΔTgLaf tissue cysts can reestablish infections in naïve mice

To determine if the loss of TgLaforin impacted the overall viability/infectivity of *in vivo*

tissue cysts, we examined the disease progression in WT, ΔTgLaf, and COMP infected

animals following injection of 20 tissue cysts i.p. Consistent with prior data (52), infection

with tissue cysts results in markedly lower pathology and consequent mortality during the

acute phase for WT as well as both the ΔTgLaf and COMP lines (**Figure 7A, B**). Mortality

from cyst infections did not differ statistically among the three lines (**Figure 7A, B)**.

Twenty-eight days post-infection, cyst burdens were again enumerated for each line.

ΔTgLaf parasites were once again much less competent at forming cysts *in vivo* (**Figure**

7C). However, unlike the tachyzoite infection, the COMP line was unable to rescue this

defect in cyst formation (**Figure 7C**), suggesting that physiological and metabolic changes

associated with the loss of TgLaforin manifest differently based on the life cycle stage,

impacting their capacity to be complemented.

Loss of TgLaforin results in a delayed starch-excess phenotype within encysted

bradyzoites

We leveraged the inherent fluorescence of PAS coupled with an optimized labeling

protocol to determine the mean intensity of WT, ΔTgLaf and complemented (COMP)

tissue cyst using Image J (**Figure. 8A**). Similar to the pattern observed with in vitro

generated bradyzoites (**Figure. 5**), *in vivo* derived tissue cysts harvested at 28 days

(Week 4) post infection failed to demonstrate a starch-excess phenotype in the ΔTgLaf

mutant (**Figure 8A**). Notable here, is the fact that at week 4 post infection, bradyzoites

are in a state of low AG accumulation (14), and presumably, low turnover, potentially

minimizing the impact of the loss of TgLaf. Our prior data indicate that tissue cysts

harvested at week 6 post infection are in markedly more dynamic state noted by increased

accumulation of AG, higher levels of active mitochondria and increased replicative activity

(14). Quantification of mean PAS intensity exposed a dramatic increase in AG levels

comparing WT and ΔTgLaf tissue cysts, with overall levels between WT and COMP cysts

remaining not significantly different (**Figure 8A**). This points to the effect of the loss of

TgLaf demonstrating a delayed starch excess phenotype, indicating the penetrance of

the mutation is potentially connected to achieving a threshold level of AG as a part of an

emerging AG cycle (14).

The development of AmyloQuant, an imaging-based application, permits the direct

measurement of relative AG levels within tissue cysts (14). With is application, individual

PAS-labeled pixels are classified into 4 bins defining background pixels (black: 0-10

grayscale), low intensity (blue: 10-25), intermediate (green: 25-50) and high (red: > 50

grayscale). Thirty randomly acquired tissue cysts, (the mean intensity of which were

reported in **Figure 8A**) from WT, ΔTgLaf and COMP mice infected for 4 weeks (**Figure**

8B, top row) and 6 weeks (**Figure 8B, bottom row**) are arrayed based on their overall

mean intensity. The percentage of pixels within each of the 4 bins is presented as a

stacked plot, with the AmyloQuant generated thumbnails representing spatial heatmaps

of AG intensity at 5 cyst intervals as presented below each stacked plot. This analysis

reveals that despite the mean PAS intensities not being statistically significant (**Figure**

8A), several ΔTgLaf cysts present with varying levels of high intensity (red) PAS labeling.

Notable here is that with this cohort of cysts, effective complementation of AG levels is

not observed.

As expected for wild type Week 6 tissue cysts (14), markedly increased AG

accumulation is evident compared to Week 4 cysts (**Figure 8B**). This phenotype is grossly

exaggerated in the case of ΔTgLaf tissue cysts (**Figure 8B**), displaying the expected

starch excess phenotype. Even so, not all ΔTgLaf cysts present uniformly high levels AG

based on PAS intensity (**Figure 8B**). Notable here, the levels and distribution of AG levels

within the COMP line appear to be intermediate to that observed for the WT and ΔTgLaf,

pointing to the plasticity of the phenotype.

Several recent studies targeting activities connected with AG metabolism report

differences in cyst yield as well as cyst size (21-23, 41). We therefore examined if the

loss of TgLaf resulted in any significant impact of cyst size. Contrary to other reports

however no differences in cyst diameter were noted for either week 4 or week 6 harvested

tissue cysts (**Figure S3A**). Furthermore, we found no strong correlation between cyst

size and mean AG levels as defined by PAS labeling intensity (**Figure S3B**).

Aberrant AG accumulation in ∆TgLaf tissue cysts promotes bradyzoite death.

The inherent harshness of PAS labeling causes differential labeling intensity related

issues with regard to the labeling of bradyzoite nuclei (14) as well as labeling for TgIMC3

a marker of recent replication (14). As a result, the relationship between AG levels and

replication associated outputs (packing density and recency of replication) could not be

reliably implemented in the context of the TgLaf mutant. We therefore resorted to the

direct examination of purified tissue cysts by transmission electron microscopy.

We adapted a protocol designed to capture and image low abundance cells by TEM by

making it compatible with our tissue cyst purification protocol (see Methods) (71, 72).

TEM imaging revealed that while WT parasites formed largely normal/canonical AGs *in*

vivo as seen *in vitro* (**Figure 9** [compare with **Figure 5A**])*,* ∆TgLaf parasites contained

almost exclusively aberrant AGs that mirrored the same morphological defects seen *in*

vitro (**Figure 9** and **Figure S4**). ∆TgLaf AGs were irregularly sharpened with a flat, multi-

lobed appearance. Importantly, COMP parasites neither over-accumulated nor formed

aberrant AGs, demonstrating that this defect is specific to loss of TgLaforin (**Figure 9**).

In addition to containing aberrant AGs, the internal morphology of ∆TgLaf parasites

appeared to be altered by the presence of the AGs, as significant organelle displacement

was also noted (**Figure 9, S4**). Many of the ∆TgLaf parasites also appeared hollow or

"ghost-like" in appearance resulting from their unstained/absent cytoplasm, which

included the apparent loss of the nucleus in several bradyzoites (**Figure 9, S4**). These

data suggest that a significant number of the ∆TgLaf bradyzoites were inviable within the

cyst (**Figure 9, S4**). Together, these direct and selective impacts of TgLaforin's loss on

both bradyzoite viability and growth *in vivo* establish TgLaforin as a potentially druggable

target.

DISCUSSION

The asexual life cycle of *Toxoplasma gondii* is defined by two fundamentally distinct

forms: the rapidly replicating tachyzoite and the slowly growing bradyzoite. These forms

represent distinct physiological states that can be further subdivided, particularly within

encysted bradyzoites (73). Insights into these physiological states, inferred from

transcriptomic analyses, are reinforced in emerging metabolomic studies (74, 75). In

these studies, glucose and glutamine, which are both linked to energetics, biosynthesis,

and intermediary metabolism, appear as critical metabolites. Importantly, glucose can be

stored in polymers like glycogen and amylopectin during times of low energy need. Stored

glucose can be present in an accessible and labile form for rapid mobilization such as

glycogen, or in a less soluble forms as AGs from which it can be accessed more slowly

(76, 77). Reversible glucan phosphorylation facilitates the breakdown of such glucan

polymers by disrupting the crystalline helices on the glucan surface (78). *T. gondii*

encodes the capacity for reversible glucan phosphorylation (27, 36, 45, 46). The

contribution of this process to tachyzoite and bradyzoite biology was evaluated through

targeted manipulation of the glucan phosphatase, TgLaforin (36).

The accumulation of AGs within bradyzoites and their apparent absence in tachyzoites

has been used as discriminator between these life cycle stages (16, 79). Detailed

examination, however, presents a considerably more nuanced picture, alongside

emerging evidence that points to rapid glucan turnover within tachyzoites (20). In the

current study, using an IgM monoclonal antibody (IV58B6) that specifically recognizes

glycogen-like glucose polymers (47, 48), we demonstrate that the stored glucan within

tachyzoites is structurally closer to animal glycogen (47, 48) than the insoluble plant-like

AG granules found in bradyzoites (17, 19). This duality between life stages may be

additionally reflected in the observation that TgLaforin, the glucan phosphatase, and

TgGWD (TGME49_214260), the partner kinase that is predicted to phosphorylate *T.*

gondii glucan polymers, trace their structural lineages to animals and plants, respectively

(45, 46). The glycogen-like glucan polymer appears to be specific to tachyzoites as its

levels decrease upon *in vitro* differentiation while the overall PAS intensity increases

(**Figure 1A**). This suggests that the tachyzoite glucan and bradyzoite AG are

architecturally distinct polymers with respect to both branching frequency, solubility, and

phosphorylation status (**Figure 2D**). The glucose-based nature of both particles is

supported by the elimination of both IV58B6 and PAS staining with α-amyloglucosidase

treatment (**Figure 1B**).

To address the contribution of stored glucans in both tachyzoites and bradyzoites, we disrupted the glucan phosphatase TgLaforin. This enzyme preferentially removes phosphate groups from the C3 carbon on glucose facilitating access to enzymes that release glucose (36). The loss of other glucan phosphatases such as SEX4 in *A. thaliana* and laforin in mammals is accompanied by excessive accumulation of aberrant starch and hyperphosphorylated glycogen in plants and animals (37, 39, 40, 80, 81). Surprisingly, given these penalties in other systems, ΔTgLaf tachyzoites exhibited no gross morphological changes in glucan content, consistent with a recent study in which TgGWD was knocked out (27). These observations contrasts with other KO studies of glucan pathway associated proteins such as CDPK2 (20), glycogen phosphorylase (41), the PP2A holoenzyme (44), and α-amylase (22)which all reported exaggerated glucan accumulation in tachyzoites.

Despite the absence of glucan accumulation in ΔTgLaf tachyzoites, the loss of this gene exerts an effect on tachyzoite glucan metabolism. Consistent with the metabolic defects associated with the loss of laforin in humans (60), loss of TgLaforin in *T. gondii* resulted in altered central carbon metabolism that manifested as ΔTgLaf parasites' dependence on glutamine (**Figure 4, S2D,E**). ΔTgLaf tachyzoite dependence on glutamine supports recent studies demonstrating that tachyzoites utilize storage glucans for glucose allocation (23) because the presumed loss of efficient glucan degradation results in tachyzoite dependence on glutamine (**Figures 4B,C**). As *T. gondii* tachyzoites primarily utilize glucose and glutamine to support their rapid growth, this dependence on glutamine reinforces previous observations that glutamine can substitute for glucose in this altered

metabolic landscape (62, 82). These results are consistent with many previous studies

that disrupt glucose and/or glucan metabolism, but contrast with others. Disruption of

TgGT1 (TGME49_214320), the only plasma-membrane glucose transporter in *T. gondii*

(62, 83), or TgHK (TGME49_265450), the *T. gondii* hexokinase, resulted in upregulation

of gluconeogenesis, and parasite growth was highly attenuated with glutamine depletion

(82). Most strikingly, parasites lacking starch synthase (TgSS; TGME49_222800)

displayed no dependence on glutamine and, in fact, grew faster than WT parasites when

both glucose and glutamine were removed from the culture media (23). Interestingly,

however loss of TgSS did result in lower glucose flux through glycolysis (23), consistent

with our findings that demonstrate a role for the tachyzoite glucan in glucose allocation.

This finding may suggest that while the absence of AG in the ΔTgSS parasites may not

be detrimental, overaccumulation as observed in ΔTgLaf cysts can contribute to toxicity

(**Figure 9**). Perturbations of other glycolytic enzymes also demonstrated varied effects

related to the presence of glutamine: loss of the glycolytic enzyme glyceraldehyde-3-

phospahte dehydrogenase (GAPDH1) could be rescued with high levels of glutamine (84)

(84), but glutamine could not rescue pyruvate kinase (TgPYK1) knockdown parasites

(85). Our data indicate that loss of access to key nutrients such as glucose and glutamine

has a profound impact on the repeated rounds of the lytic cycle without being attributable

to one specific process within the cycle, suggesting that the penetrance of the phenotypic

defect manifests cumulatively over time, rather than being hard wired in each infection

cycle (**Figures 4D,E, S2D,E**).

Despite glucan metabolism being historically viewed as being important in the chronic

infection, TgLaforin protein expression decreased during the tachyzoite to bradyzoite

conversion *in vitro* even though its transcript levels do not change (**Figures 2C,D**). This

could be a transient observation as the downregulation of glucan catabolism during

conversion would facilitate accumulation of AGs for the chronic infection. We therefore

examined how the loss of TgLaforin affected the capacity of ΔTgLaf parasites to

differentiate *in vitro*.   The ΔTgLaf parasites exhibited no defect in AG-accumulation

kinetics, detected by PAS staining, or in cyst wall formation, detected with DBA lectin over

the course of the in vitro conversion assay (**Figure 5A, B**). The lack of difference in PAS

labeling between both the WT and the ΔTgLaf lines, however, did not reveal the

differences noted by TEM (**Figure 5C-E**). As initially hypothesized would be the case in

both tachyzoites and bradyzoites, loss of TgLaforin resulted in aberrant AG accumulation

within *in vitro* bradyzoites that is marked by changes in both level and morphology

(**Figures 5C-E**), as seen in plants and vertebrates (37-40, 81).   AGs in the ΔTgLaf

parasites were not only present at higher levels but were potentially chemically distinct

considering their differential binding to TEM contrast metals (**Figure 5E**). Given that

TgLaforin is a confirmed glucan phosphatase (36), we speculate that AG

hyperphosphorylation may account for both altered morphology and appearance by TEM.

These context-specific phenotypes suggested that the ΔTgLaf mutant would manifest

phenotypic differences in both the acute and chronic phases on infection *in vivo*. Indeed,

the loss of TgLaforin was associated with a markedly reduced symptomology and

associated mortality compared to both the parental and complemented parasites during

acute tachyzoite-initiated infection (**Figures 6A,B**). Not only was there a delay in

symptomatic disease, but also a reduction in disease severity and overall cyst burden.

Symptomology in the acute infection is driven by an increasing parasite burden driving an

overexuberant host inflammatory response (4, 86). The delayed symptom onset suggests

growth inhibition by the stringent *in vivo* environment that more effectively controls ∆TgLaf

parasite infection with less robust inflammation. Notably, the delayed and milder course

of the tachyzoite infection resulted in a lower overall cyst burden in surviving animals

compared to infection with both WT and COMP parasites (**Figure 6C**).

Infection with ∆TgLaf tachyzoites resulted in fewer tissue cysts being generated relative

to WT and COMP parasite (**Figure 6C**) The basis for this is not clear, but suggestive of

a regulatory imbalance that is masked in tachyzoites but evident in tissue cyst-initiated

infections which have the additional burden of converting from bradyzoites to tachyzoites,

surviving the chronic infection before forming new tissue cysts. This could potentially

parallel the observations with plaque formation in vitro (**Figure 4, S2**).

Our ability to quantify and map relative AG levels within tissue cysts using AmyloQuant

(14), confirm that the loss of TgLaf does not fundamentally alter the initiation and early

progression of an AG cycle as evidenced by the effect of the mutation not being evident

in Week 4 tissue cyst in the form of the expected starch excess phenotype (**Figure 8**).

The effect of the ∆TgLaf mutation becomes evident in week 6 tissue cysts where a

massive increase in AG accumulation (**Figure 8**) is likely due to an imbalance caused by

the predicted defect in AG turnover. As with other phenotypes associated with this

mutation, the effect continues to exhibit phenotypic variation noted by roughly a third of

the imaged cysts lacking a significant proportion of high intensity pixels (red), while others

are completely oversaturated (**Figure 8**). These distinct populations are not reflected in

any way based on the size of the cyst (**Figure S3**). In fact, contrary to other reports

regarding AG metabolism associated genes, the TgLaf mutation has no significant effect

on tissue cyst size (**Figure S3**).

While we were not able to accurately quantify either the nuclear number or TgIMC3

intensity distributions as markers of packing density and replicative activity in PAS stained

tissue cysts (14), TEM analysis exposed the true consequence of the loss of TgLaf within

tissue cysts in vivo (**Figure 9, S4**). Wild type tissue cysts presented with clusters of small

amylopectin granules as well as evidence of active endodyogeny (**Figure 9**). In contrast

ΔTgLaf cysts presented bradyzoites laden with AG to the point where other organellar

structures were obscured (**Figure 9, S4**). Several bradyzoites lacked nuclei while others

lacked the cytoplasm (**Figure 9, S4**). Together, these features are incompatible with

viability. The presence of such features does not apply to all bradyzoites within the cyst

accounting for the fact that these cysts are able to initiate a new infection (**Figure 7**.),

albeit with lower virulence and cystogenic potential. These finding suggest that AG

metabolism is under tight control as dysregulated accumulation can result in cumulative

defects resulting in toxicity and death. The high frequency of these abnormal parasites

suggests that reversible glucan phosphorylation and TgLaforin specifically represent

legitimate bradyzoite specific drug targets. We recently described a small molecule that

inhibits recombinant TgLaforin (36) which serves as a potential starting point in the

development of a new class of anti-*Toxoplasma* therapeutic agents. Particularly exciting

in this context is the fact that a class of drugs exhibiting efficacy with tissue cyst clearance

(atovaquone (87, 88) endochin-like quinolones (89-91) and JAG21 (92)), all target

mitochondrial respiration. When glucose is limiting, mitochondrial respiration can be

driven by glutamine. This provides an opportunity for combination therapy to promote the

clearance of toxoplasma tissue cysts as a means of mitigating the risk of reactivation.

METHODS

Fibroblast and parasite culture and maintenance

All parasite lines were maintained in human foreskin fibroblasts (HFFs; ATCC) in Minimal

Essential Media-α (MEM-α; Gibco) supplemented with 7% heat-inactivated fetal bovine

serum (FBS; Gemini Bio), 100 U/mL penicillin, 100 μg/mL streptomycin, and an additional

2 mM L-glutamine (Gibco; 4 mM total L-glutamine). Cells and parasites were incubated

at 37°C and 5% CO_2 in a humidified incubator. Genetically modified parasites were

maintained in MEM-α containing 7% dialyzed FBS (Gemini Bio) and either pyrimethamine

(1 μM), mycophenolic acid/xanthine (MPA: 25 μg/mL, xanthine: 50 μg/mL), or 6-

thioxanthine (6-Tx: 80 μg/mL).

Assays analyzing the effects of glutamine deprivation used Dulbecco's Modified Eagle

Medium (DMEM). Both glutamine-replete (Gibco, 11966025) and depleted (Gibco,

11054020) DMEM were supplemented with 7% dialyzed FBS. Glutamine-replete media

from the supplier lacked other key nutrients and was modified to contain 5 mM glucose,

1 mM sodium pyruvate, and 4 mM L-glutamine.

Generation of T. gondii mutant lines

Type II ME49ΔHXGPRT ("WT"—the parental line utilized to generate all other lines in this study): This line was generated in a previous study using CRISPR/Cas9 targeting of TgHXGPRT and selection with 6-Thioxanthine (52).

TgLaforin-3xHA-HXGPRT: TgLaforin was epitope tagged with HA at the C-terminus using CRISPR-Cas9 to disrupt the TgLaforin 3'UTR immediately downstream of the endogenous stop codon as has been previously described (53). Briefly, a sgRNA immediately downstream of the TgLaforin stop codon was designed using the EuPaGDT design tool (http://grna.ctegd.uga.edu). The top hit was selected (**Table S1**) and used to replace the sgRNA sequence in pSAG1::CAS9-U6::sgUPRT, a plasmid containing both Cas9-green fluorescent protein (GFP) and an interchangeable sgRNA scaffold (56); (**Table S2**). Replacement of the interchangeable sgRNA was accomplished using a Q5 site-directed mutagenesis kit (**Table S3**) (New England BioLabs). The TgLaforin-HA tagging construct was generated by amplifying the 3' end of the TgLaforin-HA construct generated for complementation (see generation of COMP line below and **Tables S2 and S3**) along with the connected HXGPRT selectable marker. Both the TgLaforin-HA PCR-amplicon and the CRISPR-Cas9-GFP were transfected into 1.4×10^7 *T. gondii* ME49ΔHXGPRT parasites (2:1 insert:plasmid molar ratio; 30 μg DNA total) by electroporation with a time constant between 0.16 and 0.20 msec (BioRad Gene Pulser II). After 24 h, surviving parasites were syringe-passaged from infected HFFs with a 27 G needle to lyse host cells, and gravity-filtered through a 10 μm filter to remove host-cell debris. Successful transformants were then enriched by use of fluorescence-activated

cell sorting (FACS; Sony SY3200, installed in a biosafety level II cabinet) to select

parasites expressing Cas9-GFP from the transfected plasmid by isolating GFP+

parasites. HFFs were infected with GFP+ parasites, and then placed in media containing

MPA/xanthine 24 h later to select for restoration of HXGPRT. MPA/xanthine-resistant

parasites were cloned by limiting dilution into a 96 well plate. Wells containing single

plaques were picked 7 days later and expanded. Genomic DNA was extracted from

clones using a Proteinase K treatment detailed elsewhere (93). Successful tagging of

TgLaforin was verified using sequencing, immunoblotting, and IFA.

ME49ΔHXΔTgLaforin ("ΔTgLaf"): TgLaforin was disrupted using a CRISPR-Cas9

mediated strategy as detailed above, with several differences. Briefly, a single sgRNA

was designed to target the first exon of TgLaforin with the top hit from EuPaGDT (**Table**

S1). To disrupt TgLaforin with a selectable drug marker, DHFR-TS*, a pyrimethamine-

resistant mutant of the DHFR gene, containing a 5'-NcGra7 promotor and DHFR 3'UTR

was amplified from pJET-NcGra7_DHFR (**Table S2**). Amplification utilized primers

containing 40 nt extensions homologous to the 5'- and 3'-UTR of TgLaforin to encourage

homologous recombination-mediated whole-gene replacement with the drug cassette

(**Table S3**). Both the PCR-amplified DHFR* homology cassette and the CRISPR-Cas-

GFP plasmid were transfected and FACS-sorted as described above. GFP+ parasites

underwent drug selection in pyrimethamine. Parasites were then cloned and expanded

as detailed above. Successful integration of the DHFR* cassette into the TgLaforin locus

was verified using PCR with inside/out primer pairs to the chimeric, interrupted gene

(**Table S3**). Loss of TgLaforin transcription was verified by purifying RNA from TgLaforin

clones on RNeasy spin columns (Qiagen). Using the Promega Reverse Transcriptase

System, cDNA was synthesized from RNA extracts. Primers designed for full-length

TgLaforin amplification were then used to verify loss of TgLaforin cDNA in knockout lines.

ME49ΔHXΔTgLaforin+ChrVI-TgLaforin ("COMP"): Complementation of TgLaforin was

also executed using a CRISPR-mediated strategy. A sgRNA to a neutral locus on

chromosome VI identified previously (58) was generated using the same mutagenesis

strategy as above (**Table S1 and S3**). A full length TgLaforin cDNA containing its

endogenous 5'UTR (2000 bp upstream from gDNA) was synthesized by GenScript and

inserted into a pHA3x-LIC vector (**Table S2**) containing a C-terminal HA tag and a DHFR

3'UTR, linked to the HXGPRT selectable marker (named "TgLaforin-HA3x-LIC"; also

used above for endogenous tagging to create the TgLaforin-HA line). The entire construct

(5'UTR:TgLaforin-cDNA:DHFR-3'UTR:HXGPRT) was amplified from the vector and co-

transfected into ΔTgLaf parasites with the CRISPR-Cas9 plasmid as done above.

Successful transformants that received the HXGPRT marker were selected with

MPA/xanthine. Successful insertion of TgLaforin along with its promoter was verified

using PCR (**Table S3**), immunoblotting, and IFA with an anti-HA antibody (Abcam).

Immunofluorescence (IF) staining

HFFs were grown on glass coverslips until confluent and subsequently infected. Infected

HFFs were fixed with either methanol (MeOH) (100%, -20°C) or methanol-free

paraformaldehyde (PFA) (4% in phosphate-buffered saline (PBS); Electron Microscopy

Sciences) as indicated below for each antibody. Infected HFFs fixed with PFA were

permeabilized in 0.1% TritonX-100 in PBS++ (PBS containing 0.5 mM $CaCl_2$ and 0.5 mM

MgCl$_2$) for 10 min at room temperature (RT). Primary and secondary antibodies were

diluted in 3% (w/v) bovine serum albumin (BSA; Fisher) in PBS++. Samples were first

incubated with the primary antibody (αHA-1:1,000; αSAG-1:10,000; αGAP45-1:5,000;

αGRA3-1:1500; IV58B6-1:50) at RT for 45 min, washed 3x with PBS++, and then

incubated with fluorescent secondary antibodies (1:2,000) and 4',6-diamidino-2-

phenylindole (DAPI; 300 nM) for 45 min. Secondary antibodies (Invitrogen) were

conjugated to either Oregon Green or Texas Red fluorophores and specific to the species

and class of primary antibody used. Samples were then washed 3x with PBS++ before

mounting the coverslip on a glass slide using MOWIOL mounting media.

Immunofluorescence staining was visualized using a Zeiss AxioVision upright microscope

with a 100X 1.4 numerical-aperture oil immersion objective, and images were acquired

using a grayscale Zeiss AxioCam MRM digital camera. Grayscale images were pseudo-

colored in ImageJ using magenta (Texas Red), yellow (Oregon Green), and cyan (DAPI),

and further alterations to brightness and contrast were also made in ImageJ when

deemed appropriate. For all assays in which staining intensity was compared across

treatments and parasite lines, concentrations of antibodies, exposure times, and

alterations to brightness/contrast were identical. Colocalization of fluorescent

antibodies/reagents was quantified using Pearson's coefficient calculated with the JACoP

plugin on ImageJ (94).

***PAS staining*-tachyzoite and in vitro bradyzoite**

PAS staining of tachyzoites and in vitro bradyzoites was done on infected HFFs fixed in

4% PFA and permeabilized as above. Coverslips were then washed 3x in tap water before

the addition of 1% periodic acid (Sigma-Aldrich) for 5 min. Coverslips were then washed

with three changes of tap water. Schiff's reagent (diluted 1:4 in tap water) was added for

15 min. Coverslips were subsequently washed 10x with tap water to develop stain before

being incubated with DAPI for 10 min and then mounted as above. PAS-stained samples

were visualized using fluorescence microscopy (excitation: 545 nm, emission: 605 nm).

When PAS was co-stained with antibodies, primary antibodies were incubated with PAS-

stained slides overnight in BSA at 4 °C before standard secondary staining.

PAS labeling of methanol fixed tissue cysts was performed as described elsewhere, using

Schiff reagent diluted 1:10 in tap water as described elsewhere ().

Samples treated with acid-α-amyloglucosidase (GAA) (from *Aspergillus niger*, >260

U/mL, Sigma) were incubated with GAA after permeabilization. GAA was diluted 1:50 in

50 mM sodium phthalate buffer, pH 5.5, and samples were treated for 24 h at room

temperature. Untreated controls were incubated in phthalate buffer without GAA.

Samples were then stained with PAS or IV58B6 as described in the IF-staining workflow

above.

In vitro bradyzoite conversion assay

Tachyzoites were converted to bradyzoites *in vitro* using alkaline stress as has been done

previously with several modifications (95). HFFs grown were infected with tachyzoites in

standard cell culture media. 4 h later, media was replaced with RPMI 1640 (Gibco

31800022) supplemented with 50 mM HEPES and adjusted to pH 8.2 with NaOH.

Parasites were then cultured for 2-6 days at 37°C, ambient CO_2, and sealed in Parafilm.

Media was replaced every other day to maintain the basic pH. Parasites were fixed in

PFA and stained with fluorescein conjugated *Dolichos biflorus* agglutinin (DBA; 1:1000,

Vector Laboratories) and PAS. Images were obtained in grayscale on a Zeiss AxioVision

upright microscope as described above. To determine the degree of labeling with DBA

or PAS, the Fiji/ImageJ (96) was used to create a binary mask outlining cysts that was

applied to the PAS-stained image to measure the greyscale intensity of each ROI (i.e.

each individual vacuole/*in vitro* cyst).

Transmission electron microscopy of in vitro tachyzoites and bradyzoites

Transmission-electron microscopy (TEM) was performed as done previously (97). Blocks

were stained at the University of Kentucky's Imaging Center in the College of Arts and

Sciences. Blocks were trimmed and sectioned on an ultramicrotome with a diamond knife.

Sections were placed on copper grids and then contrast stained with lead citrate.

Micrographs were collected at the University of Kentucky's Electron Microscopy Center

on a Talos F200X TEM (Thermo) operated at 200 kV accelerating voltage with a 50 μm

objective aperture inserted to enhance contrast using a 16M pixel 4k x 4k CMOS camera

(Ceta, Thermo Scientific). AG size and grayscale values were measured in ImageJ.

Immunoblotting

Parasites were syringe lysed from host cells, pelleted, and $2x10^6$ parasites were

resuspended in SDS-PAGE sample buffer and boiled for 10 min before being run on a

single lane of a 10% polyacrylamide gel. The gel was then transferred to a 0.2 μm PVDF

membrane (BioRad) using a Turbotransfer System (BioRad) for 7 min at 25 V. The PVDF

membrane was blocked in 5% (w/v) non-fat milk in Tris-buffered saline plus Tween-20

detergent (TBST; 0.1% Tween-20) for 20 min before being probed with a primary antibody

(αHA-1:1,000; αGAP45-1:5,000; αSRS9-1:1,000; αSAG1-1:10,000) in non-fat milk

overnight at 4^{o}C (Cell Signaling C29F4). The blot was washed 3x with TBST before

probing with either HRP-conjugated α-rabbit or α-mouse-IgG (Jackson Laboratories). Blot

was washed and developed for 5 min using SuperSignal^{TM} West Pico PLUS (Thermo

Scientific) and visualized on a GelDoc station (BioRad).

Steady state polar metabolite analysis

Parasites were prepared as previously described (61). Confluent HFFs were infected with

parasites at a multiplicity of infection (MOI) of 2 to achieve a high density of parasites

after 48 h of growth (>80% cells containing >32 parasites each). Plates containing

infected HFFs were placed on ice, media removed, and the monolayer was washed 2X

with ice-cold PBS. Parasites were harvested on ice in a 4 ^{o}C cold-room. Cells were

scraped from plate surface, resuspended in PBS (8 plates/50 mL PBS), and centrifuged

at 1000g for 10 min at 4 ^{o}C. PBS was removed, the cell pellet was resuspended in 2 mL

PBS, and syringe passaged successively in 23 G and 27 G needles. The soluble host cell

lysate was removed by centrifugation (1000g). The pellet was resuspended in 5 mL PBS

and host-cell debris was removed by syringe-filtering the suspension through a 3 μm filter

(Whatman). Filtered parasites were then pelleted, resuspended in 1 mL PBS, and

counted on a hemacytometer. Parasites were pelleted a final time at 14,000g for 30 s at

4 °C, supernatant was removed, and pelleted parasites were flash frozen in liquid nitrogen

and stored at -80 °C until metabolite extraction.

Polar metabolite extraction: Polar metabolites were extracted in 0.5 mL -20 °C 50%

methanol (MeOH) containing 20 μM L-norvaline (procedural control) for 30 min on ice.

During the 30 min incubation, samples were regularly vortexed. Samples were then

centrifuged at 14000*g* for 10 min to pellet insoluble material (protein, DNA, RNA, and

glycans). Supernatant containing polar metabolites and pellet were dried separately on a

SpeedVac (Thermo) at 10^{-3} mBar until methanol (MeOH) was completely sublimated and

only dried pellet remained.

Pellet hydrolysis and extraction: Dried fraction containing protein was hydrolyzed by

resuspending the pellet in 2 N HCl (final concentration) at 95°C for 2 h. Hydrolysis was

quenched, and hydrolyzed amino acids were extracted by the addition of an equal volume

of 100% MeOH with 40 μM L-norvaline such that the final concentration was 50% and 20

μM, respectively. Extraction and drying then proceeded as described above.

Sample derivatization: Dried samples (both polar metabolites and hydrolyzed protein)

were derivatized in 70 μL 20 mg/mL methoxyamine hydrochloride in pyridine for 90 min

at 30 °C. Samples were then centrifuged at 14000*g* for 10 min to remove any particulate,

and 50 μL of the methoxyamine supernatant was mixed with 80 μL *N*-methyl-*N*-

trimethylsilyl trifluoroacetamide (MSTFA) and incubated for 30 min at 37°C. Samples

were then transferred to amber glass chromatography vials and analyzed by GC/MS.

GC/MS analysis: Metabolites were analyzed on an Agilent 7800B GC coupled to a 5977B

MS detector using a previously established protocol (98). Automated Mass Spectral

Deconvolution and Identification System (AMDIS) was used to analyze metabolites by

matching metabolites to the FiehnLib metabolomics library via retention time and

fragmentation pattern. Quantification of metabolite levels was performed in Mnova.

Sample abundance was normalized to L-norvaline (procedural control) and protein from

the protein pellet (experimental control). Steady state metabolites are presented as the

mean of three independent replicates.

Conventional plaque assays

HFFs were grown in 12-well plates until confluent. HFFs were subsequently infected with

200 parasites/well under standard cell culture conditions. Wells were washed with PBS

to remove residual invasion media, and media was changed to glutamine replete or

depleted media 4 h post-infection to allow for invasion. Plates remained undisturbed for

10 days before the infected HFFs were fixed with 100%, -20°C MeOH for 20 min, stained

with 1% crystal violet solution for 20 min, and then the plaques were de-stained with

repeated tap water washes. Zones of lysis (white clearings) could be visualized against

intact cells (purple). Images of plaques were obtained by scanning plates on an Epson

Perfection V600 photo scanner at a resolution of 600 dpi. The plaques were measured

by pixel area using ImageJ. Plaque assays were conducted on three independent

replicates, and plaque size from these experiments were aggregated to highlight

variability in plaque sizes.

Modified plaque assays

A modified plaque assay was developed to visualize foci of infection at higher

magnification where zones of clearing were not readily evident. Confluent HFF

monolayers on glass coverslips in 24 well plates were infected with WT, ΔTgLaf, and

COMP parasites in replete media for 4 hours to allow for invasion. The monolayers were

washed gently 3 times with PBS and either fresh replete media or glutamine-depleted

media were added for 3 or 6 days to appropriate wells. Infected monolayers were washed

and fixed with MeOH (-20 °C) and subjected to IF using DAPI and GRA3, an antibody

that detects the PVM. Individual plaques were imaged on coded blinded slides using a

10X objective, and their perimeters and encompassed areas were measured using Image

J. Host cell clearance within plaques was similarly measured, and the extent of clearance

was represented as a percentage for each individual plaque. A total of three independent

replicates were performed.

Egress assays

HFFs were grown to confluency in 35 mm glass bottom dishes (MatTek, P35G-0-14-C).

Two days before infecting HFFs on glass bottom dishes, both HFFs and parasites were

independently pre-treated in either glutamine-replete or -depleted media (see above for

media formulations). After 48 h pre-treatment, 10^5 parasites of each line (WT, ΔTgLaf,

and COMP) in each pre-treatment (gln+/-) were added to fresh dishes and allowed to

grow for 48 h so that most vacuoles contained >32 parasites each. Several hours before

egress, media in each infected plate was adjusted to 1.5 mL and allowed to equilibrate at

37°C in 5% CO_2. The calcium ionophore A23187 (Cayman Chemical Company) was

prepared as a 2 mM stock in DMSO and diluted in (+/-) gln media to make a 4X

concentration of 12 μM and maintained at 37°C throughout the assay. Zaprinast was

likewise prepared as a 100 mM stock in DMSO and diluted into media at a 4X

concentration of 2 mM. Egress was triggered by the addition of 0.5 mL 4X A23187 to

infected HFFs (3.0 μM final concentration) or 0.5 mL 4X-zaprinast (500 μM final

concentration). Egress was monitored on a Nikon Eclipse Ti2 inverted microscope with a

40X phase air objective modified with a 1.5X optivar. Several fields containing vacuoles

were selected from each plate, and an image was obtained 10 s after triggering egress

from each field once every 5 s for 5 mins (61 images/field) on a Nikon DS-Ri2 color

camera. Videos of each field were assembled on NIS Elements software. Egress was

monitored using standard deviation of pixel intensity and determined by inflection point of

change in standard deviation of pixel intensity. Inflection point was calculated by fitting a

gaussian curve to the first derivative of the standard deviation in pixel intensity and

calculating the mean of the curve. Technical replicates (fields on each plate) were

averaged for each biological replicate (average of fields from each plate).

Replication assays

HFFs were grown on glass coverslips in a 24-well plate until confluent. Two days before

infecting HFFs on coverslips, both HFFs and parasites were independently pre-treated in

either glutamine-replete or -depleted media. After pre-treatment, 10^4 parasites of each

line (WT, ΔTgLaf, and COMP) in each condition (gln+/-) were added to 3 coverslips each.

24 hours later, infected HFFs were fixed in MeOH and stained with Rb-α-SAG1 (1:10,000)

and DAPI for ease of visualization. Counting of parasites/vacuole was performed for each

line/condition on coded blinded slides with the identity of samples revealed upon

completion of the counting. A total of three independent replicates were performed.

Mouse infection studies

4- to 6-week old CBA/J mice of both sexes (Jackson Laboratories, Bar Harbor, ME) were

injected intraperitoneally (i.p.) with either 100 WT, ΔTgLaf, or COMP tachyzoites, or with

20 tissue-cysts from brain homogenates derived from previously infected mice. In either

case, parasites/cysts were suspended in a final volume of 0.2 mL serum-free, Opti-MEM

media (Gibco). Mice were then monitored and assigned a body index score daily.

Monitoring frequency increased to twice a day once symptomatic throughout the course

of infection as previously described (52). When symptomatic, mice were administered a

gel diet and wet chow on the cage floor and given 0.25-0.5 mL saline solution

subcutaneously as needed. Moribund mice were humanely euthanized. Euthanasia of

both moribund mice and mice sacrificed at the time of tissue cyst harvest was performed

by CO_2 asphyxiation, followed by cervical dislocation. The number of mice used for each

experiment is indicated within relevant figures. All protocols were carried out under the

approval of the University of Kentucky's Institutional Animal Care and Use Committee

(IACUC).

Tissue cyst purification

Tissue cysts were purified as previously described using discontinuous Percoll gradients

(11, 70). Processing of two sex/infection-matched brains was performed on each

gradient. Cysts were collected in 1 mL fractions from the bottom of the centrifuged Percoll

gradient using a peristaltic pump adjusted to a flow rate of 2 mL/min. To quantify tissue

cysts, 10-20 μL of each fraction was placed into 100 μL PBS in the well of a 96-well plate,

pelleted and directly enumerated at 20X magnification in each well. Total cysts per mouse

were calculated by summing the total number of cysts in each fraction and dividing the

total by two to adjust for brain homogenization in pairs. Each pair of mice was presented

as a single averaged data point. Tissue cysts were pelleted onto slides using a Cytospin

centrifuge and fixed and stored in 100% MeOH (-20°C) until staining.

Quantification of AG levels in tissue cysts based on PAS labeling intensity.

Tissue cysts deposited on glass slides were fixed and stored in methanol at -20°C. Slides

were equilibrated to room temperature and stained with PAS using conditions optimized

for staining tissue cysts (14).   Cysts were additionally stained with DBA lectin to

demarcate the cyst boundary. Thirty randomly acquired cysts were imaged using a fixed

exposure as a z-stack with a $0.24\mu m$ step, with the center slice used for quantification

without deconvolution using AmyloQuant, a purpose developed quantitative imaging

based application (14).   The distribution of pixel intensities was defined in 4 bins

representing the background (black: 0-10 grayscale), low (blue: 10-25), intermediate

(green: 25-50) and high (red: >50) proportion of pixels (14). The relative distribution of

PAS intensities across the 4 bins is represented using a stacked plot with the distribution

patterns arrayed from lowest to highest intensity for each cohort. AmyloQuant additionally

presents a spatial heat map to allow for the distribution of PAS intensity (AG) to be

revealed within the imaged cyst. The mean pixel intensity of each cyst was additionally

determined following the definition of the ROI defined by the boundary of the cyst wall

using Image J. Measurements of cyst diameters in microns was achieved using the Zeiss

Zen imaging software.

Preparation of in vivo tissue cysts for TEM imaging

To prepare *T. gondii* tissue cysts generated *in vivo* for TEM, cysts were isolated from the

brains of infected mice as detailed above through the counting step. The Percoll fraction

containing mouse red blood cells (RBCs) was also recovered. After combining Percoll

fractions containing tissue cysts and diluting with PBS to a volume of 15 mL (maximum

of two 1-mL fractions were combined before dilution), cysts were pelleted for 15 min at

1000g at 4 °C. To maximize cyst recovery, 10 mL supernatant was removed, and the

remaining 5 mL was divided into 1 mL fractions for the top 4 mL, and the bottom 1 mL

directly above the pellet was sub-fractionated into 100 μL volumes. Typically, the majority

of cysts were localized to within 300 μL of the Percoll pellet, rather than in the pellet itself.

Sub-fractions containing cysts were once again combined and diluted (typically 200-300

μL diluted with 1-mL PBS) in a 1.5 mL Eppendorf tube, and then pelleted in a swinging-

bucket rotor for 10 min at 1000g and 4 °C. Leaving the pellet undisturbed, all but 50 μL of

the supernatant was removed. A small volume (~5-10 μL) of the reserved RBC fraction

was added to the remaining volume for ease of visualizing the pellet throughout the

remaining processing steps.

To ensure the detection of the relatively rare cyst population, a previously described

protocol (71, 72) was adapted to concentrate the cysts into a small agarose block. A

1.33X fixative solution of glutaraldhehyde (GA) in cacodylate buffer was prepared

containing 4% GA and 133 mM sodium cacodylate. 150 μL fixative solution was then

added to the 50 μL sample, bringing the total volume to 200 μL such that the final

concentration of GA was 3% and sodium cacodylate was 100 mM. Cysts were then

incubated at room temperature for 1 h in fixative. While cysts were in fixative, 4% low-

melt agarose (BioRad) was prepared in 100 mM sodium cacodylate buffer and kept liquid

at 70 °C until needed. After fixation, cysts were pelleted again at 1000g for 10 min at room

temperature in a table-top centrifuge. All but 50 μL supernatant was once again removed,

and 200 μL warm low-melt agarose was slowly added on top of fixed, pelleted cysts (3.2%

agarose, final concentration). Suspension was then centrifuged again at 1000g for 10 min

at 30 °C to keep the agarose semi-liquid, and then placed on ice for 20 min to solidify

agarose. After solidification, entire agarose plug was removed from tube with a small a

wooden dowel that had been whittled into a thin scoop. This agarose plug was placed in

a Petri dish, and the pellet was carefully cut out of the plug with a razor blade to create a

1 mm^3 block. The agarose block was then stored in 1X GA/cacodylate buffer overnight at

4°C. Processing of the block from post-stain onward was then identical to TEM processing

described above. During sectioning, thick sections cut on a glass knife were stained with

toluidine blue and examined for cysts using a light microscope prior to ultra-thin sectioning

once the cyst containing later was identified.

Data analyses

All data analyses, including graph preparation and statistics, were performed using

GraphPad Prism 9 or 10. Details on statistical tests applied are presented in the specific

figure legends.

1000

1001

Figure Legends.

Figure 1. Glucan dynamics in *T. gondii* ME49 tachyzoites and in vitro bradyzoites.

A, Microscopy-based glucan evaluation of *T. gondii* tachyzoites and bradyzoites using

PAS (left), IV58B6 (middle; α-glycogen IgM mAb), and TEM (right). **B,** GAA digest of AGs

in tachyzoites and bradyzoites confirms specificity of IV58B6 antibody and PAS staining.

All scale bars = 5 μm.

1008

Figure 2. Endogenous epitope tagging and localization of TgLaforin.

A, Schematic depicting the TgLaforin 3xHA-epitope tagging strategy. X = stop codon. **B,**

Successful tagging of TgLaforin (62 kDa) was verified using immunoblot analysis with an

α-HA antibody with SAG1 used as a loading control. **C,** IFA of Wild type (WT) and

TgLaforin-3XHA (TgLaf-HA) tagged parasites with α-HA antibody under tachyzoite and

in vitro bradyzoite conditions. Stage conversion is confirmed by Dolichos lectin (DBA)

staining. **D,** Western blot analysis of TgLaforin expression levels in tachyzoites (T) and in

vitro bradyzoites (B). Expression of TgLaf-HA is absent under in vitro bradyzoite

conditions. GAP45 is the loading control. Decrease in SAG1 alongside increase in SRS9

confirms tachyzoite to bradyzoite conversion. **E**, IFA of TgLaforin-HA colocalization with

PAS. Pearson's coefficient: 0.765. **F**, IFA of TgLaforin-HA colocalization with IV58B6.

Pearson's coefficient: 0.737. All scale bars = 5 μm.

1021

Figure 3. Loss of TgLaforin results in no gross glucan abnormalities under glucose

replete conditions. **A,** Schematic of TgLaforin KO strategy: the pyrimethamine-resistance

DHFR* gene containing 40-nt homologous arms (dark gray boxes on either end of gene)

was inserted into the TgLaforin locus via homologous recombination. A double stranded

break was induced using CRISPR/Cas9-GFP with a PAM site in the first exon. **B,**

Inside/out PCR verification of DHFR integration into the TgLaforin locus. Amplicons

(PCR1-4) are illustrated in (A). **C,** Loss of TgLaforin mRNA was confirmed by amplifying

full-length TgLaforin cDNA generated from both WT and ΔTgLaf strains. Actin cDNA

amplification serves both as a loading control and as a control to verify the absence of

gDNA. **D,** Analysis of glucan levels in ΔTgLaf tachyzoites using three different

approaches: PAS and IV58B6 immunofluorescence staining, and TEM.

1033

Figure 4. ΔTgLaf parasites are dependent on glutamine for normal plaque formation.

A, Steady-state metabolomics suggests upregulation of glutaminolysis in ΔTgLaf

parasites. Metabolite levels of intracellular tachyzoites were analyzed after 48 hours of

growth in HFFs by GC/MS analysis. Data were collected from 3 independent replicates.

Statistical comparisons were done using unpaired two-tailed t-tests. Statistical

significance is as follows: **$p<0.01$, ns=$p>0.05$, nd=not detected.

B, Representative plaque assays following staining of infected HFF monolayers with

crystal violet under nutrient replete (+gln) and glutamine free (gln-) conditions. WT,

ΔTgLaf (KO) and complemented (COMP) lines were evaluated.

C, Quantification of visible plaques for WT, DTgLaf and COMP lines under nutrient replete

and glutamine free conditions represented as the plaque area (pixels) was measured

across three independent replicates 6 days following infection.

D, Pixel area of nascent plaques/vacuoles after 3 days of growth as monitored by IFA

microscopy following staining with the PVM maker GRA3 under both glutamine replete

(+) and depleted conditions (-). **E**, Pixel area of plaques monitored as in (D) after 6 days

of growth. Statistical comparisons for C-E were done using an ordinary one-way ANOVA

using Tukey's post-hoc test to correct for multiple comparisons. Statistical significance is

indicated as follows: *$p < 0.05$, ****$p < 0.0001$, ns=$p > 0.05$.

1052

Figure 5. Loss of TgLaforin results in aberrant glucan morphology and accumulation in

in vitro bradyzoites only visible by TEM. **A,** *In vitro* tachyzoite to bradyzoite conversion

efficiency of ΔTgLaf vs WT parasites as measured by DBA intensity confirms no defect in

induced stage conversion in ΔTgLaf parasites.　**B,** Change in AG levels detected using

the intensity of PAS labeling during bradyzoite conversion reveals no statistically different

levels in AG accumulation based on PAS intensity. **C,** Representative TEM images of

bradyzoites from each indicated parasite line. At 4300x magnification, scale bar = 2 μm;

at 8600x magnification, scale bar = 1 μm. Arrowhead = canonical AG (white, round/ovoid);

Arrow = aberrant AG (grey, flattened, multi-lobed). **D,** Quantification of relative parasite

AG content and **E,** AG grayness across parasite lines using 8-bit grayscale (0=black,

255=white). Statistical comparisons were done using an ordinary one-way ANOVA with

Tukey's post-hoc test to correct for multiple comparisons. Statistical significance is

indicated as follows: *$p < 0.05$, ***$p < 0.001$, ****$p < 0.0001$, ns=$p > 0.05$.

1066

Figure 6.　Loss of TgLaforin attenuates tachyzoite virulence and cyst burden in mice.

Equal numbers of male and female CBA/J mice were infected i.p. with 100 tachyzoites

from each group and monitored for both symptom progression and mortality.

A, Symptomology throughout the acute (< day 20) and early chronic phases of infection

(day 21-28). Mice were monitored 1-2x/day and assigned a body score index ranging

from asymptomatic (Stage 0) to moribund/deceased. Staging of disease progression is

discussed in (52). **B,** Kaplan-Meier curve of mouse survival throughout acute and early

chronic infection following tachyzoite infection. **C,** A cohort of mice that survived 4-weeks

were euthanized, and the cyst burden determined as done previously described (70).

Error bars depict SD from the mean. Statistical comparison for Kaplan Meier curves is

indicated on plot, and statistical comparison of cyst burden was done using unpaired two-

tailed t-tests. Statistical significance: *$p < 0.05$, ***$p < 0.0002$.

1079

Figure 7. Mouse infection with ΔTgLaf tissue cysts results in milder illness and lower cyst

burden. CBA/J mice were infected i.p. with 20 cysts taken from previously infected mouse

brains and monitored for symptomology and death. Tissue cysts were harvested from

mice infected for 4 or 6 weeks respectively. **A,** Symptomology throughout the acute and

early chronic phase of infection as described in Figure 6B. **B,** Kaplan-Meier curve of

mouse survival during the acute and early chronic infection reveal low overall mortality

associated with cyst initiated infections for all parasite lines. **C,** Tissue cyst burden

following a 4-week bradyzoite initiated infection. Cyst numbers were determined as

described in Figure 6C. Statistical comparison of cyst burden was done using unpaired

two-tailed t-tests. Statistical significance: *$p < 0.05$, ns=$p > 0.05$.

1090

1091

1092

Figure 8. Quantification of amylopectin levels in tissue cysts based on PAS intensity.

A. The mean intensity of PAS labeled tissue cysts measured using Image J for WT,

∆TgLaf and COMP tissue cysts harvested at 4 and 6 weeks post infection reveal no

difference in AG levels across these parasite lines at week 4. Highly significant

differences are evident between WT/COMP and DTgLaf tissue cysts at week 6 post

infection indicating a temporal component for the emergence of the starch excess

phenotype. n=30 for each parasite line at each time point of harvest. Statistical analysis

was done using Tukey's multiple comparison test. P values: * p <0.05, **** p< 0.0001

B. Analysis of WT, DTgLaf and COMP tissue cysts harvested at weeks 4 and 6 using

AmyloQuant. The 30 tissue cysts per sample presented in A. were analyzed using

AmyloQuant. While acquired at random, the data are arrayed from low to high PAS

intensity presented as the percentage of pixels in each classification bin. The

classification bins defined as: background: Black: 0-10 grayscale, low: Blue: 10-25

grayscale, intermediate: Green 25-50 grayscale and high intensity pixels: Red >50

grayscale. The higher overall sensitivity of AmyloQuant reveals increased accumulation

of AG at week 4 in the DTgLAf and COMP lines. This phenotype is greatly exaggerated

at week 6 post infection consistent with a temporal component for the phenotypic

manifestation of the starch excess phenotype. The pattern in the COMP parasites is

intermediate between the WT and ∆TgLaf cysts. The distribution of PAS intensity levels

within the imaged tissue cysts generated by AmyloQuant for cysts in 5 cyst intervals is

presented under each set of stacked plots.

1114

Figure 9. Transmission electron microscopy of purified tissue cyst from infected mouse

brains confirms the presence of aberrant AG accumulation within ΔTgLaf cysts. The

accumulation of amylopectin granules within the cytoplasm of WT bradyzoites with

parasites exhibiting different levels. Additionally, evidence of active endodyogeny is

present. ΔTgLaf tissue cysts contain a mix of bradyzoites with expected cytoplasmic and

organellar contrast as well as others with grossly exaggerated AG levels that and the

apparent loss of both the cytoplasmic and organellar contents. The COMP line exhibits

AG accumulation levels similar to that observed in WT parasites with the tissue cyst itself

appearing to be very tightly packed, with high levels of granular material between

individual bradyzoites. Upper panels: scale bar = 5 μm; lower panels (zoom of boxed

region from upper panel): scale bar = 2 μm.

1126

Supplemental Files

Supplemental File 1.

Figure S1. Schematic of TgLaforin complementation strategy and confirmation of

successful expression of TgLaforin. **A.** Schematic of TgLaforin complementation into

ΔTgLaf parasites in which a PAM site was chosen at a neutral locus previously identified

in chromosome VI (99) to insert TgLaforin cDNA under its endogenous promoter. The

TgLaforin construct was connected to the HXGPRT selectable drug marker and inserted

using NHEJ. **B.** PCR confirmation of integration of TgLaforin construct into chromosome

VI. Primer sets are indicated above amplicons. WT primers amplify the same locus as in

Figure 3 ("PCR 1"), also present in the COMP line. KO primers amplify the chimeric locus

depicted in Figure 3 ("PCR 2"). Presence of the KO amplicon confirms that KO locus

remains intact in COMP line. COMP primers amplify the chimeric locus generated upon

insertion of the complementation construct. VI primers amplify the native chromosome VI

locus, which is lost only in the COMP line. **C,** Western blot confirms expression of

TgLaforin-HA in complemented parasites. Tagged LAF-HA parasites serve as a

comparison to confirm the correct MW (62 kDa) and expression level. **D,** IFA

demonstrates restoration of cytoplasmic, punctate localization of TgLaforin. Scale bar =

5 μm.

1145

Supplemental File 2

Figure S2. Loss of TgLaforin results in cumulative defects that cannot be pinpointed to a

single aspect of lytic cycle. **A.** Calcium ionophore-stimulated egress assay in which

parasites were pre-starved of glutamine for 48 hours, seeded onto HFFs and allowed to

grow for 48 hours to produce vacuoles containing >16 parasites, and stimulated with 3

μM A23187. Egress was monitored by video microscopy, and time to egress was

monitored as described in Materials and Methods. Data is the average of 3 biological

replicates that each consist of 4-5 technical replicates. COMP experiments measured 2

biological replicates. **B.** Zaprinast stimulated egress assay performed as described for

ionophore, however 500-μM zaprinast was used to stimulate egress. Data is the average

of 3 biological replicates that each consist of 2-5 technical replicates. COMP experiments

measured 2 biological replicates. **C.** Replication assay in which parasites were pre-

starved of glutamine for 48 hours, re-seeded into HFFs, and counted after 24 hours of

growth. Numbers (2, 4, or 8) indicate the number of tachyzoites counted per vacuole.

Data is the average of 3 biological replicates with at least 70 vacuoles counted per

replicate. **D.** Representative images of plaque formation at days 3 and 6, +/- glutamine.

Images were taken at 10X magnification using a SAG1 antibody to visualize vacuoles

and developing plaque size. The boundaries of vacuoles/ plaques were traced manually

to define their area. Regions of cell clearance was additionally scored. **E,** Percent of

plaques cleared was measured at both days 3 and 6 by dividing the area of the clearing

by total plaque size. Statistical comparisons were done using an ordinary one-way

ANOVA using Tukey's post-hoc test to correct for multiple comparisons. Error bars depict

SD from the mean. Statistical significance is indicated as follows: ns=p>0.05.

1169

Supplemental File 3

Figure S3. The loss of TgLaforin does not have any impact on distribution of tissue cyst

sizes. **A**. Tissue cyst diameters of 30 cysts each per line (WT, ΔTgLaf, COMP) per time

point (Week 4 and Week 6) were measured using DBA labeled cyst wall as the delimiter

using the Zeiss Zen software functionality. While all lines exhibit considerable variability,

no statistically significant differences in cyst size were noted in the DTgLaf mutant relative

to WT and COMP cysts. **B**. The relationship between tissue cyst size and AG

accumulation defined by mean PAS intensity was not found to have any significant

correlation.

1179

Supplemental File 4

Figure S4. Additional images of *in* vivo tissue cysts. **A.** A WT tissue cyst with two zoomed

in areas (orange and yellow dashed line boxes) exhibit variability in the levels of AG and

present with well defined cytoplasmic and organellar contents. **B.** A ΔTgLaf tissue cyst

presents with a mix of bradyzoites with both cytoplasmic and organelles as well as

bradyzoites lacking clearly defined organelles and/or cytoplasm. In addition enucleated

bradyzoites are evident (asterisks) as well as the displacement of rhoptries by elevated

levels of AG in the cytoplasm. Left column scale bar = 5 μm; right column zoom scale

bars = 2 μm.

1189

1190 **Supplemental File 5**

1191 **Supplemental Table 1**. sgRNA sequences used in this study.

1192

1193 **Supplemental File 6**

1194 **Supplemental Table 2.** Plasmid constructs used in this study.

1195

Supplemental File 7

Supplemental Table 3. DNA primer sequences and their application in this study

----

Acknowledgements: The authors wish to acknowledge Dr. Elizabeth Watts who

conducted early exploratory studies that encouraged the pursuit of this project, Jim

Begley (Imaging Center, University of Kentucky) for preparation of EM samples and

willingness to explore new methodologies for isolating rare *in vivo* cysts, Jillian Cramer

(Electron Microscopy Center, University of Kentucky) for acquisition of TEM images, and

Jennifer Strange (Flow Cytometry and Immune Monitoring Core Facility, University of

Kentucky) for assistance with isolation of GFP+ *T. gondii* mutants using flow cytometry.

We additionally thank Dr's David Sibley (Washington University) and Peter Bradley

(UCLA) for their gifts of DNA constructs and antibodies.

1208

Author contributions:

Conceptualization (R.D.M., A.P.S., M.S.G.), Methodology (R.D.M., A.D., C.O.B., A.T.,

A.P.S., M.S.G.), Formal analysis (R.D.M., C.A.T., J.S.M., L.E.A.Y., R.C.S., A.T, A.P.,

C.W.V.K., A.P.S.), Investigation (R.D.M., C.A.T., J.S.M., L.E.A.Y., A.P.S.), Resources

(M.S.G., A.P.S.), Writing—original draft (R.D.M., A.P.S.), Writing—review and editing (all

authors), Visualization (R.D.M., C.A.T., J.S.M., A.P.S.), Supervision (M.S.G., A.P.S.),

Project administration (R.D.M., M.S.G., A.P.S.), Funding acquisition (R.D.M., C.W.V.K.,

M.S.G., A.P.S.).

1217

1218   ***Funding:*** This work was supported by: GRFP 1247392 to R.D.M., NIH grants R21

1219   AI150631 to A.P.S and R01 AI145335 to A.P.S. and A.P, and NIH grants R35 NS116824,

1220   P01 NS097197, and NSF CHE 1808304 to C.W.V.K. and MCB 2308488 to M.S.G.

1221

1222   ***Conflicts of Interest:*** none

1223

Abbreviations: AG, amylopectin granule; CDPK2, calcium dependent protein kinase2;

COMP, TgLaforin-complemented parasite line; CRISPR, clustered regularly interspersed

short palindromic repeats; DBA, *Dolichos biflorus* agglutinin lectin; ΔTgLaf, TgLaforin-KO

parasite line; DHFR, dihydrofolate reductase; FACS, fluorescence activated cell sorting;

GAA,   acid-α-amyloglucosidase;   Gal-NAc,   N-acetylgalactosamine;   GAPDH1,

glyceraldehyde-3-phosphate dehydrogenase; GC/MS, gas chromatography/mass

spectrometry; GFP, green fluorescent protein; GT1, glucose transporter1; GWD, glucan,

water di-kinase; HA, hemagglutinin; HFF, human foreskin fibroblasts; HK, hexokinase;

HR, homologous recombination; HXGPRT, hypoxanthine-xanthine-guanine

phosphoribosyl transferase; IF, immunofluorescence; i.p., intraperitoneally; KO,

knockout; MPA, mycophenolic acid; PAM, protospacer adjacent motif; PAS, periodic acid-

Schiff; PV, parasitophorous vacuole; PWD, phospho-glucan, water di-kinase; PYK1,

pyruvate kinase1; RBC, red blood cell; RT, room temperature; SEX4, starch-excess4;

SS, starch/glycogen synthase; TCA, tricarboxylic acid; TEM, transmission electron

microscopy; UTR, untranslated region; WT, parental ME49ΔHXGPRT parasite line used

in this study.

1240

Literature Cited

1. Flegr J, Prandota J, Sovickova M, Israili ZH. 2014. Toxoplasmosis--a global threat.

Correlation of latent toxoplasmosis with specific disease burden in a set of 88

countries. PLoS One 9:e90203.

2. Tenter AM, Heckeroth AR, Weiss LM. 2000. *Toxoplasma gondii*: from animals to

humans. Int J Parasitol 30:1217-58.

3. Jones JL, Dubey JP. 2012. Foodborne toxoplasmosis. Clin Infect Dis 55:845-51.

4. Derouin F, Garin YJ. 1991. *Toxoplasma gondii*: blood and tissue kinetics during acute

and chronic infections in mice. Exp Parasitol 73:460-8.

5. Pittman KJ, Knoll LJ. 2015. Long-Term Relationships: the Complicated Interplay
between the Host and the Developmental Stages of *Toxoplasma gondii* during
Acute and Chronic Infections. Microbiol Mol Biol Rev 79:387-401.

6. Zhao XY, Ewald SE. 2020. The molecular biology and immune control of chronic
Toxoplasma gondii infection. J Clin Invest 130:3370-3380.

7. Luft BJ, Hafner R, Korzun AH, Leport C, Antoniskis D, Bosler EM, Bourland DD, 3rd,
Uttamchandani R, Fuhrer J, Jacobson J, et al. 1993. Toxoplasmic encephalitis in
patients with the acquired immunodeficiency syndrome. Members of the ACTG
077p/ANRS 009 Study Team. N Engl J Med 329:995-1000.

8. Luft BJ, Remington JS. 1992. Toxoplasmic encephalitis in AIDS. Clin Infect Dis
15:211-22.

9. Cerutti A, Blanchard N, Besteiro S. 2020. The Bradyzoite: A Key Developmental Stage
for the Persistence and Pathogenesis of Toxoplasmosis. Pathogens 9.

10. Sinai AP, Watts EA, Dhara A, Murphy RD, Gentry MS, Patwardhan A. 2016.
Reexamining Chronic *Toxoplasma gondii* Infection: Surprising Activity for a
"Dormant" Parasite. Curr Clin Microbiol Rep 3:175-185.

11. Watts E, Zhao Y, Dhara A, Eller B, Patwardhan A, Sinai AP. 2015. Novel Approaches
Reveal that *Toxoplasma gondii* Bradyzoites within Tissue Cysts Are Dynamic and
Replicating Entities In Vivo. MBio 6:e01155-15.

12. Place BC, Troublefield C, Murphy RD, Sinai AP, Patwardhan A. 2021. Computer
Aided Image Processing to Facilitate Determination of Congruence in Manual
Classification of Mitochondrial Morphologies in *Toxoplasma gondii* Tissue Cysts.
Annu Int Conf IEEE Eng Med Biol Soc 2021:3509-3513.

13. Place BC, Troublefield CA, Murphy RD, Sinai AP, Patwardhan AR. 2023. Machine

learning based classification of mitochondrial morphologies from fluorescence

microscopy images of *Toxoplasma gondii* cysts. PLoS One 18:e0280746.

14. Tripathi A, Donkin RW, Miracle JS, Murphy RD, Patwardhan A, Sinai AP. 2024.

Dynamics of amylopectin granule accumulation during the course of the chronic

Toxoplasma infection is linked to intra-cyst bradyzoite replication. bioRxiv

doi:10.1101/2024.09.02.610794:2024.09.02.610794.

15. MacNeill GJ, Mehrpouyan S, Minow MAA, Patterson JA, Tetlow IJ, Emes MJ. 2017.

Starch as a source, starch as a sink: the bifunctional role of starch in carbon

allocation. J Exp Bot 68:4433-4453.

16. Dubey JP, Lindsay DS, Speer CA. 1998. Structures of *Toxoplasma gondii* tachyzoites,

bradyzoites, and sporozoites and biology and development of tissue cysts. Clin

Microbiol Rev 11:267-99.

17. Coppin A, Dzierszinski F, Legrand S, Mortuaire M, Ferguson D, Tomavo S. 2003.

Developmentally regulated biosynthesis of carbohydrate and storage

polysaccharide during differentiation and tissue cyst formation in *Toxoplasma*

gondii. Biochimie 85:353-61.

18. Guerardel Y, Leleu D, Coppin A, Lienard L, Slomianny C, Strecker G, Ball S, Tomavo

S. 2005. Amylopectin biogenesis and characterization in the protozoan parasite

Toxoplasma gondii, the intracellular development of which is restricted in the

HepG2 cell line. Microbes Infect 7:41-8.

19. Coppin A, Varre JS, Lienard L, Dauvillee D, Guerardel Y, Soyer-Gobillard MO, Buleon

A, Ball S, Tomavo S. 2005. Evolution of plant-like crystalline storage

polysaccharide in the protozoan parasite *Toxoplasma gondii* argues for a red alga

ancestry. J Mol Evol 60:257-67.

20. Uboldi AD, McCoy JM, Blume M, Gerlic M, Ferguson DJ, Dagley LF, Beahan CT,

Stapleton DI, Gooley PR, Bacic A, Masters SL, Webb AI, McConville MJ, Tonkin

CJ. 2015. Regulation of Starch Stores by a Ca(2+)-Dependent Protein Kinase Is

Essential for Viable Cyst Development in *Toxoplasma gondii*. Cell Host Microbe

18:670-81.

21. Yang J, He Z, Chen C, Zhao J, Fang R. 2022. Starch Branching Enzyme 1 Is Important

for Amylopectin Synthesis and Cyst Reactivation in *Toxoplasma gondii*. Microbiol

Spectr 10:e0189121.

22. Yang J, Yang C, Qian J, Li F, Zhao J, Fang R. 2020. *Toxoplasma gondii* alpha-

amylase deletion mutant is a promising vaccine against acute and chronic

toxoplasmosis. Microb Biotechnol 13:2057-2069.

23. Lyu C, Yang X, Yang J, Hou L, Zhou Y, Zhao J, Shen B. 2021. Role of amylopectin

synthesis in *Toxoplasma gondii* and its implication in vaccine development against

toxoplasmosis. Open Biol 11:200384.

24. Blume M, Nitzsche R, Sternberg U, Gerlic M, Masters SL, Gupta N, McConville MJ.

2015. A *Toxoplasma gondii* Gluconeogenic Enzyme Contributes to Robust Central

Carbon Metabolism and Is Essential for Replication and Virulence. Cell Host

Microbe 18:210-20.

25. Ralton JE, Sernee MF, McConville MJ. 2021. Evolution and function of carbohydrate

reserve biosynthesis in parasitic protists. Trends Parasitol 37:988-1001.

26. Sernee MF, Ralton JE, Nero TL, Sobala LF, Kloehn J, Vieira-Lara MA, Cobbold SA,

1319   Stanton L, Pires DEV, Hanssen E, Males A, Ward T, Bastidas LM, van der Peet

1320   PL, Parker MW, Ascher DB, Williams SJ, Davies GJ, McConville MJ. 2019. A

1321   Family of Dual-Activity Glycosyltransferase-Phosphorylases Mediates Mannogen

1322   Turnover and Virulence in Leishmania Parasites. Cell Host Microbe 26:385-399

1323   e9.

1324   27. Chen P, Lyu CC, Wang YD, Pan M, Lin XY, Shen B. 2023. Key roles of amylopectin

1325   synthesis and degradation enzymes in the establishment and reactivation of

1326   chronic toxoplasmosis. Animal Diseases 3.

1327   28. Stitt M, Zeeman SC. 2012. Starch turnover: pathways, regulation and role in growth.

1328   Curr Opin Plant Biol 15:282-92.

1329   29. Kotting O, Kossmann J, Zeeman SC, Lloyd JR. 2010. Regulation of starch

1330   metabolism: the age of enlightenment? Curr Opin Plant Biol 13:321-9.

1331   30. Silver DM, Kotting O, Moorhead GB. 2014. Phosphoglucan phosphatase function

1332   sheds light on starch degradation. Trends Plant Sci 19:471-8.

1333   31. Ritte G, Lloyd JR, Eckermann N, Rottmann A, Kossmann J, Steup M. 2002. The

1334   starch-related R1 protein is an alpha -glucan, water dikinase. Proc Natl Acad Sci

1335   U S A 99:7166-71.

1336   32. Kotting O, Pusch K, Tiessen A, Geigenberger P, Steup M, Ritte G. 2005. Identification

1337   of a novel enzyme required for starch metabolism in Arabidopsis leaves. The

1338   phosphoglucan, water dikinase. Plant Physiol 137:242-52.

33. Edner C, Li J, Albrecht T, Mahlow S, Hejazi M, Hussain H, Kaplan F, Guy C, Smith

SM, Steup M, Ritte G. 2007. Glucan, water dikinase activity stimulates breakdown

of starch granules by plastidial beta-amylases. Plant Physiol 145:17-28.

34. Hejazi M, Fettke J, Haebel S, Edner C, Paris O, Frohberg C, Steup M, Ritte G. 2008.

Glucan, water dikinase phosphorylates crystalline maltodextrins and thereby

initiates solubilization. Plant J 55:323-34.

35. Hansen PI, Spraul M, Dvortsak P, Larsen FH, Blennow A, Motawia MS, Engelsen SB.

2009. Starch phosphorylation--maltosidic restrains upon 3'- and 6'-phosphorylation

investigated by chemical synthesis, molecular dynamics and NMR spectroscopy.

Biopolymers 91:179-93.

36. Murphy RD, Chen T, Lin J, He R, Wu L, Pearson CR, Sharma S, Vander Kooi CD,

Sinai AP, Zhang ZY, Vander Kooi CW, Gentry MS. 2022. The Toxoplasma glucan

phosphatase TgLaforin utilizes a distinct functional mechanism that can be

exploited by therapeutic inhibitors. J Biol Chem 298:102089.

37. Kotting O, Santelia D, Edner C, Eicke S, Marthaler T, Gentry MS, Comparot-Moss S,

Chen J, Smith AM, Steup M, Ritte G, Zeeman SC. 2009. STARCH-EXCESS4 Is a

Laforin-Like Phosphoglucan Phosphatase Required for Starch Degradation in

Arabidopsis thaliana. Plant Cell 21:334-46.

38. Minassian BA, Lee JR, Herbrick JA, Huizenga J, Soder S, Mungall AJ, Dunham I,

Gardner R, Fong CY, Carpenter S, Jardim L, Satishchandra P, Andermann E,

Snead OC, 3rd, Lopes-Cendes I, Tsui LC, Delgado-Escueta AV, Rouleau GA,

Scherer SW. 1998. Mutations in a gene encoding a novel protein tyrosine

phosphatase cause progressive myoclonus epilepsy. Nat Genet 20:171-4.

39. Gentry MS, Guinovart JJ, Minassian BA, Roach PJ, Serratosa JM. 2018. Lafora

disease offers a unique window into neuronal glycogen metabolism. J Biol Chem

293:7117-7125.

40. Duran J, Hervera A, Markussen KH, Varea O, Lopez-Soldado I, Sun RC, Del Rio JA,

Gentry MS, Guinovart JJ. 2021. Astrocytic glycogen accumulation drives the

pathophysiology of neurodegeneration in Lafora disease. Brain 144:2349-2360.

41. Sugi T, Tu V, Ma Y, Tomita T, Weiss LM. 2017. *Toxoplasma gondii* Requires Glycogen

Phosphorylase for Balancing Amylopectin Storage and for Efficient Production of

Brain Cysts. MBio 8.

42. Cao XZ, Wang JL, Elsheikha HM, Li TT, Sun LX, Liang QL, Zhang ZW, Lin RQ. 2019.

Characterization of the Role of Amylo-Alpha-1,6-Glucosidase Protein in the

Infectivity of *Toxoplasma gondii*. Front Cell Infect Microbiol 9:418.

43. Wang JL, Li TT, Elsheikha HM, Liang QL, Zhang ZW, Wang M, Sibley LD, Zhu XQ.

2022. The protein phosphatase 2A holoenzyme is a key regulator of starch

metabolism and bradyzoite differentiation in *Toxoplasma gondii*. Nat Commun

13:7560.

44. Zhao M, Yang Y, Shi Y, Chen X, Yang Y, Pan L, Du Z, Sun H, Yao C, Ma G, Du A.

2023. PP2Acalpha-B'/PR61 Holoenzyme of *Toxoplasma gondii* Is Required for the

Amylopectin Metabolism and Proliferation of Tachyzoites. Microbiol Spectr

11:e0010423.

45. Gentry MS, Dowen RH, 3rd, Worby CA, Mattoo S, Ecker JR, Dixon JE. 2007. The

phosphatase laforin crosses evolutionary boundaries and links carbohydrate

metabolism to neuronal disease. J Cell Biol 178:477-88.

46. Gentry MS, Pace RM. 2009. Conservation of the glucan phosphatase laforin is linked

to rates of molecular evolution and the glycogen metabolism of the organism. BMC

Evol Biol 9:138.

47. Baba O. 1993. [Production of monoclonal antibody that recognizes glycogen and its

application for immunohistochemistry]. Kokubyo Gakkai Zasshi 60:264-87.

48. van de Weerd R, Berbis MA, Sparrius M, Maaskant JJ, Boot M, Paauw NJ, de Vries

N, Boon L, Baba O, Canada FJ, Geurtsen J, Jimenez-Barbero J, Appelmelk BJ.

2015. A murine monoclonal antibody to glycogen: characterization of epitope-fine

specificity by saturation transfer difference (STD) NMR spectroscopy and its use

in mycobacterial capsular alpha-glucan research. Chembiochem 16:977-89.

49. Oe Y, Baba O, Ashida H, Nakamura KC, Hirase H. 2016. Glycogen distribution in the

microwave-fixed mouse brain reveals heterogeneous astrocytic patterns. Glia

64:1532-45.

50. Skurat AV, Segvich DM, DePaoli-Roach AA, Roach PJ. 2017. Novel method for

detection of glycogen in cells. Glycobiology 27:416-424.

51. Guimaraes EV, de Carvalho L, Barbosa HS. 2003. An alternative technique to reveal

polysaccharides in *Toxoplasma gondii* tissue cysts. Mem Inst Oswaldo Cruz

98:915-7.

52. Troublefield CA, Miracle JS, Murphy RD, Donkin RW, Sinai AP. 2023. Factors

Influencing Tissue Cyst Yield in a Murine Model of Chronic Toxoplasmosis. Infect

Immun 91:e0056622.

53. Sidik SM, Hackett CG, Tran F, Westwood NJ, Lourido S. 2014. Efficient genome

engineering of *Toxoplasma gondii* using CRISPR/Cas9. PLoS One 9:e100450.

54. Fritz HM, Buchholz KR, Chen X, Durbin-Johnson B, Rocke DM, Conrad PA,

Boothroyd JC. 2012. Transcriptomic analysis of toxoplasma development reveals

many novel functions and structures specific to sporozoites and oocysts. PLoS

One 7:e29998.

55. Choi CP, Moon AS, Back PS, Jami-Alahmadi Y, Vashisht AA, Wohlschlegel JA,

Bradley PJ. 2019. A photoactivatable crosslinking system reveals protein

interactions in the *Toxoplasma gondii* inner membrane complex. PLoS Biol

17:e3000475.

56. Shen B, Brown KM, Lee TD, Sibley LD. 2014. Efficient gene disruption in diverse

strains of *Toxoplasma gondii* using CRISPR/CAS9. mBio 5:e01114-14.

57. Sidik SM, Huet D, Ganesan SM, Huynh MH, Wang T, Nasamu AS, Thiru P, Saeij JP,

Carruthers VB, Niles JC, Lourido S. 2016. A Genome-wide CRISPR Screen in

Toxoplasma Identifies Essential Apicomplexan Genes. Cell 166:1423-1435 e12.

58. Markus BM, Bell GW, Lorenzi HA, Lourido S. 2019. Optimizing Systems for Cas9

Expression in *Toxoplasma gondii*. mSphere 4.

59. Sun RC, Young LEA, Bruntz RC, Markussen KH, Zhou Z, Conroy LR, Hawkinson TR,

Clarke HA, Stanback AE, Macedo JKA, Emanuelle S, Brewer MK, Rondon AL,

Mestas A, Sanders WC, Mahalingan KK, Tang B, Chikwana VM, Segvich DM,

Contreras CJ, Allenger EJ, Brainson CF, Johnson LA, Taylor RE, Armstrong DD,

Shaffer R, Waechter CJ, Vander Kooi CW, DePaoli-Roach AA, Roach PJ, Hurley

TD, Drake RR, Gentry MS. 2021. Brain glycogen serves as a critical glucosamine

cache required for protein glycosylation. Cell Metab 33:1404-1417 e9.

60. Brewer MK, Uittenbogaard A, Austin GL, Segvich DM, DePaoli-Roach A, Roach PJ,

McCarthy JJ, Simmons ZR, Brandon JA, Zhou Z, Zeller J, Young LEA, Sun RC,

Pauly JR, Aziz NM, Hodges BL, McKnight TR, Armstrong DD, Gentry MS. 2019.

Targeting Pathogenic Lafora Bodies in Lafora Disease Using an Antibody-Enzyme

Fusion. Cell Metab 30:689-705 e6.

61. King EFB, Cobbold SA, Uboldi AD, Tonkin CJ, McConville MJ. 2020. Metabolomic

Analysis of *Toxoplasma gondii* Tachyzoites. Methods Mol Biol 2071:435-452.

62. Nitzsche R, Zagoriy V, Lucius R, Gupta N. 2016. Metabolic Cooperation of Glucose

and Glutamine Is Essential for the Lytic Cycle of Obligate Intracellular Parasite

Toxoplasma gondii. J Biol Chem 291:126-41.

63. Oppenheim RD, Creek DJ, Macrae JI, Modrzynska KK, Pino P, Limenitakis J,

Polonais V, Seeber F, Barrett MP, Billker O, McConville MJ, Soldati-Favre D. 2014.

BCKDH: the missing link in apicomplexan mitochondrial metabolism is required for

full virulence of *Toxoplasma gondii* and Plasmodium berghei. PLoS Pathog

10:e1004263.

64. Endo T, Sethi KK, Piekarski G. 1982. *Toxoplasma gondii*: calcium ionophore A23187-

mediated exit of trophozoites from infected murine macrophages. Exp Parasitol

53:179-88.

65. Lourido S, Tang K, Sibley LD. 2012. Distinct signalling pathways control Toxoplasma

egress and host-cell invasion. EMBO J 31:4524-34.

66. Hotchin J, Kinch W, Benson L. 1971. Lytic and turbid plaque-type mutants of

lymphocytic choriomeningitis virus as a cause of neurological disease or persistent

infection. Infect Immun 4:281-6.

67. Fiala M. 1968. Plaque formation by 55 rhinovirus serotypes. Appl Microbiol 16:1445-
1454   50.

1455   68. Ferguson DJ. 2004. Use of molecular and ultrastructural markers to evaluate stage
1456   conversion of *Toxoplasma gondii* in both the intermediate and definitive host. Int J
1457   Parasitol 34:347-60.

1458   69. Sinai AP. 2008. Biogenesis of and activities at the *Toxoplasma gondii* parasitophorous
1459   vacuole membrane. Subcell Biochem 47:155-64.

1460   70. Watts EA, Dhara A, Sinai AP. 2017. Purification *Toxoplasma gondii* Tissue Cysts
1461   Using Percoll Gradients. Curr Protoc Microbiol 45:20C 2 1-20C 2 19.

1462   71. Kumar S, Ciraolo G, Hinge A, Filippi MD. 2014. An efficient and reproducible process
1463   for transmission electron microscopy (TEM) of rare cell populations. J Immunol
1464   Methods 404:87-90.

1465   72. Kumar S, Filippi MD. 2016. An Alternative Approach for Sample Preparation with Low
1466   Cell Number for TEM Analysis. J Vis Exp doi:10.3791/54724.

1467   73. Sinai AP, Suvorova ES. 2020. The RESTRICTION checkpoint: a window of
1468   opportunity governing developmental transitions in *Toxoplasma gondii*. Curr Opin
1469   Microbiol 58:99-105.

1470   74. Xue Y, Theisen TC, Rastogi S, Ferrel A, Quake SR, Boothroyd JC. 2020. A single-
1471   parasite transcriptional atlas of *Toxoplasma gondii* reveals novel control of antigen
1472   expression. Elife 9.

1473   75. Olson WJ, Martorelli Di Genova B, Gallego-Lopez G, Dawson AR, Stevenson D,
1474   Amador-Noguez D, Knoll LJ. 2020. Dual metabolomic profiling uncovers

Toxoplasma manipulation of the host metabolome and the discovery of a novel

parasite metabolic capability. PLoS Pathog 16:e1008432.

76. Zeeman SC, Delatte T, Messerli G, Umhang M, Stettler M, Mettler T, Streb S, Reinhold

H, Kotting O. 2007. Starch breakdown: recent discoveries suggest distinct

pathways and novel mechanisms. Funct Plant Biol 34:465-473.

77. Zeeman SC, Smith SM, Smith AM. 2007. The diurnal metabolism of leaf starch.

Biochem J 401:13-28.

78. Mahlow S, Orzechowski S, Fettke J. 2016. Starch phosphorylation: insights and

perspectives. Cell Mol Life Sci 73:2753-64.

79. Ferguson DJ, Hutchison WM. 1987. An ultrastructural study of the early development

and tissue cyst formation of *Toxoplasma gondii* in the brains of mice. Parasitol Res

73:483-91.

80. Niittyla T, Comparot-Moss S, Lue WL, Messerli G, Trevisan M, Seymour MD,

Gatehouse JA, Villadsen D, Smith SM, Chen J, Zeeman SC, Smith AM. 2006.

Similar protein phosphatases control starch metabolism in plants and glycogen

metabolism in mammals. J Biol Chem 281:11815-8.

81. Zeeman SC, Northrop F, Smith AM, Rees T. 1998. A starch-accumulating mutant of

Arabidopsis thaliana deficient in a chloroplastic starch-hydrolysing enzyme. Plant

J 15:357-65.

82. Shukla A, Olszewski KL, Llinas M, Rommereim LM, Fox BA, Bzik DJ, Xia D, Wastling

J, Beiting D, Roos DS, Shanmugam D. 2018. Glycolysis is important for optimal

asexual growth and formation of mature tissue cysts by *Toxoplasma gondii*. Int J

Parasitol 48:955-968.

83. Blume M, Rodriguez-Contreras D, Landfear S, Fleige T, Soldati-Favre D, Lucius R,

Gupta N. 2009. Host-derived glucose and its transporter in the obligate intracellular

pathogen *Toxoplasma gondii* are dispensable by glutaminolysis. Proc Natl Acad

Sci U S A 106:12998-3003.

84. Dubey R, Staker BL, Foe IT, Bogyo M, Myler PJ, Ngo HM, Gubbels MJ. 2017.

Membrane skeletal association and post-translational allosteric regulation of

Toxoplasma gondii GAPDH1. Mol Microbiol 103:618-634.

85. Xia N, Ye S, Liang X, Chen P, Zhou Y, Fang R, Zhao J, Gupta N, Yang S, Yuan J,

Shen B. 2019. Pyruvate Homeostasis as a Determinant of Parasite Growth and

Metabolic Plasticity in *Toxoplasma gondii*. mBio 10.

86. Suresh K, Mak JW, Yong HS. 1991. Immune response in acute Toxoplasma infection

of Balb/C, ICR and CBA/J mice. Southeast Asian J Trop Med Public Health 22:452-

4.

87. Araujo FG, Huskinson-Mark J, Gutteridge WE, Remington JS. 1992. In vitro and in

vivo activities of the hydroxynaphthoquinone 566C80 against the cyst form of

Toxoplasma gondii. Antimicrob Agents Chemother 36:326-30.

88. Araujo FG, Lin T, Remington JS. 1993. The activity of atovaquone (566C80) in murine

toxoplasmosis is markedly augmented when used in combination with

pyrimethamine or sulfadiazine. J Infect Dis 167:494-7.

89. Doggett JS, Nilsen A, Forquer I, Wegmann KW, Jones-Brando L, Yolken RH, Bordon

C, Charman SA, Katneni K, Schultz T, Burrows JN, Hinrichs DJ, Meunier B,

Carruthers VB, Riscoe MK. 2012. Endochin-like quinolones are highly efficacious

against acute and latent experimental toxoplasmosis. Proc Natl Acad Sci U S A

109:15936-41.

90. Doggett JS, Schultz T, Miller AJ, Bruzual I, Pou S, Winter R, Dodean R, Zakharov LN,

Nilsen A, Riscoe MK, Carruthers VB. 2020. Orally Bioavailable Endochin-Like

Quinolone Carbonate Ester Prodrug Reduces *Toxoplasma gondii* Brain Cysts.

Antimicrob Agents Chemother 64.

91. Vidadala RS, Rivas KL, Ojo KK, Hulverson MA, Zambriski JA, Bruzual I, Schultz TL,

Huang W, Zhang Z, Scheele S, DeRocher AE, Choi R, Barrett LK, Siddaramaiah

LK, Hol WG, Fan E, Merritt EA, Parsons M, Freiberg G, Marsh K, Kempf DJ,

Carruthers VB, Isoherranen N, Doggett JS, Van Voorhis WC, Maly DJ. 2016.

Development of an Orally Available and Central Nervous System (CNS) Penetrant

Toxoplasma gondii Calcium-Dependent Protein Kinase 1 (TgCDPK1) Inhibitor with

Minimal Human Ether-a-go-go-Related Gene (hERG) Activity for the Treatment of

Toxoplasmosis. J Med Chem 59:6531-46.

92. McPhillie MJ, Zhou Y, Hickman MR, Gordon JA, Weber CR, Li Q, Lee PJ,

Amporndanai K, Johnson RM, Darby H, Woods S, Li ZH, Priestley RS, Ristroph

KD, Biering SB, El Bissati K, Hwang S, Hakim FE, Dovgin SM, Lykins JD, Roberts

L, Hargrave K, Cong H, Sinai AP, Muench SP, Dubey JP, Prud'homme RK, Lorenzi

HA, Biagini GA, Moreno SN, Roberts CW, Antonyuk SV, Fishwick CWG, McLeod

R. 2020. Potent Tetrahydroquinolone Eliminates Apicomplexan Parasites. Front

Cell Infect Microbiol 10:203.

93. Brown KM, Long S, Sibley LD. 2018. Conditional Knockdown of Proteins Using Auxin-

inducible Degron (AID) Fusions in *Toxoplasma gondii*. Bio Protoc 8.

94. Bolte S, Cordelieres FP. 2006. A guided tour into subcellular colocalization analysis

in light microscopy. J Microsc 224:213-32.

95. Mayoral J, Di Cristina M, Carruthers VB, Weiss LM. 2020. *Toxoplasma gondii*:

Bradyzoite Differentiation In Vitro and In Vivo. Methods Mol Biol 2071:269-282.

96. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch

S, Rueden C, Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V, Eliceiri

K, Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-

image analysis. Nat Methods 9:676-82.

97. Sinai AP, Webster P, Joiner KA. 1997. Association of host cell endoplasmic reticulum

and mitochondria with the *Toxoplasma gondii* parasitophorous vacuole

membrane: a high affinity interaction. J Cell Sci 110 (Pt 17):2117-28.

98. Andres DA, Young LEA, Veeranki S, Hawkinson TR, Levitan BM, He D, Wang C, Satin

J, Sun RC. 2020. Improved workflow for mass spectrometry-based metabolomics

analysis of the heart. J Biol Chem 295:2676-2686.

99. Markus BM, Bell GW, Lorenzi HA, Lourido S. 2019. Optimizing systems for Cas9

expression in *Toxoplasma gondii*. MSphere 4:e00386-19.

1559

Murphy et al. 2024 Figure 1

A.

gDNA (WT)

TgLaforin | X | 3'UTR

PAM

+

Donor patch | HA | X | UTR (DHFR) | HX

HA-tagged TgLaforin

TgLaforin | HA | X | UTR (DHFR) | HX | UTR

B.

kDa

100
75 — α-HA

37
25 — α-SAG

C.

TgLaf-HA

WT | Tachy | Brady

SAG
HA

DBA
HA

D.

kDa | T | B

75 — α-HA

50
40 — α-GAP45

37
25 — α-SAG1

E.

TgLaforin/PAS colocalization

TgLaforin | PAS | Merge | Phase

5 μm

F.

TgLaforin/IV colocalization

TgLaforin | IV | Merge | Phase

Murphy et al 2024 Figure 2

A.

WT

PCR 1

5'UTR

PCR 3

3'UTR

PAM

pyrimethamine
resistance

DHFR*

ΔTgLaf

PCR 2

5'UTR

DHFR*

PCR 4

3'UTR

B.

Diagnostic PCRs

bp

PCR 1   PCR 2   PCR 3   PCR 4

WT KO   WT KO   WT KO   WT KO

C.

	PAS		IV58B6		TEM
WT	PAS DAPI	Merge	IV58B6 GAP45 DAPI	Phase	
ΔTgLaf					5-μm / 2-μm

Murphy et al. 2024 Figure 3

Murphy et al. 2024 Figure 4

Murphy et al. 2024 Figure 5

Murphy et al. 2024 Figure 6

Murphy et al. 2024 Figure 7

Murphy et al. 2024 Figure 8

WT **ΔTgLaf** **COMP**

5 µm

2 µm

Murphy et al. 2024 Figure 9