DIMINISHING GROWTH OF ZOOLOGICAL NATURAL HISTORY

COLLECTIONS THREATENS FUTURE RESEARCH IN THE LIFE

SCIENCES AND BEYOND

Daren C. Card[1,2,*]

[1] Department of Organismic & Evolutionary Biology, Harvard University, Cambridge,

Massachusetts, USA

[2] Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts, USA

[*] Correspondence:

Daren C. Card: dcard@fas.harvard.edu

Abstract

Natural history museums comprise a unique and important component of biological

infrastructure worldwide, and underlie diverse research, education, and outreach in the natural

sciences. Each museum is built around one or more biological collections that serve as a

repository of important materials for study, which have contributed to scientific research in

increasingly important areas ranging from understanding global climate change to developing

new biotechnological applications. However, despite centuries of existence, global collections

sizes have only been recently estimated, and aside from analyses of certain institutions or well-

studied clades, little is known about how patterns of collecting activity across institutions,

geographic regions, or taxonomic clades, or how these patterns vary across time. To address this

important gap in our understanding of critical life sciences infrastructure, I gathered and

analyzed zoological records associated with preserved specimens collected between 1900 and

2015 and housed in worldwide natural history collections using data from the Global

Biodiversity Information Facility. My analysis indicated that global, museum-associated

collecting activity focused on animals has varied greatly over time, peaking around 2000 and

declining significantly since. By stratifying data by institution, nation, and taxonomic phylum

and class, I also illuminated how individual data series contribute to these global patterns.

Institutions and nations had either concentrated or dispersed periods of relatively high collecting

activity that occurred over different time periods, although most growth occurred in the second

half of the 20th century for both many individual institutions and globally. Certain taxonomic

clades comprise the largest proportions of collections records over this time period, namely

arthropods (especially insects) and chordates (especially vertebrates), underscoring the

taxonomically biased collecting histories of many institutions. Altogether, my analyses provide a

critical, early view of historical zoological collecting activity and, to help other museum

stakeholders to explore the data and results in more depth, I also describe and release

NHMinformatics, an interactive dashboard built using R Shiny. These resources, when combined

with recommendations I have made for sustainable collections growth, will be helpful for

establishing policy goals, provisioning museum infrastructure, and training curatorial personnel.

Keywords: biological collections, digitization, GBIF, historical, informatics, infrastructure,

policy, time-series analysis

Running header: Future research is threatened by declining natural history museum growth

Introduction

Natural history museums are important institutions in the life and earth sciences and are

vital cultural centers with a worldwide distribution. Their primary mandate – largely established

when advances in seafaring technology catalyzed worldwide exploration and the need to build

repositories to store materials for future study – is to house valuable specimens and artifacts

collected from across the globe. Therefore, these institutions form a foundation for diverse

research areas in biology, geology, and other disciplines, but have also adopted the mission of

championing education and public outreach in the natural sciences. Research and education in

natural history museums has been generally influential in addressing diverse societal issues,

including combating global change, human disease, and food insecurity, and bolstering

environmental conservation, sustainable harvest of natural resources, and the emerging

bioeconomy (National Academies of Sciences, Engineering, and Medicine 2020). More directly,

research leveraging natural history museums has illuminated the link between El Niño-driven

precipitation, increased density of rodents in the US Southwest, and heightened risk for human

contraction of Hantavirus Pulmonary Syndrome in this region (Yates et al. 2002), elucidated the

impact of a century of climate change on altitudinal distributions and ecologically-relevant

genetic diversity of small mammal populations in Yosemite National Park, California, USA

(Moritz et al. 2008; Bi et al. 2019), and inspired engineering solutions for noise reduction in

Japanese high-speed trains based on owl and kingfisher morphology (McKeag 2012).

Considering the importance of natural history museums for modern research, education,

and outreach, one would expect that museum holdings would be enumerated and well

understood. However, unfortunately, a complete accounting of the collections of natural history

museums does not exist, leaving scientists with largely anecdotal information about the sizes and

compositions of the world's natural history museums. Only recently have estimates of worldwide

collection holdings been amassed, after years of dedicated collection digitization that have

resulted in only about 30% of U.S. natural history samples being digitized and made available in

electronic databases (National Academies of Sciences, Engineering, and Medicine 2020). Since

2004, the Global Biodiversity Information Facility (GBIF) has aggregated billions of organism

occurrence records, including, as of 2023, over 200 million occurrences linked to preserved

specimens stored in natural history museums, herbaria, and other biorepositories (GBIF.org

2023). Recently, a targeted survey of 73 of the world's largest natural history museums and

herbaria located in 28 countries determined that there are more than 1.1 billion specimens or

artifacts in natural history collections, which were composed of 19 collection types sourced from

16 global geographic areas (Johnson et al. 2023). While significant work remains in digitizing

and integrating natural history collections data worldwide, this survey is the first to use full data

from the largest museums to estimate the minimum size of worldwide collections holdings,

making it critically important for deciding on future research funding levels, infrastructure

investments, and workforce development needs.

Estimates of the sizes of natural history museum collections are valuable for creating

policy that helps ensure that the infrastructure and personnel needs of natural history museums

are being met. However, collection size data comprises contributions to natural history museums

over their entire history, providing relatively coarse understanding of overall collections activity.

Estimates of collection sizes are valuable for understanding the scale and cost of curating

existing collections into the future but provides essentially no information about temporal

patterns of collecting activities across various institutions and taxonomic groups, or how such

activity may continue. Information on temporal collecting activity over the combined and

individual histories of natural history museums provides a much richer picture of an institution's

status that is theoretically more useful for making policy decisions. For example, funding needs

may differ between natural history museums that maintain high collecting activity and growth

versus historically more active institutions that are now growing much more slowly, or even

between collections with different taxonomic or geographic foci with distinct collection and

curation considerations. Moreover, understanding global patterns of collections growth across

natural history collections is vital for assessing the health of natural history museum collections,

which would ideally have sustainable collecting activity over time, and the utility of these

resources for addressing a range of biological questions.

While data-informed estimates of the sizes of natural history museum collections are

beginning to emerge, there is little publicly available information on the sizes of natural history

collections that can be used to understand temporal patterns of collecting activity and museum

growth. Using data from GBIF for vertebrates, Rohwer et al. (2022) found declines in the

deposition of vertebrate specimens to natural history collections since peaks in collection activity

that occurred in the 1970s and 1980s, depending on the taxonomic class. Collecting trends in

Western Hemisphere mammal collections have been periodically assessed by the American

Society of Mammalogists, most recently around 2018, with noticeable declines in collecting

activity observed in recent years (Dunnum et al. 2018; Malaney & Cook 2018; Cook & Light

2019). In contrast, a recent evaluation of collections holdings and digitization efforts of North

American arthropod collections found that the number of specimens in collections has increased

by approximately 1% annually over 30 years, although the authors note that this rate of increase

is insufficient for addressing many biodiversity research needs (Cobb et al. 2019). Moreover,

data from five major U.S. natural history museums indicated recent growth of cryogenetic

collections, a more recent form of collection that includes important materials for molecular

investigations comprising the emerging field of museum genomics (Card et al. 2021). However,

for most natural history museums, individual collections, time periods, and taxonomic groups,

information on collecting activity and collection growth is lacking, leaving natural history

collections poorly equipped to assess curatorial costs of existing collections, predict future

collecting targets and trends, and advocate for sustainable levels of financial and infrastructure

support. Anecdotal evidence also suggests that collecting activity at natural history museums has

declined in recent years, although no published studies have confirmed whether this is true.

119   To begin addressing this knowledge gap, I filtered and standardized global, publicly

available records for animals derived from preserved specimens in natural history collections.

Based on these data, I evaluate historical patterns of collecting activity across well-represented

taxa in the largest natural history museums with catalog data available through GBIF. I also

describe a new informatics dashboard built upon these filtered, standardized museum specimen

records that will facilitate additional investigations of these data by institutions and individuals

across the globe. After characterizing global collection trends and the temporal accumulation of

zoological specimens across different nations, institutions, and taxonomic units, I conclude with

several recommendations for ensuring sustainable growth and general vitality of natural history

museum collections.

Materials & Methods

130   Publicly available records for animals deposited into natural history collections were

retrieved from GBIF (N = 99,513,214 records; retrieved 2023-01-31;

https://doi.org/10.15468/dl.uxhkny). These records represent the subset of natural history

museum holdings that have been digitized and required filtering and standardization for

comparison and visualization. I filtered the records as follows: (1) records where the Kingdom

was not 'Animalia' were removed; (2) records not identified to the level of genus or below were

removed; and (3) records with missing or undefined data for year, institution code, or collection

catalog number were removed (N = 61,248,185 records retained). I summarized the number of

records for each taxonomic phylum and class, collection code, and year, associated each

individual collection code with a parent institution or publisher (i.e., some institutions publish

subcollections data to GBIF separately), and focused on records with a defined taxonomic

phylum and class that were collected between 1900 and 2015. To avoid potential biases in

taxonomic clades and institutions with small numbers of records, I focused on well-represented

phyla and institutions by selecting phyla with at least 5,000 records worldwide and institutions

with a minimum of 100,000 records amassed between 1900 and 2015. I also ensured that phyla

were consistently collected between 1900 and 2015 (at least 100 records per year for at least 50

years) and institutions had relatively consistent collecting activity between 1900 and 2015 (at

least 100 records per year for at least 50 years). 66.6% of records (N = 40,816,928) were retained

after applying these filters. Given that the number of records for collections and taxonomic

groups can differ significantly, making comparison and visualization difficult, I standardized the

number of records for each institution and year by the total number of records amassed per

institution between 1900 and 2015, producing an annual measure of collecting activity that is a

proportion of the institutional collection growth over the 1900 to 2015 timespan. Using the

resulting dataset, I assessed overall global collections growth and temporal, institutional, and

taxonomic patterns of collecting activity between 1900 and 2015. I also used Shiny v. 1.7.4

(Chang et al. 2023) in R v. 4.2.3 (R Core Team 2023) to build a supplemental interactive

dashboard that will allow others to explore natural history collections growth.

Results

The 40,816,928 zoological records surviving quality filtering comprised 72 taxonomic

classes and 10 phyla and represent the holdings of 102 institutions from 21 countries and all

inhabited continents (Australia was included with Oceania). Although the total and relative size

of natural history museums was specifically addressed in Johnson et al. (2023), my data provided

similar, although less complete, information on the relative sizes of the largest institutions, which

were concentrated in Europe and, especially, North America. In the following sections, I briefly

describe major patterns of historical.

Global zoological collections growth has peaked

The most glaring and unfortunate result of investigating historical patterns of zoological

collection growth is that global zoological collections growth has peaked and is currently in

steady decline (Figure 1). Collecting activity was relatively modest in the early decades of the

20th century before noticeable growth beginning around 1920 that continued through the late

1930s. World War II had a marked, negative impact on collecting activity in the 1940s and it was

not until the around 1950 that collections growth matched pre-war levels. Beginning in the late

1950s and continuing until about 1965, there was significant growth of natural history

collections. Growth then plateaued or modestly declined through the late 1980s before another

era of substantial growth that extended through the early 2000s, which represents the period

when collecting activity and growth of natural history collections globally was greatest. Since

this peak, activity has declined precipitously – at a rate similar to declines observed during

World War II – through 2015, the final year evaluated (Figure 1). Activity likely remains

diminished to date, although I avoided assessing patterns after 2015 due to potentially

incomplete data from recent years. Annual collections growth in 2015, and presumedly today,

matches growth in around 1960, which is the approximate start of an era of unprecedented

growth during the second half of the 20th century. Therefore, investments and concerted effort

are needed to prevent collections growth from falling to levels not seen since before the modern

era of science research that began in the wake of World War II.

Domain-specific patterns of collections growth provides nuanced understanding of collecting

activity

In much the same way that temporal patterns of collecting activity provides richer

information than total collection sizes alone, assessing patterns of collections growth for

individual taxonomic units, natural history museums, and nations leads to even deeper

understanding of the health and vitality of global biodiversity collection activity and the

biorepositories where specimens are stored. My analysis of taxonomic, national, and institutional

patterns of zoological collecting activity between 1900 and 2015 indicates biased accumulation

of taxa and varying national and institutional growth patterns. At the level of phylum,

standardized collecting activity indicates that most taxa have been most heavily collected since

1950, although spikes in collecting activity for Platyhelminthes and Nematoda occurred around

1900 and the 1930s saw an untick in the rate of collection for Bryozoa (Figure 2A). Based on

raw number of records, Arthropoda and Chordata are by far the most collected phyla (at least

100,000 records for each 5-year period since 1900), although Mollusca, Cnidaria,

Echinodermata, and Annelida have also been collected in relatively high numbers (at least

10,000 records for each 5-year period since ~1960; Figure 3A). Porifera, specifically, shows a

different pattern where collecting activity is concentrated in the period since 1980. Collecting

patterns in Arthropoda and Chordata drive much of the overall temporal trends across taxa

(~55% and ~34% of all records, respectively), and within these phyla, certain taxonomic classes,

namely Insecta and various vertebrates, represent most of the collecting activity (Figure 2A and

Figure 3A). Collecting activity focused on Chordata peaked in the 1960s and was relatively

healthy into the 1990s, but has since waned significantly, especially in the case of amphibians

and reptiles, mirroring vertebrate-specific results presented in Rohwer et al. (2022). Vertebrates

are the most collected chordates and the collecting trends of each vertebrate taxonomic class

generally mirror overall patterns in Chordata with high collecting activity in the 1960s through

the 1990s, except in the case of Aves, which were collected in relatively high numbers only in

the 1960s, but also exhibited a burst in collecting activity far earlier than other vertebrates (1905-

1940; Figure 2A). Aves is also the only taxonomic class outside of Insecta where at least 100,000

specimens were collected during each 5-year period between 1900 and 2015 (Figure 3A),

although consistently large collections of mammals have accumulated since 1930 and gastropod

molluscs have been well represented in collections since 1960. In contrast, collecting activity in

arthropods has continued to grow over time, but has leveled off in recent years. Consistent

collections of insects dominate these trends, although Malacostraca was relatively consistently

collected between 1900 and 2015 and Arachnida and Copepoda are characterized by relatively

high collecting activity after 1990 (Figure 3A). Indeed, if not for significant growth in collecting

activity in Arthropoda beginning around 1990, which is well justified by the proportions of

known and unknown biodiversity in this clade, overall collections growth likely would have

peaked much earlier in the 1960s and the subsequent decline in activity would have been far

more precipitous (Figures 1 and 2A).

Given that national zoological collecting trends, in most cases, integrate over multiple

collections, they are relatively homogenous, with most records accumulating since about 1960,

although several European nations (Austria, Belgium, Denmark, and Estonia) had bursts of

relatively high collecting activity between 1920 and 1960. In contrast, collecting activity in

Mexico has been concentrated in the period since 1980 (Figure 2B). Collections growth patterns

in the United States dominate overall trends due to the concentration of large natural history

museums (~51.5% of records are from U.S. museums). At least 100,000 records have been

amassed every 5-year period since 1900 and since 1960 and the U.S. has accumulated more than

1 million records over most 5-year periods. Only Australia, Japan, the United Kingdom, Mexico,

Canada, and Sweden have had multi-decade sustained growth of collections exceeding 100,000

records per 5-year period, and Norway, Brazil, and Poland have also had periods of relatively

high, though less sustained, collections growth (Figure 3B). On the other hand, individual

institutions show more marked differences in the accumulation of natural history records.

Collecting activity was relatively high before 1950 for certain collections, such as the American

Museum of Natural History, Natural History Museum London, University of Michigan Museum

of Vertebrates, Field Museum, Museum of Comparative Zoology at Harvard University, Illinois

Natural History Survey, Museum of Vertebrate Zoology, and Royal Belgian Institute of Natural

Sciences (Figure 2C). Most other institutions had periods of relatively high collecting activity

that began in the 1950s and, especially, the 1960s, which corresponded with the significant

overall growth of natural history collections in this period. For many of these institutions, these

periods of relatively high collecting activity have persisted beyond 1970 and, in some cases, to

the final period evaluated in this study, such as the National Museum of Nature and Science –

Japan, University of Kansas Biodiversity Institute, the Marine Biological Association, the

Florida Museum of Natural History, Adam Mickiewicz University in Poznan, and the Muséum

National d'Histoire Naturelle (Figure 2C). Other institutions – namely the Comisión National

para el Conocimientio y Uso de la Biodiversidad (Mexico), California Academy of Sciences,

Swiss National Biodiversity Data and Information Centres, SLU Artdatabanken, and Queensland

Museum – have had most of their collection growth occur since 1970 (Figure 2C). However,

based on raw record counts, only a small number of the largest institutions had relatively

consistent, high collecting activity (at least 100,000 records) for at least two 5-year periods

between 1900 and 2015: National Museum of Natural History – Smithsonian Institute, National

Museum of Nature and Science Japan, Comisión National para el Conocimientio y Uso de la

Biodiversidad (Mexico), University of Kansas Biodiversity Institute, California Academy of

Sciences, Australian Museum, Marine Biological Association, and Texas A&M University

Insect Collection (Figure 3C). National and institutional collection trends underscore how

variable zoological collecting activity can be and provide richer understanding of the health of

global biorepositories individually and collectively.

An interactive R Shiny application for exploring patterns of collections growth

The FAIR Data Principles (Wilkinson et al. 2016) of ensuring that data are **F**indable,

Accessible, **I**nteroperable, and **R**eusable are increasingly important in the current era of "big

data". Although natural history museums remain incompletely digitized (National Academies of

Sciences, Engineering, and Medicine 2020) and have only recently begun to integrate data

summarizing their vast collective holdings (e.g., Johnson et al. 2023), in many ways, these

institutions were one of the pioneers of the FAIR Data Principles, at least in the non-digital

realm. To continue this tradition and following the recent example of Johnson et al. (2023), I

used R Shiny to build a simple, interactive application that will allow anyone to explore the

filtered, standardized data that have been summarized in this article. This application has been

provided as an R Markdown document and contains a summary of the filtered, standardized data

and embedded Shiny elements that facilitate interactive exploration of these data according to

user interests. The document also includes a detailed supplement with the full details – including

embedded code, plots, and tables – of retrieving, filtering, and standardizing the raw GBIF data,

an overview of the full exploratory analyses and visualization that formed the basis for this

investigation, and all code and documentation necessary to reproduce the analyses and figures

presented in this article. The resources produced as part this article have been made permanently

available through a Zenodo-archived repository (https://doi.org/10.5281/zenodo.8393331), which

contains a partially filtered intermediate dataset used for data summarization and exploration in

R, a final dataset summarizing the records of different taxonomic groups and institutions, and an

R Markdown document with embedded Shiny elements that enables user exploration of the data.

In the spirit of FAIR, interested users will always be able to download the permanently archived

repository and use R to run the R Markdown document to interactively explore the data on their

local computer. Additionally, instructions for accessing an internet-hosted version of the R

Markdown document are available at https://github.com/darencard/NHMinformatics to allow

interested users to explore the collections data more easily. Any future updates to these resources

will be made available through the Zenodo-linked GitHub repository

(https://github.com/darencard/NHMinformatics).

Discussion

Using data from over 40 million cataloged zoological records from global natural history

museums, I provided the first published summary of temporal collections growth based on

records from 10 well-collected phyla and the 102 largest institutions who have contributed data

to GBIF. The growth of these natural history collections has varied significantly over time,

reflecting distinct eras of global natural history collecting activity. Beginning in 1900, the first

year evaluated in this study, natural history museums saw approximately 70 years of consistent

growth only interrupted by a sharp downturn and subsequent recovery during and after World

War II. This pattern is a marked departure from the trends observed since around 1970, and

following the peak of collections growth in 2000-2001, collecting activity has fallen

precipitously. Variable patterns of collecting activity across nations, institutions, and taxonomic

clades underly these complex temporal trends and highlight the unique contributions individual

institutions have made to global natural history museum holdings. Inference of these patterns

relied upon rigorous filtering and standardization that resulted in over 40 million records,

encompassing institutions that are known to be among the largest in the world, which provides

confidence that the true temporal patterns in global collecting activity have been captured.

However, 73 of the world's largest natural history museums and herbaria house more than 1.1

billion specimens and artifacts (including plant specimens, fossils, and cultural artifacts not

assessed here; Johnson et al. 2023), reflecting the incomplete availability of natural history

collection catalog data in GBIF, which suggests that further data and research are needed to

precisely understand global collections growth. Although interesting for a variety of reasons, I

also ignored the source locations for individual records when aggregating the data due to

difficulties validating coordinates, inferring coordinates based on locality descriptions, and

associating records with a nation or region. But anecdotal information and published data

(Johnson et al. 2023) indicate that the source nations of museum specimens are less concentrated

and geographically biased than the institutions that house them, which stems largely from

historical practices during the colonial era that persist today, presenting unfortunate obstacles for

nations and populations wishing to derive cultural, intellectual, and economic benefits from their

natural heritage (Sheets-Pyenson 1987; Das & Lowe 2018; Ashby & Machin 2021; Nicolaï

2022). Overall, given that GBIF represents science's best effort towards natural history museum

collection digitization and integration, this research provides a meaningful, early representation

of these important data trends that will form an important foundation for future collection data

integration and analysis. Indeed, by making an interactive R Shiny application available for

others to interactively explore museum collections growth trends, I continue a long tradition in

natural history museum science of making data or materials publicly available for reuse.

Although it is now possible to summarize specimen collecting trends across individual countries,

institutions, and taxonomic clades – which should assist individual institutions and government

institutions or interest groups in defining strategic collections priorities – the global trend of

declining contributions to natural history museum collections uncovered in this study is

concerning and undermines the mission of these important institutions.

Diverse options are available to overcome the concerning trends evident in my temporal

analysis of natural history collections growth, and I will make several higher-level

recommendations for how we can ensure sustainable growth of natural history collections.

Natural history collection holdings continue to be digitized and amassed in data portals like

GBIF, but until now, there has been no central dashboard summarizing natural history collecting

activity across taxonomic groups, nations, or institutions. Concerted effort is needed to improve

digitization of individual collections and aggregate these data, but more specifically, natural

history museums must invest in infrastructure and personnel that allows them to informatically

explore existing data, as I have done here (see also Johnson et al. [2023]), to better guide data

digitization and aggregation and new specimen and artifact collection. Armed with such real-

time information, governments, funding agencies, and individual institutions should consider

implementing collecting quotas or recommendations for collecting that ensures sustainable

growth, which should be informed both by our understanding of biodiversity across different

taxonomic clades and our knowledge of the geographic or taxonomic specializations of

institutions or nations. Today, most collecting activity is driven by the interests of individual

curators and the funding they secure to pursue individual research programs. However, natural

history collections are emerging as critical infrastructure for diverse areas of research beyond

systematic investigations of the tree of life, which has historically been the focus of many

museum practitioners. This trend, plus knowledge of best practices in experimental design that

dictate the utility of natural history collections for experimental science, strongly argues for

prescribed minimum collecting activity that is coordinated across various hierarchical levels.

Finally, funding levels for natural history collections need to be increased to ensure continued

curation of existing holdings and sustainable growth through new collecting initiatives. While

my data and analyses are insufficient to assess whether funding levels and priorities have

contributed to diminished collections growth, documented closures and personnel layoffs

(Dunnum et al. 2018) provide anecdotal evidence that financial austerity has negatively impacted

natural history museums in recent years. Without a recommitment of funding agencies

(governmental and philanthropic) and individual institutions to the mission of natural history

museums and biorepositories through increased and sustainable funding, we risk diminished

utility of natural history collections at a time where they are increasingly valuable for the

scientific enterprise in the life sciences and beyond.

Conclusions

This study presents the first detailed investigation of temporal patterns of global

zoological natural history collections growth between 1900 and 2015. Collecting activity varied

greatly over time, both globally and based on individual patterns of collections, nations, and

taxonomic clades, and peaked around 2000, with recent years characterized by precipitous

declines in collections growth. I also released an interactive data dashboard built using R Shiny

to allow natural history museum stakeholders to explore these historical collections growth

patterns in greater detail. In response to recent downward trends in collecting activity, I

recommend that (1) institutions and other stakeholders invest more in informatic exploration of

natural history museum holdings to guide future collecting, digitization, and aggregation efforts,

(2) governments, funding agencies, and individual institutions implement collecting quotas or

recommendations to ensure consistent, sustainable collecting activity in the future, and (3)

government and philanthropic funding agencies increase financial commitments to natural

history collections to catalyze more consistent future collections growth.

Acknowledgements

I am grateful for constructive comments from Scott Edwards, Michaël Nicolaï, and anonymous

reviewers. I also acknowledge Vanya Rohwer and Matt Strimas-Mackey for advice on gathering

natural history collection data from GBIF and Rhett Rautsaw for advice on building an

interactive data dashboard in R Shiny. Postdoctoral support for D.C.C. has come from a

fellowship from the United States National Science Foundation (DEB-1812310). The

computations in this paper were run on the FASRC Cannon cluster supported by the FAS

Division of Science Research Computing Group at Harvard University.

References

References

Ashby J, Machin R. 2021. Legacies of colonial violence in natural history collections. J. Nat. Sci.
Collect. 8:44–54.

Bi K et al. 2019. Temporal genomic contrasts reveal rapid evolutionary responses in an alpine
mammal during recent climate change. PLOS Genet. 15:e1008119. doi:
10.1371/journal.pgen.1008119.

Card DC, Shapiro B, Giribet G, Moritz C, Edwards SV. 2021. Museum genomics. Annu. Rev.
Genet. 55:633–659. doi: 10.1146/annurev-genet-071719-020506.

Chang W et al. 2023. *shiny: Web application framework for R*.

Cobb NS et al. 2019. Assessment of North American arthropod collections: prospects and
challenges for addressing biodiversity research. PeerJ. 7:e8086. doi: 10.7717/peerj.8086.

Cook JA, Light JE. 2019. The emerging role of mammal collections in 21st century mammalogy.
J. Mammal. 100:733–750. doi: 10.1093/jmammal/gyy148.

Das S, Lowe M. 2018. Nature Read in Black and White: decolonial approaches to interpreting
natural history collections. J. Nat. Sci. Collect. 4–14.

Dunnum JL, McLean BS, Dowler RC, Systematic Collections Committee of the American
Society of Mammalogists. 2018. Mammal collections of the Western Hemisphere: a survey and
directory of collections. J. Mammal. 99:1307–1322. doi: 10.1093/jmammal/gyy151.

GBIF.org. 2023. GBIF Occurrences Page. https://www.gbif.org/occurrence/ (Accessed January
31, 2023).

Johnson KR, Owens IFP, The Global Collection Group. 2023. A global approach for natural
history museum collections. Science. 379:1192–1194. doi: 10.1126/science.adf6434.

Malaney JL, Cook JA. 2018. A perfect storm for mammalogy: declining sample availability in a
period of rapid environmental degradation. J. Mammal. 99:773–788. doi:
10.1093/jmammal/gyy082.

McKeag T. 2012. Auspicious Forms: Designing the Sanyo Shinkansen 500-Series Bullet Train.
Zygote Quarterly.

Moritz C et al. 2008. Impact of a Century of Climate Change on Small-Mammal Communities in
Yosemite National Park, USA. Science. 322:261–264. doi: 10.1126/science.1163428.

National Academies of Sciences, Engineering, and Medicine. 2020. *Biological Collections:*
Ensuring Critical Research and Education for the 21st Century. The National Academies Press:
Washington, DC doi: 10.17226/25592.

Nicolaï MPJ. 2022. Boost funds to make museum specimens available to all. Nature. 612:631–
631. doi: 10.1038/d41586-022-04473-y.

R Core Team. 2023. *R: A language and environment for statistical computing*. R Foundation for
Statistical Computing: Vienna, Austria http://www.R-project.org/.

Rohwer VG, Rohwer Y, Dillman CB. 2022. Declining growth of natural history collections fails
future generations. PLOS Biol. 20:e3001613. doi: 10.1371/journal.pbio.3001613.

Sheets-Pyenson S. 1987. Cathedrals of Science: The Development of Colonial Natural History
Museums during the Late Nineteenth Century. Hist. Sci. 25:279–300. doi:
10.1177/007327538702500303.

Wilkinson MD et al. 2016. The FAIR Guiding Principles for scientific data management and
stewardship. Sci. Data. 3:160018. doi: 10.1038/sdata.2016.18.

Yates TL et al. 2002. The Ecology and Evolutionary History of an Emergent Disease: Hantavirus
Pulmonary Syndrome. BioScience. 52:989–998. doi: 10.1641/0006-
3568(2002)052[0989:TEAEHO]2.0.CO;2.

Figures

Figure 1. Assessment of historical growth of natural history collections indicates a peak and

subsequent decline in collecting activity. Temporal patterns of the number of specimens

deposited in worldwide natural history collections between 1900 and 2015, as based on data

from the Global Biodiversity Information Facility (GBIF). Annual activity is reported as a

percentage of total collections made across all clades and institutions over the entire time period

after rigorous data filtering.

Figure 2. Idiosyncratic patterns of historical natural history collections growth across

taxonomic clades, nations, and natural history museums/institutions. Sliding window

measures of standardized collecting activity over 5-year intervals for (**A**) consistently-collected

taxonomic phyla (see above) and classes (at least 100,000 global records) as a percentage of total

global collections for each clade between 1900 and 2015, (**B**) 21 nations with at least one

museum/institution as a percentage of total collections for each nation between 1900 and 2015,

443   and (**C**) 26 of the largest natural history museums/institutions with total collection sizes greater

444   than 500,000 records each as a percentage of total collections for each museum/institution

445   between 1900 and 2015.

Figure 3. Variable growth of natural history collections records for different taxonomic clades, nations, and natural history museums/institutions. Sliding window measures of collecting activity over 5-year intervals based on absolute counts of records of (**A**) consistently-collected taxonomic phyla (see above) and classes (at least 100,000 global records), (**B**) 21 nations with at least one museum/institution, and (**C**) 26 of the largest natural history museums/institutions with total collection sizes greater than 500,000 records each.