***Salmonella* Typhimurium employ spermidine to exert protection against ROS-**

mediated cytotoxicity and rewires host polyamine metabolism to ameliorate its

survival in macrophages

Abhilash Vijay Nair[a], Anmol Singh[a], R. S. Rajmani[b] , Dipshikha Chakravortty[a,c]#

[a]Department of Microbiology and Cell Biology, Division of Biological Sciences, Indian Institute

of Science, Bengaluru, India

[b]Molecular Biophysiscs Unit, Indian Institute of Science, Bangalore, India

[c]Adjunct Faculty, School of Biology, Indian Institute of Science Education and Research,

Thiruvananthapuram

Running Head: *Salmonella* mounts an antioxidative response using de-novo and host-acquired

spermidine

#Address correspondence to Dipshikha Chakravortty, dipa@ iisc.ac.in, Tel: 0091 80 2293 2842,

Fax: 0091 80 2360 269

Keywords: Spermidine, Macrophages, Antioxidative response, Glutathionyl-spermidine

synthetase, D,L-α-difluoromethylornithine

19
20

Abstract

Salmonella infection involves a cascade of attacks and defence measures. After breaching the intestinal epithelial barrier, *Salmonella* is phagocytosed by the macrophages, inside which, the bacteria face multiple stresses and, consequently, employ appropriate countermeasures. We show that, in *Salmonella*, the polyamine spermidine activates a stress response mechanism by regulating critical antioxidant genes. *Salmonella* Typhimurium mutants for spermidine transport and synthesis cannot mount an antioxidative response, resulting in high intracellular ROS levels. These mutants are also compromised in their ability to be phagocytosed by macrophages. Furthermore, it regulates a novel enzyme in *Salmonella*, Glutathionyl-spermidine synthetase (GspSA), which is known to prevent the oxidation of proteins in *E.coli*. Moreover, the spermidine mutants and the GspSA mutant show significantly reduced survival in the presence of hydrogen peroxide *in vitro*, and lesser organ burden in the mouse model of *Salmonella* infection. Conversely, in macrophages isolated from *gp91phox*[-/-] mice, we observed a rescue in the attenuated fold proliferation previously observed upon infection. Interestingly, *Salmonella* upregulates polyamine biosynthesis in the host, which also solves the mystery of the attenuated proliferation observed in spermidine transport mutants. Thus, inhibition of this pathway in the host abrogates the proliferation of *Salmonella* Typhimurium in macrophages. From a therapeutic perspective, inhibiting host polyamine biosynthesis using an FDA-approved chemopreventive drug, D,L-α-difluoromethylornithine (DFMO), reduces *Salmonella* colonization and tissue damage in the mouse model of infection, while enhancing the survival of infected mice. Therefore, our work provides a mechanistic insight into the critical role of spermidine in stress resistance of *Salmonella*. It also reveals a strategy of the bacteria in modulating host metabolism to promote their intracellular survival and shows the potential of DFMO to curb S*almonella* infection.

Significance

A substantial duration of the infection cycle of *Salmonella* involves the macrophages, which present a very hostile environment to the bacteria. However, *Salmonella* is able to survive and proliferate within host macrophages and utilizes it to disseminate to secondary sites of infection. Our study identifies a novel strategy employed by *Salmonella* Typhimurium to counteract oxidative and nitrosative stress within the host. We demonstrate that spermidine is a critical regulatory molecule in *Salmonella* that regulates multiple antioxidative pathways along with a novel antioxidative enzyme (GspSA) in *Salmonella,* to prevent oxidative damage and assist in its virulence in mice. It further rewires host polyamine metabolism to prevent nitric oxide production and enhance its survival. In the era of anti-microbial resistance, this study further recognizes an FDA -approved chemopreventive drug, DFMO, which inhibits the host-polyamine metabolism, as a prospective antidote to treat *Salmonella* infection.

Introduction

A condition with ample nutrients, and desired temperature, pH, oxygen concentration, and osmolarity is usually considered optimal for the growth of microbes. An imbalance or any alteration in these parameters impedes the growth and survival of the microbes, considered as stress conditions. However, these parameters keep fluctuating in nature [1]. Hence, to persist and survive in the natural environment, with unforeseen stressful or lethal conditions, they must be adept in sensing, responding and adapting to them [2]. In the case of food-borne pathogens, they face an array of various stresses in multiple environments in which they dwell, such as from the natural habitat to commercial settings and inside the host system [3-5]. *Salmonella* is a food-borne pathogen that lives in aquatic bodies and soil, and enters the host system through contaminated food and water. In the host system, as it traverses to the intestine, *Salmonella* faces multiple

stressful conditions such as low pH, nutrient deprivation, bile salt stress, competition with the resident microbes of the gut and immunoglobulins, etc. Once it breaches the epithelial barrier, it is taken up by the macrophages at the lamina propria, through which it disseminates throughout the host system. Macrophages are phagocytic immune cells where *Salmonella* encounters a very hostile environment. Entry of the pathogen into the cell cytoplasm leads to a burst of reactive oxygen species (ROS) and reactive nitrogen species (RNS). In macrophages, *Salmonella* resides in a specialised niche called the *Salmonella*-containing vacuole (SCV), which presents multiple other stresses to the bacteria, such as acidification, nutrient limitation and attack by the antimicrobial peptides. However, *Salmonella* employs numerous weapons from its arsenal to counteract the stresses it faces within the host macrophages.

Polyamines are a group of primordial stress response molecules in prokaryotes [6]. Multiple research groups have elucidated the link between polyamines and bacterial stress response. In *Shigella spp.* the polyamine spermidine accumulates during its infection into macrophages, which increases the expression of *katG* and helps in bacterial antioxidant response [7]. Moreover, spermidine localises to the surface of *Pseudomonas aeruginosa* and bind to the lipopolysaccharide (LPS) to protect the cells from oxidative damage [8]. In the Gram-positive bacteria, *Streptococcus pyogenes,* extracellular spermidine enhance the survival of the bacteria by upregulating oxidative response genes [9]. A research group has shown that polyamines are vital in resistance against nitrosative stress in *Salmonella* Typhimurium. Further, the group showed that spermidine is required for the systemic infection of *Salmonella* Typhimurium in mice [10]. Thus, it is conceivable that *Salmonella* Typhimurium utilises polyamines, such as spermidine, as a stress response molecule; however, the mechanism remains elusive.

Here, we show that spermidine biosynthesis and transport mutants of *Salmonella* Typhimurium

exhibit reduced survival upon infection in RAW264.7 cells. This diminished proliferation is also

observed in mice models of *Salmonella* infection, which is rescued in *gp91phox*[-/-] mice. We

demonstrate that spermidine orchestrates the various arms of antioxidative response and aids in

tightly regulating intracellular ROS levels. We further identify a novel antioxidative enzyme, Gsp,

in *Salmonella* Typhimurium, which is regulated by spermidine. The fascinating question that arises

is why the transporter mutant shows reduced survival. To this we find that *Salmonella*

Typhimurium harnesses the host polyamine biosynthesis for its survival. Furthermore, for the first

time, we show that an FDA-approved chemopreventive and anti-African Human Trypanosomiasis

drug that inhibit the polyamine biosynthesis in the host, is able to curb *Salmonella* infection in

mice models.

Material and methods

Bacterial strains and growth condition

Salmonella enterica serovars Typhimurium (STM WT) wild type strain ATCC 14028 was used in

all experiments which was a kind gift from Prof. Michael Hensel, Abteilung Mikrobiologie,

Universität Osnabrück, 273 Osnabrück, Germany. The bacterial strain was cultured in Luria broth

(LB-Himedia) with constant shaking (175rpm) at 37°C orbital-shaker. Kanamycin,

Chloramphenicol and Ampicillin antibiotics (Kanamycin-50μg/ml, Cholramphenicol-20μg/ml and

Ampicillin-50μg/ml) were used wherever required. (Bacterial strain list in Supplementary table **S-**

Table 1)

Bacterial gene knock-out and strain generation

The generation of gene knock-out in bacteria was done using the One-step chromosomal gene

inactivation protocol [11]. Briefly, the Kanamycin resistance gene and the Chloramphenicol

resistance gene   amplified products were purified using chloroform-ethanol precipitation.

Followed by electroporation into the STM WT cells (expressing PKD-46 plasmid which provides

the λ-Red recombinase system) by a single pulse of 2.25 kV separately for the Kan^R and $Chlm^R$.

The transformant colonies were selected and patched on fresh plates and confirmed for knock-out

using PCR with primers designed corresponding to the ~100bp upstream and downstream of the

genes (knocked out) for the knock-out strains to observe a difference in PCR product size upon

STM *ΔpotCD* and STM *ΔspeED* knockout genration. For the generation of the double knock-out

strain (STM *ΔpotCDΔspeED*), the STM *ΔpotCD* (resistant to Kanamycin) was first transformed

with the plasmid pKD46 which provides the λ-Red recombinase system. To this transformed strain,

the purified PCR product to knock-out *speED* was electoporated to generate the STM

ΔpotCDΔspeED (resistant to Kanamycin and Chloramphenicol)*.* For the generation of STM *Δgsp,*

Kanamycin resistance gene was amplified from pKD4 plasmid, and a similar protocol was used,

followed by selection on Kanamycin containing LB agar plates. (Knockout generation Primer list

in Supplementary table **S-Table 2**)

Cell culture and maintenance

129   RAW264.7 cells (murine macrophage cell line) were cultured in DMEM - Dulbecco's Modified

130   Eagle Medium (Lonza) supplemented with 10% FBS (Gibco) and 1% Penicillin-streptomycin

131   (Sigma- Aldrich) at 37°C humidified chamber (Panasonic) with 5% CO_2. For each experiment, the

132   cells were seeded onto the appropriate treated cell culture well plate at a confluency of 80% for

133   intracellular survival assay, and expression studies.

Gentamicin protection assay

The cells were infected with STM WT, STM *ΔpotCD,* STM *ΔspeED,* STM *ΔpotCDΔspeED,* STM

Δgsp and STM *ΔkatG* at MOI of 10 (for intracellular survival assay) and MOI 25 (for qRT-PCR).

After infecting the cell line with STM WT and the mutants, the plate was centrifuged at 700-900

rpm for 10 minutes to facilitate the proper adhesion. The plate was then incubated for 25 minutes

at 37°C humidified chamber and 5% CO_2. Then the media was removed from the wells and washed

with 1X PBS. Fresh media containing 100 µg/mL gentamicin was added and again incubated for

60 minutes at 37°C and 5% CO_2. The media was then removed, cells were washed with 1X PBS

twice, and fresh media containing 25µg/mL gentamicin was added. The plate was incubated at

37°C and 5% CO_2 till the appropriate time. For the intracellular survival assay, two time points

were considered 2 hours and 16 hours, and for qRT-PCR three time points were considered 2

hours, 6 hours and 16 hours. For phagocytosis assay upon opsonisation, all the strains were washed

with 1XPBS and incubated at 37°C with mouse complement sera for 1 hour before gentamicin

protection assay

Intracellular survival assay and phagocytosis assay

At the appropriate time post-infection, the cells were lysed using 0.1% Triton X followed by

addition of more 1X PBS and samples were collected. The collected samples were plated at the

required dilutions on LB agar plates and kept at 37°C. 12 hours post incubation the Colony forming

units (CFU) were enumerated for each plate.

The fold proliferation and invasion were determined as follows

Fold Proliferation = (CFU at 16 hours post-infection)/(CFU at 2 hours post-infection)

Percentage Phagocytosis = [(CFU at 2 hours post-infection)/(CFU of the Pre-inoculum)]×100

RNA isolation and qRT-PCR

RNA isolation was performed from infected cells after appropriate hours of infection with STM

WT, STM *ΔpotCD,* STM *ΔspeED* by RNA isolation was performed using TRIzol (Takara) reagent

according to manufactures' protocol RNA was quantified using Thermo-fischer scientific Nano

Drop followed by running on 2% agarose gel for checking the quality. For cDNA synthesis, first

DNase I treatment with 3μg of isolated RNA was done at 37°C for 60 minutes, which was then

stopped by heating at 65°C for 10 minutes. Then RNA (free of DNA) was subjected to Reverse

transcription using Random hexamer, 5X RT buffer, RT enzyme, dNTPs and DEPC treated water

at 37°C for 15 minutes, followed by heating at 85°C for 15 seconds. Quantitative real-time PCR

was done using SYBR green RT-PCR kit in BioRad qRT-PCR system. A 384 well plate with three

replicates for each sample was used. The expression levels of the gene of interest were measured

using specific RT primers. Gene expression levels were normalised to 16SrDNA primers of *S.*

Typhimurium. Gene expression levels of eukaryotic gene of interest were normalised to beta-actin

of mouse/human as required. For expression studies in bacteria grown in LB media, the bacterial

samples were harvested at 3 hours, 6 hours, 9 hours and 12 hours post subculture in fresh LB media

in 1:100 ratio and 1mM H_2O_2 was added to the broth, to study the gene expression in the presence

of oxidative stress. Then similar protocol was used to isolate total RNA using TRIzol (Takara)

reagent according to manufactures' protocol. (Expression Primer list in Supplementary table **S-**

Table 2)

Primary macrophages isolation and infection

Primary macrophages were isolated from C57BL/6 mice . The mice were intraperitoneally injected

with 8% Brewer's Thioglycolate. Five days post injection the primary macrophages were

aseptically isolated by injecting ice-cold 1xPBS into the peritoneal cavity the peritoneal lavage

was collected. Any residual erythrocytes were lysed using RBC lysis buffer, and the isolated cells

were maintained in complete RPMI 1640 media for further experiments.

Intracellular Reactive oxygen species determination using H2DCFDA staining

Overnight cultures were sub-cultured in fresh LB media. Once the cultures reached OD 0.1 then

10^8 CFU/ml of each strain was incubated with 10μM of 2',7'-dichlorodihydrofluorescein diacetate

(H2DCFDA) in 1xPBS (pH 7.2) at 37°C for 30 minutes. The bacterial cells were centrifuged and

the cells were resuspended in 1xPBS (pH 7.2) with Hydrogen peroxide of different concentrations,

and incubated 37°C (orbital shaker) for 2 hours. The samples were transferred to a 96 well ELISA

plate and fluorescence was determined in Tecan-ELISA plate reader (Ex- 490nm/ Em- 520nm).

Intracellular redox status determination

The STM WT, STM *ΔpotCD,* STM *ΔspeED* and STM *Δgsp* were transformed with pQE60-Grx1-

roGFP2 (a gift from Prof. Amit Singh, CIDR, IISc). Overnight cultures were sub-cultured in fresh

LB media. Once the cultures reached OD 0.1 then 10^8 CFU/ml of each strain was incubated with

Hydrogen peroxide of different concentrations in 1xPBS (pH 7.2) with, and incubated 37°C

(orbital shaker) for 2 hours. The samples were centrifuged and resuspended in fresh 1xPBS (pH

7.2). And tubes were analysed for GFP fluorescence at 408nm and 488nm respectively using BD-

FACS Verse flowcytometer (total 10,000 events for each sample).

In vitro **sensitivity assays**

Overnight cultures were sub-cultured in fresh LB media. Once the cultures reached OD 0.1 then

10^8 CFU/ml of each strain was incubated with Hydrogen peroxide or sodium nitrite of different

concentrations in 1xPBS(pH 7.2) (1xPBS of pH 5.4 was used for nitrite sensitivity assay) with,

and incubated 37°C (orbital shaker) for 2 hours. The samples were plated on SS agar to enumerate

the CFU and the percentage survival was determined as:

Percentage Survival : [CFU/ml for treated with H_2O_2/ $NaNO_2$]/ [CFU/ml for untreated] $\times 100$

Transfection for knockdown

RAW 264.7 cells were seeded at a confluency of 50-60% 12 hours prior to transfecting using either

PEI (1:2 -DNA: PEI). We used two different constructs for knock-down of Odc1, A7 and F8 and

both as a mixed construct (A7:A8 at 1:1 ratio) and similarly for Srm , E9 and F1 and both as a

mixed construct (E9:F1 at 1:1 ratio) from the Sigma Mission shRNA library. 400ng of plasmid

DNA/well (ratio 260/280 ~1.8- 1.9) was used for transfection in a 24well plate. Cells were then

incubated for 8 hours at 37°C in a humidified incubator with 5% CO2; after that, the media

containing transfecting DNA and reagents were removed, and cells were further incubated for 48

hours in complete media DMEM +10% FBS. Cells were harvested for further analysis or infected

with the required MOI. (shRNA sequence list in Supplementary table **S-Table 3**)

Immunofluorescence

After the appropriate incubation time, the media was removed, and the cells were washed with 1X

PBS. The cells were then fixed with 3.5% Paraformaldehyde for 10 minutes. The cells were then

washed with 1X PBS and incubated with the required primary antibody (anti- *Salmonella* LPS) in

a buffer containing 0.01% saponin and 2% BSA and incubated at room temperature for 45-60

minutes. The primary antibody was then removed by washing with 1X PBS and then incubated

with the appropriate secondary antibody tagged to a fluorochrome. The coverslips were then

washed with PBS and mounted on a clean glass slide using mounting media containing an anti-

fade reagent. The coverslips were sealed with clear nail polish and observed under the confocal

microscope. For studying histopathology samples Zeiss 880 microscope 40X oil immersion 2x

323   3x zoom was used.

Intracellular RNS determination

For determination of intracellular RNS, a cell permeable nitric-oxide probe, 4, 5-diaminofluorescein diacetate (DAF2A) was used. The protocol has been followed as Roy Chowdhury A et. al. [12]. Briefly, After 16 hours of infection of RAW264.7 cells(transiently knock down of *Odc1*) at MOI 10 with STM WT, the cells were incubated with fresh DMEM containing 5µM of DAF2DA. The cells were incubated at 37°C in a humidified incubator with 5% CO_2 for 30 minutes. The media containing dye was removed and the cells were washed with 1xPBS and the cells were acquired immediately for flow-cytometry (BD FACS Verse) (Ex-491nm/Em- 513nm).

Intracellular Glutathione determination

The intracellular reduced Glutathione (GSH) concentration was determined by modification of a pubmished protocol [13]. Briefly, a standard curve with known concentration of GSH was prepared. Reaction mixture for each contained 600µL of phosphate buffer (0.1M, pH7), 40µL of 0.4% w/v 5,5-dithiobis(2-nitrobenzoic acid) (DTNB, from Sigma), 100 µL of the standard solutions of GSH (0mM-1mM range) and autoclaved MilliQ water to make up the volume to 1000µL.The mixture was incubated at room temperature for 5 minutes and absorbance was measured at 412nm using tecan plate reader. The bacterial strains were subcultured in fresh LB media and grown till OD 0.1 (exponential phase) and washed with buffer (Tris, Sodium chloride and EDTA) and lysed using sonication. The supernatant after sonication was used as the sample for GSH detection. As previously described for standard curve. From the standard curve the intracellular concentrations were interpolated.

In silico analysis

The *in silico* protein structure determination was performed using the SWISS-MODEL software (https://swissmodel.expasy.org/), where we supplied the protein sequence of GspSA of *Salmonella* Typhimurium from UniProt. We analysed the model with highest sequence identity and maximum coverage with the Gsp from *E.coli*. The structure depicts a Homo-dimer with a GMQE of 0.93 and QMEANisCo Global of 0.88 ± 0.05.

In vivo animal experiment

5-6weeks old C57BL/6 mice were infected by orally gavaging 10^7 CFU of STM WT, STM *ΔpotCD,* STM *ΔspeED,* STM *ΔpotCDΔspeED,* STM *Δgsp* and STM *ΔkatG*. To study the colonisation in organs, the intestine (Peyer's patches), MLN, spleen and liver were isolated aseptically, 5 days post-infection, and the CFU was enumerated on differential and selective SS agar by serial dilution. For intraperitoneal infection 5-6weeks old C57BL/6 mice were infected by intraperitoneally injecting 10^3 CFU of STM WT, STM *ΔpotCD,* STM *ΔspeED,* STM *ΔpotCDΔspeED,* STM *Δgsp* and STM *ΔkatG.* To study the colonisation in organs, spleen and liver were isolated aseptically 3 days post-infection. Blood was isolated by heart puncture 3 days post-infection. The CFU was enumerated on differential and selective SS agar by serial dilution. Organs were stored in 3.5%PFA before histopathogical sample preparation.

For inhibitor treatment, 5-6weeks old C57BL/6 mice were infected by orally gavaging 10^7 CFU of STM WT. The inhibitor DFMO was intraperitoneally injected every alternate day from day 1 at two doses 2mg/kg and 1mg/kg of body weight of mice. To study the colonisation in organs, the intestine (Peyer's patches), MLN, spleen and liver were isolated aseptically, 5 days post-infection, and the CFU was enumerated on differential and selective SS agar by serial dilution. For survival assay of mice 5-6weeks old C57BL/6 mice were infected by orally gavaging 10^8 CFU of STM

WT. The inhibitor DFMO was intraperitoneally injected every alternate day from day 1 at two

doses 2mg/kg and 1mg/kg of body weight of mice. The survival was monitored till 10 days.

Likewise, organs were stored in 3.5% PFA before histopathogical sample preparation

2.2.12 Statistical Analysis

Statistical analyses were performed with GraphPad Prism software. The Student's t-test was

performed as indicated. The results are expressed as mean ± SD or mean ± SEM. Group sizes,

experiment number, and p values for each experiment are described in figure legends.

Results

Loss of spermidine transporter and biosynthesis genes in *Salmonella* Typhimurium

compromises its ability to be phagocytosed by the macrophages

The pathoadaptation in *Salmonella* involves multiple players, which counteracts the stressful

condition it encounters in the host macrophages. Polyamines being a group of well-studied stress

response molecules, we were interested in determining the expression of the spermidine transporter

and biosynthesis genes in *Salmonella* Typhimurium. Our previous study shows that *Salmonella*

upregulates the spermidine transporter genes (*potA, potB, potC* and *potD*) and the biosynthesis

genes (*speE* and *speD*) during the log phase of growth *in vitro*. Here we assessed the mRNA levels

of *potA, potB, potC* and *potD* in STM WT upon infection into the RAW264.7 macrophage cell

line. We noted that all the genes showed a 1.5-2 fold upregulation post 6 hours of infection into

macrophages till 16 hours (**Fig 1 A)**. Further, our results showed that the spermidine biosynthesis

enzymes *speE* and *speD* were upregulated 1.5 to 2 folds post 6 hours to 16 hours post-infection

into macrophages (**Fig 1 B)**. These results indicate that *Salmonella* Typhimurium enhances its

intracellular spermidine biosynthesis and imports from the extracellular milieu. It might be a

strategy of the pathogen to increase the intracellular pool of stress response molecules like the

polyamine spermidine, as it encounters the hostile environment of host macrophages.

We then determined how the spermidine transporter and biosynthesis mutants (STM Δ*potCD*, STM

Δ*speED* and STM Δ*potCD*Δ*speED*) behave upon infection into RAW264.7 macrophages. We

infected STM WT, STM Δ*potCD*, STM Δ*speED* and STM Δ*potCD*Δ*speED* into RAW264.7 cells

and observed that STM Δ*potCD*, STM Δ*speED* and STM Δ*potCD*Δ*speED* showed reduced

phagocytosis by the macrophages compared to the wild type (**Fig 1 C**). Interestingly, STM

Δ*potCD*, STM Δ*speED* and STM Δ*potCD*Δ*speED* showed a compromised ability to proliferate

intracellularly in the RAW264.7 cells further validating a previous study [10](**Fig S1 A**). To further

validate our results, we assessed the behaviour of the spermidine transporter and biosynthesis gene

mutants in primary macrophages isolated from the peritoneal lavage of C57BL/6 mice. Likewise,

the mutants showed attenuated proliferation and uptake by phagocytosis into the peritoneal

macrophages (**Fig 1 D and S1 B**). Collectively these results suggest that spermidine is a critical

molecule in *Salmonella* Typhimurium to infect and survive in macrophages. To further investigate

the reason behind the reduced ability to be taken up by macrophages upon loss of spermidine

biosynthesis and transport, we treated STM WT, STM Δ*potCD*, STM Δ*speED* and STM

Δ*potCD*Δ*speED* with mouse complement-sera. Mouse complement-sera acts as an opsonin and

thus potentiates the interaction of the bacteria with the macrophages. Upon pre-treatment with

mouse complement-sera, we noted a rescue in the reduced uptake of the mutants by peritoneal

macrophages isolated from C57BL/6 mice (**Fig 1 E**). A study on *Salmonella* Typhimurium

revealed that the aflagellate and non-motile *Salmonella* collide less frequently with macrophages

and gets merest time to maintain contact with the macrophages, thereby showing decreased

phagocytosis [14]. Our group previously showed that loss of spermidine production and import in

Salmonella Typhimurium results in the loss of flagella formation on the bacterial cell surface.

Thus, we determined the percentage phagocytosis for a flagellin-deficient strain of *Salmonella*

Typhimurium, STM Δ*fliC* and observed that STM Δ*potCD*, STM Δ*speED*, STM Δ*potCD*Δ*speE*

and STM Δ*fliC* exhibited a significantly decreased ability to be taken up by the RAW264.7

macrophages (**Fig 1 F**). Furthermore, we observed a rescue of the attenuated percentage

phagocytosis and fold proliferation of only STM Δ*speED* upon supplementation of spermidine

(100μM) during the growth of bacteria prior to infection (**Fig 1 G and S1C**). Hence, the plausible

explanation is that the reduced ability to form flagella in the spermidine mutants causes less

frequent interaction with the macrophages and provides minimal contact time for infection, leading

to reduced phagocytosis by the macrophages.

Figure 1: Loss of spermidine transporter and biosynthesis genes in *Salmonella* Typhimurium compromises its ability to be phagocytosed by the macrophages

Figure 1: Loss of spermidine transporter and biosynthesis genes in *Salmonella* Typhimurium compromises its ability to be phagocytosed by the macrophages

A.The mRNA expression of spermidine transporter genes *potA, potB, potC* and *potD* in STM WT upon infection into RAW264.7 cells, B. The mRNA expression of spermidine biosynthesis genes *speE* and *speD* in STM WT upon infection into RAW264.7 cells, C. The percentage phagocytosis of the spermidine mutants in RAW264.7 cells, D. The percentage phagocytosis in primary macrophages isolated from peritoneal lavage of C57BL/6 mice, E. The percentage phagocytosis upon pre treatment with mouse-complement sera which act as an opsonin, F. The percentage phagocytosis in RAW264.7 cells with flagellin mutant (STM *ΔfliC*), G. The percentage phagocytosis in RAW264.7 cells with the spermidine mutants grown in media supplemented with 100μM spermidine (SPD). Student's t-test was used to analyze the data; p values ****<0.0001, ***<0.001, **<0.01, *<0.05. Two-way Anova was used to analyze the grouped data; p values ****<0.0001, ***<0.001, **<0.01, *<0.05.

Spermidine provides stress resistance in *Salmonella* Typhimurium by regulation of the expression of numerous antioxidative enzymes

The loss of spermidine transport and biosynthesis function in *Salmonella* Typhimurium renders it incapable of proliferation and survival in macrophages. In the host macrophages, the bacteria encounter numerous threats, of which the foremost is the rapid oxidative burst mediated by the NOX2. The reactive oxygen species superoxide radical can easily diffuse through the bacterial membrane and pose a major threat to the pathogen. ROS acts on multiple molecules such as nucleic acids, proteins and lipids, thus damaging the cell membranes, DNA and proteins within the bacteria [15]. Spermidine has been linked to stress response against oxidative stress and protects bacteria

in *E. coli* and *Pseudomonas aeruginosa.* Thus, we were intrigued to understand the role of spermidine in antioxidative response in *Salmonella* Typhimurium. We examined the survival of STM WT, STM Δ*potCD*, STM Δ*speED*, STM Δ*potCD*Δ*speE* upon exposure to the oxidative agent hydrogen peroxide *in vitro.* We noticed that at high concentrations of hydrogen peroxide, 5mM and 10mM, the STM Δ*potCD*, STM Δ*speED* and STM Δ*potCD*Δ*speE* showed significantly lesser survival than STM WT(**Fig 2 A**). We further determined the expression of *potA, potB, potC, potD, speE* and *speD* in STM WT upon exposure to 1mM hydrogen peroxide. There was a 2-3 fold upregulation in the mRNA expression of the transporter genes over the untreated during the early log phase of growth (6 hours) and the late log phase of growth (12 hours) *in vitro* (**Fig S2 A-D**). Similarly, the biosynthesis genes *speE* and *speD* were 4-6 fold upregulated in their corresponding mRNA expressions during their early log phase of growth (6 hours) and the late log phase of growth (12 hours) *in vitro* (**Fig S2 E and F**). Our results show that *Salmonella* Typhimurium upregulates spermidine transport and biosynthesis upon oxidative stress suggesting that spermidine mounts a protective function in such a stressful condition to aid bacterial survival.

Bacteria sense the environmental changes and cues to respond and adapt to the altered environment. They use the two-component systems, transcriptional activators and repressors to alter gene expression in response to a stimulus [16]. Polyamines in *E. coli* regulate multiple genes at the transcription and translation together, referred to as the "Polyamine modulon". These involve the numerous mRNAs, tRNAs, sigma factors, translational factors and two-component systems during the bacterial growth as well as in stress conditions [17-19]. *Salmonella* harbors multiple antioxidative enzymes to detoxify the ROS intracellularly [20]. Our study so far show that STM Δ*potCD*, STM Δ*speED* and STM Δ*potCD*Δ*speE* is attenuated in survival under *in vitro* oxidative stress than STM WT. To gain mechanistic insight into the attenuated survival of *Salmonella*

Typhimurium we determined the mRNA expression of the critical transcription factor *rpoS*, which

activates the expression of the catalase enzymes (*katG and katE*) in order to detoxify hydrogen

peroxide to water in the bacteria enzymatically. In both STM Δ*potCD* and STM Δ*speED* the mRNA

expression of *rpoS* is significantly down regulated from 6 hours post infection in RAW264.7 cells

(**Fig 2 B and C, S3 A**). Further, its downstream target *katG* was also down regulated in STM

Δ*potCD* and STM Δ*speED* from 6 hours post infection into RAW264.7 cells (**Fig 2 D and E, S3**

B). We further assessed the mRNA expression of the transcription factor *soxR*, which regulates the

expression of superoxide dismutases (*sodA* and *sodB*). Upon infection in RAW264.7 cells,

expression of *soxR* was significantly downregulated in STM Δ*potCD* and STM Δ*speED* with

respect to STM WT (**Fig 2 F and G, S3 C**). Superoxide dismutases act on superoxide radicals, the

potent ROS encountered in macrophages, converting to hydrogen peroxide. The mRNA expression

of both *sodA* and *sodB* were likewise downregulated in STM Δ*potCD* and STM Δ*speED* upon

infection into RAW264.7 macrophages (**Fig 2 H-K, S3 D and E**). A major antioxidant in most

living organisms is glutathione (GSH), which directly acts as a quencher of ROS. GSH is

synthesized by Glutathione synthase (GshA) which in turn is regulated by EmrR transcription

factor. We observed that the mRNA expression of *emrR* is downregulated in STM Δ*potCD* and

STM Δ*speED* upon infection into RAW264.7 macrophages (**Fig 2 L and M, S3 F**). Similarly, *gshA*

transcript expression is downregulated as well (**Fig 2 N and O, S3 G**). To further validate the down

regulation of the glutathione synthesis arm in spermidine transport and biosynthesis mutants, we

determined the intracellular GSH levels and noted the in STM Δ*potCD,* STM Δ*speED* and STM

Δ*potCD*Δ*speED* the levels were significantly less (**Fig S3H and I).** Taken together these results

indicate that spermidine regulates the transcription of multiple transcription factors involved in

oxidative stress response in *Salmonella* Typhimurium. Importantly, we found a mechanism of

oxidative stress resistance in *Salmonella* Typhimurium regulated by the spermidine, potentiating

the survival of the bacteria.

Figure 2: Spermidine provides stress resistance in *Salmonella* Typhimurium by regulation of the expression of numerous antioxidative enzymes

Figure 2: Spermidine provides stress resistance in *Salmonella* Typhimurium by regulation of the expression of numerous antioxidative enzymes

A.The *in vitro* hydrogen peroxide sensitivity assay with the spermidine transport and biosunthesis mutants, B. The mRNA expression of stress responsive transcription factor *rpoS* in STM *ΔpotCD* upon infection into RAW264.7 cells, C. The mRNA expression of stress responsive transcription factor *rpoS* in STM *ΔspeED* upon infection into RAW264.7 cells, D. The mRNA expression of *katG* in STM *ΔpotCD* upon infection into RAW264.7 cells, E. The mRNA expression of *katG* in STM *ΔspeED* upon infection into RAW264.7 cells, F. The mRNA expression of oxidative stress responsive transcription factor *soxR* in STM *ΔpotCD* upon infection into RAW264.7 cells, G. The mRNA expression of oxidative stress responsive transcription factor *soxR* in STM *ΔspeED* upon infection into RAW264.7 cells, H. The mRNA expression of *sodA* in STM *ΔpotCD* upon infection into RAW264.7 cells, I. The mRNA expression of *sodA* in STM *ΔspeED* upon infection into RAW264.7 cells, J. The mRNA expression of *sodB* in STM *ΔpotCD* upon infection into RAW264.7 cells, K. The mRNA expression of *sodB* in STM *ΔspeED* upon infection into RAW264.7 cells, L. The mRNA expression of glutathione synthetase specific transcription factor *emrR* in STM *ΔpotCD* upon infection into RAW264.7 cells, M. The mRNA expression of glutathione synthetase specific transcription factor *emrR* in STM *ΔspeED* upon infection into RAW264.7 cells, N. The mRNA expression of *gshA* in STM *ΔpotCD* upon infection into RAW264.7 cells, O. The mRNA expression of *gshA* in STM *ΔspeED* upon infection into RAW264.7 cells, Student's t-test was used to analyze the data; p values ****<0.0001, ***<0.001, **<0.01, *<0.05.

Spermidine controls a novel enzyme Glutathionyl-spermidine synthetase in *Salmonella* Typhimurium, and together mount an intracellular antioxidative response

The spermidine synthesized from putrescine has two fates. It is either acetylated by the enzyme SpeG or it is covalently conjugated to GSH to form Glutathionyl spermidine (GS-sp) catalysed by the enzyme Glutathionyl-spermidine synthetase (Gss). Tabor (1974) first discovered the existence of this enzyme in *E. coli* [21]. In *E. coli* GS-sp is generated at a higher level in the stationary phase and very less in the late exponential phase. It also interacts and modifies the thiol containing proteins under high H_2O_2-containing media leading to the formation of Gsp-thiolated proteins (PSSGsp). Certain *in vitro* experiments with dehydro-ascorbate suggested that Gsp might have higher antioxidant properties than GSH [22] and may be more effective in protecting against DNA damage by free radicals [23]. However, in *E. coli* Gsp could not be linked to its pathogenicity [24]. Among *Enterobacteriaceae*, *Salmonella* was found to possess this unique enzyme. Our *in silico* analysis suggested that the enzyme in *Salmonella* Typhimurium (GspSA, encoded by *gsp*) has 90% identity with *E. coli gss* and the SWISS MODEL predicts it to be a homo-dimeric protein.(**Fig S4 A-C**). Also, the spermidine synthesised by SpeE in *Salmonella* Typhimurium is directly fed into the pathway to synthesise GS-sp. We thus, investegated biological role of this novel enzyme in *Salmonella* Typhimurium. We noted that the mRNA expression of *gsp* is significantly upregulated at the late-log phase of growth of STM WT in LB media in the presence of H_2O_2 (**Fig S4 D**). Upon infection into RAW264.7 cells, STM WT upregulates the mRNA expression of *gsp* at 6 hours and 16 hours post-infection into RAW264.7 cells (**Fig 3 A**). Further, the *Salmonella* Typhimurium mutant of *gsp* (STM Δ*gsp*) showed attenuated proliferation in RAW264.7 cells similar to STM Δ*katG,* which has reduced capability to detoxify ROS (**Fig 3 B**). Thus, our results suggest that *gsp* is a critical in *Salmonella* Typhimurium to survive and cope with the oxidative stress and hostile

environment of macrophages. Interestingly, the mRNA expression of *gsp* was found to be

downregulated in both STM Δ*potCD* and STM Δ*speED* at 16 hours post-infection into RAW264.7

macrophages (**Fig 3 C and D**). These findings show that spermidine maintains its intracellular

levels by regulating the flux of spermidine into other pathways, such as the GspSA pathway. Also,

it further potentiates the ability of *Salmonella* Typhimurium to mount an antioxidative response by

regulating *gsp* expression

GspSA enzyme in Trypanosomes was named Trypanothione synthetase, while the conjugate is

called Trypanothione (TSH). In trypanosomes TSH is essential, and these organisms entirely rely

on TSH and possess no GSH reductase. Also, they have evolved to use TSH reductase instead of

GSH reductase, Glutaredoxins or Thioredoxins [25]. In *E. coli* Gsp forms bonds with the Cysteine

thiol groups of numerous proteins and protect them from oxidation under oxidative stress. Cysteine

thiol groups are highly prone to attack by ROS and get oxidised to sulphinic and sulphonic acids.

In *Salmonella* Typhimurium we observe that loss of *gsp* results in attenuated proliferation and

survival in macrophages. Thus, we investegated the survival of STM Δ*gsp* upon exposure to

hydrogen peroxide *in vitro*. STM Δ*gsp* exhibited reduced survival in the presence of high 5mM

and 10mM concentrations of H_2O_2 (**Fig S4 E**). Likewise, upon exposure to agents of oxidative

stress and nitrosative stress H_2O_2 and $NaNO_2$ together, we observe that STM Δ*gsp* exhibited

reduced survival at higher concentrations such as 5mM and 10mM (**Fig 4 E**). Thus, *gsp* is critical

in *Salmonella* Typhimurium to shield the bacteria from the action of ROS and RNS. As we

observed that spermidine transporter and biosynthesis mutants and *gsp* mutant of *Salmonella*

Typhimurium are compromised in their survival under oxidative stress and in macrophages thus,

we were interested to assess the intracellular ROS detoxification abilities of the strains. We

determined the intracellular ROS in STM WT, STM Δ*potCD*, STM Δ*speED*, STM Δ*potCD*Δ*speE,*

STM Δ*gsp* and STM Δ*katG*. Interestingly, STM Δ*potCD,* STM Δ*speED*, STM Δ*potCD*Δ*speE,*

STM Δ*gsp* and STM Δ*katG* showed significantly higher intracellular ROS levels when they were

exposed to 5mM and 10mM H_2O_2 (**Fig 3 F**). These suggest that lower intracellular levels of

spermidine correlate to higher intracellular ROS levels in *Salmonella* Typhimurium. To validate

our observed results, we utilised a genetically engineered tool to sense the redox status of the

bacterial cytosol, roGFP2 is a genetically modified form of GFP, and we use pQE60-Grx1-roGFP2

plasmid. The glutaredoxin (Grx1) fused to the roGFP2, reversibly transfers electrons between the

cytosolic pool of GSH/GSSG and the thiol group of roGFP2, and the ratio of fluorescence ratio at

408nm and 488nm determine the redox status of bacterial cytoplasm [12]. We observed that STM

Δ*potCD*, STM Δ*speED* harbouring the pQE60-Grx1-roGFP2 showed higher ratio of

405nm/488nm compared to STM WT in the presence of 5mM H_2O_2 (**Fig 3 G**). However, STM

Δ*gsp* did not show a significantly higher ratio of 405nm/488nm. Moreover upon supplementation

of the growth media with 100 μM spermidine, only in STM Δ*speED* we observed a lower

intracellular ROS and lesser 405nm/488nm in higher concentration of H_2O_2 (**Fig S4 F and**

G).Thus, our results indicate that spermidine is critical in mounting an antioxidative response to

detoxify the intracellular ROS, by regulating multiple antioxidant genes in *Salmonella*

Typhimurium. To further dissect the role of spermidine in protection of *Salmonella* Typhimurium

from oxidative stress, we infected STM WT, STM Δ*potCD*, STM Δ*speED*, STM Δ*potCD*Δ*speE,*

STM Δ*gsp* and STM Δ*katG* in primary macrophages isolated from the peritoneal lavage of

gp91phox -/- mice. Gp91Phox is the major subunit of the NOX2 complex, that aids in the catalysis

of oxygen to superoxide radical. Interestingly we observe a rescue in the attenuated fold

proliferation of STM Δ*potCD*, STM Δ*speED*, STM Δ*potCD*Δ*speE,* STM Δ*gsp* and STM Δ*katG* in

peritoneal macrophages isolated from *gp91phox -/-* mice (**Fig 3 H**). Our results demonstrate the

vital role of spermidine in oxidative stress resistance in *Salmonella* Typhimurium.

Figure 3: Spermidine controls a novel enzyme Glutathionyl-spermidine synthetase in *Salmonella* Typhimurium, and together mount an intracellular antioxidative response

Figure 3: Spermidine controls a novel enzyme Glutathionyl-spermidine synthetase in *Salmonella* Typhimurium, and together mount an intracellular antioxidative response

A.The mRNA expression of *gsp* in STM WT upon infection into RAW264.7 cells, B The fold proliferation of STM WT, STM *Δgsp* and STM *Δgsp:gsp* in RAW264.7 cells, C. The mRNA expression of *gsp* in STM *ΔpotCD* upon infection into RAW264.7 cells, D. The mRNA expression of *gsp* in STM *ΔspeED* upon infection into RAW264.7 cells, E The *in vitro* Hydrogen peroxide and nitic oxide sensitivity assay, F. Intracellular reactive oxygen species determination using the cell permeable H2DCFDA dye, G. The intracellular degree of oxidation using pQE60-grx1-roGFP2 construct, H. The fold proliferation in primary macrophages isolated from wild type C57bl/6 mice and *gp91phox-/-* mice, Student's t-test was used to analyze the data; p values ****<0.0001, ***<0.001, **<0.01, *<0.05. Two-way Anova was used to analyze the grouped data; p values ****<0.0001, ***<0.001, **<0.01, *<0.05.

Spermidine is critical for *Salmonella* Typhimurium to colonise the primary and secondary sites of infection in mice

Salmonella infects the host and breaching the epithelial cells at the Peyer's patches in the distal ileum, it disseminates to the secondary sites of infection namely the Mesentric Lymph node (MLN), spleen and liver. From the basolateral surface of the epithelial cells at the lamina propria, *Salmonella* is taken by the macrophages and polymorphonuclear cells (PMN). We observed that the spermidine transporter and biosynthesis mutants show attenuated survival in macrophages and under oxidative stress *in vitro*. We were intrigued to study the behaviour of the mutants during *in vivo* colonisation in the mouse model of *Salmonella* infection. We infected C57BL/6 mice by orally gavaging with STM WT, STM *ΔpotCD*, STM *ΔspeED*, STM *ΔpotCDΔspeE,* STM *Δgsp* and

STM Δ*katG* at a CFU of 10^7 per mice. We noted that STM Δ*potCD*, STM Δ*speED*, STM

Δ*potCD*Δ*speE,* STM Δ*gsp* and STM Δ*katG* showed a significantly lower organ burden in Peyer's

patches, MLN, spleen and liver upon oral gavage (**Fig 4 A-E)**. Our previous study showed that the

spermidine transporter and biosynthesis mutants exhibit a poor invasiveness into IECs, which can

explain our *in vivo* colonisation upon oral gavage. Oral gavage mimics the physiological route of

Salmonella infection into its host. Hence, it requires to be able to breach the intestinal barrier

successfully. Incapability to invade the IECs in STM Δ*potCD*, STM Δ*speED*, STM Δ*potCD*Δ*speE*

and STM Δ*gsp* explains the diminished colonisation in the organs. To dissect the role of spermidine

in *in vivo* colonisation we infected C57Bl/6 mice intraperitoneally, by bypassing the entry by

breaching epithelial barrier. We observed that upon infecting intraperitoneally, STM Δ*potCD*, STM

Δ*speED*, STM Δ*potCD*Δ*speE,* STM Δ*gsp* and STM Δ*katG* exhibited reduced colonisation in

spleen and liver and less dissemination in blood compared to STM WT (**Fig 4 F-I)**. Also, the

histopathological sections show significantly less liver tissue damage with STM Δ*potCD*, STM

Δ*speED*, STM Δ*potCD*Δ*speE,* STM Δ*gsp* and STM Δ*katG* , which is validated by disease scoring

of the same(**Fig 4 J and K)**. Our results, thus show that spermidine aids in the *in vivo* pathogenesis

and virulence of *Salmonella* Typhimurium.

Figure 4: Spermidine is critical for *Salmonella* Typhimurium to colonise the primary and secondary sites of infection in mice

Figure 4: Spermidine is critical for *Salmonella* Typhimurium to colonise the primary and secondary sites of infection in mice

A.The experimental protocol for organ burden in C57BL/6 mice by orally gavaging 10^7 CFU per mice, B. The organ burden post 5 days of oral gavage in intestine, C. in Mesentric lymph node (MLN), D. in spleen, E. in liver, F. The experimental protocol for organ burden in C57BL/6 mice upon intraperitoneal (I.P) infection with 10^3 CFU per mice, G. The organ burden 3 days post I.P infection in spleen, H. in liver, I. dissemination in blood, J. The hematoxylin and eosin staining of the sections of liver 3 days post I.P infection of C57Bl/6 mice, K. Disease score from the histopathogical sections of liver, Here, Small necrotic areas (N), Congestion and damage to the endothelial lining of the central vein (C), Congestion of the hepatic portal vein and poor hepatic architecture (HPV), Inflammatory immune cells (IC). The disease score is as :0 for normal pathology, 1 for mild/ minor pathology, 2 for moderate pathology, and 3 for severe pathological changes. Mann Whitney test was used to analyse organ burden in mice; p values ****<0.0001, ***<0.001, **<0.01, *<0.05. Student's t-test was used to analyze the data; p values ****<0.0001, ***<0.001, **<0.01, *<0.05.

Salmonella rewires host polyamine metabolism to potentiate its survival within host macrophages

Most of the intracellular pathogens establish their persistence in the phagocytic cells and are often found to be associated with different population of the macrophages. Like *Brucella abortus* preferentially resides in the Alternatively activated macrophages (AAM), where it survives and replicates by exploiting the host polyamines. A research group has shown that the metabolism of

AAM is shifted to increase polyamine biosynthesis by *Brucella abortus* and thereby promote the

bacterial survival [26]. Similarly, *Salmonella* also resides in macrophages, a way to lead chronic

infections. We were interested to know whether *Salmonella* exploits the host polyamines and leads

to a rewiring of host cell metabolism. To understand the host-pathogen relationship in depth, we

assessed the mRNA expression of Ornithine decarboxylase 1 (*mOdc1*) which catalyses the rate-

determining step of polyamine biosynthesis and Spermidine synthase (*mSrm*) that synthesises

spermidine from putrescine by transferring the aminopropyl group from decarboxylated-S in

RAW264.7 cells upon infection with STM WT. We observed that mRNA expression of *mOdc1*

and *mSrm* were upregulated at 6 and 16 hours post-infection (**Fig 5 A and B**). Further, the mRNA

expression of *mOdc1* and *mSrm* were enhanced in the spleen and liver of C57BL/6 mice 5 days

post oral gavage with STM WT (**Fig C- F**). Our results show that *Salmonella* Typhimurium upon

infection into the host enhances the expression of host polyamine biosynthesis genes. Moreover,

we have previously observed that the *Salmonella* Typhimurium that cannot import spermidine

cannot survive and proliferate as much as STM WT in macrophages. To further delve into the role

of host acquired polyamines, we knocked down *Odc1* in RAW264.7 cells (**Fig S5 A**). Upon

knockdown of *mOdc1*, *Salmonella* Typhimurium showed significantly attenuated proliferation in

RAW264.7 cells (**Fig 5 G**). However, the knockdown of *mOdc1* did not alter the percentage of

phagocytosis by the macrophages (**Fig 5 H**). Similarly we knocked down *Srm* in RAW264.7 cells

and observed that knock-down of spermidine synthase in the host compromises the ability of STM

WT to proliferate and get phagocytosed by the macrophages (**Fig S5 B-D**).

Figure 5: *Salmonella* **rewires host polyamine metabolism to potentiate its survival within host macrophages**

Figure 5: *Salmonella* **rewires host polyamine metabolism to potentiate its survival within host**

macrophages

A.The mRNA expression of *mOdc1* (mouse ornithine decarboxylase) in RAW264.7 cells upon

infection with STM WT, B. The mRNA expression of *mSrm* (mouse spermidine synthase) in

RAW264.7 cells upon infection with STM WT, C. The mRNA expression of *mOdc1* (mouse

ornithine decarboxylase) in spleen of C57BL/6 mice 5 days post infection with STM WT by oral

gavage, D. The mRNA expression of *mOdc1* (mouse ornithine decarboxylase) in liver of C57BL/6

mice 5 days post infection with STM WT by oral gavage, E. The mRNA expression of *mSrm*

(mouse spermidine synthase) in spleen of C57BL/6 mice 5 days post infection with STM WT by

oral gavage, F. The mRNA expression of *mSrm* (mouse spermidine synthase) in liver of C57BL/6

mice 5 days post infection with STM WT by oral gavage, G. The fold proliferation of STM WT in

RAW264.7 cells upon transient knockdown of *mOdc1,* H. The percentage phagocytosis of STM

WT in RAW264.7 cells upon transient knockdown of *mOdc1.* Here SCR is Scrambled (no target

for knock-down), two different targeted shRNA were used for knock-down purposes, Sh1is

shRNA-1 for knock-down, Sh2 is shRNA-2 for knock-down, and Sh1+Sh2 indicates where both

the shRNAs were used to obtain the knock-down Student's t-test was used to analyze the data; p

values ****<0.0001, ***<0.001, **<0.01, *<0.05.

The chemopreventive drug DFMO, reduces *Salmonella* Typhimurium burden in the host by enhancing nitric oxide production

Polyamines are essential molecules in eukarotes, with multiple roles in differentiation,

proliferation and development. Many studies have shown that polyamine levels are upregulated in

cancer cells and elevated levels of polyamines are associated with breast cancer, neuroblastoma,

hepatocellular carcinoma, prostate cancer, lung cancer, colorectal cancer and lukemia [27-31].

D,L-α-difluoromethylornithine (DFMO) , an inhibitor of ODC was developed as a potent drug to

treat cancer in the year 1970 [32]. DFMO irreversibly binds to the active site of ODC and acts as

a suicide inhibitor, thereby reducing polyamine levels and having a cytostatic effect. As a single therapeutic agent it was found to be effective only in neuroblastoma , and cinical trial for other cancer type was unsatisfactory [33, 34]. However, DFMO has been successfully developed as a chemopreventive drug, with FDA approval for treatment of Human African Trypanosomiasis (HAT) [35, 36]. To test whether DFMO can be used as a therapeutic drug against *Salmonella* infection, we treated RAW264.7cells with DFMO during the infection with *Salmonella* Typhimurium, and observed a significant attenuation in the fold proliferation of STM WT (**Fig 6 A)**. Studies have shown that DFMO binds to ODC to prevent further production of putrescine from ornithine and also acts on Arginase1 and reduces the available pool of ornithine for polyamine biosynthesis [37, 38]. This, ensures the flux of arginine to be fed into the nitric oxide synthase (NOS2) pathway and leads to elevated levels of nitric oxide in the cell, which in turn negatively acts on ODC to further block polyamine synthesis [39].We assessed the mRNA expression of *mNos2* upon knockdown of *mOdc1* in RAW264.7 cells followed by infection with STM. We observed an upregulation of *mOdc1* mRNA levels at 6 hours and 16 hours post infection in RAW264.7 cells (**Fig 6 B)**. Further we determined the levels of nitric oxide using a cell permeable dye DAF-2DA, and noted that upon treatment with DFMO there was higher production of nitric oxide (**Fig 6 C and D)**. Taken together our results demonstrate the role of DFMO in diminishing the survival and proliferation of *Salmonella* Typhimurium by blocking Odc1 and enhanced production of nitric oxide in murine macrophages.

To further address whether DFMO also act as potent anti-*Salmonella* drug in mice model of infection, we infected C57BL/6 mice with 10^7 CFU per mice by oral gavage. Every alternate day the mice were injected with DFMO in two different doses for two different cohorts (2mg/kg of body weight of mice and 1mg/kg of body weight of mice). Five days post oral gavage we observed

that DFMO treatment of 2mg/kg body weight of mice significantly reduced the colonisation of

STM WT in intestine, MLN, spleen and liver than in the untreated mice (**Fig 6 E-I**). Further, using

immunofluorescence we observed DFMO significantly reduced the colonisation of STM and and

the levels of spermidine in mouse ileum (**Fig S6 A-D**). Moreover, treatment of mice with DFMO

at a dose of 2mg/kg body weight of mice, increased the survival of mice upon infection with STM

WT (**Fig 6 J and K**). Also, the weight reduction in mice treated with with DFMO at a dose of

2mg/kg body weight, was less upon infection with STM WT (**Fig S6 E**). Next, we evaluated the

tissue damage upon infection of STM in mice liver. The results show that DFMO treatment of

mice at a dose of 2mg/kg of body weight significantly lowered the disease score, suggesting lesser

tissue damage compared to untreated (**Fig 6 L and M**). Thus, DFMO serves as a potential drug to

treat *Salmonella* infection in mice by reducing the bacterial burden and tissue damage in mice and

enhancing the survival of mice upon infection with STM.

Figure 6: The chemopreventive drug DFMO, reduces *Salmonella* Typhimurium burden in the host by enhancing nitric oxide production

Figure 6: The chemopreventive drug DFMO, reduces *Salmonella* Typhimurium burden in the host by enhancing nitric oxide production

A. The intracellular fold proliferation of STM WT in RAW264.7 cells upon treatment with Difluoromethyl ornithine (DFMO) (5μM), quantified by Gentamicin protection assay, B. The mRNA expression of *mNos2*(mouse Nitric oxide synthase) in RAW264.7 cells, upon transient knockdown of *Odc1* followed by infection with STM WT, the expression is normalized to beta-actin as an internal control, C. The representative scatter plot for DAF2DA dye positive population of RAW264.7 cells, upon transient knockdown of *Odc1* followed by infection with STM WT, the data is determined by flow-cytometry post staining the infected macrophages with the dye, D. The quantification of (C), E. The experimental procedure to study the organ burden of STM WT in C57BL/6 mice upon treatment with DFMO, F-I. The organ burden of STM WT in Intestine, MLN, Spleen and Liver of C57BL/6 mice upon intraperitoneal treatment of DFMO (2mg/kg and 1mg/kg of body weight) as mentioned in (E), J. The Experimental procedure to study the survival of C57BL/6 mice upon infection with STM WT and treatment with DFMO, K.The survival of C57BL/6 mice upon infection with STM WT and upon intraperitoneal treatment of DFMO (2mg/kg and 1mg/kg of body weight) as mentioned in (J), L. Hematoxylin and eosin staining of histopathological sections of liver upon DFMO treatment to C57BL/6 mice. Here, Small necrotic areas (N), Congestion and damage to the endothelial lining of the central vein (C), Congestion of the hepatic portal vein and poor hepatic architecture (HPV), Inflammatory immune cells (IC), M. The disease score index for liver tissue damage upon STM WT infection in C57BL/6 mice with DFMO treatment. The disease score is as :0 for normal pathology, 1 for mild/ minor pathology, 2 for moderate pathology, and 3 for severe pathological changes, Student's t-test was used to analyze

652 the data; p values ****<0.0001, ***<0.001, **<0.01, *<0.05. Two-way Anova was used to

653 analyze the grouped data; p values ****<0.0001, ***<0.001, **<0.01, *<0.05.

Discussion

A significant part of the life cycle of *Salmonella* during its pathogenesis involves its stay inside

the macrophages. The Gram-negative bacteria come across multiple host environmental stress

conditions inside the macrophages [40]. A key mechanism by which the host macrophages try to

limit the invading pathogen is by the rapid oxidative burst and ROS production. ROS includes

superoxide radicals, hydroxyl radicals, peroxy-nitrites, peroxy-chlorides and hydrogen peroxide.

ROS can pass through the bacterial cell wall and act on lipids, proteins and nucleic acids by

oxidising them and leading to cellular damage. *Salmonella* is often referred to as a smart pathogen.

Over the years, it has evolved to possess multiple strategies to combat the host-derived stresses

[41, 42]. To combat the oxidative burst generated NOX2 in macrophages, *Salmonella* possesses

multiple antioxidant enzymes such as the catalases KatE and KatG, the superoxide dismutases

SodA and SodB, the Alkyl hydroperoxide reductase, the glutaredoxins and thioredoxins and Hrg

transcriptional regulator [43]. Polyamines assist in *Salmonella* virulence and aid in stress

resistance. However, the mechanism behind the role of polyamines in *Salmonella* stress resistance

and virulence remains less appreciated.

Our study identifies spermidine as a stress-responsive regulatory molecule in *Salmonella*

Typhimurium. We show spermidine is critical for the survival and proliferation of STM in

macrophages and in the presence of oxidative stress *in vitro*. The spermidine transporter and

biosynthesis mutants show significant reduced capability to be phagocytosed by the macrophages.

Findings from our previous study (unpublished) showed that the intracellular level of spermidine

is significantly less in both the spermidine transport and biosynthesis mutants, which further

explains the attenuated proliferation of both the mutants in macrophages. The previous findings

also showed that in absence of the transport genes in *Salmonella* Typhimurium the synthesis genes

are down regulated and vice versa. The absence of spermidine transport and biosynthesis

diminishes the mRNA expression of multiple arms of oxidative stress response in *Salmonella.*

Multiple studies show the role of polyamines in the regulation of transcription of multiple genes

by interacting with DNA in eukaryotes. They bind to the DNA and change conformation as in C-

MYC transcription, and in other cases, enhance DNA-protein binding affinities like for Estrogen-

response elements [44, 45]. Thus, in *Salmonella* Typhimurium *rpoS, soxR* and *emrR* fall under the

"Polyamine modulon".

We further characterise a novel enzyme GspSA in *Salmonella* Typhimurium, which synthesises a

conjugated product of glutathione (GSH) and spermidine called GS-sp. GspSA is critical for

Salmonella Typhimurium to survive and proliferate in macrophages. Our study shows that the

absence of GspSA attenuates the survival of STM under oxidative stress conditions *in vitro,*

suggesting a vital role of Gsp in protecting STM from oxidative damage. Gsp in *E. coli* carries out

its function by modifying protein thiol groups under oxidative stress to protect the proteins from

getting oxidised and damaged. In *Salmonella* Typhimurium, we expect GspSA to generate GS-sp

under oxidative stress conditions and inside the macrophages, and likewise interact with cysteine-

thiol groups in proteins to shield them from oxidative damage. MK Chattopadhyay (2013) showed

that this enzyme (GspSA) is specifically present in two groups of organisms namely bacteria and

kineto-plastids respectively and completely absent in other organisms such as human, rats,

drosophila etc., among the bacteria group in all *enterobacteria* showed 27-100% homology and

>65% identity in around 50% of the species, with *E. coli* [24]. Thus, the absence of GspSA in

eukaryotes makes it a potent drug target for treatment of *Salmonella* infections. Thus, spermidine

exerts pleiotropic effects in *Salmonella*, by orchestrating the multiple arms of antioxidative

response. It strengthens the bacteria to cope with hostile host environments. Our studies in the *in*

vivo model of *Salmonella* Typhimurium infection further dissect the role of spermidine in assisting

in the pathogenesis by enhancing its virulence properties. Moreover, it is at the nexus of multiple

oxidative stress response arms in *Salmonella*, thereby assisting in mounting an antioxidative

response to promote its survival in macrophages.

Studies over the past years have given insights into the function of polyamines in the pathogenesis

of multiple virulent bacteria. Many human pathogenic bacteria have developed ways to exploit,

interfere and manipulate the polyamine metabolism of the host to enhance their fitness within the

niche. As in *Shigella* and *Vibrio spp.* polyamines produced by the bacteria are critical in

determining virulence [46, 47]. While bacteria such as *Legionella spp.*, which does not synthesise

polyamines, depend on the host-acquired polyamines for their pathogenesis [48]. Another unique

mechanism is observed in *H. pylori* which activates polyamine oxidation, thereby dysregulation

the innate-immune response [49]. Also, in response to *H. pylori* infection, the host macrophages

increase the arginase activity and ornithine decarboxylase activity to produce polyamines [50]. A

recent study showed that SARS-Cov-2 hijacks the host polyamines for its reproduction and

infection [51].

Our findings also reveal that *Salmonella* Typhimurium enhances the polyamine production in the

host upon infection, which might be a strategy to hijack the host polyamines for its survival. This

further explains another reason for the reduced proliferation observed for the spermidine transport

mutant in macrophages. Also, the knockdown of host ornithine decarboxylase attenuates

Salmonella proliferation in macrophages. The upregulation of Odc1 activity manages to feed the

amino acid, L-arginine, into the polyamine biosynthesis and prevent nitric oxide production, which

otherwise would be detrimental for the pathogen. Polyamines have been a well-studied as a drug

target for treatment of multiple cancers. The oncogene MYC is upregulated in 70 percent of the

cancer types. Ornithine decarboxylase (Odc1) rate limiting enzyme of polyamine biosynthesis is

transcriptionally activated, directly by MYC, thereby increasing Odc1 levels in cancer [52]. RAS

is another essential factor in cell growth and cancer development, and it directly acts on Odc1 to

translationally activate Odc1 in cancer cells [53]. Polyamines biosynthesis is also associated with

multiple other oncogenes such as AKT1 and mTORC1 [54, 55]. We further show that upon using

Odc1 suicide inhibitor, DFMO, *Salmonella* proliferation could be diminished, and it reduces the

bacterial burden in mice. The FDA-approved chemopreventive drug, DFMO, for Human African

Trypanosomiasis treatment serves to be a potential drug to cure *Salmonella* infection. In the

devastating age of increasing antibiotic resistance, such a drug promises to effectively combat

deadly pathogens like *Salmonella.*

Availability of data and materials

All data generated and analyzed during this study, including the supplementary information files,

are incorporated in this article. The data is available from the corresponding author on request.

Ethics statement

All the animal experiments were approved by the Institutional Animal Ethics Committee, and the

Guidelines provided by National Animal Care were strictly followed during all experiments.

(Registration No: 435 48/1999/CPCSEA).

Acknowledgement

Prof. G. Subba Rao from MCB, IISc is duly acknowledged for providing the for knockdown

generation. Departmental Confocal Facility, Departmental Real-Time PCR Facility, Divisional

Flowcytometry facility and Central Animal Facility at IISc are duly acknowledged. Mr Sumith and

Ms Navya are acknowledged for their help in image acquisition. Mr. Amartya Mukherjee, Mr.

Sushovan Bhattacharyya and Ms Bhavya Joshi are also acknowledged for technical help.

Author contribution statement

748 AVN and DC conceived the study. AVN and DC designed experiments. AVN and AS performed

749 experiments. RSR analysed for the disease score of tissue samples. AVN, analyzed the data,

750 prepared the figures and wrote the manuscript draft. AVN, AS and DC reviewed and edited the

751 manuscript. DC supervised the work. All the authors read and approved the manuscript.

Funding

This work is supported by the Department of Biotechnology (DBT), Ministry of Science and

Technology, the Department of Science and Technology (DST), Ministry of Science and

Technology. DC acknowledges DAE-SRC ((DAE00195) outstanding investigator award and funds

and ASTRA Chair Professorship funds. The authors jointly acknowledge the DBT-IISc partnership

program. Infrastructure support from ICMR (Center for Advanced Study in Molecular Medicine),

DST (FIST), UGC-CAS (special assistance), and TATA fellowship is duly acknowledged. AVN

acknowledges the IISc-MHRD for financial assistance. AS acknowledges UGC for the financial

assistance. RSR acknowledges IISc for the financial help.

Conflict of Interest

762 The authors declare no conflict of interest.

Reference

1. Hernandez, D.J., et al., *Environmental stress destabilizes microbial networks.* The ISME Journal, 2021. **15**(6): p. 1722-1734.
2. Gray, D.A., et al., *Extreme slow growth as alternative strategy to survive deep starvation in bacteria.* Nature communications, 2019. **10**(1): p. 890.
3. Winfield, M.D. and E.A. Groisman, *Role of nonhost environments in the lifestyles of Salmonella and Escherichia coli.* Applied and environmental microbiology, 2003. **69**(7): p. 3687-3694.
4. Dodd, C.E., P.J. Richards, and T.G. Aldsworth, *Suicide through stress: a bacterial response to sub-lethal injury in the food environment.* International journal of food microbiology, 2007. **120**(1-2): p. 46-50.
5. Marmion, M., et al., *Survive and thrive: Control mechanisms that facilitate bacterial adaptation to survive manufacturing-related stress.* International Journal of Food Microbiology, 2022. **368**: p. 109612.
6. Rhee, H., E.J. Kim, and J. Lee, *Physiological polyamines: simple primordial stress molecules.* Journal of cellular and molecular medicine, 2007. **11**(4): p. 685-703.
7. Barbagallo, M., et al., *A new piece of the Shigella pathogenicity puzzle: spermidine accumulationby silencing of the speG gene.* PLoS one, 2011. **6**(11): p. e27226.
8. Johnson, L., et al., *Surface-localized spermidine protects the Pseudomonas aeruginosa outer membrane from antibiotic treatment and oxidative stress.* Journal of bacteriology, 2012. **194**(4): p. 813-826.
9. Banerji, R., P. Iyer, and S.D. Saroj, *Spermidine enhances the survival of Streptococcus pyogenes M3 under oxidative stress.* Molecular Oral Microbiology, 2022. **37**(2): p. 53-62.
10. Espinel, I.C., P.R. Guerra, and L. Jelsbak, *Multiple roles of putrescine and spermidine in stress resistance and virulence of Salmonella enterica serovar Typhimurium.* Microbial pathogenesis, 2016. **95**: p. 117-123.
11. Datsenko, K.A. and B.L. Wanner, *One-step inactivation of chromosomal genes in Escherichia coli K-12 using PCR products.* Proceedings of the National Academy of Sciences, 2000. **97**(12): p. 6640-6645.
12. Roy Chowdhury, A., et al., *Salmonella Typhimurium outer membrane protein A (OmpA) renders protection from nitrosative stress of macrophages by maintaining the stability of bacterial outer membrane.* PLoS Pathogens, 2022. **18**(8): p. e1010708.
13. Salbitani, G., C. Bottone, and S. Carfagna, *Determination of reduced and total glutathione content in extremophilic microalga Galdieria phlegrea.* Bio-protocol, 2017. **7**(13): p. e2372-e2372.
14. Achouri, S., et al., *The frequency and duration of Salmonella–macrophage adhesion events determines infection efficiency.* Philosophical Transactions of the Royal Society B: Biological Sciences, 2015. **370**(1661): p. 20140033.
15. Kim, J.-S., et al., *Oxidative stress activates transcription of Salmonella pathogenicity island-2 genes in macrophages.* Journal of Biological Chemistry, 2022. **298**(7).
16. Miller, J.F., J.J. Mekalanos, and S. Falkow, *Coordinate regulation and sensory transduction in the control of bacterial virulence.* Science, 1989. **243**(4893): p. 916-922.
17. Chattopadhyay, M.K., et al., *Polyamines stimulate the level of the σ38 subunit (RpoS) of Escherichia coli RNA polymerase, resulting in the induction of the glutamate decarboxylase-dependent acid response system via the gadE regulon.* Journal of Biological Chemistry, 2015. **290**(29): p. 17809-17821.

809   18.   Kashiwagi, K., et al., *Identification of the polyamine-induced protein as a periplasmic oligopeptide*
810   *binding protein.* Journal of Biological Chemistry, 1990. **265**(15): p. 8387-8391.
811   19.   Sakamoto, A., et al., *Three members of polyamine modulon under oxidative stress conditions:*
812   *two transcription factors (SoxR and EmrR) and a glutathione synthetic enzyme (GshA).* PLoS One,
813   2015. **10**(4): p. e0124883.
814   20.   Farr, S.B. and T. Kogoma, *Oxidative stress responses in Escherichia coli and Salmonella*
815   *typhimurium.* Microbiological reviews, 1991. **55**(4): p. 561-585.
816   21.   Bollinger, J.M., et al., *Glutathionylspermidine Metabolism in Escherichia coli.: PURIFICATION,*
817   *CLONING, OVERPRODUCTION, AND CHARACTERIZATION OF A BIFUNCTIONAL*
818   *GLUTATHIONYLSPERMIDINE SYNTHETASE/AMIDASE∗.* Journal of Biological Chemistry, 1995.
819   **270**(23): p. 14031-14041.
820   22.   Krauth-Siegel, R.L. and H. Lüdemann, *Reduction of dehydroascorbate by trypanothione.*
821   Molecular and biochemical parasitology, 1996. **80**(2): p. 203-208.
822   23.   Ariyanayagam, M.R. and A.H. Fairlamb, *Ovothiol and trypanothione as antioxidants in*
823   *trypanosomatids.* Molecular and biochemical parasitology, 2001. **115**(2): p. 189-198.
824   24.   Chattopadhyay, M.K., W. Chen, and H. Tabor, *Escherichia coli glutathionylspermidine*
825   *synthetase/amidase: phylogeny and effect on regulation of gene expression.* FEMS microbiology
826   letters, 2013. **338**(2): p. 132-140.
827   25.   Wyllie, S., et al., *Dissecting the essentiality of the bifunctional trypanothione synthetase-amidase*
828   *in Trypanosoma brucei using chemical and genetic methods.* Molecular microbiology, 2009.
829   **74**(3): p. 529-540.
830   26.   Kerrinnes, T., et al., *Utilization of host polyamines in alternatively activated macrophages*
831   *promotes chronic infection by Brucella abortus.* Infection and immunity, 2018. **86**(3): p.
832   10.1128/iai. 00458-17.
833   27.   Kingsnorth, A., A.B. Lumsden, and H.M. Wallace, *Polyamines in colorectal cancer.* Journal of
834   British Surgery, 1984. **71**(10): p. 791-794.
835   28.   Evageliou, N.F. and M.D. Hogarty, *Disrupting polyamine homeostasis as a therapeutic strategy for*
836   *neuroblastoma.* Clinical Cancer Research, 2009. **15**(19): p. 5956-5961.
837   29.   Tsoi, T.-H., et al., *Urinary polyamines: a pilot study on their roles as prostate cancer detection*
838   *biomarkers.* PLoS One, 2016. **11**(9): p. e0162217.
839   30.   Choi, Y., et al., *Targeting ODC1 inhibits tumor growth through reduction of lipid metabolism in*
840   *human hepatocellular carcinoma.* Biochemical and biophysical research communications, 2016.
841   **478**(4): p. 1674-1681.
842   31.   Soda, K., *The mechanisms by which polyamines accelerate tumor spread.* Journal of Experimental
843   & Clinical Cancer Research, 2011. **30**: p. 1-9.
844   32.   Weeks, C., et al., *alpha-Difluoromethylornithine, an irreversible inhibitor of ornithine*
845   *decarboxylase, inhibits tumor promoter-induced polyamine accumulation and carcinogenesis in*
846   *mouse skin.* Proceedings of the National Academy of Sciences, 1982. **79**(19): p. 6028-6032.
847   33.   Horn, Y., P.J. Schechter, and L.J. Marton, *Phase I–II clinical trial with alpha-*
848   *difluoromethylornithine—an inhibitor of polyamine biosynthesis.* European Journal of Cancer and
849   Clinical Oncology, 1987. **23**(8): p. 1103-1107.
850   34.   Lewis, E.C., et al., *A subset analysis of a phase II trial evaluating the use of DFMO as maintenance*
851   *therapy for high-risk neuroblastoma.* International journal of cancer, 2020. **147**(11): p. 3152-
852   3159.
853   35.   Roberts, S. and B. Ullman, *Parasite polyamines as pharmaceutical targets.* Current
854   Pharmaceutical Design, 2017. **23**(23): p. 3325-3341.
855   36.   Eperon, G., et al., *Treatment options for second-stage gambiense human African*
856   *trypanosomiasis.* Expert review of anti-infective therapy, 2014. **12**(11): p. 1407-1417.

37. Li, H., et al., *Activities of arginase I and II are limiting for endothelial cell proliferation.* American
Journal of Physiology-Regulatory, Integrative and Comparative Physiology, 2002. **282**(1): p. R64-
R69.
38. Bojarska, J., et al., *The first insight into the supramolecular system of D, L-α-*
difluoromethylornithine: A new antiviral perspective. Frontiers in Chemistry, 2021. **9**: p. 679776.
39. Bauer, P.M., et al., *Nitric Oxide Inhibits Ornithine Decarboxylase viaS-Nitrosylation of Cysteine*
360 in the Active Site of the Enzyme. Journal of Biological Chemistry, 2001. **276**(37): p. 34458-
34464.
40. HaragaA, O.M., *MillerSI.* Salmonellaeinterplaywith hostcells, 2008. **6**(1): p. 53G66.
41. Buchmeier, N.A. and F. Heffron, *Induction of Salmonella stress proteins upon infection of*
macrophages. Science, 1990. **248**(4956): p. 730-732.
42. Shen, S. and F.C. Fang, *Integrated stress responses in Salmonella.* International journal of food
microbiology, 2012. **152**(3): p. 75-81.
43. Rhen, M., *Salmonella and reactive oxygen species: a love-hate relationship.* Journal of innate
immunity, 2019. **11**(3): p. 216-226.
44. Kumar, N., R. Basundra, and S. Maiti, *Elevated polyamines induce c-MYC overexpression by*
perturbing quadruplex–WC duplex equilibrium. Nucleic acids research, 2009. **37**(10): p. 3321-
3331.
45. Thomas, T., et al., *Polyamine-mediated conformational perturbations in DNA alter the binding of*
estrogen receptor to poly (dG-m5dC). poly (dG-m5dC) and a plasmid containing the estrogen
response element. The Journal of Steroid Biochemistry and Molecular Biology, 1995. **54**(3-4): p.
89-99.
46. Prosseda, G., et al., *Shedding of genes that interfere with the pathogenic lifestyle: the Shigella*
model. Research in microbiology, 2012. **163**(6-7): p. 399-406.
47. Lee, J., et al., *An alternative polyamine biosynthetic pathway is widespread in bacteria and*
essential for biofilm formation in Vibrio cholerae. Journal of Biological Chemistry, 2009. **284**(15):
p. 9899-9907.
48. Nasrallah, G.K., et al., *Legionella pneumophila requires polyamines for optimal intracellular*
growth. Journal of bacteriology, 2011. **193**(17): p. 4346-4360.
49. Chaturvedi, R., et al., *Spermine oxidase mediates the gastric cancer risk associated with*
Helicobacter pylori CagA. Gastroenterology, 2011. **141**(5): p. 1696-1708. e2.
50. Chaturvedi, R., et al., *Induction of polyamine oxidase 1 by Helicobacter pylori causes macrophage*
apoptosis by hydrogen peroxide release and mitochondrial membrane depolarization. Journal of
Biological Chemistry, 2004. **279**(38): p. 40161-40173.
51. Makhoba, X.H. and S. Makumire, *The capture of host cell's resources: The role of heat shock*
proteins and polyamines in SARS-COV-2 (COVID-19) pathway to viral infection. Biomolecular
Concepts, 2022. **13**(1): p. 220-229.
52. Bachmann, A.S. and D. Geerts, *Polyamine synthesis as a target of MYC oncogenes.* Journal of
Biological Chemistry, 2018. **293**(48): p. 18757-18769.
53. Origanti, S. and L.M. Shantz, *Ras transformation of RIE-1 cells activates cap-independent*
translation of ornithine decarboxylase: regulation by the Raf/MEK/ERK and phosphatidylinositol
3-kinase pathways. Cancer research, 2007. **67**(10): p. 4834-4842.
54. Gomes, A.P., T. Schild, and J. Blenis, *Adding polyamine metabolism to the mTORC1 toolkit in cell*
growth and cancer. Developmental cell, 2017. **42**(2): p. 112-114.
55. Dai, F., et al., *Extracellular polyamines-induced proliferation and migration of cancer cells by*
ODC, SSAT, and Akt1-mediated pathway. Anti-Cancer Drugs, 2017. **28**(4): p. 457-464.

903