TITLE
In vitro models for liver organogenesis and synthetic tissues including assembloid formation and multiple modes of collective migration

AUTHORS AND AFFILIATIONS:
Simran Kumar[1], Jenna Venturo[1], Helly Patel[1], Natesh Parashurama[1,2,3,4,5]

[1] Department of Biomedical Engineering, University at Buffalo (State University of New York), Furnas Hall, Buffalo, NY 14260
[2] Clinical and Translation Research Center (CTRC), University at Buffalo (State University of New York), 875 Ellicott St., Buffalo, NY 14203
[3] Department of Chemical and Biological Engineering, University at Buffalo (State University of New York), Furnas Hall, Buffalo, NY 14260
[4] Center for Cell, Gene and Tissue Engineering, University at Buffalo, State University of New York, Furnas Hall Buffalo NY 14260
[5] Khufu Therapeutics, Inc. Buffalo, NY, 14260

Corresponding Author:
Natesh Parashurama, 906 Furnas Hall, Buffalo, NY 14260; Tel: 716-645-1201; Fax: 716-645-3822; e-mail: nateshp@buffalo.edu

KEYWORDS:
organogenesis, liver development, collective cell migration, epithelial to mesenchymal transition, liver diverticulum, liver bud

SUMMARY:
Organoids revolutionize personalized tissue modeling for organ development, drug discovery, and disease research. Organoid engineering extends this to create more extensive synthetic tissues. We aim to merge morphogenesis, assembloid technology, and biomatrices to advance tissue engineering. Our methods aid in modeling liver organogenesis and establishing guidelines for synthetic tissue construction.

ABSTRACT:
Chronic liver disease has reached epidemic proportions, affecting over 800 million people globally. The current treatment, orthotopic liver transplantation, has several limitations. Promising solutions have emerged in the field of liver regenerative medicine, with liver organogenesis holding significant potential. Early liver organogenesis, occurring between E8.5 and 11.5, involves the formation of epithelial-mesenchymal interactions leading to morphogenesis, hepatic cord formation, and collective migration. However, there is a lack of methods for in vitro modeling of this process. In this study, we present a detailed series of methods enabling the modeling of various stages and aspects of liver organogenesis. In one method series, we utilize assembloid technology with hepatic and mesenchymal spheroids, which replicate early structures found in liver organogenesis, model early morphogenesis, and demonstrate interstitial cell migration as seen in vivo. These innovative assembloid systems help identify factors influencing assembloid formation and migration. Hepatic spheroid cultivation systems were also employed to model collective

47 migration and branching morphogenesis. Fibroblast-conditioned media play a significant
48 role in initiating dose-dependent branching migration. Future work will involve high
49 temporal and spatial resolution imaging of hepatic and mesenchymal interactions to
50 determine the cascade of cellular and molecular events involved in tissue formation,
51 morphogenesis, and migration.
52
INTRODUCTION:
Liver cell migration plays a significant role in liver organogenesis, disease, and cell
therapy. During liver organogenesis (E8.5-9.0, mouse), the ventral foregut pre-hepatic
epithelium begins to express liver genes, due to the inductive signals emanating from the
surrounding mesenchyme and heart. At E9.0, the foregut epithelium thickens as the cells
transition from a cuboidal to a pseudostratified columnar morphology, to form the liver
diverticulum (**Gualdi, Bossard et al. 1996**); (**Bort, Signore et al. 2006**). At this critical
stage, the liver diverticulum is comprised of only ~1,500 cells. Next, the hepatic
endoderm lining the liver diverticulum thickens, delaminates, and forms cords of
hepatoblasts that co-migrate with endothelial cells and mesenchymal cells and branch
into the surrounding mesenchymal tissue, thus initiating three-dimensional collective cell
migration to form the liver bud (**Ogoke, Oluwole et al. 2017**); (**Ogoke O. 2022**). In fact,
during this stage, the cells collectively undergo; 1) co-migration, or movement together
with other cell types, 2) branching morphogenesis or formation of branching tube-like
structures, and, 3) interstitial migration, or migration on top of other cells. By E11.5,
migration ceases, the primitive liver has formed and has expanded 10^3-fold (**Ogoke O.
2022**). Liver cell migration may also be required in later stages of liver organogenesis,
as rat fetal hepatoblasts (HBs) expression have shown evidence of highly upregulated
genes associated with 3D collective cell migration, morphogenesis, and extracellular
matrix remodeling **(Petkov, Kim et al. 2000**). In addition to its role in early liver
organogenesis, 3D collective migration is intricately linked to the local spread and
metastasis of advanced hepatocellular carcinoma (HCC), ultimately leading to worsened
prognosis and increased treatment resistance **(Yang, Chen et al. 2009**). Adult and fetal
hepatocytes also employ collective migration when moving from the spleen to within the
liver during liver repopulation; *in vivo* imaging studies have demonstrated that
transplanted hepatocytes enter the portal vein and then the capillaries within hours,
migrate across the liver sinusoids, and through the liver tissue (**Rajvanshi, Kerr et al.
1996**);(**Gupta, Rajvanshi et al. 1999**);(**Koenig, Stoesser et al. 2005**). Finally, recent
studies demonstrate that migrating hepatoblasts arise during murine and human liver
regeneration with some evidence of movement in sheets (**Matchett KP 2023**). Overall,
liver collective migration, capable of multiple modes of morphogenesis, plays a significant
role in organogenesis, cancer, hepatocyte cell therapy, and liver regeneration.
85
Numerous genetic studies have investigated the molecular pathways that drive 3D liver
collective cell migration. These studies demonstrate that ablation of the hepatic cords
blocks liver formation and demonstrates that therefore, formation of hepatic cords and
their ensuing interactions with supporting cells are required for liver formation (**Bort,
Signore et al. 2006**); (**Suzuki, Sekiya et al. 2008**); (**Sosa-Pineda, Wigle et al. 2000,
Matsumoto, Yoshitomi et al. 2001**). These studies also demonstrate that liver growth is
initiated by fibroblast growth factor 2 (FGF2) secreted from the cardiac mesoderm, BMP4

secreted from the surrounding mesenchyme, HGF, endothelial cell interactions, and migration-associated transcription factors including HEX, PROX1, and TBX3 (**Gualdi, Bossard et al. 1996**); (**Rossi, Dunn et al. 2001**). Overall, genetic studies support the fact that soluble factor signaling with transcription factor expression is responsible for driving migration, signaling, and molecular interactions between hepatoblasts and their surrounding mesenchyme.

Although cell migration in early liver organogenesis has been extensively investigated, the current *in vitro* hepatic migration studies frequently utilize 2D assays consisting of highly migratory HCC cells combined with *in vivo* tumor models (**Ng, Tung-Ping Poon et al. 2013**). These studies have provided insight into several factors that play a role in hepatic migration including TGFB1 (**Fransvea, Angelotti et al. 2008**) c-Myc (**Zhao, Jian et al. 2013**), Yes associate protein (YAP) (**Fitamant, Kottakis et al. 2015**), goosecoid (**Xue, Ge et al. 2014**), actopaxin (**Binamé, Lassus et al. 2008**), and miRNAs (**Zeng, Liang et al. 2016**);(**Chen, Liang et al. 2017**);(**Yang, Xu et al. 2017**). Despite the advancements in understanding the molecular mechanisms in 3D hepatic cell migration, the fundamental mechanisms between 2D and 3D cellular migration are distinct which suggests 2D assays have their limitations. Furthermore, these models typically do not implement mesenchymal cell types, which are essential to migration/growth. There has been progress in the development of 3D models for liver migration that incorporate the supporting mesenchyme, however, they are solely focused on co-migration rather than the different modes of collective migration.

The ability to form tissues from spheroids through various self-assembly and morphogenetic processes enables the scientific study of synthetic tissues for applications for drug development and screening, disease modeling, therapy, and other biomedical and biotechnological applications. Here we present methodological details for several 3D *in vitro* cultivation systems which were engineered to exhibit different modes of liver 3D collective migration. These systems include the following: (1) co-spheroid culture with hepatic and mesenchymal-derived spheroids in matrix, (2) spheroid matrix droplet cultured with fibroblast conditioned medium, and (3) mixed spheroids (hepatic and mesenchymal-derived cells). These systems enable robust modeling of liver 3D collective migration which will improve our molecular and cellular understanding of liver organogenesis, cancer, and therapy.

PROTOCOL:

1. Preparation of 1% Low EEO Agarose Solution

1.1. Measure 2.5 g of agarose powder (low EEO) and transfer it to a beaker.

NOTE: The beaker should be at least twice the size of the desired volume to account for the bubbling of the solution.

1.2. Use a graduated cylinder to measure 250 mL of distilled water (DI) water and transfer it to the beaker to dilute the agarose to obtain a final concentration of 1%.

1.3. Cover the mouth of the beaker with plastic wrap and make a small hole. Heat the
beaker in the microwave.

1.4. After 30 seconds, remove the beaker and swirl until uniform. Repeat every 30
seconds, until the agarose completely dissolves.

CAUTION: Microwaved glassware should be handled very carefully by wearing proper
gloves. The solution should be watched closely to avoid overheating or boiling over.

1.5. Remove the beaker from the microwave and gently swirl. Transfer the solution to a
pre-sterilized bottle and autoclave the solution. Store the agarose solution at room
temperature until ready to use.

2. Coating 96-Well Plate

2.1. Loosen the cap of the bottle containing the 1% agarose solution. Warm the solution
in the microwave until the solution is in the liquid phase and tighten the cap.

CAUTION: Microwaved glassware should be handled very carefully by wearing proper
gloves. The solution should be watched closely to avoid overheating or boiling over.

NOTE: Perform these steps under a sterile tissue culture laminar flow hood.

2.2. Use 55-65 µL of the sterile 1% agarose solution per well to coat the 96-well tissue
cultured plate and immediately rotate the plate.

2.3. Once the 1% agarose solution has been transferred to the desired number of wells,
allow the agarose to solidify by allowing the plates to cool for 20-30 minutes in a 4℃
fridge. Prior to use, bring the plate to room temperature (**Figure 1A**).

3. Preparation of HepG2-WT Spheroids

3.1. Cultivate HepG2-WT cells in a T-75 flask with completed growth medium (cDMEM)
containing Dulbecco's Modified Eagle Medium (DMEM), supplemented with 10%
Fetal Bovine Serum (FBS) and 1% Penicillin-Streptomycin (Pen-Strep). Incubate the
cell culture at 37℃ and 5% CO_2 with medium changes every day.

3.2. Once the cell culture reaches 80% confluency, add 0.05% of Trypsin-EDTA to the
flask for 5-10 minutes. Add equal amounts of cDMEM and wash the cells off the
flask.

3.3. Once the cells have detached, transfer the mixture to a 15 mL sterile conical
centrifuge tube and centrifuge the cell suspension at 300 $x\ g$ for 5 minutes.

3.4. Re-suspend the cell pellet in sterile 1X Phosphate Buffered Saline (PBS) and
centrifuge the cell suspension at 300 $x\,g$ for 5 minutes.
186
187 3.5. Based on the cell count, suspend the cell suspension to obtain a final concentration
188   of 1x 10^6 cells/mL (**Figure 1B**).
189
190 3.6. Dye-labeling of cells
191
192 NOTE: This is an optional step.
193
194 3.6.1. Transfer the desired amount of cell suspension to a 15 mL sterile conical centrifuge
195   tube and centrifuge the cell suspension at 300 $x\,g$ for 3 minutes.
196
197 3.6.2. Re-suspend the cell pellet in a serum-free growth medium to obtain a final
198   concentration of 1 x 10^6 cells/mL. Add 5 µL of Vybrant Cell-Labeling Solution per
199   mL of cell suspension and incubate the cell suspension on rotation for 20 minutes,
200   preferably at 37°C.
201
202 NOTE: Serum-free growth medium is DMEM only supplemented with 1% Pen-strep.
203 Different densities of the cell suspension may require longer incubation time for uniform
204 staining.
205
206 3.6.3. Once the incubation is completed, centrifuge the cell suspension at 450 $x\,g$ for 5
207   minutes and resuspend the cell pellet in fresh cDMEM. Repeat this wash process
208   two more times (**Figure 1C**).
209
210 3.7. Spheroid formation
211
212 3.7.1. Suspend the cells in fresh cDMEM to obtain a final concentration of 5.0 x 10^4
213   cells/mL. Mix the cell suspension very well and transfer 100 µL of cell suspension
214   per well to the agarose-coated 96-well plate.
215
216 NOTE: The density of cell suspension is to obtain a density of 5,000 cells per well in the
217 agarose-coated 96-well plate.
218
219 3.7.2. Centrifuge the plate at 340 $x\,g$ for 10 minutes and incubate at 5% CO_2 at 37°C for
220   5-9 days (**Figure 1D**).
221
222 NOTE: Change medium every other day after plating with gentle removal of 50% of
223 cDMEM and replacement.
224
225 NOTE: HepG2-WT spheroids can be used for the HEP-MES assembloid model or M-CM
226 model.
227
228 **4. Preparation of HFF/MRC-5 Spheroids**
229

230   4.1. Cultivate HFF/MRC-5 cells in a T-175 flask with completed growth medium (cDMEM)
231   containing Dulbecco's Modified Eagle Medium (DMEM), supplemented with 10%
232   Fetal Bovine Serum (FBS) and 1% Penicillin-Streptomycin (Pen-Strep). Incubate the
233   cell culture at 37℃ and 5% CO_2 with medium changes every other day.
234
235   4.2. Once the cell culture reaches 80% confluency, add 5 mL of 0.25% of Trypsin-EDTA
236   to the flask for 5-10 minutes. Add equal amounts of cDMEM and wash the cells off
237   the flask.
238
239   4.3. Once the cells have detached, transfer the mixture to a 15 mL sterile conical
240   centrifuge tube and centrifuge the cell suspension at 300 $x\ g$ for 3 minutes.
241
242   4.4. Re-suspend the cell pellet in sterile 1X Phosphate Buffered Saline (PBS) and
243   centrifuge the cell suspension at 300 $x\ g$ for 3 minutes.
244
245   4.5. Based on the cell count, suspend the cell suspension to obtain a final concentration
246   of 1x 10^6 cells/mL (**Figure 1B**).
247
248   4.6. Dye-labeling of Cells
249
250   NOTE: This is an optional step.
251
252   4.6.1. Transfer the desired amount of cell suspension to a 15 mL sterile conical centrifuge
253   tube and centrifuge the cell suspension at 300 $x\ g$ for 3 minutes.
254
255   4.6.2. Re-suspend the cell pellet in a serum-free growth medium to obtain a final
256   concentration of 1 x 10^6 cells/mL. Add 5 µL of Vybrant Cell-Labeling Solution per
257   mL of cell suspension and incubate the cell suspension on rotation for 20 minutes,
258   preferably at 37°C.
259
260   NOTE: Serum-free growth medium is DMEM only supplemented with 1% Pen-strep.
261   Different densities of the cell suspension may require longer incubation time for uniform
262   staining.
263
264   4.6.3. Once the incubation is completed, centrifuge the cell suspension at 450 $x\ g$ for 5
265   minutes and resuspend the cell pellet in fresh cDMEM. Repeat this wash process
266   two more times (**Figure 1C**).
267
268   4.7. Spheroid Formation
269
270   4.7.1. Suspend the cells in fresh cDMEM to obtain a final concentration of 5.0 x 10^4
271   cells/mL. Mix the cell suspension very well and transfer 100 µL of cell suspension
272   per well to the agarose-coated 96-well plate.
273
274   NOTE: The density of cell suspension is to obtain a density of 10,000 cells per well in the
275   agarose-coated 96-well plate.

276
277 4.7.2. Centrifuge the plate at 340 *x g* for 10 minutes and incubate at 5% CO_2 at 37°C for
278   5-9 days (**Figure 1D**).
279
280 NOTE: Change medium every other day after plating with gentle removal of 50% of
281 cDMEM and replacement.
282
283 NOTE: HFF/MRC-5 spheroids can be used for the HEP-MES assembloid model.
284
285 **5. HepG2-WT and HFF/MRC-5 Assembloid Formation**
286
287 NOTE: Refer to Sections 3 and 4 for formation of HepG2 (**Figure 2A)** and HFF/MRC-5
288 spheroids (**Figure 2B**).
289
290 5.1. Individually collect HFF/MRC-5 spheroids using a pipette from the 96-well plate and
291   transfer them to a 15 mL sterile conical centrifuge tube. Allow the spheroids to settle
292   and gently rinse with warm cDMEM.
293
294 NOTE: This rinsing process should be done very gently and carefully.
295
296 5.2. Transfer a single HFF/MRC-5 spheroid to a well containing a HepG2-WT spheroid
297   and add MG/CG between a 1:1 and 1:5 dilution and incubate at 37°C at 5% CO_2 for
298   3 hours.
299
300 5.3. Add 75 uL of cDMEM to each well and incubate at 37°C at 5% CO_2 for 2-3 days
301   (**Figure 2C**).
302
303 NOTE: Change medium every other day after plating with gentle removal of 50% of
304 cDMEM and replacement. Assembloids will still form without media changes for up to 3-
305 4 days.
306
307 NOTE: Assembloid formation can occur without the use of matrix.
308
309 **6. HepG2-WT Spheroid Droplet Formation**
310
311 NOTE: Refer to Section 3 for formation of HepG2 spheroids (**Figure 3A**).
312
313 NOTE: Two different materials can be used for suspending the HepG2-WT spheroids in
314 droplets. The two methods are provided below.
315
316 6.1. Matrigel (MG) droplets
317
318 6.1.1. Mix 1 mL of ice-cold diluted MG and control growth medium at a 1:1 dilution. Mix
319   the spheroid/MG suspension and distribute it evenly inside the MG solution.
320

321 6.1.2. Collect the HepG2-WT spheroids in a 15 mL sterile conical centrifuge tube on ice
322   and allow the spheroids to settle. Aspirate the medium and keep it on ice.
323
324 6.1.3. Using a 200 μL pipette, collect a 15 μL volume of one spheroid in MG solution and
325   seed onto a 60 mm petri dish (**Figure 3B**).
326
327 6.2. Collagen (CG) Droplets
328
329 NOTE: All collagen preparation should be done on ice.
330
331 6.2.1. In a microcentrifuge tube, add 358.8 μL of de-ionized water, 100 μL of 10X PBS,
332   12.1 μL of 1 N NaOH, and 529.1 μL of stock rat tail CG for a total volume of 1 mL.
333   Mix the spheroid/CG suspension and distribute it evenly inside the CG solution.
334
335 NOTE: Stock rat tail CG should always be added at the end.
336
337 6.2.2. Collect the HepG2-WT spheroids in a 15 mL sterile conical centrifuge tube on ice
338   and allow the spheroids to settle. Aspirate the medium and keep it on ice (**Figure
339   3B**).
340
341 6.2.3. Using a 200 μL pipette, collect a 15 μL volume of one spheroid in CG solution and
342   seed onto a 60 mm petri dish.
343
344 NOTE: If more than one spheroid is seeded per droplet, it is removed and reseeded
345 properly.
346
347 NOTE: Spheroid/CG solutions are pipetted slowly onto the 60 mm petri dish to avoid air
348 bubbles.
349
350 6.3 Incubate the droplet at 37°C for 60 minutes before the addition of the growth medium.
351
352 6.4 Slowly add 5 mL of desired growth medium to the petri dish and incubate in at 37°C
353   and 5% CO_2 with medium changes every three days (**Figure 3C**).
354
355 NOTE: HFF/MRC-5 conditioned-media was used in the droplet formation assay.
356
357 **7. Preparation of HFF/MRC-5 Conditioned Media (M-CM)**
358
359 7.1. Seed HFF/MRC-5 into a T-75 tissue culture-treated flask at a seeding density of
360   5,000 cells/cm^2 and incubate the flask for 72 hours in 15 mL of cDMEM.
361
362 NOTE: Flask should be checked daily during this period to ensure the maintenance of
363 cell health.
364

365 7.2. After the 72-hour incubation, collect the fibroblast-conditioned growth medium in a
366   15 mL sterile conical centrifuge tube. Centrifuge the fibroblast-conditioned growth
367   medium at 290 *x g* for 5 minutes and filter using a 0.2 µm filter (**Figure 4A**).

369 7.3. Dilute the fibroblast-conditioned growth medium with complete growth medium at a
370   1:1 to 1:7 dilution ratio and add to the desired experiment (**Figure 4B**).

8. HepG2-WT and HFF/MRC-5 Mixed Spheroid Formation

NOTE: Refer to Sections 3 and 4 for formation of HepG2 (**Figure 5A)** and HFF/MRC-5
spheroids (**Figure 5B**).

8.1. Transfer HepG2-WT and HFF/MRC-5 cell suspension to a 15 mL sterile conical
centrifuge tube at a 1:1 ratio to obtain a final concentration of 2.0 x 10^5 cells/mL. Mix
the cell suspension very well and transfer 100 µL of cell suspension per well to the
agarose-coated 96-well plate.

NOTE: The density of cell suspension is to obtain a density of 20,000 cells per well in the
agarose-coated 96-well plate.

8.2. Centrifuge the plate at 340 *x g* for 10 minutes and incubate at 5% CO_2 at 37°C for 1-
2 days (**Figure 5C**).

NOTE: Change medium every other day after plating with gentle removal of 50% of
cDMEM and replacement.

REPRESENTATIVE RESULTS:
Currently, there is increased interest in synthetic tissues for various biomedical
applications, including modeling disease, discovering drugs, and tissue engineering
(**Figure 6**). In this field, hPSC-derived organoids or spheroids, and cells can be converted
into more synthetic, complex, and larger tissues. To accomplish this, principles of
morphogenesis, tools like microfabrication, and biomatrices can be applied to cells and
spheroids to engineer these synthetic tissues more precisely (**Figure 6**). We present
several methods here with this theme in mind.

Effects of clustering on spheroid formation
The methods developed here were contingent upon successful 3D spheroid formation.
Spheroid formation is considered successful if cells fuse to form a full spheroid within five
to nine days. An early indication of successful spheroid formation is clustering of the cells
in the center of the well after centrifugation of the cultivation plate. Despite the significance
of clustering, spheroid formation did occur, but less frequently, when cells were initially
scattered rather than clustered, therefore demonstrating that successful spheroid
formation could still occur. Spheroid formation is considered unsuccessful if cells do not
spread and fuse together within nine days after plating.
Spheroids were cultured at two different sizes to perform this experiment; small
spheroids (S) were plated at a concentration of 1,500 cells per well and large spheroids

(L) were plated at a concentration of 3,000 cells per well. Hepatic spheroids compacted
and fully cultured by day five, irrespective of the spheroid size. The cell density per well
did not impact the rate at which the spheroid formed and had a significant difference in
spheroid size (**Figure 7**). In this case, it is important to observe an increase in opacity
which demonstrates thickening of the initial disc-shaped tissue, which is more translucent,
to a spheroid configuration, which has increased opacity. Unlike hepatic spheroids,
mesodermal-derived spheroids compacted within 24 hours irrespective of cell seeding
density as well. Notably, HFF/MRC-5 cells compact much tighter than HepG2-WT cells,
likely resulting in cell density having little impact on spheroid size.
420

Factors that affect hepatic and mesenchymal (mesodermal-derived) assembloid formation

Hepatic and mesodermal-derived spheroids of varying sizes were cultured to determine
the effect of size on the compaction time of spheroid formation. Hepatic and mesodermal-
derived spheroids were co-cultured in Matrigel (MG) or Collagen Gel (CG) at a 1:5 dilution
with complete growth medium or fibroblast-conditioned medium, to determine if matrix
and conditioned medium influences assembloid formation. Both cell types were dye-
labeled prior to spheroid formation to demonstrate the interaction between spheroids. It
was observed that assembloid formation occurs irrespective of matrix and medium
(**Figure 8**). We studied the effects of CG, effects of MG, and effects of conditioned
medium (MRC-5 conditioned medium or MCM-5). We observed assembloid formation in
all cases, although the morphological details varied slightly (**Figure 8**). Details regarding
these images will be re-used in later figures. To determine the effects of inter-spheroid
distance on assembloid formation, distance was measured together with success of
assembloid formation. It was observed that the compaction time of an assembloid is
directly proportional to the initial distance of the hepatic and mesodermal-derived
spheroids (**Table 1**).
438

Building more complex assembloids with arm-like structures

Methods were also developed to build assembloids that have branching cords (**Figure**
9A). To accomplish this experimentally, hepatic spheroids are mixed with biomatrix (MG),
in 384-well plate, and surrounded by single fibroblasts at high density. These fibroblasts
cluster and provide guides to which hepatic cells migrate towards and thicken, forming
thick cords over time. Hepatic spheroids were cultivated in the MG droplet system
containing a high density of MRC-5 cells (300,000 cells) in 384-wells and demonstrated
small clusters of MRC-5 cells that formed in the MG (**Figure 9B, days 3-4**). Next, the liver
spheroids formed thick migrating strands protruding out to the fibroblast clusters forming
thick strands containing both cell lines (**Figure 9B, days 9-12**). This approach led to
longer arms or cords, likely containing a mix of hepatic and fibroblast cells. Another
approach involved building bridges or small interconnections (arms) between spheroids.
To build small armed structures, a larger HEP spheroid can be co-cultured in 384-well
plate with a smaller mixed spheroid (**Figure 9C**). This leads to small, knob-like arms.
Overall, we present two approaches for forming additional arms to spheroids.
454

Spot-welding (fused edges) of complex assembloids

456 Methods were used to build assembloid with fused edges. Hepatic, and mesodermal-

457 derived spheroids are placed in a stiff environment of CG (2 mg/mL) **(Figure 9D)**.
458 Furthermore, in this stiff environment, rather than cupping, we observe spheroids fuse at
459 the edges to form assembloids with evidence of short arms, or spot-welding (**Figure 9E**).
460 We see similar data with MRC5 Fibroblasts in CG (**Figure 9F**).
461
462 ***Infiltrating and layering of complex assembloids***
463   During liver organogenesis, in the developing liver diverticulum, HEP cells are
surrounded by mesenchyme, and ultimately, they migrate or infiltrate into the
mesenchyme (**Figure 9G**). In M-CM, and CG conditions, the addition of MES and HEP
spheroids results in a different type of fusion in which we observe an infiltrative pattern
(**Figure 9H**). Further, we can obtain a layering pattern by placing single MES (human
mesenchymal stem cells (hMSC)) in MG at high density and allowing them to migrate
towards a HEP spheroid and layer on the surface without infiltration, as shown with dye
labeling of the hMSC (**Figure 9I**). We note that this is a different phenotype than when
we employed HFF. This latter arrangement is observed in the liver diverticulum stage and
many other endoderm-derived tissues. Here we present two approaches that are relevant
for modeling of the developing liver diverticulum and for creating aspects of synthetic
tissues.
475
476 ***Fused hepatic and mesenchymal (mesodermal-derived) assembloid formation***
477   The fusion of two spheroids to form assembloids are critical for modeling the liver
bud. This is because when two spheroids fuse, the cells likely migrate on top or between
other cells, which we term interstitial migration. Interstitial migration occurs in the liver
bud, when early migrating hepatoblasts migrate through mesenchyme, potentially on top
of other cells. Therefore, fusion of two spheroids is a model of interstitial migration which
occurs during liver organogenesis. Methods were developed to build assembloids that
fuse completely with separate layers (**Figure 10A**). HEP and MES (MRC5 fibroblasts)
spheroids in MG form a fused assembloid by day 9 (**Figure 10B**). Dye-labeling analysis
demonstrated that MES tissue remained inside, while the HEP tissue remained outside
(**Figure 10C**). Importantly, the final spheroid is approximately the same size as the
original spheroids. This suggests that the cells are packed at high density. Notably, the
same phenomena occur in the absence of MG in 384-well plates, when multiple MES
spheroids are placed with a single HEP spheroid (**Figure 10D**). HEP-MES assembloids,
in the absence of matrix, also form under low serum (2% FBS) and extremely low serum
(0.2% FBS) conditions (**Figure 10E**). We then determined that large distances
(approximately 3 diameters of the spheroid), assembloids did not form (**Figure 10F**). To
determine how spheroid composition determines spheroid fusion, we demonstrated that
mixed spheroids (containing MES and HEP cells) can fuse with MES spheroids by day 5,
accompanied by an increased packing density, as expected (**Figure 10G**).
496
497 ***Partially fused or cupping in complex assembloid formation***
498   Successful assembloid formation in MG demonstrates several phenotypes,
including the observation that the HEP spheroid undergoes "cupping" of the MES
spheroid to form a partially fused assembloid (**Figure 10H**). This also establishes a visual
model of interstitial migration, and points toward the current mechanism of fused
assembloid formation. Time series studies of HEP and MES spheroids demonstrate with

in CG, fusion occurs via a cup-like mechanism (**Figure 10I**). This is further demonstrated
using dye-labeling (**Figure 10J**). Finally, the same phenomena is clearly illustrated when
multiple HEP spheroids are used with a large, single MES spheroid (**Figure 10K**). The
data suggests that cupping without fusion can occur in MG or in cases where the MES
spheroid is much larger than the HEP spheroids.
508
Induced hepatic branching and linear migration via mesenchymal (mesodermal-
derived) conditioned growth medium
Collective migration is a key morphogenetic process during liver organogenesis.
Here we describe tools for inducing collective migration *in vitro*. Successful droplet
formation of HEP spheroids and utilization of M-CM (MES-conditioned media)
demonstrate outgrowth branching from the HEP spheroids (**Figure 11A-B**). HEP
spheroids cultivated in the MG droplet system demonstrated that M-CM induces collective
migration, and the data demonstrates a concentration-dependent effect (**Figure 11B,**
right). Cellular strands protruding from the hepatic spheroid are present with small
branching, thick strands, and multiple levels of branching (**Figure 11B, left**). On day 11,
the protrusions were increased with inter-connections and sheet formation (**Figure 11B,**
right). M-CM is potent, as a 1:7 dilution or M-CM present for one day still leads to
migration (**Figure 11B, right**). It was hypothesized that the M-CM induced cell migration
via TGFβ signaling pathway. A83-01, a TGFβ pathway inhibitor, was incorporated into
the M-CM at varying concentrations, and migration was significantly inhibited in a dose-
dependent manner (**Figure 11C**). To determine the effects of extracellular matrix on
collective migration, HEP spheroids cultivated in the CG droplet system demonstrate that
M-CM induces cell migration. However, the protrusions were thin linear strands and less
branching by day 7 compared to the migration observed in MG (**Figure 11D**). We also
tested fibrin hydrogels, and we observed thin, hair-like, narrow, radial protrusions of both
HEP and MES cells (**Figure 11E**).
530
Inducing hepatic co-migration via MES conditioned medium
Mixed spheroids (HEP-MES spheroids) were also employed for collective
migration, as a model of co-migration which occurs during early liver organogenesis
(**Figure 11F**). These mixed spheroids in MG result in migration, and when TGFβ1 growth
factor was added, it resulted in significantly increased collective migration (**Figure 11G**).
Thus, co-migration can also be modeled with mixed spheroids.
537
FIGURE AND TABLE LEGENDS:
Figure 1. Schematic of spheroid formation assay. (A) Coating of a 96-well plate with
1% agarose solution. The agarose solution is warmed in the microwave until it is in the
liquid phase and cooled prior to use. 55-65 µL of agarose solution is transferred to each
well and the plate is cooled in the fridge for 20-30 minutes. **(B)** Passaging of cell culture
at 80% confluency. **(C)** Dye labeling of cell suspension. 5 µL of cell-labeling solution is
added per mL of cell suspension and incubated for 20 minutes. The cell suspension is
centrifuged at 450 *x g* for 5 minutes and washed with warm cDMEM three times prior to
plating the cells. The cell suspension is diluted to the desired cell density per well. **(D)**
Plating of cell suspension for spheroid formation. 100 µL of cell suspension is added to
each well and centrifuged at 340 *x g* for 10 minutes. Mesodermal-derived spheroids are

incubated for 1-3 days until spheroid formation is complete. Hepatic spheroids are
incubated for 5-9 days until spheroid formation is complete.
551
Figure 2. Schematic of assembloid formation assay with hepatic and mesodermal-
derived spheroids. (A) Hepatic spheroid formation. HepG2-WT cells are dye-labeled
with a cell-labeling solution and plated for spheroid formation. **(B)** Mesodermal-derived
spheroid formation. HFF/MRC-5 cells are dye-labeling solution and plated for spheroid
formation. **(C)** Transferring mesodermal-derived spheroid to hepatic spheroid for
assembloid formation. Matrix (Matrigel or Collagen) is added to the well at a 1:1-1:5
dilution range and incubated for 3 hours. 75 µL of cDMEM is added to the well and is
incubated for 2-3 days until assembloid formation is complete.
560
Figure 3. Schematic of Droplet Formation Assay. (A) Hepatic spheroid formation.
HepG2-WT cells are dye-labeled with a cell-labeling solution and plated for spheroid
formation. **(B)** Hepatic spheroids are collected and gently rinsed with cDMEM. The
spheroids are suspended in a matrix (Matrigel or Collagen). **(C)** Hepatic spheroids are
seeded onto a 60mm petri dish and incubated for 60 minutes. 5 mL of fibroblast-
conditioned media is added to the petri dish and incubated with media changes every
three days.
568
569 **Figure 4. Schematic of Preparation of Fibroblast-Conditioned Media. (A)** Cells are
570 seeded at 5,000 cells/cm2 in a T-75 and incubated for 72 hours. The media is collected
571 and sterilized prior to use. **(B)** Sterilized conditioned media is diluted with cDMEM at a
572 1:1-1:7 dilution and added to a hepatic spheroid in a pre-existing well. The spheroid is
573 incubated, and the media is changed every three days.
574
Figure 5. Schematic of Hepatic and Mesodermal-derived Mixed Spheroid Formation
Assay. (A) Preparation of Hepatic cell suspension. HepG2-WT cells are harvested and
dye-labeled with a cell-labeling solution. **(B)** Preparation of Mesodermal-derived cell
suspension. HFF/MRC-5 cells are harvested and dye-labeled with a cell-labeling solution.
(C) Mixed Spheroid Formation. HepG2-WT and HFF/MRC-5 cells are mixed at a 1:1 ratio
at a final concentration of 2.0 x 105 cells/mL and plated for spheroid formation. The plate
is incubated for 1-2 days until mixed spheroid formation is complete.
582
583 **Figure 6. Overview.** Based in the science of morphogenesis (top), stem cells will be
584 implemented to build assembloids. The toolbox (middle), through the use of cells, tools,
585 and biomatrix, and together with aspects of morphogenesis, can together be used to
586 generate new synthetic tissues and assembloids.
587
588 **Figure 7. Compaction time of varying HepG2-WT spheroid size.** Progression of
589 hepatic spheroid formation of varying sizes over a five-day period. Spheroids were plated
590 at 1,500 cells per well (small) and 3,000 cells per well (large) and observed for five days.
591 Spheroid formation was observed to occur under both conditions within five days.
592
Figure 8. The effect of media and matrix on assembloid formation. Hepatic spheroids
and mesodermal-derived spheroids were transferred to an agarose-coated well. The cells

were dye-labeled prior to spheroid formation to demonstrate the interaction between
spheroids. The spheroids were suspended in a 1:5 solution of the matrix (Matrigel or
Collagen) and cDMEM or fibroblast-conditioned media (M-CM) and observed for four
days (MRC-5 is orange, HepG2-WT is green). It was observed that assembloid formation
occurred under the four conditions.

Figure 9. Engineering assembloids arms, junctions, and layers. (A) Building of large
and small arms onto hepatic spheroids. **(B)** Phase-contrast microscopy images of hepatic
spheroids containing growth factor-free (GFR) cultured in the MG droplet system, bearing
high density (30,000 cells) of MRC-5 cells were cultured for 12 days. Top row, left to right:
days 3, 4, 9, and 12. Day 3: MRC-5 cells initially after seeding (arrow). Bar = 1,000 µm.
Day 4: MRC-5 cells spreading and interconnecting (arrows). Bar = 1,000 µm. Day 9: Thick
hepatic cord (arrows). Bar = 400 µm. Day 12: multiple, thick hepatic cords. Bar = 400 µm.
Bottom row, left to right: each image in a separate experimental replicate on day 12
demonstrating thick hepatic cord formation (arrows). Bar = 400 µm. **(C)** Fluorescent
images of hepatic spheroid (figure 1) co-cultured with a HEP-MES mixed spheroid (figure
5) on days 0 and 5. It was observed on day five, that the HEP-MES mixed spheroid fused
with the edge of the hepatic spheroid to form a bridge. **(D)** Fused HEP-MES assembloids,
which are the same size as original spheroids, with the edges of the two spheroids fused
to create an assembloid. These spheroids create a higher packing density once combined
into an assembloid. **(E)** Phase-contrast images of hepatic and mesodermal-derived
spheroids in the CG droplet system cultivated in M-CM. Collagen provides stiff conditions
during assembloid formation. By day 4, it was observed that HEP-MES cultured in M-CM
fuses the edges of the spheroids and does not form cupping. **(F)** In the CG droplet system,
the HEP-MES assembloid edges are fused together. The left image is a phase-contrasted
image, and the right is a fluorescent image to show the fusion of hepatic and mesodermal-
derived spheroids. The hepatic spheroid is dye-labeled green, and the MRC-5 spheroid
is dye-labeled orange in the figure. **(G)** The left image shows HEP-MES infiltrating
assembloid and the right image demonstrates a surface-MES layered assembloid. **(H)**
Infiltration of HEP-MES spheroids in collagen matrix with MES-conditioned media (M-
CM). Left- phase-contrast image of assembloid. Right-fluorescent image showing
infiltration of HEP and MES spheroids forming assembloid. **(I)** Phase (left), double
fluorescent (red/green) images (right) of days 4, 5, and 6 LD models, bearing a HepG2-
GFP spheroid (green) and MSC (red) in MG. The right columns on the right are replicates
1 and 2 with double fluorescent images on days 4, 5, and 6. This is an example of layering
without infiltration of the spheroids to form an assembloid. Data was replicated to prove
the outcome described. Bar = 500 µm. Adopted with permission of publisher from Ogechi,
Parashurama et al., 2021 (Ogechi et al., 2021).

Figure 10. Partially and completely fused HEP-MES assembloids. (A) Fused HEP-
MES assembloids, which are the same size as the original spheroids, thus increasing
packing density. **(B)** Phase-contrast images of HEP and MES spheroids cultivated in MG
droplet system. It was observed that by day 11, assembloid formation had occurred which
typically occurs in 2-5 days. **(C)** Fluorescent images of HEP-MES assembloid on day 13.
It was observed that assembloid formation had occurred with separate layers, MES (red)
surrounded by the HEP (green). **(D)** Phase-contrast images of hepatic spheroid co-

cultured with multiple mesodermal-derived spheroids. By day 5, it was observed that the HEP-MES assembloid did not increase in volume. **(E)** HEP-MES assembloids fuse under low serum. HepG2-WT spheroids and HFF/MRC-5 cells were plated (figure 1) and suspended in low serum FBS (2% and 0.2%). Top- 2% FBS to fuse HEP-MES assembloid. Bottom- 0.2% FBS to fuse HEP-MES assembloid. It was observed that HEP-MES assembloid formation occurred over a period of two days. **(F)** Phase-contrast images of hepatic and mesodermal-derived spheroids in the CG droplet system cultivated in M-CM. When the spheroids are placed at large distances in the CG droplet system, assembloid formation does not occur. **(G)** Fluorescent images of hepatic spheroid (**Figure 1**) co-cultured with a HEP-MES mixed spheroid (**Figure 5**) on days 0 and 5. It was observed on day five, that the HEP-MES mixed spheroid fused with the edge of the hepatic spheroid to form a bridge. **(H)** Cupping or partially fused HEP-MES assembloid. The packing density of the spheroid is increased and has separate layers. **(I)** HEP tissue (H) forms cup structure around MES tissue cultivated in the CG droplet system before fusing. By day 4, we see assembloid formation occur in a cup-like mechanism. **(J)** HEP-MES cupping prior to fusion in Matrigel. The left images are phase contrast, and the right-side images are fluorescent. Top- day two of formation. Bottom- day four of partial fusion is noticeable. **(K)** HEP-MES assembloids with multiple hepatic spheroids combine through the cupping mechanism. Day 5 hepatic spheroids form a cup around mesodermal tissue. Adopted with permission of publisher from Ogechi, Parashurama et al., 2021 (Ogechi et al., 2021).

Figure 11. Methods for modulating collective migration from spheroids. (A) Branching and linear migration in hepatic spheroids in the presence of M-CM. **(B)** Phase-contrast images of day 7 and day 11 hepatic spheroids in the Matrigel (MG) droplet system cultivated in varying dilutions of M-CM. By day seven, the hepatic spheroid demonstrated 3D collective cell migration. The area was measured for varying dilution ratios. Analysis of Matrigel droplet system with M-CM. A plot of fold change in the area across different M-CM dilutions (1:1, 1:7, M-CM 1 day, and M-CM). Comparison of M-CM with 1:1 condition (P = 0.15, n = 3 for both conditions), M-CM with 1:7 condition (P = 0.0056, n = 3 for both conditions), and M-CM with M-CM 1-day-only condition (P = 0.019, n = 3 for both conditions). **(C)** Phase-contrast images on day 7 of Hepatic spheroids in MG droplet system cultivated in M-CM alone, M-CM with A83-01 (10 nM), and A83-01 (20 nM). Bar graph analysis comparing day 7 M-CM and M-CM + A83-01 (20 nM) (P = 0.047, n = 3). Plotted means ± SD. Significance is defined as P ≤ 0.05. **(D)** Phase-contrast images on days 4 and 7 of the hepatic spheroid in the collagen (CG) droplet system cultivated in M-CM medium. Arrows specify thin filopodia-like extensions into the collagen. Bar graph analysis of Hepatic spheroids in collagen (CG) in cDMEM (control) and M-CM conditions comparing protrusion length (P = 0.012, n = 3). **(E)** Fluorescent images of day 5 HEP-MES mixed spheroid in Fibrin gels cultivated in M-CM. **(F)** Star-shaped migration of HEP-MES mixed spheroids. **(G)** Fluorescent images of day 4 of HEPG2-GFP (HepG2 cells expressing green fluorescent protein) and MES mixed spheroids in the MG droplet system after treatment with TGFβ1 (20 ng/ml). From left to right: HepG2 (green) cells and combined HepG2 (red) and MRC-5 (yellow) images. Replicates 2 (above) and 1 (below) are shown. Arrows show HepG2 and MRC-5 migration. Bar graph comparing the area of fibroblast migration in the negative control

687 (HepG2-GFP/MRC-5) and (HepG2-GFP/MRC-5 + TGFβ1, 20 ng/ml), P = 0.012, n = 3.
688 Plotted means ± SD. Significance is defined as P ≤ 0.05. ∗ is used to denote the
689 significance of experimental data. Adopted with permission of publisher from Ogechi,
690 Parashurama et al., 2021 (Ogechi et al., 2021).

691
Table 1. The effect of distance on assembloid formation. Hepatic spheroids and
mesodermal-derived spheroids were transferred to an agarose-coated well. The cells
were dye-labeled prior to spheroid formation to demonstrate the interaction between
spheroids. The spheroids were suspended in a 1:5 solution of the matrix (Matrigel or
Collagen) and cDMEM or fibroblast-conditioned media (M-CM). The initial distance
between the spheroids was measured using ImageJ and the corresponding compaction
time of the assembloid formation was observed. The compaction time of assembloid
formation is directly proportional to the initial distance between the spheroids, irrespective
of the matrix or media.

701
DISCUSSION:
In this protocol, several methods are presented for cultivating simple and complex
assembloids, and methods for inducing 3D collective cell migration in early liver
organogenesis. We have presented several protocols many of which have critical steps.
Spheroid formation is a critical step in the process in all these methods. Spheroid
formation can be accomplished using microwells (96- or 384-wells) with either non-
adherent or agarose-coated plates. Considerable expertise is needed to handle
organoids regarding formation of spheroids (or organoids), transferring between wells,
addition of biomatrices upon spheroids, and addition of multiple spheroids per well.
Formation of the spheroids requires critical attention to repeatable cell counting and
seeding, agarose coating (or non-adherent), regular medium changes, dye-labeling
methods, and gentle handling of spheroids and plates together with microscopy. We also
note that dye-labeling should be experimentally determined in terms of cell number and
type, and amount of labeling time.

716
For example, preparation of agarose solution must be stringent in terms of maintaining
appropriate concentrations and seeding wells with appropriate volumes, such that a
meniscus (curvature develops) and enables collection of cells in the center. We
recommend careful attention to agarose concentration, volumes, and solidification of gel.
Determining correct micropipette tip size and transferring techniques is critical for proper
handling of spheroids. Glass wells can be used for improved visualization. As we have
noted in the paper, spheroids can first appear as translucent discs that become more
opaque over time, which can be used to monitor spheroid formation, and several factors
affect this time. Finally, assembloid formation requires careful transfer of spheroids, and
they need to be placed within a couple of spheroid diameters or less to effectively observe
changes.

728
729 Here, using cell lines, a toolbox of methods to develop complex assembloids and
730 migrating spheroids have been created. It is important to note that cell lines were
731 employed. While this enabled focus upon methods, using primary cells or hPSC-derived
732 cells would be advantageous with the use of human personalized tissues. We can

validate that our assembloid techniques work with hPSC-derived HEP cells. A significant limitation of these systems is the matrix. Matrigel, a mouse tumor extracellular matrix protein mixture, is primarily used as the matrix in these cultivation models. However, it is a major limitation due to its tumor-derived origin and high cost. Furthermore, the droplet formation system can only last 2 weeks before significantly degrade, and thus alternative gels could be implemented such as collagen mixtures or sodium alginate. Furthermore, the current systems can only grow to on the order of dimensions of the spheroids, but not beyond. Thus, determining and addressing limitations to growth is critical.

There is a lack of methods to specifically study early events in early liver organogenesis, and more generally, methods to generate synthetic tissues. The methods developed here address this gap. With further study and characterization, he assembloids and tissues we generate can be used to build larger tissues that can be used for improved modeling of solid tissues and organs. This is not currently possible using current techniques.

These methods provide a toolbox that can be used to further build assembloids and assemble larger, more complex tissues for better *ex vivo* organ modeling, and potentially for *in vivo* therapeutic approaches, in addition to modeling structures during organogenesis (liver bud) and being employed for biopharma applications.

The protocols we develop here enable models of different modes of liver cell migration such as, co-migration, interstitial migration, and branching morphogenesis, as well as assembloids that, together, can be used to build more complex tissues for various biomedical and biopharma applications.

ACKNOWLEDGMENTS:
NP was supported by the UB CBE startup funds, a Mark Diamond Fellowship, the New York State Stem Cell Science C024316, the UB the Stem cells in regenerative medicine (ScIRM) center, the University at Buffalo Center for Cell, Gene, Tissue Engineering (CGTE).

DISCLOSURES:
NP is the founder of Khufu Therapeutics, an organoid engineering company that develops treatments for acute and chronic liver disease.

REFERENCES:
Binamé, F., P. Lassus and U. Hibner (2008). "Transforming growth factor beta controls the directional migration of hepatocyte cohorts by modulating their adhesion to fibronectin." Mol Biol Cell **19**(3): 945-956.
Bort, R., M. Signore, K. Tremblay, J. P. Martinez Barbera and K. S. Zaret (2006). "Hex homeobox gene controls the transition of the endoderm to a pseudostratified, cell emergent epithelium for liver bud development." Dev Biol **290**(1): 44-56.
Chen, J. S., L. L. Liang, H. X. Xu, F. Chen, S. L. Shen, W. Chen, L. Z. Chen, Q. Su, L. J. Zhang, J. Bi, W. T. Zeng, W. Li, N. Ma and X. H. Huang (2017). "miR-338-3p inhibits epithelial-mesenchymal transition and metastasis in hepatocellular carcinoma cells." Oncotarget **8**(42): 71418-71429.

Fitamant, J., F. Kottakis, S. Benhamouche, H. S. Tian, N. Chuvin, C. A. Parachoniak, J. M. Nagle, R. M. Perera, M. Lapouge, V. Deshpande, A. X. Zhu, A. Lai, B. Min, Y. Hoshida, J. Avruch, D. Sia, G. Campreciós, A. I. McClatchey, J. M. Llovet, D. Morrissey, L. Raj and N. Bardeesy (2015). "YAP Inhibition Restores Hepatocyte Differentiation in Advanced HCC, Leading to Tumor Regression." Cell Rep **10**(10): 1692-1707.

Fransvea, E., U. Angelotti, S. Antonaci and G. Giannelli (2008). "Blocking transforming growth factor-beta up-regulates E-cadherin and reduces migration and invasion of hepatocellular carcinoma cells." Hepatology **47**(5): 1557-1566.

Gualdi, R., P. Bossard, M. Zheng, Y. Hamada, J. R. Coleman and K. S. Zaret (1996). "Hepatic specification of the gut endoderm in vitro: cell signaling and transcriptional control." Genes Dev **10**(13): 1670-1682.

Gupta, S., P. Rajvanshi, R. Sokhi, S. Slehria, A. Yam, A. Kerr and P. M. Novikoff (1999). "Entry and integration of transplanted hepatocytes in rat liver plates occur by disruption of hepatic sinusoidal endothelium." Hepatology **29**(2): 509-519.

Koenig, S., C. Stoesser, P. Krause, H. Becker and P. M. Markus (2005). "Liver Repopulation after Hepatocellular Transplantation: Integration and Interaction of Transplanted Hepatocytes in the Host." Cell Transplantation **14**(1): 31-40.

Matchett KP, W.-K., Portman jr, Kapourani CA, Feroq F, May S, Mackey JBG, Brice M, Zajdel E, Beltran M, Sutherland EF, Wilson GC, Wallace SJ, Kitto L, Youngeer NT, Dobie R, Oniscu GC, Wigmore SJ, Ramchandran P, Vallejos CA, Carragher NO, Simpson KJ, Kendall TJ, Acute Liver Failure Study Group; Rule JA, Lee WM, Hoare M, Weston CJ, Marioni JC, Teichmann ST, Bird TG, Carlin LM, Henderson NC. (2023). "Multimodal decoding of human liver regeneration." bioRxiv Pre-print Server.

Matsumoto, K., H. Yoshitomi, J. Rossant and K. S. Zaret (2001). "Liver organogenesis promoted by endothelial cells prior to vascular function." Science **294**(5542): 559-563.

Ng, L., R. Tung-Ping Poon, S. Yau, A. Chow, C. Lam, H. S. Li, T. Chung-Cheung Yau, W. L. Law and R. Pang (2013). "Suppression of actopaxin impairs hepatocellular carcinoma metastasis through modulation of cell migration and invasion." Hepatology **58**(2): 667-679.

Ogoke, O., J. Oluwole and N. Parashurama (2017). "Bioengineering considerations in liver regenerative medicine." J Biol Eng **11**: 46.

Ogoke O., G. D., Thompson S., Chiang A., Ngyuen TH., Berke D., Ott C., Kalinousky A., Shamul C., Chen P., Ross S., Chen Z., Srivastava P., Mahajan S., Zhao R., Gunawan R., Parashurama N. (2022). "Dynamic changes in the niche and transcription trigger early murine and human pluripotent stem cell-derived liver organogenesis." BioRxiv (Pre-print server) **10.1101/2022.07.24.501313**.

Petkov, P. M., K. Kim, J. Sandhu, D. A. Shafritz and M. D. Dabeva (2000). "Identification of Differentially Expressed Genes in Epithelial Stem/Progenitor Cells of Fetal Rat Liver." Genomics **68**(2): 197-209.

Rajvanshi, P., A. Kerr, K. K. Bhargava, R. D. Burk and S. Gupta (1996). "Studies of liver repopulation using the dipeptidyl peptidase IV-deficient rat and other rodent recipients: cell size and structure relationships regulate capacity for increased transplanted hepatocyte mass in the liver lobule." Hepatology **23**(3): 482-496.

Rossi, J. M., N. R. Dunn, B. L. Hogan and K. S. Zaret (2001). "Distinct mesodermal signals, including BMPs from the septum transversum mesenchyme, are required in combination for hepatogenesis from the endoderm." Genes Dev **15**(15): 1998-2009.

Sosa-Pineda, B., J. T. Wigle and G. Oliver (2000). "Hepatocyte migration during liver
development requires Prox1." Nat Genet **25**(3): 254-255.
Suzuki, A., S. Sekiya, D. Büscher, J. C. Izpisúa Belmonte and H. Taniguchi (2008). "Tbx3
controls the fate of hepatic progenitor cells in liver development by suppressing p19ARF
expression." Development **135**(9): 1589-1595.
Xue, T. C., N. L. Ge, L. Zhang, J. F. Cui, R. X. Chen, Y. You, S. L. Ye and Z. G. Ren (2014).
"Goosecoid promotes the metastasis of hepatocellular carcinoma by modulating the
epithelial-mesenchymal transition." PLoS One **9**(10): e109695.
Yang, C., Y. Xu, F. Cheng, Y. Hu, S. Yang, J. Rao and X. Wang (2017). "miR-1301 inhibits
hepatocellular carcinoma cell migration, invasion, and angiogenesis by decreasing
Wnt/beta-catenin signaling through targeting BCL9." Cell Death Dis **8**(8): e2999.
Yang, M. H., C. L. Chen, G. Y. Chau, S. H. Chiou, C. W. Su, T. Y. Chou, W. L. Peng and
J. C. Wu (2009). "Comprehensive analysis of the independent effect of twist and snail in
promoting metastasis of hepatocellular carcinoma." Hepatology **50**(5): 1464-1474.
Zeng, Y. B., X. H. Liang, G. X. Zhang, N. Jiang, T. Zhang, J. Y. Huang, L. Zhang and X.
C. Zeng (2016). "miRNA-135a promotes hepatocellular carcinoma cell migration and
invasion by targeting forkhead box O1." Cancer Cell Int **16**: 63.
Zhao, Y., W. Jian, W. Gao, Y.-X. Zheng, Y.-K. Wang, Z.-Q. Zhou, H. Zhang and C.-J. Wang
(2013). "RNAi silencing of c-Myc inhibits cell migration, invasion, and proliferation in
HepG2 human hepatocellular carcinoma cell line: c-Myc silencing in hepatocellular
carcinoma cell." Cancer Cell International **13**(1): 23.
846

A Heat 1% agarose solution in microwave and allow to cool prior to use

Transfer 55-65 μL of agarose solution per well

Cool the plate for 20-30 mins at 4°C and bring to room temperature

B Cell culture has reached 80% confluency

Passage cells using Tryspin-EDTA

Suspend cells at a density of 1 x 10⁶ cells/mL of serum-free medium and add 5 uL of cell-labeling solution

C Add 5 uL of cell-labeling solution per mL and incubate for 20 mins

Centrifuge at 450 x g for 5 mins and wash cells with cDMEM. Repeat this wash process two more times.

D Add 100 μl of cell suspension to per agarose-coated well and centrifuge at 340 x g for 10 mins

agarose

mesodermal-derived spheroid

Incubate for 1-3 days until spheroid formation is complete (mesodermal-derived spheroid)

agarose

hepatic spheroid

Incubate for 5-9 days until spheroid formation is complete (hepatic spheroid)

agarose

A

Passage HepG2-WT cell culture. Suspend cells at a density of 1 x 10^6 cells/mL

Dye-label the cells and suspend at a density of 5 x 10^4 cells/mL

Add 100 μl of cell suspension to per well and centrifuge 340 *x g* for 10 mins

Incubate for 5-9 days until spheroid formation is complete

B

Passage HFF/MRC-5 cell culture. Suspend cells at a density of 1 x 10^6 cells/mL

Dye-label the cells and suspend at a density of 1 x 10^5 cells/mL

Add 100 μl of cell suspension to per well and centrifuge 340 *x g* for 10 mins

Incubate for 1-3 days until spheroid formation is complete

C

Transfer a single spheroid to a pre-existing well

Add matrix (MG or CG) at a 1:1-1:5 dilution and incubate for 3 hours

Add 75 μL of cDMEM and incubate for 2-3 days for assembloid formation

A

Passage HepG2-WT cell culture. Suspend cells at a density of 1 x 10⁶ cells/mL

Dye-label the cells and suspend at a density of 5 x 10² cells/mL

Add 100 µl of cell suspension to per well and centrifuge 340 x g for 10 mins

Incubate for 5-9 days until spheroid formation is complete

B

Collect the HepG2-WT spheroids and gently rinse with cDMEM

Suspend the spheroids in matrix (MG or CG)

C

Seed one spheroid in 15 uL of matrix (MG or CG) onto 60 mm petri dish and incubate at 37°C

Slowly add 5 mL of growth medium and incubate at 37°C

A Seed 5,00 cells/cm² in 15 mL of cDMEM (T-75 flask) → 72 hour incubation → Collect conditioned media and centrifuge at 290 x g for 5 mins

B Filter conditioned media using a 0.2 µm syringe filter → Dilute conditioned media with cDMEM (1:1-1:7)

A

Passage HepG2-WT cell culture. Suspend cells at a density of 1 x 10^6 cells/mL

Dye-label the cells and re-suspend in cDMEM

B

Passage HFF/MRC-5 cell culture. Suspend cells at a density of 1 x 10^6 cells/mL

Dye-label the cells and re-suspend in cDMEM

C

Suspend HepG2-WT and HFF/MRC-5 cells at 1:1 ratio at a density of 2.0 x 10^5 cells/mL

Add 100 μl of cell suspension to per well and centrifuge 340 x g for 10 mins

agarose

Incubate for 1-2 days for mixed spheroid formation

agarose

HepG2-WT

| | Day 2 | Day 2 (Dye labeled) | Day 4 |

A. Building Large or Small Arms

B. HEP-MES Assembloids demonstrate branching arms

Day 3 Liver/MRC5
Day 4 Liver/MRC5
Day 9 Liver/MRC5
Day 12 Liver/MRC5
Day 12 Liver /MRC5

Day 0
1000 HepG2 cells GFP/Fluc or WT 384 well RB plate

Day 3

Day 4-6
GFR Matrigel 20,000 MRC-5

C. Mixed spheroid form bridges with HEP spheroids

HepG2-WT

Mixed Spheroid

Day 0

HepG2-WT

Bridge

Mixed Spheroid

Day 5

D. Spot-Welding (fused edges)

E. HEP-MES in collagen with MES-conditioned medium fuse edges and don't form cupping

Day 0   M-CM
M   H
CG

Day 1
M   H
CG

Day 4   M-CM
M   H
CG

F. In Collagen 2 mg/ml (stiff), HEP-MES edges fuse

Collagen 2 mg/ml   HEPG2-WT
MRC5
Day 3

Collagen 2 mg/ml
H
M
Day 3

G. Infiltrating and Layering

H. HEP-MES infiltrate in Collagen, with MES- conditioned medium

Collagen 2 mg/ml MES-condition medium MCM
Day 3

Collagen 2 mg/ml MCM
M
H
Day 3

I. HEP-inside, MES-outside leads to layered assembloid

Day 6

Replicate 1
Day 6
HepG2/MSC

Replicate 2
Day 6
HepG2/MSC

A

Fused HEP-MES assembloids (increased packing density, separate layers)

B

HEP and MES spheroids fuse in matrigel

Day 3 — L MRC5 200 µm
Day 5 — L MRC5 200 µm
Day 7 — L MRC5 200 µm
Day 9 — L MRC5

C

Day 13 MRC5 L 200 µm MRC5 L

D

HEP-MES assembloids with multiple MES spheroids fuse but overall volume does not increase

Day 0 H M M M
Day 1 H M M M
Day 5 M

E

HEP-MES assembloids form under low serum

2% FBS M H Day 0
2% FBS M H Day 1
2% FBS M H Day 2
0.2% FBS H M Day 0
0.2%FBS H Day 1
0.2%FBS Day 2

F

HEP-MES don't form at large distances in collagen with MES-conditioned medium

Day 1 M-CM M H CG
Day 5 M-CM M H CG

G

Mixed spheroids fuse with MES spheroids

Day 0 Mixed spheroid HFF
Day 5 HFF Mixed spheroid

H

Cupping, partially fused HEP-MES assembloids (increased packing density, separate layers)

I

In collagen, HEP-MES form assembloids via cup-like mechanism

Day 0 M
Day 1 M
Day 2 M H
Day 4 M H CG

J

HEP-MES cupping prior to fusion in MG

Matrigel H M Day 2
Matrigel H M Day 2
Matrigel H M Day 4
Day 4

K

HEP-MES assembloids with multiple HEP spheroids combine with cupping mechanism

Day 0 H H H M
Day 1 H M
Day 5 H H M H

A

Branching

Linear

B

M-CM | Day 7 | M-CM | Day 11

P = NS

*

*

Fold-increase in area

1:1 | 1:7 | M-CM 1 day | M-CM

C

Branching morphogenesis of HEP-spheroids inhibited by TGF-beta

M-CM | M-CM + 10 nM A83-01 | M-CM + 20 nM A83-01

Day 7 | Day 7 | Day 7

300 um

*

Max Branch Length/Mean Branch Length

D7 MCM | D7 (10 nM) | D7 (20 nM)

D

HEP spheroids demonstrate linear branching in collagen gels

M-CM | Day 4 | M-CM | Day 7

500 µm | CG | CG

*

Protrusion length (µm)

Cntrl | M-CM

E

HEP-MES mixed spheroids demonstrate hair-like migration in Fibrin gels

Day 4 | HepG2/HFF | Day 5 | HFF | HepG2 | HFF HepG2

500 um | Fib | Fib | Fib | Fib

F

Star-shaped

G

Increasing branch length in HEP-MES mixed spheroids with TGF-Beta

Tgfb + MRC5/ HepG2 in Matrigel

Replicate 2 | Day 4 | HepG2 | HepG2 MRC5

Replicate 1 | Day 4 | HepG2 | HepG2 MRC5

*

Area of Fibroblast Migration (µm²)

Control | TGFB (+)

Condition Media & Matrix	Initial Distances (μm)	Days to Merge
Matrigel + DMEM	492.742 – 974.051	6-12
Collagen + DMEM	413.983 – 593.872	5-8
Collagen + MCM	695.235	9