1 **TITLE**
2 *In vitro* cultivation techniques for modeling liver organogenesis, building assembloids, and
3 designing synthetic tissues
4

5 **AUTHORS AND AFFILIATIONS:**
6 Simran Kumar[1], Helly Patel[1], Natesh Parashurama[1,2,3,4,5]
7
[1] Department of Biomedical Engineering, University at Buffalo (State University of New York),
Furnas Hall, Buffalo, NY 14260
[2] Clinical and Translation Research Center (CTRC), University at Buffalo (State University of New
York), 875 Ellicott St., Buffalo, NY 14203
[3] Department of Chemical and Biological Engineering, University at Buffalo (State University of
New York), Furnas Hall, Buffalo, NY 14260
[4] Center for Cell, Gene and Tissue Engineering, University at Buffalo, State University of New
York, Furnas Hall Buffalo NY 14260
[5] Khufu Therapeutics, Inc. Buffalo, NY, 14260
17
18 **Corresponding Author:**
19 Natesh Parashurama, 906 Furnas Hall, Buffalo, NY 14260; Tel: 716-645-1201; Fax: 716-645-
20 3822; e-mail: nateshp@buffalo.edu
21
22 **Co-Authors:**
23 Helly Patel (hellypatel1999@gmail.com)
24 Simran Kumar (skumar27@buffalo.edu)
25

26 **KEYWORDS:**
27 organogenesis, liver development, collective cell migration, epithelial to mesenchymal
28 transition, liver diverticulum, liver bud
29
**ABBREVIATION LIST:**
**BMP4** Bone morphogenetic protein 4
**FGF-2** Fibroblast growth factor 2
**PROX1** Prospero homeobox 1
**HBs** hepatoblasts
**HEP** hepatic
**HGF** hepatocyte nuclear factor 4 alpha
**hMSC** human mesenchymal stem cells
**HPSC** human pluripotent stem cells
**MES** mesenchymal
**MES-CM** fibroblast-conditioned medium
**TBX3** T-box transcription box factor 3
**TGFβ1** transforming growth factor beta1
43
**SUMMARY:**

Organoids revolutionize personalized tissue modeling for organ development, drug discovery,
and disease research. Organoid engineering advances into creating intricate synthetic tissues.
The aim is to integrate morphogenesis, assembloid technology, and biomatrices to advance
tissue engineering. The presented methods aid in modeling liver organogenesis and establishing
guidelines for synthetic tissue construction.
50

**ABSTRACT:**
Chronic liver disease has reached epidemic proportions, affecting over 800 million people
globally. The current treatment, orthotopic liver transplantation, has several limitations.
Promising solutions have emerged in the field of liver regenerative medicine, with liver
organogenesis holding significant potential. Early liver organogenesis, occurring between E8.5
and 11.5, involves the formation of epithelial-mesenchymal interactions leading to
morphogenesis, hepatic cord formation, and collective migration. However, there is a lack of
methods for *in vitro* modeling of this process. In this study, a detailed series of methods are
presented enabling the modeling of various stages and aspects of liver organogenesis. In one
method series, assembloid technology with hepatic and mesenchymal spheroids is utilized,
replicating early structures found in liver organogenesis, modeling early morphogenesis, and
demonstrating interstitial cell migration as seen *in vivo*. These innovative assembloid systems
help identify factors influencing assembloid formation and migration. Hepatic spheroid
cultivation systems were also employed to model collective migration and branching
morphogenesis. Fibroblast-conditioned media (MES-CM) plays a significant role in initiating
dose-dependent branching migration. Future work will involve high temporal and spatial
resolution imaging of hepatic and mesenchymal interactions to determine the cascade of
cellular and molecular events involved in tissue formation, morphogenesis, and migration.

**INTRODUCTION:**
Liver cell migration plays a significant role in liver organogenesis, disease, and cell therapy.
During liver organogenesis (E8.5-9.0, mouse), the ventral foregut pre-hepatic epithelium begins
to express liver genes due to inductive signals emanating from the surrounding mesenchyme
and heart. At E9.0, the foregut epithelium thickens as the cells transition from cuboidal to
pseudostratified columnar morphology, forming the liver diverticulum [3, 7]. At this critical stage,
the liver diverticulum is comprised of only ~1,500 cells. Next, the hepatic endoderm lining the
liver diverticulum thickens, delaminates, and forms cords of migrating hepatoblasts that co-
migrate with endothelial cells and mesenchymal cells, branching into the surrounding
mesenchymal tissue, initiating three-dimensional collective cell migration to form the liver bud
[15, 16]. In fact, during this stage, cells collectively undergo: 1) co-migration, or movement
together with other cell types, 2) branching morphogenesis, or formation of branching tube-like
structures, and 3) interstitial migration, or migration on top of other cells. By E11.5, migration
ceases, and the primitive liver has formed, expanding $10^3$-fold [15]. Liver cell migration may also
be required in later stages of liver organogenesis, as rat fetal hepatoblasts (HBs) expression has
shown evidence of highly upregulated genes associated with 3D collective cell migration,
morphogenesis, and extracellular matrix remodeling [18]. In addition to its role in early liver
organogenesis, 3D collective migration is intricately linked to the local spread and metastasis of
advanced hepatocellular carcinoma (HCC), ultimately leading to worsened prognosis and

increased treatment resistance[25]. Adult and fetal hepatocytes also employ collective migration
when moving from the spleen to within the liver during liver repopulation. *In vivo* imaging
studies have demonstrated that transplanted hepatocytes enter the portal vein and then the
capillaries within hours, migrate across the liver sinusoids, and through the liver tissue[8, 10, 19].
Finally, recent studies demonstrate that migrating hepatoblasts arise during murine and human
liver regeneration with some evidence of movement in sheets[11]. Overall, liver collective
migration, capable of multiple modes of morphogenesis, plays a significant role in
organogenesis, cancer, hepatocyte cell therapy, and liver regeneration.
97
Numerous genetic studies have investigated the molecular pathways that drive 3D liver
collective cell migration. These studies demonstrate that ablation of hepatic cords blocks liver
formation and shows that, therefore, the formation of hepatic cords and their ensuing
interactions with supporting cells  are required for liver formation[3, 12, 21, 22]. These studies also
demonstrate that liver growth is initiated by fibroblast growth factor 2 (FGF2) secreted from
the cardiac mesoderm, bone morphogenetic protein 4 (BMP4) secreted from the surrounding
mesenchyme, hepatocyte growth factor (HGF), endothelial cell interactions, and migration-
associated transcription factors including (HEX), PROX1, and TBX3[7, 20]. Overall, genetic studies
support the fact that soluble factor signaling leading to transcription factor expression is
responsible for driving migration, signaling, and molecular interactions between hepatoblasts
and their surrounding mesenchyme.
109
Although cell migration in early liver organogenesis has been extensively investigated, current
*in vitro* hepatic migration studies frequently utilize 2D assays consisting of highly migratory HCC
cells combined with *in vivo* tumor models[13]. These studies have provided insight into several
factors that play a role in hepatic migration including TGFβ1[6], c-Myc[27], Yes-associate protein
(YAP)[5], goosecoid[23], actopaxin[2], and miRNAs[4, 24, 26]. Despite the advancements in understanding
the molecular mechanisms in 3D hepatic cell migration, the fundamental mechanisms between
2D and 3D cellular migration are distinct, suggesting 2D assays have their limitations.
Furthermore, these models typically do not implement mesenchymal cell types, which are
essential to migration/growth. There has been progress in the development of 3D models for
liver migration that incorporate the supporting mesenchyme; however, they are solely focused
on co-migration rather than the different modes of collective migration.
121
The ability to form tissues from spheroids through various self-assembly and morphogenetic
processes enables the scientific study of synthetic tissues for applications in drug development
and screening, disease modeling, therapy, and other biomedical and biotechnological
applications. Methodological details for several 3D *in vitro* cultivation systems engineered to
exhibit different modes of liver 3D collective migration are presented. These systems include
the following: (1) co-spheroid culture with hepatic (HEP) and mesenchymal-derived (MES)
spheroids in matrix, (2) spheroid matrix droplet cultured with fibroblast-conditioned medium
(MES-CM), and (3) mixed spheroids (HEP and MES cells). These systems enable robust modeling
of liver 3D collective migration, improving our molecular and cellular understanding of liver
organogenesis, cancer, and therapy.
132

**PROTOCOL:**

**1. Preparation of 1% Low EEO Agarose Solution**

1.1. Weigh 2.5 g of low EEO agarose powder and transfer it to a beaker.

NOTE: Ensure the beaker is at least twice the size of the desired volume to accommodate bubbling during the solution preparation.

1.2. Use a graduated cylinder to measure 250 mL of distilled (DI) water and pour it into the beaker, aiming for a final concentration (w/v) of 1%.

1.3. Cover the beaker's opening with plastic wrap, leaving a small hole for ventilation. Heat the beaker in the microwave for 30 seconds.

1.4. Carefully remove the beaker and swirl the contents. Repeat this process every 30 seconds until complete dissolution.

CAUTION: Handle microwaved glassware carefully, using proper gloves. Monitor the solution closely to prevent overheating or boiling over.

1.5. Once a homogeneous solution is achieved, remove the beaker from the microwave. Gently swirl the solution before transferring it to a pre-sterilized bottle. Autoclave the solution and store the agarose solution at room temperature until ready to use.

**2. Coating 96-Well Plate**

NOTE: 384-well ultra-low attached spheroid microplates can be utilized in these and further experiments.

2.1. Loosen the cap of the bottle containing the sterile 1% agarose solution. Warm the solution in the microwave in 30-second intervals until it transitions to the liquid phase; then tighten the cap.

CAUTION: Handle microwaved glassware carefully, using proper gloves. Monitor the solution closely to prevent overheating or boiling over.

NOTE: Perform these steps under a sterile tissue-culture laminar flow hood.

2.2. Transfer 55-65 µL of the 1% agarose solution per well to coat the 96-well tissue-cultured plate and immediately rotate the plate.

2.3. Once the 1% agarose solution has been transferred to the desired number of wells, place
the plates in a 4°C fridge for 20-30 minutes to allow the agarose to solidify. Before use,
bring the plate to room temperature (**Figure 1A**).

**3.  Preparation of HepG2 (HEP) Spheroids**

3.1. Cultivate HepG2 cells in a T-75 flask using completed growth medium (cDMEM) comprising
of Dulbecco's Modified Eagle Medium (DMEM), supplemented with 10% Fetal Bovine
Serum (FBS) and 1% Penicillin-Streptomycin (Pen-Strep). Maintain the cell culture at 37°C
and 5% $CO_2$, changing the medium daily.

3.2. Upon reaching 80% confluency, perform a rinse of the cells with 1X Phosphate Buffered
Saline (PBS) and discard the rinse. Add 5 mL of 0.05% of Trypsin-EDTA to the flask and
incubate at 37°C and 5% $CO_2$ for 5-10 minutes.

3.3. Add equal amounts of cDMEM and wash the cells off the flask. Once cells detach, transfer
the mixture to a 15 mL sterile conical centrifuge tube and centrifuge the cell suspension at
300 $x$ $g$ for 5 minutes.

NOTE: All centrifugation steps outlined in this protocol were performed at room temperature.
However, these centrifugation procedures can alternatively be carried out at cold
temperatures.

3.4. Re-suspend the cell suspension in fresh cDMEM and count the cells to determine the final
concentration in cells/mL (**Figure 1B**).

3.5. Dye-labeling of cells

NOTE: This is an optional step.

3.5.1. Transfer the desired amount of cell suspension to a 15 mL sterile conical centrifuge tube
and centrifuge the cell suspension at 300 $x$ $g$ for 5 minutes.

3.5.2. Re-suspend the cell pellet in a serum-free growth medium to achieve a final
concentration of 1.0 x $10^6$ cells/mL. Add 5 µL of Vybrant Cell-Labeling Solution per mL of
cell suspension and incubate the cell suspension at 37°C and 5% $CO_2$ on rotation for 20
minutes.

NOTE: Serum-free growth medium is DMEM only supplemented with 1% Pen-strep. Adjust the
incubation time for uniform staining based on the cell suspension density.

3.5.3. Once the incubation is completed, centrifuge the cell suspension at 450 $x$ $g$ for 5 minutes
and re-suspend the cell pellet in fresh cDMEM. Repeat this wash process two more times
(**Figure 1C**).

3.6. Spheroid formation

3.6.1. Suspend the cells in fresh cDMEM to achieve a final concentration of 5.0 x $10^4$ cells/mL
based on the cell count. Mix the cell suspension well and transfer 100 μL of cell
suspension per well to the agarose-coated 96-well plate.

NOTE: The cell suspension density is set to achieve 5,000 cells per well in the agarose-coated
96-well plate.

3.6.2. Centrifuge the plate at 340 $x$ $g$ for 10 minutes and incubate at 37°C and 5% CO2 for 5-9
days (**Figure 1D**).

NOTE: Change medium every other day after plating, gently removing 50% of cDMEM and
replacing it.

NOTE: HepG2 spheroids are suitable for the HepG2 and HFF/MRC-5 (HEP-MES) assembloid
model or MES-CM model.

**4. Preparation of HFF/MRC-5 (MES) Spheroids**

4.1. Cultivate HFF/MRC-5 cells in a T-175 flask using completed growth medium (cDMEM)
comprising of Dulbecco's Modified Eagle Medium (DMEM), supplemented with 10% Fetal
Bovine Serum (FBS) and 1% Penicillin-Streptomycin (Pen-Strep). Maintain the cell culture
at 37°C and 5% $CO_2$, changing the medium every other day.

4.2. Upon reaching 80% confluency, perform a rinse of the cells with 1X Phosphate Buffered
Saline (PBS) and discard the rinse. Add 5 mL of 0.25% of Trypsin-EDTA to the flask and
incubate at 37°C and 5% CO2 for 5-10 minutes.

4.3. Add equal amounts of cDMEM and wash the cells off the flask. Once cells detach, transfer
the mixture to a 15 mL sterile conical centrifuge tube and centrifuge the cell suspension at
*300 x g* for 5 minutes.

4.4. Re-suspend the cell suspension in fresh cDMEM and count the cells to determine the final
concentration in cells/mL (**Figure 1B**).

4.5. Dye-labeling of Cells

NOTE: This is an optional step.

4.5.1. Transfer the desired amount of cell suspension to a 15 mL sterile conical centrifuge tube
and centrifuge the cell suspension at 300 $x$ $g$ for 5 minutes.

4.5.2. Re-suspend the cell pellet in a serum-free growth medium to achieve a final
265            concentration of $1.0 \times 10^6$ cells/mL. Add 5 µL of Vybrant Cell-Labeling Solution per mL of
266            cell suspension and incubate the cell suspension at 37°C and 5% CO2 on rotation for 20
267            minutes.
268
269 NOTE: Serum-free growth medium is DMEM only supplemented with 1% Pen-strep. Adjust the
270 incubation time for uniform staining based on the cell suspension density.
271
272     4.5.3. Once the incubation is completed, centrifuge the cell suspension at 450 $x$ $g$ for 5 minutes
273            and re-suspend the cell pellet in fresh cDMEM. Repeat this wash process two more times
274            (**Figure 1C**).
275
276     4.6.  Spheroid Formation
277
278     4.6.1. Suspend the cells in fresh cDMEM to achieve a final concentration of $1.0 \times 10^5$ cells/mL
279            based on the cell count. Mix the cell suspension well and transfer 100 µL of cell
280            suspension per well to the agarose-coated 96-well plate.
281
282 NOTE: The cell suspension density is set to achieve a density of 10,000 cells per well in the
283 agarose-coated 96-well plate.
284
285     4.6.2. Centrifuge the plate at 340 $x$ $g$ for 10 minutes and incubate at 37°C and 5% CO2 for 5-9
286            days (**Figure 1D**).
287
288 NOTE: Change medium every other day after plating, gently removing 50% of cDMEM and
289 replacing it.
290
291 NOTE: HFF/MRC-5 spheroids are suitable for the HEP-MES assembloid model.
292
293 **5.  HepG2 and HFF/MRC-5 (HEP-MES) Assembloid Formation**
294
295 NOTE: Refer to Sections 3 and 4 for formation of HepG2 (**Figure 2A)** and HFF/MRC-5 spheroids
296 (**Figure 2B**).
297
298 5.1.  Collect HFF/MRC-5 spheroids individually using a pipette from the 96-well plate. Transfer
299         them to a 15 mL sterile conical centrifuge tube containing fresh cDMEM. Allow the
300         spheroids to settle in the tube for 5 minutes and then gently rinse with warm cDMEM.
301
302 NOTE: Perform the rinsing process gently and with care.
303
304 5.2.  Gently transfer a single HFF/MRC-5 spheroid using a pipette to a well containing a HepG2
305         spheroid. Add ice-cold growth factor reduced (GFR) Matrigel (MG)/ Collagen (CG) in a 1:1
306         and 1:5 dilution and incubate at 37°C at 5% $CO_2$ for 3 hours (**Figure 2C**).
307

NOTE: Thaw MG overnight on ice in a 2°C to 8°C fridge. Ensure the use of pre-cooled pipettes,
tips, and tubes when handling MG, as it forms a gel above 10℃.

5.3.  After incubation, add 75 µL of fresh cDMEM to each well and incubate at 37°C at 5% $CO_2$
for 2-3 days.

NOTE: Change medium every other day after plating, gently removing 50% of cDMEM and
replacing it. Assembloids will still form without media changes for up to 3-4 days.

NOTE: Assembloid formation can occur without the use of matrix.

**6.  HEP Spheroid Droplet Formation**

NOTE: Refer to Section 3 for formation of HepG2 spheroids (**Figure 3A**).

NOTE: Two different materials can be used for suspending the HepG2 spheroids in droplets. The
two methods are provided below.

6.1.  Matrigel (MG) droplets

6.1.1. Collect HepG2 spheroids individually using a pipette from the 96-well plate, allowing
them to settle in a 15 mL sterile conical centrifuge on ice.

6.1.2. In a separate 15 mL sterile conical centrifuge tube, mix 1 mL of ice-cold GFR MG and ice-
cold control growth medium at a 1:1 dilution. Aspirate the medium from the 15 mL tube
containing the HepG2 spheroids and place it on ice.

NOTE: Thaw MG overnight on ice in a 2°C to 8°C fridge. Ensure pre-cooled pipettes, tips, and
tubes when handling MG, as it forms a gel above 10℃.

6.1.3. Add the MG solution to HepG2 spheroids, ensuring even spheroid distribution inside the
MG solution.

6.1.4. Using a 200 µL pipette, collect a 15 µL volume of a single spheroid in the MG solution and
slowly seed it onto a 60 mm petri dish (**Figure 3B**).

NOTE: If more than one spheroid is seeded per droplet, remove, and reseed properly. Spheroids
are repositioned if needed to ensure they are at the center of the droplet.

NOTE: For optimal spheroid seeding, consider using reverse pipetting to minimize air bubble
formation.

6.2.  Collagen (CG) Droplets
351

NOTE: All CG preparation should be done on ice.

6.2.1. Collect HepG2 spheroids individually using a pipette from the 96-well plate, allowing
them to settle in a 15 mL sterile conical centrifuge tube on ice (**Figure 3B**).

6.2.2. In a microcentrifuge tube, add 358.8 µL of de-ionized water, 100 µL of 10X PBS, 12.1 µL
of 1 N NaOH, and 529.1 µL of stock rat tail CG for a total volume of 1 mL.

NOTE: Stock rat tail CG should always be added at the end.

6.2.3. Add the CG solution to HepG2 spheroids, ensuring even spheroid distribution inside the
CG solution.

6.2.4. Using a 200 µL pipette, collect a 15 µL volume of a single spheroid in the CG solution and
slowly seed it onto a 60 mm petri dish.

NOTE: If more than one spheroid is seeded per droplet, remove, and reseed properly. Spheroids
are repositioned to ensure they are at the center of the droplet.

NOTE: For optimal spheroid seeding, consider using reverse pipetting to minimize air bubble
formation.

6.3 Incubate the droplets at 37°C and 5% CO2 for 60 minutes before the addition of growth
medium.

6.4 Slowly add 5 mL of the desired growth medium to the petri dish and incubate at 37°C and
5% $CO_2$ with medium changes every three days (**Figure 3C**).

NOTE: HFF/MRC-5 conditioned-media was used in the droplet formation assay.

**7. Preparation of HFF/MRC-5 Conditioned Media (MES-CM)**

7.1. Harvest HFF/MRC-5 cells and seed them into a T-75 tissue culture-treated flask at a
seeding density of 5,000 cells/cm$^2$ and incubate the flask at 37°C and 5% CO2 for 72 hours
in 15 mL of cDMEM.

NOTE: Monitor the flask during this period to ensure optimal cell health.

7.2. After the 72-hour incubation, collect the fibroblast-conditioned growth medium in a 15 mL
sterile conical centrifuge tube. Centrifuge the fibroblast-conditioned growth medium at
290 $x g$ for 5 minutes and filter using a 0.2 µm filter (**Figure 4A**).

7.3. Dilute the fibroblast-conditioned growth medium with complete growth medium at a 1:1
to 1:7 dilution ratio and add to the desired experiment (**Figure 4B**).

NOTE: Utilize HFF/MRC-5 conditioned-media in the droplet formation assay.

**8. HepG2 and HFF/MRC-5 Mixed Spheroid Formation**

NOTE: Refer to steps 3.1 – 3.5. for HepG2 cell suspension preparation and steps 4.1. – 4.5. for
HFF/MRC-5 cell suspension preparation.

8.1. Suspend the HepG2 cells in fresh cDMEM to achieve a final concentration of 1.0 x 105
cells/mL for each cell suspension.

8.2. Suspend the HFF/MRC-5 cells in fresh cDMEM to achieve a final concentration of 1.0 x 105
cells/mL for each cell suspension.

8.3. Transfer HepG2 and HFF/MRC-5 cell suspensions to a 15 mL sterile conical centrifuge tube
at a 1:1 ratio to obtain a final concentration of $2.0 \times 10^5$ cells/mL. Thoroughly mix the cell
suspension and transfer 100 μL of cell suspension per well to the agarose-coated 96-well
plate.

NOTE: The cell suspension density is set to obtain 20,000 cells per well in the agarose-coated
96-well-plate.

8.4. Centrifuge the plate at 340 $x$ $g$ for 10 minutes and incubate at 37℃ and 5% CO2 for 1-2
days (**Figure 5C**).

8.5. Addition of Fibrin Hydrogels

NOTE: This is an optional step that is done in 72 hours after plating HEP-MES mixed spheroids.

8.5.1. In a 15 mL sterile conical centrifuge tube, mix fibrinogen (3.25 mg/mL) and thrombin
(12.5 units/mL) at a 4:1 ratio for the formation of fibrin and put the tube on ice.

8.6. Transfer the fibrin to each well with a pre-existing mixed spheroid at a 1:1 dilution with
current cDMEM in the well. Incubate the plate at 37℃ and 5% $CO_2$ for 30 minutes.

8.7. After incubation, add 75 μL of fresh cDMEM to each well and incubate at 37°C at 5% $CO_2$
for 1-2 days.

NOTE: Change medium every other day after plating, gently removing 50% of cDMEM and
replacing it.

**9. High Density HFF/MRC-5 Migratory Model**

NOTE: Refer to section 3 for HepG2 spheroid preparation (**Figure 6A**) and steps 4.1. − 4.5. for
HFF/MRC-5 cell suspension preparation (**Figure 6B**).

9.1. Suspend the HFF/MRC-5 cells in fresh cDMEM to achieve a final concentration of 6.0 x $10^5$
cells/mL and put on ice.

9.2. Mix the HFF/MRC-5 cell suspension and ice-cold MG/CG at a 1:1 dilution (**Figure 6C**).

NOTE: The cell suspension density is set to obtain 30,000 cells per well in the agarose-coated
96-well-plate.

NOTE: Thaw MG overnight on ice in a 2°C to 8°C fridge. Ensure pre-cooled pipettes, tips, and
tubes when handling MG, as it forms a gel above 10°C.

9.3. Transfer 100 µL of HFF/MRC-5 and MG solution using a pipette to a well containing a
HepG2 spheroid. Incubate the plate at 37°C and 5% $CO_2$ for 30 minutes.

9.4. After incubation, add 75 µL of fresh cDMEM to each well and incubate at 37°C at 5% $CO_2$
for up to 12 days.

**REPRESENTATIVE RESULTS:**
Currently, increased interest exists in synthetic tissues for various biomedical
applications, including disease modeling, drug discovery, and tissue engineering (**Figure 7**). In
this field, human pluripotent stem cell (hPSC)-derived organoids, spheroids, and cells can be
transformed into more synthetic, complex, and larger tissues. Achieving this involves applying
principles of morphogenesis, utilizing tools like microfabrication, and incorporating biomatrices
to engineer synthetic tissues more precisely (**Figure 7**). Several methods are presented here
with this theme in mind.

*Effects of clustering on spheroid formation*
The methods developed for this study were contingent upon successful 3D spheroid
formation. Successful spheroid formation is indicated by cells fusing within five to nine days,
forming a full spheroid. An early indication of success is cell clustering in the center of the well
post-centrifugation of the cultivation plate. While spheroid formation occurred less frequently
when cells were initially scattered, it still demonstrated that successful spheroid formation
could occur. Unsuccessful spheroid formation is characterized by cells not spreading and fusing
within nine days post-plating.
Spheroids of two different sizes were used to perform this experiment. Small spheroids
(S) were plated at a concentration of 1,500 cells per well, and large spheroids (L) were plated at
a concentration of 3,000 cells per well. HEP spheroids compacted and fully cultured by day five,
irrespective of the spheroid size. The cell density per well did not affect the rate at which the
spheroid formed but had a significant impact on spheroid size (**Figure 8A**). In this instance, it is
crucial to observe an increase in opacity, indicating thickening of the initial disc-shaped tissue,

which transitions from being more translucent to a spheroid configuration with increased
opacity. HEP spheroids are uniformly less round and increase in optical density (increasing
opacity) over time. If these HEP spheroids are still translucent, the spheroids should be either
discarded or the experiment should be re-done. The same experiment was performed for
HFF/MRC-5 spheroids, in which, the mesenchymal-derived spheroids compacted and fully
cultured within 24 hours, irrespective of spheroid size. The cell density per well did not affect
the rate at which the spheroid had formed (**Figure 8B**). Unlike hepatic spheroids, mesodermal-
derived spheroids compact within 24 hours, irrespective of cell seeding density. Due to their
contractile properties, MES spheroids are uniformly round and optically dense spheroids.
Notably, HFF/MRC-5 cells compact much tighter than HepG2 cells, likely resulting in little
impact of cell density on spheroid size (**Figure 8C**).
493
***Factors that affect hepatic and mesenchymal (mesodermal-derived) assembloid formation***
Hepatic and mesodermal-derived spheroids of varying sizes underwent cultivation to
investigate the impact of size on the compaction time of spheroid formation. Hepatic and
mesodermal-derived spheroids were co-cultured in GFR Matrigel (MG) or Collagen (CG) gel at a
1:5 dilution with complete growth medium (cDMEM) or fibroblast-conditioned medium (MES-
CM) to explore the potential influence of matrix and conditioned medium on assembloid
formation. Both cell types were dye-labeled before spheroid formation to illustrate the
interaction between spheroids. In all experiments in which spheroids were dye-labeled, they
were visualized using standard filters for green (488 nm), blue (450 nm), and red (594 nm).
Images of the cells were captured using a Zeiss Axio fluorescent microscope equipped with
Axiovision software to zero the background noise and images were analyzed using Image J.
Phase-contrast, individual, and overlay images were obtained as .zip files, which were
formatted into .tiff files using Image J.
It was observed that assembloid formation occurs regardless of matrix and medium
(**Figure 9**). The effects of CG, MG, and MES-CM were studied, and assembloid formation was
observed in all cases, although morphological details varied slightly (**Figure 9**). Details regarding
these images will be re-used in later figures. To assess the impact of inter-spheroid distance on
assembloid formation, distance measurements were taken along with the success of
assembloid formation in the same experimental system (MG and DMEM). It was noted that the
compaction time of an assembloid is directly proportional to the initial distance between
hepatic and mesodermal-derived spheroids (**Table 1**).
515
***Building more complex assembloids with arm-like structures***
Methods were developed to construct assembloids with branching cords (**Figure 10A**).
To perform this experiment, hepatic spheroids were combined with biomatrix (GFR MG) in 384-
well plate and surrounded by single fibroblasts at high density. These fibroblasts clustered,
providing guides toward which hepatic cells migrated, thickening over time and forming cords.
Hepatic spheroids were cultured in the MG droplet system containing a high density of MRC-5
cells (30,000 cells) in 100 μL of cDMEM and matrix in 384-wells, displaying small clusters of
MRC-5 cells in the MG (**Figure 10B, days 3-4**). Subsequently, liver spheroids formed thick
migrating strands protruding towards the fibroblast clusters, creating thick strands containing
both cell lines (**Figure 10B, days 9-12**). This approach resulted in longer arms or cords, likely

containing a mix of hepatic and fibroblast cells. Another approach involved constructing bridges
or small interconnections (arms) between spheroids. To build small armed-structures, a larger
HEP spheroid could be co-cultured in 384-well plate with a smaller mixed spheroid (**Figure 10C**),
leading to small, knob-like arms. In summary, two approaches for forming additional arms to
spheroids are presented.
531
***Spot-welding (fused edges) of complex assembloids***
Methods were employed to construct assembloids with fused edges. The methodology
to perform these experiments was HEP-MES assembloid formation assay, however, the MES
spheroid had a lower density than previous experiments. Hepatic, and mesodermal-derived
spheroids were positioned in a rigid CG environment (2 mg/mL) (**Figure 10D**). In this rigid
setting, instead of cupping, spheroids were observed to fuse at the edges, creating assembloids
with indications of short arms or spot-welding (**Figure 10E**). On the other hand, described
below, a cupping mechanism was also observed using the same culture system, in which the
HEP spheroid surrounds the MES spheroid to form a partially fused assembloid. Regarding the
spot-welding mechanism, similar data were observed with MRC-5 fibroblasts in CG (**Figure 10F**).
542
***Infiltrating and layering of complex assembloids***
During liver organogenesis, in the developing liver diverticulum, HEP cells are
surrounded by mesenchyme. Ultimately, they migrate or infiltrate into the mesenchyme (**Figure
10G**). In MES-CM and CG conditions, the addition of MES and HEP spheroids results in a
different type of fusion, where an infiltrative pattern is observed (**Figure 10H**). Furthermore, a
layering pattern can be achieved by placing single human mesenchymal stem cells (hMSC) in
GFR MG at high density. They migrate towards a HEP spheroid and layer on the surface without
infiltration, as shown with dye-labeling of the hMSC (**Figure 10I**). It is noteworthy that this is a
different phenotype than when HFF was employed. The latter arrangement is observed in the
liver diverticulum stage and many other endoderm-derived tissues. Two approaches are
presented here, relevant for modeling the developing liver diverticulum and creating aspects of
synthetic tissues.
555
***Fused hepatic and mesenchymal (mesodermal-derived) assembloid formation***
The fusion of two spheroids to form assembloids is crucial for modeling the liver bud.
This is because when two spheroids fuse, the cells likely migrate on top or between other cells,
a phenomenon referred to as interstitial migration. Interstitial migration occurs in the liver bud
when early migrating hepatoblasts move through the mesenchyme, potentially on top of other
cells. Therefore, the fusion of two spheroids serves as a model for interstitial migration during
liver organogenesis. Methods were developed to construct assembloids that fuse completely
with separate layers (**Figure 11A**). HEP and MES (MRC-5) spheroids in MG form a fused
assembloid by day 9 (**Figure 11B**). Dye-labeling analysis demonstrated that MES tissue
remained inside, while the HEP tissue remained outside (**Figure 11C**). Importantly, the final
spheroid is approximately the same size as the original spheroids, suggesting that the cells are
packed at high density. Notably, the same phenomena occur in the absence of MG in 384-well
plates when multiple MES spheroids are placed with a single HEP spheroid (**Figure 11D**). HEP-
MES assembloids, in the absence of matrix, also form under low serum (2% FBS) and extremely

low serum (0.2% FBS) conditions (**Figure 11E**). We then determined that at large distances
(approximately 3 diameters of the spheroid), assembloids did not form (**Figure 11F**). To
determine how spheroid composition determines spheroid fusion, it was demonstrated that
mixed spheroids (containing MES and HEP cells) can fuse with MES spheroids by day 5,
accompanied by an increased packing density, as expected (**Figure 11G**).
575
***Partially fused or cupping in complex assembloid formation***
Successful assembloid formation in MG demonstrates several phenotypes, including the
observation that the HEP spheroid undergoes "cupping" of the MES spheroid to form a partially
fused assembloid (**Figure 11H**). This also establishes a visual model of interstitial migration and
points toward the current mechanism of fused assembloid formation. Time-series studies of
HEP and MES spheroids demonstrate that in CG, fusion occurs via a cup-like mechanism (**Figure
11I**). This is further demonstrated using dye-labeling (**Figure 11J**). Finally, the same phenomena
are clearly illustrated when multiple HEP spheroids are used with a large, single MES spheroid
(**Figure 11K**). The significantly smaller HEP spheroid cannot completely encompass the larger
MES spheroid; therefore, it results in the same cupping mechanism. The data suggest that
cupping without fusion can occur in MG or in cases where the MES spheroid is much larger than
the HEP spheroids.
588
***Induced hepatic branching and linear migration via mesenchymal (mesodermal-derived)***
***conditioned growth medium***
Collective migration is a pivotal morphogenetic process during liver organogenesis.
Tools for inducing collective migration *in vitro* are described here. Successful droplet formation
of HEP spheroids and the use of MES-CM demonstrate outgrowth branching from the HEP
spheroids (**Figure 12A-B**). HEP spheroids cultivated in the MG droplet system show that MES-
CM induces collective migration, and the data demonstrate a concentration-dependent effect
(**Figure 12B, right**). Cellular strands protruding from the hepatic spheroid are present with small
branching, thick strands, and multiple levels of branching (**Figure 12B, left**). On day 11, the
protrusions increased with interconnections and sheet formation (**Figure 12B, right**). MES-CM is
potent, as a 1:7 dilution or MES-CM present for one day still leads to migration (**Figure 12B,
right**). It was hypothesized that the MES-CM induced cell migration via TGFβ signaling pathway.
A83-01, a TGFβ pathway inhibitor, was incorporated into the MES-CM at varying
concentrations, and migration was significantly inhibited in a dose-dependent manner (**Figure
12C**). To determine the effects of extracellular matrix on collective migration, HEP spheroids
cultivated in the CG droplet system demonstrate that MES-CM induces cell migration. However,
the protrusions were thin linear strands and less branching by day 7 compared to the migration
observed in MG (**Figure 12D**). Fibrin hydrogels were also tested, and thin, hair-like, narrow,
radial protrusions of both HEP and MES cells were observed (**Figure 12E**).
608
***Inducing hepatic co-migration via MES conditioned medium***
Mixed spheroids (HEP-MES spheroids) were utilized for collective migration, as a model
of co-migration which occurs during early liver organogenesis (**Figure 12F**). These mixed
spheroids in MG result in migration, and when TGFβ1 growth factor was added, it resulted in
significantly increased collective migration (**Figure 12G**). Thus, co-migration can also be

modeled with mixed spheroids.
615

**FIGURE AND TABLE LEGENDS:**
**Figure 1. Schematic of spheroid formation assay. (A)** Coating a 96-well plate with 1% agarose
solution. Agarose solution is melted in the microwave, 55-65 µL is transferred to each well and
the plate is cooled for 20-30 minutes. **(B)** Passaging of cell culture at 80% confluency. **(C)** Dye-
labeling of cell suspension. 5 µL of cell-labeling solution is added per mL After incubation and
centrifugation, the cell suspension is washed with warm cDMEM before plating the cells. **(D)**
Plating of cell suspension for spheroid formation. 100 µL of cell suspension is added to each
well and the plate is centrifuged. Mesodermal-derived spheroids form in 1-3 days, while hepatic
spheroids form in 5-9 days.
625

**Figure 2. Schematic of assembloid formation assay with hepatic and mesodermal-derived**
**spheroids. (A)** Hepatic spheroid formation. HepG2 cells are dye-labeled and plated for spheroid
formation. **(B)** Mesodermal-derived spheroid formation. HFF/MRC-5 cells are dye-labeled and
plated for spheroid formation. **(C)** HEP-MES assembloid formation. Transfer mesodermal-
derived spheroid to hepatic spheroid for assembloid formation. Matrix (MG or CG) is added
(1:1-1:5 dilution), incubated for 3 hours, followed by the addition of 75 µL cDMEM. Incubated
for 2-3 days until assembloid formation is complete.
633

**Figure 3. Schematic of Droplet Formation Assay. (A)** Hepatic spheroid formation. HepG2 cells
are dye-labeled. **(B)** Spheroid collection and matrix suspension. Collect and gently rinse hepatic
spheroids with cDMEM. Spheroids are suspended in MG or CG solution (1:1 – 1:5 dilution). **(C)**
Droplet seeding and addition of growth medium. Hepatic spheroids are seeded onto a 60mm
petri dish and incubated for 60 minutes, and 5 mL of MES-CM is added. Petri dish is incubated
with media changes every three days.
640

641  **Figure 4. Schematic of Preparation of MES-CM. (A)** Cell seeding and MES-CM collection.
642  HFF/MRC-5 cells are seeded at 5,000 cells/cm$^2$ in a T-75 flask and incubated for 72 hours. The
643  media is collected and centrifuged for debris removal. **(B)** MES-CM sterilization and addition.
644  The media is sterilized with a 0.2 µm filter and combined with cDMEM (1:1-1:7 dilution). Media
645  is added to a hepatic spheroid and incubated with media changes every three days.
646

**Figure 5. Schematic of Hepatic and Mesodermal-derived Mixed Spheroid Formation Assay. (A)**
Preparation of hepatic cell suspension. HepG2 cells are harvested and dye-labeled. **(B)**
Preparation of Mesodermal-derived cell suspension. HFF/MRC-5 cells are harvested and dye-
labeled. **(C)** HEP-MES mixed spheroid formation. HepG2 and HFF/MRC-5 cells are mixed at a 1:1
ratio at a final concentration of 2.0 x 10$^5$ cells/mL and plated. The plate is incubated for 1-2 days
until mixed spheroid formation is complete.
653

**Figure 6. Schematic of High Density HFF/MRC-5 Migratory Model. (A)** Hepatic spheroid
formation. HepG2 cells are dye-labeled and plated for spheroid formation. **(B)** Preparation of
Mesodermal-derived cell suspension. HFF/MRC-5 cells are harvested and dye-labeled. **(C)**
HFF/MRC-5 cell suspension is pre-mixed with MG solution and added to pre-existing HepG2

well. The plate is incubated for 30 minutes and 75 µL of cDMEM is added. The plate is
incubated for up to 12 days.
660
661 **Figure 7. Overview.** Based in the science of morphogenesis (top), stem cells will be
662 implemented to build assembloids. The toolbox (middle), through the use of cells, tools, and
663 biomatrix, and together with aspects of morphogenesis, can together be used to generate new
664 synthetic tissues and assembloids.
665
666 **Figure 8. Compaction time of varying spheroid size. (A)** Phase-contrast images of small (1,500
667 cells) and large (3,000 cells) hepatic spheroids on days 1, 3, and 5. Bar = 300 µm. **(B)** Same study
668 as **(A)** except HFF cells. Bar = 100 µm. **(C)** Bar graph comparing HepG2 and HFF spheroid
669 compaction. (n = 5). Right: bar graph plotting area of HepG2 and HFF spheroids over time. (n =
670 5).
671
**Figure 9. The effect of media and matrix on assembloid formation.** Phase contrast and
fluorescent images of fused HepG2/MRC-5 spheroids exposed to varying matrices (1:5 dilution)
and growth medium: DMEM in CG, MES-CM in CG, DMEM in MG, and MES-CM in MG. Left:
phase-contrast (day 2) images of fused HepG2/MRC-5 assembloid. Bar = 100 µm. Middle:
fluorescent (day 2) of fused HepG2/MRC-5 assembloid. Bar = 200 µm. Left: phase-contrast of
fused HepG2/MRC-5 assembloid. Bar = 200 µm. (green: HepG2; orange: MRC-5; arrows
indicating spheroid types).
679
**Figure 10. Engineering assembloids arms, junctions, and layers. (A)** Schematic of building of
large and small arms onto hepatic spheroids. **(B)** Phase-contrast images of HepG2 spheroids
suspended in the MG droplet system, bearing high density (30,000 cells) of MRC-5 cells. Top:
days 3, 4, 9, and 12. Day 3: MRC-5 cells initially after seeding (arrow). Bar = 1,000 µm. Day 4:
MRC-5 cells spreading and interconnecting (arrows). Bar = 1,000 µm. Day 9: Thick hepatic cord
(arrows). Bar = 400 µm. Day 12: multiple, thick hepatic cords. Bar = 400 µm. Bottom: images of
a separate experimental replicate on day 12 demonstrating thick hepatic cord formation
(arrows). Bar = 400 µm. Subfigures obtained with permission from Frontiers Journal[17]. **(C)**
Fluorescent images of hepatic spheroid co-cultured with a HEP-MES mixed spheroid on days 0
and 5. Day 0: HepG2 and mixed spheroid initially after plating. Bar = 100 µm. Day 5: mixed
spheroid has fused to the edge of HepG2 spheroid. Bar = 100 µm. (need to add dye-labeling
information) **(D)** Schematic of building HEP-MES assembloids with fused edges (spot-welding).
(arrows indicating spheroid type). **(E)** Phase-contrast images of HEP and MES spheroids co-
cultivated in the CG droplet system exposed to MES-CM. Day 0: HEP and MES spheroids initially
after plating. Bar = 100 µm. Day 1: HEP and MES migrating towards one another, but no fusion
has occurred. Bar = 100 µm. Day 4: HEP and MES spheroids have fused at the edges of the
spheroids. Bar = 100 µm. **(F)** Phase-contrast and fluorescent images of HEP-MES assembloid
cultivated in CG droplet system (2 mg/mL) on day 3 demonstrating fused edges. (green: HepG2;
orange: MRC-5; arrow indicating where fusion occurred). Bar = 100 µm. **(G)** Schematic of
building HEP-MES assembloids with infiltration or a surface-MES layering. **(H)** Phase-contrast
and fluorescent images of HEP-MES assembloid cultivated in CG droplet system (2 mg/mL)
suspended in MES-CM on day 3 demonstrating infiltration of spheroids. (green: HepG2; orange:

MRC-5; arrow indicating where fusion occurred). Bar = 100 μm. **(I)** Phase-contrast and fluorescent images on day 6 of HepG2 spheroid and MSC in MG droplet system. The right columns on the right are replicates 1 and 2 with fluorescent images on day 6. (green: HepG2; red: MSC). Bar = 100 μm. Subfigures obtained with permission from Frontiers Journal[17].

**Figure 11. Partially and completely fused HEP-MES assembloids. (A)** Schematic of building fused HEP-MES assembloids with increased packing density consisting of separate layers. **(B)** Phase-contrast images of HepG2 and MRC-5 spheroids co-cultivated in MG droplet system on days 3, 5, 7, 9. Days 3-5: HepG2 and MRC-5 spheroids interact. Day 7: migration at the spheroids edge and fusion occurred. Day 9: spheroid fusion and assembloid formation; arrows demonstrate protrusions. Bar = 200 μm. Subfigures obtained with permission from Frontiers Journal[17]. **(C)** Fluorescent images of HEP-MES assembloid, the same study as **(B)** on day 13. Day 9: spheroid fusion and assembloid formation; arrows demonstrate protrusions. (green: HepG2; red: MRC-5). Bar = 200 μm. Subfigures obtained with permission from Frontiers Journal[17]. **(D)** Phase-contrast images of HepG2 spheroid co-cultured with multiple mesodermal-derived spheroids on days 0, 1, and 5. Day 5: HEP-MES assembloid formation occurred but overall volume did not increase. **(E)** Phase-contrast images of HepG2 and MRC-5 spheroids co-cultivated in MG under low serum conditions (0.2% and 2% FBS) on days 0-2. Top: HEP-MES assembloid fusion occurs in 2% FBS by day 2. Bar = 100 μm. Bottom: HEP-MES assembloid fusion occurs in 0.2% FBS by day 2. Bar = 100 μm. **(F)** Phase-contrast images of HepG2 and MRC-5 spheroids co-cultured in the CG droplet system exposed to MES-CM. Day 0: HEP and MES spheroids initially after plating. Bar = 100 μm. Day 5: HEP and MES spheroids have not fused. Bar = 100 μm. **(G)** Fluorescent images of MES spheroid co-cultured with a HEP-MES mixed spheroid on days 0 and 5. Day 0: HFF and HEP-MES mixed spheroid initially after plating. Bar = 100 μm. Day 5: HFF and HEP-MES mixed spheroid fuse. Bar = 100 μm. **(H)** Schematic of building partially fused HEP-MES assembloid with cupping. **(I)** Phase-contrast images of HEP and MES spheroids co-cultured in the CG droplet system on days 0, 1, 2, and 4. Day 0: HEP and MES spheroids initially after plating. Days 1-2: HEP and MES spheroids interact. Day 4: HEP-MES assembloid formation occurs in a cup-like mechanism. Arrow indicating spheroid position during fusion. Bar = 100 μm. **(J)** Phase-contrast and fluorescent images of HEP-MES assembloid cultivated in MG droplet system on days 2 and 4. Top: HEP and MES spheroid interaction occurs. Bottom: partial fusion of HEP-MES assembloid. (green: HepG2; orange: MRC-5; arrow indicating where fusion occurred). Bar = 100 μm. **(K)** Phase-contrast images of MES spheroid co-cultured with multiple HEP spheroids on days 0, 1, and 5. Day 0: MES and HepG2 spheroids initially after platting. Day 1: HEP and MES spheroid interaction occurs. 5: HEP spheroid forms a cup around MES spheroid.

**Figure 12. Methods for modulating collective migration from spheroids. (A)** Schematic of branching and linear migration in HEP spheroids in the presence of MES-CM. **(B)** Phase-contrast images of HEP spheroid in the MG droplet system cultivated in varying dilutions of MES-CM on day 7 (low and high magnification) and 11. Day 7: HEP spheroid demonstrates various modes of 3D collective migration. Bar = 300 μm. Day 11: HEP spheroid outgrowth demonstrates interaction between branches. Bar = 500 μm. Plot comparing fold-increase in area across different dilution ratios of MES-CM (1:1, 1:7, MES-CM 1 day, and MES-CM).Comparison of MES-

CM with 1:1 condition (P = 0.15, n = 3 for both conditions), MES-CM with 1:7 condition (P =
0.0056, n = 3 for both conditions), and MES-CM with MES-CM 1-day-only condition (P = 0.019, n
= 3 for both conditions). Subfigures obtained with permission from Frontiers Journal[17]. **(C)**
Phase-contrast images of HEP spheroid cultivated MG droplet system exposed to MES-CM
alone, MES-CM with A83-01 (10 nM), and A83-01 (20 nM) on day 7. Bar = 300 µm. Bar graph
analysis comparing day 7 MES-CM and MES-CM + A83-01 (20 nM) (P = 0.047, n = 3). Plotted
means ± SD. Significance is defined as P ≤ 0.05. Subfigures obtained with permission from
Frontiers Journal[17]. **(D)** Phase-contrast images of HEP spheroid cultured in the CG droplet
system cultivated in MES-CM on days 4 and 7. Arrows specify thin filopodia-like extensions into
the CG. Bar = 500 µm. Bar graph comparing protrusion length of HEP spheroids in cDMEM and
MES-CM conditions (P = 0.012, n = 3). Subfigures obtained with permission from Frontiers
Journal[17]. **(E)** Phase-contrast and fluorescent images of HEP-MES mixed spheroid cultivated in
fibrin gels exposed to MES-CM on days 4 and 5. (green: HepG2; red: HFF; arrows indicated
protrusions). Bar = 500 µm. Subfigures obtained with permission from Frontiers Journal[17]. **(F)**
Schematic of star-shaped migration of HEP-MES mixed spheroids. **(G)** Fluorescent images of
HEPG2-GFP (HepG2 cells expressing green fluorescent protein) and HEP-MES mixed spheroids
co-cultivated in the MG droplet system after treatment with TGFβ1 (20 ng/ml) on day 4. Left:
HepG2 cells in HEP-MES mixed spheroid exposed to TGFβ1 on day 4. Right: HEP-MES mixed
spheroid exposed to TGFβ1 on day 4. Replicates 1 (below) and 2 (above) are shown. (green:
HepG2; red: MRC-5; yellow: combined; arrows show HepG2 and MRC-5 migration). Bar = 500
µm. Bar graph comparing the area of fibroblast migration when exposed to TGFβ1 (HepG2-
GFP/MRC-5 and HepG2-GFP/MRC-5 + TGFβ1, 20 ng/ml), P = 0.012, n = 3. Plotted means ± SD.
Significance is defined as P ≤ 0.05. ∗ is used to denote the significance of experimental data.
Subfigures obtained with permission from Frontiers Journal[17].

**Table 1. The effect of distance on assembloid formation.** Hepatic spheroids and mesodermal-
derived spheroids were transferred to an agarose-coated well. The cells were dye-labeled prior
to spheroid formation to demonstrate the interaction between spheroids. The spheroids were
suspended in a 1:5 solution of the MG matrix suspended in cDMEM. The initial distance
between the spheroids was measured using ImageJ and the corresponding compaction time of
the assembloid formation was observed. The compaction time of assembloid formation is
directly proportional to the initial distance between the spheroids. (n = 3).

**DISCUSSION:**
In this protocol, multiple methods are presented for cultivating both simple and
complex assembloids, as well as techniques for inducing 3D collective cell migration in early
liver organogenesis[14, 17]. Several critical steps are outlined in the presented protocols, with
spheroid formation being a key aspect across all methods. Achieving spheroid formation
involves using microwells (96- or 384-wells) with either non-adherent or agarose-coated plates.
Spheroids are abundantly produced for subsequent experimentation, accounting for those that
may not reach the expected size or have irregular shapes. Spheroids that do not meet the
desired criteria are not utilized in downstream experiments. The initial efficiency depends on
various factors, including the user. Considerable expertise is required to handle organoids,
involving aspects such as spheroid (or organoid) formation, transfer between wells, addition of

biomatrices upon spheroids, and placement of multiple spheroids per well.

Formation of the spheroids demands meticulous attention to repeatable cell counting
and seeding, agarose coating (or non-adherent), regular medium changes, dye-labeling
methods, and gentle handling of spheroids and plates, along with microscopy. Dye-labeling
specifics, including cell number and type, as well as, the labeling duration, should be
experimentally determined. Typically, concentration of MG averaged 9.02 ± 0.63 mg/mL.
Furthermore, experience suggests that minor differences do not significantly impact
phenotype, it could be highly beneficial to track protein concentrations for each MG lot in initial
experiments. In terms of the CG, the concentration was prepared to be 2 mg/mL.

For instance, preparing the agarose solution necessitates strict adherence to
maintaining appropriate concentrations, seeding wells with proper volumes to allow for the
development of a meniscus (curvature), and enable cell collection in the center. Careful
consideration of agarose concentration, volume, and gel solidification is crucial. Selecting the
correct micropipette tip size and employing proper transferring techniques are essential for
handling spheroids effectively. The use of glass wells can enhance visualization.  As observed in
the paper, spheroids initially appear as translucent discs, becoming more opaque over time,
providing a means to monitor spheroid formation. Several factors can influence this process.
Lastly, for assembloid formation[9], precise transfer of spheroids is vital, placing them within a
couple of spheroid diameters or less to effectively observe changes.

Here, a toolbox of methods has been established using cell lines to develop complex
assembloids and migrating spheroids. It is noteworthy that cell lines were employed for method
focus, but using primary cells or hPSC-derived cells would be advantageous for human
personalized tissues, and our assembloid techniques have been validated with hPSC-derived
HEP cells[14,15]. However, a significant limitation of these systems is the use of GFR MG, a mouse
tumor extracellular matrix protein mixture primarily used in these cultivation models. However,
it is a major limitation due to its tumor-derived origin and high cost. Additionally, the droplet
formation system degrades significantly after two weeks, suggesting the implementation of
alternative gels such as CG mixtures or sodium alginate. Furthermore, the current systems can
only grow to the dimensions of the spheroids, but not beyond, emphasizing the need to identify
and address limitations to growth.

The lack of methods to specifically study early events in early liver organogenesis[14] and,
more generally, methods to generate synthetic tissues[1]. These methods offer a toolbox for
building assembloids and assembling larger, more complex tissues for enhanced *ex vivo* organ
modeling. This holds potential for *in vivo* therapeutic approaches, in addition to modeling
structures during organogenesis (liver bud) and applications in biopharma.

The protocols presented enable models of different modes of liver cell migration,
including co-migration, interstitial migration, and branching morphogenesis, along with
assembloids that collectively can be used to construct more complex tissues for various
biomedical and biopharma applications.

839 **ACKNOWLEDGMENTS:**
840 Authors thank Jenna Venturo, undergraduate in the lab, for her assistance. NP was supported
841 by the UB CBE startup funds, a Mark Diamond Fellowship, the New York State Stem Cell Science
842 C024316, the UB the Stem cells in regenerative medicine (ScIRM) center, the University at
843 Buffalo Center for Cell, Gene, Tissue Engineering (CGTE).
844

845 **DISCLOSURES:**
846 NP is the founder of Khufu Therapeutics, an organoid engineering company that develops
847 treatments for acute and chronic liver disease.
848

849 **REFERENCES:**
850
1      Bayley, H., Cazimoglu, I., Hoskin, C. E. G. Synthetic tissues. *Emerg Top Life Sci.* **3** (5), 615-622
(2019).
2      Biname, F., Lassus, P., Hibner, U. Transforming growth factor beta controls the directional
migration of hepatocyte cohorts by modulating their adhesion to fibronectin. *Mol Biol Cell.* **19**
(3), 945-956 (2008).
3      Bort, R., Signore, M., Tremblay, K., Martinez Barbera, J. P., Zaret, K. S. Hex homeobox gene
controls the transition of the endoderm to a pseudostratified, cell emergent epithelium for liver
bud development. *Dev Biol.* **290** (1), 44-56 (2006).
4      Chen, J. S. et al. Mir-338-3p inhibits epithelial-mesenchymal transition and metastasis in
hepatocellular carcinoma cells. *Oncotarget.* **8** (42), 71418-71429 (2017).
5      Fitamant, J. et al. Yap inhibition restores hepatocyte differentiation in advanced hcc, leading to
tumor regression. *Cell Rep.* **10** (10), 1692-1707 (2015).
6      Fransvea, E., Angelotti, U., Antonaci, S., Giannelli, G. Blocking transforming growth factor-beta
up-regulates e-cadherin and reduces migration and invasion of hepatocellular carcinoma cells.
*Hepatology.* **47** (5), 1557-1566 (2008).
7      Gualdi, R. et al. Hepatic specification of the gut endoderm in vitro: Cell signaling and
transcriptional control. *Genes Dev.* **10** (13), 1670-1682 (1996).
8      Gupta, S. et al. Entry and integration of transplanted hepatocytes in rat liver plates occur by
disruption of hepatic sinusoidal endothelium. *Hepatology.* **29** (2), 509-519 (1999).
9      Kanton, S. Pasca, S. P. Human assembloids. *Development.* **149** (20), (2022).
10     Koenig, S., Stoesser, C., Krause, P., Becker, H., Markus, P. M. Liver repopulation after
hepatocellular transplantation: Integration and interaction of transplanted hepatocytes in the
host. *Cell Transplantation.* **14** (1), 31-40 (2005).
11     Matchett, K. et al. Multimodal decoding of human liver regeneration. *bioRxiv.*
10.1101/2023.02.24.529873 2023.2002.2024.529873 (2023).
12     Matsumoto, K., Yoshitomi, H., Rossant, J., Zaret, K. S. Liver organogenesis promoted by
endothelial cells prior to vascular function. *Science.* **294** (5542), 559-563 (2001).
13     Ng, L. et al. Suppression of actopaxin impairs hepatocellular carcinoma metastasis through
modulation of cell migration and invasion. *Hepatology.* **58** (2), 667-679 (2013).

14    Ogoke, O. et al.  10.1101/2022.07.24.501313 (2023).

15    Ogoke, O. et al. Spatiotemporal imaging and analysis of mouse and human liver bud
morphogenesis. *Dev Dyn.* **251** (4), 662-686 (2022).

16    Ogoke, O., Oluwole, J., Parashurama, N. Bioengineering considerations in liver regenerative
medicine. *J Biol Eng.* **11** 46 (2017).

17    Ogoke, O. et al. Modeling liver organogenesis by recreating three-dimensional collective cell
migration: A role for tgfbeta pathway. *Front Bioeng Biotechnol.* **9** 621286 (2021).

18    Petkov, P. M., Kim, K., Sandhu, J., Shafritz, D. A., Dabeva, M. D. Identification of differentially
expressed genes in epithelial stem/progenitor cells of fetal rat liver. *Genomics.* **68** (2), 197-209
(2000).

19    Rajvanshi, P., Kerr, A., Bhargava, K. K., Burk, R. D., Gupta, S. Studies of liver repopulation using
the dipeptidyl peptidase iv-deficient rat and other rodent recipients: Cell size and structure
relationships regulate capacity for increased transplanted hepatocyte mass in the liver lobule.
*Hepatology.* **23** (3), 482-496 (1996).

20    Rossi, J. M., Dunn, N. R., Hogan, B. L., Zaret, K. S. Distinct mesodermal signals, including bmps
from the septum transversum mesenchyme, are required in combination for hepatogenesis
from the endoderm. *Genes Dev.* **15** (15), 1998-2009 (2001).

21    Sosa-Pineda, B., Wigle, J. T., Oliver, G. Hepatocyte migration during liver development requires
prox1. *Nat Genet.* **25** (3), 254-255 (2000).

22    Suzuki, A., Sekiya, S., Büscher, D., Izpisúa Belmonte, J. C., Taniguchi, H. Tbx3 controls the fate of
hepatic progenitor cells in liver development by suppressing p19arf expression. *Development.*
**135** (9), 1589-1595 (2008).

23    Xue, T. C. et al. Goosecoid promotes the metastasis of hepatocellular carcinoma by modulating
the epithelial-mesenchymal transition. *PLoS One.* **9** (10), e109695 (2014).

24    Yang, C. et al. Mir-1301 inhibits hepatocellular carcinoma cell migration, invasion, and
angiogenesis by decreasing wnt/beta-catenin signaling through targeting bcl9. *Cell Death Dis.* **8**
(8), e2999 (2017).

25    Yang, M. H. et al. Comprehensive analysis of the independent effect of twist and snail in
promoting metastasis of hepatocellular carcinoma. *Hepatology.* **50** (5), 1464-1474 (2009).

26    Zeng, Y. B. et al. Mirna-135a promotes hepatocellular carcinoma cell migration and invasion by
targeting forkhead box o1. *Cancer Cell Int.* **16** 63 (2016).

27    Zhao, Y. et al. Rnai silencing of c-myc inhibits cell migration, invasion, and proliferation in hepg2
human hepatocellular carcinoma cell line: C-myc silencing in hepatocellular carcinoma cell.
*Cancer Cell International.* **13** (1), 23 (2013).

A

Passage HepG2-WT cell culture and suspend at a density of 1 x 10⁶ cells/mL

Dye-label the cells and suspend at a density of 5 x 10⁴ cells/mL

Add 100 μL of cell suspension to per well and centrifuge 340 *x g* for 10 min

Incubate for 5-9 days until spheroid formation is complete

B

Passage HFF/MRC-5 cell culture and suspend at a density of 1 x 10⁶ cells/mL

Dye-label the cells and suspend at a density of 1 x 10⁵ cells/mL

Add 100 μL of cell suspension to per well and centrifuge 340 *x g* for 10 min

Incubate for 1-3 days until spheroid formation is complete

C

Transfer a single spheroid to a pre-existing well

Add matrix (MG or CG) at a 1:1-1:5 dilution and incubate for 3 h

Add 75 μL of cDMEM and incubate for 2-3 days for assembloid formation

A

Passage HepG2-WT cell culture and suspend at a density of 1 x 10⁶ cells/mL → Dye-label the cells and suspend at a density of 5 x 10⁴ cells/mL → Add 100 μL of cell suspension to per well and centrifuge 340 *x g* for 10 min → Incubate for 5-9 days until spheroid formation is complete

B

Collect the HepG2-WT spheroids and gently rinse with cDMEM → Suspend the spheroids in matrix (MG or CG)

C

Seed one spheroid in 15 μL of matrix (MG or CG) onto 60mm petri dish and incubate at 37°C for 60 min → Slowly add 5 mL of growth medium and incubate at 37°C

A                                                                              B

Seed 5,00 cells/cm² in 15 mL of
cDMEM (T-75 flask)

72 h incubation

Collect conditioned media
and centrifuge at 290 x g
for 5 min

Filter conditioned media
using a 0.2 µm syringe
filter

Dilute conditioned media
with cDMEM (1:1-1:7)

A

Passage HepG2-WT cell culture and suspend at a density of 1 x 10⁶ cells/mL

Dye-label the cells and re-suspend in cDMEM

B

Passage HFF/MRC-5 cell culture and suspend cells at a density of 1 x 10⁶ cells/mL

Dye-label the cells and re-suspend in cDMEM

C

Suspend HepG2-WT and HFF/MRC-5 cells at 1:1 ratio at a density of 2.0 x 10⁵ cells/mL

Add 100 µL of cell suspension to per well and centrifuge 340 x g for 10 min

Incubate for 1-2 days for mixed spheroid formation

A

Passage HepG2-WT cell culture. Suspend cells at a density of 1 x 10⁶ cells/mL

Dye-label the cells and suspend at a density of 5 x 10⁴ cells/mL

Add 100 µl of cell suspension to per well and centrifuge 340 *x g* for 10 mins

Incubate for 5-9 days until spheroid formation is complete

B

Cell culture has reached 80% confluency

Passage cells using Tryspin-EDTA

Suspend cells at a density of 1 x 10⁶ cells/mL of serum-free medium and add 5 µL of cell-labeling solution

Add 5 µL of cell-labeling solution per mL and incubate for 20 min

Centrifuge at 450 *x g* for 5 min and wash cells with cDMEM. Repeat this washing process two more times.

C

Dilute cell suspension to 6.0 x 10⁵ cells/mL and mix with ice-cold MG at a 1:1 dilution.

Add 100 µL of HFF/ MRC-5 and MG solution to a pre-existing well and incubate for 30 min

Add 75 µL of cDMEM and incubate for up to 12 days

A HepG2-WT

Small (S)
1500 cells/well

Large (L)
3000 cells/well

D1　　　D3　　　D5

B HFF

Small (S)
1500 cells/well

Large (L)
3000 cells/well

D1　　　D3　　　D5

C

**A** Building Large or Small Arms

HFF/MRC-5
HepG2-WT

**B** HEP-MES Assembloids demonstrate branching arms

Day 0    Day 3    Day 4-6
1000 HepG2 cells GFP/Flux or WT 384 well RB plate    GFR Matrigel 20,000 MRC-5

Day 3    Liver/MRC5    1000 µm
Day 4    Liver/MRC5    1000 µm
Day 9    Liver/MRC5    400 µm
Day 12    Liver /MRC5    400 µm
Day 12    Liver /MRC5    400 µm

Mixed spheroid form bridges with HEP spheroids

**C**
HepG2-WT    100 µm
Mixed Spheroid
Day 0

HepG2-WT    Bridge
100 µm
Mixed Spheroid
Day 5

**D** Spot-Welding (fused edges)

Day 0    Day 3-4

HFF/MRC-5
HepG2-WT

**E** HEP-MES in collagen with MES-conditioned medium fuse edges and don't form cupping

Day 0    M-CM
M    H
100 µm    CG

Day 1    M    H
100 µm    CG

Day 4    M-CM
M    H
100 µm    CG

**F** In Collagen 2 mg/ml (stiff), HEP-MES edges fuse

Collagen 2 mg/ml    HEPG2-WT
MRC5
100 µm    Day 3

Collagen 2 mg/ml    H
M
100 µm    Day 3

**G** Infiltrating and Layering

HFF/MRC-5
HepG2-WT

**H** HEP-MES infiltrate in Collagen, with MES- conditioned medium

Collagen 2 mg/ml MES-condition medium MCM
100 µm    Day 3

Collagen 2 mg/ml MCM
M
H
100 µm    Day 3

**I** HEP-inside, MES-outside leads to layered assembloid

Replicate 1    Replicate 2

Day 6
100 µm

Day 6
100 µm    HepG2/MSC

Day 6
100 µm    HepG2/MSC

**A** Fused HEP-MES assembloids (increased packing density, separate layers)

Day 5 → Day 7 → Day 9

■ HFF/MRC-5
■ HepG2-WT

**B** HEP and MES spheroids fuse in matrigel

Day 3    Day 5    Day 7    Day 9
L MRC5   L MRC5   L MRC5   L MRC5
200 µm   200 µm   200 µm   200 µm

**C** Day 13
MRC5 L    MRC5    L
200 µm    200 µm    200 µm

**D** HEP-MES assembloids with multiple MES spheroids fuse but overall volume does not increase

Day 0    Day 1    Day 5
H M      H M      H M
M        M        M
100 µm   100 µm   100 µm

**E** HEP-MES assembloids form under low serum

2% FBS    2% FBS    2% FBS
M         M         M
H         H         H
100 µm    100 µm    100 µm
Day 0     Day 1     Day 2

0.2% FBS   0.2%FBS   0.2%FBS
M          M          M
H          H          H
100 µm     100 µm     100 µm
Day 0      Day 1      Day 2

**F** HEP-MES don't form at large distances in collagen with MES-conditioned medium

Day 1    M-CM          Day 5    M-CM
M        H             M        H
         CG                     CG
100 µm                 100 µm

**G** Mixed spheroids fuse with MES spheroids

Day 0              Day 5
Mixed              HFF
spheroid           Mixed
        HFF        spheroid
100 µm             100 µm

**H** Cupping, partially fused HEP-MES assembloids (increased packing density, separate layers)

■ HFF/MRC-5
■ HepG2-WT

**I** In collagen, HEP-MES form assembloids via cup-like mechanism

Day 0    M        Day 1    M
    H             H
100 µm   100 µm

Day 2    M        Day 4    M
    H             H
100 µm   100 µm          CG

**J** HEP-MES cupping prior to fusion in MG

Matrigel    H        Matrigel    H
M                    M
100 µm    Day 2      100 µm    Day 2

Matrigel    H        Matrigel    H
M                    M
100 µm    Day 4      100 µm    Day 4

**K** HEP-MES assembloids with multiple HEP spheroids combine with cupping mechanism

Day 0              Day 1              Day 5
H    H             H                  H
100 µm             H                  H
H                  M         H        M
     M             H                  H
100 µm             100 µm

**A** Branching    Linear

HepG2-WT

**E** HEP-MES mixed spheroids demonstrate hair-like migration in Fibrin gels

Day 4 HepG2/HFF    Day 5    HFF    HepG2    HFF HepG2
Fib    Fib    Fib    Fib

**B** M-CM Day 7    M-CM Day 11

P = NS

1:1  1:7  M-CM  M-CM
              3 day

Fold-increase in area

**F** Star-shaped

HFF/MRC-5
HepG2-WT

**C** Branching morphogenesis of HEP-spheroids inhibited by TGF-beta

M-CM    M-CM + 10 nM A83-01    M-CM + 20 nM A83-01
Day 7    Day 7    Day 7

Max Branch Length/Mean Branch Length

D7 MCM  D7 (10 nM)  D7 (20 nM)

**G** Increasing branch length in HEP-MES mixed spheroids with TGF-Beta

HepG2/MRC5 in Matrigel

Replicate 2    Day 4 HepG2    HepG2/MRC5
Replicate 1    Day 4 HepG2    HepG2/MRC5

Area of Fibroblast Migration (μm²)

Control    TGFB (+)

**D** HEP spheroids demonstrate linear branching in collagen gels

M-CM Day 4    M-CM Day 7
CG    CG

Protrusion length (μm)

Cntrl    M-CM

Tgfb + MRC5/ HepG2 in Matrigel

Replicate 2    Day 4 HepG2    HepG2/MRC5
Replicate 1    Day 4 HepG2    HepG2/MRC5

| Growth Medium & Matrix | Initial Distances (µm) | Days to Merge |
| --- | --- | --- |
| Matrigel + DMEM | 492.742 | 6 |
| Matrigel + DMEM | 580.142 | 8 |
| Matrigel + DMEM | 974.051 | 12 |