Sex-Specific DNA Methylation and Gene Expression Changes in Mouse Placentas After Early Preimplantation Alcohol Exposure

Lisa-Marie Legault[1,2], Thomas Dupas[1,3], Mélanie Breton-Larrivée[1,2], Fannie Filion-Bienvenue[1,2],

Anthony Lemieux[1], Alexandra Langford-Avelar[1,2], Serge McGraw[1,2,3,#]

[1]CHU Ste-Justine Research Center, 3175 Chemin de la Côte-Sainte-Catherine, Montréal, QC H3T 1C5, Canada

[2]Department of Biochemistry and Molecular Medicine, Université de Montréal, 2900 Boulevard Edouard-Montpetit, Montréal, QC H3T 1J4, Canada

[3]Department of Obstetrics and Gynecology, Université de Montréal, 2900 Boulevard Edouard-Montpetit, Montréal, QC H3T 1J4, Canada

Lisa-Marie Legault : lisa-marie.legault@umontreal.ca

Thomas Dupas : thomas.dupas@umontreal.ca

Mélanie Breton-Larrivée : melanie.breton-larrivee@umontreal.ca

Fannie Filion-Bienvenue : fannie.filion-bienvenue@umontreal.ca

Anthony Lemieux : anthony.lemieux@umontreal.ca

Alexandra Langford-Avelar : alexandra.langford.avelar@umontreal.ca

Serge McGraw : serge.mcgraw@umontreal.ca (corresponding author)

ABSTRACT

During pregnancy, exposure to alcohol represents an environmental insult capable of negatively impacting embryonic development. This influence can stem from disruption of molecular profiles, ultimately leading to manifestation of fetal alcohol spectrum disorder. Despite the central role of the placenta in proper embryonic development and successful pregnancy, studies on the placenta in a prenatal alcohol exposure and fetal alcohol spectrum disorder context are markedly lacking. Here, we employed a well-established model for preimplantation alcohol exposure, specifically targeting embryonic day 2.5, corresponding to the 8-cell stage. The exposure was administered to pregnant C57BL/6 female mice through subcutaneous injection, involving two doses of either 2.5 g/kg 50% ethanol or an equivalent volume of saline at 2-hour intervals. Morphology, DNA methylation and gene expression patterns were assessed in male and female late-gestation (E18.5) placentas. While overall placental morphology was not altered, we found a significant decrease in male ethanol-exposed embryo weights. When looking at molecular profiles, we uncovered numerous differentially methylated regions (DMRs; 991 in males; 1309 in females) and differentially expressed genes (DEGs; 1046 in males; 340 in females) in the placentas. Remarkably, only 21 DMRs and 54 DEGs were common to both sexes, which were enriched for genes involved in growth factor response pathways. Preimplantation alcohol exposure had a greater impact on imprinted genes expression in male placentas (imprinted DEGs: 18 in males; 1 in females). Finally, by using machine learning model (L1 regularization), we were able to precisely discriminate control and ethanol-exposed placentas based on their specific DNA methylation patterns. This is the first study demonstrating that preimplantation alcohol exposure alters the DNA methylation and transcriptomic profiles of late-gestation placentas in a sex-specific manner. Our findings highlight that the DNA methylation profiles of the placenta could serve as a potent predictive molecular signature for early preimplantation alcohol exposure.

KEYWORDS

Preimplantation, prenatal alcohol exposure, DNA methylation, transcriptome, placenta, machine learning

GRAPHICAL ABSTRACT

Created with BioRender.com

INTRODUCTION

The maternal environment is essential for a healthy pregnancy and embryonic development. Adverse effects of alcohol exposure at any point during pregnancy are referred to as fetal alcohol spectrum disorder (FASD) (1). FASD manifests as various symptoms, including cognitive impairments, learning difficulties, behavioral issues, and, in severe cases, profound intellectual disabilities and craniofacial abnormalities (1). Annually, approximately 630,000 newborns are affected by FASD worldwide (2). Its presentation and severity depend on multiple factors, including the timing, amount, and pattern of prenatal alcohol exposure (PAE) (1). Poor maternal health, exposure to other environmental insults (e.g. tobacco, drugs, contaminants), and nutritional deficiencies can also increase FASD severity (1). Currently, there is no molecular diagnostic tool for FASD, and most cases are not identified until school-aged children, when neurodevelopmental symptoms become noticeable (1). Early diagnosis and management can alleviate the impacts of FASD, while late detection often results in a need for lifelong health and social care (1).

The placenta is a transient organ that supplies oxygen, essential nutrients, hormones, and metabolites to the developing fetus. The placenta is of fetal lineage, stemming from the same pool of early embryonic blastomere cells that form the embryo, and shares its sex (3). During early development, the trophectoderm and inner cell mass separate into distinct lineages, and the placenta develops alongside the embryo after implantation. In mice, placental remodeling occurs as the chorion fuses with the allantois, followed by trophoblast differentiation and labyrinth structure formation (3). This remodeling process typically ends by mid-gestation, at E10.5 in mice and E18 in humans (3, 4). Proper placental development and function are crucial for the progression of pregnancy, and abnormalities lead to a range of adverse pregnancy outcomes, including embryonic growth delays, intrauterine growth restriction, preeclampsia, and preterm birth (5, 6). Interestingly, a recent study has revealed that prenatal alcohol exposure is linked to alterations in human placental gene expression (7). This underscores the potential

role of this tissue in exploring and understanding the repercussions of alcohol on embryonic development.

The preimplantation period is a critical phase in embryonic development, in which the parental DNA methylation marks are erased and the epigenome is remodeled to initiate the developmental program (8). Since pregnancy is typically detectable only after implantation, when human chorionic gonadotropin hormone production begins, there is a 6-day window of high risk for PAE during the preimplantation development stage (9). Given the increasing number of women of 18–34 years old consuming alcohol, including binge drinking (10), and the substantial number of unintended pregnancies worldwide (approximately 40%) (11), it is essential to investigate the impact of alcohol during this critical phase. Despite its importance for proper embryonic development and its heightened susceptibility to adverse environmental exposures, the preimplantation period has been understudied in relation to PAE, with only a few models focusing exclusively on this early developmental timepoint (12-15).

DNA methylation, established by DNMT3A/DNMT3B and maintained by DNMT1, is a stable modification of the DNA consisting in the covalent addition of a methyl group to cytosine (16-18). During preimplantation period, DNA methylation undergoes a wave of reprogramming with a loss and a re-establishment (16). Importantly, DNA methylation seems to act like a sensor to exposure to environmental contaminants and/or infections (19-21). In our recent study using a preclinical mouse model of early preimplantation binge-like alcohol exposure, we reported sex-specific DNA methylation changes in the forebrains of embryos at mid-gestation (E10.5) (14). We also observed an increased incidence of morphological defects, including growth delays, in embryos exposed to ethanol at E10.5 and E18.5 (14, 15). These disruptions could be potentially imputed to changes in DNA methylation at imprinting control regions (ICRs) which are associated with fetal growth restriction (22-24). Despite all these evidence, a significant knowledge gap remains regarding the effects of early preimplantation

95 alcohol exposure on the placenta. The aim of the study was to define impacts of early preimplantation

96 prenatal alcohol exposure on the late-gestation placenta using our preclinical mouse model.

METHODS

Mouse model of preimplantation alcohol exposure

Eight-week-old female and male C57BL/6 mice (Charles River Laboratories) were mated, and pregnancy

with E0.5 embryos was confirmed by the presence of a visible copulatory plug the following morning.

Pregnant females were separated from the males and housed together ($n=2–3$) in a 12-h light/dark cycle

with unrestricted access to food and water. Like in our previously published preimplantation binge-like

PAE model (14, 15), pregnant females were subcutaneously injected with two doses of either 2.5 g/kg of

50% ethanol or an equivalent volume of saline at 2-h intervals on E2.5 (the 8-cell stage), then housed

under similar conditions with limited handling during gestation.

Tissue collection and analysis

At E18.5, pregnant females were euthanized and embryos and placentas were collected. The maternal

tissue layer was then removed from the placenta. The embryos and placentas were weighed, and the

placental area was calculated using a Leica stereo microscope and Leica LasX software (v1.4.4).

Placentas were flash frozen in liquid nitrogen and stored at -80°C for subsequent DNA and RNA

extraction. The sex of each embryo and its placenta was determined by qPCR for *Ddx3x* and *Ddx3y*

(LightCycler 96, Roche) (14, 15).

Histological analysis

114 On E18.5, placentas were collected with the maternal decidua to preserve structural integrity. Placentas

115 were fixed in 4% paraformaldehyde, post-fixed in 70% ethanol for 48 h, and then divided into two

116 identical sections for paraffin embedding. Cross-sections (5 µm thick) were stained with hematoxylin

117   and eosin and imaged on a Zeiss Zen Axioscan Slide Scanner. Image processing and junctional and

118   labyrinth zone measurements were performed with ImageJ (v1.52).

DNA/RNA extraction and library preparation

We randomly selected 6 control placentas (1 male and 1 female per litter from 3 different litters) and 10

ethanol-exposed placentas (1 male and 1 female per litter from 5 different litters) with normal

morphometric measurements and no visible morphological defects. The corresponding embryos also

developed normally. Whole placentas were ground in liquid nitrogen and equivalent amounts of each

sample were used for DNA and RNA extraction using QIAamp DNA Micro (Qiagen #56304) and

RNeasy Mini (Qiagen #74004) kits, respectively, following the manufacturer's recommendations. DNA

and RNA were quantified on a QuBit Fluorimeter using Broad-Range DNA assay (ThermoFisher

#Q32853) and High Sensitivity RNA assay (ThermoFisher #Q32852) kits, respectively.

DNA (1 µg) was used to produce Methyl-Seq libraries using a SureSelectXT Methyl-Seq Target

Enrichment System with a mouse enrichment panel (Agilent #G9651B and #5191-6704) following the

manufacturer's recommendations. After final amplification/indexation, libraries were quantified using a

QuBit fluorimeter with a High Sensitivity DNA assay kit (ThermoFisher #Q32854) and quality was

assessed using a BioAnalyzer before paired-end sequencing on a NovaSeq6000 S4 at the Genome

Quebec core facility. We obtained 112–139 M reads per sequenced library.

High-quality RNA (500 ng) was used to produce mRNA-Seq libraries using NEBNext mRNA

stranded library kits at the Genome Quebec core facility, followed by paired-end sequencing on a

NovaSeq6000 S4. We obtained 26–43 M reads per sequenced library.

Bioinformatics analysis

Methyl-Seq results were analyzed using the GenPipes Methyl-Seq pipeline (v3.3.0) (25). Reads

were aligned to the mouse reference genome (mm10) and methylation calls were obtained with Bismark

(v0.18.1) (26). The R package MethylKit (version 1.8.1) (27) was used to obtain differentially methylated

regions (DMRs) based on the Benjamini–Hochberg false discovery rate procedure. We used fixed

parameters, such as 100 base pair (bp) stepwise tiling windows and a threshold of $q < 0.01$ and $\geq 10\%$

difference of methylation between conditions. DNA methylation levels were calculated as the average

methylation of all CpGs within a tile for all the samples within a condition (minimum 3

samples/condition/sex at $\geq 10X$ sequencing depth) (14, 28). The bisulfite conversion rate ($> 96\%$) and

number of CpGs per tile were obtained using a custom Perl script (14).

Annotation of all analyzed tiles was conducted using Homer (version 4.10.1) with the mouse

mm10 reference genome (29). Functional enrichment analyses were performed with Metascape (30)

using differentially methylated tiles located in genes as an input. We obtained CpG island coordinates of

the mm10 genome from the UCSC Genome Browser (31), then delineated CpG shores and shelves by

extending 0–2 and 2–4 kb from them, respectively, as described previously (14, 28).

For mRNA-Seq data, post-sequencing bioinformatics analyses were performed on the mouse

reference genome (mm10) using the GenPipes RNA-Seq pipeline (v4.1.2) (25), including tools such as

Trimmomatic (v0.36) (32), STAR (v2.5.3) (33), Picard (v2.9.0), and BWA (v0.7.12) (34). DEGs were

obtained using the R package DESeq2 (v1.24.0) (35) including multiple testing correction using the

Benjamini-Hochberg false discovery rate procedure. Genes with p-values < 0.05 were considered

significant.

To establish a molecular signature of early preimplantation PAE, we used L1-regularized linear

regression (LASSO) (36, 37) with cross-validation for the gene expression and DNA methylation

profiles. The curse of dimensionality (*i.e.*, challenges stemming from training models on datasets with a

large number of features relative to samples) was addressed using the Synthetic Minority Over-sampling

Technique to counteract sample imbalances (38). A random forest classifier was trained on these

balanced datasets to emphasize the significance of the selected DEGs and DMRs in the classification.

The specific regions used to establish the signature were among the top ranked based on feature

importance. The efficacy of the identified DEGs or DMRs to differentiate the two groups was validated

by two-sample *t*-test and Wilcoxon rank-sum test, while ROC curves were used to assess the

discriminatory capacity of each DEGs or DMRs (37).

Statistics

Statistical analyses not included in bioinformatic pipelines were performed using R (version 3.5.0) or

GraphPad Prism (version 9.5.0). Statistical significance was calculated using unpaired *t*-tests with

Welch's correction and an *F* test for variance or two-proportion *z*-test. All data represent the mean ±

standard deviation (SD). *p*-values < 0.05 were considered significant.

RESULTS

Early preimplantation alcohol exposure does not induce morphometric changes in the placenta but

induces lower embryo weight in males

In this study, we examined the effect of early preimplantation binge alcohol exposure on late placental

development using an established FASD mouse model (14, 15) . Briefly, to simulate short-term exposure

to a high level of alcohol, pregnant females were given two doses (2.5 g/kg) of ethanol at 2-hour intervals

during preimplantation (at the 8-cell stage (E2.5)). We previously showed that this increases the average

blood alcohol level to 158.31 mg/dL (14). At late gestation (E18.5), we collected the embryos and

placentas to assess the overall impacts of alcohol on gestational outcomes, focusing first on basic

morphometric measurements. All measurements were compared within the same sex.

The proportion of male embryos is unchanged between control (47/86 – 55%) and ethanol-

exposed group (62/101 – 61%) (Fig. 1A). Additionally, there were no significant effects of early

preimplantation alcohol exposure on litter size, the number of males and females per litter (Fig. 1B), or

the placental area and weight (Fig. 1C, 1D). Weight of ethanol-exposed male embryos is significantly

decreased compared to controls (1.134g ± 0.067 vs 1.195g ± 0.102, respectively; $p<0.01$), while no

difference was observed in females (Fig. 1E). Despite the reduced weights of male embryos, the placental

efficiency (*i.e.*, the embryo-to-placental weight ratio) was similar between control and ethanol-exposed

embryos for both sexes (Fig. 1F).

When investigating the impact of early alcohol exposure on the overall composition of placental

cells, we observed no significant differences in the sizes or thicknesses of the junctional and labyrinth

zones between the groups. Additionally, the ratios of their respective areas/thicknesses to the total

area/thickness showed no significant differences (Fig. S1). However, in female ethanol-exposed

placentas, the junctional zone thickness, and the ratio of the junctional zone area to the total area

presented significantly increased variance compared to control ($p<0.05$) (Fig. S1).

Overall, these findings suggest that early preimplantation alcohol exposure does not significantly

impact placental morphology during late gestation but does have a developmental effect on male

embryos.

Sex-specific DNA methylation dysregulation in late-gestation placentas following early

preimplantation alcohol exposure

To examine the impact of preimplantation alcohol exposure on the DNA methylation patterns of late-

gestation placentas, we performed genome-wide DNA methylation analyses on control and ethanol-

exposed placentas. After establishing sex-specific DNA methylation profiles in E18.5 placentas, we

compared the average DNA methylation levels between control and ethanol-exposed male or female

placentas, and identified 751,083 tiles (100bp) with sufficient coverage in both data sets, allowing us to

investigate shared and sex-specific DNA methylation differences induced by early preimplantation alcohol exposure (Fig. S2A).

Strikingly, we identified 991 and 1309 differentially methylated regions (DMRs; ≥10% difference between conditions) in response to early preimplantation alcohol exposure that were exclusive to males and females (Fig. 2A, 2C; Table 1 and S1), respectively, while only 21 DMRs were shared between the sexes (Fig. S2A; Table 2 and S1). Male placentas had nearly equal distributions of DMRs, displaying both increased (540 – 54%) and decreased (451 – 46%) methylation after ethanol exposure (Fig. 2A). In contrast, DMRs in female placentas were predominantly increased with ethanol exposure (988 – 75% vs. 321 – 25%) (Fig. 2C). Most DMRs were annotated as introns (male, 41%; female, 42%) or intergenic regions (male, 39%; female, 37%) (Fig. 2B, 2D). We observed larger DNA methylation average changes between control and ethanol-exposed placentas in DMRs located in non-coding regions (controls 44% vs ethanol-exposed 51%) and in exons (controls 47% vs ethanol-exposed 52%) in males, while in females, larger average changes were observed for DMRs located in 3′ UTRs (controls 53% vs ethanol-exposed 63%) and introns (controls 45% vs ethanol-exposed 54%) (Fig. 2B, 2D).

A detailed examination of the distributions of sex-specific DMRs across various CpG methylation levels revealed a consistent pattern between the sexes. DNA methylation tends to increase in genomic regions with higher basal methylation levels (50-100%), while it decreases in genomic regions with lower basal methylation levels (0-50%) (Fig. S3A, S3C). This trend was especially pronounced for female-specific DMRs (Fig. S3C). Although some sex-specific DMRs displayed methylation changes exceeding 20%, the majority exhibited changes in the range of 10 to 15% (Fig. S3B, S3D).

To identify pathways associated with sex-specific DMRs in ethanol-exposed placentas, we performed gene annotation enrichment analyses. First, we removed DMRs associated with intergenic regions. For males, the remaining 607 DMRs represented 542 unique genes. Enriched pathways included cell junction assembly (*Cttn*, *Mapt*), response to growth factors (*Fgf1*, *Zeb2*), negative regulation of

synaptic transmission (*Cttn*, *Mapt*), organic hydroxy compound biosynthesis, (*Slc11a2*, *Star*), and

phenol-containing compound biosynthesis (*Hdc*, *Nr4a2*) (Fig. 2E,2F). For females, the remaining 821

DMRs corresponded to 749 unique genes. The top enriched pathways differed from in males, including

plasma membrane-bounded cell projection morphogenesis (*Ank3*, *Cdh11*), regulation of inflammatory

responses (*Bcl6*, *Foxj1*), alpha-beta T cell differentiation (*Bcl6*, *Foxj1*), negative regulation of cell

migration (*Rbp4*) and negative regulation of cell population proliferation (*Foxj1*, *Wnt5a*) (Fig. 2G,2H).

These findings reveal that early preimplantation alcohol exposure results in sex-specific DNA

methylation dysregulation in late-gestation placentas, with females showing a higher susceptibility.

Furthermore, these sex-specific alterations in DNA methylation are linked to specific biological

pathways.

Female ethanol-exposed placentas tend to lose methylation in CpG-rich regions

To determine whether the local genomic sequence contributes to the inherent sex-specific susceptibility

of DMRs, we examined their CpG contexts in relation to CpG islands (CGIs). Relative to control, most

male and female ethanol-exposed placentas showing decreased and increased DMRs were located in

open sea regions, far from CGIs (Fig. 3A). Regarding the DMRs situated in CGIs, in males, we

predominantly observed DMRs with a gain of methylation (68 increased vs. 11 decreased). Conversely,

in females, a larger proportion of DMRs with methylation loss was observed (45 decreased vs. 37

increased) (Fig. 3A). For both sexes, the largest changes in methylation levels occurred in open sea

regions and shores, and the smallest changes mainly occurred in CGIs (Fig. 3B).

For a more comprehensive assessment of methylation differences in sex-specific DMRs based on

their CpG contexts, we compared the methylation profiles of DMRs in regions of different CpG density,

notably CGIs, promoters, and repeat regions (Fig. 3C, 3D). Generally, male DMRs in CGIs had higher

methylation in ethanol-exposed placentas than in controls. This difference lessened with decreasing CpG

density, consistent with the fact that DMRs in repeat regions displayed both decreased and increased

methylation levels in male ethanol-exposed placentas compared to controls (Fig. 3C). Interestingly, we

observed the opposite trend when examining the methylation profiles of female DMRs across different

annotations. While we observed generally lower methylation levels for DMRs in CGIs in ethanol-

exposed placentas, the methylation levels of DMRs in promoters and particularly in repeat regions

increased in female ethanol-exposed placentas compared to controls (Fig. 3D).

Taken together, our findings suggest that early preimplantation alcohol exposure differentially

impacts the placental epigenome depending on sex and CpG density.

DMRs shared by male and female late-gestation placentas after early preimplantation alcohol

exposure are related to growth and development

Although our primary focus was to identify sex-specific DMRs in the placenta resulting from early

preimplantation alcohol exposure, we also observed shared affected regions between sexes. Of the 21

shared DMRs (Table 2), 17 (80%) DMRs increased or decreased consistently in both sexes, while 4

(20%) displayed opposing changes (Fig. S4A). Regions of interest included DMRs located in genes such

as *Galp* (neuropeptide signaling), *Psd3* (protein transduction), or *Wwox* (apoptotic signaling) (Fig. S4B).

A total of 12 (57%) DMRs were located in promoters or intragenic regions. Pathway enrichment analysis

of these regions revealed enrichment in growth regulators (Fig. S4C, S4D).

For a broader understanding of how preimplantation alcohol exposure alters the DNA methylation

patterns of specific genes in both male and female placentas, we performed DMRs comparisons based

on gene name annotations instead of specific genomic tile-positions. After excluding intergenic DMRs,

we compared a total of 607 male-specific DMRs (542 genes) to 821 female-specific DMRs (749 genes)

located in promoters and intragenic regions. Along with the 21 shared DMRs previously described, we

uncovered 69 shared genes across 89 DMRs with DNA methylation differences in male or female

ethanol-exposed placentas (Fig. S5A). These included methylation changes that were consistent (39) and

opposing (50) between the sexes (Fig. S5A, S5B). Pathway enrichment analysis revealed pathways

related to the reproductive process (*Bmp6*, *Prdm14*, *Runx1*), cell morphogenesis (*Prdm14*, *Ptprk*),

transmembrane transport activity, myeloid cell development, and response to growth factors (*Bmp6*,

Prdm14, *Ptprk*, *Runx1*), (Fig. S5C, S5D).

These findings indicate that the dysregulated pathways shared between male and female placentas

in response to preimplantation alcohol exposure could contribute to pathogenicity of diverse symptoms

including growth defects.

Early preimplantation alcohol exposure leads to sex-specific gene expression changes in late-

gestation placentas

To further our understanding of the molecular impact of preimplantation alcohol exposure on the late-

gestation placenta, we performed gene expression profiling using mRNA-Seq of the same set of samples

used for DNA methylation analysis. We used a significance threshold of $p<0.05$ to identify differentially

expressed genes (DEGs) between ethanol-exposed and control placentas. We identified 1046 male-

specific DEGs, 340 female-specific DEGs, and 54 shared DEGs (Fig. 4A-D; Fig. S6A; Table 3 and 4).

In males, 498 genes (48%) showed decreased expression in ethanol-exposed placentas, while 548 genes

(52%) showed increased expression (Fig. 4A, 4C). Of the 340 female-specific DEGs in ethanol-exposed

placentas, 185 (54%) and 155 (46%) showed decreased and increased expression relative to the control,

respectively (Fig. 4B, 4D).

Functional enrichment analyses of male- and female-specific DEGs revealed divergent pathways.

For instance, male DEGs were related to cell migration (*Aqp1*, *Rhoc*), cytoskeletal organization (*Apoa1*,

Bmp10), cell-substrate adhesion (*Adam8*, *Utrn*), cell junction organization (*Ank6*, *Cdh6*), and blood

vessel development (*Flt4*, *Slc16a1*) (Fig. 4E); whereas the female DEGs showed enrichment in pathways

involved in cell adhesion (*Itgb3*, *Ret*), serotonin uptake (*Slc22a2*, *Snca*), PIWI-interacting RNA metabolism (*Fkbp6*, *Piwil2*), megakaryocyte development and platelet production (*Gata2*, *Gata4*), and locomotory behavior (*Chrna4*, *Efnb3*) (Fig. 4F). Examples of male- and female-specific DEGs are shown in Figure 4G and 4H.

Shared gene expression changes in late-gestation placentas after early preimplantation alcohol exposure

We also identified 54 genes (Table 4) that showed similar differential expression in both male and female ethanol-exposed placentas compared to control following early preimplantation alcohol exposure. For 93% of genes (50/54), we observed comparable changes between the sexes (Fig. S6A-C); however, 4 genes showed divergent changes in males and females, including *Ifit2* (response to interferon-alpha) and *Krt14* (cytoskeleton organization) (Fig. S6D). For *Ifit2*, the control expression level also differed between males and females (Fig. S6D).

Pathway enrichment analysis of these shared DEGs revealed enrichment in neuromuscular junction development (*Musk*, *Pmp22*), sphingolipid biosynthesis (*B3gnt5*, *St8sia5*), artery development (*Chd7*, *Luzp1*), and matrix organization (*Foxf2*, *Kif9*) (Fig. S6E). Additional interesting shared DEGs included *Bhlhe41* (associated with muscle development) and *Cited2* (involved in decidualization, placenta, cardiac, and cortex development), which both showed decreased expression in ethanol-exposed placentas, and *Scel* (which regulates Wnt signaling) and *Syt14* (involved in synaptic transmission), which exhibited increased expression (Fig. S6F). These specific DEGs highlight potential molecular mechanisms and pathways affected by preimplantation alcohol exposure in the placenta regardless of sex.

322   These findings provide insights into the biological processes that may be impacted by

323   preimplantation alcohol exposure in both male and female placentas, suggesting potential negative

324   consequences of placental dysfunction on embryonic development following early PAE (14).

325   Early preimplantation alcohol exposure does not strictly impact genes crucial for placental

326   development or placental imprinting genes at late gestation

To evaluate the impact of early preimplantation alcohol exposure on genes crucial for placental

development, we analyzed our DNA methylation and gene expression datasets using previously

published curated gene lists (39).

Of the 205 curated placental genes, 193 were present in our DNA methylation dataset, accounting

for 7327 tiles located in promoters and intragenic regions (Fig. S7A). However, only six contained

DMRs: three in males (*Epas1*, *Id1*, and *Peg10*) and three in females (*Flt1*, *Ppard*, and *Serpine1*). All 205

curated placental genes were included in our mRNA-Seq dataset, of which 19 and 8 were dysregulated

following ethanol exposure in males and females, respectively (Fig. S7B). Only one DEG, *Cited2*, was

changed in both male and female placentas (Fig. S6B, S6C, and S6F and Fig. S7B). Additionally, *Peg10*'s

intronic DMR, which overlaps with the *Peg10* imprinting control region (ICR), exhibited increased DNA

methylation and increased expression in male ethanol-exposed placentas (Fig. S7A-B).

To further investigate the impact of preimplantation alcohol exposure on imprinted genes, we

searched our datasets for a curated list of 105 genes known to be regulated by genomic imprinting in the

placenta (40-45). In male ethanol-exposed placentas, we identified 17 DMRs located in the intragenic

regions of 11 genes (Fig. S8A, S8C). Of these, 12 DMRs in 7 genes were located directly in the ICR.

Furthermore, we observed 18 imprinted DEGs in male placentas (Fig. S8B). In contrast, female ethanol-

exposed placentas showed fewer DNA methylation changes on imprinting genes, with only 9 DMRs in

9 different genes. Of these, only *Slc22a2* showed differential expression (Fig. S8A, S8B, Table S2 and

S3). Detailed examples of DMRs (*Airn*, *Commd1*, *Grb10* and *Peg10*) and DEGs (*Grb10*, *Igf2r*, *Peg10*,

Zrsr1 and *Slc22a2*) for male and female placentas are shown in Figure S8C and S8D.

These results reveal that although several placental and imprinted genes exhibited dysregulated

DNA methylation levels or expression following early preimplantation alcohol exposure, the

dysregulation was not widespread. These findings align with our morphometric analyses, which did not

reveal any significant placental malformations.

Sex-specific regions with both DNA methylation and gene expression changes in the late-gestation

following early preimplantation alcohol exposure

To explore the relationship between dysregulated DNA methylation and gene expression profiles

in the placenta after preimplantation alcohol exposure, we next compared our lists of promoter and

intragenic DMRs and DEGs by examining the expression profiles of the genes associated with 607 male

DMRs (542 unique genes), 821 female DMRs (749 unique genes), and 12 shared DMRs (11 unique

genes).

Of the 11 genes with shared DMRs, only *Clstn3* and *Tro* exhibited gene expression changes in

male ethanol-exposed placentas (Fig. S9A-B), and no expression changes were observed in females. In

males, 40 unique genes containing 46 DMRs displayed both DNA methylation and gene expression

changes in ethanol-exposed placentas (Fig. 5A). The majority of these DMRs (36; 78%) were associated

to small but significant gene expression changes (Fig. S10A). Notably, altered regions/genes included

Apoc1 (lipidic activity), *Carmil1* (implicated in actin filament processes), *Slc16a3* (encodes the MCT4

protein, important for transmembrane transport, placental organization, and vascularization) and *Zfhx3*

(a transcription factor important for development) (Fig. 5B).

In females, 15 unique genes containing 16 DMRs showed both differential methylation and gene

expression changes in ethanol-exposed placentas (Fig. 5C). Eleven of these DMRs (69%), associated to

7 genes (47%), revealed changed expression by (Fig. S10B). Noteworthy genes included *Chd7*

(implicated in cardiac, muscle, and neuronal development), *Cux2* (related to neuronal pathways and

functions, including cortical interneuron development), *Elovl2* (involved in fatty acid processes), and

Mfsd2a (implicated in the establishment of the blood-brain barrier) (Fig. 5D).

These results offer significant insights into the enduring effects of preimplantation alcohol

exposure on multiple genes, shedding light on the potential interplay between gene expression and DNA

methylation. The findings underscore the importance of understanding sex-specific DNA methylation

dysregulation as a potential mechanism shaping the distinctive gene expression patterns observed in both

male and female placentas.

Unveiling predictive molecular signatures in the placenta for early preimplantation alcohol

exposure

Next, we wanted to determine which molecular signature, either DNA methylation or gene expression

profiles, proves superior in predicting early preimplantation alcohol exposure in the mouse placenta. We

used a random forest classifier and least absolute shrinkage and selection operator (LASSO) regression,

a machine-learning technique, to create a predictive model using identifiable markers found in our

transcriptomic or DNA methylation data. To enhance accuracy and effectiveness, we trained the model

using both male and female samples together, increasing our statistical strength and refining the

analytical process.

First, we used gene expression profiles and performed LASSO regression to establish specific

transcriptomic signatures for ethanol-exposed placentas. Despite being statistically significant, the 20

DEGs identified were not different enough between the conditions to establish a good transcriptomic

signature of preimplantation alcohol exposure in late-gestation placentas, with small Euclidean distances

between samples of the two conditions and poor statistical coefficient of importance (Fig. S11A-C).

Interestingly, LASSO regression using our DNA methylation profiles identified a set of 24

statistically significant DMRs that formed a specific signature enabling the effective discrimination of

ethanol-exposed samples from controls for both principal component analysis and Euclidean distance

calculations (Fig. 6A-B). These DMRs were both intragenic (15) and intergenic (9), and each had

different levels of importance based on their statistical parameters, notably the LASSO coefficient,

Wilcoxon *p*-value, and AUC generated by LASSO analysis (Fig. 6C). The most significant DMR in the

signature was located in an intergenic region close to *Nup35*. Additionally, there were DMRs in genes

such as *B3galt1*, *Kcng1*, *Mir3960*, *Mir6940*, *Psd3*, and *Sesn1*, along with intergenic DMRs close to

Bicd1, *Gatsl2*, and *Zeb2* (Fig. 6C-D).

These findings unveil a distinctive molecular signature in placental tissue, highlighting a reliable

DNA methylation pattern that serves as a robust molecular tool for detecting early embryonic alcohol

exposure at birth.

DISCUSSION

In this study, we examined the molecular and developmental effects of PAE during early preimplantation

on the late-gestation placenta. Our analyses are the first to reveal extensive sex-specific alterations in

both DNA methylation and gene expression patterns in late-gestation mouse placentas exposed to

ethanol. Interestingly, despite these molecular changes, the overall morphological development of the

placenta appeared unaffected. Finally, employing machine learning, we demonstrate that DNA

methylation profiles in the placenta emerge as more robust indicators of early alcohol exposure compared

to gene expression signatures.

Sex-specific impacts of preimplantation alcohol exposure on embryo growth and related pathways

Prenatal alcohol exposure was previously shown to alter embryonic development in different contexts,

by inducing fetal growth restriction or growth delay (14, 15). With our model, we found that the impact

on growth and related pathways was more pronounced in male ethanol-exposed placentas, consistent with the observed male-specific changes in genes regulating growth, cytoskeletal organization, placental development, and vascularization. Furthermore, we noted a reduction in embryo weight only in male, indicating that males may be more susceptible to growth delays following preimplantation alcohol exposure. However, it is worth noting that in our previous studies using this model, we observed similar percentages of growth-delayed embryos (~5%) between males and females at mid-gestation (14). It is possible that some female embryos with growth delays at mid-gestation may catch up later in development, eventually achieving a similar embryo weight to control. Several studies have demonstrated that female embryos generally exhibit greater adaptability to adverse *in utero* environments than male embryos, facilitating their development even in the presence of altered resource availability (46-48). Male ethanol-exposed placentas displayed altered methylation and/or expression patterns in several genes that are directly linked to embryonic growth restriction. For example, a study recently showed that injecting *fibroblast growth factor* 2 (*Fgf2;* downregulated in male ethanol-exposed placentas) into a rat model of preeclampsia improved clinical outcomes, including attenuating fetal growth restriction (49). Mutation of *Fgf1* (increased methylation in male ethanol-exposed placentas) has been linked to preeclampsia in humans (50), while *Bmp6* (decreased methylation in male ethanol-exposed placentas) was downregulated in the fetal kidneys of rats with intrauterine growth restriction in a model of induced hypertension due to chronic insulin administration (51). Changes in these genes may significantly contribute to the inability of male embryos to develop normally and achieve the embryo weights of control embryos.

Distinct patterns of DNA methylation dysregulation in female and male ethanol-exposed placentas

DNA methylation analysis of late-gestation placentas following preimplantation alcohol exposure uncovered sex-specific DMRs. Moreover, we also observed sex-specific alteration patterns in DNA methylation changes as well as in the proximity of the sex-specific DMRs to CpG-rich regions,

suggesting a sex-specific effect on DNA methylation based on CpG content following preimplantation

alcohol exposure.

In a recent study, we discovered genome regions with sex-biased DNA methylation profiles in

control placentas at E18.5 (39). When we compared the DMRs observed following preimplantation

alcohol exposure with these sex-biased regions, we found that 115 male DMRs and 188 female DMRs

were also sex-biased at late gestation. Importantly, these ethanol-exposed DMRs and sex-biased regions

exhibited distinct CpG distributions based on sex. Notably, we observed a higher proportion of female-

specific DMRs located in sex-biased regions in CpG islands and shores than male-specific DMRs located

in sex-biased regions (39). These differentially methylated regions between male and female embryos

could help explain the sex-specific changes observed in both DNA methylation and gene expression. It

is known that sex-biased differences exist during early development, and that divergent DNA

methylation profiles can contribute to sex-specific DMRs in late-gestation placentas. One hypothesis is

that these specific regions are more vulnerable to harmful exposures from the maternal environment due

to their sex-specificity during development. Epigenetic dysregulation in these specific regions may

progress differently in males and females, resulting in sex-specific DMRs in the late-gestation following

preimplantation alcohol exposure.

Further studies investigating dysregulated DNA methylation in the trophectoderm and placenta

during early development could provide insights into the variability of these sex-specific regions after

preimplantation PAE. Understanding the dynamics and mechanisms underlying these sex-specific effects

will contribute to our knowledge of how PAE impacts placental function and subsequent developmental

outcomes in a sex-dependent manner.

Preimplantation alcohol exposure had a greater impact on imprinted and placental development genes in male placentas

The methylation patterns of the ICRs of imprinted genes and the DNA methylation and expression patterns of genes crucial for placental development can significantly influence proper embryo development (52, 53). While placental development being largely completed by late gestation, epigenetic and transcriptomic dysregulation of key genes can disrupt its function, especially when the dysregulation occurs at earlier developmental stages (54). Although histological analysis of the junctional and labyrinth zones showed comparable results between ethanol-exposed and control placentas in both sexes, we observed more changes in imprinted and placental development genes in male ethanol-exposed placentas compared to females.

Among the dysregulated genes, we observed differential methylation and expression of the imprinted gene *Grb10* in male ethanol-exposed placentas. *Grb10* encodes a growth restrictor, and irregular imprinting, especially in its ICR, can directly affect embryonic development (52). The increased *Grb10* expression observed in male ethanol-exposed placentas could therefore directly impact embryonic growth and development. Additionally, we observed dysregulation in both DNA methylation and gene expression in the *H19/Igf2* cluster, including increased expression of *Igf2r* in male ethanol-exposed placentas and increased methylation in the ICR of *Airn* (antisense of *Igf2r*). Like *Grb10*, *Igf2r* negatively regulates embryonic growth and development (52, 55). This imprinted cluster has been previously identified using a similar paradigm of alcohol exposure during preimplantation. In their study, Haycock and Ramsay showed decreased DNA methylation in the ICR of *H19* in the mid-gestation (12). They also observed significant decreases in the weights of both ethanol-exposed embryos and placentas at mid-gestation and found a strong correlation between DNA methylation of the paternal *H19* allele and weight loss in the ethanol-exposed group (12). The abnormal methylation and expression profiles of imprinted

genes collectively impact proper embryonic development, even though placental morphology is not affected.

Analysis of placental development genes also revealed more genes with altered expression profiles in male ethanol-exposed placentas. Notably, we observed increased expression of *Dnmt3l*, a coactivator of *Dnmt3a* and *Dnmt3b*, which catalyze *de novo* DNA methylation (56). During placental development, *Dnmt3l* is essential for establishing proper maternal gene imprinting profiles and is highly expressed in trophoblasts (56-58). While the impact of *Dnmt3l* overexpression on the placenta is unknown, studies have demonstrated morphological defects and abnormal trophoblast patterns in *Dnmt3l* knockout mice, providing further evidence of its critical role in proper embryonic development (59, 60). We cannot exclude the potential impact of dysregulated *Dnmt3l* expression on the irregular DNA methylation and transcriptomic profiles observed in certain imprinted genes, such as the maternally methylated genes *Peg10*, *Jade1*, or *Zrsr1,* in male ethanol-exposed placentas.

Dysregulation of the placental epigenetic and transcriptomic landscape could play a role in proper embryonic brain development

The placenta-brain axis has recently received considerable attention, and it is now well-established that placental dysregulation can lead to adverse effects on brain development and function (48, 61). Neurodevelopmental disorders and intellectual disabilities are commonly observed in FASD, resulting from disruptions during brain development and incomplete maturation (1). While the impact of endocrine placental dysfunction on brain development is increasingly understood, the role of placental epigenetic and transcriptomic dysregulation has been less explored. Among the many enriched pathways observed in our datasets, two main pathways emerged that could directly affect normal embryonic brain development.

First, we found that male-specific DMRs were enriched for components of a pathway related to

synaptic transmission, with dysregulated genes including *Asic1*, *Cbln1*, and *Plk2*. *Asic1* and *Cbln1* are

required for normal neuronal development (62, 63), and *Plk2* has been linked to epilepsy because of its

role in synaptic potentiation (64). Additionally, our transcriptome analysis revealed that one of the main

pathways enriched in female ethanol-exposed placentas was related to serotonin metabolism. Serotonin

is a neurotransmitter that regulates embryonic neurodevelopment, and the placenta plays a

neuroendocrine role by producing and providing serotonin to the developing forebrain (65). Placental

insufficiency, including failing to provide an adequate serotonin supply to the embryo, is associated with

the development of mental health issues and reduced cognitive abilities (65). Knockout of *Itgb3*, which

was downregulated in female ethanol-exposed placentas, had adverse effects on the placenta's

vascularization structure in a sheep model, compromising nutrient transport (66). ITGB3 is also directly

involved in serotonin uptake by interacting with different serotonin transporters (67). Interestingly,

Slc22a2, a gene implicated in serotonin clearance has been identified as overexpressed in female ethanol-

exposed placentas (68). Together, changes in these two genes could negatively modulate serotonin uptake

in the developing brain *via* altered placental endocrine function and disrupted communication, further

emphasizing the impact of the placenta on embryonic health and development.

Dysregulated placental DNA methylation profiles serve as a robust signature of early

preimplantation alcohol exposure

Our study unequivocally demonstrates that alcohol exposure as early as the 8-cell stage induces enduring

molecular alterations in both DNA methylation and gene expression profiles in the mouse placenta.

However, we show that the stability and persistence of DNA methylation alterations across ethanol-

exposed placenta profiles underscore its potential superiority as a molecular signature for capturing and

reflecting the long-term impact of alcohol exposures on developmental processes. This further supports

the ability to employ DNA methylation patterns as a viable, reliable, and precise indicator of previous prenatal exposure.

In recent years, two research groups showed DNA methylation changes in buccal epithelial cell swabs from children with confirmed FASD (69-71). Moreover, one of these studies applied a machine learning approach with the identified epigenetic signature to train and further efficiently validate the identification of confirmed FASD cases (71). Although these cells are easily accessible and require no invasive procedure to obtain, their epigenome will vary with age and may generate some bias in the generation of accurate DNA methylation signatures. Moreover, the high renewal rate of these surface cells may not homogeneously reveal PAE-induced epigenetic modifications. PAE's heterogenous paradigms and their effects on the epigenomes of buccal epithelial cells are also an obstacle in establishing a universal DNA methylation signature. Tissues or cell types more directly exposed to alcohol during gestation may be more promising alternatives. In this perspective, the placenta could be an effective means to detect PAE at birth since its development occurs rapidly during gestation. A previous study using preconception and gestational cannabis also observed shared changes in DNA methylation in the placentas and brains of exposed embryos (72), indicating that the placenta may be able to effectively mirror some DNA methylation alterations present in the brain. The short lifespan of the placenta (resulting in minimal cell renewal) makes it an optimal candidate tissue to profile epigenetic alterations caused by PAE any time during gestation. The placenta can be thought of as an organ having a high potential to teach us more about what transpired during pregnancy and how it may have affected embryonic development.

As mentioned earlier, the absence of a molecular diagnostic tool for PAE and FASD results in most cases going undetected until children reach school age, where noticeable neurodevelopmental symptoms emerge (1). Although our study focused on mice, using DNA methylation profiles and machine-learning approaches to identify robust molecular signatures for early embryonic alcohol

exposure serves as a proof of concept that could be applied to humans, enabling early diagnosis and optimal support for newborns with FASD from their early days. Nevertheless, ethical concerns surrounding PAE diagnostic tests, including the risk of stigmatization and discrimination, challenges in obtaining informed consent, and issues related to accuracy, need careful consideration. Thus, there is a crucial need to balance these concerns with initiatives aimed at enhancing our understanding of the mechanisms and risks associated with alcohol consumption during pregnancy, addressing issues like lack of awareness and unintentional exposure. Additionally, there is a call for the development of clearer guidelines to assist women and healthcare professionals in making informed decisions about alcohol consumption, prioritizing the well-being of both the mother and the child.

CONCLUSION

In this study, we showed that early preimplantation alcohol exposure can alter the molecular profile of late-gestation mouse placentas. More precisely, our study provides valuable insights into the sex-specific epigenetic and transcriptomic consequences of early PAE on the placenta, contributing to a better understanding of its potential effects on placental function and embryo development. Furthermore, this study highlights for the first time that the placental DNA methylation profiles provide robust molecular signature for early PAE detection. Further studies based on our model will help decipher the mechanisms of sex-specific placental DNA methylation and gene expression changes on embryonic development and their long-term implications.

DECLARATIONS

Ethics approval

All animal work conducted in this study was approved by the CHU Ste-Justine Research Center *Comité*

Institutionnel de Bonnes Pratiques Animales en Recherche (CIBPAR) under the guidance of the

Canadian Council on Animal Care (CCAC).

Consent for publication

Not applicable

Availability of data and materials

All sequencing datasets generated and used in this study will be publicly available via the Gene

Expression Omnibus repository.

Competing interest

The authors have declared that no conflict of interest exists.

Author contributions

LML and SM designed the study. LML and MBL conducted experiments and contributed to data

acquisition. LML, ALA, AL, and FFB participated in data analysis. LML, TD, and SM wrote the

manuscript. All authors read and approved the final manuscript.

Funding

This work was supported by a research grant to SM from the Natural Sciences and Engineering Research

Council of Canada. LML and TD are supported by Canadian Institutes of Health Research scholarship.

MBL is supported by a scholarship/fellowship from Fonds de Recherche du Québec - Santé (FRQS).

FFB is supported by a Mitacs accelerate scholarship. ALA is supported by scholarships from Université

de Montréal and Réseau Québécois en Reproduction. SM is supported by an FRQS-Junior 2 salary award.

Acknowledgements

REFERENCES

1. Popova S, Charness ME, Burd L, Crawford A, Hoyme HE, Mukherjee RAS, et al. Fetal alcohol
spectrum disorders. Nature reviews Disease primers. 2023;9(1):11.
2. Lange S, Probst C, Gmel G, Rehm J, Burd L, Popova S. Global Prevalence of Fetal Alcohol
Spectrum Disorder Among Children and Youth: A Systematic Review and Meta-analysis. JAMA
pediatrics. 2017;171(10):948-56.
3. Hemberger M, Hanna CW, Dean W. Mechanisms of early placental development in mouse and
humans. Nature reviews Genetics. 2020;21(1):27-43.
4. Simmons DG, Natale DR, Begay V, Hughes M, Leutz A, Cross JC. Early patterning of the chorion
leads to the trilaminar trophoblast cell structure in the placental labyrinth. Development
(Cambridge, England). 2008;135(12):2083-91.
5. Wadhwa PD, Entringer S, Buss C, Lu MC. The contribution of maternal stress to preterm birth:
issues and considerations. Clinics in perinatology. 2011;38(3):351-84.
6. Coussons-Read ME. Effects of prenatal stress on pregnancy and human development: mechanisms
and pathways. Obstetric medicine. 2013;6(2):52-7.
7. Deyssenroth MA, Williams RP, Lesseur C, Jacobson SW, Jacobson JL, Cheng H, et al. Prenatal
alcohol exposure is associated with changes in placental gene co-expression networks. Scientific
reports. 2024;14(1):2687.
8. Canovas S, Ross PJ. Epigenetics in preimplantation mammalian development. Theriogenology.
2016;86(1):69-79.
9. Cole LA. New discoveries on the biology and detection of human chorionic gonadotropin.
Reproductive biology and endocrinology : RB&E. 2009;7:8.
10. Organization WH. Global status report on alcohol and health 2018: World Health Organization;
2019.
11. Sedgh G, Singh S, Hussain R. Intended and unintended pregnancies worldwide in 2012 and recent
trends. Studies in family planning. 2014;45(3):301-14.
12. Haycock PC, Ramsay M. Exposure of mouse embryos to ethanol during preimplantation
development: effect on DNA methylation in the h19 imprinting control region. Biology of
reproduction. 2009;81(4):618-27.
13. Legault LM, Bertrand-Lehouillier V, McGraw S. Pre-implantation alcohol exposure and
developmental programming of FASD: an epigenetic perspective. Biochemistry and cell biology =
Biochimie et biologie cellulaire. 2018;96(2):117-30.
14. Legault LM, Doiron K, Breton-Larrivée M, Langford-Avelar A, Lemieux A, Caron M, et al. Pre-
implantation alcohol exposure induces lasting sex-specific DNA methylation programming errors
in the developing forebrain. Clinical epigenetics. 2021;13(1):164.

15. Breton-Larrivée M, Elder E, Legault LM, Langford-Avelar A, MacFarlane AJ, McGraw S.
Mitigating the detrimental developmental impact of early fetal alcohol exposure using a maternal
methyl donor-enriched diet. FASEB journal : official publication of the Federation of American
Societies for Experimental Biology. 2023;37(4):e22829.
16. Shaffer B, McGraw S, Xiao SC, Chan D, Trasler J, Chaillet JR. The DNMT1 intrinsically
disordered domain regulates genomic methylation during development. Genetics.
2015;199(2):533-41.
17. McGraw S, Zhang JX, Farag M, Chan D, Caron M, Konermann C, et al. Transient DNMT1
suppression reveals hidden heritable marks in the genome. Nucleic acids research.
2015;43(3):1485-97.
18. McGraw S, Oakes CC, Martel J, Cirio MC, de Zeeuw P, Mak W, et al. Loss of DNMT1o disrupts
imprinted X chromosome inactivation and accentuates placental defects in females. PLoS genetics.
2013;9(11):e1003873.
19. Maurice C, Dalvai M, Lambrot R, Deschênes A, Scott-Boyer MP, McGraw S, et al. Early-Life
Exposure to Environmental Contaminants Perturbs the Sperm Epigenome and Induces Negative
Pregnancy Outcomes for Three Generations via the Paternal Lineage. Epigenomes. 2021;5(2).
20. Pierre WC, Legault LM, Londono I, McGraw S, Lodygensky GA. Alteration of the brain
methylation landscape following postnatal inflammatory injury in rat pups. FASEB journal :
official publication of the Federation of American Societies for Experimental Biology.
2020;34(1):432-45.
21. Breton-Larrivée M, Elder E, McGraw S. DNA methylation, environmental exposures and early
embryo development. Animal reproduction. 2019;16(3):465-74.
22. Koukoura O, Sifakis S, Soufla G, Zaravinos A, Apostolidou S, Jones A, et al. Loss of imprinting
and aberrant methylation of IGF2 in placentas from pregnancies complicated with fetal growth
restriction. International journal of molecular medicine. 2011;28(4):481-7.
23. Koukoura O, Sifakis S, Zaravinos A, Apostolidou S, Jones A, Hajiioannou J, et al.
Hypomethylation along with increased H19 expression in placentas from pregnancies complicated
with fetal growth restriction. Placenta. 2011;32(1):51-7.
24. Bourque DK, Avila L, Peñaherrera M, von Dadelszen P, Robinson WP. Decreased placental
methylation at the H19/IGF2 imprinting control region is associated with normotensive
intrauterine growth restriction but not preeclampsia. Placenta. 2010;31(3):197-202.
25. Bourgey M, Dali R, Eveleigh R, Chen KC, Letourneau L, Fillon J, et al. GenPipes: an open-source
framework for distributed and scalable genomic analyses. GigaScience. 2019;8(6).
26. Krueger F, Andrews SR. Bismark: a flexible aligner and methylation caller for Bisulfite-Seq
applications. Bioinformatics (Oxford, England). 2011;27(11):1571-2.
27. Akalin A, Kormaksson M, Li S, Garrett-Bakelman FE, Figueroa ME, Melnick A, et al. methylKit:
a comprehensive R package for the analysis of genome-wide DNA methylation profiles. Genome
biology. 2012;13(10):R87.
28. Legault LM, Doiron K, Lemieux A, Caron M, Chan D, Lopes FL, et al. Developmental genome-
wide DNA methylation asymmetry between mouse placenta and embryo. Epigenetics.
2020;15(8):800-15.
29. Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, et al. Simple combinations of lineage-
determining transcription factors prime cis-regulatory elements required for macrophage and B cell
identities. Molecular cell. 2010;38(4):576-89.
30. Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, et al. Metascape provides
a biologist-oriented resource for the analysis of systems-level datasets. Nature communications.
2019;10(1):1523.

31. Karolchik D, Hinrichs AS, Furey TS, Roskin KM, Sugnet CW, Haussler D, et al. The UCSC Table Browser data retrieval tool. Nucleic acids research. 2004;32(Database issue):D493-6.

32. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics (Oxford, England). 2014;30(15):2114-20.

33. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq aligner. Bioinformatics (Oxford, England). 2013;29(1):15-21.

34. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics (Oxford, England). 2009;25(14):1754-60.

35. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome biology. 2014;15(12):550.

36. Xu Y, Wang X, Huang Y, Ye D, Chi P. A LASSO-based survival prediction model for patients with synchronous colorectal carcinomas based on SEER. Translational cancer research. 2022;11(8):2795-809.

37. Sohn I, Kim J, Jung SH, Park C. Gradient lasso for Cox proportional hazards model. Bioinformatics (Oxford, England). 2009;25(14):1775-81.

38. Taft LM, Evans RS, Shyu CR, Egger MJ, Chawla N, Mitchell JA, et al. Countering imbalanced datasets to improve adverse drug event predictive models in labor and delivery. Journal of biomedical informatics. 2009;42(2):356-64.

39. Legault LM, Breton-Larrivée M, Langford-Avelar A, Lemieux A, McGraw S. Sex-based disparities in DNA methylation and gene expression in late-gestation mouse placentas. Biology of sex differences. 2024;15(1):2.

40. Coan PM, Burton GJ, Ferguson-Smith AC. Imprinted genes in the placenta--a review. Placenta. 2005;26 Suppl A:S10-20.

41. Proudhon C, Bourc'his D. Identification and resolution of artifacts in the interpretation of imprinted gene expression. Briefings in functional genomics. 2010;9(5-6):374-84.

42. Wang X, Soloway PD, Clark AG. A survey for novel imprinted genes in the mouse placenta by mRNA-seq. Genetics. 2011;189(1):109-22.

43. Lefebvre L. The placental imprintome and imprinted gene function in the trophoblast glycogen cell lineage. Reproductive biomedicine online. 2012;25(1):44-57.

44. Golding MC, Magri LS, Zhang L, Lalone SA, Higgins MJ, Mann MR. Depletion of Kcnq1ot1 non-coding RNA does not affect imprinting maintenance in stem cells. Development (Cambridge, England). 2011;138(17):3667-78.

45. Okae H, Hiura H, Nishida Y, Funayama R, Tanaka S, Chiba H, et al. Re-investigation and RNA sequencing-based identification of genes with placenta-specific imprinted expression. Human molecular genetics. 2012;21(3):548-58.

46. Christians JK, Ahmadzadeh-Seddeighi S, Bilal A, Bogdanovic A, Ho R, Leung EV, et al. Sex differences in the effects of prematurity and/or low birthweight on neurodevelopmental outcomes: systematic review and meta-analyses. Biology of sex differences. 2023;14(1):47.

47. Meakin AS, Cuffe JSM, Darby JRT, Morrison JL, Clifton VL. Let's Talk about Placental Sex, Baby: Understanding Mechanisms That Drive Female- and Male-Specific Fetal Growth and Developmental Outcomes. International journal of molecular sciences. 2021;22(12).

48. Bale TL. The placenta and neurodevelopment: sex differences in prenatal vulnerability. Dialogues in clinical neuroscience. 2016;18(4):459-64.

49. Martinez-Fierro ML, Garza-Veloz I, Castañeda-Lopez ME, Wasike D, Castruita-De la Rosa C, Rodriguez-Sanchez IP, et al. Evaluation of the Effect of the Fibroblast Growth Factor Type 2 (FGF-2) Administration on Placental Gene Expression in a Murine Model of Preeclampsia Induced by L-NAME. International journal of molecular sciences. 2022;23(17).

50. Marwa BA, Raguema N, Zitouni H, Feten HB, Olfa K, Elfeleh R, et al. FGF1 and FGF2 mutations
in preeclampsia and related features. Placenta. 2016;43:81-5.
51. Bursztyn M, Gross ML, Goltser-Dubner T, Koleganova N, Birman T, Smith Y, et al. Adult
hypertension in intrauterine growth-restricted offspring of hyperinsulinemic rats: evidence of
subtle renal damage. Hypertension (Dallas, Tex : 1979). 2006;48(4):717-23.
52. Plasschaert RN, Bartolomei MS. Genomic imprinting in development, growth, behavior and stem
cells. Development (Cambridge, England). 2014;141(9):1805-13.
53. Hemberger M, Cross JC. Genes governing placental development. Trends in endocrinology and
metabolism: TEM. 2001;12(4):162-8.
54. Apicella C, Ruano CSM, Méhats C, Miralles F, Vaiman D. The Role of Epigenetics in Placental
Development and the Etiology of Preeclampsia. International journal of molecular sciences.
2019;20(11).
55. Sleutels F, Tjon G, Ludwig T, Barlow DP. Imprinted silencing of Slc22a2 and Slc22a3 does not
need transcriptional overlap between Igf2r and Air. The EMBO journal. 2003;22(14):3696-704.
56. Bourc'his D, Xu GL, Lin CS, Bollman B, Bestor TH. Dnmt3L and the establishment of maternal
genomic imprints. Science (New York, NY). 2001;294(5551):2536-9.
57. Logan PC, Mitchell MD, Lobie PE. DNA methyltransferases and TETs in the regulation of
differentiation and invasiveness of extra-villous trophoblasts. Frontiers in genetics. 2013;4:265.
58. Joh K. Transcription-dependent DNA methylation at the imprinted Zrsr1-DMR.
bioRxiv. 2018:249524.
59. Arima T, Hata K, Tanaka S, Kusumi M, Li E, Kato K, et al. Loss of the maternal imprint in
Dnmt3Lmat-/- mice leads to a differentiation defect in the extraembryonic tissue. Developmental
biology. 2006;297(2):361-73.
60. Andrews S, Krueger C, Mellado-Lopez M, Hemberger M, Dean W, Perez-Garcia V, et al.
Mechanisms and function of de novo DNA methylation in placental development reveals an
essential role for DNMT3B. Nature communications. 2023;14(1):371.
61. Rosenfeld CS. The placenta-brain-axis. Journal of neuroscience research. 2021;99(1):271-83.
62. Lim J, Tai HH, Liao WH, Chu YC, Hao CM, Huang YC, et al. ASIC1a is required for neuronal
activation via low-intensity ultrasound stimulation in mouse brain. eLife. 2021;10.
63. McCormick LE, Gupton SL. Cerebellin-1 leads the way. PLoS biology. 2022;20(11):e3001880.
64. Seeburg DP, Sheng M. Activity-induced Polo-like kinase 2 is required for homeostatic plasticity of
hippocampal neurons during epileptiform activity. The Journal of neuroscience : the official
journal of the Society for Neuroscience. 2008;28(26):6583-91.
65. Rosenfeld CS. Placental serotonin signaling, pregnancy outcomes, and regulation of fetal brain
development†. Biology of reproduction. 2020;102(3):532-8.
66. Frank JW, Steinhauser CB, Wang X, Burghardt RC, Bazer FW, Johnson GA. Loss of ITGB3 in
ovine conceptuses decreases conceptus expression of NOS3 and SPP1: implications for the
developing placental vasculature†. Biology of reproduction. 2021;104(3):657-68.
67. Whyte A, Jessen T, Varney S, Carneiro AM. Serotonin transporter and integrin beta 3 genes
interact to modulate serotonin uptake in mouse brain. Neurochemistry international. 2014;73:122-
6.
68. Bacq A, Balasse L, Biala G, Guiard B, Gardier AM, Schinkel A, et al. Organic cation transporter 2
controls brain norepinephrine and serotonin clearance and antidepressant response. Molecular
psychiatry. 2012;17(9):926-39.
69. Laufer BI, Kapalanga J, Castellani CA, Diehl EJ, Yan L, Singh SM. Associative DNA methylation
changes in children with prenatal alcohol exposure. Epigenomics. 2015;7(8):1259-74.

70. Portales-Casamar E, Lussier AA, Jones MJ, MacIsaac JL, Edgar RD, Mah SM, et al. DNA
methylation signature of human fetal alcohol spectrum disorder. Epigenetics & chromatin.
2016;9:25.
71. Lussier AA, Morin AM, MacIsaac JL, Salmon J, Weinberg J, Reynolds JN, et al. DNA
methylation as a predictor of fetal alcohol spectrum disorder. Clinical epigenetics. 2018;10:5.
72. Laufer BI, Neier K, Valenzuela AE, Yasui DH, Schmidt RJ, Lein PJ, et al. Placenta and fetal brain
share a neurodevelopmental disorder DNA methylation profile in a mouse model of prenatal PCB
exposure. Cell reports. 2022;38(9):110442.

TABLES

Table 1: Top 10 sex-specific DMRs

Male-specific DMRs			
Effect of EtOH exposure	**Gene symbol**	**Methylation change (%)**	**Annotation**
Decreased	*(Hmox2)*	-33.13	Intergenic
Decreased	*Trappc9*	-31.79	Intron
Decreased	*Arsg*	-28.90	Intron
Decreased	*Mir148b*	-28.61	Intron
Decreased	*(Cplx2)*	-28.02	Intergenic
Decreased	*Mrpl45*	-27.29	3′ UTR
Decreased	*(Slc1a5)*	-26.62	Intergenic
Decreased	*(Cdh15)*	-26.58	Intergenic
Decreased	*a (agouti)*	-26.54	Intron
Decreased	*Cacna2d3*	-25.99	Intron
Increased	*Snai2*	36.27	Intron

Increased	*Egln3*	29.58	Intron
Increased	*(Ctrc)*	28.57	Intergenic
Increased	*(Snai1)*	28.13	Intergenic
Increased	*Pik3r4*	27.20	Exon
Increased	*Me1*	27.06	Intron
Increased	*Rubcnl*	26.94	Intron
Increased	*(Chrna5)*	26.47	Intergenic
Increased	*Ilf3*	25.76	3′ UTR
Increased	*(Tmem150cos)*	25.49	Intergenic

Female-specific DMRs

Effect of EtOH exposure	Gene symbol	Methylation change (%)	Annotation
Decreased	*(Pla2g4d)*	-39.54	Intergenic
Decreased	*Mfsd2a*	-37.35	Intron
Decreased	*Pygo1*	-31.79	Intron
Decreased	*(Plau)*	-29.63	Intergenic
Decreased	*(Jun)*	-29.28	Intergenic
Decreased	*Msx1os*	-28.73	Non-coding
Decreased	*(Zfp57)*	-28.31	Intergenic
Decreased	*Capza2*	-26.06	Intron
Decreased	*(Hpf1)*	-25.90	Intergenic
Decreased	*Traf6*	-25.16	Intron
Increased	*(Ssbp3)*	37.50	Intergenic

Increased	*Jdp2*	34.63	TTS	
Increased	*(Sox5os3)*	33.91	Intergenic	
Increased	*(Fam98b)*	32.86	Intergenic	
Increased	*(Usp7)*	32.66	Intergenic	
Increased	*Fgf12*	32.61	Intron	
Increased	*(Kpna1)*	32.06	Intergenic	
Increased	*Fam105a*	31.71	Intron	
Increased	*(Ppif)*	30.73	Intergenic	
Increased	*(Pax9)*	30.45	Intergenic	

787 **Table 2: Top 10 shared DMRs between male and female placentas**

			Shared top DMRs		
Effect of EtOH (males)	**Gene symbol**	**Methylation change (%) (males)**	**Effect of EtOH (females)**	**Methylation change (%) (females)**	**Annotation**
Cdk8	Decreased	-23.33	Decreased	-12.21	Intron
(Mirt1)	Decreased	-13.04	Decreased	-12.24	Intergenic
Hs3st2	Decreased	-12.54	Decreased	-10.83	Exon
Tmem267	Decreased	-12.51	Decreased	-10.67	Promoter-TSS
Rbm3os	Decreased	-10.07	Increased	15.50	Promoter-TSS
(Ifngas1)	Increased	15.22	Decreased	-20.68	Intergenic

(*Slc22a19*)	Increased	10.02	Decreased	-10.61	Intergenic
Tro	Increased	10.12	Increased	13.77	Promoter-TSS
Galp	Increased	15.12	Increased	10.21	Exon
(*Spry1*)	Increased	15.01	Increased	15.70	Intergenic
Clstn3	Increased	13.36	Increased	15.06	Exon
(*Spry1*)	Increased	13.08	Increased	12.83	Intergenic
Spry1	Increased	13.00	Increased	16.15	Intergenic
(*Spry1*)	Increased	12.05	Increased	11.63	Intergenic
Socs5	Increased	11.55	Increased	13.67	Intron
Psd3	Increased	11.40	Increased	10.02	Intron
Socs5	Increased	11.02	Increased	11.20	Exon
Wwox	Increased	10.80	Increased	10.94	Intron
Trh	Increased	10.18	Increased	11.33	Intergenic

789 **Table 3: Top 10 decreased and increased sex-specific DEGs**

Male-specific DEGs			
Condition	**Gene symbol**	**Log2 fold change**	***p*-value**
Decreased	*Ear1*	-5.66	0.04
Decreased	*Klk15*	-4.67	0.001
Decreased	*Kap*	-4.31	0.01
Decreased	*Ang2*	-4.03	0.05
Decreased	*Pcdhb8*	-3.93	0.04

Decreased	*Jsrp1*	-3.74	0.01
Decreased	*Cdc20b*	-3.64	0.04
Decreased	*Aadacl3*	-3.53	0.04
Decreased	*Shisa3*	-3.11	0.01
Decreased	*Crabp1*	-3.05	0.001
Increased	*Lrrc4b*	4.38	0.01
Increased	*C8a*	3.51	0.01
Increased	*Alb*	3.43	0.02
Increased	*Krt77*	3.34	0.04
Increased	*Gria1*	3.24	0.03
Increased	*Stmn3*	3.08	0.03
Increased	*Fgg*	3.03	0.02
Increased	*Trpm2*	2.95	0.02
Increased	*Spink3*	2.94	0.03
Increased	*Krt6b*	2.93	0.01

Female-specific DEGs

Condition	Gene symbol	Log2 fold change	*p*-value
Decreased	*Ddn*	-3.88	0.04
Decreased	*Caly*	-3.86	0.04
Decreased	*Serpinb2*	-2.89	0.01
Decreased	*Dsc1*	-2.81	0.01
Decreased	*Cyp2b19*	-2.63	0.003
Decreased	*Cyp2ab1*	-2.46	0.02

Decreased	*Dsg3*	-1.96	0.002
Decreased	*Serpina3c*	-1.96	0.04
Decreased	*Nol4*	-1.86	0.05
Decreased	*Marco*	-1.74	0.03
Increased	*Serpina1c*	4.96	0.003
Increased	*Mir296*	4.73	0.001
Increased	*Oprm1*	4.06	0.03
Increased	*Klk1b8*	4.06	0.02
Increased	*Fgf23*	4.04	0.04
Increased	*Dmrt3*	3.86	0.03
Increased	*St6galnac1*	3.67	0.05
Increased	*Myot*	3.24	0.05
Increased	*C6*	3.10	0.03
Increased	*Pigr*	3.08	0.02

791 **Table 4: Top 10 shared DEGs**

Shared top DEGs						
Effect of EtOH (males)	**Gene symbol**	**Log2 fold change (males)**	***p*-value (males)**	**Effect of EtOH (females)**	**Log2 fold change (females)**	***p*-value (females)**
Decreased	*Dok7*	-0.92	0.00	Decreased	-0.60	0.03
Decreased	*Spata2l*	-0.36	0.04	Decreased	-0.39	0.04
Decreased	*Foxf2*	-0.55	0.03	Decreased	-0.38	0.02

Decreased	*Chchd10*	-0.41	0.01	Decreased	-0.33	0.01
Decreased	*Timp2*	-0.44	0.04	Decreased	-0.33	0.01
Decreased	*Gpr156*	-0.52	0.01	Decreased	-0.33	0.01
Decreased	*Map2k6*	-0.37	0.02	Decreased	-0.31	0.05
Decreased	*St8sia5*	-0.61	0.00	Decreased	-0.29	0.04
Decreased	*Ypel3*	-0.32	0.00	Decreased	-0.23	0.03
Decreased	*Ifit2*	-0.43	0.02	Increased	0.33	0.04
Increased	*Dsc3*	1.74	0.05	Decreased	-2.10	0.03
Increased	*Dnajc6*	1.22	0.03	Increased	0.87	0.04
Increased	*Syt14*	0.60	0.00	Increased	0.60	0.01
Increased	*Pm20d1*	0.51	0.01	Increased	0.55	0.004
Increased	*Tceal7*	0.50	0.02	Increased	0.51	0.01
Increased	*Pdzk1ip1*	0.69	0.00	Increased	0.47	0.04
Increased	*Serpinb1a*	0.59	0.03	Increased	0.46	0.02
Increased	*Musk*	0.54	0.00	Increased	0.40	0.04
Increased	*Gsta3*	0.64	0.01	Increased	0.40	0.04
Increased	*Scel*	0.53	0.00	Increased	0.28	0.04

FIGURES and LEGENDS

Figure 1: Absence of morphometric alterations in E18.5 placentas following preimplantation alcohol exposure

Evaluation of the impact of preimplantation alcohol exposure on **A)** sex ratios of litters (control: males *n*=47, females *n*=39; ethanol-exposed: males *n*=62, females *n*=39); **B) i)** total embryos, number of **ii)** male and **iii)** female embryos per litter (control: *n*=8 litters; ethanol-exposed: *n*=10 litters); **C)** placental

surface area in **i)** males and **ii)** females (control embryos: males, $n=19$, females, $n=24$; ethanol-exposed

embryos: males $n=30$, females $n=22$; 6 litters, excluded litters used for placentas with decidua); **D)**

placental weight in **i)** males and **ii)** females (control embryos: males, $n=19$, females, $n=24$; ethanol-

exposed embryos: males, $n=30$, females, $n=22$; 6 litters, excluded litters used for placentas with decidua);

E) embryo weight in **i)** males and **ii)** females (control embryos: males, $n=31$, 8 litters; females, $n=28$, 8

litters; ethanol-exposed embryos: males, n=46, 10 litters; females, $n=32$, 8 litters); **F)** placental efficiency

(embryo-to-placental weight ratio) in **i)** males and **ii)** females (control embryos: males, $n=19$, females,

$n=24$; ethanol-exposed embryos: males, $n=30$, females, $n=22$; 6 litters, excluded litters used for placentas

with decidua). All data represent the mean $\pm$ standard deviation (SD). Significant differences were

assessed by *t*-test with Welch's correction. **: $p<0.01$. Ctrl: control, EtOH: ethanol-exposed.

Figure 2: Sex-specific DNA methylation dysregulation in the placenta following early preimplantation alcohol exposure

Scatterplot of differentially methylated regions (DMRs) between control and ethanol-exposed **A)** male and **C)** female placentas. Blue (male) or green (female) dots represent tiles with ≥ 10% methylation changes in ethanol-exposed placentas compared to controls (number of DMRs: male: 991, female: 1309). CpG methylation levels in control and ethanol-exposed **B)** male and **D)** female placentas based on DMR annotations (left Y-axis) and DMR distributions across genomic annotations (right Y-axis). Top enriched pathways in **E)** male (542 unique gene DMRs) and **F)** female (749 unique gene DMRs) DMRs based on Metascape analysis. CpG methylation levels of top changed DMRs or DMRs associated with the top

827   enriched pathways shown in **E** and **F** in control and ethanol-exposed **G)** male (left; grey/blue) and **H)**

828   female placentas.

Figure 3: Female ethanol-exposed placentas tend to lose methylation in CpG rich regions

A) Distribution of DMRs between control and ethanol-exposed male (blue) and female (green) placentas

based on their proximity to CpG-rich regions (CpG islands (CGIs). Regions are defined as shores (≤ 2

kb from CGIs), shelves (2–4 kb from CGIs), and open sea (≥ 4 kb from CGIs). **B)** Methylation differences

in male- (blue) and female- (green) specific DMRs based on their proximities to CGIs. DNA methylation

distributions in different CpG density contexts in **C)** males and **D)** females control and ethanol-exposed

placentas. Ctrl: control, EtOH: ethanol-exposed.

Figure 4: Gene expression profiles are especially affected in the male placenta following preimplantation alcohol exposure

A-B) Differential expression analyses between control and ethanol-exposed placenta in **A)** males and **B)**

females. Colored dots represent statistically significant DEGs ($p<0.05$; males, 1046; females, 340).

Darker dots represent DEGs with log2 fold change values $> |0.5|$ (males, $n=359$; females, $n=157$). **C-D)**

Heatmaps of DEG expression levels (z-scores) in control and ethanol-exposed **C)** male and **D)** female

placentas. **E-F)** Top pathways for **E)** male (1046) and **F)** female (340) DEGs based on Metascape

analysis. **G-H)** Normalized read counts of selected DEGs in **G)** male and **H)** female ethanol-exposed

placentas. Data represent the mean ± SD. Ctrl: control, EtOH: ethanol-exposed.

Figure 5: Sex-specific regions show both DNA methylation and gene expression alterations in the

placenta at late gestation in response to early preimplantation alcohol exposure

A) Heatmaps of CpG methylation (left) and gene expression (z-score; right) levels of DEGs containing

intragenic DMRs between control and ethanol-exposed male (46 DMRs, 42 DEGs) **B)** Normalized read

counts of DEGs and DMRs in male placentas. **C)** Heatmaps of CpG methylation (left) and gene

expression (z-score; right) levels of DEGs containing intragenic DMRs between control and ethanol-

exposed female (16 DMRs, 15 DEGs) placentas. **D**) Normalized read counts of DEGs and DMRs in

female placentas. Data represent the mean ± SD.

Figure 6: Identification of distinctive placental DNA methylation signatures indicative of early
preimplantation alcohol exposure

A) Principal component analysis showing the clustering of control and ethanol-exposed samples based

on their CpG methylation levels in the 24 DMRs making up the signature.. **B)** Heatmap of the Euclidean

distances of each control and ethanol-exposed sample based on their CpG methylation levels in the 24

DMRs. **C)** Feature importance plot of the DMRs based on LASSO analysis. **D)** CpG methylation levels

in control and ethanol-exposed male and female placental DMRs making up the signature.. Genes in

brackets are the closest gene to the intergenic DMR. Ctrl: control, EtOH: ethanol-exposed.

SUPPLEMENTAL FIGURES and LEGENDS

Supplemental Figure 1: Increased variability in female ethanol-exposed placenta morphometric measurements

Histological morphometric measurements of E18.5 placentas based on placental cross-sections (Males : Ctrl *n* = 10 ; EtOH *n* = 13. Females Ctrl *n*= 10; EtOH *n* =12). Assessment of the impact of preimplantation alcohol exposure on **A)** junctional zone (Jcz) area; **C)** Labyrinth zone (Lbz) area; **E)** thickness of the Jcz; **G)** thickness of the Lbz and their respective ratio to the total placenta area (**B, D**) or thickness (**F, H**). **I)**

883   Representative images of hematoxylin and eosin-stained control and ethanol-exposed samples. The Jcz

884   and Lbz are delimited by black lines. All data represent the mean $\pm$ SD. Significant differences were

885   assessed by *t*-test with Welch's correction and *F* tests for the variance analysis. † Higher variance in

886   ethanol-exposed placentas; *F*-test, $p<0.05$. Ctrl: control, EtOH: ethanol-exposed.

Supplemental Figure 2: Establishing DNA methylation profiles in alcohol-exposed late-gestation placentas (E18.5)

A) Schematic of the sex-related genome-wide CpG methylation analysis in male (Ctrl, $n=3$; EtOH, $n=5$) and female (Ctrl, $n=3$; EtOH, $n=5$) placentas at E18.5 to identify all tiles (100bp, $> 10\times$ coverage) associated with either males (791780), females (774904), or both sexes (751083), as well as male-specific (991), female-specific (1309), and common ($n=21$) DMRs). **B)** Schematic of sex-related differential transcriptomic analyses in male (Ctrl, $n=3$; EtOH, $n=5$) and female (Ctrl, $n=3$; EtOH, $n=5$) placentas at E18.5. Identification of all expressed genes in both sexes (31,235) and male-specific (1046), female-specific (340), and shared (54) DEGs.

Supplemental Figure 3: DNA methylation level distribution across differentially methylated regions

Tiles distributions across DNA methylation levels in **A)** male- and **C)** female-specific control and ethanol-exposed DMRs. Proportion of **B)** male and **D)** female DMRs associated with changes in CpG methylation levels between control and ethanol-exposed placentas. Data represent the mean ± SD. *: $p<0.05$, **: $p<0.01$, ***: $p<0.001$, ****: $p<0.0001$ by two-proportions z-test.

Supplemental Figure 4: Regions presenting shared differential DNA methylation patterns in ethanol-exposed male and female placentas are enriched in genes related to growth

A) Percentage of change in the 21 shared DMRs in males and females. **B)** CpG methylation levels of shared DMRs in control and ethanol-exposed male (left; grey/blue) and female (right; grey/green) placentas. **C)** Top pathways enriched for shared DMRs (11 unique gene DMRs) based on Metascape analysis. **D)** CpG methylation levels for DMRs located in enriched growth-regulating genes (from **C**) in control and ethanol-exposed male (left dots; grey/blue) and female (right dots; grey/green) placentas. Data represent the mean ± SD. Ctrl: control, EtOH: ethanol-exposed.

Supplemental Figure 5: Sex-specific changes in different regions of the same genes in ethanol-exposed placentas

A) Heatmaps of the CpG methylation levels of DMRs within the same genes in male (blue) and female (green) placentas. **B)** Methylation differences between males and females. Connected dots represent DMRs in a shared gene. **C)** Top pathways enriched for shared genes with male and female intragenic DMRs (69 unique genes) based on Metascape analysis. **D)** CpG methylation levels of DMRs located in genes related to top enriched pathways in control and ethanol-exposed male (left dots; grey/blue) and female (right dots; grey/green) placentas. Data represent the mean ± SD. Ctrl: control, EtOH: ethanol-exposed.

Supplemental Figure 6: Shared DEGs are similarly altered in male and female ethanol-exposed placentas

A) Heatmaps showing expression level (*z*-score) of common differentially expressed genes (DEGs) between male (blue; left) and female (green; right) placentas (54) following ethanol exposure. **B-C)** Scatterplots of the normalized read counts of the 54 common DEGs in **B)** male and **C)** female control and ethanol-exposed placentas. **D)** Normalized read counts of 2/4 genes with opposing expression changes between male and female ethanol-exposed placentas. **E)** Top pathways enriched for common

956 DEGs (54) based on Metascape analysis. **F)** Normalized read counts of genes with altered expression in

957 male and female ethanol-exposed placentas. Data represent the mean ± SD. Ctrl: control, EtOH: ethanol-

958 exposed.

Supplemental Figure 7: Early preimplantation alcohol exposure selectively alters the epigenetic profiles of placental development genes

A) Scatterplot of the tiles located in placental development genes in control and ethanol-exposed **i)** male and **ii)** female placentas. Blue (male) and green (female) dots represent tiles with ≥ 10% methylation changes in ethanol-exposed placentas compared to controls (males, 3; females, 3). **B)** Differential expression analyses of placental development genes between control and ethanol-exposed **i)** male and **ii)**

972   female placentas. Blue (male) and green (female) dots represent genes that were significantly changed

973   ($p<0.05$) in ethanol-exposed placentas (males, 19; females, 8) compared to controls. Ctrl: control, EtOH:

974   ethanol-exposed.

Supplemental Figure 8: Early preimplantation alcohol exposure alters epigenetic profiles of placental imprinting genes in late-gestation placentas

A) Scatterplot of tiles located in placental imprinting genes in control and ethanol-exposed **i)** male and **ii)** female placentas. Blue (male) and green (female) dots represent tiles with ≥ 10% methylation changes

in ethanol-exposed placentas (males, 17; females, 8) compared to controls. **B)** Differential expression

analysis of placental imprinting genes between control and ethanol-exposed **i)** male and **ii)** female

placentas. Blue (male) and green (female) dots represent genes that were significantly changed ($p<0.05$)

in ethanol-exposed placentas (males, 18; females, 1) compared to controls. **C)** CpG methylation levels

in control and ethanol-exposed male (left; grey/blue) and female (right; grey/green) placentas for DMRs

located in placental imprinting genes. **D-E)** Normalized read counts of placental imprinting genes with

altered expression in **D)** male (blue) and **E)** female (green) ethanol-exposed placentas compared to

controls. Data are the mean ± SD. Ctrl: control, EtOH: ethanol-exposed.

Supplemental Figure 9: Growth-related genes with altered DNA methylation after ethanol exposure only show expression changes in male placentas

A) CpG methylation levels in control and ethanol-exposed male placentas for DMRs located in the DEGs **i)** *Clstn3* and **ii)** *Tro*. **B)** Normalized read counts of **i)** *Clsnt3* and **ii)** *Tro* in control and ethanol-exposed male placentas (*n=3–5* for each group). All data represent the mean ± SD. Ctrl: control, EtOH: ethanol-exposed.

1007

Supplemental Figure 10: Differential expression is not directly linked to differential DNA methylation in late-gestation placentas following early preimplantation alcohol exposure

1008

1009

1010 DNA methylation differences between control and ethanol-exposed (X-axis) placentas and changes in

1011 gene expression (log2 fold change; Y-axis) in regions containing both DMRs and DEGs between control

1012 and ethanol-exposed placentas in **A)** males and **B)** females. Ctrl: control, EtOH: ethanol-exposed.

1013

1014

1015

1016

1017

Supplemental Figure 11: The transcriptomic signature of PAE does not enable PAE detection

A) Principal component analysis showing the clustering of control and ethanol-exposed samples based on their gene expression levels in the 20 DEGs making up the signature.. **B)** Heatmap of the Euclidean distances of each control and ethanol-exposed sample based on their gene expression levels in the 20 DEGs. **C)** Feature importance plot of the DEGs based on LASSO analysis.

1018

1019

1020

1021

1022

1023

1024

1025 **SUPPLEMENTARY TABLES**

1028

1029   **Table S2: Male- and female-specific DMRs located in imprinting genes**

		Male-specific DMRs in imprinting genes			
Effect of EtOH exposure	**Gene symbol**	**Tile ID**	**Methylation change (%)**	**Annotation**	**In ICR?**
Decreased	*Trappc9*	chr15.72945601.72945700	-31.79	Intron	No
Decreased	*Phf17*	chr3.41556301.41556400	-13.38	Intron	No
Increased	*Peg13*	chr15.72810001.72810100	10.24	Non-coding	Yes
Increased	*Grb10*	chr11.12025901.12026000	11.29	Intron	Yes
Increased	*Peg13*	chr15.72809901.72810000	11.39	Non-coding	Yes
Increased	*H13*	chr2.152670801.152670900	11.83	Intron	No
Increased	*Airn*	chr17.12741901.12742000	11.94	Intron	Yes
Increased	*Kcnq1ot1*	chr7.143295601.143295700	13.34	Non-coding	Yes
Increased	*Airn*	chr17.12741801.12741900	13.60	Intron	Yes
Increased	*Airn*	chr17.12741701.12741800	14.11	Intron	Yes
Increased	*Nap1l5*	chr6.58906901.58907000	14.84	Exon	Yes
Increased	*Commd1*	chr11.22973201.22973300	15.38	Promoter-TSS	Yes
Increased	*Cmah*	chr13.24468301.24468400	15.96	Intron	No

Increased	Grb10	chr11.12035301.12035400	16.20	Intron	No
Increased	Peg10	chr6.4748601.4748700	16.47	Intron	Yes
Increased	Nap1l5	chr6.58907001.58907100	17.25	Promoter-TSS	Yes
Increased	Nap1l5	chr6.58907101.58907200	18.13	Promoter-TSS	Yes

Female-specific DMRs in imprinting genes

Effect of EtOH exposure	Gene Name	Tile ID	Methylation change (%)	Annotation	In ICR?
Decreased	Grb10	chr11.12025501.12025600	-13.78	Intron	Yes
Increased	Art5	chr7.102095201.102095300	10.68	Non-coding	No
Increased	Plagl1	chr10.13090701.13090800	10.95	Promoter-TSS	Yes
Increased	H13	chr2.152670901.152671000	11.87	Intron	No
Increased	Airn	chr17.12742201.12742300	12.03	Intron	Yes
Increased	Pde10a	chr17.8909701.8909800	12.48	Intron	No
Increased	Inpp5f	chr7.128690801.128690900	12.72	Intron	No
Increased	Peg3	chr7.6729401.6729500	14.97	Intron	Yes
Increased	Commd1	chr11.22973101.22973200	18.60	Promoter-TSS	Yes

1030

1031

1032

 Table S3: Male- and female-specific DEGs located in imprinting genes

Male-specific DEGs in imprinting genes			
Effect of EtOH exposure	**Gene Name**	**Log2FoldChange**	**_p_-value**
Decreased	_Htra3_	-2.220	0.002
Decreased	_Gatm_	-2.103	0.001
Decreased	_Ampd3_	-1.234	0.016
Decreased	_Dcn_	-1.068	0.006
Decreased	_Qpct_	-0.645	0.003
Decreased	_Rasgrf1_	-0.525	0.024
Decreased	_Ano1_	-0.437	0.041
Decreased	_Tnfrsf23_	-0.430	0.034
Decreased	_Osbpl5_	-0.377	0.042
Decreased	_Zrsr1_	-0.248	0.018
Increased	_Gab1_	0.142	0.042
Increased	_Ube3a_	0.183	0.045
Increased	_Peg10_	0.254	0.006
Increased	_Ppp1r9a_	0.258	0.001
Increased	_Grb10_	0.273	<0.001
Increased	_Igf2r_	0.307	0.048
Increased	_Slc22a3_	0.327	0.002
Increased	_Cobl_	0.354	0.031

Female-specific DEGs in imprinting genes			
Effect of EtOH exposure	**Gene Name**	**Log2FoldChange**	***p*-value**
Increased	*Slc22a2*	1.792	0.010

1034