Localization of Cadherins in the postnatal cochlear epithelium and their relation to space formation.

Holly J. Beaulac[1] and Vidhya Munnamalai[1*]

[1]The Jackson Laboratory, Bar Harbor, ME 04609, USA

* Correspondence:

Vidhya Munnamalai

The Jackson Laboratory

Bar Harbor, ME 04609, USA

Email: vidhya.munnamalai@jax.org

Phone: 207-288-6916

Keywords: Cochlea, Support Cells, F-actin, Cell Adhesion, Cadherin, Tunnel of Corti

Running title: Modifications in E-cadherin cell contacts form the tunnel of Corti in the cochlea.

Key findings:

F-actin remodeling occurs between E18.5 to P7 in the cochlear sensory epithelium.

Transient changes of F-actin cytoskeleton drives epithelial morphogenesis.

Fluid-filled spaces in epithelium is driven by changes in cell adhesion.

Funding: This work was supported by the National Institutes of Health (R21DC016376) to V.M.

Abstract (200 words)

The sensory epithelium of the cochlea, the organ of Corti, has complex cytoarchitecture consisting of mechanosensory hair cells intercalated by epithelial support cells. The support cells provide important trophic and structural support to the hair cells. Thus, the support cells must be stiff yet compliant enough to withstand and modulate vibrations to the hair cells. Once the sensory cells are properly patterned, the support cells undergo significant remodeling from a simple epithelium into a structurally rigid epithelium with fluid-filled spaces in the murine cochlea. Cell adhesion molecules such as cadherins are necessary for sorting and connecting cells in an intact epithelium. To create the fluid-filled spaces, cell adhesion properties of adjoining cell membranes between cells must change to allow the formation of spaces within an epithelium. However, the dynamic localization of cadherins has not been properly analyzed as these spaces are formed. There are three cadherins that are reported to be expressed during the first postnatal week of development when the tunnel of Corti forms in the cochlea. In this study, we characterize the dynamic localization of cadherins that are associated with cytoskeletal remodeling at the contacting membranes of the inner and outer pillar cells flanking the tunnel of Corti.

1. Introduction

Epithelial morphogenesis requires complex cytoskeletal remodeling and maturation [1,2] and is essential for proper organ formation and organ function. Cadherin-driven cell adhesion provides epithelial stability during morphogenesis by inducing cell-to-cell contact-mediated mechanical coupling with the cellular F-actin cytoskeleton [2,3]. Cadherins are Ca^{2+}-dependent cell adhesion molecules that lead to the formation of mature adherens junctions during tissue development [4]. Initial recruitment of cadherin molecules at the membrane is followed by cadherin clustering and is then stabilized by coupling to the actin cytoskeleton [5-7]. Maturation of adherens junctions and strengthening of cell-cell adhesion further induces tension-dependent actin remodeling and recruitment of actin binding proteins to the site of de novo actin synthesis [8,9].

In the cochlea, auditory function is dependent on proper epithelial morphogenesis of the sensory epithelium, the organ of Corti. The organ of Corti attains a highly specialized organization of mechanosensory hair cells intercalated by support cells [10,11]. At neonatal stages

of the mouse cochlea, the organ of Corti has already established its classical mosaic pattern of

hair cells intercalated by support cells (Figure 1). There are two classes of mechanosensory hair

cells: one row of inner hair cells (IHCs) and three rows outer hair cells (OHCs) (Figure 1A) that

are innervated by afferent and efferent neurons enabling the transmission of auditory information

along central auditory pathways [12,13]. The inner and outer hair cells are segregated by the IPC

and OPC support cells. The outer hair cells are intercalated by a specific subtype of support cells

known as the Deiters' cells (DCs) (Figure 1A, A'). Thus far, most studies have focused on the

cytoskeleton and cytoskeletal remodelers related to stereocilia formation in sensory hair cells;

however, the support cells have not received the same attention. Recent studies have shown that

the support cells have important electrophysiology, metabolic and sensor properties that are

necessary for proper auditory function [14-16]. The support cells also serve as important support

structures for the hair cells [17] and help to establish the overall morphology and complex

cytoarchitecture of the sensory epithelium. The organ of Corti must be stable enough to

withstand vibrational movements of the basilar membrane [16,18] and the movement of fluid

through the epithelium [19-21] to facilitate the propagation of sound information. During sensory

epithelial morphogenesis, fluid-filled spaces such as the tunnel of Corti, the Nuel space and the

outer tunnel are formed within the epithelium [22], which require stabilization of cell and epithelial

structure. We show that between late embryonic and early postnatal stages, the support cells

undergo significant dynamic morphological changes. However, thus far, no studies have

uncovered the spatiotemporal characteristics of cytoskeletal modifications and how the tunnel of

Corti is formed during maturation of the cochlear sensory epithelium.

In this study, we focus on the F-actin cytoskeletal changes that the support cells undergo

and the cell adhesion modifications that occur during a specific time in development when fluid-

filled spaces form in the sensory epithelium, particularly the pillar cells (PCs) that undergo the

most drastic structural cytoskeletal remodeling to form the tunnel of Corti. The tunnel of Corti

forms between the IPCs and the OPCs during the neonatal ages of development; however, a

close examination of the spatial changes of the F-actin cytoskeleton and cell adhesion of the

cochlea undergoing extensive remodeling and the tunnel of Corti opening, has not been

performed. Although the F-actin cytoskeleton establishes the structure and shape of all cells [23],

its expression and structural organization can vary in different cell types in the cochlea. Since

epithelial integrity is dependent on cadherins in the cochlea; for example, integrity of the stria

vascularis is dependent on cadherins [24], we also examine whether the cellular spatial localization

of cadherins play a role in establishing the integrity of the sensory epithelium with spaces

interspersed within the cochlear sensory epithelium. Since this study focuses on the support cells,

the cross-sectional view of the organ of Corti reveals structural details (Figure 1A, A') that are

difficult to appreciate in wholemount cochlear preparations where the stereocilia on hair cells

and the overall patterning of the sensory epithelium are the prime focus (Figure 1B). We can also

view the immunolabeling in the entire cell along the apico-basolateral axis with finer subcellular

differences from the apical, mid- and basolateral regions of the SCs and the membrane versus

cytoplasmic localization of proteins. The whole mount provides a single plane view through the

epithelium at a time, which may lead to missing out key observations.

Three cadherin molecules reported to be expressed in the mouse cochlea during the first

postnatal week are neural (N-), epithelial (E-), and placental (P-) cadherins [25-27]. The tunnel of

Corti forms between the inner pillar cells (IPC) and the outer pillar cells (OPCs), spaces of Nuel

and the outer tunnel [27]. A recent study suggested that the abnormal premature enlargement of the

tunnel of Corti was caused by loss of P-cadherin. From the study's wholemount images, P-

cadherin localization in the PCs appeared to be apical [25] but was not clear from single optical

planes. Cross-sectional views of cadherin localization along the apico-basolateral axis during the

narrow temporal window as the tunnel of Corti forms would provide a clearer view of how E-

cadherin or N- or P- cadherin localization influence tunnel formation under normal

developmental conditions. In this study, we analyze the spatiotemporal localization of these three

cadherins in the cochlea.

2. Results and Discussion

To identify the correct stage to observe the opening of the tunnel of Corti and

characterize the associated subcellular changes in the F-actin cytoskeletal organization, we first

analyzed sectioned cochlear tissue on E18.5, P6 and P8. On E18.5, Phalloidin labeling shows

that F-actin is not distinguishable in the pillar cells (dotted lines) but is clearly present at the

membranes adjoining both the OHCs and the DCs (Figure 2A). Thus, we examined markers for

cell adhesion along the OHC-DC membrane contacts. A prior study showed an increase in

junctional E-cadherin in the apical junctions of the PCs in wholemount mouse cochleas during

the first postnatal week after birth [28]. Thus, E-cadherin signaling is a strong candidate for PC

remodeling. Comparing F-actin with E-cadherin, we found E-cadherin localized to the OHC-DC

membrane contacts (Figure 2A'). On E18.5, E-cadherin was also localized to the membranes of

Hensen's cells (Figure 2A', green arrow). At this stage, there was no specific enrichment of F-

actin or E-cadherin in the PCs. Cell adhesion induces bi-directional signaling that promotes the

recruitment of phosphorylated actin binding proteins at their tyrosine residues (pTyr), a sign of

active actin assembly [29,30]. Immunolabeling for pTyr was undetectable on E18.5, presumably

because minimal remodeling of the F-actin cytoskeleton was required at this stage of

development (Figure 2A"). There was no notable patterns of phalloidin, E-cadherin, or pTyr

labeling in the PCs on E18.5 (Figure 2A"').

Next we analyzed cochleas on P6. The tunnel of Corti was still not formed at this stage.

However, actin filaments were now visible in the PCs, especially in the IPC and the IPC-OPC

membrane contacts (Figure 2B, arrow indicates mid-region of IPC-OPC membrane). There is a

$5e2 \pm 0.95e2$ % enrichment of F-actin at the membrane contacts of the mid-IPC-OPC region

relative to the cytoplasm (T-test, p = 1.2e-6). E-cadherin was also localized at the adjoining IPC-

OPC membranes, but E-cadherin was expressed from the apical to the mid-regions (arrow) along

the apico-basolateral axis of the IPC-OPC membranes, indicating there was an enrichment in cell

adhesion contacts associated correlating with F-actin in the mid-region of the PCs (Figure 2B',

arrow). We detected an enrichment of $2.9e2 \pm 0.52e2$ % in the mid-region of IPC-OPC

membrane contacts relative to the cytoplasm, assayed at the same region of interest as phalloidin

(p = 3.1e-6). At P6, there was minimal recruitment of pTyr actin binding proteins at the IPC-

OPC membrane contacts by $1.4e1 \pm 0.31e2$ percent relative to the cytoplasm, assayed at the

same region of interest as phalloidin p = 0.0004). This suggests that there may be a temporal

delay in pTyr protein recruitment once E-cadherin localization stabilizes (Figure 2B"). Merged

image show co-localization of Phalloidin with E-cadherin (Figure 2B"').

We then performed the same analysis in the P8 cochlea. By P8, the tunnel of Corti has

opened (Figure 2C). There is a clear F-actin bundle that has formed within the IPC (Figure 2C,

yellow arrow). However, by P8, F-actin and E-cadherin were no longer identifiable in the mid-

regions of the IPC-OPC contact membranes that were now occupied by the tunnel of Corti and

Nuel space. Despite the formation of the tunnel of Corti, the IPC and the OPC remain connected

at their adjoining apical junctions where F-actin and E-cadherin remains localized (Figure 2C,

2C', arrow). P-Tyr labeling was not present in the PCs at P8, suggesting that a critical window

was missed (Figure 2C", 2C'''). The time interval between P6 and P8 was too large to observe

the transient cytoskeletal modifications associated with the formation of the tunnel of Corti.

To capture the opening of the tunnel of Corti, we harvested P7 cochleas at 3 timepoints

throughout the day in the morning (9 am), mid-day (1 pm) and in the evening (5 pm) (Figure 3).

In the morning, the tunnel of Corti (ToC) was not yet opened (closed). We observed an

accumulation of F-actin at the mid-region of the IPC-OPC membranes (Figure 3A, arrow) along

the apico-basolateral axis, which will go onto to become the axial regions of the PCs that flank

the tunnel of Corti. We detected an enrichment of phalloidin labeling by $26.7e2 \pm 8.9e2$ % at the

mid-region of IPC-OPC membrane contacts relative to the cytoplasm (p= 5.9e-9). Coincidently,

in the same region of interest where we detected phalloidin enrichment, we also saw increases in

E-cadherin (Figure 3A', arrow) and pTyr labeling (Figure 3A", arrow) at the IPC-OPC

membrane contacts with enrichments of $23.6e2 \pm 12.2e2$ % (p = 5.9e-5) and $13.7e2 \pm 3.8e2$ % (p

= 4.1e-6) respectively relative to the cytoplasm. This indicates that F-actin, E-cadherin and pTyr

were co-localized at the same mid-region of interest at the IPC-OPC membranes (Figure 3A'''

inset, arrow). These enrichments of phalloidin, E-cadherin and pTyr at the IPC-OPC membranes

relative to the cytoplasm were higher in the closed tunnel on P7 than on P6 when the tunnel was

also closed, and E-cadherin and pTyr localization at the membrane were only 2.9 -fold and 1.4 -

fold higher than in the cytoplasm respectively (Figure 2B'-B'''). There was some pTyr labeling

at the apical junction that the mid-region (Figure 3A"), which suggests that there may be some

active F-actin assembly on the apical junctions of the PCs. Unlike the mid-apical region of the

IPC-OPC membrane contacts, there was no overlay or significant enrichment of F-actin, E-

cadherin and pTyr signals in the basolateral end of the PC membranes relative to cytoplasmic

levels (Figure 3A-A'''). We also observed pTyr labeling when the Nuel space has a very narrow

opening, which suggests there was also recruitment of actin binding proteins during initiation of

Nuel space formation (Figure 3A", red arrow). At the mid-timepoint on P7, the tunnel of Corti

was open at less than 100 μm^2 in cross-sectional area (small ToC). F-actin and E-cadherin

expression was not detectable in the mid-axial region of the IPC-OPC membranes but are present

at the apical junction (Figure 3B-B'). Interestingly, pTyr labeling remained high at the edges of

cell membranes (Figure 3B"-B'''). This suggests that these sites remained enriched for actin

binding proteins which help remodel cellular structure of the tunnel. In the final timepoint on P7,

the tunnel was large at more than 120 μm^2 (large ToC). F-actin and E-cadherin remained at the

apical regions of the PCs (Figure 3C'C'); while pTyr labeling was relatively minimal at the

apical junctions (Figure 3C"-C"'). This suggests that there was a greater recruitment of pTyr-

labeled actin binding proteins where the cochlear spaces are forming and not at stable contacts.

The variable localization of E-cadherin on the adjoining IPC-OPC membranes along the apico-

basolateral axis suggests that a reduction in cell adhesion precedes the opening of the tunnel of

Corti at the mid-region of the PC membranes and was accompanied by a large recruitment of

pTyr-labeled proteins. This data also show that P7 is critical timepoint when the support cells in

the sensory epithelium undergo significant cellular remodeling to attain its complex

cytoarchitecture. Conversely, stabilized F-actin will require less recruitment of actin binding

proteins as there are no sites for de novo F-actin synthesis e.g., in the apical junctions of the PCs.

N-cadherin is another cadherin known to be expressed in the developing cochlea [26]. We

examined whether N-cadherin is spatially positioned to influence the opening of the spaces in the

cochlear sensory epithelium. On P6, N-cadherin is visible in the inner hair cell (IHC), inner

phalangeal and inner border cell, but is clearly absent in the PCs and the DCs (Figure 4A-A').

This expression pattern remains consistent in P7 cochleas at the same three timepoints when the

tunnel of Corti was still closed (Figure 4B-B'), has a small opening (Figure 4C-C') and when the

tunnel opening was larger (Figure 4D-D'). Thus, the stable expression of N-cadherin and its

consistent absence of expression in the PCs suggests that N-cadherins do not influence the

opening of the tunnel of Corti.

A recent study showed that P-cadherin is expressed in the PCs of the cochlea during early

postnatal stages. Upon Thyroxine administration, P-cadherin was translocated from the PC

membranes to the cytoplasm, thereby reducing cell-cell adhesion [25]. The study analyzed

wholemount cochleas, showing images from a single plane; thus, the expression of P-cadherin

along the entire apico-basolateral axis of PCs was unclear. To determine its spatial localization,

we analyzed P-cadherin expression. On P6, P-cadherin was localized in the IPCs, but enriched at

the top apical end of the IPC (Figure 5A-A', white arrow). In addition to the PCs, we found P-

cadherin was also expressed in the membranes of the Hensen's cells and the cells in the outer

sulcus region (Figure 5A, green arrow). The expression of P-cadherin was much more substantial

in these lateral non-sensory cells compared with E-cadherin. On P7, while the tunnel of Corti

was still closed, P-cadherin remained enriched in the apical portion of the PCs, the Hensen's

cells and the outer sulcus cells (Figure 5B-B'). The apical spatial localization remained

unchanged while the tunnel of Corti opened (Figure 5C, 5C', 5D, 5D'). The expression of P-

cadherin remained stable in the Hensen's and outer sulcus cells as well.

Since pTyr labeled proteins were recruited to sites of F-actin and E-cadherin

accumulation at IPC-OPC membrane prior to the opening of the tunnel of Corti (Figure 3). We

postulated that in preparation for the tunnel opening, actin binding proteins are recruited to

facilitate the remodeling of F-actin in the PCs. Thus, we examined the expression of Cofilin, an

F-actin severing protein, on P6 and the three stages of tunnel of Corti formation on P7 (Figure 6).

Previous studies showed cofilin labeling in the HCs [31,32]; however, Cofilin localization in the

SCs has not been characterized thus far. Thus, we were surprised to find punctate labeling of

Cofilin in the cytoplasm of the SCs but not in the HCs (Figure 6A-D). On P6, we saw a 70 ± 10

% (p = 5.5e-7) enrichment of Cofilin localized in the mid-region of the IPC-OPC junction

relative to cofilin expression in the cytoplasm of the PCs (Figure 6A-A"). On P7, prior to

opening of the tunnel, we saw a 320 ± 40 % (p = 6.2e-7) enrichment of Cofilin localized in the

mid-region of the IPC-OPC junction relative to cofilin expression in the cytoplasm of the PCs

(Figure 6B-B", arrow). This suggests that during this critical period, Cofilin recruitment is

associated with the actin remodeling and aids in disrupting cell adhesion in the mid-region of the

IPC-OPC membranes.

The stable P-cadherin expression in the apical portion of the PCs is unlikely to influence

the opening of the tunnel of Corti under normal developmental conditions. On the other hand, the

dynamic and transient localization of E-cadherin in the mid- regions of the IPC-OPC

membranes, and its subsequent downregulation preceding the opening of the tunnel of Corti

makes the dynamic downregulation E-cadherin a much more likely mechanism for opening the

tunnel under normal developmental conditions. However, it is possible that in a disease context,

the loss of P-cadherin can exacerbate the tunnel opening and final size of the tunnel, disrupting

the overall stability of the sensory epithelium. Perhaps both E-cadherin and P-cadherin

trafficking are impacted when the cytoarchitecture of the sensory epithelium is disrupted under

disease conditions. Thus, follow up studies are needed to validate this. In this study, we show

evidence that significant actin remodeling takes place in the SCs during P6-P8 and paves way for

future investigations on actin assembly in the SCs on which little focus has been placed thus far.

Here we closely analyze the cell adhesion characteristics associated with the opening of the

tunnel of Corti. We predict that the same decrease in cadherin trafficking will promote the

formation of the Nuel space and the outer tunnel. Catenins are intracellular cell adhesion proteins

that link cadherins to the F-actin cytoskeleton [33]; thus, are critical players in stabilizing adherens

junctions and assembling epithelia, including the cochlear epithelium [26,28,34]. However, how their

localizations are modified during the critical stages of the opening of cochlear spaces would be a

good focus for future studies.

3.   Experimental Procedures

3.1 Mice

Time-mated Swiss Webster dams (Envigo, Indianapolis, IN, USA) were denoted as embryonic

day (E)0.5 on the first day a plug was observed. E18.5 embryos, and postnatal day (P)6-P8 pups

were euthanized by decapitation, in accordance with the Institutional Animal Care and Use

Committee (IACUC) guidelines at The Jackson Laboratory, Bar Harbor, and heads or ears were

harvested and fixed in 4% paraformaldehyde (PFA) (Electron Microscopy Sciences, Hatfield,

PA, USA).

3.2 Histology

After fixation, E18.5 heads were bisected and processed through a 10%, 20%, and 30% sucrose

gradient series, embedded with TFM and cryofrozen. For postnatal studies, cochleas were

extracted on P6, P7 and P8, and fixed in 4% PFA for three hours at 4°C, immersed in a 30%

sucrose and transferred to a 1:1 solution of 30% sucrose: Tissue Freezing Medium (TFM)

(General Data Healthcare, Cincinnati, OH, USA) before final embedding in 100% TFM and

cryofrozen. Tissues were cryosectioned at 12 μm for histology at all ages. To capture the

opening of the tunnel of Corti (ToC) on P7, inner ears were harvested and fixed at 3 timepoints

throughout the day in the morning (9 am), mid-day (1 pm) and in the evening (5 pm).

3.3 Immunofluorescence

Tissues were washed in PBS/0.5% Triton and blocked at for at room temperature (RT) 1 hour

with 2% Donkey serum (Jackson ImmunoResearch, West Grove, PA, USA) in PBS/0.5% Triton,

followed by overnight primary antibody incubation at 4°C. Primary antibodies included: rat anti-

E-cadherin (1:250, AB11512, Abcam, Waltham, MA, USA), sheep anti- N-cadherin (1:250,

AF6426, R&D Systems, Minneapolis, MN, USA), goat anti- P-cadherin (1:250, AF761, R&D

Systems), mouse anti-phosphoTyrosine (1:250, zms1b282, Sigma-Aldrich), anti-Cofilin (1:250,

5175, Cell Signaling Technology, Danvers, MA, USA). For N-cadherin, P-cadherin and Cofilin

immunolabeling, antigen retrieval was performed prior to primary antibody incubation with a 10

mM sodium citrate/ 0.05% Tween20 buffer at 6.0 pH for 15 min at 99°C. Since phalloidin

labeling of F-actin is incompatible with antigen retrieval, we co-labeled with an actin antibody

(1:250, MA5-11869, Invitrogen). Alexa-conjugated secondary antibodies (Invitrogen) were

incubated for 2 hours at RT. Nuclei were counter stained with DAPI (Abcam, Waltham, MA,

USA) and tissues were mounted with Fluoromount G mounting medium (Life Technologies

Corporation, Carlsbad, CA, USA).

3.4 Data Acquisition, Analysis and Image processing

60X magnification images were acquired on a Zeiss LSM 800 confocal microscope (ZEISS,

Oberkochen, BW, Germany) in The Jackson Laboratory microscopy core. Raw data were

analyzed using the FIJI software (NIH). At the morning time point, mid-turns were consistently

closed. In themed-day timepoint, we qualified small ToCs as those tunnels with a range of cross-

sectional areas between 35 μm^2 – 100 μm^2. In the afternoon timepoint, ToCs were consistently

open, and we qualified large ToCs to be greater than 120 μm^2. Data were collected and

quantified on 2 sections per cochlea for n = 6 cochleas at each time point. All 6 cochleas in each

of the presented figures were immunolabeled and data were acquired at the same time and

settings. To quantify enrichment of signals, a 0.6 μm^2 region of interest was used to measure

levels at the IPC-OPC membrane and the IPC cytoplasm within the same sample. We utilized the

same region of interest at the same coordinates across channels to define co-localized signals.

We calculated the percent difference between the membrane versus the cytoplasm. To qualify

specific localization at the membrane, student's t-test were performed to determine significant

differences in levels at the membrane versus the cytoplasm. Bon-Ferroni correction factor was

300 applied for multiple comparisons. TIFF images for figures were processed in Adobe Photoshop

301 and assembled in Adobe Illustrator (Adobe, San Jose, CA, USA).

4. Acknowledgements

The authors acknowledge the Microscopy Core at The Jackson Laboratory for equipment maintenance and upkeep, and assistance. The authors thank members of the Munnamalai lab for providing critical feedback on the manuscript.

5. Conflict of Interest

The authors declare no conflicts of interest or competing financial interests.

6. Author Contributions

H.J.B., study conceptualization and design, development of methodology, acquisition, data analysis, writing of the manuscript. V.M., funding, material support, study conceptualization and design, development of methodology, acquisition, analysis, and interpretation of data; writing and revision of the manuscript. All authors read and approved the final manuscript.

7. Ethics Approval

All experiments were performed under IACUC protocol 18013: PI Vidhya Munnamalai, in compliance with the U.S. Department of Health and Human Services and reviewed by The Jackson Laboratory Institutional Animal Care and Use Committee.

8. Data Availability

The datasets used and/or analyzed during the current study are available from the corresponding author on request.

9. References

1. Schock F, Perrimon N. Molecular mechanisms of epithelial morphogenesis. *Annu Rev Cell Dev Biol*. 2002;18:463-93. https://doi.org10.1146/annurev.cellbio.18.022602.131838.
2. Clarke DN, Martin AC. Actin-based force generation and cell adhesion in tissue morphogenesis. *Curr Biol*. May 24 2021;31(10):R667-R680. https://doi.org10.1016/j.cub.2021.03.031.
3. Samarage CR, White MD, Alvarez YD, et al. Cortical Tension Allocates the First Inner Cells of the Mammalian Embryo. *Dev Cell*. Aug 24 2015;34(4):435-47. https://doi.org10.1016/j.devcel.2015.07.004.
4. Leckband DE, de Rooij J. Cadherin adhesion and mechanotransduction. *Annu Rev Cell Dev Biol*. 2014;30:291-315. https://doi.org10.1146/annurev-cellbio-100913-013212.
5. Mege RM, Gavard J, Lambert M. Regulation of cell-cell junctions by the cytoskeleton. *Curr Opin Cell Biol*. Oct 2006;18(5):541-8. https://doi.org10.1016/j.ceb.2006.08.004.
6. Hong S, Troyanovsky RB, Troyanovsky SM. Binding to F-actin guides cadherin cluster assembly, stability, and movement. *J Cell Biol*. Apr 1 2013;201(1):131-43. https://doi.org10.1083/jcb.201211054.
7. Priest AV, Shafraz O, Sivasankar S. Biophysical basis of cadherin mediated cell-cell adhesion. *Exp Cell Res*. Sep 1 2017;358(1):10-13. https://doi.org10.1016/j.yexcr.2017.03.015.
8. Watabe-Uchida M, Uchida N, Imamura Y, et al. alpha-Catenin-vinculin interaction functions to organize the apical junctional complex in epithelial cells. *J Cell Biol*. Aug 10 1998;142(3):847-57. https://doi.org10.1083/jcb.142.3.847.
9. Yu Q, Holmes WR, Thiery JP, Luwor RB, Rajagopal V. Cortical tension initiates the positive feedback loop between cadherin and F-actin. *Biophys J*. Feb 15 2022;121(4):596-606. https://doi.org10.1016/j.bpj.2022.01.006.
10. Zine A, Van De Water TR, de Ribaupierre F. Notch signaling regulates the pattern of auditory hair cell differentiation in mammals. *Development*. Aug 2000;127(15):3373-83. https://doi.org10.1242/dev.127.15.3373.
11. McNeill H. Planar cell polarity: keeping hairs straight is not so simple. *Cold Spring Harb Perspect Biol*. Feb 2010;2(2):a003376. https://doi.org10.1101/cshperspect.a003376.
12. Munnamalai V, Fekete DM. The acquisition of positional information across the radial axis of the cochlea. *Dev Dyn*. Mar 2020;249(3):281-297. https://doi.org10.1002/dvdy.118.
13. Di Bonito M, Studer M. Cellular and Molecular Underpinnings of Neuronal Assembly in the Central Auditory System during Mouse Development. *Front Neural Circuits*. 2017;11:18. https://doi.org10.3389/fncir.2017.00018.
14. Chen T, Rohacek AM, Caporizzo M, et al. Cochlear supporting cells require GAS2 for cytoskeletal architecture and hearing. *Dev Cell*. May 17 2021;56(10):1526-1540 e7. https://doi.org10.1016/j.devcel.2021.04.017.
15. Velez-Ortega AC, Stepanyan R, Edelmann SE, et al. TRPA1 activation in non-sensory supporting cells contributes to regulation of cochlear sensitivity after acoustic trauma. *Nat Commun*. Jun 30 2023;14(1):3871. https://doi.org10.1038/s41467-023-39589-w.
16. Zhou W, Jabeen T, Sabha S, Becker J, Nam JH. Deiters Cells Act as Mechanical Equalizers for Outer Hair Cells. *J Neurosci*. Nov 2 2022;42(44):8361-8372. https://doi.org10.1523/JNEUROSCI.2417-21.2022.
17. Wan G, Corfas G, Stone JS. Inner ear supporting cells: rethinking the silent majority. *Semin Cell Dev Biol*. May 2013;24(5):448-59. https://doi.org10.1016/j.semcdb.2013.03.009.

18. Reichenbach T, Hudspeth AJ. The physics of hearing: fluid mechanics and the active process of
the inner ear. *Rep Prog Phys*. Jul 2014;77(7):076601. https://doi.org10.1088/0034-
4885/77/7/076601.
19. Zagadou BF, Barbone PE, Mountain DC. Significance of the Microfluidic Flow Inside the Organ of
Corti. *J Biomech Eng*. Aug 1 2020;142(8). https://doi.org10.1115/1.4046637.
20. Wangemann P. Supporting sensory transduction: cochlear fluid homeostasis and the
endocochlear potential. *J Physiol*. Oct 1 2006;576(Pt 1):11-21.
https://doi.org10.1113/jphysiol.2006.112888.
21. Meenderink SWF, Dong W. Organ of Corti vibrations are dominated by longitudinal motion in
vivo. *Commun Biol*. Nov 24 2022;5(1):1285. https://doi.org10.1038/s42003-022-04234-7.
22. Ulfendahl M, Scarfone E, Flock A, Le Calvez S, Conradi P. Perilymphatic fluid compartments and
intercellular spaces of the inner ear and the organ of Corti. *Neuroimage*. Sep 2000;12(3):307-13.
https://doi.org10.1006/nimg.2000.0617.
23. Pollard TD, Cooper JA. Actin, a central player in cell shape and movement. *Science*. Nov 27
2009;326(5957):1208-12. https://doi.org10.1126/science.1175862.
24. Trowe MO, Maier H, Petry M, Schweizer M, Schuster-Gossler K, Kispert A. Impaired stria
vascularis integrity upon loss of E-cadherin in basal cells. *Dev Biol*. Nov 1 2011;359(1):95-107.
https://doi.org10.1016/j.ydbio.2011.08.030.
25. Zhang H, Xie L, Chen S, Qiu Y, Sun Y, Kong W. Thyroxine Regulates the Opening of the Organ of
Corti through Affecting P-Cadherin and Acetylated Microtubule. *Int J Mol Sci*. Nov 1 2022;23(21).
https://doi.org10.3390/ijms232113339.
26. Chacon-Heszele MF, Ren D, Reynolds AB, Chi F, Chen P. Regulation of cochlear convergent
extension by the vertebrate planar cell polarity pathway is dependent on p120-catenin.
Development. Mar 2012;139(5):968-78. https://doi.org10.1242/dev.065326.
27. Whitlon DS. E-cadherin in the mature and developing organ of Corti of the mouse. *J Neurocytol*.
Dec 1993;22(12):1030-8. https://doi.org10.1007/BF01235747.
28. Luo WW, Wang XW, Ma R, et al. Junctional E-cadherin/p120-catenin Is Correlated with the
Absence of Supporting Cells to Hair Cells Conversion in Postnatal Mice Cochleae. *Front Mol
Neurosci*. 2018;11:20. https://doi.org10.3389/fnmol.2018.00020.
29. Andl CD, Rustgi AK. No one-way street: cross-talk between e-cadherin and receptor tyrosine
kinase (RTK) signaling: a mechanism to regulate RTK activity. *Cancer Biol Ther*. Jan 2005;4(1):28-
31. https://doi.org10.4161/cbt.4.1.1431.
30. Brunton VG, MacPherson IR, Frame MC. Cell adhesion receptors, tyrosine kinases and actin
modulators: a complex three-way circuitry. *Biochim Biophys Acta*. Jul 5 2004;1692(2-3):121-44.
https://doi.org10.1016/j.bbamcr.2004.04.010.
31. McGrath J, Tung CY, Liao X, et al. Actin at stereocilia tips is regulated by mechanotransduction
and ADF/cofilin. *Curr Biol*. Mar 22 2021;31(6):1141-1153 e7.
https://doi.org10.1016/j.cub.2020.12.006.
32. Matsumoto N, Kitani R, Maricle A, Mueller M, Kalinec F. Pivotal role of actin depolymerization in
the regulation of cochlear outer hair cell motility. *Biophys J*. Oct 6 2010;99(7):2067-76.
https://doi.org10.1016/j.bpj.2010.08.015.
33. Kobielak A, Fuchs E. Alpha-catenin: at the junction of intercellular adhesion and actin dynamics.
Nat Rev Mol Cell Biol. Aug 2004;5(8):614-25. https://doi.org10.1038/nrm1433.
34. Nunes FD, Lopez LN, Lin HW, et al. Distinct subdomain organization and molecular composition
of a tight junction with adherens junction features. *J Cell Sci*. Dec 1 2006;119(Pt 23):4819-27.
https://doi.org10.1242/jcs.03233.

 10. Figure Legends

Figure 1: Cross-sectional and top-down view of organ of Corti reveal cytoarchitecture of the

support cells and hair cells.

(A) Phalloidin labeling of F-actin (green) and Myo6 labeling of HCs (magenta) in a P1 sectioned

cochlea shows structural details of the F-actin cytoskeleton in the sensory epithelium along the

apico-basolateral axis. (A') Phalloidin labeling alone. White Arrow indicates the location of the

PCs.

(B) Top-down view of a Phalloidin-labeled P1 wholemount cochlea shows the overall

organization and location of the IHC, PC (black arrow) and the three rows of OHCs.

Figure 2: Significant remodeling of F-actin and E-cadherin occurs in the cochlear sensory

epithelium between late embryonic and early postnatal ages.

(A-A''') Phalloidin labeling of F-actin, and E-cadherin and pTyr immunolabeled proteins and

merged image in E18.5 cochleas. Dotted lines outline the PCs, asterisks denote the HCs.

(B-B''') Phalloidin labeling of F-actin, and E-cadherin and pTyr immunolabeled proteins and

merged image in P6 cochleas. Arrows indicate the mid-regions of IPC-OPC membranes that will

form the axial regions of the mature PCs.

(C-C''') Phalloidin labeling of F-actin, and E-cadherin and pTyr immunolabeled proteins and

merged image in P6 cochleas. Yellow arrow in C indicates actin bundle in IPC. White arrows in

C' and C'' indicate the apical regions of IPC-OPC membrane contacts. Dotted lines outline the

tunnel of Corti and the space of Nuel.

Nuclei are counter stained with DAPI (magenta).

Figure 3: Fine temporal resolution of the opening of the tunnel of Corti reveals novel

cytoskeletal details.

(A-A''') Phalloidin labeling of F-actin, and E-cadherin and pTyr immunolabeled proteins and

merged image in P7 cochleas harvested at 9 am. Arrows indicate the Arrows indicate the mid-

regions of IPC-OPC membrane contacts. A''' Inset- overlay of mid-region of IPC-OPC junction.

E-cadherin (green), pTyr (red) and Phalloidin (blue). White arrow indicates co-localization of the

three markers at the mid-region of the IPC-OPC junction. Red arrow indicates pTyr

accumulation at OPC-DC contact flanking Nuel space. Dotted lines denote the locations of

spaces that will form the tunnel of Corti and the space of Nuel

(B-B''') Phalloidin labeling of F-actin, and E-cadherin and pTyr immunolabeled proteins in P7

cochleas harvested at 1 pm. Dotted lines denote the locations of spaces that will form the tunnel

of Corti and the space of Nuel.

(C-C''') Phalloidin labeling of F-actin, and E-cadherin and pTyr immunolabeled proteins and

merged image in P7 cochleas harvested at 5 pm. Arrows indicate the apical junctional contacts

between the IPCs and the OPCs.

Asterisks denote the IHCs, and nuclei are counter stained with DAPI (magenta).

Figure 4: N-cadherin localization remains stable during postnatal ages.

(A-A') On P6, N-cadherin is localized to the membranes of the inner border cell, inner

phalangeal cell and the IHC. Merged image is counterstained with DAPI.

(B-D) On P7, N-cadherin is localized to membranes of the inner border, IHC and inner

phalangeal cell membranes remains stable as the tunnel of Corti is formed.

Asterisks denote the hair cells and actin is immunolabeled in magenta. Merged image is

counterstained with DAPI.

Figure 5: P-cadherin localization remains stable during postnatal ages.

(A-A') On P6, P-cadherin is localized to the membranes of the IPC (white arrow), and Hensen's

and outer sulcus cells (green arrow). Merged image is counterstained with DAPI.

(B-D) On P7, P-cadherin is localized to the membranes of the IPCs and the OPCs, Hensen's cells

and outer sulcus cells. Arrows indicate the apical portion of the IPC-OPC adherens junctions.

Asterisks denote the hair cells, and actin is immunolabeled in magenta. Merged image is

counterstained with DAPI.

Figure 6: Cofilin is expressed in the support cells of the sensory epithelium in the postnatal

cochlea.

(A-A'') On P6, Cofilin labeling is punctate in the cytoplasm and high at the PC membranes.

Cofilin is expressed in the SCs, but not the HCs. Merged image is counterstained with DAPI.

(B-B") On P7, before the tunnel of Corti forms, Cofilin is present in the mid-region of the IPC-

OPC junctions (arrow). Merged image is counterstained with DAPI.

(C-D) As the tunnel opens, Cofilin remains at the edge of the cell membrane that retracts to form

the tunnel of Corti. Merged image is counterstained with DAPI.

Asterisks denote the hair cells, and actin in immunolabeled in magenta. Yellow arrow indicates

cofilin enrichment.

	P7 (closed)	P7 (small ToC)	P7 (large ToC)
Phalloidin	A	B	C
E-Cadherin	A'	B'	C'
pTyr	A''	B''	C''
Merge	A'''	B'''	C'''

pTyr Phall E-cadh