Changes in cell adhesion properties are associated with the formation of fluid-filled spaces in an epithelium.

Holly J. Beaulac[1], Amy M. Segee[1] and Vidhya Munnamalai[1*]

[1]The Jackson Laboratory, Bar Harbor, ME 04609, USA

* Correspondence:

Vidhya Munnamalai

The Jackson Laboratory

Bar Harbor, ME 04609, USA

Email: vidhya.munnamalai@jax.org

Phone: 207-288-6916

Keywords: Cochlea, Support Cells, F-actin, Cell Adhesion, Cadherin, Tunnel of Corti

Running title: Modifications in E-cadherin cell contacts form the tunnel of Corti in the cochlea.

Funding: This work was supported by the National Institutes of Health (R21DC016376) to V.M.

Abstract (200 words)

The sensory epithelium of the cochlea, the organ of Corti, has complex cytoarchitecture

consisting of mechanosensory hair cells intercalated by support cells. The support cells provide

important trophic and structural support to the hair cells and must be stiff yet compliant enough

to modulate vibrations to the organ of Corti. Once the sensory cells are properly arranged, the

support cells undergo significant remodeling to help the sensory epithelium transition from a

simple epithelium into a structurally rigid epithelium with fluid-filled spaces. Cadherins are cell

adhesion molecules that are necessary for sorting and connecting cells in an intact epithelium. To

create the fluid-filled spaces, the cell adhesion properties of adjoining cell membranes between

cells must change to allow the formation of spaces within an epithelium. However, the dynamic

localization of cadherins has not been properly analyzed as these spaces are formed. There are

three cadherins that are reported to be expressed during the first postnatal week of development

when the tunnel of Corti forms in the cochlea. We characterized the dynamic localization of

cadherins that are associated with the cytoskeletal remodeling at the contacting membranes of

the inner and outer pillar cells flanking the tunnel of Corti.

1. Introduction

Epithelial morphogenesis requires complex cytoskeletal remodeling and maturation [1,2]

and is essential for proper organ formation and organ function. Cadherin-mediated cell adhesion

provides epithelial stability during morphogenesis by inducing cell-cell contact-mediated

mechanical coupling with the cellular actin cytoskeleton [2,3]. Cadherins are Ca^{2+}-dependent cell-

cell adhesion molecules that lead to the formation of adherens junctions during tissue

development [4]. Recruitment of cadherin molecules at the membrane is followed by cadherin

clustering and is stabilized by coupling to the actin cytoskeleton [5-7]. Maturation of adherens

junctions and strengthening of cell-cell adhesion further induces tension-dependent actin

remodeling and recruitment of actin binding proteins to the site of de novo actin synthesis [8,9].

In the cochlea, auditory function is also reliant on proper epithelial morphogenesis of the

sensory epithelium, the organ of Corti. The organ of Corti attains a highly specialized

organization of mechanosensory hair cells intercalated by support cells [10,11]. Thus far, most

studies have focused on the cytoskeleton and cytoskeletal remodelers in the sensory hair cells;

but recent studies have shown that the support cells have important electrophysiology, metabolic

and sensor properties that are necessary for proper auditory function [12-14]. The support cells also

serve as important support structures for the hair cells [15] and help to establish the overall

morphology and complex cytoarchitecture of the sensory epithelium. The organ of Corti must be

stable enough to withstand vibrational movements of the basilar membrane [14,16] and the

movement of fluid through the epithelium [17-19] to facilitate the propagation of sound information.

During sensory epithelial morphogenesis, fluid-filled spaces such as the tunnel of Corti, the Nuel

space and the outer tunnel are formed within the epithelium [20], which requires stabilization of the

cytoskeletal structure. Between late embryonic and early postnatal stages, the support cells

undergo significant dynamic morphological changes. However, few studies have uncovered the

spatiotemporal characteristics of cytoskeletal modifications on the cochlear sensory epithelium

and how the tunnel of Corti is formed during maturation. In this study we focus on the

cytoskeletal changes that the support cells undergo and the cell adhesion modification that occur

during a specific time in development when fluid-filled spaces form in the sensory epithelium,

particularly the pillar cells (PCs) that undergo the most drastic structural cytoskeletal remodeling

to form the tunnel of Corti. Since epithelial integrity in the cochlea is dependent on cadherins [21],

we also examine whether the cellular spatial localization of cadherins play a role in establishing

the integrity of the sensory epithelium with spaces interspersed within the cochlear sensory

epithelium.

Three cadherin molecules reported to be expressed in the mouse cochlea during the first

postnatal week are neural (N-), epithelial (E-), and placental (P-) cadherins [22-24]. The tunnel of

Corti forms between the inner pillar cells (IPC) and the outer pillar cells (OPCs), spaces of Nuel

and the outer tunnel [24]. A recent study suggested that the abnormal premature enlargement of the

tunnel of Corti was caused by loss of P-cadherin [22]; however, from wholemount analysis, P-

cadherin localization in the PCs appears to be apical. Cross-sectional views of cadherin

localization along the apico-basolateral axis during the narrow temporal window as the tunnel of

Corti forms would provide a clearer view of how E-cadherin or N- or P- cadherin localization

influence tunnel formation under normal developmental conditions. In this study, we analyze the

spatiotemporal localization of these three cadherins in the cochlea.

2.   Results and Discussion

At neonatal stages of the mouse cochlea, the organ of Corti has already established its
classical mosaic pattern of hair cells intercalated by support cells (Figure 1). There are two
classes of mechanosensory hair cells: one row of inner hair cells (IHCs) and three rows outer hair
cells (OHCs) (Figure 1A) that are innervated by afferents and efferent neurons enabling the
transmission of auditory information along the central auditory pathways [25,26]. The inner and
outer hair cells are segregated by the IPC and OPC support cells. The outer hair cells are
intercalated by a specific subtype of support cells known as Deiters' cells (DCs) (Figure 1A, A').
A prior study showed an increase in junctional E-cadherin in the apical junctions of the PCs in
wholemount mouse cochleas during the first postnatal week after birth [27]. Thus, E-cadherin
signaling is a strong candidate for PC remodeling. Since this study focuses on the support cells,
the cross-sectional view of the organ of Corti reveals structural details (Figure 1A, A') that are
difficult to appreciate in wholemount cochlear preparations where the stereocilia on hair cells
and the overall patterning of the sensory epithelium are the prime focus (Figure 1B).

A tunnel of Corti forms between the IPC and the OPC during the neonatal ages of
cochlear development; however, a close examination of the spatial changes of the F-actin
cytoskeleton and cell adhesion of the cochlea undergoing extensive remodeling and the tunnel of
Corti opening, has not been performed. Although the F-actin cytoskeleton establishes the
structure and shape of cells, its enrichment can vary in different cell types. In order to identify
the correct stage to observe the opening of the tunnel of Corti and characterize the associated
subcellular changes in the F-actin cytoskeletal organization, we first analyzed cochlear tissue on
E18.5, P6 and P8. On E18.5, although F-actin is present in all cells, Phalloidin labeling shows
that F-actin is specifically enriched at membranes contacting both the OHCs and the DCs (Figure
2A). Thus, we examined markers for cell adhesion along the OHC-DC membrane contacts.
Comparing F-actin with the cell adhesion protein, E-cadherin, we found a striking enrichment of
E-cadherin at OHC-DC membrane contacts (Figure 2B). On E18.5, E-cadherin was also
localized to the membrane of Hensen's cells (Figure 2B, green arrow). At this stage, there was
no marked enrichment of F-actin or E-cadherin in the PCs. Cell adhesion induces bi-directional
signaling that promotes the recruitment of phosphorylated actin binding proteins at their tyrosine
residues (pTyr), a sign of active actin assembly [28,29]. Immunolabeling for pTyr was undetectable
on E18.5, presumably because minimal remodeling of the F-actin cytoskeleton was required at
this stage of development (Figure 2C).

We next analyzed cochleas on P6. The tunnel of Corti was still not formed at this stage.

However, F-actin assembly is enhanced, particularly in the IPC and the IPC-OPC membrane

contacts (Figure 2D, arrow). E-cadherin was also localized at the adjoining IPC-OPC

membranes, but E-cadherin was concentrated mainly in the apical and mid-regions along the

apico-basolateral axis of the IPC-OPC membranes, indicating there was an increase in cell

adhesion contacts associated with the increase in F-actin (Figure 2E, arrow). However, there was

no apparent increase in the recruitment of pTyr actin binding proteins at the IPC-OPC

membranes (Figure 2F).

We then performed the same analysis in the P8 cochlea. At this stage, the tunnel of Corti

has fully formed (Figure 2G-I). The F-actin bundle that establishes the rigid support structure

within the IPC is much more developed (Figure 2G, green arrow) compared to the P6 stage.

However, by P8, F-actin, E-cadherin were no longer localized in the mid-regions of the IPC-

OPC contact membranes where the tunnel of Corti and Nuel space were formed. F-actin and E-

cadherin remain enriched in the adjoining apical IPC-OPC membranes (Figure 2G-H). Despite

the formation of the tunnel of Corti, the IPC and the OPC remain connected at their apical

junctions where E-cadherin remains localized (Figure 2H). P-Tyr labeling was not present in the

PCs at P8, suggesting that a critical window was missed. The time interval between P6 and P8

was too large to observe the transient cytoskeletal modifications associated with the formation of

the tunnel of Corti.

To capture finer temporal resolution of the opening of the tunnel of Corti, we analyzed P7

cochleas that were harvested at 3 timepoints throughout the day at 9 am, 1 pm and 5 pm (Figure

3). At the first timepoint, the tunnel had not yet opened (closed). We observed an accumulation

of F-actin at the mid-region of the IPC-OPC membranes (Figure 3A, arrow) along their apico-

basolateral axis, which will go onto to become the axial regions of the PCs that flank the tunnel

of Corti. Coincidently, we also saw an increase in E-cadherin cell contacts (Figure 3B, arrow).

and pTyr labeling (Figure 3C, arrow). At this stage, F-actin, E-cadherin and pTyr co-localize at

the mid-region of the IPC-OPC membranes (Figure 3C' inset, arrow). This is a notable

enrichment compared to the P6 when the tunnel was also closed but no pTyr labeling was

detectable (Figure 2D-F). P-Tyr labeling was comparatively less pronounced at the apical

junction (Figure 3C), which suggests that there comparatively less active F-actin assembly on the

apical junctions of the PCs. There was less also reduced F-actin, E-cadherin and pTyr in the

basolateral end of the PCs (Figure 3B-C). We also observed a notable increase of pTyr when the

Nuel space has a very narrow opening, which suggests a significant recruitment of actin binding

proteins during initiation of Nuel space formation (Figure 3C, red arrow). At the mid-timepoint

on P7, the tunnel of Corti was open but small (small ToC). F-actin and E-cadherin have already

decreased their expression in the mid-axial region of the IPC-OPC membranes but remain

enriched at the apical junction (Figure 3D-E). Interestingly, pTyr labeling remained high at the

edges of cell membranes (Figure 3F). This suggests that these sites remain enriched for actin

binding proteins which help remodel cellular structure of the tunnel. In the final timepoint on P7,

the tunnel was significantly larger (large ToC). F-actin and E-cadherin remained at the apical

regions of the PCs (Figure 3G-H); while pTyr labeling was relatively minimal at the apical

junctions (Figure 3I). This suggests that there is a greater recruitment of pTyr-labeled actin

binding proteins where the cochlear spaces are forming and not at stable contacts. The variable

localization of E-cadherin on the adjoining IPC-OPC membranes along the apico-basolateral axis

suggests that a reduction in cell adhesion precedes the opening of the tunnel of Corti at the mid-

region of the PC membranes and is accompanied by a large recruitment of pTyr-labeled proteins.

This data also show that P7 is critical timepoint when the support cells in the sensory epithelium

undergo significant cellular remodeling to attain its complex cytoarchitecture. Conversely,

stabilized F-actin will require less recruitment of actin binding proteins as there are no sites for

de novo F-actin synthesis e.g., in the apical junctions of the PCs.

N-cadherin is another cadherin known to be expressed in the developing cochlea [23]. We

examined whether N-cadherin is spatially positioned to influence the opening of the spaces in the

cochlear sensory epithelium. On P6, N-cadherin is visible in the inner hair cell (IHC), inner

phalangeal and inner border cell, but is clearly absent in the PCs and the DCs (Figure 4A). This

expression pattern remains consistent in P7 cochleas at the same three timepoints when the

tunnel of Corti was still closed (Figure 4B), has a small opening (Figure 4C) and when the tunnel

was fully open (Figure 4D). Thus, the stable expression of N-cadherin and its consistent absence

of expression in the PCs suggests that N-cadherins do not influence the opening of the tunnel of

Corti.

A recent study showed that P-cadherin is expressed in the PCs of the cochlea during early

postnatal stages. Upon Thyroxine administration, P-cadherin was translocated from the PC

membranes to the cytoplasm, thereby reducing cell-cell adhesion [22]. The study analyzed

wholemount cochleas; thus, the expression of P-cadherin along the entire apico-basolateral axis

of PCs was unclear. To determine its spatial localization, we analyzed P-cadherin expression. On

P6, P-cadherin was localized in the IPCs, but enriched at the top apical end of the IPC (Figure

5A, white arrow). In addition to the PCs, we found P-cadherin was also expressed in the

membranes of the Hensen's cells and the cells in the outer sulcus region (Figure 5A, green

arrow). The expression of P-cadherin was much more substantial in these lateral non-sensory

cells compared with E-cadherin. On P7, while the tunnel of Corti was still closed, P-cadherin

remained enriched in the apical portion of the PCs, the Hensen's cells and the outer sulcus cells

(Figure 5B). The apical spatial localization remained unchanged while the tunnel of Corti opened

(Figure 5C-D). The expression of P-cadherin remained stable in the Hensen's and outer sulcus

cells as well.

We found that there was significant recruitment of pTyr labeled proteins co-localized

along with F-actin and E-cadherin at IPC-OPC membrane prior to the opening of the tunnel of

Corti (Figure 3). We postulated that in preparation for the tunnel opening, actin binding proteins

are recruited to facilitate the remodeling of F-actin in the PCs. Thus, we examined the expression

of Cofilin, an F-actin severing protein, on P6 and the three stages of tunnel of Corti formation on

P7 (Figure 6). Previous studies showed cofilin labeling in the HCs [30,31], but Cofilin localization

in the SCs has not been reported. Thus, we were surprised to find a much higher punctate

labeling of Cofilin in the SCs compared to the HCs. Relative to P6 (Figure 6A), immunolabeling

shows an increase in Cofilin localization in the mid-region of the IPC-OPC junction of the closed

P7 closed tunnel of Corti (Figure 6B, arrow). This suggests that during this critical period,

Cofilin recruitment is associated with actin remodeling and will aid in disrupting cadherin

localization and consequently, disrupting cell adhesion at the mid-region of the mid-region of the

IPC-OPC walls.

Although P-cadherin was the first cell adhesion protein to be expressed in the PCs and

continued to be expressed in the apical portion on P7, the stable P-cadherin expression is

unlikely to influence the opening of the tunnel of Corti under normal developmental conditions.

On the other hand, the dynamic and transient localization of E-cadherin in the mid- regions of the IPC-OPC membranes, and its subsequent downregulation preceding the opening of the tunnel of Corti makes the dynamic downregulation E-cadherin a much more likely mechanism for opening the tunnel under normal developmental conditions. However, it is possible that in a disease context, the loss of P-cadherin can exacerbate the tunnel opening and final size of the tunnel, disrupting the overall stability of the sensory epithelium. Perhaps both E-cadherin and P-cadherin trafficking are impacted when the cytoarchitecture of the sensory epithelium is disrupted under disease conditions. Thus, follow up studies are needed to validate this. In this study, we show evidence that significant actin remodeling takes place in the SCs during P6- P8 and paves way for future investigations on actin assembly in the SCs on which little focus has been placed thus far. Here we closely analyze the cell adhesion characteristics associated with the opening of the tunnel of Corti because they are expressed at much higher levels. We predict that the same decrease in cadherin trafficking will promote the formation of the Nuel space and the outer tunnel.

3. Experimental Procedures

3.1 Mice

Time-mated Swiss Webster dams (Envigo, Indianapolis, IN, USA) were denoted as embryonic day (E)0.5 on the first day a plug was observed. E18.5 embryos, and postnatal day (P)6-P8 pups were euthanized by decapitation, in accordance with the Institutional Animal Care and Use Committee (IACUC) guidelines at The Jackson Laboratory, Bar Harbor, and heads or ears were harvested and fixed in 4% paraformaldehyde (PFA) (Electron Microscopy Sciences, Hatfield, PA, USA).

3.2 Histology

After fixation, E18.5 heads were bisected and processed through a 10%, 20%, and 30% sucrose gradient series, embedded with TFM and cryofrozen. For postnatal studies, cochleas were extracted on P6, P7 and P8, and fixed in 4% PFA for three hours at 4°C, immersed in a 30% sucrose and transferred to a 1:1 solution of 30% sucrose: Tissue Freezing Medium (TFM) (General Data Healthcare, Cincinnati, OH, USA) before final embedding in 100% TFM and

cryofrozen. Tissues were cryosectioned at 12 μm for histology at all ages. To capture the

opening of the tunnel of Corti (ToC) on P7, inner ears were harvested and fixed at 9 am, 1 pm

and 5 pm.

3.3 Immunofluorescence

Tissues were washed in PBS/0.5% Triton and blocked at for at room temperature (RT) 1 hour

with 2% Donkey serum (Jackson ImmunoResearch, West Grove, PA, USA) in PBS/0.5% Triton,

followed by overnight primary antibody incubation at 4°C. Primary antibodies included: rat anti-

E-cadherin (1:250, AB11512, Abcam, Waltham, MA, USA), sheep anti- N-cadherin (1:250,

AF6426, R&D Systems, Minneapolis, MN, USA), goat anti- P-cadherin (1:250, AF761, R&D

Systems), mouse anti-phosphoTyrosine (1:250, zms1b282, Sigma-Aldrich), anti-Cofilin (1:250,

5175, Cell Signaling Technology, Danvers, MA, USA). For N-cadherin, P-cadherin and Cofilin

immunolabeling, antigen retrieval was performed prior to primary antibody incubation with a 10

mM sodium citrate/0.05% Tween20 buffer at 6.0 pH for 15 min at 99°C. Since phalloidin

labeling of F-actin is incompatible with antigen retrieval, we co-labeled with an actin antibody

(1:250, MA5-11869, Invitrogen). Alexa-conjugated secondary antibodies (Invitrogen) were

incubated for 2 hours at RT. Nuclei were counter stained with DAPI (Abcam, Waltham, MA,

USA) and tissues were mounted with Fluoromount G mounting medium (Life Technologies

Corporation, Carlsbad, CA, USA).

3.4 Data Acquisition and Image processing

60X magnification images were acquired on a Zeiss LSM 800 confocal microscope (ZEISS,

Oberkochen, BW, Germany) in The Jackson Laboratory microscopy core. Z-stacks were

analyzed using the FIJI software (NIH). Data were collected on N=6 cochleas for each time

point. TIFF images were processed in Adobe Photoshop and assembled in Adobe Illustrator

(Adobe, San Jose, CA, USA).

4. Acknowledgements

The authors acknowledge the Microscopy Core at The Jackson Laboratory for equipment maintenance and upkeep, and assistance. The authors thank members of the Munnamalai lab for providing critical feedback on the manuscript.

5. Conflict of Interest

The authors declare no conflicts of interest or competing financial interests.

6. Author Contributions

H.J.B., study conceptualization and design, development of methodology, acquisition, data analysis, writing and revision of the manuscript. A.M.S., study design, acquisition, data analysis, and revision of the manuscript. V.M., funding, material support, study conceptualization and design, development of methodology, acquisition, analysis, and interpretation of data; writing and revision of the manuscript. All authors read and approved the final manuscript.

7. Ethics Approval

All experiments were performed under IACUC protocol 18013: PI Vidhya Munnamalai, in compliance with the U.S. Department of Health and Human Services and reviewed by The Jackson Laboratory Institutional Animal Care and Use Committee.

8. Data Availability

The datasets used and/or analyzed during the current study are available from the corresponding author on request.

9. References

1. Schock F, Perrimon N. Molecular mechanisms of epithelial morphogenesis. *Annu Rev Cell Dev Biol*. 2002;18:463-93. https://doi.org10.1146/annurev.cellbio.18.022602.131838.
2. Clarke DN, Martin AC. Actin-based force generation and cell adhesion in tissue morphogenesis. *Curr Biol*. May 24 2021;31(10):R667-R680. https://doi.org10.1016/j.cub.2021.03.031.
3. Samarage CR, White MD, Alvarez YD, et al. Cortical Tension Allocates the First Inner Cells of the Mammalian Embryo. *Dev Cell*. Aug 24 2015;34(4):435-47. https://doi.org10.1016/j.devcel.2015.07.004.
4. Leckband DE, de Rooij J. Cadherin adhesion and mechanotransduction. *Annu Rev Cell Dev Biol*. 2014;30:291-315. https://doi.org10.1146/annurev-cellbio-100913-013212.
5. Mege RM, Gavard J, Lambert M. Regulation of cell-cell junctions by the cytoskeleton. *Curr Opin Cell Biol*. Oct 2006;18(5):541-8. https://doi.org10.1016/j.ceb.2006.08.004.
6. Hong S, Troyanovsky RB, Troyanovsky SM. Binding to F-actin guides cadherin cluster assembly, stability, and movement. *J Cell Biol*. Apr 1 2013;201(1):131-43. https://doi.org10.1083/jcb.201211054.
7. Priest AV, Shafraz O, Sivasankar S. Biophysical basis of cadherin mediated cell-cell adhesion. *Exp Cell Res*. Sep 1 2017;358(1):10-13. https://doi.org10.1016/j.yexcr.2017.03.015.
8. Watabe-Uchida M, Uchida N, Imamura Y, et al. alpha-Catenin-vinculin interaction functions to organize the apical junctional complex in epithelial cells. *J Cell Biol*. Aug 10 1998;142(3):847-57. https://doi.org10.1083/jcb.142.3.847.
9. Yu Q, Holmes WR, Thiery JP, Luwor RB, Rajagopal V. Cortical tension initiates the positive feedback loop between cadherin and F-actin. *Biophys J*. Feb 15 2022;121(4):596-606. https://doi.org10.1016/j.bpj.2022.01.006.
10. Zine A, Van De Water TR, de Ribaupierre F. Notch signaling regulates the pattern of auditory hair cell differentiation in mammals. *Development*. Aug 2000;127(15):3373-83. https://doi.org10.1242/dev.127.15.3373.
11. McNeill H. Planar cell polarity: keeping hairs straight is not so simple. *Cold Spring Harb Perspect Biol*. Feb 2010;2(2):a003376. https://doi.org10.1101/cshperspect.a003376.
12. Chen T, Rohacek AM, Caporizzo M, et al. Cochlear supporting cells require GAS2 for cytoskeletal architecture and hearing. *Dev Cell*. May 17 2021;56(10):1526-1540 e7. https://doi.org10.1016/j.devcel.2021.04.017.
13. Velez-Ortega AC, Stepanyan R, Edelmann SE, et al. TRPA1 activation in non-sensory supporting cells contributes to regulation of cochlear sensitivity after acoustic trauma. *Nat Commun*. Jun 30 2023;14(1):3871. https://doi.org10.1038/s41467-023-39589-w.
14. Zhou W, Jabeen T, Sabha S, Becker J, Nam JH. Deiters Cells Act as Mechanical Equalizers for Outer Hair Cells. *J Neurosci*. Nov 2 2022;42(44):8361-8372. https://doi.org10.1523/JNEUROSCI.2417-21.2022.
15. Wan G, Corfas G, Stone JS. Inner ear supporting cells: rethinking the silent majority. *Semin Cell Dev Biol*. May 2013;24(5):448-59. https://doi.org10.1016/j.semcdb.2013.03.009.

16. Reichenbach T, Hudspeth AJ. The physics of hearing: fluid mechanics and the active process of the inner ear. *Rep Prog Phys*. Jul 2014;77(7):076601. https://doi.org10.1088/0034-4885/77/7/076601.

17. Zagadou BF, Barbone PE, Mountain DC. Significance of the Microfluidic Flow Inside the Organ of Corti. *J Biomech Eng*. Aug 1 2020;142(8). https://doi.org10.1115/1.4046637.

18. Wangemann P. Supporting sensory transduction: cochlear fluid homeostasis and the endocochlear potential. *J Physiol*. Oct 1 2006;576(Pt 1):11-21. https://doi.org10.1113/jphysiol.2006.112888.

19. Meenderink SWF, Dong W. Organ of Corti vibrations are dominated by longitudinal motion in vivo. *Commun Biol*. Nov 24 2022;5(1):1285. https://doi.org10.1038/s42003-022-04234-7.

20. Ulfendahl M, Scarfone E, Flock A, Le Calvez S, Conradi P. Perilymphatic fluid compartments and intercellular spaces of the inner ear and the organ of Corti. *Neuroimage*. Sep 2000;12(3):307-13. https://doi.org10.1006/nimg.2000.0617.

21. Trowe MO, Maier H, Petry M, Schweizer M, Schuster-Gossler K, Kispert A. Impaired stria vascularis integrity upon loss of E-cadherin in basal cells. *Dev Biol*. Nov 1 2011;359(1):95-107. https://doi.org10.1016/j.ydbio.2011.08.030.

22. Zhang H, Xie L, Chen S, Qiu Y, Sun Y, Kong W. Thyroxine Regulates the Opening of the Organ of Corti through Affecting P-Cadherin and Acetylated Microtubule. *Int J Mol Sci*. Nov 1 2022;23(21). https://doi.org10.3390/ijms232113339.

23. Chacon-Heszele MF, Ren D, Reynolds AB, Chi F, Chen P. Regulation of cochlear convergent extension by the vertebrate planar cell polarity pathway is dependent on p120-catenin. *Development*. Mar 2012;139(5):968-78. https://doi.org10.1242/dev.065326.

24. Whitlon DS. E-cadherin in the mature and developing organ of Corti of the mouse. *J Neurocytol*. Dec 1993;22(12):1030-8. https://doi.org10.1007/BF01235747.

25. Munnamalai V, Fekete DM. The acquisition of positional information across the radial axis of the cochlea. *Dev Dyn*. Mar 2020;249(3):281-297. https://doi.org10.1002/dvdy.118.

26. Di Bonito M, Studer M. Cellular and Molecular Underpinnings of Neuronal Assembly in the Central Auditory System during Mouse Development. *Front Neural Circuits*. 2017;11:18. https://doi.org10.3389/fncir.2017.00018.

27. Luo WW, Wang XW, Ma R, et al. Junctional E-cadherin/p120-catenin Is Correlated with the Absence of Supporting Cells to Hair Cells Conversion in Postnatal Mice Cochleae. *Front Mol Neurosci*. 2018;11:20. https://doi.org10.3389/fnmol.2018.00020.

28. Andl CD, Rustgi AK. No one-way street: cross-talk between e-cadherin and receptor tyrosine kinase (RTK) signaling: a mechanism to regulate RTK activity. *Cancer Biol Ther*. Jan 2005;4(1):28-31. https://doi.org10.4161/cbt.4.1.1431.

29. Brunton VG, MacPherson IR, Frame MC. Cell adhesion receptors, tyrosine kinases and actin modulators: a complex three-way circuitry. *Biochim Biophys Acta*. Jul 5 2004;1692(2-3):121-44. https://doi.org10.1016/j.bbamcr.2004.04.010.

30. McGrath J, Tung CY, Liao X, et al. Actin at stereocilia tips is regulated by mechanotransduction and ADF/cofilin. *Curr Biol*. Mar 22 2021;31(6):1141-1153 e7. https://doi.org10.1016/j.cub.2020.12.006.

372   31.   Matsumoto N, Kitani R, Maricle A, Mueller M, Kalinec F. Pivotal role of actin
373   depolymerization in the regulation of cochlear outer hair cell motility. *Biophys J*. Oct 6
374   2010;99(7):2067-76. https://doi.org10.1016/j.bpj.2010.08.015.

10. Figure Legends

Figure 1: Cross-sectional and top-down view of organ of Corti reveal cytoarchitecture of the support cells and hair cells.

(A) Phalloidin labeling of F-actin (green) and Myo6 labeling of HCs (magenta) in a P1 sectioned cochlea shows structural details of the F-actin cytoskeleton in the sensory epithelium. (A') Phalloidin labeling alone. White Arrow indicates the location of the PCs.

(B) Top-down view of a Phalloidin-labeled P1 wholemount cochlea shows the overall organization and location of the IHC, PC (black arrow) and the three rows of OHCs.

Figure 2: Significant remodeling of F-actin and E-cadherin occurs in the cochlear sensory epithelium between late embryonic and early postnatal ages.

(A-C) Phalloidin labeling of F-actin, and E-cadherin and pTyr immunolabeled proteins in E18.5 cochleas. Dotted lines outline the PCs, asterisks denote the HCs.

(D-E) Phalloidin labeling of F-actin, and E-cadherin and pTyr immunolabeled proteins in P6 cochleas. Arrows indicate the mid-regions of IPC-OPC membranes that will become the axial regions of the mature PCs.

(G-I) Phalloidin labeling of F-actin, and E-cadherin and pTyr immunolabeled proteins in P6 cochleas. Arrows indicate the apical regions of IPC-OPC membrane contacts. Dotted lines outline the tunnel of Corti and the space of Nuel.

Nuclei are counter stained with DAPI (magenta).

Figure 3: Fine temporal resolution of the opening of the tunnel of Corti reveals novel cytoskeletal details.

(A-C) Phalloidin labeling of F-actin, and E-cadherin and pTyr immunolabeled proteins in P7 cochleas harvested at 9 am. Arrows indicate the Arrows indicate the mid-regions of IPC-OPC membrane contacts. C' Inset- overlay of mid-region of IPC-OPC junction. E-cadherin (green), pTyr (red) and Phalloidin (blue). White arrow indicates co-localization of the three markers at the mid-region of the IPC-OPC junction. Red arrow indicates pTyr accumulation at OPC-DC contact flanking Nuel space. Dotted lines denote the locations of spaces that will form the tunnel of Corti and the space of Nuel

(D-F) Phalloidin labeling of F-actin, and E-cadherin and pTyr immunolabeled proteins in P7

cochleas harvested at 1 pm. Dotted lines denote the locations of spaces that will form the tunnel

of Corti and the space of Nuel.

(G-I) Phalloidin labeling of F-actin, and E-cadherin and pTyr immunolabeled proteins in P7

cochleas harvested at 5 pm. Arrows indicate the apical junctional contacts between the IPCs and

the OPCs.

Asterisks denote the IHCs, and nuclei are counter stained with DAPI (magenta).

Figure 4: N-cadherin localization remains stable from embryonic to postnatal ages.

(A) On P6, N-cadherin is localized to the membranes of the inner border cell, inner phalangeal

cell and the IHC.

(B-C) On P7, N-cadherin is localized to membranes of the inner border, IHC and inner

phalangeal cell membranes remains stable as the tunnel of Corti is formed.

Asterisks denote the hair cells and actin is immunolabeled in magenta.

Figure 5: P-cadherin localization remains stable from embryonic to postnatal ages.

(A) On P6, P-cadherin is localized to the membranes of the IPC (white arrow), and Hensen's and

outer sulcus cells (green arrow).

(B-C) On P7, P-cadherin is localized to the membranes of the IPCs and the OPCs, Hensen's cells

and outer sulcus cells. Arrows indicate the apical portion of the IPC-OPC adherens junctions.

Asterisks denote the hair cells, and actin is immunolabeled in magenta.

Figure 6: Cofilin is enriched in the support cells of the sensory epithelium in the postnatal

cochlea.

(A) On P6, Cofilin labeling is punctate in the cytoplasm and slightly enriched at the PC

membranes. Cofilin is expressed in the SCs, but not the HCs.

(B) On P7, before the tunnel of Corti forms, Cofilin is selectively enriched in the mid-region of

the IPC-OPC junctions.

(C-D) As the tunnel begins to open, Cofilin remains enriched at the edge of the cell membrane

that retracts to form the tunnel of Corti.

Arrows indicate the apical regions of the pillar cells. Asterisks denote the hair cells, and actin in

immunolabeled in magenta. Yellow arrow indicates cofilin enrichment.

A P1
IHC OHC
Phalloidin
(F-actin)
Hensen's
IPC OPC DCs Outer
sulcus

A' F-actin

B IHC OHC
F-actin P1