Phosphorylation-induced rewiring of intermolecular interactions with LC3B: local and long-range effects on short linear motifs

Oana Nicoleta Antonescu[1†], Mattia Utichi[1,2†], Valentina Sora[1,2,3], Matteo Tiberti[1], Emiliano Maiani[1,4], Matteo Lambrughi[1], Elena Papaleo[1,2]*

[1]*Cancer Structural Biology, Danish Cancer Society Research Center, Strandboulevarden 49, 2100, Copenhagen, Denmark*

[2]*Cancer System Biology, Section of Bioinformatics, Health and Technology Department, Technical University of Denmark, Lyngby, Denmark*

*Corresponding author: elenap@cancer.dk, elpap@dtu.dk

[3] Current affiliation: University of Copenhagen, Department of Computer Science, Machine Learning Section, 2100 Copenhagen, Denmark
[4] Current affiliation UniCamillus - Saint Camillus International University of Health and Medical Sciences, 00131, Roma, Italy

[†] The two authors contributed equally to this work

Keywords: short linear motifs, LC3B, molecular dynamics, binding free energies, post-translational modifications, mutations

Abstract

Short Linear Motifs (SLiMs) are pivotal in mediating interactions between intrinsically disordered proteins and their binding partners. SLiMs exhibit sequence degeneracy and undergo regulation by post-translational modifications, including phosphorylation. These modifications can either augment or attenuate the binding affinities for a SliM and its binding partner. In this study, we set out to integrate biomolecular simulations, in silico high-throughput mutational scans, and biophysical measurements to elucidate the structural details of phospho-regulation in a class of SLiMs crucial for autophagy, known as LC3 interacting regions (LIRs). As a case study, we investigated the interaction between optineurin and LC3B, a system extensively characterized by experiments. We confirmed the robustness of our computational approach through the comparison with the available experimental data. Furthermore, we unveiled the previously unexplored role of the N-terminal flanking region upstream of the LIR core motif. Our study offers an atom-level perspective on the structural mechanisms and conformational alterations induced by phosphorylation in optineurin and LC3B recognition. Additionally, we assessed the impact of disease-related mutations on optineurin, accounting for different functional features.

Notably, we established an approach based on Microfluidic Diffusional Sizing to investigate the binding affinity of SLiMs to target proteins, enabling precise measurements of the dissociation constant for a selection of variants from the in silico mutational screening. Overall, our work provides a versatile toolkit to characterize other LIR-containing proteins and their modulation by phosphorylation or other phospho-regulated SLiMs, thereby advancing the understanding of important cellular processes.

Introduction

Several proteins harbor regions that are intrinsically disordered[1], which support networks of protein-protein interactions[2,3]. Intrinsically disordered proteins or regions feature diverse interaction modes with their binding partners, including folding-upon-binding, conformational selection, formation of dynamic complexes, retained disorder in the bound state or a combination of different mechanisms at the same time[4–8].

Thanks to their pliability, intrinsically disordered proteins are often implicated in multiple biological processes[2,9]. Furthermore, post-translational modifications can alter the interactome of disordered proteins[10–13].

Disordered proteins can rely on short linear motifs (SLiMs) to mediate interactions with their biological partners[4,14–16]. SLiMs are short and degenerated sequence stretches formed by few highly conserved residues (i.e., the SLIM core motif) interspersed with less conserved positions[15]. Regions flanking the core motifs and post-translational modifications, such as phosphorylation, can play a crucial role in modulating binding specificity and affinity[17–20]. The contributions of flanking regions and phosphorylations to the binding mechanisms of SLiMs are often challenging to study both experimentally and computationally due to the transient nature of the interactions. Structure-based methods have the potential to provide an accurate understanding of SLiM modulation by phosphorylation or flanking regions, accompanied by a mechanistic understanding of their interactions with biological partners[19,21].

SLiMs involved in the cellular clearance mechanism known as autophagy represent an intriguing case study due to their central role in the interaction with autophagy receptors and adaptors and a large variety of binding modes[19,22,23]. The ATG8 (LC3/GABARAP) family of proteins has a key role in both non-selective and selective degradation through autophagy. The LC3/GABARAP proteins interact with other proteins through the recognition of a SLiM, i.e., the LC3-interacting region (LIR), which binds to two main hydrophobic pockets (HP) of LC3 or GABARAP structures, known as HP1 and HP2 pockets[19,22,23]. The LIR motif is often associated with autophagy receptors that mediate the simultaneous binding of components of the autophagy machinery and cytosolic cargo[24–26], therefore representing a hub in the autophagy process. LIR amino acid sequences vary between proteins and share a core motif [W/F/Y]-x-x-[L/I/V], whereas other residues of the LIR could contribute to the specificity toward different LC3/GABARAP family members[27–29].

Among the known LIR-containing proteins, optineurin (OPTN) acts as a selective autophagy receptor and adaptor[30–32]. OPTN also regulates several other cellular processes associated with the trafficking of membrane-delivered cargo[30], a trait common to scaffolding proteins. OPTN is a multi-domain protein (**Figure 1A**), and it harbors a LIR motif (**Figure 1B)**, which

mediates the interaction with LC3 and GABARAP proteins. The LIR motif allows OPTN to interact with LC3 and GABARAP proteins[33,34]. The LIR motif of OPTN has a role in mitophagy amplification rather than in selectivity, suggesting a general mechanism in which LC3 proteins recruit autophagy factors to drive the autophagosome growth and amplify selective autophagy in an LC3-dependent feedback loop[35]. OPTN mutations or level alterations are also associated with several different pathologies, including glaucoma, amyotrophic lateral sclerosis, Paget's disease of bone, Crohn's disease and cancer [36–38].

In light of the above observations, it is clear that the OPTN function needs to be tightly orchestrated and modulated at the post-translational level. In this context, OPTN undergoes phosphorylation, and a subset of these phosphorylations are relevant in the selective autophagy process known as xenophagy, namely the degradation of intracellular pathogens[36]. In particular, OPTN is phosphorylated by TANK-binding kinase 1 (TBK1) on S177 in its LIR domain, which results in its increased binding affinity to LC3B, enhancing the autophagy-mediated clearance of ubiquitin-coated *Salmonella enterica*[33,34].

We here show how Alphafold-based models for interaction[39,40], biomolecular simulations [41–43] and other structure-based computational approaches[44–47], together with experimental biophysical data based on Microfluidic Diffusional Sizing[48,49] are effective to study the modulation by phosphorylation of SLiMs focusing on the OPTN$_{LIR}$-LC3B complex and its disease-related mutations. In a broader context, our work provides a computational framework with in vitro experimental validation to apply to other LIRs or SLiMs regulated by phosphorylation.

Materials and Methods

Modeling of constructs for MD simulations

We used the NMR structure of OPTN LIR (residues 169-185, NpSpSGpSpSEDpSFVEIRMAE) in its pentaphosphorylated form (p5) in complex with LC3B (residues 1-119, PDB entry 2LUE[34], first conformer) as a starting structure for four MD replicates of the LC3B$_{1-119}$-p5OPTN$_{168-185}$ complex. The structure includes an N-terminal expression tag (1-GAMG-4). Thus, we mutated it into the canonical sequence of LC3B from UniProtKB (1-MPSE-4) and added the missing C-terminal residue G120 (LC3B$_{1-120}$-p5OPTN$_{169-185}$). We then model its corresponding wild-type variant (LC3B$_{1-120}$-OPTN$_{169-185}$) by retro-mutating pS170, pS171, pS173, pS174, and pS177 to non-phosphorylated serines, using the mutagenesis tool of PyMOL (http://www.pymol.org/pymol). Furthermore, we modeled a single phosphorylated variant (LC3B$_{1-120}$-p1OPTN$_{169-185}$) in which we kept only pS177.

We also used AlphaFold-multimer[39] to generate a model of the full-length complex of LC3B and OPTN and retained the first five ranked models (out of 25 models) characterized by low Predicted Aligned Error (PAE) scores. We used the AlphaFold database (version available 25/01/2023). We then retained the first ranked model to generate a trimmed version of the complex spanning the OPTN region 158-191 ($LC3B_{1-120}$-$OPTN_{158-191}$) and remodel the corresponding phosphorylated variants ($LC3B_{1-120}$-$p1OPTN_{158-191}$ and $LC3B_{1-120}$-$p5OPTN_{158-191}$) with ViennaPTM[50].

Molecular Dynamics (MD) simulations

We performed all-atom MD simulations in explicit solvent for each variant and construct (**Table S1**) using GROMACS[51]. We employed the CHARMM22* protein force field[52], which we recently shown as a good force field for LC3B[53]. We used the TIPS3P water model[54] and SP2 parameters for phosphorylated serine residues adapted from[55]. We used periodic boundary conditions and a dodecahedral box with a minimum distance between the protein surface and the box edges of 15 Å. We capped the N- and C-terminal groups of both LC3B (NH_3+ and COOH, respectively) and the LIR peptides (NH_2 and COOH). We predicted the His27, His57, and His86 tautomeric states with H++, Propka software[56], accompanied by visual inspection. We used the Nε2-H tautomeric state for all the histidine residues in the simulation. We equilibrated the systems before the production run minimization, solvent equilibration, pressurization, and thermalization. We performed productive MD simulations at 298 K with the canonical ensemble, mimicking a concentration of 150 mM NaCl and including counterions to neutralize the system.

The MD simulations of $LC3B_{1-119}$-$p5OPTN_{169-185}$ were used to compare the MD results with available NMR data to evaluate the sampling quality achieved with the selected force field since they included the same construct used for NMR. The simulations with $LC3B_{1-120}$-$OPTN_{158-191}$ and $LC3B_{1-120}$-$OPTN_{168-185}$ were used for the analyses to understand the mechanisms induced by phosphorylations or the impact of disease-related mutations.

During pre-processing of the trajectories, we verified that the minimal distance between each protein complex and its image was sufficient (i.e., higher than 15 Å) to rule out artifacts due to contacts between the protein and the corresponding image in the periodic boundary conditions. We carried out the simulations without embedding LC3B in a lipidic membrane in light of our previous study where we showed that the majority of the properties of interest were similar comparing MD simulations of LC3B in solution or associated with the membrane[53]. The details on the constructs used for MD simulations are reported in **Figure 1** and **Table S1**.

Comparison of MD replicates with NMR data

We selected 1000 frames (equally spaced in time) for the replicate1 of the LC3B$_{1-119}$-p5OPTN$_{169-185}$ to predict chemical shifts from the MD simulations and compare them to the corresponding experimental data (Biological Magnetic Resonance Bank (BMRB), entry 18518). To this goal, we applied the delta_CS Python-based pipeline[57] (https://github.com/ELELAB/delta_cs). This includes the calculation of backbone, β-carbon chemical shifts, along with chemical shifts from side-chain hydrogens for the MD ensemble with PPM [58] and the associated squared errors between the predicted and the corresponding experimentally measured chemical shifts. Furthermore, the pipeline includes CH3Shift[59] to calculate chemical shifts of side-chain methyl carbon atoms (i.e., 13C shifts). It then compares them to the experimental side-chain chemical shifts. We used a reduced χ^2 (χ^2_{red}) metric as recently applied to another case study[60]. The square deviation between the calculated and experimental values was normalized by the variance of the chemical shift predictor for each type of chemical shift and the total number of chemical shifts. Therefore, low χ^2_{red} values indicate a good agreement between experimental and predicted chemical shifts.

Protein flexibility

We calculated the average Root Mean Square Fluctuation (RMSF) for each Cα atom in the central part of the LIR peptides (residues 170-184). We did the calculations averaging the RMSF over 10-ns or 100-ns time windows as a probe of the LIR flexibility in the LC3B LIR-binding pocket and estimated the associated standard errors.

Solvent-accessible surface

We calculated the side-chain solvent-accessible surface area (SASA) for the side-chain atoms of the residues of the LIR core motifs that bind to HP1 (F178) or HP2 (I181) pockets of LC3B. We estimated the SASA for the LC3B residues belonging to HP1 (D19, I23, P32, I34, K51, L53, and F108) and HP2 (F52, V54, P55, L63, I66, and I67) pockets[19] from the MD simulations of the different variants. We also included in the analysis the LDK tripeptide (L47, D48, and K49) and the ubiquitin patch of LC3B (R10, R11, K49, and K50), which are located in the proximity of the HP1 pocket and contribute LIR-specific interactions[19,25]. We used NACCESS[61] to calculate the relative SASA for each residue using frames extracted every 20 ps from the MD trajectories.

Secondary structure propensity of the LIR peptides

We employed the DSSP dictionary[62] to estimate the different classes of secondary structures for each residue of the OPTN peptides in the MD simulations. DSSP defines eight classes of

secondary structure elements based on an estimate of patterns of hydrogen bonds: 3.10 helices, α-helices, π-helices, β-sheets, β-bridges, turns, bends, and coil. For each residue of interest, we calculated the occurrence of each structure class in the MD simulation.

Analysis of electrostatic interactions and hydrogen bonds

We used PyInteraph2[63] to calculate electrostatic interactions in the form of hydrogen bonds and salt bridges. For salt bridges, all the distances between atom pairs belonging to charged moieties of two oppositely charged residues were calculated, and the charged moieties were considered as interacting if at least one pair of atoms was found at a distance shorter than 5 Å. We used the atoms forming the carboxylic group for aspartate and glutamate. The NH3 and the guanidinium groups were employed for lysine and arginine, respectively. A hydrogen bond was identified when the distance between the acceptor atom and the hydrogen atom was lower than 3.5 Å, and the donor-hydrogen-acceptor atom angle was greater than 120°. We calculated main-chain main-chain, main-chain side-chain, and side-chain side-chain hydrogen bonds. We customized the configuration files used by PyInteraph2 for the identification of salt bridges and hydrogen bonds by adding the oxygens of the phosphate group to the charged groups (for salt bridges) and to the hydrogen bond acceptors (for hydrogen bonds) to include the phosphorylated serines. We retained only intermolecular interactions present in at least 20% of the simulation frames (*pcrit* = 20%), as previously applied to other proteins[64–67].

Atomic contact-based Protein Structure Network (PSN)

We used the PyInteraph2 implementation[63] of the PSN method based on atomic contacts as originally formulated by Vishveshwara's group[47] and applied to the MD ensemble of the different constructs used in this study. We retained the pairs of residues whose sequence distance is higher than 1 for the edge calculations and at a distance lower than 4.5 Å. In the calculation, normalization factors for each residue are applied as previously discussed[68] to calculate the interaction strength. We used an I_{min} value of 3. For phospho-serines, we used the normalization factor reported in a previous paper[69]. In addition, we retained edges with occurrences higher than 50% of the frames in the MD ensemble in the final PSN. The interaction strength associated with each edge in the final PSN corresponds to the average interaction strength for the edge over all the ensemble structures, and it is used to weight the PSN.

Changes in binding free energy

We employed Foldx5 [44] to perform *in silico* saturation mutagenesis and estimate changes in binding free energy upon mutations using the Python wrapper MutateX[70]. We estimated the average difference in binding free energy (ΔΔGs) upon mutations using 25 representative structures from the MD simulations of $LC3B_{1-120}$-$OPTN_{158-191}$, $LC3B_{1-120}$-p1$OPTN_{158-191,}$ and $LC3B_{1-120}$-p5$OPTN_{158-191}$. This strategy may help mitigate the limitation due to the backbone stiffness of FoldX[70,71]. As a reference, we also performed similar calculations using as starting structure the 20 NMR-derived structures of $LC3B_{1-119}$-p5$OPTN_{168-185}$ deposited in the PDB entry 2LUE). We calculated the average ΔΔGs for each position, and we used a cutoff of 1kcal/mol in ΔΔG to classify the effects of the mutations[70]. In particular, we defined stabilizing or destabilizing variants featuring ΔΔG values lower than -1 or higher than 1 kcal/mol, respectively.

MAVISp analyses for disease-related variants of $OPTN_{158-191}$

We used the data deposited in the MAVISp database[72] where variants found in COSMIC[73], cBioPortal[74], and ClinVar[75] have been retrieved for OPTN (Uniprot ID Q96CV9, and RefSeq ID NP_068815).

We retained only mutations in the region used for the modeling, i.e., 158-191 (i.e., V161M, E163D, L166V, N169S/T, E175Q, D176Y, F178V, E187Q, V179D, A184D/S). We then applied the MAVISp framework [72] for the assessment of the effects of these variants on: i) structural stability, ii) interactions with LC3B, iii) homodimerization, iv) interplay with post-translational modifications and v) long-range effects.

We used the model downloaded from the AlphaFold database[76] on the 23rd of March 2023 for the STABILITY module of MAVISp. We used the starting structure for the simulations of the non-phosphorylated variant ($LC3B_{1-120}$-$OPTN_{158-191}$), and 25 representative frames from the MD replicate 1 for the LOCAL INTERACTION module in the simple and ensemble mode, respectively. We also used a model generated with AlphaFold multimer and experimentally validated[77] to assess the effects of the mutations on the homodimerization of $OPTN_{1-577}$.

In addition, we retrieved (REVEL, AlphaMissense) or calculated (DeMask, EVE) the pathogenicity scores associated with the mutations using REVEL[78], DeMask[79], EVE[80] and AlphaMissense[81] with MAVISp workflows.

Structural clustering

We calculated pairwise RMSD values for each trajectory using the main-chain atoms of the complex. We performed clustering on the RMSD matrix using the GROMOS algorithm[82] and an RMSD cutoff equal to the average RMSD from each MD replicate. We then retrieved the

average structures for the first three populated clusters for the simulations of the unphosphorylated variant LC3B$_{1-120}$-OPTN$_{158-191}$. The average structure of each cluster corresponds to the structure belonging to the cluster, which features the lowest RMSD values to its neighbors in the cluster. These have been used as starting structures for the calculations in the ensemble mode with MAVISp and the LOCAL_INTERACTION module with the flexddg protocol of RosettaDDGPrediction (see above).

Protein purification

The construct encoding for GST-fused LC3B was expressed in *E. coli* BL21 (DE3) cells, and the expression of the fusion proteins was induced by adding 0.5 mM IPTG at 37 °C for 4 hours. Cells were lysed by sonication in lysis buffer (50 mM Tris-HCl pH 8, 250 mM NaCl, 1 mM DTT, cOmplete™, EDTA-free Protease Inhibitor Cocktail tablets, and PhoSTOP phosphatase inhibitor tablets). Pierce™ Glutathione Magnetic Agarose beads were used to precipitate the GST-fused protein for 2 hours at room temperature under shaking. After washing of the beads in lysis buffer, the protein was eluted in elution buffer (50 mM Tris-HCl pH 8, 10 mM reduced glutathione). The buffer was exchanged in assay buffer (10 mM Tris-HCl pH 7.4, 150 mM NaCl, 0.1 % Tween-20) using PD10 gravity columns (GE Healthcare). The protein was concentrated using Amicon Ultra-0.5 10K Centrifugal Filter Devices (Merck Millipore). The protein concentration was measured on a Nanodrop 1000 spectrophotometer at 280 nm using the extinction coefficient of Trp and Tyr residues in the GST-LC3B sequence (48820 cm-1M-1).

Peptides

N-terminally labeled FITC-Ahx OPTN LIR peptides were purchased from TAGCopenhagen in >95% purity. We have used the following peptide sequences corresponding to the 169-185 residues in full-length OPTN:

FITC-Ahx-NSSGSSEDpSFVEIRMAE (pS177),

FITC-Ahx-NSSGSSEDDFVEIRMAE (S177D),

FITC-Ahx-NSSGSSEDEFVEIRMAE (S177E),

FITC-Ahx-NSSGSSEDSWVEIRMAE (F178W),

FITC-Ahx-NSSGSSEDpSWVEIRMAE (pS177 F178W),

FITC-Ahx-NSSGSSEDpSFVEIRAAE (M183A), and

FITC-Ahx-NSSGSSEDpSAVEIRMAE (F178A),

where "pS" is phosphorylated Ser.

Sequences were confirmed by mass spectrometry and purity by RP-HPLC. The peptides were dissolved in an assay buffer and their concentration was measured on a Nanodrop 1000 spectrophotometer at 280 nm using the extinction coefficient of the FITC group (18542 cm-1M-1) and Trp residue where present..

Microfluidic Diffusional Sizing (MDS)

All MDS experiments were performed on a Fluidity One-M instrument at room temperature with commercial chips (Fluidic Analytics, Cambridge, UK). FITC-Ahx-labeled OPTN LIR peptidesat a total concentration of 1 µM, together with a varying concentration of GST-LC3B, were added and diluted in assay buffer. The samples were incubated at room temperature for at least 2 hours to reach equilibrium. The change in hydrodynamic radius with GST-LC3B concentration was fitted in Prism 9 for macOS using a one-site total binding model, to obtain Kd values.

The equation used for fitting the data was:

$$Y = Rh\_free + (Rh\_complex - Rh\_free) * ((X + n*L + Kd) - ((X + n*L + Kd)^2 - 4*n*L*X)^{0.5}) / (2*n*L)$$

where:

Rh_free = Hydrodynamic radius of the labeled peptide (nm)

Rh_complex = Hydrodynamic radius of the complex between the labeled peptide and unlabeled GST-LC3B (nm)

n = binding sites occupied on the labeled species at saturation (i.e., stoichiometry)

L = the labeled peptide concentration (nM)

Kd = the equilibrium dissociation constant or affinity of the complex (nM)

Each MDS experiment was performed in 3-5 biological replicates, each with 1-3 technical replicates The data for each replicate was normalized to 100% according to the fitted Rh_complex value, and to 0% according to the fitted Rh_free value and was plotted as % BOUND versus GST-LC3B concentration.

The differences in Kd between the tested peptide variants and the pS177 variant were assessed by an unpaired, parametric t-test. Where the tested variant had a variance of more than 4-fold of the pS177 variant, we used the t-test with Welch's correction.

Results and Discussion

Overview of the complex between OPTN and LC3B

The LIR (residues 169-185) of OPTN is located C-terminal to a coiled-coil region and in a disordered tract, according to the AlphaFold2 model deposited in the AlphaFold database (**Figure 1C**). The core region of the motif is encompassed by the sequence 178-FVEI-181, where F178 and I181 occupy the HP1 and HP2 hydrophobic pockets of LC3B, respectively (**Figure 1B**). S177 can be phosphorylated in position -1 to the phenylalanine (**Figure 1B**). Four additional serine residues (S170, S171, S173, and S174) N-terminal to the LIR core motif can also undergo phosphorylation (**Figure 1B**). The structure of the complex between an OPTN LIR variant with the five phospho-serines in complex with LC3B (LC3B$_{1-119}$-p5OPTN$_{169-185}$) has been solved by NMR[34] and used as a starting structure for the modeling and simulations in this work (**Table S1**).

Moreover, we collected one-microsecond unbiased MD simulations of the unphosphorylated (LC3B$_{1-120}$-OPTN$_{169-185}$ and LC3B$_{1-120}$-OPTN$_{158-191}$), mono-phosphorylated (LC3B$_{1-120}$-p1OPTN$_{169-185}$ and LC3B$_{1-120}$-p1OPTN$_{158-191}$) and the multiple phosphorylated variants (LC3B$_{1-120}$-p5OPTN$_{169-185}$ and LC3B$_{1-120}$-p5OPTN$_{158-191}$) all without the tag sequence used for NMR or including a N-terminal flanking region which has helical propensity (**Table S1**).

MD simulations of LC3B in complex with phosphorylated OPTN are in good agreement with NMR chemical shifts

Before proceeding with the analyses of the MD ensembles, it is essential to assess the quality of the structures collected in the simulations, i.e., the conformational sampling. Hence, we compared the MD ensembles with chemical shift data from NMR, similar to what we did for other case studies[53,57,60,83].

For this purpose, we carried out four one-μs MD replicates of LC3B$_{1-119}$-p5OPTN$_{169-185}$. The NMR data of LC3B$_{1-119}$-p5OPTN$_{169-185}$ includes the assignment of backbone and side-chain chemical shifts[34]. Chemical shifts are probes of dynamics along different time scales[84–86] and can also be predicted by MD simulations ensemble of structures[58,87,88]. The calculated chemical shifts from our simulations agree with the experimental data with low $\chi 2_{red}$ values, which converged to a plateau after 200-300 ns (**Figure 2A**).

Furthermore, we calculated the side-chain chemical shifts for methyl-containing residues (Ile, Val, and Leu), which are suitable probes of more local conformations (**Table 1**) and are also associated with preferences for different rotameric states of each residue in solution (refs to the work with the link between CS and rotameric states of chi1/chi2). Most of the chemical

shift classes featured predicted values in agreement with the experimental data, with low χ^2_{red} indicating that the MD sampling capture the overall dynamics observed experimentally in solution. The only exception is isoleucine residues with a χ^2_{red} higher than 3 for the Cδ depending on the replicate under investigation (**Table 1**).

With MD simulations of reasonable quality and in agreement with the protein dynamics in solution on the microsecond timescale, it becomes meaningful to use the MD ensembles to study the effects induced by phosphorylation and unveil new details of the underlying mechanism.

The phosphorylation reduces the flexibility of the OPTN LIR in the binding pocket of LC3B

Phosphorylation of LIR motifs can decrease the flexibility of the LIR peptide in complex with LC3B[89], stabilizing the conformation of the LIR residues in the binding pocket of LC3B. We thus investigated if the phosphorylation of the LIR of OPTN could exert a similar effect. We used the RMSF of the Cα atoms of the LIR peptide as a probe for changes in structural flexibility using the LC3B$_{1-120}$-OPTN$_{158-191}$ constructs. We observed a marked decrease in LIR flexibility upon phosphorylation of the OPTN LIR (**Figure 2B**), whereas the unphosphorylated peptide displays higher RMSF values along the entire sequence. The effect is observable at different timescales, as attested by the analyses carried out within ten and 100-ns time windows (**Figure 2B**). The phosphorylation of only S177 is not sufficient to decrease the flexibility of the LIR in the complex (**Figure 2C**).

Phosphorylation of S177 is sufficient to increase the propensity to β-strand formation in the LIR binding pocket

LIRs, like other SLiMs, can have different secondary structure propensity upon binding with an interactor. Indeed, it has been experimentally shown that several LIRs fold upon binding, inducing the formation of an intermolecular parallel β-sheet with the β2 strand of LC3B. In contrast, other LIRs retain disorder in the bound state[25,34]. We thus evaluated if the phosphorylation could affect the secondary structure propensity of the OPTN LIR in a complex with LC3B using the simulations with the LC3B$_{1-120}$-OPTN$_{158-191}$ constructs (**Figure 2D-E**, **Figure S1**).

We only observed the two residues downstream to F178 (i.e., 179-180) in the unphosphorylated variant with a propensity for β-strands (**Figure 2D**). The phosphorylation of S177 is sufficient to slightly locally increase the β-strand propensity, including F178 (**Figure S1**). The multiple phosphorylation did not increase further β-strand formations with respect to the mono-phosphorylated variant (**Figure 2E, Figure S1**). The β-strand of OPTN contributes

to the intermolecular β-sheet via interaction with the β-strand 2 of LC3B, as observed in the experimental structures of the multiple phosphorylated variants or of the phospho-mimetics[34].

Phosphorylation of OPTN does not change the solvent accessibility of the core LIR residues but influences charged residues in the LDK tripeptide and in the LC3B HP1 pocket

Next, we verified if the decreased flexibility of the peptide in the pocket caused by the phosphorylation was also causing changes in solvent accessibility of the side chains of the two hydrophobic residues of the LIR core motif (i.e., F178 and I181). We could not observe marked differences in the distribution of the SASA values of HP1 and HP2 residues (**Figures S2 and S3**). Furthermore, we evaluated if the residues lining the HP1 and HP2 pockets of LC3B, the LDK tripeptide, or the LC3B ubiquitin patch were sensitive to changes in solvent accessibility. We observed a marked effect on K51 in the HP1 pocket, where the phosphorylation of S177 is sufficient to exert the effect (**Figure 3A**). Moreover, the multiple phosphorylations also decreased the solvent accessibility of D48 and K49 in the LDK tripeptide (**Figure 3B-C**). Other residues of LC3B were not influenced by changes in the accessibility of their side chains upon phosphorylation.

In summary, the decreased structural flexibility seems not to cause changes in the residues of the LIR core motif, but it influences the accessibility of the interacting residues in LC3B.

The phosphorylation increases the interaction strength between the LIR and LC3B both locally and long-range

Furthermore, we evaluated if the changes in solvent accessibility and flexibility were associated with changes in the interaction strenght between OPTN and LC3B. We used the pairwise interaction strength as calculated from the atomic-contact-based protein structure network (see Materials and Methods, **Table S2 and Figure 4A**).

We observed that the phosphorylation of pS177 and even more of the five serine residues increased the interaction strength between F178 of OPTN and I23 of the HP1 pocket of LC3B. Moreover, it also increased the strength of interaction between V179 of OPTN and F52 of the HP2 pocket of LC3B. In contrast, the interaction strength of the LIR core residue I181 was similar in all the simulations. The phosphorylation at multiple sites also has an additional long-range effect, increasing the interaction strength in the N-terminal flanking region of the OPTN LIR between E163 and R37.

A rewiring of intermolecular salt bridges upon phosphorylation of OPTN with both local and long-range effects

To evaluate if phosphorylation of the OPTN LIR could affect electrostatic interactions, we estimated salt bridges (**Table S3, Figure 4B-D**) from the MD simulations of the LC3B$_{1-120}$-OPTN$_{158-191}$ variants (see Materials and Methods).

The interaction between E180 of OPTN and K30 of LC3B is the only salt bridge that is conserved at high occurrence (> 90%) in all the complexes. The unphosphorylated variant has a different pattern of salt bridges with occurrence higher than 20% depending on the MD replicate under consideration, changing from only E180-K30 in replicate 3 to some more sparse and lower occurrence salt bridges involving different residues at the interaction interface in replicates 1 and 2 (**Table S3**). Interestingly, E175 and D176 of OPTN can interchange in their role of partners of interactions for K49 and K51 of OPTN in a fluid salt-bridge network.

The single phosphorylation of S177 rewired the electrostatic interactions toward a less dynamic and heterogeneous behavior. The high-occurrence salt bridges were strengthened and concentrated in the central part of the interface with only three main players, i.e., D176, pS177, and E180, stably interacting with K51, K49, and K30, respectively (**Table S3**, **Figure 4D-F**). The interaction between K51 and pS177 agrees with what was observed with NMR experiments and changes in affinity upon a K51A mutation on shorter constructs of OPTN[34].

We observed the same rewiring of interactions for the MD simulations of the pentaphosphorylated variant. Moreover, pS170 and pS171 contributed to stabilizing and expanding the network of electrostatic interactions involving R10 and R11 or LC3B. In addition, in agreement with the long-range effects observed above, the multiple phosphorylations promoted more stable electrostatic interactions in the N-terminal region of OPTN, where E163 and K167 also participated in salt bridges, and pS174 contributed to strengthening the interaction with K49 (**Table S3, Figure 4B-D**). The other phospho-residues were not involved in stable salt bridges with LC3B. The results above agree and provide a mechanistic explanation for the decreased solvent accessibility of K49 and K51 in the phosphorylated variants observed in **Figure 3**.

Interestingly, K49 also engages in a salt bridge with the residue corresponding to S177 in the LIR motif of p62 (i.e., D337), suggesting that this lysine may participate in the LIR-binding process by recruiting negatively charged residues from different positions of the N-terminal region of a LIR peptide[89].

The analysis of intermolecular hydrogen bonds involving either main chain or side chain atoms in the different MD replicates confirmed a similar rewiring of interactions (**Table S4**).

In a previous study with shorter constructs for the OPTN LIR, the phosphorylated structure was studied using NMR for the phosphorylated variants. Moreover, an additional structure of a penta-phosphomimetic replacing the serines with glutamates was solved by X-ray[34]. In the

X-ray structure, the authors observed an inward flip of R11 to make hydrogen bonds with the side chain of E177 of LC3B. The authors could not detect the amide proton of R11 side chain with NMR. We thus evaluated if a similar mechanism was occurring in our simulations with longer constructs for the interaction between R11 and pS177. We collected the distances over time between the side chains of the two residues in the penta-phosphorylated simulations and evaluated their probability distribution. We found the two residues at distances larger than 10 Å in our simulations, whereas we noticed that R11 mostly sense another phospho-site, i.e., pS171. The mechanism suggested here could still explain the changes in the chemical shift for R11 upon phosphorylation observed with NMR. The control simulation with a shorter construct (LC3B[1-119]-p5OPTN[169-185]) featured the interaction between pS177 and R11, as observed in the model derived by the NMR data and deposited in PDB.

High-throughput mutational scans to identify binding hotspots and residues conferring specificity for the binding mode of the phosphorylated variant

The integration of MD-derived ensembles of conformations with empirical free energy calculations[53,70,71,83,90,91] could shed light on hotspots for the binding between LIR and LC3B proteins. According to a structure-based framework that we recently developed, i.e., MAVISp framework [72], we rely on a consensus approach based on FoldX and Rosetta. However, the protocols for Rosetta-based free energy calculations do not support phospho-residues, and one should rely on phosphomimetics. To determine the effectiveness of phosphomimetics in replicating the effects of phosphorylation, we synthesized OPTN LIR peptides, introducing a phosphoserine at position 177 (pS177), as well as phosphomimicking substitutions (S177D and S177E). Subsequently, we conducted MDS experiments to evaluate the binding affinity of these peptides to GST-LC3B (**Figure 5A**). The affinity of the pS177 peptide was measured at 13.7 µM (**Figure 5A1**), while the S177D and S177E variants exhibited affinities of 35.5 µM (**Figure 5A2**) and 41.1 µM (**Figure 5A3**), respectively. The difference between pS177 and S177D does not seem significant (p = 0.123), possibly due to the high variability of the S177D data, while the difference between pS177 and S177E is statistically significant (p = 0.001). Existing literature data indicates that the affinity of the wild-type peptide falls within the range of 40-60 µM[33,34]. Therefore, it appears that the single phosphomimetic mutation at position 177 might confer a modest increase in binding affinity compared to the wild-type sequence. Still, the mimetics do not reach the level observed with authentic phosphorylation. Consequently, we decided not to include Rosetta-based calculations in this study and relied on FoldX calculations, using 25 representative structures from each of the MD replicates of the constructs LC3B[1-120]-OPTN[169-185] and LC3B[1-120]-OPTN[158-191] from **Table S1**.

We had three main goals: i) evaluate if, with the mutational scan, we can predict the experimentally known stabilizing effect of the phosphorylations on the binding [33,34] to provide a baseline for similar future studies on other phosphorylatable LIRs; ii) to use the data to understand in more details the structural mechanisms promoted by the phosphorylation.

At first, we designed the calculations to evaluate the effect of introducing the phosphorylations on the structure of the unphosphorylated complex. In addition, we also used the structures from the phosphorylated complex and assessed the effect of removing the phosphorylation, which should destabilize the binding in the case of FoldX. We used this 'reverse approach' since FoldX has limitations in predicting stabilizing effects[92,93]. Thus, we speculate that it could be an effective alternative strategy to apply for the de novo discovery effects induced by phosphorylations on SLiMs. In more detail, the reverse approach entails a mutational scan with Foldx on ensembles of the phospho-variant, where the assessment is on the effect induced by substituting the mutation to the original unphosphorylated amino acid. In this kind of comparison, for example, if one observes only a marginal effect when a PTM/mutation is introduced starting from the wild-type ensemble of conformations but a clearly destabilizing effect when this PTM/mutation is removed in the ensemble of the modified variant, it can be more confidently predicted that the mutation is stabilizing the wild-type variant of the protein. In our comparison with experimental data, we also added the variant F178W of the OPTN LIR, which should mimic the interaction occurring for p62 with the HP1 pocket of LC3B.

At first, we observed that we could not recapitulate the stabilizing effects induced on the binding by the mutations F178W of the complex between LC3B and the unphosphorylated OPTN construct since the mutation was predicted with a neutral effect in most of the structures with regards to binding free energy. However, we observed, for some of the structures of the MD replicate2, a stabilizing effect induced by F178W with changes in binding ΔΔG of approximately -2 kcal/mol. F178W is of particular interest since it should mimic the interaction of the conventional p62 LIR using the wild-type ensemble of the complex.

When we assessed the effect of the phosphorylation starting from the unphosphorylated complex, we obtained average ΔΔGs values lower than 0, with pS177 and pS174 as the most important modifications to increase the binding affinity. Although the average differences were lower than the threshold of 1 kcal/mol generally used to pinpoint stabilizing or destabilizing variants for a protein-protein interaction. A group of conformations from the MD simulations featured values lower than the threshold, as observed for the F178W mutation introduced in the unphosphorylated variant. With the reverse approach, i.e., using the phospho-variant and measuring the effect of reintroducing a serine, we also observed a small average destabilizing effect (i.e., average ΔΔG values of 0.4-0.8 kcal/mol). The small changes might be due to the fact that the MutateX protocol only allows considering one mutation at a time[70]. This implies that the other phosphorylations were still included in the model when assessing a change from

phospho-serine to serine at one site. We did not include calculations using multiple mutations of the phospho-sites simultaneously since empirical free energy functions, such as the one applied here with local sampling, are often not able to capture with accuracy the contribution of multiple mutations and would rather provide predictions that are the results of additive contributions from the single mutations.

Furthermore, we used the data from the saturation scan to identify residues that could contribute specifically to the binding affinity upon phosphorylation due to the rewiring of the interactions at the binding interface in the presence of phosphorylation (**Figure 4B-D**). To this goal, we selected residues that could serve as binding hotspots in the LIR (or LC3B) only in its pentaphosphorylated form and with marginal contributions in the unphosphorylated variant (**Table S5**, **Figure 5B-C**). We also verified if the mono-phosphorylation of S177 was sufficient to promote the same effects (**Table S5, Figure 5D**).

Apart from confirming the importance of the phosphorylation sites, we identified three sites in the N-terminal flaking region as key only for the phosphorylated variant, i.e., E163 (when mutated to S, G, T, H, or R), different mutations at the positions L166 (especially substitutions to I, M, F or R), and K167 when mutated to small, aromatic, or negatively charged residues. The single phosphorylation does not seem to be sufficient to exert the same N-terminal effects as the ones observed for the penta-phosphorylated variant (**Table S5**). On the other side, E175 seems to have a more central role in the interaction of the unphosphorylated variant than the penta-phosphorylated one, in agreement with the changes observed in salt-bridge interactions (**Figure 4B-D**). D176 and M183 are sensitive to changes in binding free energies upon mutation both in the unphosphorylated and phosphorylated variants.

We also noticed different amino acid preferences upon phosphorylation for the LC3B sites at the interface. For example, mutating D19 to aromatic or positively charged amino acids could further stabilize the complex with the pentaphosphorylated OPTN peptide. K49 and K51 are slightly stronger interaction hotspots for the phospho-peptide than the unphosphorylated variant. R37 is instead a centra hotspot only for the phosphorylated peptide. In fact, R37 can only tolerate substitutions to lysines in the LC3B-p5OPTN complex, in agreement with its importance in mediating the interaction strength and salt bridges with the N-terminal region of OPTN.

Effects of disease-related mutations of OPTN on the binding with LC3B

The methodology provided here could also be applied to scrutinize the effects of disease-related variants, which still have uncertain significance or conflicting evidence and pose a challenge to biomedical research [94].

We identified twelve variants reported in cancer databases or in ClinVar (**Table 2, Figure 5E-F**) located in the OPTN LIR or its flanking regions (residues 158-191).

None of the variants seem to affect OPTN homodimerization, and thus, potentially on its oligomerization, even if E175Q, F178V, and V179D, remains uncertain according to MAVISp (**Figure 5E**).

The MAVISp approach and the pathogenicity scores agree in the classification of only mutations at the HP1-binding residue F178 (i.e., F178V) as damaging for the protein for its interaction with LC3B and possibly pathogenic (**Figure 5E-F**).

D176Y is predicted to damage mostly the binding with LC3B for the unphosphorylated forms of OPTN using the ensemble mode of MAVISp, i.e., accounting for protein dynamics (**Figure 5F** and **Table 2**).

L166V seems to be damaging for the binding only for the phosphorylated form of the OPTN-LC3B complex, but its damaging effect cannot be recapitulated by the single phosphorylation at S177 (**Table 2**).

The different effects that disease-related mutations can exert depending on the phosphorylation status of the protein are interesting from a broader point of view. Approaches to predicting the effects of mutations often neglect the interplay between the mutation itself and the phosphorylation status of the target protein. We will use this example to design and implement a new module for the ensemble mode of the MAVISp framework, where we can also capture the differential effects of mutations depending on a PTM.

Of interest, V179D, V161M, and E163D are reported as a variant of uncertain significance in glaucoma and amyotrophic lateral sclerosis in ClinVar. Even though EVE and AlphaMissense predict a possible pathogenic effect for E163D and V179D (**Figure 5E-F, Table 2**), our structure-based assessment suggests that they have neutral effects. Instead, the MAVISp assessment and all the pathogenicity scores agree on the neutral effect of V161M, in agreement with what was observed in a population study for the same variant[95].

Microfluidic Diffusional Sizing can accurately estimate changes in Kd upon substitutions in the phosphorylated variant of OPTN

To further evaluate stabilizing or destabilizing effects of amino acid substitutions on the interaction between LC3B and phosphorylated optineurin, we established an approach based on Microfluidic Diffusional Sizing (MDS)[48,49,96] to study the binding affinity between GST-tagged LC3B and different peptide variants corresponding to the 169-185 residues of the full-length OPTN. We have purchased FITC-labeled peptides and titrated them with increasing concentrations of GST-LC3B to follow changes in hydrodynamic radius that could be fitted to obtain a Kd value for each peptide. The low sample volume of GST-LC3B needed per binding

curve (8 µL) allowed us to upconcentrate GST-LC3B to the hundreds of micromolar levels needed to saturate the binding signals for most peptide. Due to synthesis issues with penta-phosphorylated peptides, we introduced substitutions in the pS177 mono-phosphorylated peptide of OPTN. Luckily, we observed in the simulations and the free energy scans discussed above that many of the structural effects observed in the penta-phosphorylated variant are also recapitulated by the single phosphorylation. At first, we measured a Kd of 13.7 µM (Figure 5A1) for the mono-phosphorylated variant, in good agreement with what was previously shown with isothermal titration calorimetry[34]. Moreover, we added as a control the unphosphorylated peptide bearing the F178W substitution that mimics the p62 LIR and resulted in a Kd of 18.9 µM, in range with the pS177 variant (**Figure 6A**, p = 0.214) as previously reported[34]. An alanine substitution at position 178 (F178A) of OPTN abolishes the hydrophobic residue for the interaction with the HP1 pocket of LC3B. This variant combined with the phosphorylation at position 177 caused an increase of Kd of 5-fold (**Figure 6B**, p = 0.0019). This agrees with what we observed in the scans in silico where the introduction of F178A in the mono-phosphorylated construct resulted in average changes in binding ΔΔG of approximately 4.5 kcal/mol. This suggests that the phosphorylation alone cannot compensate for the loss of binding affinity caused by removing the central residue in the core motif. This result is also aligned with the possible damaging impact of the shortening of the side-chain length for the disease-related F178V variant discussed above (**Figure 5E-F**).

On the contrary, the F178W substitution that mimics a W-type LIR combined with the pS177 phosphorylation can increase the binding affinity 7-fold compared to the phosphorylation alone (**Figure 6C**, p = 0.003). To evaluate the effects of variants in the LIR flanking region, as well as to test the limit of the MDS approach in capturing smaller changes in binding affinity, we assessed the M183A substitution, which is predicted to destabilize the interaction for both the phosphorylated and unphosphorylated variants of OPTN with approximately 2 kcal/mol. We observed similar Kd values within the experimental error, from combining the phosphorylation at S177 and M183A compared to the phosphorylation alone (**Figure 6D, p = 0.25**). Our results collectively suggest that the MDS binding assay could be used to identify and validate predicted changes in binding affinity of more than 4 kcal/mol. Thus, we did not work experimentally with the variants reported in **Table 2 or S4**, since all the predicted average changes for mutations in residues outside the core motif were mostly 1-2.5 kcal/mol (**Table S5**).

Conclusions

Phosphorylation can modulate LIR motifs and compensate for the lower binding affinity in LIRs. Phosphorylatable sites have been reported and experimentally verified for other LIR

motifs with an impact on binding affinity and recognition of LC3/GABARAP proteins[19]. It is thus of fundamental importance to investigate the role of these phosphorylations on the structure and function of LIR-containing proteins and their complexes.

We here employed the OPTN LIR as a model system of phospho-switches for the LIR-ATG8 recognition, thanks to the availability of biophysical data on its phosphorylations and mutations that can be used to compare and validate our simulation data[33,34]. Moreover, we generated new experimental data on different OPTN variants applying for the first time a technique based on Microfluidic Diffusional Sizing to protein-SLiM interactions. In addition, we showed for the first time the role of the flanking-region N-terminal to the core LIR motif of OPTN as a sensor of long-range effects induced by OPTN phosphorylation.

We show, together with previous results on another LIR-containing protein[89], that phosphorylation can exert this effect by impacting the flexibility of the entire LIR peptide once the complex is formed. Moreover, the post-translational modification can induce long-range effects stabilizing the electrostatic interactions in the flanking region of the peptide, rewiring the network of intermolecular interactions. This rewiring also causes changes in hotspots for binding in the flanking regions and their contribution to the binding with LC3B. The phosphorylation of the residue pS177 alone seems to be sufficient to promote changes in solvent accessibility for the LC3B residues involved in interactions with the central part of the OPTN LIR motif and to expand the β-sheet structure in the binding cleft. On the other hand, the mono-phosphorylation is not sufficient to decrease the protein flexibility of OPTN in the LC3B binding pocket, along with eliciting long-range effects to recruit residues from the N-terminal flanking regions into the complex formation.

Phosphorylation of OPTN is important to regulate its cellular function and the structural changes induced by it should also be considered in the context of understanding the impact of disease-related mutations. The approach used in this study to investigate the effects of disease-related variants on the phosphorylated form of the protein can serve to design future studies on other disease-related targets for which phosphorylation events have been experimentally characterized.

Tables

Table 1. Comparison between experimental and predicted side chain chemical shifts for methyl groups. The results for the four MD replicates of LC3B$_{1-119}$-p5OPTN$_{169-185}$ are reported in the table. We report the reduced χ^2 for each replicate and residue. Low χ^2_{red} values indicate a good agreement with the experimental data.

LC3B residue type	Replicate 1	Replicate 2	Replicate 3	Replicate 4
ILE (Cδ)	3.75	4.16	2.81	3.31
ILE (Cγ2)	0.73	0.72	0.56	0.64
LEU (Cδ1)	1.35	1.34	1.36	1.34
LEU (Cδ2)	0.34	0.36	0.46	0.42
VAL(Cγ1)	0.59	0.61	0.59	0.60
VAL(Cγ2)	0.29	0.24	0.39	0.30

Table 2. Disease-related mutations and pathogenicity scores. If the source for the variant is ClinVar we reported the review status and the classification. DeMask generally classifies as loss-of-function variants with a fitness score < 0 and gain-of-function > 0. We used as a threshold for damaging variants a DeMask score < -0.3 or > 0.3 to decrease the number of false positives. A similar strategy is applied to the REVEL score.

variant	source	REVEL	DeMask delta fitness	EVE	Alphamissense	ΔΔG binding - p5	ΔΔG binding - p1
V161M	ClinVar (1, VUS)	0.435.	-0.201	Benign	Ambiguous	Neutral	Neutral
E163D	ClinVar (1,VUS)	0.404	-0.264	Pathogenic	Pathogenic	Neutral	Neutral
L166V	cBioPortal	0.378	-0.132	Benign	Benign	Potentially Damaging (0.8-1.6 kcal/mol)	Neutral
N169S	cBioPortal	0.161	-0.010	Benign	Benign	Neutral	Neutral
N169T	cBioPortal	0.227	-0.059	Uncertain	Benign	Neutral	Neutral
E175Q	cBioPortal COSMIC	0.309	-0.092	Pathogenic	Benign	Potentially Neutral (0.2-1.3 kcal/mol)	Damaging (1.0-145 kcal/mol)
D176Y	cBioPortal COSMIC	0.442	-0.317	Pathogenic	Ambiguous	Potentially Neutral (0.2-1.2 kcal/mol)	Potentially Damaging (0.9-1.6 kcal/mol)
F178V	cBioPortal ,COSMIC	0.58	-0.341	Pathogenic	Pathogenic	Damaging (3.0-4.5 kcal/mol)	Damaging (3.0-.3.5 kcal/mol)
V179D	ClinVar (1,VUS)	0.482	-0.389	Pathogenic	Pathogenic	Neutral	Neutral
A184D	cBioPortal	0.245	-0.091	Benign	Benign	Neutral	Neutral
A184S	cBioPortal COSMIC	0.159	0.042	Benign	Benign	Neutral	Neutral

Figure legends

Figure 1. OPTN LIR region and constructs used in the molecular dynamics (MD) simulations. The figure illustrates the location of the LIR motif of OPTN and the hydrophobic residues for binding to the HP1 and HP2 pockets of LC3B. We also summarized the constructs used for the different MD simulations. The list of simulations collected in the study can be found in Table S1.

Figure 2. Comparison with experimental data and changes in protein flexibility and secondary structures induced by phosphorylation. A) For each replicate of the $LC3B_{1-119}$-$p5OPTN_{169-185}$ construct, we estimated a reduced χ^2 (χ^2_{red}) over the simulation time. This value provides a measurement of the agreement between the experimental chemical shifts from NMR and the calculated chemical shifts from the MD trajectories. The metric is calculated for each atom type. Low values of χ^2_{red} indicate better agreement between experiments and simulations. We reported the example of the MD replicate 1 of $LC3B_{1-119}$-$p5OPTN_{169-185}$ for the sake of clarity. **B-C)** The average Cα RMSF for each replicate, calculated using 100-ns time windows is showed for $LC3B_{1-120}$-$p5OPTN_{158-191}$ (B) and $LC3B_{1-120}$-$p1OPTN_{158-191}$ (C) compared to the unphosphorylated construct, respectively. **D-F)** The bar plots show the secondary structure content according to the DSSP dictionary using a concatenated trajectory with all the MD replicates for $LC3B_{1-120}$-$OPTN_{158-191}$ (D), $LC3B_{1-120}$-$p1OPTN_{158-191}$ (E) and $LC3B_{1-120}$-$p5OPTN_{158-191}$ (F). It can be observed that the phosphorylation of S177 is sufficient to increase the propensity to β-strand formation.

Figure 3. Phosphorylated $OPTN_{158-191}$ causes changes in solvent accessibility of charged residues of LC3B at the interaction interface. The relative solvent accessibility of side chain atoms is reported for K51 (A), K49 (B), and D48 (C) in the different replicates of $LC3B_{1-120}$-$OPTN_{158-191}$ and $LC3B_{1-120}$-$OPTN_{168-185}$ variants.

Figure 4. Changes in intermolecular interactions upon phosphorylation. A) The histograms report the results of pairwise Interaction Strength as calculated using an atomic-contact-based protein structure network for the pairs more sensitive to changes upon phosphorylation for the $LC3B_{1-120}$-$OPTN_{158-191}$, $LC3B_{1-120}$-$p5OPTN_{158-191}$. and $LC3B_{1-120}$-$p1OPTN_{158-191}$ constructs. The full set of data on interaction strengths is reported in Table S2. B-D) The networks of salt bridges are reported on the 3D structure of the first frame of the MD simulations of $LC3B_{1-120}$-$OPTN_{158-191}$ (B), $LC3B_{1-120}$-$p1OPTN_{158-191}$ (C) and $LC3B_{1-120}$-$p5OPTN_{158-191}$ (D) using the MD replicate 1 for sake of clarity. The salt bridges are depicted as gray sticks, whose size is proportional to their occurrence in the simulation. The full set of data is reported in Table S3.

Figure 5. High-throughput mutational scans. A) Binding isotherms of 1) pS177, 2) S177D, and 3) S177E FITC-labeled peptide variants to GST-LC3B. All biological and technical replicates are plotted, and fitted Kd values represent the global fit ± standard error across biological replicates. p values are generated through t-tests or t-tests with Welch's correction, as described in Methods, to assess differences to the pS177 variant. B-D) The heatmaps for the saturation scan in silico for $LC3B_{1-120}$-$OPTN_{158-191}$ (B), $LC3B_{1-120}$-$p5OPTN_{158-191}$ (C), and $LC3B_{1-120}$-$p1OPTN_{158-191}$ (D) are illustrated with a truncation of the ΔΔG values at 3 kcal/mol for visualization purposes. We can observe that the pentaphosphorylation activates binding hotspots in the N-terminal flanking region and decreases the importance of the negatively

charged residues in the proximity of the core motif (E175 and D176) for the binding to LC3B (B-C). The effect cannot be recapitulated by the single phosphorylation (C-D). E-F) The dotplot summarized the results of the MAVISp assessment in the simple (E) and ensemble (F) modes. Black, white, and grey dots represent mutations predicted with a damaging, neutral, or uncertain effect for one of the MAVISp modules, respectively.

Figure 6. Equilibrium binding curves by Microfluidic Diffusional Sizing. Binding isotherms of A) F178W, B) pS177 F178W, C) pS177 M183A and D) pS177 F178A FITC-labeled peptide variants to GST-LC3B. All biological and technical replicates are plotted, and fitted Kd values represent the global fit ± standard error across biological replicates. p values are generated through t-tests or t-tests with Welch's correction, as described in Methods, to assess differences to the pS177 variant.

References

1.   Oldfield, C. J. & Dunker, A. K. Intrinsically Disordered Proteins and Intrinsically Disordered Protein Regions. *Annu Rev Biochem* **83**, 553–584 (2014).
2.   Wright, P. E. & Dyson, H. J. Intrinsically disordered proteins in cellular signalling and regulation. *Nat Rev Mol Cell Biol* **16**, 18–29 (2014).
3.   Teilum, K., Olsen, J. G. & Kragelund, B. B. On the specificity of protein-protein interactions in the context of disorder. *Biochemical Journal* vol. 478 2035–2050 Preprint at https://doi.org/10.1042/BCJ20200828 (2021).
4.   Van Der Lee, R. *et al.* Classification of intrinsically disordered regions and proteins. *Chem Rev* **114**, 6589–6631 (2014).
5.   Dogan, J., Gianni, S. & Jemth, P. The binding mechanisms of intrinsically disordered proteins. *Physical Chemistry Chemical Physics* vol. 16 6323–6331 Preprint at https://doi.org/10.1039/c3cp54226b (2014).
6.   Schuler, B. *et al.* Binding without folding – the biomolecular function of disordered polyelectrolyte complexes. *Curr Opin Struct Biol* (2020) doi:10.1016/j.sbi.2019.12.006.
7.   Malagrinò, F. *et al.* Unveiling induced folding of intrinsically disordered proteins – Protein engineering, frustration and emerging themes. *Current Opinion in Structural Biology* vol. 72 153–160 Preprint at https://doi.org/10.1016/j.sbi.2021.11.004 (2022).
8.   Barrera-Vilarmau, S., Teixeira, J. M. C. & Fuxreiter, M. Protein interactions: anything new? *Essays Biochem* (2022) doi:10.1042/EBC20220044.
9.   Bondos, S. E., Dunker, A. K. & Uversky, V. N. Intrinsically disordered proteins play diverse roles in cell signaling. *Cell Communication and Signaling* vol. 20 Preprint at https://doi.org/10.1186/s12964-022-00821-7 (2022).
10.   Darling, A. L. & Uversky, V. N. Intrinsic disorder and posttranslational modifications: The darker side of the biological dark matter. *Front Genet* **9**, 1–18 (2018).
11.   Phillips, A. H. & Kriwacki, R. W. Intrinsic protein disorder and protein modifications in the processing of biological signals. *Current Opinion in Structural Biology* vol. 60 1–6 Preprint at https://doi.org/10.1016/j.sbi.2019.09.003 (2020).
12.   Kliche, J. *et al.* Large-scale phosphomimetic screening identifies phospho-modulated motif-based protein interactions. *Mol Syst Biol* **19**, (2023).
13.   Newcombe, E. A., Delaforge, E., Hartmann-Petersen, R., Skriver, K. & Kragelund, B. B. How phosphorylation impacts intrinsically disordered proteins and their function.

Essays in Biochemistry vol. 66 901–913 Preprint at https://doi.org/10.1042/EBC20220060 (2022).

14. Tompa, P., Davey, N. E., Gibson, T. J. & Babu, M. M. A Million peptide motifs for the molecular biologist. *Molecular Cell* vol. 55 161–169 Preprint at https://doi.org/10.1016/j.molcel.2014.05.032 (2014).

15. Davey, N. E. *et al.* Attributes of short linear motifs. *Mol Biosyst* **8**, 268–281 (2011).

16. Cermakova, K. & Hodges, H. C. Interaction modules that impart specificity to disordered protein. *Trends in Biochemical Sciences* vol. 48 477–490 Preprint at https://doi.org/10.1016/j.tibs.2023.01.004 (2023).

17. Palopoli, N., Foutel, N. S. G., Gibson, T. J. & Chemes, L. B. Short linear motif core and flanking regions modulate retinoblastoma protein binding affinity and specificity. *Protein Engineering, Design and Selection* vol. 31 69–77 Preprint at https://doi.org/10.1093/protein/gzx068 (2018).

18. Bugge, K. *et al.* Interactions by Disorder – A Matter of Context. *Front Mol Biosci* **7**, 110 (2020).

19. Sora, V. *et al.* Structure and Dynamics in the ATG8 Family From Experimental to Computational Techniques. *Frontiers in Cell and Developmental Biology* vol. 8 Preprint at https://doi.org/10.3389/fcell.2020.00420 (2020).

20. Kliche, J. & Ivarsson, Y. Orchestrating serine/threonine phosphorylation and elucidating downstream effects by short linear motifs. *Biochemical Journal* vol. 479 1–22 Preprint at https://doi.org/10.1042/BCJ20200714 (2022).

21. Sora, V. & Papaleo, E. Structural Details of BH3 Motifs and BH3-Mediated Interactions: an Updated Perspective. *Frontiers in Molecular Biosciences* vol. 9 Preprint at https://doi.org/10.3389/fmolb.2022.864874 (2022).

22. Johansen, T. & Lamark, T. Selective Autophagy: ATG8 Family Proteins, LIR Motifs and Cargo Receptors. *Journal of Molecular Biology* vol. 432 80–103 Preprint at https://doi.org/10.1016/j.jmb.2019.07.016 (2020).

23. Rogov, V. V. *et al.* Atg8 family proteins, LIR/AIM motifs and other interaction modes. *Autophagy Reports* **2**, (2023).

24. Gatica, D., Lahiri, V. & Klionsky, D. J. Cargo recognition and degradation by selective autophagy. *Nat Cell Biol* **20**, 233–242 (2018).

25. Birgisdottir, Å. B., Lamark, T. & Johansen, T. The LIR motif - crucial for selective autophagy. *Journal of Cell Science* vol. 126 3237–3247 Preprint at https://doi.org/10.1242/jcs.126128 (2013).

26. Adriaenssens, E., Ferrari, L. & Martens, S. Orchestration of selective autophagy by cargo receptors. *Current Biology* vol. 32 R1357–R1371 Preprint at https://doi.org/10.1016/j.cub.2022.11.002 (2022).

27. Rogov, V. V *et al.* Structural and functional analysis of the GABARAP interaction motif (GIM). *EMBO Rep* **18**, 1382–1396 (2017).

28. Joachim, J. *et al.* Centriolar Satellites Control GABARAP Ubiquitination and GABARAP-Mediated Autophagy. *Current Biology* **27**, 1–14 (2017).

29. Holdgaard, S. G. *et al.* Selective autophagy maintains centrosome integrity and accurate mitosis by turnover of centriolar satellites. *Nat Commun* **10**, 1–19 (2019).

30. Ryan, T. A. & Tumbarello, D. A. Optineurin: A coordinator of membrane-associated cargo trafficking and autophagy. *Front Immunol* **9**, 1–16 (2018).

31. Wong, Y. C. & Holzbaur, E. L. F. Optineurin is an autophagy receptor for damaged mitochondria in parkin-mediated mitophagy that is disrupted by an ALS-linked mutation. *Proc Natl Acad Sci U S A* **111**, E4439–E4448 (2014).

32.   Ying, H. & Yue, B. Y. J. T. Optineurin: The autophagy connection. *Exp Eye Res* **144**, 73–80 (2016).

33.   Wild, P. *et al.* Phosphorylation of the Autophagy Receptor Optineurin Restricts Salmonella Growth. *Science (1979)* **333**, 228–233 (2011).

34.   Rogov, V. V. *et al.* Structural basis for phosphorylation-triggered autophagic clearance of Salmonella. *Biochemical Journal* **454**, 459–466 (2013).

35.   Padman, B. S. *et al.* LC3/GABARAPs drive ubiquitin-independent recruitment of Optineurin and NDP52 to amplify mitophagy. *Nat Commun* **10**, (2019).

36.   Qiu, Y. *et al.* Emerging views of OPTN (optineurin) function in the autophagic process associated with disease. *Autophagy* **18**, 73–85 (2022).

37.   Weil, R., Laplantine, E., Curic, S. & Génin, P. Role of optineurin in the mitochondrial dysfunction: Potential implications in neurodegenerative diseases and cancer. *Frontiers in Immunology* vol. 9 Preprint at https://doi.org/10.3389/fimmu.2018.01243 (2018).

38.   Swarup, G. & Sayyad, Z. Altered functions and interactions of glaucoma-associated mutants of optineurin. *Front Immunol* **9**, (2018).

39.   Evans, R. *et al.* Protein complex prediction with AlphaFold-Multimer. *biorxiv* (2022) doi:10.1101/2021.10.04.463034.

40.   Ibrahim, T. *et al.* AlphaFold2-multimer guided high-accuracy prediction of typical and atypical ATG8-binding motifs. *PLoS Biol* **21**, (2023).

41.   Thomasen, F. E. & Lindorff-Larsen, K. Conformational ensembles of intrinsically disordered proteins and flexible multidomain proteins. *Biochemical Society Transactions* vol. 50 541–554 Preprint at https://doi.org/10.1042/BST20210499 (2022).

42.   Rizzuti, B. Molecular simulations of proteins: From simplified physical interactions to complex biological phenomena. *Biochimica et Biophysica Acta - Proteins and Proteomics* vol. 1870 Preprint at https://doi.org/10.1016/j.bbapap.2022.140757 (2022).

43.   Hollingsworth, S. A. & Dror, R. O. Molecular Dynamics Simulation for All. *Neuron* **99**, 1129–1143 (2018).

44.   Delgado, J., Radusky, L. G., Cianferoni, D. & Serrano, L. FoldX 5.0: Working with RNA, small molecules and a new graphical interface. *Bioinformatics* **35**, 4168–4169 (2019).

45.   Barlow, K. A. *et al.* Flex ddG: Rosetta Ensemble-Based Estimation of Changes in Protein-Protein Binding Affinity upon Mutation. *Journal of Physical Chemistry B* **122**, 5389–5399 (2018).

46.   Tiberti, M. *et al.* PyInteraph: a framework for the analysis of interaction networks in structural ensembles of proteins. *J Chem Inf Model* **54**, 1537–1551 (2014).

47.   Ghosh, A. & Vishveshwara, S. A study of communication pathways in methionyl-tRNA synthetase by molecular dynamics simulations and structure network analysis. *Proc Natl Acad Sci U S A* **104**, 15711–15716 (2007).

48.   Watkin, S. A. J. *et al.* On the utility of microfluidic systems to study protein interactions: advantages, challenges, and applications. *European Biophysics Journal* Preprint at https://doi.org/10.1007/s00249-022-01626-9 (2022).

49.   Otzen, D. E., Buell, A. K. & Jensen, H. Microfluidics and the quantification of biomolecular interactions. *Current Opinion in Structural Biology* vol. 70 8–15 Preprint at https://doi.org/10.1016/j.sbi.2021.02.006 (2021).

50. Margreitter, C., Petrov, D. & Zagrovic, B. Vienna-PTM web server: a toolkit for MD simulations of protein post-translational modifications. *Nucleic Acids Res* **41**, W422-6 (2013).

51. Abraham, M. J. *et al.* GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. *SoftwareX* **2**, 19–25 (2015).

52. Piana, S., Lindorff-Larsen, K. & Shaw, D. E. How robust are protein folding simulations with respect to force field parameterization? *Biophys J* **100**, (2011).

53. Fas, B. A. *et al.* The conformational and mutational landscape of the ubiquitin-like marker for autophagosome formation in cancer. *Autophagy* **17**, 2818–2841 (2021).

54. Mackerell, A. D. *et al.* *All-Atom Empirical Potential for Molecular Modeling and Dynamics Studies of Proteins †*. https://pubs.acs.org/sharingguidelines (1998).

55. Khoury, G. A., Thompson, J. P., Smadbeck, J., Kieslich, C. A. & Floudas, C. A. Forcefield-PTM: Ab initio charge and AMBER forcefield parameters for frequently occurring post-translational modifications. *J Chem Theory Comput* **9**, 5653–5674 (2013).

56. Olsson, M. H. M., Søndergaard, C. R., Rostkowski, M. & Jensen, J. H. PROPKA3: Consistent Treatment of Internal and Surface Residues in Empirical pKa Predictions. *J Chem Theory Comput* **7**, 525–37 (2011).

57. Sora, V., Sanchez, D. & Papaleo, E. Bcl-xL Dynamics under the Lens of Protein Structure Networks. *J Phys Chem B* **125**, 4308–4320 (2021).

58. Li, D.-W. W. & Brüschweiler, R. PPM: A side-chain and backbone chemical shift predictor for the assessment of protein conformational ensembles. *J Biomol NMR* **54**, 257–265 (2012).

59. Sahakyan, A. B., Vranken, W. F., Cavalli, A. & Vendruscolo, M. Structure-based prediction of methyl chemical shifts in proteins. *J Biomol NMR* **50**, 331–46 (2011).

60. Papaleo, E., Camilloni, C., Teilum, K., Vendruscolo, M. & Lindorff-Larsen, K. Molecular dynamics ensemble refinement of the heterogeneous native state of NCBD using chemical shifts and NOEs. *PeerJ* **6**, e5125 (2018).

61. Hubbard, S. J. & Thornton, J. M. ,. NACCESS. Preprint at Department of Biochemistry and Molecular Biology, University College London (1993).

62. Kabsch, W. & Sander, C. Dictionary of protein secondary structure: Pattern recognition of hydrogen-bonded and geometrical features. *Biopolymers* **22**, 2577–2637 (1983).

63. Sora, V. *et al.* PyInteraph2 and PyInKnife2 to Analyze Networks in Protein Structural Ensembles. *J Chem Inf Model* **63**, 4237–4245 (2023).

64. Papaleo, E. *et al.* Loop 7 of E2 Enzymes : An Ancestral Conserved Functional Motif Involved in the E2-Mediated Steps of the Ubiquitination Cascade. *PLoS One* **7**, e40786 (2012).

65. Papaleo, E., Renzetti, G. & Tiberti, M. Mechanisms of Intramolecular Communication in a Hyperthermophilic Acylaminoacyl Peptidase: A Molecular Dynamics Investigation. *PLoS One* **7**, e35686 (2012).

66. Papaleo, E., Parravicini, F., Grandori, R., De Gioia, L. & Brocca, S. Structural investigation of the cold-adapted acylaminoacyl peptidase from Sporosarcina psychrophila by atomistic simulations and biophysical methods. *Biochim Biophys Acta Proteins Proteom* **1844**, 2203–2213 (2014).

67. Lambrughi, M., Lucchini, M., Pignataro, M., Sola, M. & Bortolotti, C. A. The dynamics of the β-propeller domain in Kelch protein KLHL40 changes upon nemaline myopathy-associated mutation. *RSC Adv* **6**, 34043–34054 (2016).

68. Kannan, N. & Vishveshwara, S. Identification of side-chain clusters in protein structures by a graph spectral method. *J Mol Biol* **292**, 441–464 (2002).

69. Felline, A., Seeber, M. & Fanelli, F. webPSN v2.0: A webserver to infer fingerprints of structural communication in biomacromolecules. *Nucleic Acids Res* **48**, W94–W103 (2020).

70. Tiberti, M. *et al.* MutateX: an automated pipeline for in silico saturation mutagenesis of protein structures and structural ensembles. *Brief Bioinform* **23**, bbac074 (2022).

71. Degn, K. *et al.* Cancer-related Mutations with Local or Long-range Effects on an Allosteric Loop of p53. *J Mol Biol* **434**, 167663 (2022).

72. Arnaudi, M. *et al.* MAVISp: Multi-layered Assessment of VarIants by Structure for proteins. *bioRxiv* (2022) doi:10.1101/2022.10.22.513328.

73. Tate, J. G. *et al.* COSMIC: The Catalogue Of Somatic Mutations In Cancer. *Nucleic Acids Res* **47**, D941–D947 (2019).

74. Gao, J. *et al.* Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal. *Sci Signal* **6**, pl1 (2013).

75. Landrum, M. J. *et al.* ClinVar: improvements to accessing data. *Nucleic Acids Res* **48**, D835–D844 (2020).

76. Varadi, M. *et al.* AlphaFold Protein Structure Database: Massively expanding the structural coverage of protein-sequence space with high-accuracy models. *Nucleic Acids Res* **50**, D439–D444 (2022).

77. Burke, D. F. *et al.* Towards a structurally resolved human protein interaction network. *Nat Struct Mol Biol* **30**, (2023).

78. Ioannidis, N. M. *et al.* REVEL: An Ensemble Method for Predicting the Pathogenicity of Rare Missense Variants. *Am J Hum Genet* **99**, 877–885 (2016).

79. Munro, D. & Singh, M. DeMaSk: A deep mutational scanning substitution matrix and its use for variant impact prediction. *Bioinformatics* **36**, 5322–5329 (2020).

80. Frazer, J. *et al.* Disease variant prediction with deep generative models of evolutionary data. *Nature* **599**, 91–95 (2021).

81. Cheng, J. *et al.* Accurate proteome-wide missense variant effect prediction with AlphaMissense. *Science (1979)* **381**, eadg7492 (2023).

82. Daura, X. *et al.* Peptide folding: When simulation meets experiment. *Angewandte Chemie - International Edition* **38**, 236–240 (1999).

83. Nygaard, M. *et al.* The mutational landscape of the oncogenic MZF1 SCAN domain in cancer. *Front Mol Biosci* **3**, (2016).

84. Robustelli, P., Stafford, K. A. & Palmer, A. G. Interpreting Protein Structural Dynamics from NMR Chemical Shifts. *J Am Chem Soc* **134**, 6365–6374 (2012).

85. Case, D. a. Chemical shifts in biomolecules. *Curr Opin Struct Biol* **23**, 172–179 (2013).

86. Palmer, A. G. Enzyme dynamics from NMR spectroscopy. *Acc Chem Res* **48**, 457–65 (2015).

87. Berjanskii, M. V. & Wishart, D. S. Unraveling the meaning of chemical shifts in protein NMR. *Biochim Biophys Acta Proteins Proteom* **1865**, 1564–1576 (2017).

88. Li, D. W. & Brüschweiler, R. Certification of molecular dynamics trajectories with NMR chemical shifts. *Journal of Physical Chemistry Letters* **1**, 246–248 (2010).

89. Di Rita, A. *et al.* HUWE1 E3 ligase promotes PINK1/PARKIN-independent mitophagy by regulating AMBRA1 activation via IKKα. *Nat Commun* **9**, 3755 (2018).

90. Christensen, N. J. & Kepp, K. P. Accurate stabilities of laccase mutants predicted with a modified FoldX protocol. *J Chem Inf Model* **52**, 3028–3042 (2012).

91. Caldararu, O., Blundell, T. L. & Kepp, K. P. A base measure of precision for protein stability predictors: structural sensitivity. *BMC Bioinformatics* **22**, 1–14 (2021).
92. Pucci, F., Bernaerts, K. V., Kwasigroch, J. M. & Rooman, M. Quantification of biases in predictions of protein stability changes upon mutations. *Bioinformatics* **34**, 3659–3665 (2018).
93. Buß, O., Rudat, J. & Ochsenreither, K. FoldX as Protein Engineering Tool: Better Than Random Based Approaches? *Comput Struct Biotechnol J* **16**, 25–33 (2018).
94. Burke, W., Parens, E., Chung, W. K., Berger, S. M. & Appelbaum, P. S. The Challenge of Genetic Variants of Uncertain Clinical Significance: A Narrative Review. *Annals of Internal Medicine* vol. 175 Preprint at https://doi.org/10.7326/M21-4109 (2022).
95. Naruse, H. *et al.* Mutational analysis of familial and sporadic amyotrophic lateral sclerosis with OPTN mutations in Japanese population. *Amyotrophic Lateral Sclerosis* **13**, 562–566 (2012).
96. Zijlstra, N. *et al.* Rapid Microfluidic Dilution for Single-Molecule Spectroscopy of Low-Affinity Biomolecular Complexes. *Angewandte Chemie* **129**, 7232–7235 (2017).

Author Contributions (CRediT Classification)

Conceputalization: EP. Data Curation: EP, MU, OA, VS, ML. Formal Analysis: MU, OA, EP Funding Acquisition: EP. Investigation: EP, OA, MU, VS, ML. Methodology: OA, EM, ML, MT. Project administration: EP. Resources: EP. Software: MT, ML, VS, MU, EP. Supervision: EP, ML, EM. Validation: MU, ML, EP. Visualization: MU, ML, OA, VS, EP. Writing – Original Draft: EP, OA, MU, VS, ML, EM. Writing – Review and Editing: All the coauthors.

Acknowledgments

Our research has been supported by Danmarks Grundforskningsfond (DNRF125), Carlsberg Foundation Distinguished Fellowship (CF18-0314), Hartmanns Fond (R241-A33877) and NovoNordisk Fonden Bioscience and Basic Biomedicine (NNF20OC0065262) to EP group. Part of the calculations have been supported by a ISCRA CINECA Grant, DECI 16h and a EuroHPC Regular Access Grant (EHPC-REG-2023R01-051) on Discoverer. The calculations described in this paper were performed using the DeiC National Life Science Supercomputer at DTU. The funders had no role in study design, data collection or analysis, decision to publish, or manuscript preparation.
We would like to thank Kirsten Frederisken for statistical advice, as well as Burcu Aykac Fas, Mukesh Kumar, and Željka Sanader Maršić for their involvement in the early stages of this project.

A

OPTN

31 — 97 NEMO-like
1

LZ 141 161
169 177 185
LIR
CC 445 — 502 CC
UBD 547 577
ZnF

I181
E180
F178
HP2
V179
HP1
pS177
pS174
pS173
pS171
pS170

B

LIR constructs used for MD simulations

LC3B$_{1-120}$-OPTN$_{169-185}$ 169-NSSGSSEDSFVEIRMAE-185
LC3B$_{1-120}$-OPTN$_{158-191}$ 158-LGIVSELQLKLNSSGSSEDSFVEIRMAEGEAEGS-191
LC3B$_{1-120}$-p1OPTN$_{169-185}$ 169-NSSGSSEDpSFVEIRMAE-185
LC3B$_{1-120}$-p1OPTN$_{158-191}$ 158-LGIVSELQLKLNSSGSSEDpSFVEIRMAEGEAEGS-191
LC3B$_{1-120}$-p5OPTN$_{169-185}$ 169-NpSpSGpSpSEDpSFVEIRMAE-185
LC3B$_{1-120}$-p5OPTN$_{158-191}$ 158-LGIVSELQLKLNpSpSGpSpSEDpSFVEIRMAEGEAEGS-191

C

pS171
pS170
pS173
pS174
pS177
F178
I181