Full Title: *Wolbachia* **enhances the survival of** *Drosophila* **infected with fungal pathogens**

Authors:

Jessamyn I. Perlmutter[a*], Aylar Atadurdyyeva[a], Margaret E. Schedl[a], & Robert L. Unckless[a]

Affiliations:

[a]Department of Molecular Biosciences, University of Kansas, Lawrence, Kansas, USA

*Correspondence to:

Jessamyn I. Perlmutter (jessie.perlmutter@gmail.com)

Abstract:

    Wolbachia bacteria of arthropods are at the forefront of basic and translational research on multipartite host-symbiont-pathogen interactions. These microbes are vertically inherited from mother to offspring via the cytoplasm. They are the most widespread endosymbionts on the planet due to their infamous ability to manipulate the reproduction of their hosts to spread themselves in a population, and to provide a variety of fitness benefits to their hosts. Importantly, some strains of *Wolbachia* can inhibit viral pathogenesis within and between arthropod hosts. Mosquitoes carrying the *w*Mel *Wolbachia* strain of *Drosophila melanogaster* have a greatly reduced capacity to spread viruses like dengue and Zika to humans. Therefore, *Wolbachia* are the basis of several global vector control initiatives. While significant research efforts have focused on viruses, relatively little attention has been given to *Wolbachia*-fungal interactions despite the ubiquity of fungal entomopathogens in nature. Here, we demonstrate that *Wolbachia* increase the longevity of their *Drosophila melanogaster* hosts when challenged with a spectrum of yeast and filamentous fungal pathogens. We find that this pattern can vary based on host genotype, sex, and fungal species. Further, *Wolbachia* correlates with higher fertility and reduced pathogen titers during initial fungal infection, indicating a significant fitness benefit. This study demonstrates *Wolbachia*'s role in diverse fungal pathogen interactions and determines that the phenotype is broad, but with several variables that influence both the presence and strength of the phenotype. These results enhance our knowledge of the strategies *Wolbachia* uses that likely contribute to such a high global symbiont prevalence.

Importance:

Wolbachia bacteria of arthropods are at the forefront of global initiatives to fight arthropod-borne viruses. Despite great success in using the symbiont to fight viruses, little research has focused on Wolbachia-fungal interactions. Here, we find that Wolbachia of Drosophila melanogaster, the same strain widely used in antiviral initiatives, can also increase the longevity of flies systemically infected with a panel of yeast and filamentous fungal pathogens. The symbiont also partially increases host fertility and reduces fungal titers during early infection, indicating a significant fitness benefit. This represents a major step forward in Wolbachia research since its pathogen blocking abilities can now be extended to a broad diversity of another major branch of microbial life. This discovery may inform basic research on pathogen blocking and has potential translational applications in areas including biocontrol in agriculture.

Introduction:

Microbe-host symbioses are ubiquitous in nature and exhibit a broad range of relationships from facultative parasitism to obligate mutualism[1,2]. Microbial symbionts of arthropods in particular exhibit a striking array of phenotypes in their hosts[2], ranging from provision of nutrients[3] to protection from parasitoids[4] to death of the host's offspring[5]. One microbial symbiont, Wolbachia pipientis, is an exemplary case of a microbe with diverse symbiont-host interactions. Wolbachia are obligate intracellular bacteria found in germline and somatic tissues of diverse arthropods and are almost exclusively inherited vertically through the cytoplasm of infected mothers[6]. They are found in an estimated 40-52% of all arthropod species on Earth[7,8], making them the most widespread endosymbiont and "the world's greatest

pandemic"[9,10]. There is such genetic diversity that there are 18 recognized *Wolbachia*

supergroups[11-13]. Some can act as "reproductive parasites" that manipulate host reproduction to

facilitate their spread by enhancing the relative fitness of infected female transmitters[14]. Others

are obligate mutualists necessary for host oogenesis or early development[15]. Depending on

context, *Wolbachia* can use their diverse genetic toolkit to engage in a variety of interactions

with their hosts. These interactions have had immense impacts on both basic and applied

research in many fields, including utility in fighting human diseases vectored or caused by

insects and nematodes and to an understanding of the role of symbionts in shaping host

evolutionary processes[6,16-18].

65   *Wolbachia*'s employment of such diverse host interactions has been critical to its global

success, however, these phenotypes do not fully explain how widespread *Wolbachia* is. Indeed,

while some strains are reproductive manipulators (enhancing the fitness of the infected

matriline)[5,10,19-21] or obligate mutualists (enhancing the fitness of all hosts)[12,22-24], but many are

not, even among organisms that have been phenotypically assessed[25]. Some strains also exhibit

no reproductive parasitism in and provide no currently known fitness benefit[26,27]. Further, those

that are reproductive manipulators can vary both in the effect size of their phenotype (either

weak or strong induction[28-31]) and in their frequency in the population (high or low[32-35]). Even

when reproductive phenotypes or benefits are known, they are often context-dependent and vary

based on factors such as temperature[36-39], symbiont density[40,41], or host genetic background[42].

Further, in the wild, vertical transmission fidelity of *Wolbachia* is not 100%[27,43,44], making the

basis of the symbiont's maintenance in populations even less clear. For many years, a question of

significant focus in the field has been how it is that *Wolbachia* is so widespread[45], particularly

given the fact that we have not identified a clear host fitness benefit of the symbiont for all

strains or contexts. Research over the years has identified some contributing factors such as

nutritional contributions of the symbiont to the host[46,47], as well as rescue of host deficiencies

like mutations in the key sex development regulator *sex-lethal*[48,49] and germline stem cell self-

renewal and differentiation deficiencies[50]. Yet these contributing factors do not fully answer the

question, and other factors must be involved.

One such crucial and somewhat common beneficial *Wolbachia*-host interaction was

discovered through work on an early theory that *Wolbachia*'s prevalence could be based on an

ability to inhibit pathogens, thereby conferring a significant fitness benefit to the host[51-53]. The

rationale was based partially on the observation that facultative infection (as opposed to obligate

mutualism) is relatively common with *Wolbachia* infections, but with few accompanying known

benefits to explain their frequency. It was also partially based on an observation that *Wolbachia*

infection correlated with host resistance to infection with the common *Drosophila* C virus

(DCV)[51]. Two foundational early studies on this topic demonstrated that *Drosophila*

melanogaster flies with their native *Wolbachia* strain exhibit greater longevity on the order of

days to weeks of increased life when infected with several common arthropod RNA viruses[51,54].

This coincides with reduced viral load in *Wolbachia*-viral co-infection, which increases host

fitness and survival likelihood though reduced pathogen burden. These and latter studies also

demonstrated that the phenotype could be induced by some additional *Wolbachia* strains or in

additional host genetic backgrounds or species, but that the effect was largely restricted to RNA

viruses (not DNA viruses)[55]. Finally, and crucially, some *Wolbachia* strains are also able to

inhibit the transmission of viral (and some other) pathogens to new host individuals, including

pathogens spread by mosquitoes to humans[51,54,56,57]. This ability of the symbiont to protect its

host from viruses is considered a major factor contributing to *Wolbachia*'s success.

Virus pathogen blocking has therefore become an eminent area of *Wolbachia* research

not only for its broad applicability across the symbiont genus and importance to basic biology,

but also for its translational potential. For example, *Aedes aegypti* mosquitoes and other common

human disease vectors exhibit significantly reduced capacities to transmit parasites like malaria[57]

or viruses like Zika[56], dengue[58,59], yellow fever[60], or chikungunya[61] to humans when they carry

certain strains of *Wolbachia*. This feature has made *Wolbachia* central to global efforts to reduce

disease through groups like MosquitoMate[62] and the World Mosquito Program[63]. These

programs rear *Wolbachia*-positive mosquitoes on a massive scale and release mosquitoes into the

wild. One strategy is to release infected females that then outcompete local *Wolbachia*-negative

counterparts and replace them with a disease-resistant population. Collaborative efforts through

this program across four continents have resulted in stable, wild *Wolbachia*-positive populations

in many locations and significant reductions in disease[58,64]. Arthropod vector-borne diseases are

responsible for millions of illnesses, deaths, and contribute to significant inequality around the

world[65], and the use of *Wolbachia*-positive mosquitoes is one of our most promising solutions[66-

68].

In contrast with all of this progress on viruses, comparatively little research has been

done on *Wolbachia* interactions with non-viral pathogens[57,69]. This is despite the extraordinary

genetic and phenotypic diversity of *Wolbachia* symbioses that indicate the likelihood of broader

protective abilities. Early theory predicted that pathogen protection could increase the relative

fitness of hosts with *Wolbachia* compared to those without, contributing to maintenance and

spread of the symbiont[32], and this was one of the original bases for investigations into viral

pathogen blocking, and could apply to many other types of pathogens too[51,54]. However, one

particular gap in the research is the potential for *Wolbachia* to inhibit fungal pathogens. Fungal

pathogens of arthropods are common in the wild[70], yet few studies have investigated the

interactions between *Wolbachia*, hosts, and fungal pathogens, and the studies that do present

different results. One early study showed no effect of *w*Ri *Wolbachia* strain infection on survival

from topical cuticle infection of the common insect fungal pathogen, *Beauveria bassiana*, in *D.*

simulans male flies[71]. Another reported higher survival of *D. melanogaster* female flies with

their native *w*Mel *Wolbachia* symbiont after immersion in a suspension of *B. bassiana*[72].

Conversely, a third study on infection of female spider mites in topical contact with *B. bassiana*

or *Metarhizium* fungal pathogens indicated that *Wolbachia* may actually increase mortality of the

host with fungal infection[73]. A fourth investigated the effect of *Wolbachia* on injection with two

Beauveria pathogens on *Aedes albopictus* and *Culex pipiens* mosquitoes[74]. This study found no

enhancement in host survival with the symbiont, but reported some putative differences in host

immune gene expression and reduced fungal load in some contexts. Finally, a recent study

indicates that the *w*Pni strain of *Pentalonia* aphids may result in increased survival of hosts

infected topically with the specialized fungal pathogen, *Pandora neoaphidis*[75]. Thus, there have

been several investigations, with some prior reports indicating that *Wolbachia* may interact with

fungal pathogens in some contexts.

Despite this research, the question of *Wolbachia*'s ability to interact with fungal

pathogens on a larger scale remains unanswered. It is unclear how broad the fungal blocking

ability is in terms of host, symbiont, and pathogen factors, and if the phenotype is likely to be

common or not. This difficulty is because the studies draw different conclusions from different

contexts. These prior reports have used different host species, host sexes, *Wolbachia* strains,

pathogen species, pathogen concentrations, routes of pathogen infection, and been measured by

different host fitness and health assays or conducted over different lengths of time[71-75]. These

factors make it difficult to compare across studies, as there are multiple variables between any

two publications. Further, due to the small number of studies, limited parameters have been

tested thus far. Thus, the breadth of *Wolbachia*-fungal interactions is unclear, as comparison

between studies is difficult and there is limited published data.

To begin to fill this gap in knowledge, we conducted a series of systemic fungal infection

assays using *D. melanogaster* flies with the *w*Mel *Wolbachia* symbiont in the context of several

host and pathogen variables. Notably, *w*Mel is the initial strain that was reported to inhibit

viruses and mosquitoes transinfected with this symbiont strain are the basis of many of the global

vector control initiatives[51,54,58]. This approach addresses several outstanding research questions

in this area: (i) can *Wolbachia* inhibition of fungal pathogenesis be confirmed when tested in

various contexts, (ii) how broad is this protective phenotype within one *Wolbachia* strain, and

(iii) do factors such as fungal pathogen species, fungal pathogen types (filamentous vs yeast),

host sex, and host genetic background contribute to the *Wolbachia*-fungal pathogen interaction.

Here we report that *Wolbachia* is indeed capable of significantly increasing the longevity and

reproductive fitness of flies infected with a wide variety of fungal pathogens, and the phenotype

is influenced by several host and pathogen factors.

Results:

***Wolbachia's* association with an increase in longevity of flies infected with filamentous**

fungi is dependent on genetic background and host sex

To test the breadth and ability of *Wolbachia* to inhibit fungal pathogenesis in flies, a

series of systemic infection assays were conducted. Experiments were performed with two

different *Drosophila melanogaster* host background lines infected with their native *w*Mel

Wolbachia. The host strains themselves have diverse origins: the w^{1118} line was collected in

California, USA and was reported in 1985[76], and the w^k line was collected in 1960 in Karsnäs,

Sweden[77]. Different collection origins together with Illumina sequencing showing a high number

of SNPs between the *D. melanogaster* lines indicate the lines represent genetically diverse host

backgrounds. Each strain has its own natural *Wolbachia* along with genetically identical

counterpart strains that were previously treated with antibiotics to remove the symbiont. Thus,

we tested four strains total: w^{1118} with *Wolbachia*, w^{1118} without *Wolbachia*, w^k with *Wolbachia*,

and w^k without *Wolbachia*. Whole genome sequencing of the *Wolbachia* symbionts of each

strain indicates that they are highly similar despite disparate origins, with only a single divergent

SNP across the entire genome. This SNP is a silent (synonymous) polymorphism in a membrane

transporter of the major facilitator superfamily, which transports small solutes[78]. Thus, the vast

majority of genetic differences between strains can be attributed to the host, and most phenotypic

differences are therefore likely due to the host as well.

To determine if *Wolbachia* can increase the longevity of flies infected with fungi as

hypothesized, systemic infections were performed with both sexes of all four strains against a

variety of pathogens. We started with several *Aspergillus* and *Fusarium* filamentous fungal

species that infect both arthropods and humans: *Aspergillus fumigatus*, *Aspergillus flavus*,

Fusarium oxysporum, and *Fusarium graminaerum* (Figure 1). Survival was scored daily for

three weeks, as differences in survival were broadly apparent across treatment groups for most

pathogens by this point. The data revealed several key results. First, *Wolbachia* was associated

with significantly greater survival across the trial period in many contexts. In the w^k background,

Wolbachia-positive flies had higher survival for all pathogens except *Fusarium oxysporum*,

which was only significant when comparing within just males (Figure 1). Second, genetic

backgrounds played a significant role in the infection outcomes. Indeed, *Wolbachia* was not a

significant predictor of increased longevity for any of the pathogens in the w^{1118} host

background, except when considering sex (Figure S1). Third, sex is repeatedly a significant

factor in survival outcomes for some pathogens. Males alone had a significant increase in

longevity for *Aspergillus fumigatus* and *Fusarium oxysporum* for both genetic backgrounds

(Figures 1a,c & S1a,c), with a statistically significant *Wolbachia* x sex interaction for *A.*

fumigatus in the w^k background and *Fusarium oxysporum* in the w^{1118} background (Figures 1a,

S1c). Fourth, the host strains had generally different overall susceptibilities to fungal infection,

with w^k generally having lower survival than w^{1118} in both *Wolbachia*-positive and -negative

contexts (Figures 1 & S1, mean 51.1% death for all pathogen infections combined in the w^{1118}

background by day 21, 60.4% death in the w^k background). In particular, there is a significant

Wolbachia x genotype interaction for *Aspergillus flavus* (*p=0.043, Table S1).

Figure 1. *Wolbachia* **increases the longevity of flies of the *w^k* background line infected with several filamentous fungal pathogens.** Flies of each given background and sex were systemically infected with the indicated pathogen. Infections were performed with either (a) *Aspergillus fumigatus*, (b) *Aspergillus flavus*, (c) *Fusarium oxysporum*, or (d) *Fusarium graminaerum*. Infections of all groups were performed side-by-side, along with those of the *w^1118* background line (Figure S1), with at least two blocks of infections performed on different days. Each line represents a total of 60 flies. Sham controls were performed with sterile 20% glycerol. Full statistics, available in Table S1, were done with a Cox mixed effects model. Controls are the same in all panels and in Figure 2a because they were performed concurrently in the same background.

Figure S1. *Wolbachia* does not increase the longevity of flies of the *w^1118^* background line infected with several filamentous fungal pathogens. Flies of each given background and sex were systemically infected with the indicated pathogen. Infections were performed with either (a) *Aspergillus fumigatus*, (b) *Aspergillus flavus*, (c) *Fusarium oxysporum*, or (d) *Fusarium graminaerum*. Infections of all groups were performed side-by-side, along with those of the *w^k^* background line (Figure 1), with at least two blocks of infections performed on different days. Each line represents a total of 60 flies. Sham controls were performed with sterile 20% glycerol. Full statistics, available in Table S1, were done with a Cox mixed effects model. Controls are the same in all panels and in panel S2a because they were performed concurrently in the same background.

Wolbachia can increase the longevity of flies infected with filamentous fungal

entomopathogens

To determine if *Wolbachia* could also increase longevity of flies infected with common

filamentous fungal insect pathogens (entomopathogens), we performed systemic infections with

Beauveria bassiana, *Metarhizium anisopliae*, *Clonostachys rosea*, and *Trichoderma atroviride*.

Beauveria and *Metarhizium* in particular are ubiquitous insect pathogens and are the subject of

extensive research in biocontrol of pests in particular[79], while *Clonostachys* and *Trichoderma* are

also globally widespread and have received recent attention in biocontrol as well[80-82]. The latter

two were collected from mosquitoes, and are thus of potential relevance to mosquito biology

(Table S2). Similar to the results of the pathogens in Figures 1 & S1, *Wolbachia* increased

longevity in many, but not all fungal infection contexts (Figures 2 & S2). Namely, *Wolbachia*

significantly increased longevity for *Beauveria bassiana* and *Clonostachys rosea* in the w^k

background (Figure 2a,c), and *Beauveria bassiana* and *Metarhizium anisopliae* in the w^{1118}

background (Figure S2a,b). Thus, there is some positive longevity effect of the symbiont in

either background, not just w^k, but the effect depends on the pathogen. Further, sex was also a

factor with a significant effect for *Beauveria bassiana* and *Metarhizium anispoliae* in the w^k

background (Figure 2a,b) and *Metarhizium anisopliae* and *Trichoderma atroviride* in the w^{1118}

background (Figure S2b,d). Additionally, as with previous infections, w^k was broadly more

susceptible to infection as flies generally died earlier and at higher rates than their w^{1118}

counterparts (Figures 2 & S2, mean 70.3% death for all entomopathogen infections combined in

the w^{1118} background by day 21, 85.8% death in the w^k background).

Figure 2. *Wolbachia* increases the longevity of flies of the *w^k* background line infected with certain filamentous fungal entomopathogens. Flies of each given background and sex were systemically infected with the indicated pathogen. Infections were performed with either (a) *Beauveria bassiana*, (b) *Metarhizium anisopliae*, (c) *Clonostachys rosea*, or (d) *Trichoderma atroviride*. Infections of all groups were performed side-by-side, along with those of the *w^1118* background line (Figure S2), with at least two blocks of infections performed on different days. Each line represents a total of 60 flies. Sham controls were performed with sterile 20% glycerol. Full statistics, available in Table S1, were done with a Cox mixed effects model. Controls for panel 2a are the same for Figure 1, and the panels in 2b-d are the same because they were performed concurrently in the same background.

255
256
Figure S2. *Wolbachia* increases the longevity of *w[1118]* background line flies infected with certain filamentous fungal
entomopathogens. Flies of each given background and sex were systemically infected with the indicated pathogen. Infections
were performed with either (a) *Beauveria bassiana*, (b) *Metarhizium anisopliae*, (c) *Clonostachys rosea*, or (d) *Trichoderma*
atroviride. Infections of all groups were performed side-by-side, along with those of the *w[k]* background line (Figure 2), with at
least two blocks of infections performed on different days. Each line represents a total of 60 flies. Sham controls were performed
with sterile 20% glycerol. Full statistics, available in Table S1, were done with a Cox mixed effects model. Controls for panel
S2a are the same for Figure S1, and the panels in S2b-d are the same because they were performed concurrently in the same
background.
265

***Wolbachia* can increase the longevity of flies infected with yeasts**

To test if *Wolbachia* could also increase the longevity of flies infected with yeast, we

performed systemic infections using *Candida auris, Candida glabrata*, and *Galactomyces*

pseudocandidus. For *Candida* pathogens, *Wolbachia* significantly increased longevity of w^k

background flies. In contrast, *Wolbachia* did not significantly increase longevity for any of the

yeast pathogens in the w^{1118} background. Further, sex was not a significant factor in any of the

yeast infections for either background. However, flies of the w^k background again were more

broadly susceptible to infection based on higher overall mortality (mean 40% death for all yeast

infections combined in the w^{1118} background by day 21, 58.3% death in the w^k background).

Key

🟢 *Wolbachia* (+) Male

🟩 *Wolbachia* (-) Male

🟣 *Wolbachia* (+) Female

🟪 *Wolbachia* (-) Female

⚪ Sham Control

Figure 3. *Wolbachia* increases the longevity of flies of the w^k background line infected with yeast pathogens. Flies of each given background and sex were systemically infected with the indicated pathogen. Infections were performed with either (a) *Candida auris*, (b) *Candida glabrata*, or (c) *Galactomyces pseudocadidus*. Infections of all groups were performed side-by-side, along with those of the w^{1118} background line (Figure S3), with at least two blocks of infections performed on different days. Each line represents a total of 60 flies. Sham controls were performed with sterile 20% glycerol. Full statistics, available in Table S1, were done with a Cox mixed effects model. Controls are the same in all panels and because they were performed concurrently in the same background.

Key

● *Wolbachia* (+) Male
● *Wolbachia* (-) Male
● *Wolbachia* (+) Female
● *Wolbachia* (-) Female
● Sham Control

287
Figure S3. *Wolbachia* increases the longevity of flies of the *w^1118* background line infected with yeast pathogens. Flies of
each given background and sex were systemically infected with the indicated pathogen. Infections were performed with either (a)
Candida auris, (b) *Candida glabrata*, or (c) *Galactomyces pseudocadidus*. Infections of all groups were performed side-by-side,
along with those of the *w^1118* background line (Figure 3), with at least two blocks of infections performed on different days. Each
line represents a total of 60 flies. Sham controls were performed with sterile 20% glycerol. Full statistics, available in Table S1,
were done with a Cox mixed effects model. Controls are the same in all panels and because they were performed concurrently in
the same background.
295

Wolbachia can partially rescue female fertility reduction after infection

To assess whether *Wolbachia* impacts fitness of hosts early in fungal infection, female

flies were systemically infected with *B. bassiana* because *Wolbachia* significantly increased

longevity for all treatment groups with this pathogen (Figures 2a, S2a). Egg laying and egg

hatching rates were quantified for the first 3 days post infection for flies with either the infection

or a sham control (Figures 4, S4). Although both *Wolbachia*-positive and *Wolbachia*-negative

flies laid similar numbers of eggs in the w^k background without treatment, and although the

overall egg-laying was lower in *B. bassiana*-infected flies, *Wolbachia* significantly increased

egg-laying with fungal infection (Figure 4). This was also true in the w^{1118} background (Figure

S4). In contrast, the percentage of eggs hatched was not greatly impacted by either *Wolbachia* or

fungal infection in either background (Figures 4b, S4b).

**Figure 4. *Wolbachia* increases the number of eggs laid but not the percentage of eggs hatched post-*B. bassiana* infection in
the w^k background line.** Female flies were systemically infected with *B. bassiana* or treated with a sham control. The flies then
laid eggs for 3 days post-infection. (a) Numbers of eggs laid. (b) Proportion of eggs hatched. Each dot represents the total
offspring of a single female, with an overall mean of 35 eggs laid. The boxes indicate the interquartile range. Outer edges of the
box indicate 25th (lower) and 75th (upper) percentiles and the middle line indicates 50th percentile (median). Whiskers represent
maximum and minimum ranges of data within 1.5 times the interquartile range of the box. Statistics are based on a logistic
regression (Table S1). The entire experiment was performed twice, and graphs represent a combination of data from both blocks.
315
316
317

318
319
Figure S4. ***Wolbachia* increases the number of eggs laid but not the percentage of eggs hatched post-*B. bassiana* infection**
in the *w¹¹¹⁸* background line. Female flies were systemically infected with *B. bassiana* or treated with a sham control. The flies
then laid eggs for 3 days post-infection. (a) Numbers of eggs laid. (b) Proportion of eggs hatched. Each dot represents the total
offspring of a single female, with an overall mean of 48 eggs laid. The boxes indicate the interquartile range. Outer edges of the
box indicate 25th (lower) and 75th (upper) percentiles and the middle line indicates 50th percentile (median). Whiskers represent
maximum and minimum ranges of data within 1.5 times the interquartile range of the box. Statistics are based on a logistic
regression (Table S1). The entire experiment was performed twice, and graphs represent a combination of data from both blocks.
327

***Wolbachia* associates with reduced fungal titer after infection**

To determine if enhanced longevity is likely based on killing or reduction of pathogen

(immune resistance) vs tolerance and maintenance of the pathogen (immune tolerance), and to

determine if reproductive benefits with fungal infection in Figures 4 & S4 can be attributed to

reduced pathogen load, we measured fungal and *Wolbachia* titers over time in *B. bassiana-*

infected females (Figure 5). We measured over the first 24 h because this is before flies begin to

die and many essential early host molecular responses to pathogen infection begin by this

timepoint during infection[83,84]. We find that *Wolbachia* titer stays constant over the 24 h period

(Figure 5a) and that pathogen load is not significantly different between lines immediately post-

infection (Figure 5b). Thus, both *Wolbachia*-positive and -negative flies are receiving similar

starting amounts of pathogen. However, by 24 h post-infection, we see that pathogen load is

reduced in the *Wolbachia*-positive flies compared to those without *Wolbachia*. This trend holds

true in the w^{1118} background as well.

Figure 5. *Wolbachia* associates with reduced pathogen titer after infection with no significant change in *Wolbachia* titer in
w^k **flies.** Female flies were systemically infected with the indicated fungal pathogen and pathogen titers were measured both
immediately after infection and 24 h post-infection. Dots represent pools of 3 infected females. (a) *Wolbachia* titers. (b) *B.*
bassiana titers. The boxes indicate the interquartile range. Outer edges of the box indicate 25th (lower) and 75th (upper)
percentiles and the middle line indicates 50th percentile (median). Whiskers represent maximum and minimum ranges of data
within 1.5 times the interquartile range of the box. Statistics are based on a logistic regression (Table S1). The entire experiment
was performed twice, and graphs represent a combination of data from both blocks.
350
351

a. *Wolbachia* titers

b. Fungal titers

Time: ns, p=0.244

Wolbachia: ns, p=0.70
Time: *, p=0.027
Wolbachia x Time: **p=0.0012

Key

● *Wolbachia*

● No *Wolbachia*

Figure S5. *Wolbachia* associates with reduced pathogen titer after infection with no significant change in *Wolbachia* titer in *w^{1118}* flies. Female flies were systemically infected with the indicated fungal pathogen and pathogen titers were measured both immediately after infection and 24 h post-infection. Dots represent pools of 3 infected females. (a) *Wolbachia* titers. (b) *B. bassiana* titers. The boxes indicate the interquartile range. Outer edges of the box indicate 25th (lower) and 75th (upper) percentiles and the middle line indicates 50th percentile (median). Whiskers represent maximum and minimum ranges of data within 1.5 times the interquartile range of the box. Statistics are based on a logistic regression (Table S1). The entire experiment was performed twice, and graphs represent a combination of data from both blocks.

Discussion:

In the 15 years since the discovery of *Wolbachia*-based virus inhibition, there has been significant research into the mechanism and translational applications of the phenotype[51,54,55,64]. However, comparatively little attention has been given to the potential for *Wolbachia* to interact with other types of pathogens, including fungi. Prior research gave contrasting results either suggesting there was a *Wolbachia*-fungal infection interaction[72,75] or not[71,73,74]. However, these previous studies were performed in different contexts with many different variables between them. Thus, the breadth of *Wolbachia*'s ability to interact with fungal pathogens as well as

identification of factors that influence the putative phenotype have remained unclear. Given the

likely importance of fungal interactions to the basic biology of *Wolbachia* and potential

applications in areas like agriculture, these are important research topics to address. For example,

the large field trials that release *Wolbachia*-positive mosquitoes to combat arthropod-transmitted

viruses rely on *Wolbachia*'s reproductive manipulations of the host to help spread itself in the

wild[64]. The *Wolbachia*-positive mosquitoes must reach a sometimes unstable equilibrium level to

reliably spread[85], which could be altered by fitness impacts induced through fungal infection.

Further, many agricultural fungal diseases are vectored by arthropods and Wolbachia could be

used as a tool to combat disease spread. To begin filling this gap, we sought here to test

Wolbachia-fungus interactions by systemically infecting the model host *Drosophila*

melanogaster with a panel of fungal pathogens and measuring host longevity. We included

several variables that we hypothesized might be important factors in any potential pathogen-

blocking phenotype, including host genotype, host sex, and pathogen species. We then tested the

effect of *Wolbachia* on host fertility and pathogen load when infected or not with fungus.

The main conclusions that can be drawn from the results are that the *w*Mel strain of *D.*

melanogaster has a broad, but variable ability to inhibit fungal pathogenesis and that both host

and pathogen variables significantly contribute to infection outcomes. Across the systemic

infection assays (Figures 1-3, S1-S3), we found a variety of patterns in the results. There are

cases where *Wolbachia*-positive flies live significantly longer with fungal infection in all tested

contexts, such as *B. bassiana* (Figures 2a, S2a). Notably, this is in agreement with one prior

study that showed *D. melanogaster* females with *Wolbachia* lived longer when dipped in a

suspension of the same pathogen[72], suggesting that the phenotype may hold with multiple

different infection routes as well. There were also cases where *Wolbachia* significantly increased

host longevity in only one host background, such as the *Aspergillus* and *Fusarium* pathogens

(Figures 1, S1), *C. rosea* (Figures 2c, S2c), and *Candida* pathogens (Figures 3, S3), examples for

which *Wolbachia* was only significant in the w^k background. In contrast, *Wolbachia* was

significant in only the w^{1118} background for *M. anisopliae* infection (Figures 2b, S2b), so either

host genotype can result in a statistically significant outcome while the other does not. However,

and on a related note, the effect size of *Wolbachia* on host survival may be small in a given

context and may lead to lower power to detect the differences with our sample sizes, like *M.*

anisopliae in w^k (Figure 2b) or *F. graminaerum* in w^{1118} (Figure S1d). In contrast, there was one

case where the infection outcome was not significant in any context, with the *T. atroviride*

pathogen (Figures 2d, S2d), so there may not be an interaction with all pathogens. Further, there

were no cases of increased mortality with *Wolbachia*-fungal co-infection, as was suggested in a

prior study with fungal pathogens in *Wolbachia*-positive spider mites[73]. Thus, broadly speaking,

both pathogen species and host genetics are factors that significantly associate with *Wolbachia*-

fungus co-infection outcomes. These patterns suggest that the mechanism(s) of protection are

likely not universal to fungal infection, and that host factors are likely involved.

Notably, host sex was a significant predictor of infection outcome in several cases as a

standalone variable. For example, females had increased longevity compared to males with *B.*

bassiana and *M. anisopliae* infection in w^k hosts (Figures 2a,b) and *M. anisopliae* infection in

w^{1118} hosts (Figure S2b), regardless of *Wolbachia* status. In one case, however, male w^{1118} flies

survived at higher rates than females for *T. atroviride* infection (Figure S2d), so the pattern of

higher female survival is not always true. Broadly speaking, sex differences in infection

outcomes have long been noted in the literature, and are conserved across diverse host and

pathogen species[86-88]. Some of the results presented here are also in line with observations that

males of many species are often more susceptible to infection than females[89]. Within *Drosophila*,

prior research has shown sex differences in infection are common, can favor either males or

females, and depend on many different factors[90]. Indeed, infectious challenge with a broad

spectrum of bacterial pathogens in *D. melanogaster* demonstrated that females were more

broadly susceptible to infection[91], while another study showed greater female survival with *E.*

coli challenge[92]. Those studies identified specific regulators or sensors in both the IMD and Toll

pathways that are sexually dimorphic in their expression or activation, contributing to differential

immune responses. Sex differences in gut pathology[93], sexual antagonism in immune resistance

and tolerance mechanisms[94], sex chromosome regulation of immune responses[95], and sex

differences in behavior symptoms[96] have all been reported for bacterial or viral infections in

Drosophila. Reports on sex differences in fungal infection have shown mixed results. Notably,

several studies have examined sex-specific outcomes of *B. bassiana* infection in *D.*

melanogaster. One study showed no sex differences in *D. melanogaster* cuticle infection with *B.*

bassiana[97], another showed higher male survival with *B. bassiana* cuticle infection[98], and a third

also showed higher male survival with *B. bassiana* infection introduced either by spray method

or injection[99]. In the third case, removal of various Toll and Imd genes ablated the dimorphism,

indicating their role in the phenotype[99]. Notably, the results herein differed, with females

showing marginally higher survival with *B. bassiana* infection in the w^k line (Figure 2a), and no

sex differences in the w^{1118} line (Figure S2a). This could be due to differences in the host genetic

background strains used in this vs other studies in addition to differences in pathogen infection

method or pathogen strain. Thus, sex differences in infection, favoring males or females, are

common and the result of many different factors. The fact that we observe sex differences in our

results here, but to different extents and in different directions in various contexts, is largely in

line with the literature. Future work will be needed to determine basis of these sex differences.

Sex was not only significant predictor of host outcomes alone, but also in combination

with *Wolbachia* presence or absence. One particularly interesting case was the significant

Wolbachia x sex interaction with *F. oxysporum* infection in the w^{1118} background (Figure S1c).

In this case, only *Wolbachia*-positive males survived significantly longer with fungal infection,

not females. A similar trend was seen in the w^k background, where statistical significance was

evident only when specifically testing within males (Figure 1c). The interaction term of

Wolbachia x sex was not significant, but these sorts of interactions also suffer from low power.

Thus, the mechanism of *Wolbachia* protection from fungal pathogenesis may partially depend on

host factors that differ between the sexes, at least in *F. oxysporum* infection. As for why

Wolbachia may protect males despite transmission mainly through females, it may be due to the

dependency of the symbiont on males to induce reproductive parasitism in this species[100].

Notably, the literature investigating *Wolbachia* blocking of viruses and bacteria in arthropods

often focuses on one specific sex as opposed to both together, particularly for mosquito research,

where viruses are transmitted through female bloodmeals[54,56,101-106]. However, at least one study

reports that female *D. melanogaster* infections with Drosophila C Virus are similar to males[51].

Due to few studies comparing the sexes, it is unclear if there are sexually dimorphic outcomes in

other cases of *Wolbachia* pathogen blocking or what the molecular and genetic bases of putative

Wolbachia x sex interactions may be. However, some possibilities include sex differences in

Wolbachia density, tissue tropism, or dependency on sexually dimorphic host immune responses

to inhibit pathogenesis. Future research will be required to investigate this more fully.

Additionally, there was variation in the size of survival differences between *Wolbachia*-

positive and -negative flies. In some cases, the difference was small but significant, as with *B.*

bassiana (Figures 2a, S2a). In others, the difference was large, such as the *Candida* infections in

the w^k background, (Figures 3a,b). Further, there were differences in longevity based on host

genetic background, with the w^k flies often succumbing to death earlier, or with fewer overall

survivor by the end of the trial period. These results indicate that *Wolbachia*'s impact on fly

survival during fungal infection can have a wide range, from only a slight increase in longevity

to a much larger one, and that host genetics alone (both sex and genetic background) still

significantly influence infection outcomes regardless of *Wolbachia* status. However, even with a

modest increase in longevity of a few days for *B. bassiana*-infected flies with *Wolbachia* as an

example, the fitness benefits in early stages of infection are significant too (Figures 4, S4).

Indeed, the observed increase in early fertility is likely due to reduced pathogen load during

initial infection (Figures 4, S4, 5b, S5b). Notably, the lower fungal titers are not due to

fluctuating *Wolbachia* titers, as they remain the same during infection (Figures 5a, S5a). This

indicates that the symbiont would likely confer a high fitness benefit to a host infected with

fungus in the wild due to the combined effects of laying more eggs per day and living more days.

The potential mechanism of fungal pathogen blocking will be the subject of future study.

From the reduced pathogen load, it is likely to be an immune resistance mechanism as opposed

to tolerance, either of which are known in flies[84,94,107]. In addition, since factors like host sex and

genetic background are significant variables, this suggests that the mechanism is likely at least

partially mediated through the host. Importantly, the *Wolbachia* strains from each background

are nearly genetically identical, with only one single identifiable SNP segregating between the

two strains. Although this does not rule out the possibility of differences due to factors like

different tissue tropism or DNA structural differences not uncovered by Illumina sequencing, it

suggests that differences in phenotypes are likely due to the host rather than symbiont. They do

appear to have similar whole-body titers (Figures 5a, S5a), so overall titer probably does not

explain any differences. However, future research will need to investigate the relative roles of

host and symbiont further. Notably, there is likely to be some overlap in the mechanism(s) of

viral and fungal pathogen blocking in *Drosophila*. First, *w*Mel can block both types of pathogens

based on the results here and shown elsewhere[51,54,72]. Second, some of the molecular

mechanisms contributing to viral blocking could also ostensibly apply to fungal pathogens, such

as immune priming[108], increased ROS production[109], or competition for resources between

symbiont and pathogen[110-112].

Based on the results, we draw several main conclusions: 1) *w*Mel can confer broad, but

not universal, protection against fungal pathogenesis, 2) fungal pathogen blocking by *Wolbachia*

is highly context-dependent, with host sex, genetics, and pathogen species being significant

determinants of host outcomes, and 3) inhibition of fungal pathogenesis can have positive fitness

impacts on the host from early during infection, likely due to reduced pathogen load. Many

questions remain unanswered and future work will be needed to investigate this further. For

example: How broad is the phenotype in terms of symbiont strains, fly species and strains, and

pathogen species? How do other host variables like age impact the phenotype? How do symbiont

density and tissue tropism impact the phenotype? Are the results applicable to other insect

species for potential translational use in agriculture or other fields? What is the mechanism of

fungal pathogen blocking, and can it help inform the mechanism of viral pathogen blocking?

How prevalent is fungal pathogen blocking in the wild? This and prior studies pave the way to

answering these and other important questions.

Materials and Methods:

Fly strains and husbandry

Fly strains include *Drosophila melanogaster w*[1118] (one strain with *Wolbachia*, one cured

of *Wolbachia* via tetracycline) and *D. melanogaster w*[k] (one strain with *Wolbachia*, one cured of

Wolbachia via tetracycline). The w^k line was isolated in Karsnäs, Sweden in 1960 (*white* allele

named for location of isolation)[77] and the w^{1118} line was isolated in California and described in

1985 (*white* allele named for date of isolation)[76]. Both were maintained in various labs since their

isolation. Flies were reared on CMY media: 64.3 g/L cornmeal (Flystuff Genesee Scientific, San

Diego CA), 79.7 mL/L molasses (Flystuff Genesee Scientific), 35.9 g/L yeast (Genesee Scientific

inactive dry yeast nutritional flakes), 8 g/L agar (Flystuff Genesee Scientific *Drosophila* type II

agar), 15.4 mL of antimicrobial mixture [50 mL phosphoric acid (Thermo Fisher, Waltham MA),

418 mL propionic acid (Thermo Fisher), 532 mL deionized water], and 1g/L tegosept (Genesee

Scientific). Flies were kept at 25°C on a 16h light/8 h dark light cycle.

Microbial strains and growth conditions for fly infections

522   The microorganisms used in this study are summarized in Table S2.

523   **Table S2. Microorganisms used in this study.**

Species (strain)	Microbial Classification	Isolation Source	Stock Number or Isolated/Gifted By
Candida glabrata (CBS 138)	Yeast	Feces	ATCC 2001
Candida auris	Yeast	Clinical isolate	CDC B11903
Galactomyces pseudocandidus	Yeast	*Drosophila*	Isolated by I. Nevarez-Saenz
Fusarium oxysporum (f. sp. Lycopersici)	Filamentous fungus	Tomato	FGSC 9935

Beauveria bassiana (GHA)	Filamentous fungus	*Locusta migratoria*	Gift from P. Shahrestani
Aspergillus fumigatus	Filamentous fungus	Clinical isolate	FGSC 1100
Aspergillus flavus (NRRL 3357)	Filamentous fungus	Peanut	FGSC A1446
Metarhizium anisopliae (recently renamed *Metarhizium robertsii*)	Filamentous fungus	Insect	ARSEF 23
Clonostachys rosea	Filamentous fungus	*Aedes albopictus* (mosquito) L4 larvae, Manhattan, KS	Isolated by P. Tawidian & gifted by K. Michel
Trichoderma viride	Filamentous fungus	*Aedes albopictus* (mosquito) L4 larvae, Manhattan, KS	Isolated by P. Tawidian & gifted by K. Michel

Yeast colonies were grown for 16 h on potato dextrose (PD) agar at 30°C. To grow cultures

for fly infections, yeast isolates were grown overnight for 16 h from a single colony in 2 mL PD

broth (BD, Sparks MA) with shaking at 225 rpm. Isolates were then prepared as described below.

Filamentous fungi were prepared by purifying conidia grown on PD agar at 30°C (*Fusarium*,

Aspergillus, and *Beauveria*) or 25°C (*Metarhizium*, *Clonostachys*, and *Trichoderma*) for 1-2 weeks.

Autoclaved DI water was poured over each plate and the conidia were suspended in the liquid.

This was then poured over a filter (Millipore Sigma, Burlington MA, Miracloth 22-25 µm pore

size) and the filtrate was placed into a 50 mL falcon tube. This was then centrifuged at 1000 rpm

for 5 min and the supernatant was discarded. The conidia were then resuspended in sterile 20%

glycerol and were counted using a hemocytometer. The conidia concentrations used in this study

were (conidia/mL): *Aspergillus fumigatus* (1.75×10^9), *Aspergillus flavus* (1.18×10^8), *Fusarium*

oxysporum (9.65×10^7), *Fusarium graminaerum* (1.24×10^8), *Beauveria bassiana* (4.38×10^8),

Metarhizium anisopliae (1.5×10^7), *Clonostachys rosea* (1×10^8), and *Trichoderma atroviride*

(7.2×10^7).

Fly infections

Yeast cultures were grown overnight in the conditions described above. Yeasts *C. glabrata*, *C. auris*, and *G. pseudocandidus* were diluted in PD broth to an optical density (OD) value of $A_{600}= 200$ +/- 5 for *Candida auris* and *Galactomyces pseudocandidus*, and an OD value of $A_{600}= 220$ +/- 5 for *Candida glabrata*. Filamentous fungi were prepared as described above. Mated males or females 4-6 days old of a given genotype were pierced in the thorax just beneath the wing using a 0.15 mm dissecting pin (Entosphinx, Czech Republic, No. 15 Minuten pins 12 mm long 0.15 mm diameter) dipped into the diluted culture or control. Controls were the growth broth for yeasts (PD broth) or sterile 20% glycerol for the filamentous fungi. Flies were then placed in groups of 10 per food vial. 20-30 individuals of each treatment x sex x genotype group were infected in each block, and at least two blocks of infections were performed on separate days for every experiment. Flies were counted for survival daily for 21 days.

Fertility assay

To measure fertility post-infection, 32 virgin 3-5 day old females were collected from each fly strain (w^{1118} and w^{k}, with or without *Wolbachia*). Half of the samples of each strain was infected with *B. bassiana*, as described above. The other half was given 20% glycerol control treatments, also as described above. They were then immediately crossed to 2-4 day old males of the same genotype. Eggs were collected by placing single male-female pairs into a 6 oz. square bottom *Drosophila* bottle (Fisher Scientific, Hampton NH) covered with a grape juice agar plate [100% concord grape juice (Welch's, MA), tegosept (Genesee Scientific, San Diego CA), 200-proof ethanol (Decon Laboratories Inc, PA), agar (Teknova, Hollister CA), DI water] with yeast paste

(Fleischmann's Active Dry Yeast, Heilsbronn Germany, mixed 1:1 volume with water). These

bottles were placed at 25°C incubator overnight. Grape plates were swapped the next morning (16

hr later) with fresh plates and yeast. The bottles were placed back in the incubator and flies were

allowed to lay eggs for 72 h. Plates were then removed and eggs were counted immediately. Plates

were then kept covered for 24 h and egg hatching was recorded.

DNA Extractions

569   DNA extractions were performed with a modified protocol using reagents from the

Qiagen Puregene Cell Core Kit (cat. #158046). Cells from samples were lysed by adding 100 µL

chilled Cell Lysis Solution to each tube, homogenizing the sample with a pestle, incubating at

65°C for 15 min, then cooling on ice. To precipitate protein, 33 µL Protein Precipitation Solution

was added to each sample followed by vortexing for 10 s. Samples were cooled on ice for 5 minutes,

and then centrifuged at 14,000 rpm for 3 min. To precipitate DNA, the supernatant was removed

and mixed with 100 µL pure isopropanol per sample and each sample was inverted 50 times to

mix. The samples were centrifuged 5 min at 14,000 rpm, and supernatant was discarded. Then,

100 µL 70% ethanol was added to each sample and tubes were inverted several times to wash the

DNA pellet. Samples were centrifuged 1 min at 14,000 rpm and supernatant was discarded. Tubes

were inverted over a paper towel for 10 minutes to dry. DNA was then resuspended with 30 µL

DNA Hydration Solution per sample, left at room temperature overnight to allow resuspension,

and then frozen and kept at -20°C the next day until use.

Wolbachia **and fungal titers**

To measure microbial titers post-infection, virgin 3-5 day old females were collected

from each fly strain. Flies were then given the indicated treatment, either *B. bassiana* or 20%

glycerol sham control. They were then collected at 0 and 24 hr post infection. Samples were

flash frozen at their given time point. This led to 10 samples of 3 flies per treatment x time

group. This was done for each of the four fly strains.

qPCR was then performed using the Bio-Rad SsoAdvanced Universal SYBR Green

Supermix (cat. #1725270) according to manufacturer instructions. Primers are listed in Table S3.

qPCR was then performed using a Bio-Rad CFX Connect System with the following conditions:

50°C 10 min, 95°C 5 min, 40x (95°C 10 s, 55°C 30 s), 95°C 30 s. Differences in gene expression

were done by calculating $2^{-\Delta ct}$.

Table S3. Primers used in this study.

Gene	Primer Name	Sequence
Wolbachia groEL	groEL_F	CTAAAGTGCTTAATGCTTCACCTTC
	groEL_R	CAACCTTTACTTCCTATTCTTG
Drosophila rp49	Rp49_F	CGGTTACGGATCGAACAAGC
	Rp49_R	CTTGCGCTTCTTGGAGGAGA
Beauveria bassiana gamma-tubulin	Bbas_F	CAGAGCGACGACACACGC
	Bbas_R	CCCACGCCATTCTTGCCAATG

Drosophila and **Wolbachia** **sequencing and analysis**

For the comparison of the *Wolbachia* from the w^{1118} and w^{k} strains, DNA from 3 female

flies each of each strain with *Wolbachia* was extracted as described above. Samples were prepared

for whole genome sequencing with the xGen™ DNA Library Prep EZ Kit (Integrated DNA

Technologies, #10009821) with a protocol modified to 1/4 reaction volumes. Briefly, 100 ng of

DNA from each sample was buffer exchanged via Ampure XP bead purification (Beckman Coulter

Life Sciences product number A63881) into the low EDTA TE buffer needed for the xGen™ kit,

resulting in a starting input volume of 5 μL. Genomic DNA was enzymatically fragmented to an

expected 350 bp insert size, end repaired, and A-tailed in one reaction step. Stubby Y adapters

were then ligated onto the fragmented DNA, and reactions were bead-purified following adapter

ligation. Unique dual indexes were added to each sample with eight cycles of PCR amplification

of the program provided in the xGen™ DNA Library Prep EZ Kit protocol. The libraries were

then bead-purified twice, first by a 0.6X purification ratio, followed by a 1.2X purification ratio to

provide adapter and primer dimer free libraries. Library quantity was determined with the broad

range dsDNA Qubit Assay on the Qubit 1 Fluorometer (Thermofisher Scientific), and the library

quality and median library size was assessed with a D1000 screen tape on the TapeStation 4150

(Agilent Technologies). Nanomolar concentrations were determined for each library based on their

Qubit concentration in ng/μL and an averaged 442 bp library size. Libraries were pooled at 3 nM

concentration along with another set of libraries for a different project. The libraries were

sequenced at the University of Kansas Medical Center Genome Sequencing Facility on a NovaSeq

6000 S2 150PE flowcell (Illumina Technologies).

Raw reads were trimmed and filtered using fastp[113] with default parameters and removing

the first and last 5 bases from each sequence. Reads were then mapped to a chimeric assembly of

D. melanogaster (Release 6 plus ISO1 MT from NCBI) and *w*Mel *Wolbachia* (ASM1658442v1

from NCBI) using bwa[114] and samtools[115] with default parameters. SNPs were called using

Freebayes[116] with ploidy set to 1 since the host was inbred and *Wolbachia* is haploid, and filtered

with vcffilter[117] with depth greater than 10 and quality greater than 30.

Data visualization and statistical analyses

Data analysis and figure generation were performed in R[118] version 4.2.2, using several

packages: coxme[119] (version 2.2.18.1), ggplot2[120] (version 3.4.0), cowplot[121] (version 1.1.1), car

(version 3.1.1)[122], SurvMiner[123] (version 0.4.9), and SurvMisc[124] (version 0.5.6). Dot plots were

analyzed with a logistic regression. Longevity plots with infection were analyzed using a Cox

proportional hazard model with no *Wolbachia* as the reference.

Data Availability:

633   All data will be deposited in Dryad upon publication of this manuscript.

Acknowledgments:

636   We would like to thank P. Shahrestani and K. Michel for providing certain microbial strains,

637   as well as J. Blumenstiel for providing fly lines. This work was supported by two National

638   Institutes of Health (NIH) K-INBRE P20 GM103418 postdoctoral awards (to JIP), National

639   Science Foundation (NSF) Postdoctoral Fellowship in Biology (PRFB) DBI 2109772 to JIP, NIH

640   K-INBRE P20 GM103418 student award to AA, and NIH grant R01 AI139154 to RLU.

Contributions:

643   JIP and RLU conceived, designed, and analyzed experiments and wrote the manuscript.

644   JIP and AA performed fly experiments. MES performed DNA sequencing. All authors approved

645   of the final version of the manuscript.

References:

648
649
1   Buchner, P. Endosymbiosis of animals with plant microorganisms. (1965).
2   Perlmutter, J. I. & Bordenstein, S. R. Microorganisms in the reproductive tissues of
arthropods. *Nature Reviews Microbiology*, doi:10.1038/s41579-019-0309-z (2020).
3   Douglas, A. E. Nutritional interactions in insect-microbial symbioses: Aphids and
their symbiotic bacteria *Buchnera*. *Annual Review of Entomology* **43**, 17-37,
doi:10.1146/annurev.ento.43.1.17 (1998).
4   Xie, J., Butler, S., Sanchez, G. & Mateos, M. Male killing *Spiroplasma* protects
Drosophila melanogaster against two parasitoid wasps. *Heredity* **112**, 399 (2014).
5   Yen, J. H. & Barr, A. R. New hypothesis of the cause of cytoplasmic incompatibility in
Culex pipiens L. *Nature* **232**, 657-658, doi:10.1038/232657a0 (1971).
6   Kaur, R. *et al.* Living in the endosymbiotic world of *Wolbachia*: A centennial review.
Cell Host & Microbe (2021).
7   Weinert, L. A., Araujo-Jnr, E. V., Ahmed, M. Z. & Welch, J. J. The incidence of bacterial
endosymbionts in terrestrial arthropods. *Proceedings of the Royal Society B:*
Biological Sciences **282**, 20150249 (2015).
8   Zug, R. & Hammerstein, P. Still a host of hosts for *Wolbachia*: analysis of recent data
suggests that 40% of terrestrial arthropod species are infected. *PLoS One* **7**, e38544,
doi:10.1371/journal.pone.0038544 (2012).
9   LePage, D. & Bordenstein, S. R. *Wolbachia*: Can we save lives with a great pandemic?
Trends in Parasitology **29**, 385-393, doi:10.1016/j.pt.2013.06.003 (2013).
10   Werren, J. H., Baldo, L. & Clark, M. E. *Wolbachia*: master manipulators of invertebrate
biology. *Nature Reviews Microbiology* **6**, 741-751, doi:10.1038/nrmicro1969 (2008).
11   Wang, G.-H., Jia, L.-Y., Xiao, J.-H. & Huang, D.-W. Discovery of a new *Wolbachia*
supergroup in cave spider species and the lateral transfer of phage WO among
distant hosts. *Infection, Genetics and Evolution* **41**, 1-7 (2016).
12   Taylor, M., Bordenstein, S. & Slatko, B. Microbe Profile: *Wolbachia*: a sex selector, a
viral protector and a target to treat filarial nematodes. *Microbiology* **164**, 1345-1347
(2018).
13   Slatko, B. *et al.* Pseudoscorpion *Wolbachia* symbionts: Diversity and evidence for a
new supergroup S. (2020).
14   Hurst, G. D. & Frost, C. L. Reproductive parasitism: maternally inherited symbionts
in a biparental world. *Cold Spring Harbor Perspectives in Biology* **7**, a017699 (2015).
15   Dedeine, F., Bouletreau, M. & Vavre, F. *Wolbachia* requirement for oogenesis:
occurrence within the genus *Asobara* (Hymenoptera, Braconidae) and evidence for
intraspecific variation in *A. tabida*. *Heredity* **95**, 394 (2005).
16   Manoj, R. R. S., Latrofa, M. S., Epis, S. & Otranto, D. *Wolbachia*: endosymbiont of
onchocercid nematodes and their vectors. *Parasites & Vectors* **14**, 1-24 (2021).
17   Gong, J.-T., Li, T.-P., Wang, M.-K. & Hong, X.-Y. *Wolbachia*-based strategies for control
of agricultural pests. *Current Opinion in Insect Science*, 101039 (2023).
18   Ant, T. H., Mancini, M. V., McNamara, C. J., Rainey, S. M. & Sinkins, S. P. *Wolbachia*-
virus interactions and arbovirus control through population replacement in
mosquitoes. *Pathogens and Global Health* **117**, 245-258 (2023).

19   Hurst, G. D. D. *et al.* Male–killing *Wolbachia* in two species of insect. *Proceedings of*
the Royal Society of London. Series B: Biological Sciences **266**, 735-740,
doi:10.1098/rspb.1999.0698 (1999).
20   Schilthuizen, M. O. & Stouthamer, R. Horizontal transmission of parthenogenesis–
inducing microbes in *Trichogramma* wasps. *Proceedings of the Royal Society of*
London. Series B: Biological Sciences **264**, 361-366 (1997).
21   Bouchon, D., Rigaud, T. & Juchault, P. Evidence for widespread *Wolbachia* infection
in isopod crustaceans: molecular identification and host feminization. *Proceedings of*
the Royal Society of London. Series B: Biological Sciences **265**, 1081-1090 (1998).
22   Hosokawa, T. *et al.* Obligate bacterial mutualists evolving from environmental
bacteria in natural insect populations. *Nature Microbiology* **1**, 15011 (2016).
23   Taylor, M. J., Bandi, C. & Hoerauf, A. *Wolbachia* bacterial endosymbionts of filarial
nematodes. *Advances in Parasitology* **60**, 245-284, doi:10.1016/s0065-
308x(05)60004-8 (2005).
24   Dedeine, F. *et al.* Removing symbiotic *Wolbachia* bacteria specifically inhibits
oogenesis in a parasitic wasp. *Proceedings of the National Academy of Sciences* **98**,
6247-6252 (2001).
25   Hoffmann, A. A., Clancy, D. & Duncan, J. Naturally-occurring *Wolbachia* infection in
Drosophila simulans that does not cause cytoplasmic incompatibility. *Heredity* **76**, 1-
8 (1996).
26   Hamm, C. A. *et al. Wolbachia* do not live by reproductive manipulation alone:
infection polymorphism in *Drosophila suzukii* and *D. subpulchrella*. *Molecular*
Ecology **23**, 4871-4885 (2014).
27   Meany, M. K. *et al.* Loss of cytoplasmic incompatibility and minimal fecundity effects
explain relatively low *Wolbachia* frequencies in *Drosophila mauritiana*. *Evolution* **73**,
1278-1295 (2019).
28   Reynolds, K. T. & Hoffmann, A. A. Male age, host effects and the weak expression or
non-expression of cytoplasmic incompatibility in *Drosophila* strains infected by
maternally transmitted *Wolbachia*. *Genetics Research* **80**, 79-87 (2002).
29   Yamada, R., Floate, K. D., Riegler, M. & O'Neill, S. L. Male development time influences
the strength of *Wolbachia*-induced cytoplasmic incompatibility expression in
Drosophila melanogaster. *Genetics* **177**, 801-808, doi:10.1534/genetics.106.068486
(2007).
30   Hague, M. T., Mavengere, H., Matute, D. R. & Cooper, B. S. Environmental and genetic
contributions to imperfect *w*Mel-like *Wolbachia* transmission and frequency
variation. *Genetics* **215**, 1117-1132 (2020).
31   Narita, S., Shimajiri, Y. & Nomura, M. Strong cytoplasmic incompatibility and high
vertical transmission rate can explain the high frequencies of *Wolbachia* infection in
Japanese populations of *Colias erate* poliographus (Lepidoptera: Pieridae). *Bulletin*
of Entomological Research **99**, 385-391 (2009).
32   Hoffmann, A. A., Hercus, M. & Dagher, H. Population dynamics of the *Wolbachia*
infection causing cytoplasmic incompatibility in *Drosophila melanogaster*. *Genetics*
148, 221-231 (1998).
33   Dyer, K. A. & Jaenike, J. Evolutionary dynamics of a spatially structured host-parasite
association: *Drosophila innubila* and male-killing *Wolbachia*. *Evolution* **59**, 1518-
1528 (2005).

34   Kittayapong, P., Baimai, V. & O'Neill, S. L. Field prevalence of *Wolbachia* in the
739   mosquito vector *Aedes albopictus*. *American Journal of Tropical Medicine and Hygiene*
740   **66**, 108-111 (2002).

741   35   Jiggins, F. M., Bentley, J. K., Majerus, M. E. & Hurst, G. D. How many species are
742   infected with *Wolbachia*? Cryptic sex ratio distorters revealed to be common by
743   intensive sampling. *Proceedings of the Royal Society of London. Series B: Biological*
744   *Sciences* **268**, 1123-1126 (2001).

745   36   Hurst, G. D., Johnson, A. P., Schulenburg, J. H. & Fuyama, Y. Male-killing *Wolbachia* in
746   *Drosophila*: a temperature-sensitive trait with a threshold bacterial density. *Genetics*
747   **156**, 699-709 (2000).

748   37   Murdock, C. C., Blanford, S., Hughes, G. L., Rasgon, J. L. & Thomas, M. B. Temperature
749   alters *Plasmodium* blocking by *Wolbachia*. *Scientific Reports* **4**, 3932 (2014).

750   38   Bordenstein, S. R. & Bordenstein, S. R. Temperature affects the tripartite interactions
751   between bacteriophage WO, *Wolbachia*, and cytoplasmic incompatibility. *PLoS One*
752   **6**, e29106, doi:10.1371/journal.pone.0029106 (2011).

753   39   Chrostek, E., Martins, N., Marialva, M. S. & Teixeira, L. *Wolbachia*-conferred antiviral
754   protection is determined by developmental temperature. *mBio* **12**, 10.1128/mbio.
755   02923-02920 (2021).

756   40   Osborne, S. E., Iturbe-Ormaetxe, I. a., Brownlie, J. C., O'Neill, S. L. & Johnson, K. N.
757   Antiviral protection and the importance of *Wolbachia* density and tissue tropism in
758   *Drosophila simulans*. *Applied and Environmental Microbiology* **78**, 6922-6929 (2012).

759   41   Ikeda, T., Ishikawa, H. & Sasaki, T. Infection density of *Wolbachia* and level of
760   cytoplasmic incompatibility in the Mediterranean flour moth, *Ephestia kuehniella*.
761   *Journal of Invertebrate Pathology* **84**, 1-5 (2003).

762   42   Hughes, G. L. & Rasgon, J. L. Transinfection: a method to investigate *Wolbachia*-host
763   interactions and control arthropod-borne disease. *Insect Molecular Biology* **23**, 141-
764   151, doi:10.1111/imb.12066 (2014).

765   43   Hoffmann, A. A., Turelli, M. & Harshman, L. G. Factors affecting the distribution of
766   cytoplasmic incompatibility in *Drosophila simulans*. *Genetics* **126**, 933-948 (1990).

767   44   Unckless, R. L., Boelio, L. M., Herren, J. K. & Jaenike, J. Wolbachia as populations
768   within individual insects: causes and consequences of density variation in natural
769   populations. *Proceedings of the Royal Society B: Biological Sciences* **276**, 2805-2811
770   (2009).

771   45   Jiggins, F. M., Randerson, J. P., Hurst, G. D. & Majerus, M. E. How can sex ratio
772   distorters reach extreme prevalences? Male-killing *Wolbachia* are not suppressed
773   and have near-perfect vertical transmission efficiency in *Acraea encedon*. *Evolution*
774   **56**, 2290-2295 (2002).

775   46   Lindsey, A. R. *et al. Wolbachia* is a nutritional symbiont in *Drosophila melanogaster*.
776   *bioRxiv*, 2023.2001. 2020.524972 (2023).

777   47   Hosokawa, T., Koga, R., Kikuchi, Y., Meng, X.-Y. & Fukatsu, T. *Wolbachia* as a
778   bacteriocyte-associated nutritional mutualist. *Proceedings of the National Academy*
779   *of Sciences* **107**, 769-774 (2010).

780   48   Starr, D. J. & Cline, T. W. A host–parasite interaction rescues *Drosophila* oogenesis
781   defects. *Nature* **418**, 76-79 (2002).

49 Ote, M., Ueyama, M. & Yamamoto, D. *Wolbachia* protein TomO targets nanos mRNA
and restores germ stem cells in *Drosophila sex-lethal* mutants. *Current Biology* **26**,
2223-2232 (2016).
50 Russell, S. L., Castillo, J. R. & Sullivan, W. T. *Wolbachia* endosymbionts manipulate
GSC self-renewal and differentiation to enhance host fertility. *bioRxiv*, 2022.2012.
2015.520626 (2022).
51 Teixeira, L., Ferreira, A. & Ashburner, M. The bacterial symbiont *Wolbachia* induces
resistance to RNA viral infections in *Drosophila melanogaster*. *PLoS Biol* **6**, e2,
doi:10.1371/journal.pbio.1000002 (2008).
52 Lipsitch, M., Nowak, M. A., Ebert, D. & May, R. M. The population dynamics of
vertically and horizontally transmitted parasites. *Proceedings of the Royal Society of*
London. Series B: Biological Sciences **260**, 321-327 (1995).
53 Harcombe, W. & Hoffmann, A. *Wolbachia* effects in *Drosophila melanogaster*: in
search of fitness benefits. *Journal of Invertebrate Pathology* **87**, 45-50 (2004).
54 Hedges, L. M., Brownlie, J. C., O'neill, S. L. & Johnson, K. N. *Wolbachia* and virus
protection in insects. *Science (New York, N.Y.)* **322**, 702-702 (2008).
55 Lindsey, A. R., Bhattacharya, T., Newton, I. L. & Hardy, R. W. Conflict in the
intracellular lives of endosymbionts and viruses: a mechanistic look at *Wolbachia*-
mediated pathogen-blocking. *Viruses* **10**, 141 (2018).
56 Dutra, H. L. *et al.* *Wolbachia* blocks currently circulating Zika Virus isolates in
Brazilian *Aedes aegypti* mosquitoes. *Cell Host & Microbe* **19**, 771-774,
doi:10.1016/j.chom.2016.04.021 (2016).
57 Hughes, G. L., Koga, R., Xue, P., Fukatsu, T. & Rasgon, J. L. *Wolbachia* infections are
virulent and inhibit the human malaria parasite *Plasmodium falciparum* in *Anopheles*
gambiae. *PLoS Pathogens* **7**, e1002043 (2011).
58 Hoffmann, A. A. *et al.* Successful establishment of *Wolbachia* in *Aedes* populations to
suppress dengue transmission. *Nature* **476**, 454-457, doi:10.1038/nature10356
(2011).
59 Walker, T. *et al.* The *w*Mel *Wolbachia* strain blocks dengue and invades caged *Aedes*
aegypti populations. *Nature* **476**, 450-453, doi:10.1038/nature10355 (2011).
60 Van den Hurk, A. F. *et al.* Impact of *Wolbachia* on infection with chikungunya and
yellow fever viruses in the mosquito vector *Aedes aegypti*. *PLoS Neglected Tropical*
Diseases **6**, e1892 (2012).
61 Aliota, M. T. *et al.* The *w*Mel strain of *Wolbachia* reduces transmission of
chikungunya virus in *Aedes aegypti*. *PLoS Neglected Tropical Diseases* **10**, e0004677
(2016).
62 MosquitoMate, I. *MosquitoMate: Environmentally friendly innovative mosquito*
control, <https://mosquitomate.com> (2022).
63 Program, W. M. *The World Mosquito Program*,
<https://www.worldmosquitoprogram.org> (2022).
64 O'Neill, S. L. *et al.* Scaled deployment of *Wolbachia* to protect the community from
dengue and other *Aedes* transmitted arboviruses. *Gates Open Research* **2** (2018).
65 World Health Organization. A global brief on vector-borne diseases. (WHO, 2014).
66 O'Neill, S. L. The use of *Wolbachia* by the World Mosquito Program to interrupt
transmission of *Aedes aegypti* transmitted viruses. *Dengue and Zika: Control and*
Antiviral Treatment Strategies, 355-360 (2018).

67   Mains, J. W., Kelly, P. H., Dobson, K. L., Petrie, W. D. & Dobson, S. L. Localized control
of *Aedes aegypti* (Diptera: Culicidae) in Miami, FL, via inundative releases of
Wolbachia-infected male mosquitoes. *Journal of Medical Entomology* (2019).
68   Beebe, N. W. *et al.* Releasing incompatible males drives strong suppression across
populations of wild and *Wolbachia*-carrying *Aedes aegypti* in Australia. *Proceedings*
of the National Academy of Sciences **118**, e2106828118 (2021).
69   Ye, Y. H., Woolfit, M., Rancès, E., O'Neill, S. L. & McGraw, E. A. *Wolbachia*-associated
bacterial protection in the mosquito *Aedes aegypti*. *PLoS Neglected Tropical Diseases*
7, e2362 (2013).
70   Vega, F. E., Meyling, N. V., Luangsa-ard, J. J. & Blackwell, M. Fungal entomopathogens.
Insect Pathology, 171-220 (2012).
71   Fytrou, A., Schofield, P. G., Kraaijeveld, A. R. & Hubbard, S. F. *Wolbachia* infection
suppresses both host defence and parasitoid counter-defence. *Proceedings.*
Biological Sciences **273**, 791-796, doi:10.1098/rspb.2005.3383 (2006).
72   Panteleev, D. *et al.* The endosymbiotic bacterium *Wolbachia* enhances the
nonspecific resistance to insect pathogens and alters behavior of *Drosophila*
melanogaster. *Genetika* **43**, 1277-1280 (2007).
73   Zélé, F., Altıntaş, M., Santos, I., Cakmak, I. & Magalhães, S. Population‐specific effect
of *Wolbachia* on the cost of fungal infection in spider mites. *Ecology and Evolution*
10, 3868-3880 (2020).
74   Ramirez, J. L., Schumacher, M. K., Ower, G., Palmquist, D. E. & Juliano, S. A. Impacts of
fungal entomopathogens on survival and immune responses of *Aedes albopictus* and
Culex pipiens mosquitoes in the context of native *Wolbachia* infections. *PLoS*
Neglected Tropical Diseases **15**, e0009984 (2021).
75   Higashi, C. *et al.* ANOTHER TOOL IN THE TOOLBOX: *Wolbachia*-mediated protection
against a specialized fungal pathogen of aphids. *bioRxiv*, 2023.2007. 2024.550390
(2023).
76   Levis, R., Hazelrigg, T. & Rubin, G. M. Effects of genomic position on the expression of
transduced copies of the white gene of *Drosophila*. *Science (New York, N.Y.)* **229**,
558-561 (1985).
77   Luning, K. Genetics of inbred *Drosophila melanogaster*. *Hereditas* **95**, 181-188
(1981).
78   Law, C. J., Maloney, P. C. & Wang, D.-N. Ins and outs of major facilitator superfamily
antiporters. *Annual Reviews in Microbiology* **62**, 289-305 (2008).
79   Islam, W. *et al.* Insect-fungal-interactions: A detailed review on entomopathogenic
fungi pathogenicity to combat insect pests. *Microbial Pathogenesis* **159**, 105122
(2021).
80   Poveda, J. *Trichoderma* as biocontrol agent against pests: New uses for a
mycoparasite. *Biological Control* **159**, 104634 (2021).
81   Peng, Y. *et al.* Research progress on phytopathogenic fungi and their role as
biocontrol agents. *Frontiers in Microbiology* **12**, 670135 (2021).
82   Sun, Z.-B. *et al.* Biology and applications of *Clonostachys rosea*. *Journal of Applied*
Microbiology **129**, 486-495 (2020).
83   Schlamp, F. *et al.* Dense time-course gene expression profiling of the *Drosophila*
melanogaster innate immune response. *BMC Genomics* **22**, 1-22 (2021).

Duneau, D. *et al.* Stochastic variation in the initial phase of bacterial infection predicts the probability of survival in *D. melanogaster*. *eLife* **6**, e28298 (2017).

Turelli, M. Cytoplasmic incompatibility in populations with overlapping generations. *Evolution* **64**, 232-241 (2010).

Klein, S. L. & Flanagan, K. L. Sex differences in immune responses. *Nature Reviews Immunology* **16**, 626-638 (2016).

Lotter, H. & Altfeld, M. Sex differences in immunity. in *Seminars in Immunopathology.* 133-135 (Springer).

vom Steeg, L. G. & Klein, S. L. SeXX matters in infectious disease pathogenesis. *PLoS Pathogens* **12**, e1005374 (2016).

Zuk, M. The sicker sex. *PLoS Pathogens* **5**, e1000267 (2009).

Belmonte, R. L., Corbally, M.-K., Duneau, D. F. & Regan, J. C. Sexual dimorphisms in innate immunity and responses to infection in *Drosophila melanogaster*. *Frontiers in Immunology* **10**, 3075 (2020).

Duneau, D. F. *et al.* The Toll pathway underlies host sexual dimorphism in resistance to both Gram-negative and Gram-positive bacteria in mated *Drosophila*. *BMC Biology* **15**, 1-17 (2017).

Vincent, C. M. & Dionne, M. S. Disparate regulation of IMD signaling drives sex differences in infection pathology in *Drosophila melanogaster*. *Proceedings of the National Academy of Sciences* **118**, e2026554118 (2021).

Regan, J. C. *et al.* Sex difference in pathology of the ageing gut mediates the greater response of female lifespan to dietary restriction. *eLife* **5**, e10956 (2016).

Vincent, C. M. & Sharp, N. P. Sexual antagonism for resistance and tolerance to infection in *Drosophila melanogaster*. *Proceedings of the Royal Society B: Biological Sciences* **281**, 20140987 (2014).

Kutch, I. C. & Fedorka, K. M. Y-linked variation for autosomal immune gene regulation has the potential to shape sexually dimorphic immunity. *Proceedings of the Royal Society B: Biological Sciences* **282**, 20151301 (2015).

Vale, P. F. & Jardine, M. D. Sex-specific behavioural symptoms of viral gut infection and *Wolbachia* in *Drosophila melanogaster*. *Journal of Insect Physiology* **82**, 28-32 (2015).

Kraaijeveld, A. R., Barker, C. L. & Godfray, H. C. J. Stage-specific sex differences in *Drosophila* immunity to parasites and pathogens. *Evolutionary Ecology* **22**, 217-228 (2008).

Taylor, K. & Kimbrell, D. Host immune response and differential survival of the sexes in *Drosophila*. *Fly* **1**, 197-204 (2007).

Shahrestani, P. *et al.* Sexual dimorphism in *Drosophila melanogaster* survival of *Beauveria bassiana* infection depends on core immune signaling. *Scientific Reports* **8**, 1-9 (2018).

Hoffmann, A. A., Clancy, D. J. & Merton, E. Cytoplasmic incompatibility in Australian populations of *Drosophila melanogaster*. *Genetics* **136**, 993-999 (1994).

Mancini, M. V., Herd, C. S., Ant, T. H., Murdochy, S. M. & Sinkins, S. P. *Wolbachia* strain *w*Au efficiently blocks arbovirus transmission in *Aedes albopictus*. *Plos Neglected Tropical Diseases* **14**, e0007926 (2020).

Moreira, L. A. *et al.* A *Wolbachia* symbiont in *Aedes aegypti* limits infection with dengue, Chikungunya, and Plasmodium. *Cell* **139**, 1268-1278 (2009).

103   Mousson, L. *et al. Wolbachia* modulates Chikungunya replication in *Aedes albopictus.*
Molecular Ecology **19**, 1953-1964 (2010).

104   Cogni, R., Ding, S. D., Pimentel, A. C., Day, J. P. & Jiggins, F. M. *Wolbachia* reduces virus
infection in a natural population of *Drosophila. Communications Biology* **4**, 1327
(2021).

105   Osborne, S. E., Leong, Y. S., O'Neill, S. L. & Johnson, K. N. Variation in antiviral
protection mediated by different *Wolbachia* strains in *Drosophila simulans. PLoS*
Pathogens **5**, e1000656 (2009).

106   Wong, Z. S., Hedges, L. M., Brownlie, J. C. & Johnson, K. N. *Wolbachia*-mediated
antibacterial protection and immune gene regulation in *Drosophila. PloS One* **6**,
e25430 (2011).

107   Chambers, M. C., Jacobson, E., Khalil, S. & Lazzaro, B. P. Consequences of chronic
bacterial infection in *Drosophila melanogaster. PloS One* **14**, e0224440 (2019).

108   Rancès, E., Ye, Y. H., Woolfit, M., McGraw, E. A. & O'Neill, S. L. The relative importance
of innate immune priming in *Wolbachia*-mediated dengue interference. *PLoS*
Pathogens **8**, e1002548 (2012).

109   Pan, X. *et al. Wolbachia* induces reactive oxygen species (ROS)-dependent activation
of the Toll pathway to control dengue virus in the mosquito *Aedes aegypti.*
Proceedings of the National Academy of Sciences **109**, E23-E31 (2012).

110   Caragata, E. P. *et al.* Dietary cholesterol modulates pathogen blocking by *Wolbachia.*
PLoS Pathogens **9**, e1003459 (2013).

111   Caragata, E. P., Rancès, E., O'Neill, S. L. & McGraw, E. A. Competition for amino acids
between *Wolbachia* and the mosquito host, *Aedes aegypti. Microbial Ecology* **67**,
205-218 (2014).

112   Molloy, J. C., Sommer, U., Viant, M. R. & Sinkins, S. P. *Wolbachia* modulates lipid
metabolism in *Aedes albopictus* mosquito cells. *Applied and Environmental*
Microbiology **82**, 3109-3120 (2016).

113   Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics (Oxford, England) **34**, i884-i890 (2018).

114   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler
transform. *Bioinformatics (Oxford, England)* **25**, 1754-1760 (2009).

115   Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, giab008
(2021).

116   Garrison, E. & Marth, G. Haplotype-based variant detection from short-read
sequencing. *arXiv preprint arXiv:1207.3907* (2012).

117   Garrison, E., Kronenberg, Z. N., Dawson, E. T., Pedersen, B. S. & Prins, P. Vcflib and
tools for processing the VCF variant call format. *bioRxiv*, 2021.2005.2021.445151,
doi:10.1101/2021.05.21.445151 (2021).

118   R: A language and environment for statistical computing. (R Foundation for
Statistical Computing, Vienna, Austria, 2020).

119   Therneau, T. Package 'coxme': Mixed Effects Cox Models. (R Foundation for
Statistical Computing, Vienna, Austria, 2022).

120   Villanueva, R. A. M. & Chen, Z. J. ggplot2: elegant graphics for data analysis. (Taylor
& Francis, 2019).

121   Wilke, C. O., Wickham, H. & Wilke, M. C. O. Package 'cowplot'. *Streamlined Plot Theme*
and Plot Annotations for 'ggplot2 (2019).

122   Fox, J. & Weisberg, S. *An R Companion to Applied Regression*. Third edn, (Sage,
966   Thousand Oaks CA, 2019).
967   123   Kassambara, A., Kosinski, M., Biecek, P. & Fabian, S. Package 'survminer'. *Drawing*
968   *Survival Curves using 'ggplot2'(R package version 03 1)* (2017).
969   124   Dardis, C. & Dardis, M. C. Package 'survMisc'.   (2018).
970