Time Tiling of Learned Stimulus-Based Decision Making in Dorsal CA1 in the Hippocampus

Ma, M.[1,8], Simoes de Souza, F.[1,2,8], Futia, G.L.[3], Anderson, S.R.[4], Riguero, J. [4,5], Tollin, D.[4,5], Gentile-Polese, A.[1], Platt, J.P.[6], Hiratani, N.[7], Gibson, E.[3,5] and Restrepo, D[1,5,*].

[1]Department of Cell and Developmental Biology, University of Colorado Anschutz Medical Campus, Aurora, CO, 80045, USA.
[2]Center for Mathematics, Computation and Cognition, Federal University of ABC, Sao Bernardo do Campo, SP, Brazil.
[3]Department of Bioengineering, University of Colorado Anschutz Medical Campus, Aurora, CO, 80045, USA.
[4]Department of Physiology and Biophysics, University of Colorado Anschutz Medical Campus, Aurora, CO, 80045, USA.
[5]Neuroscience Graduate Program, University of Colorado Anschutz Medical Campus, Aurora, CO 80045, USA.
[6]Department of Neurosurgery, University of Colorado Anschutz Medical Campus, Aurora, CO, 80045, USA.
[7]Department of Neuroscience, Washington University, St. Louis, MO 63110, USA

[8]These authors contributed equally to this work
*Correspondence: diego.restrepo@cuanschutz.edu

SUMMARY

Hippocampal neural activity encodes for external cues and their variable temporal

relationships, which is thought to be necessary to construct maps of sequential

experiences. Here we used multiphoton microscopy to image neural activity in

pyramidal cells of dorsal CA1 (dCA1) in mice undergoing go-no go associative learning.

We found that pyramidal cells responded differentially to the rewarded and unrewarded

stimuli. Most pyramidal cells responded to the valence of the stimuli (is the stimulus

rewarded?) and some cells responded to stimulus identity. The stimulus was decoded

accurately from the activity of the neuronal ensemble within each session, and decoding

accuracy increased substantially as the animal learned to differentiate the stimuli.

Decoding the stimulus from individual pyramidal cell activity revealed time-tiling of the

time for discrimination of the stimuli. This time-tiled representation of stimulus

divergence in dCA1 by sequential neural dynamics likely constitutes a map used for

decision-making in associative learning.

INTRODUCTION

The hippocampus provides real time encoding and retrieval of detailed context memories that enables reactions to a constantly changing environment (Wiltgen et al., 2010). This is accomplished by diverse and distributed neuronal activity encoding external features such as contextually referenced space and time and sensory stimuli, as well as features influenced by the animal's behavior such as speed and direction of motion (Buzsáki and Llinás, 2017; Buzsaki and Moser, 2013; Mau et al., 2018; Stefanini et al., 2020). This is a demanding task mediated by plastic circuits involved in associative learning (Gruart et al., 2015; Neves et al., 2008). For example, imagine you are cooking your favorite dish, paella, for a gathering with friends. You carefully set up your paella before your friends arrive relying on your sensory systems and memory, re-reading your recipe, sensing the aroma of the saffron, visually inspecting the arrangement of the dish and checking the temperature. Your friends arrive, and while the paella is cooking you need to rely on periodic reminders to check the dish, walk back to the kitchen where you evaluate the aroma, whether the rice is sticking to the paella pan, and assess when it is ready (a go-no go decision). This behavior takes place within a highly distracting context where you interact with your friends while making the decision whether the dish is ready.

Here we address the neural representation of associative learning in dorsal CA1 (dCA1), an area of the hippocampus involved in spatial learning and working memory tasks. The involvement of dCA1 in associative learning of stimulus discrimination in

head-fixed animals is still unclear (Biane et al., 2023; Li et al., 2017; Moser et al., 1993;

Rangel et al., 2016; Strange et al., 2014; Taxidis et al., 2020). Responsiveness of

dCA1 pyramidal cells to sensory stimuli have been reported to change with learning in

animals undergoing an odorant discrimination associative learning go-no go task (Biane

et al., 2023; Li *et al.*, 2017). Li and co-workers showed that dCA1 pyramidal cells

receiving connections conveying olfactory information from lateral entorhinal cortex

(LEC) develop more selective spiking responses to odor cues during learning for mice

engaged in an odor discrimination go-no go task and showed that optogenetic

inactivation of the LEC to dCA1 connections slows learning in this task (Li *et al.*, 2017).

In addition, Biane and colleagues find a learning-related increase in the proportion of

cells responsive to rewarded (S+) odorant, but not to the unrewarded (S-) odorant,

suggesting that stimulus representations in dCA1 are sensitive to perceived value.

However, learning did not alter decoding accuracy that was already high before training

in dCA1 (Biane *et al.*, 2023). Importantly, temporal patterning of dCA1 neural activity

plays a key role in odorant working memory tasks (MacDonald et al., 2013; Taxidis *et*

al., 2020) and has been postulated to play a role in organizing memories (Eichenbaum,

2017) raising the question whether it contributes to the neural mechanism for go-no go

associative learning in this brain region. Hippocampal "time cells" participate in

conveying memory of the temporal order of events, working memory and signaling

changes in temporal context (Eichenbaum, 2017; Shimbo et al.; Taxidis *et al.*, 2020). It

has been hypothesized that the precise temporally structured activity of neurons make

perceptually related responses coherent in time (Singer and Lazar, 2016), but it

unknown whether there is time-tiled representation of *stimulus divergence* in dCA1 that

could contribute though sequential neural dynamics to behavioral responses reflecting

decision-making in associative learning.

Here we use two-photon calcium imaging to evaluate ensemble decoding of stimuli from

temporally patterned neural activity in dCA1 in mice engaged in an olfactory go-no go

task where they receive a water reward after licking on a spout when presented with the

S+ odorant and do not receive a water reward for the S- odorant (Ma et al., 2020)

(Figure 1A). We characterized stimulus divergence of calcium responses of different

cells (regions of interest, ROIs) from the calcium video by multivariate robust estimation

(Hakan et al., 2021) and we assessed changes in neuronal activity after switching

rewarded with unrewarded stimuli to determine whether the cells respond to the identity

of the stimulus as opposed to its valence (is the stimulus rewarded?). Finally, we

assessed the accuracy of decoding the stimuli with different subsets of cells and

determined whether the onset of stimulus decoding reflecting divergence of stimulus

responses was time-tiled.

RESULTS

Two-Photon Calcium Imaging in Dorsal CA1 in Head-Fixed Mice undergoing a Go-

No Go Olfactory Associative Learning Task

Calcium imaging was performed through a GRIN lens in adult Thy1-GCaMP6f mice

(n=4) expressing the calcium indicator GCaMP6f in pyramidal neurons in dCA1 (Figure

1A, Figure S1). Water-deprived mice started the trial by licking on the water spout. One

of two odorants was delivered in random order 1-1.5 sec after the start of the trial and

the odorant was presented for 4 sec. Mice were trained to respond to the rewarded

odorant (S+) by licking at least once during two 2 sec response areas to obtain a water

reward. The water reward was not delivered regardless of licking when the unrewarded

odorant (S-) was presented. Two-photon calcium imaging movies were denoised (Li et

al., 2023), motion-corrected (Pnevmatikakis and Giovannucci, 2017) and time-binned

traces of nonnegative changes in fluorescence intensity (ΔF/F) were obtained for

multiple ROIs using EXTRACT, an algorithm based on robust estimation theory (Hakan

et al., 2021)(Figure 1B). Figures 1C,E show calcium imaging data from 153 ROIs

recorded in dCA1 during a 20 min session for a mouse that was proficient in the go-no

go task (proficient stage is defined as percent correct >= 80%). Figure S1A shows the

differential lick response to S+ vs. S- in this session. As expected the fluorescence

traces displayed calcium spikes whose timing was heterogeneous when evaluated by

the cross correlation between the traces of all ROIs (Figure 1D). In this session a subset

of the ROIs responded differentially during S+ vs. S- odorant trials (Figures 1C,E).

Consistent with the heterogeneous response there was little difference between S+ and

S- trials when the ΔF/F was averaged over all ROIs (Figure S1C). However, there was

an increase in mean ΔF/F following trial start. These results suggest the existence of a

complex time-structure underlying the calcium responses from a population of dCA1

neurons in the go-no go task.

Mice Learn to Respond Differentially to the Rewarded and Unrewarded Odorants in the Go-No Go Task

In the first training session the percent correct response to the odorants was between 45% and 65% correct for three of four mice (Figure S2 A-C)(S+: 1% heptanal, HEP, S-: mineral oil, MO) Percent correct was calculated as 100*(hits + CR)/number of trials, we name the session as "learning stage" when percent correct behavior is between 45% and 65% correct. We then trained the mice in 3-6 training sessions per day. As shown in Figure S2 A-D the mice gradually reached the proficient stage (percent correct behavior >80%). The fourth mouse became proficient in the first session (Figure S2D). We recorded from a total of 18 learning stage sessions in three mice and a total of 66 proficient sessions in four mice. Once the mice had stable proficient performance the odorants were reversed (S+: mineral oil, MO, S-: 1% HEP) to investigate whether dCA1 calcium responses are encoding the identity or valence of odorants. The percent correct behavior dropped immediately below 50% and gradually recovered to proficient (>80%) (Figure S2A-D). We recorded from a total of 25 forward proficient stage sessions and 41 reversed proficient stage sessions in four mice).

The go-no go task used in this study is an olfactory cued task where the animal starts a trial by licking on the water spout. At the start of the trial a diverter valve turns on directing the 2 lt/min air flow away from the nose cone towards an exhaust line (Figure 1A) and one of two odorant valves (S+ or S-) opens allowing the odorant, delivered at 50 ml/min, to equilibrate with the 2 lt/min background air flow. 1-1.5 sec after trial start

the diverter valve brings the air back to the nose cone resulting in a sharp increase in

odorant concentration providing the mouse with an odorant cue with an onset ~100

msec after reset of the diverter valve (Figure S3A). In the rest of the manuscript the time

when the diverter valve turns back to the odorant cone is time zero. Finally, turning on

the odorant valves at the start of the trial delivers a clicking sound that can be used to

decode the stimuli (see Methods and Figure S3 B-D). However, mice do not cue on the

click as shown by the fact that when the odorant is removed (but the odor valve still

clicks) the performance drops to 50% (Figure S2 E-G, consistent with previous studies

with the same olfactometer (Slotnick and Restrepo, 2005)).

Figure S4 illustrates lick behavior in learning and proficient stages for the four mice in

this study. Figures in S4A show the per trial time course for mean lick fraction for S+

and S- trials calculated for all sessions (18 learning stage and 66 proficient stage, 4

mice). The lick fraction is defined per time point as the fraction of trials within a session

when the lick is detected and ranges from zero (no licks detected) to one (a lick was

detected in every trial). Mice start the trial by licking on the water spout 1 to 1.5 sec

before odorant delivery evidenced by an increase in mean lick fraction before odorant

delivery (arrow in Figure S4A right panel). Mice in the learning stage start licking again

shortly before the odorant is delivered and the mean lick fraction starts diverging

between S+ and S- trials after about one second (Figure S4A, left panel). In order to

quantify when the lick behavior differs between S+ and S- we computed a p value of the

difference in S+ vs S- lick fraction using a ranksum test. The left panel in Figure S4B

shows that for mice in the learning stage the p value for lick divergence decreases

monotonically below 0.05 (black dashed line) when the water reward is delivered (first

red vertical line). In contrast, for the proficient mice (Figure S4B, right panel) the mean

lick fraction diverges between S+ and S- a fraction of a second after odorant delivery.

Figure S4C shows the summary of mean lick fraction for different time windows within

the trial. We find a sharp increase for proficient mice in the difference between S+ and

S- trials in lick fraction during the odor period. We do not find a difference in lick fraction

between S+ and S- trials during the pre time period (-1.5 to 0 sec) when the animal

could use auditory cues. A generalized linear model (GLM) analysis yielded a

statistically significant difference for the interaction between learning vs. proficient, time

window and S+ vs S- for the odor and reinforcement windows (vs. baseline time

window) (p<0.001) but there was no difference for this interaction for the pre time

window (p>0.05), 744 observations, 725 d.f., 4 mice, GLM F-statistic=153, p<0.001

(Table S1). This lick behavior that becomes divergent between S+ and S- trials a

fraction of a second after odorant delivery is consistent with a previous study performed

with the same olfactometer (Ma *et al.*, 2020).

Taken together the data on 50% mouse performance in the absence of odorants (Figure

S2E-G) and lick behavior (Figure S4) indicate that mice cue on odorants in this

multisensory go-no go task.

n-Dimensional ΔF/F Space Expands before the Animal Starts the Trial and the

Dimensionality of Neural Space Increases as the Animal Learns

The highly complex dynamics of dCA1 network provide a high-dimensional space for

the accommodation of an immense repertoire of potential states. Presumably, the n-

dimensional network dynamics collapses into stimulus specific metastable subregions

detectable as manifolds that constrain the dynamic space in anticipation to the

presentation of the stimulus reflected by anticipatory changes in dimensionality (Nieh et

al., 2021; Singer and Lazar, 2016). We proceeded to evaluate changes in n-

dimensional neural space where each dimension is the neural activity for each ROI

quantified by z-scored ΔF/F calcium changes as a function of time within each trial

(zΔF/F, Figure 2A). As a first step we evaluated overall changes in mean zΔF/F

averaged among all ROIs at each time step. Interestingly in this task where the mouse

starts the trial by licking on the lick spout we found a substantial increase in the overall

mean ΔF/F for both S+ and S- *before* the animal started the trial (pre window compared

to baseline increased by ~50%, Figures 2B and C). There appeared to be a difference

in the mean zΔF/F between S+ and S- for proficient mice (Figure 2B). However, there

was no statistical difference between S+ and S- trials in the mean zΔF/F calculated in

different time windows (Figure 2C). A GLM analysis yielded a statistically significant

difference between the time windows (compared to baseline, $p<0.05$) but there was no

difference between S+ and S- or between learning and proficient stages ($p>0.05$), 744

observations, 725 d.f., 4 mice, F-statistic=34.6, $p<0.001$ (Table S2).

We then assessed whether there were changes in the overall volume of n-dimensional

space where each dimension is defined as the axis depicting zΔF/F for each ROI at

each time point evaluated for each trial (Figure 2A). To assess changes in n-

dimensional volume we calculated at each time point the Euclidean distance (Edelman,

1998) between zΔF/F for each S+ and all other S- trials (we name this distance

"between S+ and S-") as opposed to the Euclidean distance between zΔF/F for each

pair of S+ trials or between zΔF/F for each pair of S- trials ("within S+ or S-"). The

increase in mean zΔF/F shown in Figure 2B and C was mirrored by an increase in the

mean Euclidean distance calculated either "between S+ and S-" or "within S+ or S-"

(Figure 2D). This increase in Euclidean distance had an onset slightly before trial start

indicating that there was an expansion of n-dimensional neural space as the animal

prepared to start the trial. This increase started decreasing when the odorant was

presented and subsided by the end of the trial ~15 seconds after trial start. A GLM

analysis of the Euclidean distance calculated for the different time windows (Figure 2E)

yielded a statistically significant difference for Euclidean distance calculated in the

different time windows (compared to baseline, $p < 0.05$), but there was no difference

comparing "between S+ and S-" vs "within S+ or S-" or between learning and proficient

stages ($p > 0.05$), 744 observations, 725 d.f., 4 mice, GLM F-statistic=36.6, $p < 0.001$

(Table S2).

Finally, in order to understand changes in overall n-dimensional space information

content we estimated the dimensionality of the n-dimensional neural space (Litwin-

Kumar et al., 2017). Dimensionality is a convenient measure to quantify representation

of information in a neural layer (in this case CA1) that received mixed sensory and

motor inputs. Its value ranges from the number of ROIs if all display independent

uncorrelated activity to one if all ROIs display correlated activity. Dimensionality is

closely related to classification performance of a decoder. Figures 2F-G show that the

dimensionality is constant within trials and that learning elicits an increase in

dimensionality. A GLM analysis of the per time window dimensionality data shown in

Figure 2G yielded a statistically significant difference for dimensionality for learning

stage vs. proficient stage ($p<0.05$), but there was no difference comparing the pre (-1 to

0 sec) and odor (3.1 to 4.1 sec) time windows ($p>0.05$), 184 observations, 177 d.f., 4

mice, GLM F-statistic=15.4, $p<0.001$ (Table S3). These results indicate that

dimensionality of the initially unconstrained n-dimensional neural space increases as

the mouse's dCA1 network changes with learning.

Demixed Principal Component Analysis (dPCA) Reveals Increased Differential

Response to Stimuli upon Learning

Mean zΔF/F did not differ between S+ and S- trials, and did not change with learning

(Figures 2B-C). This is not surprising given the heterogeneity of CA1 neural activity (e.g.

Figure 1D). In order to perform an assessment of changes in neural activity with

learning and differences in activity between rewarded (S+) and unrewarded (S-)

stimulus trials we proceeded to assess changes in neural activity after dimensionality

reduction. We performed a demixed principal component analysis (dPCA) (Kobak et al.,

2016) of the zΔF/F population data per mouse to evaluate changes in principal

components quantifying neural activity in the axis maximizing stimulus variance. An

example of dPCA for a proficient mouse is shown in Figure S5. dPCA estimates

stimulus components (Figure S5E) and stimulus-independent components (Figure S5F).

Figure S5A and B show the explained variance of the first 10 components. We used the

sum of stimulus components to estimate the divergence of S+ vs. S- trial component

and their changes with learning. Figure S5Di shows that the sum of stimulus

components diverged shortly after trial start. As expected, the sum of independent

components did not differ between S+ and S- trials (Figure S5Dii).

Figure 3 shows the results of dPCA analysis for all mice. The sum of the stimulus

principal components diverged between S+ and S- trials shortly after trial start and the

magnitude of the divergence increased as the animal learned (Figure 3A and B). The

sum of stimulus-independent components did not differ between stimuli and did not

display changes as a function of time (Figure 3C and D). A GLM analysis yielded for the

sum of the stimulus principal components a statistically significant difference for the

interaction between the learning vs. proficient stages and the odor time window (or the

reinforcement time window) vs. the baseline time window and for the interaction

between S+ and S-, the odor time window (or reinforcement window) vs. the baseline

time window and the learning vs. proficient stages (p<0.01), 48 observations, 32 d.f., 4

mice, GLM F-statistic=10.6, p<0.001 (Table S4). The GLM analysis did not yield

statistically significant changes between S+ and S- or for the odor time window for

stimulus-independent components (Table S4).

Stimulus-divergent z∆F/F Responses are Heterogeneous, Divergence Increases

with Learning and the Onset of Divergence Takes Place at Discrete Times

The results of dPCA shown in Figure 3 indicate that there is a substantial increase in

the divergence between S+ and S- as the animal learns. We proceeded to ask whether

this divergence of the responses is heterogeneous when assessed on a per ROI basis.

We classified $z\Delta F/F$ responses in S+ or S- trials within a session when the difference in

$z\Delta F/F$ between a baseline time window (-3 to -1.5 sec) and a response window (-1 to

5.5 sec) was found to be significantly different when tested with a GLM with time and

time window as the two independent variables. The GLM was calculated for each ROI

within all sessions belonging to the learning or proficient stages for the four mice in this

study and the p value for significance was corrected for multiple comparisons using a

false discovery rate (Curran-Everett, 2000). Figure 4A shows the time courses for $z\Delta F/F$

for S+ and S- trials for all divergent ROIs for the proficient stage and Figure 4C shows

the cross correlogram of the $z\Delta F/F$ time courses. We sorted the time courses into three

clusters using a hierarchical binary cluster tree. The average $z\Delta F/F$ time courses for the

three clusters are shown in Figure 4E. Cluster 1 displayed a bimodal time course where

$z\Delta F/F$ for S+ increased above S- shortly after trial start and this reversed after the

reward when S- increased above S+. Cluster 2 displayed an increase in $z\Delta F/F$ for S+

and a decrease for $z\Delta F/F$ for S- shortly after trial start. Cluster 3, that was the cluster

with the largest number of ROIs displayed a robust increase in $z\Delta F/F$ for S- and a

smaller increase for S+. Therefore, time courses for $z\Delta F/F$ for divergent ROIs for the

proficient stage display substantial heterogeneity.

Analysis of $z\Delta F/F$ divergent ROIs for the learning stage also displayed a similar

heterogeneity in time courses (Figure S6B). The percent of divergent ROIs had a

tendency to be smaller for the learning stage, but was not significantly different

compared to the proficient stage (Figure 4D, t-test, p>0.05, 3 mice for learning, 4 mice

for proficient). However, when the zΔF/F was tested with GLM analysis for statistically

significant differences using learning stage and S+ vs. S- as independent variables

(Figure S6D) there were statistically significant differences for learning stage and the

interaction of learning stage and S+ vs. S- for the peak of S- for cluster 1, for the

interaction of learning stage and S+ vs. S- for cluster 2 and for learning stage and the

interaction of learning stage and S+ vs. S- for cluster 3 (p<0.05, number of

observations, d.f. and F statistics are in Table S5). This indicates that there are changes

in zΔF/F time course elicited by learning.

Finally, responses of a subset of CA1 pyramidal cells that have been named odor-

specific time cells take place at discrete time points in the delay period in a delayed

non-match to sample task where the animal receives a reward when they respond in

trial where the second odor differs from the first odor (Taxidis *et al.*, 2020)(also see

(MacDonald *et al.*, 2013). Interestingly, perusal of the pseudocolor time courses for

zΔF/F in Figure 4A appeared to show differences in time onset of the divergence. We

assessed the divergence time for each divergent ROI. Figure S6A shows examples of

the time course for zΔF/F for a subset of ROIs. The divergence takes place at discrete

time points after the start of the trial ranging from onset of divergence close to the start

of the trial (Figure S6A ii) to onset close to the time when the reward was delivered

(Figure S6C vi). Figure 4B shows histograms of the divergence times for the learning

and proficient stages. The divergence times spanned the time period from trial onset to

times after the reward is delivered. The divergence times for the learning stage took

place at earlier times than those for the proficient stage (ranksum p<0.001, 99

divergence time bins in the histograms).

Switching the Rewarded Stimulus to Unrewarded Elicits Changes in zΔF/F

Stimulus Responses Consistent with Response to the Valence of the Stimulus

In order to determine whether stimulus-divergent zΔF/F responses are divergent

responses to the stimulus identity (odorant identity or odorant valve click) vs. stimulus

valence (are the stimuli rewarded?) we switched the rewarded S+ stimulus (1%

heptanal, HEP) with the unrewarded S- stimulus (mineral oil, MO) after the animal

became proficient. We call this a switch from a forward go-no go task when HEP was

S+ to a reversed go-no go task where MO became S+. As shown in Figures S2A-D

after this stimulus reversal the percent correct behavior decreased below 50%, and

subsequently recovered eventually reaching >80% indicating that the animal learned the

new reversed valence of the stimulus. Figure 5A shows the zΔF/F time course for ROIs

that responded with divergence to the stimuli when the animal was proficient in the

forward task. As performed for zΔF/F time courses in Figure 4 we performed a cross

correlation analysis for this forward task (shown in Figure S7C) and we sorted the time

courses into three clusters using a hierarchical binary cluster tree. The average zΔF/F

time courses for cluster 2 that responded with a large increase to S- is shown in Figure

5C and the average time courses for the other clusters are shown in Figure S7A and B.

As in Figure 4 the responses to S- (MO) tended to be larger than responses to S+

(HEP) in the most abundant cluster of responses (cluster 3) in this forward task (Figure

5A).

Interestingly, Figures 5A-D show that when we reversed the stimuli the majority of the

divergent zΔF/F time courses still responded with larger increases for the S- stimulus

(cluster 2, now switched to HEP, also see mean zΔF/F time courses for each cluster in

Figures 5D and S7A,B). Therefore in both the forward and reversed tasks the

unrewarded odorant stimuli elicited a larger zΔF/F response in a large portion of cluster

2 ROIs when the animal was proficient. We quantified this shift in zΔF/F time courses

with reversal by determining the peak value for stimulus-induced changes in zΔF/F.

Figure 5D shows a bar graph of the peak changes in zΔF/F for forward and reversed

runs for cluster 2 and Figure S7Aiii and Biii shows the results for clusters 1 and 3. When

the zΔF/F peak values were tested with GLM analysis for statistically significant

differences using odorant and forward vs. reversed as independent variables there were

statistically significant differences for both odorant and forward vs. reversed ($p < 0.001$,

number of observations, d.f. and F statistics are in Table S6). These data suggest that

the majority of the zΔF/F stimulus responses are responses to the odorant value (is the

odorant rewarded?) as opposed to the odorant identity.

We proceeded to determine for the mouse whose percent correct behavior is shown in

Figure S2D the zΔF/F divergent responses for ROIs whose location was matched

across the last proficient forward vs. the first reversed proficient session. We were able

to match the ROIs because all of these sessions were recorded the same day. Figure

S8A shows divergent zΔF/F responses for matched ROIs in forward and reversed

reward proficient sessions and Figure S8C shows the d' time course calculated for HEP

vs. MO for these divergent zΔF/F responses. d' was calculated on a per time point basis

for zΔF/F time courses for HEP vs. MO. As defined (see methods), a change in d'

polarity indicates that the response is a response to the valence. Figure 5E shows a plot

of the peak value of d' in the odorant period for the forward and reversed runs for all the

matched ROI divergent zΔF/F responses shown in Figure S8C. Most of the d' values

reverse polarity with the forward to reversed switch indicating that these ROIs represent

the odorant value. However, a smaller number kept polarity of the response as the

odorant reward was reversed indicating that a smaller number of ROIs does respond to

the stimulus identity. In conclusion, for proficient sessions (>80% correct behavior) most

of the zΔF/F divergent ROIs represent stimulus valence, and a smaller number of ROIs

respond to the odorant identity.

The Accuracy for Decoding the Stimulus Increases with Learning and is

Dependent on the Fit Window

We asked whether the information embedded within zΔF/F activity of all ROIs can be

used to decode the stimulus (S+ vs. S-). Figure 6 shows the time courses for GLM

decoding of stimulus identity (other decoding algorithms yielded similar results, see

Figure S9). GLM was trained with different time windows using zΔF/F activity of all ROIs

in all S+ and S- trials within each session to evaluate the dependence of decoding

accuracy per trial on the information embedded within particular time windows. As a

control time courses for accuracy are shown where the trial identity (S+ or S-) was

shuffled (red time courses in i and ii panels and light grey bars in iii panels in Figure 6).

When we trained the GLM using a broad fit time window spanning the odorant period

and the beginning of the reward period (0.5 to 5.5 sec) decoding accuracy started

increasing above 0.5 slightly after trial start (~-1 sec) and reached ~0.8 through a

window spanning the later 3 sec of the odorant window for the proficient animal (Figure

6Aii). In contrast, the decoding accuracy for the learning stage only reached ~0.65

(Figure 6Ai). The bar graph in Figure 6Aiii shows the data for pre (-1 to 0), odor (3.1 to

4.1) and reinforcement (4.5 to 5.5) time windows. A GLM analysis yielded a statistically

significant difference for odor and reinforcement time windows vs. pre window and for

the interaction between these time window comparisons and proficient vs. learning

stages (p<0.05-0.001), 279 observations, 270 d.f., 4 mice, GLM F-statistic=37.7,

p<0.001 (Table S7).

We then asked how the decoding accuracy time course varies depending on the fit time

window. As expected when we the fit window spanned a time period before stimulus

presentation (-5 to -3 sec) accuracy did not increase above shuffled (Figure 6C, the p

value for GLM was >0.5 for all comparisons, see Table S7). In contrast, when the

window spanned the reinforcement period (4.5 to 6.5 sec) decoding increased above

0.5 for the learning stage after the mouse was given the reinforcement (Figure 6Bi) and

the increase in decoding accuracy above 0.5 shifted to the time when the odorant was

presented when the mouse became proficient (Figure 6Bii). GLM analysis of the bar

graph in Figure 6Biii yielded a statistically significant difference for odor and

reinforcement time windows vs. pre window and for the interaction between odor

window vs. pre window and proficient vs. learning stages (p<0.05-0.001), but was not

significant for the interaction between reinforcement window vs. pre window and

proficient vs. learning stages 279 observations, 264 d.f., 4 mice, GLM F-statistic=45,

p<0.001 (Table S7). Finally, when the fit window was a narrow window at the end of the

odorant period (2 to 4 sec) the increase in accuracy for the proficient mice did not reach

a plateau during the odorant window (Figure 6D, see GLM results in Table S7). The

results indicate that zΔF/F activity of all ROIs contains information sufficient to decode

the stimuli and that, as expected this information differs when different time windows are

used to fit the GLM decoding algorithm.

Between Trials Stimulus Decoding is Biased to Predict the Unrewarded Stimulus

and there are Sudden Brief Replays of Rewarded Stimulus Prediction

Hippocampal offline replay of neural patterns plays a role in spatial learning (Michon et

al., 2019; Singer et al., 2013). We explored whether replay of neural activity for stimulus

decoding happened between trials for proficient mice engaged in the go-no go task.

Figure 7A shows a time course for prediction of the stimuli by GLM decoding calculated

during an imaging session. As in Figure 6A, GLM was fit to neural activity for all ROIs

during the 0.5 to 5.5 sec spanning most of the odorant window. A prediction value of 1

corresponds to S+ prediction and a value of 0 represents S-. The blue shade shows the

5th to 95th percentile band for prediction calculated after shuffling the stimulus labels. As

expected prediction increases to 1 for a period of time longer than the odorant window

during S+ trials while it increases briefly at the start of S- trials but returns to zero before

the end of the odorant period. Figure 7B shows the mean prediction time course for all

trials in proficient sessions for the 4 mice (a total of 1848 trials). For this within trial

decoding the prediction starts increasing before trial start for both the S+ and S- trials

and it diverges between S+ and S- at the start of the trial. The bar graph in Figure 7C

shows the prediction for S+ and S- in a baseline period (-2.5 to -1.5) and an odor period

(2 to 4.1 sec). A GLM analysis yields significant differences for baseline vs. odor time

windows and for the interaction between S+ vs. S- and time windows, $p<0.001$, 4 mice,

220 observations, 213 d.f., F-statistic 194, $p <0.001$ (Table S8).

Interestingly, the average prediction is lower than shuffled in between trials (Figure 7A

and 7D) indicating that the default decoding is S- for most of the time between trials

(GLM p value for between vs. shuffled <0.001, 110 observations, 105 d.f., F-statistic

70.7, $p<0.001$, Table S8). However, there are clear brief shifts of prediction to 1

(predicting S+, e.g. arrow in Figure 7A). We identified these spontaneous shifts of

prediction from 0 to 1 in between trials by finding increases beyond the 95 percentile of

the shuffled prediction values (blue shade in Figure 7A). We found a total number of

1635 spontaneous shifts to S+ prediction. The duration of these spontaneous shifts in

prediction varied between a fraction of a second to two seconds (example in Figure 7A).

The blue bounded line in Figure 7E shows the mean time course for the between trial

spontaneous shifts in prediction compared to an adjacent time period with no

spontaneous changes in prediction ("no change", magenta). The bar graph in Figure 7F

shows a large difference between the prediction before the onset of the spontaneous

increase in prediction to the peak of prediction for the spontaneous shift. A GLM

analysis yields significant differences for spontaneous vs. no change and for the

interaction between spontaneous vs. no change and baseline vs. peak time windows,

p<0.001, 4 mice, 192 observations, 185 d.f., F-statistic 703, p <0.001 (Table S8).

Finally, we calculated the correlation between zΔF/F time courses for spontaneous shift

and zΔF/F time courses for either S+ or S- trials for all ROIs within each session to

determine whether the neural activity underlying the spontaneous prediction shifts from

S- to S+ are more correlated with S+ vs S-. Figure 7G shows the time courses for the

average correlations. The correlation of spontaneous zΔF/F with S+ diverges from the

correlation with S- shortly after the start of the trial and remains high for a few seconds

after reward. A GLM analysis yields significant differences in average correlations for

comparison of the time windows (baseline vs. odor) and for the interaction between S+

vs. S- and time window, p<0.001, 4 mice, 216 observations, 209 d.f., F-statistic 15.7, p

value <0.001 (Table S8). The sudden shift in prediction from S- to S+ in between trials

and the fact that zΔF/F activity during these spontaneous shifts corelates with zΔF/F

neural activity for S+ suggest that these events are replay of neural activity for the

rewarded stimulus.

Decoding of Decision Making in the Go-No Go Task from CA1 Neural Activity is

Time Tiled

Divergence of z∆F/F responses between S+ and S- stimuli takes place at different times

after trial start (Figure 4C) suggesting that stimulus decoding from neural activity is time

tiled. In order to assess time tiling of the time for decision making (is the stimulus S+ or

S-), we performed an analysis of the time course of stimulus decoding from z∆F/F

responses of *subsets of ROIs* per session for mice proficient in the go-no go task. The

number of ROIs used to calculate GLM decoding per session was varied from 1, 2, 5 to

15 ROIs per session and the results were compared to decoding with all ROIs in each

session. We first performed a visual survey of the within trial dynamics of stimulus

decoding accuracy when the decoding was performed with z∆F/F responses of single

ROIs (Figure S10). We found that decoding accuracy for different ROIs increased at

different times after trial start and reached accuracy values above 0.65. Three examples

of decoding accuracy time courses and their corresponding z∆F/F time courses for S+

and S- trials are shown Figure S10 A-C. In Figure S10 A accuracy increases to ~0.6 at

the start of the trial (immediately following valve click and before odorant addition) and

increases to a higher value (~0.8) after odorant presentation. In Figures S10 B and C

there is no increase in accuracy at the start of the trial and accuracy increases either

shortly after odorant addition (Figure S10 B) or ~3 sec after odorant addition (Figure

S10 C). A histogram of average accuracy values computed during the odor window (3.1

to 4.1 sec) showed a clear difference between S+ and S- trial z∆F/F for these three

examples (Figures S10 A-Ciii). In addition, we found a subset of ROIs whose decoding

yielded decreases in accuracy below 0.5. An example is shown in Figure S10 D where

the accuracy decreased to ~0.1. These decreases in accuracy took place for ROIs with

largely overlapping S+ and S- odorant period z∆F/F values with the exception of one or

two trials where the zΔF/F deviated from the other trials as evidenced in the histogram

in Figure S10 Diii.

We then performed stimulus decoding analysis with subsets of ROIs for the entire

dataset for mice proficient in the go-no go task (four mice, 66 sessions). Figure 8A

shows histograms of decoding accuracy calculated in the odor window (3.1 to 4.1 sec)

for stimulus decoding performed with zΔF/F from subsets of ROIs. As the number of

ROIs per decoding run is decreased stimulus decoding accuracy decreases (compare

Figure 8Av with 8Ai). In addition, accuracies calculated in the pre window (-1 sec to 0

sec) are smaller than accuracies calculated in the odor window (compare Figures 8A

and 8B) and the pre accuracies also appear to decrease as the number of ROIs per

decoding run is decreased (compare Figure 8Bv with 8Bi). A GLM analysis yields

significant differences in decoding accuracy for comparisons between the number of

ROIs and the time windows, $p<0.001$, 4 mice, 27344 observations, 27344 d.f., F-statistic

798, p value <0.001 (Table S9).

In the histograms shown in Figures 8A and B a subset of the accuracy values fall below

0.5 for decoding runs with smaller numbers of ROIs (see for example Figure 8 Ai). As

explained above and shown in Figure S10 Diii for these accuracy decreases the zΔF/F

odorant period values overlap for many of the S+ and S- trials, while they diverge for

one or two trials. However, for the odorant window a large number of decoding accuracy

values remain above 0.65 for decoding with small ROI numbers. For example for

decoding with a single ROI 5.8% out of 6713 ROIs yield odor window accuracy above

0.65 (Figure 8 Ai). We proceeded to quantify the divergence time for each decoding

accuracy time course with odor window decoding accuracy >0.65. Divergence time was

calculated as the time after trial start when accuracy increased above 0.55 for at least

0.2 sec. Figure S11 shows for one session the divergence times estimated using this

method for all decoding accuracy time courses calculated with zΔF/F from single ROIs

that yield odor accuracy above 0.65. Histograms of these divergence times are shown

in Figure 8C for the different multiple ROI decoding sessions. The divergence times

decrease as the number of ROIs used for decoding calculation is increased (Figure 8C).

A GLM analysis of divergence times yields significant differences for comparisons

between accuracies calculated with different numbers of ROIs, $p<0.001$, 4 mice, 27354

observations, 27349 d.f., F-statistic 1400, p value <0.001 (Table S9). Finally, Figure 8D

shows all the decoding accuracy time courses with odorant decoding accuracy >0.65

calculated with a single ROI (Figure 8 Di) or all ROIs (Figure 8 Diii) sorted by

divergence time. For the single ROI decoding there is a large shift in the start of the

increase in accuracy (Figure 8 Di). Computing average accuracy time courses for

different divergence time windows shows that the onset of the increase in decoding

accuracy is time tiled and that decoding accuracy calculated for the earliest divergence

time window is biphasic with a small increase at trial start followed by a larger increase

after odor onset (Figure 8 Dii). These data show that the onset of increases in decoding

accuracy, quantifying time of decision making, is time tiled in hippocampal CA1

suggesting that these are decision making time cells.

Lick Decoding and Stimulus Decoding Differ in their Relationship to Lick

Behavior

Next we asked how stimulus decoding prediction is related to lick behavior (quantified as lick fraction defined per time point as the fraction of trials when the animal was making contact with the lick tube). We expected to find differences between stimulus decoding and lick fraction because the accuracy of stimulus decoding starts increasing above shuffled at trial start, before odorant is presented (Fig. 6A) and, in contrast, licking starts diverging between S+ and S- after the odorant is delivered (Figs. S4A and 9A). In addition, we asked whether it was possible to decode lick behavior from neural activity and whether the relationship between lick prediction and lick fraction differed from the relationship between stimulus prediction and lick fraction. We performed this analysis separately for different trial types classified on behavioral outcome: correct responses (hits and correct rejections, CRs) and incorrect responses (miss and false alarms, FAs)(Figure 1A).

Figure 9A shows the time course for lick fraction for the four types of trials. For all trial types there was a transient increase in lick fraction when the animal started the trial by licking on the water spout (arrow in Figure 9A). For hits (red) this transient increase was followed by a steady increase in lick fraction during the odor period (0-4 sec) while for CR trials (blue) there was a decrease toward zero shortly after odorant presentation. In contrast, for miss trials lick fraction did not increase during the first second of the odorant period (cyan), and for FA trials lick fraction was high for the first half of the

odorant period (magenta). Figure 9B is a bar graph showing the mean lick fraction

during the first and second 2 sec response areas (RAs) during the odorant period (RAs

are defined in Figure 1A). A GLM analysis yields significant differences in mean lick

fraction for comparisons between CR or miss vs. FA or hit and for the interaction of miss

or hit and FA and the RAs, $p<0.001$, 4 mice, 452 observations, 441 d.f., F-statistic 36.7,

p value <0.001 (Table S10).

The accuracy for decoding lick fraction from $z\Delta F/F$ for all ROIs per session when the

mice were proficient (four mice, 66 sessions) is shown in Figure S12. In contrast with

accuracy for stimulus decoding that starts diverging from shuffled at the trial start before

the odorant is presented (Fig. 6A) lick decoding accuracy starts diverging from shuffled

after odorant presentation (Fig. S12A). The bar graph in Figure S12B shows the mean

lick decoding accuracy data for pre (-1 to 0), odor (3.1 to 4.1) and reinforcement (4.5 to

5.5) time windows. A GLM analysis yielded a statistically significant difference for odor

and reinforcement time windows vs. pre window and for the interaction between these

time window comparisons and shuffling ($p<0.001$), 378 observations, 369 d.f., 4 mice,

GLM F-statistic=62.1, $p<0.001$ (Table S10). Figures 9C and D show the lick prediction

time course and mean lick prediction calculated in the RA windows for the different trial

types (hit, miss, CR and FA). A GLM analysis yields significant differences in mean lick

prediction for comparisons between hit or miss and FA and for the interaction of hit and

FA and the RAs, $p<0.001$, 4 mice, 488 observations, 477 d.f., F-statistic 24.3, p value

<0.001 (Table S10).

Figure 9E shows the time course for stimulus prediction for the four types of trials. For

hits (red) there was a steady increase in stimulus prediction that started at trial onset

and leveled to a value of ~0.8 during the odor period for hits (red) while for CR trials

(blue) there was an initial increase after trial start that decreased to ~0.2 during the odor

period. For miss trials stimulus prediction appeared to increase to a value below hit

prediction (cyan), and for FA trials stimulus prediction started increasing at trial start, but

did not decrease to ~0.3 during the odor period (magenta). Figure 9D is a bar graph

showing the mean stimulus prediction during the first and second 2 sec response areas

(RAs) during the odorant period. A GLM analysis yields significant differences in mean

stimulus prediction for comparisons between CR, miss or hit and FA, $p<0.05$, 4 mice,

452 observations, 441 d.f., F-statistic 57.3, p value <0.001 (Table S10).

The time course for lick fraction (Figure 9A) appears to be similar to the time course for

lick prediction (Figure 9C) and appears to differ from stimulus prediction (Figure 9E). In

order to get information on whether there is a common monotonic relationship between

lick fraction and stimulus prediction we plotted the mean lick fraction and mean stimulus

prediction calculated during each second of the odor period. The relationship between

lick fraction and lick prediction was linear (Figure 9G). In contrast, the relationship

between lick fraction and stimulus prediction differed between hit/miss and CR/FA

(Figure 9H). For hits and miss the stimulus/lick fraction relationship was linear (Figure

9E, cyan and red circles) while CR/FA points (Figure 9E, magenta and blue circles) the

relationship appeared to be quadratic and the points did not fall along the hit/miss line.

Therefore, the relationship between lick fraction and stimulus prediction in CA1 does not

follow the simple linear relationship that is expected from neural activity closely related

to motor action. Rather, the relationship differs strikingly for S+ vs. S- stimuli. However,

decoding lick behavior from the same neural activity did yield a prediction that showed a

linear relationship with lick fraction.

DISCUSSION

In this study head-fixed mice engage in an associative learning go-no go task where

they decide whether they lick to obtain a reward when presented with rewarded (S+) or

unrewarded (S-) odorants at concentrations below irritant (trigeminal) threshold

(Schaper, 1993). Unlike other sensory systems, the efferent axons from the olfactory

bulb bypass the thalamus and synapse (directly or through relay in piriform cortex) onto

the LEC that constitutes a major sensory input to dCA1 (Imai, 2014; Witter et al., 2017).

Indeed, before it was realized that the hippocampus plays a key role in learning, Ramón

y Cajal called the hippocampal formation the "quaternary" olfactory system because of

this lack of relay through thalamus (Cajal, 1901). Olfactory responses mediated by LEC

input are particularly prominent in the mouse dCA1 (Igarashi et al., 2014; Manns et al.,

2007; Wood et al., 1999) and odorant responses in dCA1 are known to be involved in

tasks such as delayed match/non match to sample (working memory)(MacDonald *et al.*,

2013; Taxidis *et al.*, 2020), odor place associative learning (Igarashi *et al.*, 2014),

odorant gradient navigation (Radvansky and Dombeck, 2018) and odorant sequence

evaluation (Martinez et al., 2019).

Dorsal CA1 is arguably one of the most studied areas of the brain involved in learning

and memory (Basu and Siegelbaum, 2015; Citri and Malenka, 2008; Fanselow and

Dong, 2010) and it is well documented to play a role in spatial and episodic learning and

memory (Eichenbaum, 2017; Moser *et al.*, 1993). Yet, its role in go-no go associative

learning has been called into question. Li et al. found that calbindin positive pyramidal

neurons in CA1 became more selective for odorants as animals became proficient in

the go-no go task and optogenetic inhibition of these cells slowed learning (Li *et al.*,

2017). In contrast, a study by Biane and co-workers found that the accuracy of odor

decoding in dCA1 was not altered by learning in the go-no go task (Biane *et al.*, 2023).

Here we find that there is a substantial increase in stimulus decoding accuracy when

the animal becomes proficient in the go-no go task (Figure 6); that learning results in

increased stimulus divergence of zΔF/F responses for single cells (Figure 4) and that

learning elicits an increase in the stimulus divergence of stimulus components in

demixed PCA (Figure 3). Our results are consistent with a role for dCA1 pyramidal cells

in go-no go associative learning proposed by Li et al. and with local field potential

recordings in dCA1 that find increased beta power (Martin et al., 2007) and increased

theta-referenced beta and gamma power (Ramirez-Gordillo et al., 2022) upon learning

in dCA1 in the go-no go associative learning task.

The go-no go task in this study also presents the animal with an auditory cue delivered

by opening one of two odorant valves at the start of the trial, 1 to 1.5 sec before the

odorant is delivered (Figures 1A and S3). Using dCA1 neural activity we were able to

decode, albeit with low accuracy, this differential stimulus at the start of the trial (arrow

in Figure 8Dii, Figure S10Ai). The hippocampus is known to respond to auditory cues

(Aronov et al., 2017). Interestingly, even though we find that the information on the odor

valve click is represented by neural activity in dCA1, when we removed the odorants the

behavioral performance of the mice dropped to 50% (Figure S2E-G). Therefore,

although this auditory information is present in CA1 it is not used by these animals as a

stimulus cue in the go-no go task. In addition, for proficient mice lick fraction prediction

in error trials in lick decoding differs from prediction for stimulus decoding (compare

Figures 9D and F, and Figures 9G and H). Indeed, prediction for stimulus decoding

tends to represent the correct stimulus even when the animal makes a mistake (Figure

9F) whereas for lick decoding the predictions for errors align closer with the mistakes

the animal makes in the lick response (compare Figure 9D with Figure 9B). Likely the

animal does perceive the odorant, but is satiated and makes a mistake on purpose. This

indicates that CA1 encodes for different information manifolds that can be used

differentially depending on the context and behavioral purpose.

We asked whether dCA1 pyramidal cells represent the stimulus identity (HEP vs. MO

for odorant identity) or its valence (is this the rewarded stimulus?, S+ vs. S-). We found

that prediction of the stimulus decoded from zΔF/F ensemble neural activity in dCA1

was biased towards the unrewarded stimulus (S-) between trials (Figure 7D). This is

unlike stimulus identity decoding in areas such as auditory cortex where there is no bias

in stimulus prediction (Libby and Buschman, 2021) suggesting that dCA1 ensemble

activity represents stimulus value (S+ vs. S-) as opposed to odorant identity (HEP vs.

MO). Indeed, when the stimuli were reversed the responses were consistent with

representation of valence for a large portion of the ROIs (Figure 5). This is consistent

with valence representation found in olfactory bulb single units after reversal in the go-

no go learning task (Doucette and Restrepo, 2008; Li et al., 2015), functional plasticity

of sensory inputs elicited by learning in the olfactory bulb (Abraham et al., 2014; Chu et

al., 2016), and with decoding of stimulus identity after stimulus reversal for

measurements of theta-referenced beta and gamma LFP power in dCA1 (Ramirez-

Gordillo *et al.*, 2022). Thus, pyramidal cells represent both odorant valence and odorant

identity.

Time-tiled odorant responses in by "time-cells" in dCA1 are thought to be involved in

delayed match/non match to sample where they presumably provide short term working

memory of the identity of the first odorant that is presented in the task (MacDonald *et*

al., 2013; Taxidis *et al.*, 2020). Interestingly, we find that for the go-no go task the

divergence between S+ and S- stimuli in $z\Delta F/F$ odorant responses of individual

pyramidal cells takes place at different times between trial start and reinforcement

(Figure 4) and, consistent with these results, stimulus decoding accuracy decoded from

single neuron activity starts diverging from accuracy computed from shuffled stimulus

labels (50%) at discrete times ranging from trial start to several seconds after

reinforcement (Figures 8, S10 and S11). Therefore, decision making for the difference

between stimuli based on single cell dCA1 neural activity takes place at different

discrete times in dCA1. This finding of what we call "decision-making time-cells" is novel

and complementary to stimulus time-cells engaged in delayed match/non match to

sample tasks (MacDonald *et al.*, 2013; Taxidis *et al.*, 2020). We hypothesize that this

discrete time tiling of information for decision making in dCA1 by sequential neural

dynamics plays a role in go-no go associative learning by representing a memory of

whether the rewarded odorant is presented to the animal that the mouse uses to decide

whether it should sustain licking to get a reward. Whether this is the case will be

determined in future studies.

Finally, our data raise the question on what the specific role is for sequential activity in

associative learning in dCA1. On one hand, sequential activity in dCA1 is often

associated with transition in the state space where each state encodes for a predictive

representation of future states given the current state (Nieh *et al.*, 2021; Stachenfeld et

al., 2017). Under this interpretation, time-tiling represents state transition in a cognitive

map representing the temporal task structure. On the other hand, sequential activity

potentially encodes a single working memory that makes the animal more likely to

choose a certain outcome as time passes. Previous modeling work revealed that

depending on the model configuration, a recurrent neural network learns to represent

working memory either by sequential or persistent activity (Orhan and Ma, 2019). It

indicates that time-tiling might be encoding a binary decision (go or no-go) with spatial-

temporal patterns. Sequential working memory representation is also beneficial for

binding multiple information, such as stimulus identity, valence, and context, into a

single working memory (Feldman, 2013; Hiratani and Sompolinsky, 2023). Future work

will make it clear whether time-tiling in the decision period represents a transition in the

cognitive map or a go/no-go decision variable.

ACKNOWLEDGEMENTS

We would like to thank Brooke Baxter for technical help and Kira Steinke for discussions and feedback on the manuscript. This research was supported by the US National Institutes of Health (NIH UF1 NS116241 and NIH R01 DC000566), and the National Science Foundation (NSF BCS-1926676).

CONTRIBUTIONS

M.M., F.S.de S., E.G. and D.R. conceived and designed the experiments, F.S.de S. performed the surgeries, acquired experimental data and did data analysis, M.M. performed experiments and data analysis, E.G. and G.F. designed and implemented optical instrumentation, G.F. provided technical assistance with multiphoton microscopy, S.A. and D. T. performed sound recordings and analysis, J.R. assisted with immunohistochemistry, A.G.-P. provided technical help, generated illustrations and did immunohistochemistry, J.P. generated code for NWB data sharing, N.H. provided advice on data analysis and writing, M.M., F.S.de S. and D.R. wrote the manuscript and all authors edited the manuscript.

DECLARATION OF INTERESTS

The authors declare no competing interests

STAR * METHODS

Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Donkey anti-rabbit 594 secondary antibody	Jackson Immunoresearch	11-585-152
Rabbit anti GFAP antibody	Abcam	ab7260
EXPERIMENTAL MODELS: ORGANISMS/STRAINS		
Thy1-GCaMP6f mouse, C57BL/6J-Tg(Thy1-GCaMP6f)GP5.17Dkim/J	The Jackson Laboratory	025393
SOFTWARE AND ALGORITHMS		
Demixed PCA	(Kobak *et al.*, 2016)	http://github.com/machenslab/dPCA
Matlab 2022b and 2023a	https://www.mathworks.com/products/matlab.html	N/A
OTHER		
GRIN lenses	GRINTECH GmbH	NEM-050-25-10-860-DS
Metabond	Thermo Fisher	NC0877382, NC2048900, NC2048904
Paraformaldehyde	Electron Microscopy Sciences	15714-S
DAPI	Thermo Fisher	62248
Buprenorphine SR/ER	Zoopharm/Wedgewood	
Vetbond	MWI	006245
Light mineral oil	Fisher	O121-1
Heptanal	Sigma-Aldrich	H-6129
OCT	Sakura Finetech	4583

Animals

All experimental protocols were approved by the Institutional Animal Care and Use

Committee of the University of Colorado Anschutz Medical Campus, in accordance to

NIH guidelines. Mice were bred in the animal facility and we used both males and

females, a total of four adult Thy1-GCaMP6f mouse (Jax 025393 C57BL/6J-Tg(Thy1-

GCaMP6f)GP5.17Dkim/J) for head-fixed awake behaving two photon imaging of CA1

pyramidal cells through a GRIN lens. We used the 5.17 strain of Thy1-GCaMP6f mice

because of their strong expression of GCaMP6f in CA1(Dana et al., 2014). The animals

were housed in a vivarium with a 14/10 h light/dark cycle. Food was available ad libitum.

Access to water was restricted in behavioral training session according to protocols, all

mice were weighed daily and received sufficient water during behavioral sessions to

maintain >80% of original body weight.

Surgical Procedures

Mice were injected with carprofen (5 mg/kg, IP) one hour prior to surgery and were

anesthetized with ketamine/xylazine (IP, 100 mg/kg and 10 mg/kg, respectively).

Anesthetic redosing was assessed by pinching the hind feet every 5 min. A craniotomy

of diameter ~1.2 mm was made over the right hippocampus and a GRIN lens of 4 mm

length and 1 mm diameter (GRINTECH, NEM-050-25-10-860-DS) was implanted at 1.8

mm lateral and 2.4 mm caudal of Bregma and 1.25 mm below dura (Radvansky and

Dombeck, 2018). The GRIN lens edges were sealed with Vetbond glue (3M, USA) and

a custom-made steel head bracket was glued to the skull with Metabond (Parker, USA)

for head-fixing and imaging. The mice were treated with buprenorphine SR/ER for two

days after surgery (0.05 mg/kg, SQ). dCA1 imaging was initiated two weeks after

surgery.

Immunohistochemistry and CT-Scan

To visualize GCaMP6f expression in hippocampus CA1 region and GRIN lens track,

immunohistochemistry was performed after animals finished all the training. Mice were

sacrificed and transcardially perfused with ice cold 4% paraformaldehyde (PFA),

followed by equilibration in 30% sucrose. After the brain sank, the tissue was embedded

in optimal cutting temperature (OCT) compound and were frozen at -80°C. Slices of 60

μm thickness were cut with a Leica cryostat. The slices were stained with rabbit anti-

GFAP primary antibody (Ab7260, Abcam, USA) and Alexa Fluor 594 donkey anti-rabbit

secondary antibody (11-585-152, Jackson Immunoresearch, USA). The slices were

counterstained with DAPI (Thermo Fisher Scientific, USA) and imaged using a confocal

laser scanning microscope (Nikon A1R, Japan). To further verify the 3D position of the

GRIN lens on the skull and in the brain, we performed microCT-Scan recording. We

perfused the mice with PBS and the ex-vivo PFA fixed head was used for acquiring the

microCT imaging on a Siemens Inveon microCT system (Siemens, Germany).

Go-No Go training

Mice were water deprived by restricting daily consumption of water to 1-1.5 ml. Mice were

monitored for signs of dehydration or a decrease in body weight below 80% of the initial

weight. If either condition occurred the animals received water ad-lib until they recovered.

When the animals were water-deprived, they were trained in a head-fixed olfactory go-no

go task with 1% HEP vs MO odorant application (Sigma-Aldrich, USA)(Gire et al., 2013;

Li *et al.*, 2015; Ma *et al.*, 2020). Licks were monitored by an electrical circuit monitoring

the resistance between the lick spout and recorded with Intan RHD2000 software (Intan,

USA). The floor in an olfactometer that controlled valves to deliver a 1:40 dilution of

odorant at a rate of 2 L min^{-1}. Lick rate was calculated from the lick records and the time

course was convolved with a two second Gaussian for the experiments where we

performed multiphoton calcium imaging. The water-deprived mice started the trial by

licking on the water port. The odorant was delivered after a random time interval ranging

from 1 to 1.5 seconds. In S+ trials, the mice needed to lick at least once in two 2 sec lick

segments to obtain a reward (0.1 g ml^{-1} sucrose water) (Figure 1A). In S- trials, the mice

need to refrain licking one of the two 2 sec segments to avoid a longer inter-trial interval

(+ 10 sec). The animal's behavior performance was evaluated in a sliding window of 20

trials and the calculated value was assigned to the last trial in the window. Therefore, it

estimated the performance in the last 20 trials. The percent correct value represents the

percent of trials in which the animal successfully performed appropriate actions

(100*(Hits+CRs)/number of trials), and we considered the animal proficient if percent

correct performance is above 80%. In reverse go-no go training sessions the S+ and S-

odorants were switched after reaching proficient level in forward training.

Acoustic recordings of auditory cue at the start of the trial in the go-no go task

In addition to the odorant cue the olfactometer emits a click at the start of the trial when

one of the two odorant valves opens 1-1.5 sec before odorant onset. Acoustic

recordings of the olfactometer were made using a GRAS 46BH-1 omnidirectional

microphone and GRAS 26AC-1 preamplifier with Type 12AA power module, M-Audio

Pro interface, and Audacity software. The microphone was positioned to approximate

the location of the mouse's head during experiments. Recordings were time-aligned to

experimental hardware using a trigger pulse. Recordings were windowed 7 seconds

prior to 15 seconds following stimulus offset and separated according to S- or S+

conditions in Matlab. Spectrograms of each stimulus were then generated to identify

potential confounding auditory events.

Figures S3B-C show a sound spectrogram of the click recorded with the ultrasound

microphone at the location of the mouse's head in the two-photon microscope aligned

with opening of the odorant valve. Figure S3D shows that the identity of the odorant

valve (S+ vs S-) can be decoded from the sound click using a generalized linear model

(GLM) decoding algorithm, albeit with low accuracy (~ 65%) (ranksum test $p<0.05$, 58

trials for decoding stimuli vs. 290 trials for shuffled stimulus decoding).

While this task provides both olfactory and sound sensory inputs the mouse cues on the

olfactory stimulus as shown by the fact that percent correct behavior decreases close to

50% when the odor cues are removed while the valve opening (and consequent sound)

are not altered (Figure S2E-G, consistent with previous studies with the same

olfactometer (Slotnick and Restrepo, 2005)).

Two photon imaging of dorsal hippocampus CA1 activity in animals undergoing the Go-No Go task

All the animals were first habituated to the setup to minimize stress during the imaging experiments. All the imaging sessions started at least 10 minutes after mice had been head-fixed. The head fixed two photon imaging system consisted of a movable objective microscope (MOM, Sutter Instrument Company, USA) paired with a 80 MHz, ~100 femtosecond laser (Mai-Tai HP DeepSee, Spectra Physics, USA) centered at 920 nm. The MOM was fitted with a single photon epifluorescence eGFP filter path (475 nm excitation/500-550 nm emission) used for initial field targeting followed by switching to the two photon laser scanning path for imaging GCaMP at the depth of the CA1 cell body layer. Either the galvometric laser scanning system or a resonant scanner were driven by SlideBook 6.0 (Intelligent Imaging Innovations, USA). The two photon time lapses were acquired at 395 x 380 pixels using a 0.4 NA/10x air objective (Olympus, Japan) at either 5.3 Hz (galvo) or 20 Hz (resonant). On the day of initial imaging, a FOV was selected to image a large number of active hippocampal CA1 neurons through the GRIN lens, and GCaMP6f movies for several sessions (12-40 trials each) were collected each day. After two photon imaging a second image of the vasculature was captured with wide field epifluorescence to reconfirm the field.

Statistical analysis

Statistical analysis was performed in Matlab R2022b or R2023a (Mathworks, USA). Statistical significance for changes in measured parameters for multiple factors such as

learning and odorant identity (S+ vs. S-) was estimated using a generalized linear model

(GLM)(Yu et al., 2022) with post-hoc comparisons performed either with a two-sided t

test, or a ranksum test, depending on the result of an Anderson-Darling test of normality.

Post-hoc tests were corrected for multiple comparisons using the false discovery rate p

value (pFDR) (Curran-Everett, 2000). Asterisks in bar graphs specify statistically

significant differences (p<pFDR) on post-hoc tests. Not all statistically significant

differences are shown with an asterisks because this makes the figures too complicated.

All pairwise post-hoc test p values and pFDR are shown in the supplementary tables.

Finally, we provide 95% confidence intervals (CIs)(Halsey et al., 2015) estimated by

bootstrap analysis of the mean by sampling with replacement 1000 times using the bootci

function in MATLAB (shown in the figures as vertical black lines for bar graphs or shade

bounding of mean value lines for time courses).

Data analysis

Raw imaging data was first surveyed in ImageJ 1.52 (NIH, USA) to exclude image

sequences exhibiting axial movement. We did not find evidence of axial movement while

the animal was engaged in the go-no go task. The raw imaging data was processed with

DeepCAD-RT to denoise time-lapse images (https://github.com/cabooster/DeepCAD-

RT)(Li *et al.*, 2023), and then motion corrected with NoRMCorre algorithm (Pnevmatikakis

and Giovannucci, 2017). The nonegative fluorescence traces were generated with

EXTRACT (https://github.com/schnitzer-lab/EXTRACT-public)(Hakan *et al.*, 2021) with

appropriate parameters. After Extract analysis, the ΔF/F traces of the spatial components

were sorted and we assigned trial traces to different behavioral events (Hit, CR, Miss and

CR) and aligned them to trial start, odorant onset or reinforcement delivery for further

analysis.

Data analysis was performed using custom code in Matlab R2022b or R2023a

(https://github.com/restrepd/CaImAnDR). Four mice were trained in 3-6 training sessions

per day. Analysis was performed per session classified into either learning stage with

percent correct behavior within the 45-65% range (blue in Figure S2, 20 sessions) or

proficient stage with percent correct behavior in the 80-100% range (red in Figure S2, 55

sessions). For each ROI we z-scored ΔF/F by dividing by the standard deviation of ΔF/F

calculated over the entire session (we call this z scored ΔF/F zΔF/F). Below we describe

the different data analysis methods.

Dimensionality

Following Litwin-Kumar et al. (2017) we defined the dimension of the system (dim) with

M inputs as the square of the sum of the eigenvalues of the covariance matrix of the

measured ΔF/F for all ROIs in the FOV divided by the sum of each eigenvalue squared:

$$\dim\left(\frac{\Delta F}{F}\right) = \left(\sum_{i=1}^{M} \lambda_i\right)^2 \Big/ \left(\sum_{i=1}^{M} \lambda_i^2\right) \qquad (1)$$

where λ_i are the eigenvalues of the covariance matrix of ΔF/F computed over the

distribution of ΔF/F signals measured in the FOV. If the components of ΔF/F are

independent and have the same variance, all the eigenvalues are equal and dim(ΔF/F) =

M. Conversely, if the ΔF/F components are correlated so that the data points are

distributed equally in each dimension of an *m*-dimensional subspace of the full *M*-

dimensional space, only *m* eigenvalues will be nonzero and dim(ΔF/F) = *m*.

Demixed principal component analysis

Demixed principal component analysis (dPCA) is a dimensionality reduction method

decomposing population activity into components related to task parameters such as

stimuli, decisions, or rewards (Kobak *et al.*, 2016). dPCA was calculated for S+ or S-

trials using code provided by Kobak et al. modified to yield stimulus and independent

components using data from all ROIs from all sessions belonging to either the learning or

proficient stages.

Divergent cell detection and analysis

Whether a cell's zΔF/F diverged between S+ and S- (Figures 4 and 5) was determined

as follows: Only sessions that included a minimum of 12 trials were included. ROIs with

calcium spikes in fewer than 25% of the trials were excluded. For the rest of the ROIs the

zΔF/F in the session was considered to be divergent if the GLM p value for S+ vs. S- was

below the pFDR. The time for divergence was calculated as the time after trial onset when

the S+ and S- 95% CIs did not overlap. Examples of the time course zΔF/F for divergent

ROIs are shown in Figure S6A.

As shown in Figures 4A, 5A and B and S8A the per trial time courses for z∆F/F were

heterogeneous. In order to quantify the heterogeneity we calculated within trial cross

correlation coefficients between all divergent z∆F/F time courses (including both S+ and

S- in the calculation) for all divergent ROIs (e.g. Figure 4B). The z∆F/F time courses for

the different ROIs were then separated into different clusters by estimating an

agglomerative hierarchical cluster tree using the linkage function of MATLAB. The

number of clusters was specified arbitrarily.

In order to quantify the divergence in z∆F/F between HEP and MO trials for Figure S8 we

calculated d', which is a measure of the difference in between two distributions.

971
$$d' = \frac{\mathrm{mean}(z\Delta\mathrm{F/F(HEP)}) - \mathrm{mean}(z\Delta\mathrm{F/F(MO)})}{\sqrt{(SD(z\Delta\mathrm{F/F(HEP)})^2 + SD(z\Delta\mathrm{F/F(MO)})^2)/2}} \qquad (2)$$

where SD is the standard deviation. As done for z∆F/F time courses we used cross

correlation coefficients and an agglomerative hierarchical cluster tree to sort the time

courses shown for d' in Figure S8C,D.

Stimulus decoding

Stimulus decoding was performed using GLM (fitglm in MATLAB). The algorithm was

trained per session for all time points and all trials within time periods and number of

ROIs per session indicated in the text. Only sessions with 16 or more trials were

processed. The predicted stimulus was assessed using leave one trial out and winner

takes all procedures. We report the results of GLM decoding, but we obtained similar

results with linear discriminant analysis, artificial neural network decoding, binary

decision tree decoding and support vector machine decoding (Figure S9).

For GLM stimulus decoding performed with a subset of 2, 5 or 15 ROIs (Figure 8)

decoding was performed in separate runs with 40 unique subsets of ROIs drawn at

random from the total number of ROIs per session.

Lick fraction decoding

Lick fraction decoding was performed using a neural network classification model

(fitcnet in MATLAB). The algorithm was trained per session for one second windows

covering from -5 to + 10 sec in each trial. Only sessions with 16 or more trials were

processed. The predicted stimulus was assessed using leave one trial out and winner

takes all procedures.

Analysis of replay of S+ prediction for stimulus decoding between trials

Sudden changes in stimulus prediction from S- to S+ between trials were determined in

decoding prediction moving window averages (with windows of 10 time points) such as

the prediction time course shown in Figure 7A. We searched for a replay (arrow in

Figure 7A) by searching for a sudden shift in prediction from below the 5 percentile of

the shuffled stimulus control for decoding prediction (lower edge of the blue shade in

Figure 7A) to above 95 percentile for the shuffled stimulus decoding prediction (upper

edge of the blue shade in Figure 7A). We compared the average time courses for these

sudden between trial increase to S+ (blue line in Figure 7E) to the time course of

decoding prediction centered in the middle of adjacent between trial intervals where we

did not find a sudden increase to S+ (magenta line in Figure 7E).

1007

1008   **REFERENCES**

1009

Abraham, N.M., Vincis, R., Lagier, S., Rodriguez, I., and Carleton, A. (2014). Long term
functional plasticity of sensory inputs mediated by olfactory learning. Elife *3*, e02109.
10.7554/eLife.02109.

Aronov, D., Nevers, R., and Tank, D.W. (2017). Mapping of a non-spatial dimension by
the hippocampal–entorhinal circuit. Nature *543*, 719-722. 10.1038/nature21692.

Basu, J., and Siegelbaum, S.A. (2015). The corticohippocampal circuit, synaptic
plasticity, and memory. Cold Spring Harbor perspectives in biology *7*, a021733.

Biane, J.S., Ladow, M.A., Stefanini, F., Boddu, S.P., Fan, A., Hassan, S., Dundar, N.,
Apodaca-Montano, D.L., Zhou, L.Z., Fayner, V., et al. (2023). Neural dynamics
underlying associative learning in the dorsal and ventral hippocampus. Nature
Neuroscience *26*, 798-809. 10.1038/s41593-023-01296-6.

Buzsáki, G., and Llinás, R. (2017). Space and time in the brain. Science *358*, 482-485.
10.1126/science.aan8869.

Buzsaki, G., and Moser, E.I. (2013). Memory, navigation and theta rhythm in the
hippocampal-entorhinal system. Nat Neurosci *16*, 130-138. 10.1038/nn.3304.

Cajal, S. (1901). Estudios sobre la corteza cerebral humana. IV. Estructura de la
corteza cerebral olfativa del hombre y mamíferos. Trab Lab Invest Biol 1:1-140.

Chu, M.W., Li, W.L., and Komiyama, T. (2016). Balancing the Robustness and
Efficiency of Odor Representations during Learning. Neuron *92*, 174-186.
10.1016/j.neuron.2016.09.004.

Citri, A., and Malenka, R.C. (2008). Synaptic plasticity: multiple forms, functions, and
mechanisms. Neuropsychopharmacology *33*, 18-41. 10.1038/sj.npp.1301559.

Curran-Everett, D. (2000). Multiple comparisons: philosophies and illustrations.
Am.J.Physiol Regul.Integr.Comp Physiol *279*, R1-R8.

Dana, H., Chen, T.-W., Hu, A., Shields, B.C., Guo, C., Looger, L.L., Kim, D.S., and
Svoboda, K. (2014). Thy1-GCaMP6 Transgenic Mice for Neuronal Population Imaging
In Vivo. PLOS ONE *9*, e108697. 10.1371/journal.pone.0108697.

Doucette, W., and Restrepo, D. (2008). Profound context-dependent plasticity of mitral
cell responses in olfactory bulb. PLoS Biol *6*, e258. 08-PLBI-RA-1962 [pii]
10.1371/journal.pbio.0060258.

Edelman, S. (1998). Representation is representation of similarities. Behavioral and
Brain Sciences *21*, 449-467. 10.1017/S0140525X98001253.

Eichenbaum, H. (2017). On the Integration of Space, Time, and Memory. Neuron *95*,
1007-1018. 10.1016/j.neuron.2017.06.036.

Fanselow, M.S., and Dong, H.-W. (2010). Are the Dorsal and Ventral Hippocampus
Functionally Distinct Structures? Neuron *65*, 7-19. 10.1016/j.neuron.2009.11.031.

Feldman, J. (2013). The neural binding problem(s). Cognitive Neurodynamics *7*, 1-11.
10.1007/s11571-012-9219-8.

Gire, D.H., Whitesell, J.D., Doucette, W., and Restrepo, D. (2013). Information for
decision-making and stimulus identification is multiplexed in sensory cortex. Nat
Neurosci *16*, 991-993. 10.1038/nn.3432.

Gruart, A., Leal-Campanario, R., Lopez-Ramos, J.C., and Delgado-Garcia, J.M. (2015).
Functional basis of associative learning and their relationships with long-term
potentiation evoked in the involved neural circuits: Lessons from studies in behaving
mammals. Neurobiol Learn Mem *124*, 3-18. 10.1016/j.nlm.2015.04.006.

Hakan, I., Claudia, S., Tugce, T., Biafra, O.A., Oscar, H., Jérôme, L., Fatih, D., Mark,
J.W., Murat, A.E., and Mark, J.S. (2021). Fast and statistically robust cell extraction
from large-scale neural calcium imaging datasets. bioRxiv, 2021.2003.2024.436279.
10.1101/2021.03.24.436279.

Halsey, L.G., Curran-Everett, D., Vowler, S.L., and Drummond, G.B. (2015). The fickle
P value generates irreproducible results. Nature methods *12*, 179-185.
10.1038/nmeth.3288.

Hiratani, N., and Sompolinsky, H. (2023). Optimal quadratic binding for relational
reasoning in vector symbolic neural architectures. Neural Computation *35*, 105-155.

Igarashi, K.M., Lu, L., Colgin, L.L., Moser, M.-B., and Moser, E.I. (2014). Coordination of
entorhinal–hippocampal ensemble activity during associative learning. Nature *510*, 143-
147. 10.1038/nature13162.

Imai, T. (2014). Construction of functional neuronal circuitry in the olfactory bulb.
Seminars in Cell & Developmental Biology *35*, 180-188.
https://doi.org/10.1016/j.semcdb.2014.07.012.

Kobak, D., Brendel, W., Constantinidis, C., Feierstein, C.E., Kepecs, A., Mainen, Z.F.,
Qi, X.L., Romo, R., Uchida, N., and Machens, C.K. (2016). Demixed principal
component analysis of neural population data. Elife *5*. 10.7554/eLife.10989.

Li, A., Gire, D.H., and Restrepo, D. (2015). Y spike-field coherence in a population of olfactory bulb neurons differentiates between odors irrespective of associated outcome. J Neurosci *35*, 5808-5822. 10.1523/JNEUROSCI.4003-14.2015.

Li, X., Li, Y., Zhou, Y., Wu, J., Zhao, Z., Fan, J., Deng, F., Wu, Z., Xiao, G., He, J., et al. (2023). Real-time denoising enables high-sensitivity fluorescence time-lapse imaging beyond the shot-noise limit. Nature Biotechnology *41*, 282-292. 10.1038/s41587-022-01450-8.

Li, Y., Xu, J., Liu, Y., Zhu, J., Liu, N., Zeng, W., Huang, N., Rasch, M.J., Jiang, H., Gu, X., et al. (2017). A distinct entorhinal cortex to hippocampal CA1 direct circuit for olfactory associative learning. Nature Neuroscience *20*, 559-570. 10.1038/nn.4517.

Libby, A., and Buschman, T.J. (2021). Rotational dynamics reduce interference between sensory and memory representations. Nature Neuroscience *24*, 715-726. 10.1038/s41593-021-00821-9.

Litwin-Kumar, A., Harris, K.D., Axel, R., Sompolinsky, H., and Abbott, L.F. (2017). Optimal Degrees of Synaptic Connectivity. Neuron *93*, 1153-1164 e1157. 10.1016/j.neuron.2017.01.030.

Ma, M., Futia, G.L., de Souza, F.M.S., Ozbay, B.N., Llano, I., Gibson, E.A., and Restrepo, D. (2020). Molecular layer interneurons in the cerebellum encode for valence in associative learning. Nature Communications *11*, 4217. 10.1038/s41467-020-18034-2.

MacDonald, C.J., Carrow, S., Place, R., and Eichenbaum, H. (2013). Distinct Hippocampal Time Cell Sequences Represent Odor Memories in Immobilized Rats. The Journal of Neuroscience *33*, 14607. 10.1523/JNEUROSCI.1537-13.2013.

Manns, J.R., Zilli, E.A., Ong, K.C., Hasselmo, M.E., and Eichenbaum, H. (2007). Hippocampal CA1 spiking during encoding and retrieval: relation to theta phase. Neurobiol Learn Mem *87*, 9-20. 10.1016/j.nlm.2006.05.007.

Martin, C., Beshel, J., and Kay, L.M. (2007). An olfacto-hippocampal network is dynamically involved in odor-discrimination learning. J.Neurophysiol. *98*, 2196-2205.

Martinez, R.H., Lansner, A., and Herman, P. (2019). Probabilistic associative learning suffices for learning the temporal structure of multiple sequences. PLOS ONE *14*, e0220161. 10.1371/journal.pone.0220161.

Mau, W., Sullivan, D.W., Kinsky, N.R., Hasselmo, M.E., Howard, M.W., and Eichenbaum, H. (2018). The Same Hippocampal CA1 Population Simultaneously Codes Temporal Information over Multiple Timescales. Current Biology *28*, 1499-1508.e1494. https://doi.org/10.1016/j.cub.2018.03.051.

Michon, F., Sun, J.-J., Kim, C.Y., Ciliberti, D., and Kloosterman, F. (2019). Post-learning Hippocampal Replay Selectively Reinforces Spatial Memory for Highly Rewarded

Locations. Current Biology *29*, 1436-1444.e1435.
https://doi.org/10.1016/j.cub.2019.03.048.

Moser, E., Moser, M.B., and Andersen, P. (1993). Spatial learning impairment parallels the magnitude of dorsal hippocampal lesions, but is hardly present following ventral lesions. The Journal of Neuroscience *13*, 3916. 10.1523/JNEUROSCI.13-09-03916.1993.

Neves, G., Cooke, S.F., and Bliss, T.V.P. (2008). Synaptic plasticity, memory and the hippocampus: a neural network approach to causality. Nature Reviews Neuroscience *9*, 65-75. 10.1038/nrn2303.

Nieh, E.H., Schottdorf, M., Freeman, N.W., Low, R.J., Lewallen, S., Koay, S.A., Pinto, L., Gauthier, J.L., Brody, C.D., and Tank, D.W. (2021). Geometry of abstract learned knowledge in the hippocampus. Nature *595*, 80-84. 10.1038/s41586-021-03652-7.

Orhan, A.E., and Ma, W.J. (2019). A diverse range of factors affect the nature of neural representations underlying short-term memory. Nature Neuroscience *22*, 275-283. 10.1038/s41593-018-0314-y.

Pnevmatikakis, E.A., and Giovannucci, A. (2017). NoRMCorre: An online algorithm for piecewise rigid motion correction of calcium imaging data. J Neurosci Methods *291*, 83-94. 10.1016/j.jneumeth.2017.07.031.

Radvansky, B.A., and Dombeck, D.A. (2018). An olfactory virtual reality system for mice. Nat Commun *9*, 839. 10.1038/s41467-018-03262-4.

Ramirez-Gordillo, D., Bayer, K.U., and Restrepo, D. (2022). Hippocampal-Prefrontal θ Coupling Develops as Mice Become Proficient in Associative Odorant Discrimination Learning. eneuro *9*, ENEURO.0259-0222.2022. 10.1523/ENEURO.0259-22.2022.

Rangel, L.M., Rueckemann, J.W., Riviere, P.D., Keefe, K.R., Porter, B.S., Heimbuch, I.S., Budlong, C.H., and Eichenbaum, H. (2016). Rhythmic coordination of hippocampal neurons during associative memory processing. Elife *5*. 10.7554/eLife.09849.

Schaper, M. (1993). Development of a database for sensory irritants and its use in establishing occupational exposure limits. Am.Ind.Hyg.Assoc.J. *54*, 488-544.

Shimbo, A., Izawa, E.-I., and Fujisawa, S. Scalable representation of time in the hippocampus. Science Advances *7*, eabd7013. 10.1126/sciadv.abd7013.

Singer, Annabelle C., Carr, Margaret F., Karlsson, Mattias P., and Frank, Loren M. (2013). Hippocampal SWR Activity Predicts Correct Decisions during the Initial Learning of an Alternation Task. Neuron *77*, 1163-1173.
https://doi.org/10.1016/j.neuron.2013.01.027.

Singer, W., and Lazar, A. (2016). Does the cerebral cortex exploit high-dimensional,
non-linear dynamics for information processing? Frontiers in computational
neuroscience *10*, 99.

Slotnick, B.M., and Restrepo, D. (2005). Olfactometry with mice. In Current Protocols in
Neuroscience, J.N. Crawley, C.R. Gerefen, M.A. Rogawski, D.R. Sibley, P. Skolnick,
and S. Wray, eds. (John Wiley and Sons, Inc), pp. 1-24
1150

Stachenfeld, K.L., Botvinick, M.M., and Gershman, S.J. (2017). The hippocampus as a
predictive map. Nature Neuroscience *20*, 1643-1653. 10.1038/nn.4650.

Stefanini, F., Kushnir, L., Jimenez, J.C., Jennings, J.H., Woods, N.I., Stuber, G.D.,
Kheirbek, M.A., Hen, R., and Fusi, S. (2020). A Distributed Neural Code in the Dentate
Gyrus and in CA1. Neuron *107*, 703-716.e704.
https://doi.org/10.1016/j.neuron.2020.05.022.

Strange, B.A., Witter, M.P., Lein, E.S., and Moser, E.I. (2014). Functional organization
of the hippocampal longitudinal axis. Nature Reviews Neuroscience *15*, 655-669.
10.1038/nrn3785.

Taxidis, J., Pnevmatikakis, E.A., Dorian, C.C., Mylavarapu, A.L., Arora, J.S., Samadian,
K.D., Hoffberg, E.A., and Golshani, P. (2020). Differential Emergence and Stability of
Sensory and Temporal Representations in Context-Specific Hippocampal Sequences.
Neuron *108*, 984-998.e989. https://doi.org/10.1016/j.neuron.2020.08.028.

Wiltgen, B.J., Zhou, M., Cai, Y., Balaji, J., Karlsson, M.G., Parivash, S.N., Li, W., and
Silva, A.J. (2010). The Hippocampus Plays a Selective Role in the Retrieval of Detailed
Contextual Memories. Current Biology *20*, 1336-1344.
https://doi.org/10.1016/j.cub.2010.06.068.

Witter, M.P., Doan, T.P., Jacobsen, B., Nilssen, E.S., and Ohara, S. (2017).
Architecture of the Entorhinal Cortex A Review of Entorhinal Anatomy in Rodents with
Some Comparative Notes. Frontiers in Systems Neuroscience *11*.

Wood, E.R., Dudchenko, P.A., and Eichenbaum, H. (1999). The global record of
memory in hippocampal neuronal activity. Nature *397*, 613-616. 10.1038/17605.

Yu, Z., Guindani, M., Grieco, S.F., Chen, L., Holmes, T.C., and Xu, X. (2022). Beyond t
test and ANOVA: applications of mixed-effects models for more rigorous statistical
analysis in neuroscience research. Neuron *110*, 21-35.
https://doi.org/10.1016/j.neuron.2021.10.030.
1177

1178

1179
1180

Figure 1. Two-photon Ca^{2+} imaging of pyramidal cells in CA1 in head-fixed mice

undergoing the go-no go associative learning task.

(A) Go-no go task. Left: Two-photon imaging of a head-fixed mouse responding to

odorants by licking on a water spout in response to the rewarded odorant in the go-no go

task. Center: Scoring of decision making. Right: time course for the trial. For water reward

in Hit trials the animal must lick at least once in each of the two 2 second lick segments.

(B) Two photon microscopy images of GCaMP6f fluorescence recorded from CA1

pyramidal cells through a GRIN lens in a proficient Thy1-GCaMP6f mouse engaged in

1191   the go-no go task. Left: Activity at rest. Right: Activity during rewarded odorant trial. Video

1192   1 shows fluorescence changes in this group of cells.

1193

1194   (C) ΔF/F traces are shown for a subset of the regions of interest (ROIs) for the figure

1195   shown in B. The magenta vertical lines are the on and off times for the odorant in the

1196   unrewarded (S-) odorant trials and the cyan vertical lines are on and off times for the

1197   rewarded (S+) odorant. The blue lines at the bottom are lick recordings.

1198

1199   (D) Cross correlation and hierarchical clustering of the ΔF/F traces for 153 ROIs for the

1200   entire 20 min session corresponding to B and C shows substantial heterogeneity in the

1201   calcium responses.

1202

1203   (E) Pseudocolor plot of the time course for ΔF/F for the 153 hierarchically clustered ROIs.

1204   Figure S1 shows the time courses for S+ and S- trials for the mean ΔF/F traces, lick

1205   fraction and a p value quantifying the difference between S+ and S- licks.

1206

1207

1208

Figure 2. Changes in n-dimensional z∆F/F space start before the animal starts the

trial and dimensionality increases with learning.

1211

(A) Diagram illustrating the trajectory for the time course of S+ and S- trials in n-

dimensional z∆F/F space and how dimensionality reduction by demixed PCA (dPCA,

1214   Figure 3) reduces the dimensions to stimulus-relevant components. zΔF/F is z-scored

1215 ΔF/F (see methods).

1216

1217   (B) Mean of per trial zΔF/F time course for S+ and S- trials for all learning and proficient

1218   sessions. Left: learning stage, right: proficient stage.

1219

1220   (C) Bar graph showing the mean value of zΔF/F calculated in the following time windows:

1221   Base: -4.5 to -3.5 sec, Pre: -1 to 0 sec, Odor: 3.1 to 4.1 sec, Reinf: 4.5 to 5.5 sec. Dots

1222   are zΔF/F values per session and grey lines are zΔF/F values per mouse.

1223

(D) Mean per trial Euclidean distance calculated in n dimensional zΔF/F ROI space. The

Euclidean distance was calculated: 1) At each time point for each pair of S+ vs. S- trials

and we plot the mean of all of the pair Euclidean distances (we call this "Between"

Euclidean distance). 2) At each time point for each pair of S+ vs. S+ or S- vs. S- trials and

we plot the mean of all of the pair Euclidean distances (we call this "Within" Euclidean

distance).

1230

1231   (E) Bar graph showing the mean value of between and within Euclidean distances

1232   calculated in the following time windows: Base: -4.5 to -3.5 sec, Pre: -1 to 0 sec, Odor:

1233   3.1 to 4.1 sec, Reinf: 4.5 to 5.5 sec. Dots are Euclidean distance values per session and

1234   grey lines are Euclidean distance values per mouse. Hatch bar colors are the same as

1235   the colors for between and within in panel D.

1236

1237   (F) Mean per trial dimensionality calculated in n dimensional zΔF/F ROI space.

1238

1239   (G) Bar graph showing the mean per trial dimensionality calculated in the following time

1240   windows: Base: -4.5 to -3.5 sec, Pre: -1 to 0 sec, Odor: 3.1 to 4.1 sec, Reinf: 4.5 to 5.5

1241   sec. Dots are dimensionality values per session and grey lines are dimensionality values

1242   per mouse.

1243

1244   GLM statistical analysis and post hoc statistics for significance of differences for the bar

1245   graphs are shown in Tables S2 and S3.

1246

1247

1248   For all figures in the manuscript: The bounded lines such as those in B, D and F represent

1249   mean and 95% CI. The light grey horizontal bar is the period between the earliest trial

1250   start and odorant presentation (at 0 sec), the black horizontal bar is the time period for

1251   odorant presentation and the red horizontal bar is the period for water reward delivery.

1252

1253

1254

Figure 3. Demixed PCA reveals a substantial increase elicited by learning in the divergence between S+ and S- stimulus components.

(A) Time courses for the sum of demixed PCA (dPCA) stimulus components for the learning (i, 20 sessions, 4 mice) and proficient (ii, 55 sessions, 4 mice) stages.

(B) Bar graph showing the mean value of the sum of stimulus dPCA components calculated in the following time windows: Base: -4.5 to -3.5 sec, Pre: -1 to 0 sec, Odor: 3.1 to 4.1 sec, Reinf: 4.5 to 5.5 sec.

(C) Time courses for the sum of independent dPCA components for the learning (i, 20 sessions, 4 mice) and proficient (ii, 55 sessions, 4 mice) stages.

(D) Bar graph showing the mean value of the sum of independent dPCA components calculated in the same time windows as in (B).

1270

1271   dPCA was performed following the procedure in (Kobak *et al.*, 2016). An example for all

1272   components and dPCA explained variance for one mouse is shown in Figure S5. GLM

1273   statistical analysis and post hoc statistics for significance of differences for the bar graphs

1274   are shown in Table S4.

1275

1276

Figure 4. The magnitude of divergence between S+ and S- zΔF/F responses increases with learning and the onset of divergence takes place at discrete times that differ between the learning and proficient stages.

(A) zΔF/F time courses for single ROIs that were divergent between S+ and S- trials in the proficient sessions (66 sessions, 4 mice). Time courses were sorted by estimating an agglomerative hierarchical cluster tree shown on the left that was calculated using the

1285 cross correlation coefficients between all divergent zΔF/F time courses shown in C. The

1286 red vertical lines show (in order): trial start, odorant on, odorant off, reinforcement on and

1287 reinforcement off.

1288

1289 (B) Cross correlation coefficients computed between all per trial zΔF/F time courses

1290 shown in A. The coefficients were sorted by the agglomerative hierarchical cluster tree

1291 shown in A.

1292

1293 (C) Histograms for divergence times for all the ROIs for the proficient stage shown in A

1294 (dark blue) and for all divergent ROIs for learning stage sessions (20 sessions, light blue).

1295 Divergence time histograms are significantly different between learning and proficient

1296 (ranksum $p<0.001$, n=99 divergence time bins).

1297

1298 (D) i. Percent divergent ROIs per mouse. ii and iii. Percent responsive ROIs per mouse

1299 for S+ and S- trials. The percent divergent (i) and percent responsive (ii and ii) differences

1300 between learning and proficient stages were not statistically significant (two tailed t test

1301 $p>0.05$, n=3 for learning and 4 for proficient, 5 d.f.).

1302

1303 (E) Mean zΔF/F time courses for the three clusters in the hierarchical tree shown in A.

1304

1305 (F) Peak zΔF/F calculated per session for each cluster. Note that cluster 1 has two peaks

1306 (see arrows in Ei).

1307

1308   Table S5 shows the GLM statistical analysis for Figure 4F.

1309

1310

1311

Figure 5. Reversal of odorant valence reveals that a substantial number of dCA1 cells respond to stimulus valence.

1312

1313

1314

(A and B) zΔF/F time courses for single ROIs that were divergent between S+ and S-

1315

trials for proficient mice. (A) shows zΔF/F time courses for the forward proficient sessions

1316

where the rewarded odorant (S+) was HEP and the unrewarded odorant (S-) was MO (25

1317

sessions, 4 mice) and (B) shows time courses for the reversed proficient sessions where

1318

the rewarded odorant (S+) was MO and the unrewarded odorant (S-) was HEP (41

1319

sessions, 4 mice). For both A and B time courses were sorted by estimating an

1320

agglomerative hierarchical cluster tree shown on the left that was calculated using the

1321

cross correlation coefficients between all divergent zΔF/F time courses shown in Figure

1322

1323 S7 C. The red vertical lines in A and B show (in order): trial start, odorant on, odorant off,

1324 reinforcement on and reinforcement off.

1325

1326 (C) Mean $z\Delta F/F$ time courses for cluster 2 (blue cluster in A and B). Mean $z\Delta F/F$ time

1327 courses for clusters 1 and 3 are shown in Figure S7.

1328

1329 (D) Peak $z\Delta F/F$ calculated per session for cluster 2. Table S6 shows the GLM statistical

1330 analysis for these data.

1331

(E) Relationship for recordings in one mouse between forward and reversed peak d' for

HEP vs. MO for all forward stimulus divergent $z\Delta F/F$ time courses for a set of

forward/reversed sessions where ROIs were matched from session to session. The per

ROI $z\Delta F/F$ and d' time courses for these forward and reversed sessions are shown in

Figure S8. The line shown is d' reversed = d' forward that would be followed if the ROIs

represent stimulus identity.

1338

1339

1340

1341

1342

1343 **Figure 6. Learning elicits an increase in stimulus decoding accuracy.**

1344

(A to D). Each panel shows the accuracy of GLM decoding of the stimulus (S+ vs. S-)

from per trial z∆F/F responses for all trials and all ROIs per session for all learning stage

sessions (i) and proficient sessions (ii) (4 mice, 18 learning sessions, 66 proficient

sessions). The bounded black line represents the mean accuracy bounded by the 95%

CI. The grey lines are per mouse accuracies. The red line is the stimulus decoding

accuracy calculated after shuffling the stimulus labels (S+ vs S-). A to D differ by the time

period used to train the GLM decoder. The A training period (0.5 to 5.5 sec) covers most

of the odorant training period, the B training period takes place before trial onset (-5 to -3

sec), the period for C included the reward delivery period (4.5 to 6.5 sec) and the training

period for D was the latter half of the odorant period (2-4 sec). The bar graphs in iii show

the mean accuracy for different accuracy periods (Pre -1 to 0, Odor 3.1 to 4.1 and Reinf

4.5 to 5.5). Light gray bars are the shuffled stimulus accuracies. Points are per session

accuracies and bars are 95% CIs.

1358

1359   *$p < 0.05$ for a pFDR-corrected t-test or ranksum tests, GLM statistics are in Table S7.

1360

1361

1362　**Figure 7. Replay of prediction of S+ for stimulus decoding between trials.**

1363

(A) Example of the time course for stimulus decoding prediction during a session when

the animal was proficient in the go-no go task. Prediction ranges from 0 (S-) to 1 (S+).

Cyan vertical bars are the odor on and odor off times for the S+ trials and magenta vertical

bars are the odor on and odor off times for the S- trials. The shaded blue area is the 5 to

95 percentile decoding prediction calculated for decoding with shuffled trial labels. The

arrow points to a sudden shift in prediction above shuffled 95 percentile that took place

between trials (we call this a replay).

1371

1372　(B) Mean label prediction time course within trials calculated for S+ and S- trials for all

1373　sessions when the mice were proficient (66 sessions, 4 mice).

(C) Bar graph quantifying mean label prediction for S+ and S- trials in two time windows: Base (-2.5 to -1.5 sec) and odor (2 to 4.1 sec). Dots are per session mean label prediction and grey lines are per mouse mean label prediction.

(D) Bar graph quantifying the mean label prediction between trials compared to the mean label prediction between trials for shuffled label decoding.

(E) Mean label prediction time course for between trial replays ("replay") and for time courses in between trials periods where we did not find spontaneous shifts to S+ ("no change") (66 sessions, 4 mice).

(F) Bar graph quantifying mean label prediction for between trial spontaneous prediction shifts and for no change between trial prediction time courses calculated in two time windows: Base (-2.5 to -1.5 sec) and odor (2 to 4.1 sec). Dots are per session mean label prediction and grey lines are per mouse mean label prediction.

(G) Correlation between label prediction calculated at each time point in the prediction time course for all S+ or S- trials within a session and the label prediction value found at the point of spontaneous shift for spontaneous shifts in prediction found in between trials in the same session (66 sessions, 4 mice).

*p<0.05 for a pFDR-corrected t-test or ranksum tests, GLM statistics are in Table S8.

1396

Figure 8. GLM stimulus decoding accuracy calculated with different subsets of ROIs per session reveals time tiling of the onset of the increase in stimulus decoding accuracy.

(A) Histogram for stimulus decoding accuracy calculated in the odor period (3.1 to 4.1 sec) for all proficient sessions (66 sessions) for 4 mice. i to v show the histogram for an increasing number of ROIs per decoding session. Blue is stimulus decoding accuracy, brown is stimulus decoding accuracy calculated after shuffling the labels. Histograms were normalized by dividing by the total number of counts.

(B) Histograms calculated in the Pre period (-1 to 0 sec).

(C) Histograms for the onset of the increase in decoding accuracy for decoding runs that achieved at least 0.65 accuracy after trial start.

1411

(D) (i). Time courses for decoding accuracies calculated with one ROI per session that

reach at least 0.65 after trial start. (ii) Mean accuracy time courses calculated for one ROI

accuracy time courses shown in Di with accuracy increase onsets in the following time

windows: -1.5 to -1, -1 to 0, 0 to 1, 1 to 2, 2 to 3, >3 sec. (iii) Time courses for decoding

accuracies calculated all ROIs per session. (iv) Mean accuracy time course for the time

courses shown in Diii.

1418

1419   GLM analysis indicates that all histograms in A and B differ from each other and all

1420   histograms in C differ from each other (Table S9).

1421

1422

Figure 9. The relationship between lick fraction and prediction differs between

stimulus decoding and decoding of lick fraction.

(A) Mean lick fraction time course calculated for hits (red), miss (cyan), correct rejections (CR, blue) or false alarm (FA, magenta) trials for all sessions when the mice were proficient (66 sessions, 4 mice).

(B) Bar graph quantifying mean lick fraction for proficient mice (66 sessions, 4 mice) in the two 2 second response areas of the odor period where the mouse must lick at least once to get water in a Hit trial (Figure 1A). Dots are per session lick fraction and grey lines are per mouse averages.

(C) Mean prediction time course for lick fraction decoding within trials calculated for hits (red), miss (cyan), correct rejections (CR, blue) or false alarm (FA, magenta) trials for all sessions when the mice were proficient (66 sessions, 4 mice).

(D) Bar graph quantifying mean lick fraction prediction for proficient mice (66 sessions, 4 mice) in the two 2 second response areas of the odor period where the mouse must lick at least once to get water in a Hit trial (Figure 1A). Dots are per session prediction and grey lines are per mouse averages.

(E) Mean stimulus prediction time course within trials calculated for hits (red), miss (cyan), correct rejections (CR, blue) or false alarm (FA, magenta) trials for all sessions when the mice were proficient (66 sessions, 4 mice).

1448   (F) Bar graph quantifying mean stimulus prediction for proficient mice (66 sessions, 4

1449   mice) in the two 2 second response areas of the odor period where the mouse must lick

1450   at least once to get water in a Hit trial (Figure 1A). Dots are per session prediction and

1451   grey lines are per mouse averages. The training period for the GLM decoding algorithm

1452   was 0.5 to 5.5 sec.

1453

1454   (G) Relationship between mean lick fraction and mean lick fraction prediction (calculated

1455   from per mouse values). The bars are 95% CIs calculated per mouse. Lick fraction and

1456   lick fraction prediction were calculated in two second time windows spanning from odor

1457   onset to 6 sec after odor onset. The line is a linear fit.

1458

1459   (H) Relationship between mean lick fraction and mean stimulus prediction (calculated

1460   from per mouse values). The bars are 95% CIs calculated per mouse. Lick fraction and

1461   stimulus prediction were calculated in two second time windows spanning from odor onset

1462   to 6 sec after odor onset. The data for Hit and Miss is fit by a linear fit and the data for CR

1463   and FA is fit by a second order polynomial.

1464

1465   For all graphs the data are shown separately for S+ Hits (red) and Miss (cyan) and S- CR

1466   (blue) and FA (magenta) trials. *p<0.05 for a pFDR-corrected t-test or ranksum tests,

1467   GLM statistics are in Table S10.

1468

1469

1470

Figure S1. Supplementary data for Figure 1.

(A) Mean of lick fraction time course for S+ and S- trials for the session for the data shown

in Figure 1.

1471

1472

1473

1474

1475

1476
1477

1478   (B) p value for a ranksum test comparing S+ vs. S- per trial lick time courses for the

1479   session for the data shown in Figure 1. The horizontal red line is p=0.05.

1480

1481   (C) Mean of per trial zΔF/F time course for S+ and S- trials for the session for the data

1482   shown in Figure 1.

1483

1484   (D) Immunofluorescence and CT scan showing the location of the GRIN lens in one

1485   mouse.

1486

1487

Figure S2. Behavioral performance of mice in the go-no go task.

1488

1489

(A to D) show the percent correct behavior calculated in a moving window of 20 trials for

the entire training period for the four mice used in this study. Percent correct behavior is

calculated as 100*(Hit trials+CR trials)/20. The trials with performance between 45 and

65 % correct (shown in blue) were classified as learning stage trials while the trials with

performance at or above 80 % correct were classified as proficient (red). The task was

started with heptanal (HEP) as S+ and mineral oil (MO) as S- (forward sessions). The

vertical black line shows the point where these stimuli were reversed (MO became S+

and HEP became S-, reversed sessions).

1498

1499   (E, F and G) show for three mice that the behavioral performance drops to 50% when the

1500   odorants are not delivered, but the odorant delivery valves still produce a click sound at

1501   the start of the trial (vertical line, no odor).

1502

1503

1504

Figure S3. Time course for odorant onset and auditory cues.

1505

1506

1507 (A) Increase in odorant concentration measured with a photoionization detector (PID).

1508 The inset shows that the onset of the increase in odorant concentration takes place ~100

1509 msec after the diverter valve is turned off to deliver the odorant to the nose cone (time

1510 zero).

1511

(B) Spectrogram time course for the mean sound power (in dB) for the clicks produced

by the olfactometer for S+ and S- trials measured at the location where the mouse's head

is in the 2P imaging enclosure. Time is referenced to the time point where the diverter

and odor valves are turned on simultaneously (the start of the trial). Sound recording was

performed in 30 S+ and 28 S- trials. The average power before diverter valve activation

was subtracted from the power at each time point.

1518

1519 (C) Spectrogram for the mean sound power (in dB) measured at the time bin when the

1520 click occurred for the clicks produced by the olfactometer for S+ and S- trials measured

1521 at the location where the mouse's head is in the 2P imaging enclosure.

1522

1523 (E) Accuracy for decoding of trial type (S+ or S-) from sound power measured at the click.

1524 Decoding of trial type from sound power for clicks recorded in 30 S+ and 28 S- trials was

1525 performed using GLM with a leave one out procedure. As a control the labels (S+ or S-)

1526 were shuffled. A ranksum test yields a p value of 0.01 for the difference in per trial label

1527 prediction between shuffled and original (no shuffling) (n=58 trials).

1528

1529

1530

1531
1532
1533 **Figure S4. Lick responses for learning and proficient stages in the go-go task.**
1534
1535

1536 (A) Time course for lick fraction mean $\pm$ 95% CI for S+ and S- trials for all learning and

1537 proficient sessions (4 mice). Left: learning stage, right: proficient stage. Arrow points to

1538 the increase in lick fraction that takes place at the start of the trial.

1539

(B) p value (mean $\pm$ 95% CI) calculated by a per time point ranksum test of the difference

in lick fraction between S+ and S- trials (evaluated per session). The p value was

calculated for all learning and proficient sessions (4 mice). The horizontal dashed line is

p=0.05. The arrow points to the time when the p value decreases monotonically below

0.05 for proficient mice.

(C) Bar graph showing the mean value of lick fraction calculated in the following time

windows: Base: -4.5 to -3.5 sec, Pre: -1 to 0 sec, Odor: 3.1 to 4.1 sec, Reinf: 4.5 to 5.5

sec. Dots are lick fraction values per session and grey lines are lick fraction values per

mouse.

1549

1550   *$p<0.05$ for a pFDR-corrected t-test or ranksum tests, GLM statistics are in Table S1.

1551

1552

1553

1554
1555 **Figure S5. Example of demixed PCA for one mouse.**

1556

1557 (A) Cumulative variance explained by PCA (black) and dPCA (red). Demixed PCA

1558 explains almost the same amount of variance as the standard PCA.

1559

1560 (B) Variance of the individual demixed principal components. Each bar shows the

1561 proportion of total variance, and is composed out of two stacked bars of different color:

1562 blue for

stimulus variance and red for stimulus independent variance. Pie chart shows how the

total signal variance is split among parameters.

(C) Upper-right triangle shows dot products between all pairs of the first 15 demixed

principal axes. Stars mark the pairs that are significantly and robustly non-orthogonal (see

Materials and methods). Bottom-left triangle shows correlations between all pairs of the

first 15 demixed principal components.

(D) Time course for the sum of all stimulus (i) or independent (ii) dPCA components

(E) Time course for the three top stimulus dPCA components

(F) Time course for the three top independent dPCA components

1577

Figure S6. Stimulus-divergent zΔF/F responses for the learning stage and examples of zΔF/F responses that diverge between S+ and S- at different discrete time points after the start of the trial.

1581

(A) Examples of zΔF/F time courses for single ROIs that differ in onset time for divergence between S+ and S- trials

1584

(B) zΔF/F time courses for single ROIs that were divergent between S+ and S- trials in the learning sessions (18 sessions, 4 mice). Time courses were sorted by estimating an agglomerative hierarchical cluster tree shown on the left that was calculated using the cross correlation coefficients between all divergent zΔF/F time courses (not shown). The

1589   red vertical lines show (in order): trial start, odorant on, odorant off, reinforcement on and

1590   reinforcement off.

1591

1592   (C) Mean $z\Delta F/F$ time courses for the three clusters in the hierarchical tree shown in B.

1593

1594

1595

1596

1597

1598

Figure S7. Supplemental data for Figures 5A and B characterizing changes in stimulus divergent z∆F/F responses after reversal of odorant valence.

(A and B) Mean z∆F/F time courses (i,ii) and peak z∆F/F (iii) for clusters 1 and 3 of the hierarchical cluster trees shown in Figures 5 A and B. Mean z∆F/F time courses for cluster 2 is Figure 5C. Table S6 shows the GLM statistical analysis for iii.

1606 (C) Cross correlation coefficients computed between all per trial zΔF/F time courses

1607 shown in Figures 5A and B. The coefficients were sorted by the agglomerative

1608 hierarchical cluster tree shown in those figures.

1609

1610 (D) i. Percent divergent ROIs per mouse for the data in Figures 5A and B. ii and iii. Percent

1611 responsive ROIs per mouse for S+ and S- trials. The percent divergent (i) and percent

1612 responsive (ii and ii) differences between learning and proficient stages were not

1613 statistically significant (two tailed t test p>0.05, n=3 for learning and 4 for proficient, 5 d.f.).

1614

1615

1618 **Figure S8. Supplemental figure for Figure 5D characterizing zΔF/F for reversal**

1619 **calculated for one mouse with matched ROIs between forward and reversed**

1620 **sessions.**

1621

(A) zΔF/F time courses for single ROIs that were divergent between S+ and S- trials. (i)

shows zΔF/F time courses for HEP in the forward and reversed proficient sessions (1

session) and (ii) shows zΔF/F time courses for MO in the forward and reversed proficient

sessions (2 sessions). Time courses were sorted by estimating an agglomerative

hierarchical cluster tree shown on the left that was calculated using the cross correlation

coefficients between all divergent zΔF/F time courses (panel B). The red vertical lines

show (in order): trial start, odorant on, odorant off, reinforcement on and reinforcement

off.

1630

1631 (B) Cross correlation coefficients computed between all per trial zΔF/F time courses

1632 shown in Figure S8A. The coefficients were sorted by the agglomerative hierarchical

1633 cluster tree shown in that figure.

1634

(C) zΔF/F d' time courses for single ROIs that were divergent between S+ and S- trials.

(i) forward (1 session) and (ii) reversed (2 sessions) (d' was calculated for HEP vs. MO,

see methods). Time courses were sorted by estimating an agglomerative hierarchical

cluster tree shown on the left that was calculated using the cross correlation coefficients

between all d' zΔF/F time courses (panel D). The red vertical lines show (in order): trial

start, odorant on, odorant off, reinforcement on and reinforcement off.

1641

1642   (D) Cross correlation coefficients computed between all d' zΔF/F time courses shown in

1643   Figure S8C. The coefficients were sorted by the agglomerative hierarchical cluster tree

1644   shown in that figure.

1645

1646

1647

1648

1649

A

1650

Figure S9. Stimulus decoding accuracy obtained by training in the wide odorant window with different decoding algorithms.

1651

1652

The bar graph shows the mean stimulus decoding accuracy obtained in the 3.1 to 4.1 sec

window for different decoding algorithms trained with data in the 0.5 to 5.5 sec window

for all proficient sessions (66 sessions, 4 mice, related to Figure 6). The different

algorithms are: GLM: generalized linear model, SVM: support vector machine, BDT:

binary decision tree, ANN: artificial neural network and LDA: linear discriminant analysis.

A GLM analysis yields a statistically significant difference between GLM vs. SVM and

GLM vs. ANN (n=684, 671 d.f., F-statistic 203, p-value <0.001, Table S11).

1660

1661

1662

1663 **Figure S10. Examples of single ROI stimulus decoding.**

1664 (A to D) show four examples of single ROI stimulus decoding. i. Decoding accuracy

1665 (black) vs. shuffled (red). ii. zΔF/F time courses for S+ (cyan) and S- (magenta). iii.

1666 Histogram of mean zΔF/F computed between 2 and 4.1 sec. This session had 62 ROIs

1667 with 18 S+ trials and 19 S- trials.

1668

1669

1670

A

Figure showing Discrimination time, Accuracy (y-axis), Time(sec) (x-axis) with time points -4, 0, 4, 8, 12, and a 0.5 Accuracy scale bar.

1671

1672 **Figure S11. Examples of stimulus decoding accuracy time courses for one session.**

1673 The figure shows stimulus decoding accuracy time courses for all ROIs that yielded

1674 accuracies > 0.65 for the proficient session of the examples shown in Figure S10.

1675

1676

Figure S12. Lick decoding accuracy.

A. Time course per trial for the accuracy for ANN decoding of lick fraction from per trial

z∆F/F responses for all trials and all ROIs per session for all proficient sessions (4 mice,

66 proficient sessions). The bounded black line represents the mean accuracy bounded

by the 95% CI. The grey lines are per mouse accuracies. The red line is the stimulus

decoding accuracy calculated after shuffling the stimulus labels (S+ vs S-).

B. Bar graph showing the mean accuracy for different accuracy periods (Pre -1 to 0, Odor

3.1 to 4.1 and Reinf 4.5 to 5.5). Light gray bars are the shuffled stimulus accuracies.

Points are per session accuracies and bars are 95% CIs.

1686

1687 *p<0.05 for a pFDR-corrected t-test or ranksum tests, GLM statistics are in Table S10.

1688

1689

1690   **Video 1. Movie of fluorescence changes for CA1 pyramidal cells**

1691

Go-no Go Associative Learning

Time-Tiled Stimulus Decoding

Multiphoton Calcium Imaging

Heterogeneous dorsal CA1 pyramidal cell activity

1692

1693

In Brief

Ma et al. show that the times for stimulus divergence of pyramidal cell neural activity in

dorsal CA1 are time tiled during the mouse's response in a go-no go associative

learning odorant discrimination task. This sequential neural dynamics of decoding of

odorant discrimination in dorsal CA1 would provide a cognitive time map for decision

making in associative learning.

1700

Highlights

• Spiking of pyramidal cells encodes for stimulus identity and lick behavior in

dorsal CA1 for a mouse engaged in a go-no go odorant associative learning task

• The time of onset of stimulus divergence in neural activity of individual pyramidal

cells in dorsal CA1 is time-tiled

• Learning increases the accuracy of decoding stimulus identity in this task

• The majority of pyramidal cells encode for odorant valence as opposed to

odorant identity

1709

1710

1711