A weak coupling mechanism mediates the recovery stroke of myosin VI: a free energy simulation and string method analysis

Florian E.C. Blanc ● [1,2¤,*], Anne Houdusse ● [2], Marco Cecchini ● [1,*]

1 Molecular Function Engineering, Institut de Chimie de Strasbourg, UMR 7177 CNRS, Université de Strasbourg, F-67083 Strasbourg Cedex, France
2 Structural Motility, Institut Curie, CNRS, UMR 144, PSL Research University, F-75005 Paris, France

¤Current Address: Biophysics of Complex Systems, Institut des Sciences Analytiques, UMR 5280, Université Claude Bernard Lyon 1, CNRS, F-69100 Villeurbanne, France
* florian.blanc@normalesup.org * mcecchini@unistra.fr

Abstract

Myosin motors use the energy of ATP to produce force and directed movement on actin by a swing of the lever arm. ATP is hydrolysed during the off-actin re-priming transition termed recovery stroke. To provide an understanding of chemo-mechanical transduction by myosin, it is critical to determine how the reverse swing of the lever arm and ATP hydrolysis are coupled. Previous studies concluded that the recovery stroke of myosin II is initiated by closure of the Switch II loop in the nucleotide-binding site. Recently, we proposed that the recovery stroke of myosin VI starts with the spontaneous re-priming of the converter domain to a putative pre-transition state (PTS) intermediate that precedes Switch II closing and ATPase activation. Here, we investigate the transition from the pre-recovery, post-rigor (PR) state to PTS in myosin VI using geometric free energy simulations and the string method. First, our calculations rediscover the PTS state agnostically and show that it is accessible from PR via a low free energy transition path. Second, separate path calculations using the string method illuminate the mechanism of the PR to PTS transition with atomic resolution. In this mechanism, the initiating event is a large movement of the converter/lever-arm region that triggers rearrangements in the Relay-SH1 region and the formation of the kink in the Relay helix with no coupling to the active site. Analysis of the free-energy barriers along the path suggests that the converter-initiated mechanism is much faster that the one initiated by Switch II closure, which supports the biological relevance of PTS as a major on-pathway intermediate of the recovery stroke in myosin VI. Our analysis suggests that lever-arm re-priming and ATP hydrolysis are only weakly coupled, so that the myosin recovery stroke is mostly driven by thermal fluctuations and stabilised by nucleotide consumption via a ratchet-like mechanism.

Author summary

Myosin is an ATP-powered motor protein that is crucial for cellular functions like muscle contraction, cell division, and the transport of molecular cargos. Understanding how myosin transforms chemical energy from ATP hydrolysis into

mechanical work is a central open question in structural bioenergetics, which could guide the design of next-generation synthetic nanomachines. In myosin, ATP hydrolysis is coupled to the re-priming of the lever-arm during the recovery stroke transition. Using advanced molecular dynamics simulations, we described the sequence of events and the energetics of the recovery stroke of myosin VI with an unprecedented level of detail. The results support a mechanism in which the re-priming of the mechanical amplifier region of the protein is almost entirely driven by thermal fluctuations and stabilized by ATP hydrolysis. This weakly-coupled mechanism suggests that myosin can "rectify" thermal fluctuations to work efficiently in an isothermal environment dominated by stochastic fluctuations like the cell.

Introduction

Myosins are a superfamily of actin-based motor proteins involved in cellular functions such as cargo transport, endocytosis, cell division and motility [1]. Force and directional movement are produced upon strong binding of the motor domain to F-actin, which triggers the sequential release of the ATP hydrolysis products and the forward movement of the lever-arm domain, or powerstroke [1–3], see Fig 1a. Binding of a new ATP molecule promotes detachment from F-actin and initiates the recovery stroke, *i.e.*, the transition during which the lever-arm is re-primed to its armed configuration. In absence of actin, this transition is rather fast ($\approx 1\,\mathrm{ms}$) and reversible. Interestingly, ATP is hydrolyzed at the end of the recovery stroke rather than during the powerstroke. Therefore, the mechanistic elucidation of this complex, $\sim 1\,\mathrm{ms}$ conformational transition is necessary to understand the structural basis for chemo-mechanical transduction in myosin. Myosin VI (Myo6) is the only-known minus-directed motor, and fulfills specific cellular roles [4,5]. Myo6 harbors two unique inserts including one responsible for directionality reversal, but otherwise shares the same design plan as other myosins [6–8].

High-resolution crystal structures of the initial state (post-rigor state, PR) and final state (pre-powerstroke state, PPS) of the recovery stroke of *Dictyostelium discoideum* Myosin 2 (Dd Myo2) as well as Myo6 have revealed the main subdomain rearrangements, which are similar for both motors. Chief among them are: the re-priming of the converter/lever-arm; the formation of a kink halfway along the conserved Relay Helix (RH); and the closure of the Switch II loop over the γ-phosphate of ATP [7,9]. This latter rearrangement turns on ATPase function by stabilizing an active site geometry conducive to hydrolysis [10–13]. A key open

question is how Switch II closure couples to converter re-priming. To date, most $_{25}$
computational analyses of the PR→PPS transition mechanism have predicted that $_{26}$
Switch II closure initiates the recovery stroke [14–25]. Among them, the popular model $_{27}$
by Fischer and co-workers posits a strongly, or *mechanically* coupled mechanism where $_{28}$
Switch II closure directly drives the rotation of the converter [14, 17]. In this model, $_{29}$
structural changes occur in progressive, concerted fashion and involve a two-step $_{30}$
rearrangement in the RH, which sits between the active site and the converter. $_{31}$

In sharp contrast, we have recently proposed a mechanism for the recovery stroke $_{32}$
of Myo6 in which the rotation of the converter precedes the closure of Switch II and is $_{33}$
uncoupled from it until the later stages of the transition [26]. Because this mechanism $_{34}$
entails near complete converter re-priming driven purely by random fluctuations, we $_{35}$
named it the "ratchet-like" model. It involves a putative structural intermediate called $_{36}$
pretransition state (PTS), which we characterized by X-ray crystallography and $_{37}$
all-atom Molecular Dynamics (MD), see Fig 1a. The PTS structure (PDB: 5O2L) $_{38}$
exhibits hallmarks of the post-recovery state, namely a (nearly) re-primed converter $_{39}$
and a kink in the Relay helix (Figure 1b-e), yet with an open Switch II. The $_{40}$
ratchet-like model of the recovery stroke therefore comprises two major conformational $_{41}$
transitions, the PR → PTS transition in which (most of) the converter re-priming $_{42}$
takes place, and the PTS → PPS transition in which Switch II closes. In this scenario, $_{43}$
converter swing is weakly, or *statistically*, coupled to Switch II closure. $_{44}$

Whether the recovery stroke of Myo6 proceeds by a strongly-coupled, Switch $_{45}$
II-initiated or statistically-coupled, ratchet-like mechanism remains an open question. $_{46}$
To explore the recovery stroke mechanism with atomic resolution, we turn to all-atom $_{47}$
MD simulations with enhanced sampling strategies. Geometric free energy calculations $_{48}$
(including the adaptive biasing force method, or ABF) evaluate the potential of mean $_{49}$
force (PMF) along low-dimensional collective variables (CVs) describing the $_{50}$
transition [27–30]. The related string method in collective variables (CVSM) identifies $_{51}$
the minimum free energy path (MFEP) in a CV-space of arbitrary dimension [31, 32], $_{52}$
which makes it ideal to elucidate the transition mechanism between two known $_{53}$
structures. The string method has been used to describe conformational transitions in $_{54}$
molecular machines, see for example [33–35]. For myosin, Ovchinnikov et al. applied it $_{55}$
to the internal conformational transition of the Myo6 converter [36, 37] and Cui et al. $_{56}$

used it in combination with a quantum potential to map ATP hydrolysis pathways in Dd Myo2 [12]. To our knowledge, the string method has not been applied yet to the elucidation of conformational transitions of the full-length myosin motor domain.

We report here on a computational analysis of the mechanism and energetics of the transition from PR to PTS, *i.e.*, the first major step of the recovery stroke in the ratchet-like model. First, we use ABF to map a coarse-grained free energy landscape of the transition. Consistent with the ratchet-like prediction, we identify agnostically a state corresponding to PTS, and show that it is accessible from PR by a low free energy path, supporting its relevance as an on-pathway intermediate. Second, we perform path optimizations with the CVSM to determine a higher-resolution description of the transition. These independent calculations agree remarkably well with the ABF map and reveal a plausible mechanism by which near-complete re-priming of the mechanical amplifier region is triggered by a spontaneous converter/lever-arm swing, eventually resulting in the formation of the kink in the Relay helix while Switch II remains open. Despite some uncertainty as to the nature of the rate-limiting step (lever-arm movement vs formation of the kink), free energy barriers suggest that the ratchet-like pathway is explored much faster than the Switch-II initiated one. Therefore, our results support the PTS state as a functional intermediate of the myosin VI cycle, and the ratchet-like model for the myosin VI recovery stroke.

Results

To investigate the recovery stroke of Myo6, we combine extended ABF (eABF, [38]) and the CVSM to take advantage of their respective strengths. We introduce CVs X_c (Fig. 1d) to measure the position of the converter relative to the motor domain and $\Delta RMSD$ (Fig. 1e) to describe the local conformation of the RH (*i.e.,* the formation of the kink); see Methods for a precise definition. Together, these CVs describe two characteristic rearrangements from PR to PTS. We perform an eABF calculation over $(X_c, \Delta RMSD)$ to determine the coarse-grained free energy landscape of early recovery stroke in the ratchet-like model, starting from the PR state. Then, we carry out string optimizations in a higher-dimensional space encompassing degrees of

Fig 1. Overview of the PR → PTS transition. (a) Simplified Myosin VI functional cycle including the putative PTS. (b) Equilibrated PR structure highlighting the structural elements involved in the recovery stroke. (c) Equilibrated PTS structure, colored as in b. (d) Comparison between PR (transparent) and PTS (opaque) showing the movement of the converter, the tilting of the SH1 helix, and the kink and re-orientation of the C-terminal fragment of the Relay Helix. Typical values for the corresponding CVs are given. (e) Comparison of the backbone structure of residues 485-491 of the Relay helix before and after kink formation. Typical values of the ΔRMSD are given.

freedom left out of the eABF calculation, to determine a high-resolution picture of the transition mechanism from PR to PTS. This strategy yields an integrated structural and energetic description of the early stages of the recovery stroke and enables consistency checks between separate sets of simulations.

Coarse-grained free energy landscape of the PR $\rightarrow$ PTS transition

The full coverage of configurational space, smooth convergence of the free energy gradient estimates and low estimated statistical error suggest that the eABF calculations are converged (see Supplementary Text 1).

eABF discovers the PTS state without prior knowledge

The free energy landscape computed by eABF reveals 5 local free energy minima, including the PR basin (upper right corner), see Fig. 2a. Strikingly, a wide basin encompassing 2 sub-minima is found in the lower left corner of the free energy landscape, corresponding to a kinked Relay helix and (partially) re-primed converter. Interestingly, we observe that the active site -as characterized by the d_1 and d_γ distances, see Methods- remains open throughout eABF sampling, including in the identified metastable states (Supplementary Figure 9a). Therefore, the calculations indicate that there exists a metastable Myo6 configuration with open active site, nearly re-primed converter and kinked Relay helix. We note that if the closure of Switch II were strongly coupled to converter movement and/or Relay helix kink formation, closed active-site conformers would have been preferentially sampled in the lower-left region of the free energy landscape, which is not the case. Additionally, this basin's characteristics are consistent with the PTS state, as evidenced by comparison to previously reported unbiased MD simulations of Myo6 [26], see Supplementary Fig. 9b. We find that both sub-basins (labelled PTS1 and PTS2 in Fig 2a) correspond to converter positions and Relay helix conformations explored in unbiased PTS simulations. Finally, representative conformers extracted from the PTS1 basin by clustering demonstrate marked structural similarity to the PTS crystal structure (Fig. 2b). Taken together, these findings imply that not only a partially re-primed state

with open active site exists, but also that the PTS structure is representative of this state, indicating that eABF has discovered the PTS state without prior knowledge of the PTS structure (see Methods). The finer structure of the free energy landscape suggests -also in agreement with earlier unbiased MD [26]- that the PTS state encompasses at least two distinct metastable states, PTS1 and PTS2, separated by a low, $3 \, \text{kcal} \, \text{mol}^{-1}$ free energy barrier, and which correspond to slightly different RH conformers and average converter positions. The PTS2 converter position is closer to PPS, suggesting it is on the recovery stroke pathway. From PTS1, the transition to PTS2 entails a further $7 \, \text{Å}$ converter movement associated with a slight re-orientation of the RH. This transition corresponds to rapid fluctuations within the PTS basin, as demonstrated by their reversible sampling on the $\sim 100 \, \text{ns}$ timescale in earlier unbiased MD simulations, and agrees with our previous observation that the converter is highly dynamic in the PTS state [26].

Transition tube and mechanism

The free energy landscape shows that a low free energy transition tube connects PR to PTS. This indicates that the PTS state is kinetically accessible from PR and supports the hypothesis that it is an on-pathway intermediate along the recovery stroke. Two local free energy minima are identified within the transition tube, labelled I_A and I_B on Fig 2a. We use structural clustering to reveal their structural features. This analysis indicates that I_A corresponds to an intermediate along the PR$\rightarrow$ PTS transition where the converter has moved by $7.8 \, \text{Å}$ in the re-priming direction, promoting partial tilting of SH1 and bending of the RH in response, but not yet formation of the kink (Fig 2c). In I_B, the converter has moved by only $6.5 \, \text{Å}$ and its orientation is quite different from I_A, and closer to PR ((Fig 2d and Supp. Fig. 10a). Moreover, in I_B the RH backbone differs both from the regular helical structure observed in PR and the kink observed in PTS (and PPS) crystal structures. Rather, the RH exhibits local unfolding and bending near residues 490 to 494 (*i.e*, ~ 5 residues in C-ter of the canonical kink), see Fig 2e and Supp. Fig. 10b.

Although the transition tube is suggestive of a sequential mechanism with I_A, then I_B as on-pathway intermediates, the differences in converter position/orientation and RH structure are also consistent with I_A and I_B belonging to two different pathways

(Paths A_{ABF} and B_{ABF}, respectively in cyan and pink on Fig. 2). In Path A_{ABF}, the initiating event is a $\approx 8\,\text{Å}$ movement of the converter which crosses the highest free energy barrier along this path ($6 \pm 1\,\text{kcal}\,\text{mol}^{-1}$, see SI for statistical error estimation), suggesting it may represent the rate-limiting step. The second step to reach PTS1 is the formation of the kink in the RH, which contributes an additional $\sim 2\,\text{Å}$ of converter swing. In Path B_{ABF}, the first step is a $6.5\,\text{Å}$ movement of the converter associated with local disruption of the RH backbone, and the second step corresponds to the formation of the "canonical" kink in the RH along with a further $3.5\,\text{Å}$ converter swing to PTS1. Despite their differences, both paths predict an early converter movement as the first step of the recovery stroke in Myo6. Notably, our previous unbiased simulations of the PR state did capture spontaneous converter uncoupling without Relay helix kinking [26], which may indicate that Path A_{ABF} is more likely to represent the dominant mechanism to reach PTS (Supplementary Figure 9, simulation PR (3)).

The analysis of the free energy landscape allows us to 1) validate the relevance of PTS as an accessible metastable state from PR and 2) draw a first mechanistic picture of this transition. However, it does not reveal how other important structural rearrangements, such as changes in the SH1 helix, are coupled to the transition. We now address this with the string method in collective variables.

String method analysis of the PR $\rightarrow$ PTS transition

First, we performed preliminary string optimizations in $(X_c,\ \Delta RMSD)$, which showed that the minimum free energy path (MFEP) for the PR $\rightarrow$ PTS transition lies within the transition tube predicted by eABF (Supp. Fig. 12, Supp. Fig. 13, Supplementary Text 2). This independent agreement confirms that the general features of the transition predicted by eABF are robust. Then, we set out to describe the transition mechanism with higher resolution. To this end, we used the string method with 12 collective variables giving a more detailed insight into the rearrangements involved in the PR $\rightarrow$ PTS transition (see Table 1; we note that $\Delta RMSD$ was not used as a supporting CV for 12D CVSM calculations because the 4 kink distances serve the same purpose, but with better descriptive power).

Fig 2. Analysis of the PR → PTS transition by eABF. (a) Free energy landscape obtained by eABF with metastable states labelled. The dotted green (respectively orange) line materializes the value of X_c in the PR crystal structure (respectively PTS crystal structure). The arrows indicate putative transition pathways, namely Path A_{ABF} (cyan) and Path B_{ABF} (pink). (b-d). Representative conformations of the converter and Relay and SH1 helices of the PTS1 (b, dark red), I_A (c, cyan) and I_B (d, bright pink) metastable states, compared to PR (transparent green) and PTS (transparent orange) equilibrated structures. Structures are aligned onto the N-terminal fragment of the Relay helix (residues 468-482) to better isolate changes in Relay and SH1 helix orientation and converter position. (e) Close-up on the disruption of the Relay helix in I_B.

.

The starting (or guess) path for string optimizations was obtained by "uplifting" and regularizing the 2-dimensional string optimized over the eABF free energy landscape to the 12-dimensional space (see Supplementary Text 2). Two separate string optimizations, with slightly different parameters, were performed until convergence (Supp. Fig. 14, Supplementary Text 2). Visual analysis of the structures along averaged strings over the last 50 iterations shows that the two calculations converged towards a similar sequence of events, which represents the most likely pathway for the PR → PTS transition of Myo6, and which we call Path A. Backprojection of the two converged Path A strings (Strings A1 and A2) onto the $(X_c, \Delta RMSD)$ plane reveals that they both lie within the eABF-predicted transition tube. Interestingly, the two converged strings visit I_A, but not I_B, see Supp. Fig. 15. To obtain an energetic view of the transition mechanism, we then performed an umbrella sampling calculation along string A1 with a finer discretization (from 32 to 128 equally-spaced images) and computed the free energy profile along the string, and 2D free energy maps along pairs of observables (see Methods and Supplementary Text 3). From the analysis of string A1, we now describe the sequence of events, structural

couplings, and energetics governing the PR → PTS transition. ₁₉₃

Overview of the mechanism

Free energy profile and comparison with eABF The free energy profile along
string A1 computed by Umbrella Integration [39] shows 4 metastable states (Fig. 3a),
indicating a 3-stage mechanism. Projection of the umbrella sampling simulations along
string A1 onto the $(X_c, \ \Delta RMSD)$ with MBAR [40, 41] reveals a free energy
landscape very similar to the eABF one, but with a number of remarkable differences
(Fig. 3b). Free energy minima corresponding to states PR, I_A and PTS1 are identified.
PTS2 is not detected because the string does not extend to this basin. I_B is not seen
despite its position in configurational space being clearly sampled. This confirms the
proposal that I_A and I_B do not belong to the same transition pathway. Comparison of
I_A representative structures sampled independently in the eABF and string
calculations reveals striking similarity (Supplementary Figure 11), which confirms that
the same metastable intermediate is picked up by both calculations and justifies using
the same denomination indifferently. We conclude that Path A_{ABF} and Path A are
virtually identical, and we can now detail the transition mechanism.

Step 1: PR → I_A: converter rotation re-orients Relay and SH1 helices
Starting from the PR state, the first step is a thermally-activated (stochastic) ≈8 Å
movement of the converter leading the motor domain into the I_A intermediate (Fig
3c,d, Supp. Fig. 16). This is consistent with the eABF prediction, but the free energy
barriers differ, with umbrella sampling indicating a ≈1 kcal mol^{-1} barrier (Fig 3a),
much smaller than the 6 kcal mol^{-1} eABF barrier (Fig 2). The converter displacement
reduces the total number of contacts it makes with the N-terminal subdomain (Supp.
Fig. 17a,b,e) which increases the converter's positional freedom relative to PR,
because it is attached to the main body only through the flexible Relay and SH1
helices. The most energetically costly event in the PR → I_A step seems to be the
disruption of the converter/N-terminal contacts. Since these are mostly non-specific
hydrophobic interactions, the corresponding free energy barrier is rather small and can
probably be crossed reversibly on relatively short timescales. This is consistent with
spontaneous ∼100 ns converter uncoupling in unbiased MD of PR we previously

Fig 3. Free energy profiles and mechanism of the PR → PTS transition.
(a) PMF along the path obtained by Umbrella Integration. (b) 2D PMF along
$(X_c, \Delta RMSD)$ obtained by MBAR. Arrows indicate transition steps between
metastable states. (c-f) Sequence of metastable states in the PR → PTS transition
from string calculations. Arrows indicate the movements of structural elements. For
easier comparison, the PR structure is shown in transparent grey. (g-j) Mechanism of
formation of the kink in the RH. Arrows indicate backbone and side chain movements.
H-bonds are shown as green dotted lines, shading indicating weak binding. (c,g) From
PR, in response to the first converter movement, the RH C-ter shifts and rotates,
which extends the N485O:L489N H-bond and brings L489 closer to L700. (d,h) In I_A,
the second converter movement translates into corkscrew movement of the RH along
its axis, shifting K490N and E491N by a quarter of helical turn, but complete rotation
of RH is prevented by the L489/L700 contact. Instead, the I488-L489 peptide bond
flips, which breaks the I488O:E492N H-bond. This deprives R487O and I488O from
H-bond partners, and weakens the N485O:L489N H-bond because of unfavorable angle.
In counterpart, non-canonical H-bonds E486O:E491N and N485O:K490N form. (e,i)
In I_A2, structural tension is relieved by movements of the SH1 helix and the Relay
loop. This frees steric constraints on the L489 and L700 side chains, which exchange
their positions while remaining in contact. In addition, I488-L489 flips back, allowing
the I488O:E492N H-bond to reform. (f,j) In PTS1, the hydrophobic switch is
rearranged, stabilizing the kink in the Relay helix. R487O is free of H-bond partner.

reported [26]. Although I_A is in fast interconversion with PR, it represents an ²²³
important intermediate because converter uncoupling from the main body is a ²²⁴
pre-requisite to facilitate its swing. ²²⁵

Both the RH C-ter moiety (residues 489 to 499) and the SH1 helix re-orient ²²⁶
concurrently with the converter movement (Fig 3c,d, Supp. Fig. 16). The C-ter ²²⁷
moiety of the RH is strongly bound to the converter through specific contacts, and its ²²⁸
N-ter moiety is anchored within the main body. As a result, the RH bends following ²²⁹
the movement of the converter. RH bending entails only limited backbone ²³⁰
perturbation, with a transient extension of the N485O:L489N hydrogen bond close to ²³¹
the bending point (Fig 3g,h, Supplementary Figure 16). This enables a ²³²
"corkscrew"-like rotation of the RH C-ter moeity around the helical axis, which brings ²³³
the L489 side chain in contact with L700 on the SH1 helix, resulting in tilting of the ²³⁴
SH1 helix. This steric clash constitutes a first blocking point to the free movement of ²³⁵
the converter, which is later relieved through extensive conformational changes. ²³⁶

Step 2: $I_A \rightarrow I_A2$: converter movement drives formation of the kink At the ²³⁷
beginning of Step 2, a second movement of the converter is translated into further ²³⁸
angular re-orientation of RH and SH1 (Fig 3d,e, Supp. Fig. 16). However, unlike Step ²³⁹
1, the RH backbone can no longer accommodate the strain only through bending, ²⁴⁰
because this movement is blocked by the clash of L489/L700 side chains. Therefore, ²⁴¹
the "corkscrew"-like motion of the RH backbone is coupled with dramatic alterations ²⁴²
to the H-bonding pattern, namely the formation of the E486O:E491N and the ²⁴³
disruption of the E486O:K490N, R487O:E491N, I488O:E492N H-bonds (Fig 3h,i). ²⁴⁴
N485O:L489N, which was extended by about 0.5 Å in I_A, returns to perfect geometry. ²⁴⁵
And, atoms N485O and K490N are brought closer but not yet quite enough to form a ²⁴⁶
H-bond. The cost of breaking these backbone H-bonds likely explains why this step ²⁴⁷
would be rate-limiting for the PR $\rightarrow$ PTS transition, as suggested by the free energy ²⁴⁸
profile (highest free energy barrier, $\approx 4\,\mathrm{kcal\,mol^{-1}}$, Fig. 3a). Upon reaching I_A2, the ²⁴⁹
Relay helix is clearly kinked, even though the H-bonding pattern differs slightly from ²⁵⁰
what is observed in PTS. Notably, I_A2 features outward-pointing, unbound R487O ²⁵¹
and I488O backbone atoms, which may explain why I_A2 is rather high in free energy ²⁵²
(Figure 3a and Fig 3i), even if the formation of at least one non-canonical H-bond in ²⁵³

the RH backbone (E486O:E491N) makes this state metastable.   254

The analysis of the $I_A \to I_A2$ step indicates that the main driving force for the   255
formation of the RH kink is the movement of the converter. It is plausible that   256
capturing a large enough converter movement to drive the costly kink formation would   257
require first putting the converter in an uncoupled state. There, the converter can   258
efficiently harness fluctuations from the bath and amplify them into functional   259
rearrangements within the motor domain. This shows why it is important to first visit   260
I_A as an easily accessible uncoupled state.   261

Step 3: $I_A2 \to$ PTS1: seclusion of the SH1 helix and rearrangement of the   262
hydrophobic switch   The tight packing of the L489/L700 side chains is preserved   263
throughout the transition to I_A2, including upon formation of the kink. This builds up   264
structural frustration at the Relay/SH1 interface. During the $I_A2 \to$ PTS1 transition,   265
this frustration is relieved through two consecutive rearrangements. First, the SH1   266
helix moves by $\approx 1\,\text{Å}$ parallel to the RH in a "piston-like" motion, and re-orients   267
upward by $\approx 5\,\text{deg}$ (Fig. 3e,f, Supplementary Figure 16). This creates enough space to   268
allow packing L489 against the SH1 helix. Second, a movement of the Relay loop   269
relieves the steric hindrance from the Y508 side chain (Fig 3i,j). This allows the   270
L489 and L700 side chains to eventually exchange their positions (Fig 3i,j). In the   271
process, the I488-L489 peptide bond rotates, allowing I488O to re-form a H-bond with   272
E492N while the N485O:L489N hydrogen bond breaks. Fischer et al. and Baumketner   273
independently proposed that a similar transition for the Dd Myo2-homologous side   274
chains (Dd Myo2 F487, F506 and I687) stabilizes the kink [14, 24]. By analogy to the   275
"aromatic switch" of Fischer et al., we call this triplet of side chains the "hydrophobic   276
switch" for Myo6. The sequence of events captured by our calculations for L489 and   277
L700 is reminiscent of the mechanism proposed by these investigators, but not entirely   278
consistent with it (see Discussion). Because the Relay loop is very flexible, its   279
movement is likely to be a stochastic, low-barrier event weakly coupled to the rest of   280
the transition. Thus, the movement of the SH1 helix, driven by the push of L489,   281
likely accounts for most of the $\approx 3\,\text{kcal}\,\text{mol}^{-1}$ barrier (Fig. 3).   282

No coupling to the active site is detected

We searched for a structural response of the active site to the rearrangements of the $\quad$ 284
PR $\rightarrow$ PTS transition. First, we computed the free energy map along the distances $\quad$ 285
defining the catalytically-competent configuration from umbrella sampling (Fig. 4a,c), $\quad$ 286
which shows that the position of Switch II is nearly unaffected by the transition, like $\quad$ 287
our observations from eABF. Then, we looked into a possible communication pathway $\quad$ 288
between SH1 and the active site proposed by Fischer et al. for Dd Myo2 [17]. These $\quad$ 289
authors proposed that the "piston-like" motion of the SH1 helix translates into the $\quad$ 290
formation of a hydrogen bond between P-loop and Switch II, which may contribute to $\quad$ 291
ATPase activation (see also Discussion). To assess whether a similar mechanism is at $\quad$ 292
play during the PR $\rightarrow$ PTS transition, we computed the free energy map along the $\quad$ 293
SH1 helix longitudinal projection Z_{SH1} (accounting for the piston-like motion) versus $\quad$ 294
the S153N:F460O hydrogen bond distance between P-loop and Switch II, see Fig. $\quad$ 295
4b,d. Although the free energy map suggests that the piston movement brings S153 $\quad$ 296
and F460 slightly closer, this does not result in the formation of a hydrogen bond $\quad$ 297
between these two residues. The contact between the wedge loop and the SH1-SH2 $\quad$ 298
junction was proposed as the key element mediating structural communication $\quad$ 299
between Switch II and the SH1 helix by Fischer et al. [17]. In our simulations, F582 on $\quad$ 300
the wedge loop is brought somewhat closer to the SH1-SH2 junction as the transition $\quad$ 301
proceeds (Fig. 4e), but this is not sufficient to trigger the global movement of the $\quad$ 302
wedge loop which would drive the formation of the P-loop / Switch II hydrogen bond. $\quad$ 303
We conclude that the near-complete rearrangement of the mechanical amplifier $\quad$ 304
element (Relay/SH1/Converter region), which is required to reprime the lever arm, $\quad$ 305
proceeds without detectable effect on the active site. $\quad$ 306

Alternate mechanism for the PR $\rightarrow$ PTS transition

To assess the robustness of our findings to the starting point of string optimizations, $\quad$ 308
we carried out a second set of string calculations where the guess path was obtained as $\quad$ 309
the straight path joining PR to PTS in 12D CV-space (Supplementary Text 2). This $\quad$ 310
resulted in the so-called Path B, which is also consistent with the eABF transition $\quad$ 311
tube, but does not overlap with Path A. Instead, Path B admits I$_B$ as an intermediate $\quad$ 312

Fig 4. The active site remains open throughout the PR→PTS transition.
(a) Close-up on the active site for a representative PTS configuration from umbrella sampling. The critical salt-bridge (R205CZ-E461CD, distance d_1) and the Switch II-ATP hydrogen bond (G459N:ATPO1G, distance d_γ) are clearly not formed, corresponding to an ATPase-incompetent active site. (b) Same configuration seen from a different angle, highlighting the SH1-SH2 junction, the wedge-loop and the P-loop. Although F582 is close to G693 at the SH1-SH2 junction, the P-loop-Switch II hydrogen bond (S153N:F460O) is clearly not formed. (c) 2D PMF over d_1 and d_γ computed along the transition. The closed Switch II state is not detected. (d) 2D PMF over the piston-like movement of the SH1 helix (Z_{SH1}) and the P-loop-Switch II hydrogen bond distance (S153N:F460O) computed along the transition. The formed hydrogen bond is not detected. (e) 2D PMF over the piston-like movement of the SH1 helix (Z_{SH1}) and the distance between the wedge loop and the SH1-SH2 junction (F582-G693 distance). The completion of the piston-like movement of SH1 favors closer approach between the wedge loop and the SH1-SH2 junction.

and is virtually identical to Path B_{ABF}, *i.e.,* a mechanism in which the RH transiently ₃₁₃ develops a secondary kink (see Supplementary Text 4 and Supplementary Figure 19). ₃₁₄ Because Path B entails more structural perturbation than Path A (two kinks instead ₃₁₅ of one), it is less likely to represent the main sequence of events of the PR→PTS ₃₁₆ transition, hence why our focus is on Path A. Path B may contribute marginally to the ₃₁₇ transition flux, but this does not change our conclusions that lever arm priming comes ₃₁₈ from harnessing converter fluctuations without immediate coupling to the active site. ₃₁₉

Discussion ₃₂₀

Molecular motors like myosin convert chemical energy from ATP hydrolysis into ₃₂₁ mechanical work following established thermodynamic principles [42]. However, how ₃₂₂ these principles are implemented at the molecular level remains unclear. We have ₃₂₃ tackled this question using all-atom MD simulations and free energy calculations. In ₃₂₄ particular, we focused on the recovery stroke transition that couples the re-priming of ₃₂₅ the myosin lever arm to ATP hydrolysis. Our results shed light on the ₃₂₆ chemo-mechanical transduction mechanism and illuminate structural intermediates ₃₂₇ visited during this off-actin transition, which may be targeted by allosteric modulatory ₃₂₈ ligands [43, 44]. ₃₂₉

Using advanced free energy simulations started from the X-ray structure of the ₃₃₀ post-rigor (PR) state of myosin VI, we illuminated a low-energy transition pathway ₃₃₁ connecting PR to a previouly characterized structural intermediate, named PTS, with ₃₃₂ an unprecedented level of detail. Consistent with previous observations [26], our ₃₃₃ calculations suggest that the recovery stroke of myosin VI is initiated by a movement ₃₃₄ of the converter by thermal fluctuations accompanied by the kinking of the Relay ₃₃₅ helix, which corresponds to the highest energy barrier, tilting of the SH1 helix and ₃₃₆ stabilization via a switch of hydrophobic interactions between residues at the ₃₃₇ Relay/SH1 interface. Remarkably, these calculations not only "re-discover" PTS ₃₃₈ agnostically but also show that the same sequence of events could be obtained using ₃₃₉ two different free energy simulation methods, which strenghtens the relevance of these ₃₄₀ findings. Most importantly, the striking observation is that repriming of the converter ₃₄₁ subdomain, *i.e.,* a key element of the mechanical amplifying region of the motor ₃₄₂

protein, precedes closure of Switch II at the nucleotide-binding site. In addition, our ₃₄₃ free energy analysis predicts that the rate-limiting step to PTS is $\leq 6\,\text{kcal}\,\text{mol}^{-1}$ with ₃₄₄ Switch II open, whereas the free energy barrier for closing Switch II from PR was ₃₄₅ $12\,\text{kcal}\,\text{mol}^{-1}$ [26]. Assuming similar pre-exponential factors, these results suggest that ₃₄₆ the converter-initiated mechanism would be $\approx 23\,000$ times faster than the Switch ₃₄₇ II-initiated scenario. Intriguingly, these observations suggest that lever-arm repriming ₃₄₈ and ATP hydrolysis are only weakly coupled, so that the recovery stroke transition ₃₄₉ can be initiated by thermal fluctuations and driven via modulation of kinetic barriers. ₃₅₀ These observations thus support the conclusion that the recovery stroke in Myo6 is ₃₅₁ mediated by a "ratchet-like" mechanism. As compared to a Switch II-initiated ₃₅₂ mechanism, a ratchet-like mechanism would increase the motor efficiency by ensuring ₃₅₃ that ATP hydrolysis takes place only once the mechanical amplifier region is ₃₅₄ re-primed,+ avoiding futile ATP consumption [45], and allowing the motor to ₃₅₅ effectively harness and "rectify" ubiquitous Brownian fluctuations. ₃₅₆

Our mechanistic proposal contrasts with previous models of the recovery stroke, ₃₅₇ most of whom predicted Switch II closure as the initiating event. The mechanism ₃₅₈ proposed by Fischer and co-workers in Dd Myo2 consists of two phases [14, 17]. In ₃₅₉ phase I, a movement of the converter couples to Switch II closure through a rigid-body ₃₆₀ "seesaw" movement of the RH. In phase II, an additional movement of the converter ₃₆₁ couples to SH1 helix tilting, RH kink formation, and the establishment of Switch ₃₆₂ II/P-loop interactions. This scenario is plausible and consistent with Dd Myo2 ₃₆₃ mutants, but it derives from zero-temperature calculations, which are inherently ₃₆₄ biased towards strong coupling. Elber and West calculated a zero-temperature ₃₆₅ transition path similar to Fischer's model and used finite-temperature MD to evaluate ₃₆₆ the transition rate along this path [22]. Although the predicted rate is consistent with ₃₆₇ experimental estimates, these simulations are likely too short (a few ps) to appreciably ₃₆₈ deviate from the zero-temperature transition path. Woo and Harris explored the ₃₆₉ recovery stroke of Scallop myosin II by umbrella sampling. These calculations ₃₇₀ suggested a weak coupling between Switch II closure and converter rotation, but it is ₃₇₁ unclear whether the short simulation trajectories (<6 ns) could faithfully reveal the ₃₇₂ sequence of events and the energetics [18, 19]. Using Targeted MD (TMD) simulations ₃₇₃ from the PR structure of Dd Myo2, Cui and co-workers predicted an early movement ₃₇₄

of the converter followed by RH kinking and Switch II closure [21], which is reminiscent ₃₇₅ of the mechanism proposed here. However, partial Switch II closure (critical ₃₇₆ salt-bridge formation) was observed at the beginning of the transition consistent with ₃₇₇ an early Switch II closure. Baumketner and Nemeslov came to the same conclusion ₃₇₈ using high-temperature simulations of the PR state in implicit solvent [23]. Later, ₃₇₉ using enhanced sampling Baumketner convincingly showed that rearrangements within ₃₈₀ the mechanical amplifier region are statistically coupled, but its fragmental model ₃₈₁ lacking Switch II prevented him from exploring the consequences at the nucleotide ₃₈₂ binding site [24, 25]. In brief, methodological and sampling limitations (with respect to ₃₈₃ current state-of-art) may explain why earlier studies did not detect PTS. In addition, ₃₈₄ the usage of different myosin isoforms might also contribute to diverging outcomes. ₃₈₅

The ratchet-like model presented here shares features with the more broadly ₃₈₆ accepted Switch II-initiated mechanism by Fischer's and Baumketner's although with ₃₈₇ important differences. First, in the Switch II-initiated scenario, the seesaw motion of ₃₈₈ the Relay helix precedes its kinking, as the latter involves threading the bulky Dd ₃₈₉ Myo2:F487 between the RH backbone and the Relay loop, which would be sterically ₃₉₀ prevented until the RH seesaw motion opens the way. By contrast, we find that a ₃₉₁ movement of the Relay loop is sufficient to enable threading of the homologous L489 ₃₉₂ in Myo6 before the RH seesaw motion, thus resulting in an opposite order of events. ₃₉₃ Second, in the Switch II-initiated scenario it was found that SH1 tilting is coupled to ₃₉₄ the formation of a H-bond between Switch II and P-loop at the nucleotide-binding site ₃₉₅ (DdMyo2:S183N-F458O) via the wedge loop. Yet, we do not observe the formation of ₃₉₆ the corresponding H-bond in Myo6 (Myo6:S153N-F460O) and the PTS crystal ₃₉₇ structure does not exhibit it either, suggesting that it forms during the PTS → PPS ₃₉₈ transition instead. Third, in the Switch II-initiated scenario, the tilting of the SH1 ₃₉₉ helix drives the converter rotation, whereas we come to the opposite conclusion. ₄₀₀ Fourth, in the converter-initiated scenario, the seesaw motion of the Relay helix ₄₀₁ (coupled to an additional converter movement and Switch II closure) occurs in the ₄₀₂ PTS → PPS transition, which will be detailed elsewhere. ₄₀₃

Based on these observations, the Switch II- and converter-initiated mechanisms ₄₀₄ emerge as plausible alternative scenarios depending on whether RH kinking occurs ₄₀₅ before (converter-initiated) or after (Switch II-initiated) the seesaw motion (Fig. 5). ₄₀₆

In this interpretation, the PTS state would be naturally described as the post-kink, pre-seesaw intermediate along the converter-initiated pathway. A different intermediate with a partially re-primed converter, an unkinked but "seesawed" Relay helix, and a closed Switch II should be observed along the Switch II-initiated pathway [17]. To the best of our knowledge, the latter has not been observed experimentally yet. Whether the PTS state and ratchet-like recovery stroke are common to all myosins remains an open question. Dd Myo2 mutants interpreted as supporting the Switch II-initiated recovery stroke do not disprove the ratchet-like pathway [26]. The same is true for pathogenic mutations disrupting the converter-Relay interactions in Human β-cardiac myosin, which would perturb the recovery stroke regardless of its mechanism, as supported by recent free energy calculations in Dd Myo2 [46] and β-cardiac myosin [47]. In addition, recent free energy calculations on β-cardiac myosin also identified a PTS-like intermediate [44]. Last, the ratchet-like mechanism is consistent with the structural intermediate of smooth muscle myosin II trapped off-actin by the CK-571 drug [43], which exhibits a partially re-primed converter, an "un-kinked" Relay helix and an open Switch II. To advance on this question, detailed computational investigations of the recovery stroke mechanism in other myosin isoforms [48] are needed. As a bonus, these analyses would enable the characterization of cryptic binding pockets for the design of myosin's allosteric modulators [44, 49, 50].

Materials and methods

Preparation of structural models for Molecular Dynamics simulations

Structural models of the Myo6 PR (PDB ID: 2VAS, [8]) and PTS (PDB ID: 5O2L) states were prepared from the corresponding crystal structures according to the protocol reported in [26]. Briefly, missing loops were built with MODELLER, then each structure was placed in an orthorhombic box of TIP3P water molecules supplemented with Na^+ and Cl^- ions to reach a $150\,mM$ salt concentration. Each system was energy-minimized with harmonic restraints on the protein atoms, followed

Fig 5. Unified mechanistic framework for the myosin recovery stroke.

by a 1 ns-long heating MD simulation to 300 K, then a 2 ns-long equilibration MD at 436
constant temperature (300 K) and pressure (1 bar) during which harmonic restraints 437
were smoothly scaled down. In the present work, only the PR system was used to 438
launch new simulations. Simulations used NAMD 2.13 [51] or Gromacs 2021.5 [52]. 439

Collective variables 440

Collective variables (CVs) were used to characterize, and in some cases bias, the 441
structural dynamics of Myo6, focusing on the structural elements involved in the 442
recovery stroke (Fig 1b-e). The complete list of CVs is given in Table 1. All CVs were 443
computed with the *colvars* module [53]. 444

Converter 445

The Cartesian coordinates of the converter's center of geometry in the frame formed 446
by the principal axes of the motor domain are denoted as X_c, Y_c, Z_c (X', Y' Z' 447
in [26]). They capture the movements of the converter relative to the motor domain 448
during the recovery stroke. The reader is referred to reference [26], SI Appendix, for 449
the full description of the converter CVs. The interactions of the converter with the 450
main body of the motor domain are described by a set of interatomic distances i_j (see 451

Table 1 for details).

Relay/SH1 elements

Angular descriptors of the orientation of the Relay helix C-terminal fragment θ_{RH} and the SH1 helix θ_{SH1} with respect to the PR crystal structure were defined as described in [26], SI Appendix. Following Baumketner [24, 25] we introduced the distance $d_{R/SH1}$ between the centers of geometry of the CA atoms of residues 469-482 (N-terminal fragment of the Relay helix) and 693-703 (SH1 helix). To characterize the "piston"-like motion of SH1 highlighted by Fischer and co-workers, we defined Z_{SH1} as the orthogonal projection of the SH1 helix' center of geometry onto the longitudinal principal axis of the motor domain (*i.e.*, the same axis as for Z_c). The local conformation of the Relay helix backbone was described by a $\Delta RMSD$ CV defined as:

$$\Delta RMSD(x) \equiv RMSD(x, PTS) - RMSD(x, PR) \tag{1}$$

where the $RMSD$ values are computed after optimal translation/rotation and with respect to the corresponding crystal structures. For the kink of the Relay helix, $\Delta RMSD$ was computed on atoms C, CA, O, N (backbone heavy atoms) of residues 485 to 493. This CV takes values $\approx 1.4\,\text{Å}$ in PR (straight Relay helix) and $\approx -1.4\,\text{Å}$ in PTS/PPS (kinked Relay helix), Fig 1e. Although $\Delta RMSD$ is defined using the PTS crystal structure as a reference, we note that the local backbone configuration of the kinked Relay helix is virtually indistinguishable between PTS and PPS ($0.32\,\text{Å}$ RMSD). For all intents and purposes, using a CV based on the PPS crystal structure [7] would have given identical results. To analyze the backbone hydrogen-bonding pattern of the Relay helix with a higher resolution than offered by $\Delta RMSD$, we introduced a set of distances k_j (see Table 1 for details). And, we computed dihedral angles χ_{11} and χ_{12} (see Table 1) to describe side-chain rotameric transitions associated with the formation of the Relay helix kink, as pointed out previously by Fischer et al. and independently by Baumketner [14, 24].

ATPase site

The opening state of the Switch II loop in the active site was characterized by
distances d_1 and d_γ as defined previously [26]. d_1 describes the critical salt-bridge
between Switch I (R205) and Switch II (E461). d_γ describes the Switch II-ATP
hydrogen bond. When both interactions are formed, Switch II is closed and the motor
is considered ATPase-competent.

Extended Adaptive Biasing Force (eABF) calculations

Two-dimensional eABF calculations were run along $X_c \times \Delta RMSD$ using NAMD 2.13
with the *colvars* module. We prioritized X_c for the eABF calculation as it is the
converter component which exhibits the largest variation from PR to PTS, with
typical values of $5\,\text{Å}$ in PR and $-10\,\text{Å}$ in PTS/PPS. Briefly, ABF records the average
generalized force applied by the system onto the collective variables, then applies an
exactly opposite bias to enhance the sampling by flattening the free energy landscape.
The free energy profile is recovered by numerical integration of the free energy gradient
estimate, which is the opposite of the collected average force. In extended ABF, an
extended degree of freedom is harmonically coupled to the collective variable [38]. The
generalized thermodynamic force felt by the extended degree of freedom is the
harmonic coupling force, which can be easily evaluated. eABF then applies the
standard ABF dynamics on the extended degree of freedom. eABF does not require
collective variable component orthogonality, which allows us to use a $\Delta RMSD$ CV.
Moreover, it favors smooth convergence of the biasing force estimate [38]. We used the
corrected-z-averaged restraint (CZAR) estimator to deconvolve the free energy
gradient estimate [38] and the *abf_integrate* tool to integrate it. eABF calculations
were run with SHAKE active and a $2\,\text{fs}$ timestep. The X_c collective variable was
coupled to its extended degree of freedom using a $10\,\text{kcal/mol/Å}^2$ harmonic force
constant; a $125\,\text{kcal/mol/Å}^2$ was used for $\Delta RMSD$. Force constants were determined
by trial-and-error as the lowest force constant sufficient to elicit the corresponding
conformational change in short Steered Molecular Dynamics simulations (data not
shown). This ensures a coupling that is both strong enough to bias the dynamics, and
soft enough to promote smooth exploration of the free energy surface [38]. Both

Collective Variable	Rearrangement	Biased in	Force constant (kcal/mol/U^2)
Orthogonal projection X_c	Converter swing	eABF, string (2d, 12d, 20d)	10
Orthogonal projection Y_c	Converter swing	string (12d, 20d)	10
Orthogonal projection Z_c	Converter swing	string (12d, 20d)	10
Distance i_1 (65O:707OG)	Converter/main body interaction	string (20d)	20
Distance i_2 (66CE:F763 phenyl ring)	Converter/main body interaction	string (20d)	20
Distance i_3 (65O:702NE2)	Converter/main body interaction	string (20d)	20
Distance i_4 (66O:702NE2)	Converter/main body interaction	string (20d)	20
Distance i_5 (64O:761N)	Converter/main body interaction	string (20d)	20
Orientation angle θ_{RH}	Orientation of the Relay helix	string (12d, 20d)	5
Orientation angle θ_{SH1}	Orientation of the SH1 helix	string (12d, 20d)	5
Distance $d_{R/SH1}$ (469-482CA:693-703CA)	Relay/SH1 seclusion	string (12d, 20d)	20
Orthogonal projection Z_{SH1}	SH1 piston motion	None	N.A.
Relay helix $\Delta RMSD$	Relay helix kink	eABF, string (2d)	125
Distance k_1 (486O:490N)	Relay helix kink	string (12d, 20d)	50
Distance k_2 (485O:489N)	Relay helix kink	string (12d, 20d)	50
Distance k_3 (485O:490N)	Relay helix kink	string (12d, 20d)	50
Distance k_4 (486O:491N)	Relay helix kink	string (12d, 20d)	50
Distance k_5 (490O:494N)	Relay helix secondary kink	string (20d)	50
Distance k_6 (491O:495N)	Relay helix secondary kink	string (20d)	50
Distance k_7 (492O:496N)	Relay helix secondary kink	string (20d)	50
Dihedral χ_{11} (L489 N:CA:CB:CG)	Hydrophobic switch	string (12d, 20d)	0.01
Dihedral χ_{12} (L700 N:CA:CB:CG)	Hydrophobic switch	string (12d, 20d)	0.01
Distance d_1 (R205CZ:E461CD)	Critical salt-bridge	None	N.A.
Distance d_γ (G459N:ATPO1G)	Switch II-ATP hydrogen bond	None	N.A.

Table 1. List of collective variables defined in the study of the myosin VI recovery stroke. All distances, orthogonal projections and RMSD are in Å. All angles are in deg.

extended degrees of freedom were thermostatted to 300 K with a Langevin thermostat ₅₀₇

with a $10\,\text{ps}^{-1}$ friction constant. The $X_c \times \Delta RMSD$ configurational space was ₅₀₈

discretized into a $0.75\,\text{Å} \times 0.1\,\text{Å}$ grid for the calculation. The *fullSamples* parameter ₅₀₉

was set to 2000 to limit non-equilibrium effects [54]. ₅₁₀

Similar to our previous work [26, 55], a two-step strategy was used to promote ₅₁₁

convergence of the eABF calculations. First, a 700 ns exploratory eABF run was ₅₁₂

performed starting from the equilibrated PR model, yielding near complete sampling ₅₁₃

of the configurational space. Second, the configurational space was divided into 12 ₅₁₄

non-overlapping windows separated by harmonic walls. Starting configurations for ₅₁₅

each window were extracted from the previous 700 ns run; each window was then ₅₁₆

simulated with eABF (starting with the gradient estimate obtained in the exploratory ₅₁₇

run) for 500 ns for a total of 6.7 μs of eABF simulation. This improves the sampling ₅₁₈

and refines the gradient estimate. Convergence was assessed as described in SI. ₅₁₉

String Method in Collective Variables (CVSM) calculations ₅₂₀

String method optimizations [31] with the "swarms-of-trajectories" variant [32] were ₅₂₁

used to compute MFEPs connecting the PR and PTS states. Strings were defined in ₅₂₂

either a 2-,12-, or 20-dimensional CV space designed to capture the essential features ₅₂₃

of the transition while remaining much simpler than full Cartesian space. The ₅₂₄

supporting CVs are listed in Table 1 along with their harmonic coupling force constant. ₅₂₅

The CVSM relaxes a discretized path in CV-space following the average drift ₅₂₆

computed from swarms of short MD simulations; up to unimportant dynamical effects, ₅₂₇

this average drift occurs along the direction of highest free energy gradient [32, 56, 57]. ₅₂₈

After each move, the string is reparametrized by enforcing equal spacing of the images, ₅₂₉

which projects out image displacement colinear to the string and ensures that string ₅₃₀

evolution takes place only along the orthogonal direction of the free energy gradient. ₅₃₁

Convergence is reached when orthogonal displacement is zero, that is, when the only ₅₃₂

image displacement is colinear to the string and thus, exactly cancelled by ₅₃₃

reparametrization. The converged string then represents the MFEP between the ₅₃₄

end-states in the supporting CV-space [31]. String calculations were performed using ₅₃₅

the Tcl scripting interface in NAMD 2.13 and the *colvars* module. Each CV was ₅₃₆

normalized between 0 and 1 by its total variation to ensure balanced ₅₃₇ reparametrization. Additional details on the string optimization procedure, ₅₃₈ parameters, and convergence analyses are reported in SI. Averaged, reparametrized ₅₃₉ strings over the last 50 iterations were considered for analysis. ₅₄₀

Umbrella Sampling along the converged string

To gain insight into the free energy changes along the converged string A1, we performed Umbrella sampling simulations. First, we used B-spline interpolation to increase the resolution of the converged string from 32 to 128 equally-spaced images in normalized CV-space. Then, each image was simulated for 120 ns under harmonic restraints applied to the same set of CVs and force constants as used for string optimization, except that χ_{11} and χ_{12} were excluded from the set of supporting CVs. Umbrella sampling simulations were run with Gromacs 2021.5 patched with *colvars* [52]. The PMF along the string was computed using Umbrella Integration (UI) [39] with the chain rule as described in [55], see SI. 1- and 2-dimensional PMFs along arbitrary CVs were computed using the Multistate Bennett Acceptance Ratio (MBAR) method [40] with Gaussian kernel density estimation, as implemented in *pymbar* version 4.0 [41].

Structural clustering

Structural clustering was performed using agglomerative hierarchical clustering with scikit-learn [58]. We used the RMSD over the CA atoms of the Relay and SH1 helices (after optimal translational and rotational fit) as the distance metric. Numbers of clusters were chosen to ensure that at least 2 or 3 clusters with appreciable occupancy (>10 %) were obtained.

Acknowledgments

This project was primarily supported by PRACE allocation ra4332, which allowed priority access to the Curie and Irene supercomputers. Additionally, this work was granted access to the High Performance Computing (HPC) resources of Centre de Calcul Recherche et Technologie (CCRT)/Centre Informatique National de

l'Enseignement Supérieur (CINES) under Allocation 2017-[076644] made by Grand Equipement National de Calcul Intensif, and also benefited from computational resources provided by the Mésocentre de Calcul de l'Université de Strasbourg and the Myria supercomputer of the Centre Régional Informatique et d'Applications Numériques de Normandie (CRIANN). The A.H. and M.C. teams were jointly supported by the Fondation pour la Recherche Médicale (Grant DBI20141231319). M.C. was supported by the Agence Nationale de la Recherche (ANR) through the LabEx Chemistry of Complex Systems (Project CSC-MCE-13), and the International Center for Frontier Research in Chemistry. A.H. was supported by a grant from the Association Française Contre les Myopathies 17235. The A.H. team is part of LabEx CelTisPhyBio:11-LBX-0038, which is part of the Initiative d'Excellence Paris Sciences et Lettres (Grant ANR-10-IDEX-0001-02 PSL).

Supporting information

Supplementary Text 1: Convergence and error analysis for eABF calculations

As seen on Supplementary Figure 6, all relevant regions of configurational space are visited multiple times by the eABF simulations. Convergence of eABF calculations was assessed by monitoring the per-window RMSD of the free energy gradient estimate (Supplementary Figure 7). Most windows representative of the transition tube ($W_{11}, W_{21}, W_{22}, W_{32}, W_{33}, W_{43}$) show a plateau close to $0.5\,\mathrm{kcal\,mol^{-1}\,\mathring{A}^{-1}}$ RMSD over the last few tens of ns, suggesting that the generalized force estimate has converged. Bootstrap-like statistical error analysis was performed as in our previous work [26, 59] and shows that statistical error on the free energy values is small ($<$ $1.0\,\mathrm{kcal\,mol^{-1}}$) everywhere in the relevant regions of 2D free energy landscape, and does not go above $2.0\,\mathrm{kcal\,mol^{-1}}$ even at the irrelevant margins (Supplementary Figure 8). The statistical error at the first free energy barrier is of order $1.0\,\mathrm{kcal\,mol^{-1}}$.

Supplementary Text 2: String method calculations

Procedure

An iteration of the string method with swarms of trajectories consists of the following operations, performed for each image:

1. Equilibration of length t_{eq} of the system around its image along the string in CV-space using harmonic restraining potentials.

2. Parallel, unrestrained simulation of n_{swarm} replicas for t_{free} initiated from the equilibrated system

For each image, the average drift in CV-space measured over the swarm of unrestrained trajectories is used to update the string. Then, the string is smoothed by local averaging, and reparametrized [31, 32]. This process is iterated until convergence is reached. Convergence of string optimizations is assessed by monitoring the Root Mean Square Deviation (RMSD) of the string in normalized CV-space with respect to

Fig 6. Sampling of the configurational space in eABF simulations. The number of counts, *i.e.*, the number of times a given grid point has been visited, is shown in logarithmic scale. Black dotted lines indicate the window boundaries in stratified simulation. Window labels are shown.

Fig 7. Convergence of generalized force estimate in eABF simulations. Shown is the RMSD of the gradient estimate for each window during the stratified run. X-axis, time of stratified simulation in ns; Y-axis, gradient RMSD, in $\mathrm{kcal\,mol^{-1}\,\mathring{A}^{-1}}$. The red dotted lines show the $0.5\,\mathrm{kcal\,mol^{-1}\,\mathring{A}^{-1}}$ cut-off which is used to evaluate convergence.

Fig 8. Statistical error estimate for the free energy landscape from bootstrap-like analysis (in $\mathrm{kcal\,mol^{-1}}$).

Fig 9. eABF calculations sample open Switch II conformers consistent with independent PTS simulations. (a) Distances d_1 and d_γ never sample the closed Switch II region in stratified eABF simulations. Shown are d_1, d_γ scatter plots colored per window. (b) The PTS basin identified in eABF calculation is consistent with independent, unbiased MD simulations of Myo6. Shown are density lines for X_c and $\Delta RMSD$ from the Myo6 simulations we reported in [26]. Of note are: the remarkable agreement of the PTS unbiased simulations with the free energy landscape; the sampling of a converter movement in simulation PR+ATP (3) which is roughly consistent with metastable state I_A; and the clear separation of PTS from PPS.

Fig 10. Statistical distribution of selected observables in substates identified in eABF calculations. (a) Distribution of converter coordinates X_c and Y_c. (b) Distribution of Relay helix backbone hydrogen bonds k_3 and k_5. The difference in converter position between I_A and I_B, along with the atypical hydrogen bonding pattern of the Relay helix in I_B, are consistent with I_A and I_B not belonging to the same transition pathway.

Fig 11. The same I_A intermediate is captured in eABF and string/US calculations. (a) RMSD of structures sampled along String A1 with respect to the representative structure of I_A sampled in eABF, computed after optimal fit on the CA atoms of the Relay and SH1 helices. The RMSD clearly is minimum and below 1 Å when the progress along the path is roughly between 0.24 and 0.35, which corresponds to the I_A basin as identified in string/US calculations. The red dot marks the minimal RMSD value, achieved at image 36 along the string. (b) Structural comparison of the Relay helix, SH1 helix and converter between the representative structure of state I_A sampled in eABF (cyan) and in string/US (image 36, red). The conformation and orientation of the structural elements are extremely close. These results demonstrate that the same metastable basin is sampled in both independent calculations, which justifies referring to them as I_A regardless of their provenance.

the initial string. The string is considered converged when this value stabilizes. String optimization is akin to a gradient-based high-dimensional optimization, for which a well-known issue is the existence of multiple locally optimal solutions. A careful choice of the initial "guess" path from which string iterations are initialized mitigates this issue by making it more likely to converge to a biologically meaningful solution. Comparing converged strings from different guess path initializations also allows one to assess the robustness of the findings. For these reasons, we consider several different guess paths in our string calculations.

String optimizations in $(X_c, \Delta RMSD)$ space are consistent with eABF results

To validate the transition tube predicted by eABF, we first performed string optimizations in $(X_c, \Delta RMSD)$-space.

Stability of the eABF-predicted MFEP under explicit CVSM dynamics

First, we sought to assess whether the MFEP computed analytically over the eABF free energy landscape were stable under explicit (*i.e.*, atomistic) CVSM dynamics. We

Calculation	Guess path	Ends	t_{eq} (ps)	n_{swarm}	t_{free} (ps)	n_{iter}
2D String (1)	eABF-MFEP	Fixed	1	10	0.5	201
2D String (2)	eABF-MFEP	Fixed	1	10	0.5	116
2D String (3)	Straight	Free	1	10	0.5	127
2D String (4)	Straight	Fixed	1	10	1	119

Table 2. CVSM simulations in 2D CV space

constructed this path (hereafter eABF-MFEP) by 1) evaluating analytical derivatives   619
of the eABF free energy landscape and 2) relaxing a 2D string on this free energy   620
landscape using the zero-temperature string method [60] in normalized coordinates.   621
These operations were implemented in a Python/Scipy script [61, 62]. Then, we   622
prepared equilibrated Myo6 conformers along this path in $(X_c, \Delta RMSD)$-space as   623
follows. We "pulled" Myo6 (from the equilibrated PR model) along the   624
eABF-predicted MFEP using a 120 ns-long Steered MD simulation on X_c and   625
$\Delta RMSD$ with the harmonic force constants reported in Table 1. 32 conformers   626
equally-spaced in normalized CV-space were extracted from the Steered MD trajectory   627
and further equilibrated by 20 ns of umbrella sampling with the same force constants.   628
The resulting conformers were then used to run 2 separate CVSM calculations in 2D   629
space (Supplementary Table 2). In both cases, the path obtained by averaging the   630
strings over the last 50 iterations is remarkably close to the eABF-predicted MFEP.   631
Therefore, up to equilibrium fluctuations, the 2D transition path predicted by eABF   632
calculations is essentially invariant under the string method dynamics, independently   633
confirming that it lies close to the MFEP (Supplementary Figures 12, 13).   634

Relaxation of the straight guess path towards the eABF-predicted MFEP   635
under explicit CVSM dynamics   636

Using a similar protocol, we prepared equilibrated Myo6 conformers equally-spaced   637
along the straight path joining PR and PTS in 2D CV-space. Relaxation of this   638
straight guess path by two independent CVSM runs shows that it consistently evolves   639
towards the eABF-predicted transition tube (Supplementary Figure 12,   640
Supplementary Table 2). We conclude that independent string optimizations in 2D   641
space confirm the important features of the transition mechanism predicted by eABF.   642

Fig 12. CVSM calculations in 2D CV-space are consistent with the eABF free energy landscape.

Fig 13. Convergence of CVSM calculations in 2D CV-space. Shown is the RMSD in normalized CVs from the initial path. The greyed area shows the last 50 iterations, over which the average strings pictured in Supplementary Figure 12 are computed.

String method calculations in 12D normalized CV-space

Generation of a 12-dimensional guess path from the 2D eABF-MFEP by uplifting

The eABF-MFEP is representative of the general features of the PR → PTS transition, but the 2D CV-space is too coarse-grained to capture the finer details of the mechanism. The 2D MFEP and surrounding transition tube correspond to projections of a set of transition pathways lying in a space of higher dimensionality. Refined approximations of the true MFEP could be obtained by backmapping, or "uplifting", the 2D MFEP into a higher-dimensional space provided that the new coordinates offer a better descriptive power of the transition mechanism, that is, if they capture relevant aspects of the transitions overlooked by the coarser description. We developed a method to uplift the 2D eABF-MFEP by embedding it into a 12D CV-space designed to provide a higher-resolution representation of the recovery stroke by 1) including additional degrees of freedom and 2) replacing the coarse-grained $\Delta RMSD$ Relay helix kink descriptor by 4 separate distance CVs representing local backbone rearrangements upon the formation of the kink. This uplifted string provides the starting point for a new series of CVSM calculations, this time in 12D space.

Such an uplifting requires accounting for the missing information of the additional degrees of freedom. Here, we used the average values of the additional CVs measured from the pre-equilibration Umbrella Sampling simulations along the 2D eABF-MFEP reported above. These provide reasonable initial estimates of the most likely value taken by the additional CVs when the system is restrained in the vicinity of the MFEP. However, the resulting path in 12D CV-space was found to be very rugged, which we reasoned would impede convergence of the string calculations. To alleviate this issue, we regularized the uplifted path by averaging it with a straight path connecting PR to PTS in normalized 12-dimensional CV space. The resulting uplifted regularized path retains the global shape of the uplifted path before regularization, but is much smoother.

Calculation	Guess path	Ends	t_{eq} (ps)	n_{swarm}	t_{free} (ps)	n_{iter}
12D String A1	uplifted/regularized eABF-MFEP	Fixed	100	20	0.5	179
12D String A2	uplifted/regularized eABF-MFEP	Fixed	100	20	0.5	119
12D String B1	Straight (12D)	Fixed	100	20	0.5	199
12D String B2	Straight (12D)	Fixed	100	20	0.5	119

Table 3. CVSM simulations in 12D CV space

Fig 14. Convergence of CVSM calculations in 12D CV-space. Shown is the RMSD in normalized CVs from the initial path. The greyed area shows the last 50 iterations, over which the average strings are computed.

CVSM calculations from the regularized uplifted string 671

The 32 Myo6 conformers previously pre-equilibrated in 2D along the eABF-MFEP 672
were further relaxed along the regularized uplifted string in 12D by 1 ns of US. Then, 673
they were used to perform two independent CVSM calculations to reveal the MFEP in 674
12D CV-space, yielding Strings A1 and A2. 675

12D CVSM calculations from the straight guess path 676

To assess the robustness of the string calculations, we also ran independent string 677
calculations initiated from the linear guess path connecting PR to PTS in normalized 678
12-dimensional CV-space. 32 equally-spaced conformers along this path were prepared 679
by relaxing the conformers previously pre-equilibrated along the 2D linear path to 680
their corresponding images in the 12D linear string using 1 ns of US. Then, there were 681
used to perform two separate CVSM calculations, yielding Strings B1 and B2. The 682
corresponding transition path, or Path B, is described in Supplementary Text 4. 683

Fig 15. Projections of the 12D converged strings onto 2D space reveal two alternate pathways lying within the transition tube predicted by eABF.

Fig 16. Evolution of biased structural observables along converged, averaged Strings A1 and A2.

Fig 17. Contacts between the N-terminal subdomain and the converter during the PR→ PTS transition. (a-d) Close-up on the converter/N-ter interface in representative frames of the metastable states sampled in umbrella sampling along String A1. Residues involved in contacts are shown as sticks. For clarity, non-polar hydrogens are not shown. (e) PMF along X_c and the number of contacts estimated with MBAR from umbrella sampling along string A1. A contact is defined as two heavy atoms being closer than 4.5 Å.

Calculation	Guess path	Ends	t_{eq} (ps)	n_{swarm}	t_{free} (ps)	n_{iter}
20D String (1)	Averaged converged string from 12D String (3) uplifted/regularized to 20D space	Fixed	1	10	0.5	20

Table 4. CVSM simulation in 20D CV space

String method calculations in 20D normalized CV-space

The analysis of the transition pathway emerging from the 12D string optimizations initialized with the straight guess path (*i.e.*, Strings B1 and B2) revealed 8 additional potentially relevant degrees of freedom, see Supplementary Text 4. To better reflect their involvement in the mechanism, we uplifted and regularized the converged String B1 to the 20-dimensional CV-space using the procedure described above. The 20D regularized/uplifted string was then relaxed to convergence by 39 iterations of CVSM.

Projection of the high-dimensional strings onto 2D $(X_c,\ \Delta RMSD)$-space

We projected the high-dimensional (12 or 20 dimensions) strings onto 2D $(X_c,\ \Delta RMSD)$-space to compare them with the eABF results. Since the $\Delta RMSD$

Fig 18. Evolution of biased structural observables along converged, averaged Strings B1 and B2.

CV is not a component of the high-dimensional strings, we evaluated its value as the per-image average over the last 50 string iterations. Since $\Delta RMSD$ is not explicitly biased nor involved in string reparametrization, the projected strings have a rugged aspect.

Fig 19. A secondary kink in the Relay helix forms in Path B. a. Initial state (PR-like): the Relay helix is straight and intact, the converter interacts with the N-terminal subdomain. c. $\alpha = 0.25$: a secondary kink in the Relay helix (red arrow) accomodates converter movement. The converter is still in contact with the N-terminal subdomain. c. Final state (PTS-like): the converter has moved and broken the contacts, and the canonical kink (red arrow) in the Relay helix has formed.

Fig 20. Re-optimization of String B1 in 20-dimensional space. (a) Convergence of the CVSM calculation in 20D CV-space. (b-f) Evolution of converter/N-terminal distances along the converged, averaged 20d string. (g-i) Evolution of the RH backbone distances describing the secondary kink along the converged, averaged 20D string. Panels b and c show that contacts between the converter and the N-terminal subdomain are preserved until about halfway along the transition, while panels g-i show that a secondary kink forms in the RH from about 0.25 to 0.50 progress along the string.

Supplementary Text 3: free energy profiles along the string from umbrella sampling

Umbrella sampling

An "on-the-path" Umbrella Sampling calculation was performed along the averaged string from the first string optimization initiated from the eABF-regularized guess, or String A1. First, χ_{11} and χ_{12} were excluded from the set of biased CVs because we reasoned that the sharp nature of rotameric transitions could hinder sampling around the transition state regions. Then, we fitted a collection of B-splines on each CV along the string using Scipy. Finally, 128 images equally-spaced along the string in normalized CV space were extracted and used as restraint centers for umbrella sampling. Umbrella sampling simulations were run with GROMACS 2021.5 patched with *colvars*, using the same force constants as for string optimizations, and were initiated from structures sampled in the last iteration of the string optimization. To equilibrate the images with the restraining potentials, the first 1 ns were discarded for each window. Each window was then simulated for 120 ns, resulting in a total of 15.36 μs of umbrella sampling simulation. For analysis, CV data were saved every 10 ps. Free energies along the string were then computed using two different approaches.

Free energy profile along the string by umbrella integration

To obtain a PMF along the string, we used the procedure described in [55]. Introducing α as a non-dimensional progress variable along the path, such that $\alpha = 0$ at the beginning of the path and $\alpha = 1$ at the end, one can write for the free energy $F(\alpha)$ along the path:

$$\frac{\mathrm{d}F(\alpha)}{\mathrm{d}\alpha} = \sum_i \frac{\partial F}{\partial \xi_i} \cdot \frac{\mathrm{d}\xi_i}{\mathrm{d}\alpha} \tag{2}$$

where ξ_i refers to CV i along the string, and the sum runs over all supporting CVs. The $\frac{\mathrm{d}\xi_i}{\mathrm{d}\alpha}$ are evaluated by analytical differentiation of the fitted splines. To estimate the free energy gradients with respect to CV components, we used the Umbrella Integration estimator [39] with a number of bins equal to the number of windows, *i.e.*

128.

MBAR reweighting 726

Separately from the UI calculations, free energy profiles along arbitrary CVs (possibly 727
including unrestrained ones) were computed using the Multistate Bennett Acceptance 728
Ratio (MBAR) method [40]. These PMFs represent the effective free energy landscapes 729
when the progress coordinate along the PR $\rightarrow$ PTS transition is averaged out, and 730
provide insight into the thermodynamic coupling between elementary rearrangements. 731
MBAR calculations were performed using *pymbar* 4.0.1. We used the recently 732
published Kernel Density Estimation (KDE)-based procedure to obtain smooth free 733
energy estimates [41]. In so doing, we found that the choice of the KDE bandwidth 734
parameter could strongly affect the value of the free energy barriers, but only 735
marginally affected the position and relative stability of free energy basins. Therefore, 736
the MBAR-computed free energy maps should be considered semi-quantitative. 737

Supplementary Text 4: Mechanism along Path B 738

From the straight guess path, two independent string optimizations were performed 739
until they converged towards a common sequence of events, which we name Path B 740
(Supplementary Figure 14). Similar to the situation for Path A, Path B from string 741
calculations is virtually identical to the Path B_{ABF}, as we justify below. Projection of 742
the 2 Path B 12D-strings onto the (X_c, $\Delta RMSD$) plane shows that they are 743
consistent with the shape and location of the transition tube (Supplementary Figure 744
15). Interestingly, Paths A and B do not overlap, and Path B visits I$_B$. Thus, even if 745
they share the same gross features, the structural mechanism predicted from Path B is 746
a bit different from that of Path A (Supplementary Figure 18). In Path B, the kink in 747
the RH forms with a different timing and mechanism. There, the initial movement of 748
the converter is not accomodated by the disruption of its interactions with the 749
N-terminal subdomain, but by the transient formation of a secondary kink in the RH 750
at the level of residues 490 to 496 (Supplementary Figure 19). This is similar to Path 751
B_{ABF} from eABF calculations (see notably Fig. 2e) and supports the idea that Paths 752
B_{ABF} and B are in fact representative of the same mechanism. As the 753

hydrogen-bonding distances describing the formation of the secondary kink are not   754
explicitly biased in the 12D string optimizations, it is difficult to know whether the   755
mechanism observed in Path B does correspond to an MFEP also along these degrees   756
of freedom. To clarify this matter, we extended the CV-space to 20 dimensions by   757
adding 3 distances describing the secondary kink and 5 distances describing   758
converter/N-ter interactions, uplifted the first converged Path B string to this space,   759
and relaxed it with the string method (Supplementary Text 2, Supplementary Figure   760
20a). The resulting pathway still exhibits the same sequence of events, supporting its   761
relevance as a locally optimal MFEP in the PR $\rightarrow$ PTS transition.   762

The extended, highly bent configuration of the Relay helix enables the rotation of   763
the converter while some of its contacts with the N-terminal subdomain are preserved   764
(Supplementary Fig. 20b-i, Supplementary Fig. 19). This configuration corresponds to   765
the I_B basin identified in eABF, which may represent a metastable intermediate state   766
along this path. Next, upon going from I_B to PTS1, this secondary kink disappears   767
and the canonical kink in the RH forms concurrently with continued movement of the   768
converter in a fashion similar to Path A.   769

The sensitivity to the initial guess of the converged path is not overly surprising, as   770
multiple pathways are expected for a system the size of myosin's even after reducing   771
its dimensionality to 12 (or 20). It is plausible that both families of pathways may   772
actually be explored in the recovery stroke of Myo6. Thus, the Relay helix exhibits   773
significant plasticity throughout its length, which may allow it to bend significantly   774
and even develop transient kinks to accomodate the extensive movement of the   775
converter through a diversity of mechanisms. Nevertheless, Path A is likely to be more   776
representative of the dominant mechanism because it entails less structural   777
perturbation (*i.e.,* only one kink forms), which arguably is a desirable property for an   778
MFEP.   779

References

1. Sweeney HL, Houdusse A. Structural and Functional Insights into the Myosin
   Motor Mechanism. Annual Review of Biophysics. 2010;39(1):539–557.
   doi:10.1146/annurev.biophys.050708.133751.

2. Geeves MA, Holmes KC. Structural Mechanism of Muscle Contraction. Annual Review of Biochemistry. 1999;68(1):687–728. doi:10.1146/annurev.biochem.68.1.687.

3. Robert-Paganin J, Pylypenko O, Kikuti C, Sweeney HL, Houdusse A. Force Generation by Myosin Motors: A Structural Perspective. Chemical Reviews. 2020;120(1):5–35. doi:10.1021/acs.chemrev.9b00264.

4. Sweeney HL, Houdusse A. What Can Myosin VI Do in Cells? Current Opinion in Cell Biology. 2007;19(1):57–66. doi:10.1016/j.ceb.2006.12.005.

5. de Jonge JJ, Batters C, O'Loughlin T, Arden SD, Buss F. The MYO6 Interactome: Selective Motor-Cargo Complexes for Diverse Cellular Processes. FEBS Letters. 2019;593(13):1494–1507. doi:10.1002/1873-3468.13486.

6. Ménétrey J, Bahloul A, Wells AL, Yengo CM, Morris CA, Sweeney HL, et al. The Structure of the Myosin VI Motor Reveals the Mechanism of Directionality Reversal. Nature. 2005;435(7043):779–785. doi:10.1038/nature03592.

7. Ménétrey J, Llinas P, Mukherjea M, Sweeney HL, Houdusse A. The Structural Basis for the Large Powerstroke of Myosin VI. Cell. 2007;131(2):300–308. doi:10.1016/j.cell.2007.08.027.

8. Ménétrey J, Llinas P, Cicolari J, Squires G, Liu X, Li A, et al. The Post-Rigor Structure of Myosin VI and Implications for the Recovery Stroke. The EMBO Journal. 2008;27(1):244–252. doi:10.1038/sj.emboj.7601937.

9. Fisher AJ, Smith CA, Thoden J, Smith R, Sutoh K, Holden HM, et al. X-Ray Structures of the Myosin Motor Domain of Dictyostelium Discoideum Complexed with MgADP.BeFx and MgADP.AlF4-. Biochemistry. 1995;34(28):8960–8972. doi:10.1021/bi00028a004.

10. Kiani FA, Fischer S. Catalytic Strategy Used by the Myosin Motor to Hydrolyze ATP. Proceedings of the National Academy of Sciences. 2014;111(29):E2947–E2956. doi:10.1073/pnas.1401862111.

11. Kiani FA, Fischer S. Stabilization of the ADP/Metaphosphate Intermediate during ATP Hydrolysis in Pre-power Stroke Myosin: QUANTITATIVE

ANATOMY OF AN ENZYME. Journal of Biological Chemistry. 2013;288(49):35569–35580. doi:10.1074/jbc.M113.500298.

12. Lu X, Ovchinnikov V, Demapan D, Roston D, Cui Q. Regulation and Plasticity of Catalysis in Enzymes: Insights from Analysis of Mechanochemical Coupling in Myosin. Biochemistry. 2017;56(10):1482–1497. doi:10.1021/acs.biochem.7b00016.

13. Geeves MA. Review: The ATPase Mechanism of Myosin and Actomyosin: The ATPase Mechanism of Myosin and Actomyosin. Biopolymers. 2016;105(8):483–491. doi:10.1002/bip.22853.

14. Fischer S, Windshügel B, Horak D, Holmes KC, Smith JC. Structural Mechanism of the Recovery Stroke in the Myosin Molecular Motor. Proceedings of the National Academy of Sciences of the United States of America. 2005;102(19):6873–6878. doi:10.1073/pnas.0408784102.

15. Koppole S, Smith JC, Fischer S. Simulations of the Myosin II Motor Reveal a Nucleotide-state Sensing Element That Controls the Recovery Stroke. Journal of Molecular Biology. 2006;361(3):604–616. doi:10.1016/j.jmb.2006.06.022.

16. Mesentean S, Koppole S, Smith JC, Fischer S. The Principal Motions Involved in the Coupling Mechanism of the Recovery Stroke of the Myosin Motor. Journal of Molecular Biology. 2007;367(2):591–602. doi:10.1016/j.jmb.2006.12.058.

17. Koppole S, Smith JC, Fischer S. The Structural Coupling between ATPase Activation and Recovery Stroke in the Myosin II Motor. Structure. 2007;15(7):825–837. doi:10.1016/j.str.2007.06.008.

18. Woo HJ. Exploration of the Conformational Space of Myosin Recovery Stroke via Molecular Dynamics. Biophysical Chemistry. 2007;125(1):127–137. doi:10.1016/j.bpc.2006.07.001.

19. Harris MJ, Woo HJ. Energetics of Subdomain Movements and Fluorescence Probe Solvation Environment Change in ATP-bound Myosin. European Biophysics Journal. 2008;38(1):1–12. doi:10.1007/s00249-008-0347-3.

20. Yu H, Ma L, Yang Y, Cui Q. Mechanochemical Coupling in the Myosin Motor Domain. I. Insights from Equilibrium Active-Site Simulations. PLoS Computational Biology. 2007;3(2):e21. doi:10.1371/journal.pcbi.0030021.

21. Yu H, Ma L, Yang Y, Cui Q. Mechanochemical Coupling in the Myosin Motor Domain. II. Analysis of Critical Residues. PLoS Computational Biology. 2007;3(2):e23. doi:10.1371/journal.pcbi.0030023.

22. Elber R, West A. Atomically Detailed Simulation of the Recovery Stroke in Myosin by Milestoning. Proceedings of the National Academy of Sciences. 2010;107(11):5001–5005. doi:10.1073/pnas.0909636107.

23. Baumketner A, Nesmelov Y. Early Stages of the Recovery Stroke in Myosin II Studied by Molecular Dynamics Simulations. Protein Science. 2011;20(12):2013–2022. doi:10.1002/pro.737.

24. Baumketner A. Interactions between Relay Helix and Src Homology 1 (SH1) Domain Helix Drive the Converter Domain Rotation during the Recovery Stroke of Myosin II. Proteins: Structure, Function, and Bioinformatics. 2012;80(6):1569–1581. doi:10.1002/prot.24051.

25. Baumketner A. The Mechanism of the Converter Domain Rotation in the Recovery Stroke of Myosin Motor Protein. Proteins: Structure, Function, and Bioinformatics. 2012;80(12):2701–2710. doi:10.1002/prot.24155.

26. Blanc F, Isabet T, Benisty H, Sweeney HL, Cecchini M, Houdusse A. An Intermediate Along the Recovery Stroke of Myosin VI Revealed by X-ray Crystallography and Molecular Dynamics. Proceedings of the National Academy of Sciences. 2018; p. 201711512. doi:10.1073/pnas.1711512115.

27. Torrie GM, Valleau JP. Nonphysical Sampling Distributions in Monte Carlo Free-Energy Estimation: Umbrella Sampling. Journal of Computational Physics. 1977;23(2):187–199. doi:10.1016/0021-9991(77)90121-8.

28. Laio A, Parrinello M. Escaping Free-Energy Minima. Proceedings of the National Academy of Sciences. 2002;99(20):12562–12566. doi:10.1073/pnas.202427399.

29. Comer J, Gumbart JC, Hénin J, Lelièvre T, Pohorille A, Chipot C. The Adaptive Biasing Force Method: Everything You Always Wanted To Know but Were Afraid To Ask. The Journal of Physical Chemistry B. 2015;119(3):1129–1151. doi:10.1021/jp506633n.

30. Chipot C, Pohorille A, editors. Free Energy Calculations: Theory and Applications in Chemistry and Biology. No. 86 in Springer Series in Chemical Physics. Berlin ; New York: Springer; 2007.

31. Maragliano L, Fischer A, Vanden-Eijnden E, Ciccotti G. String Method in Collective Variables: Minimum Free Energy Paths and Isocommittor Surfaces. The Journal of Chemical Physics. 2006;125(2):024106. doi:10.1063/1.2212942.

32. Pan AC, Sezer D, Roux B. Finding Transition Pathways Using the String Method with Swarms of Trajectories. The Journal of Physical Chemistry B. 2008;112(11):3432–3440. doi:10.1021/jp0777059.

33. Ma W, Schulten K. Mechanism of Substrate Translocation by a Ring-Shaped ATPase Motor at Millisecond Resolution. Journal of the American Chemical Society. 2015;137(8):3031–3040. doi:10.1021/ja512605w.

34. Lev B, Murail S, Poitevin F, Cromer BA, Baaden M, Delarue M, et al. String Method Solution of the Gating Pathways for a Pentameric Ligand-Gated Ion Channel. Proceedings of the National Academy of Sciences. 2017;114(21):E4158–E4167. doi:10.1073/pnas.1617567114.

35. Singharoy A, Chipot C, Moradi M, Schulten K. Chemomechanical Coupling in Hexameric Protein–Protein Interfaces Harnesses Energy within V-Type ATPases. Journal of the American Chemical Society. 2017;139(1):293–310. doi:10.1021/jacs.6b10744.

36. Ovchinnikov V, Karplus M, Vanden-Eijnden E. Free Energy of Conformational Transition Paths in Biomolecules: The String Method and Its Application to Myosin VI. The Journal of Chemical Physics. 2011;134(8):085103. doi:10.1063/1.3544209.

37. Ovchinnikov V, Cecchini M, Vanden-Eijnden E, Karplus M. A Conformational Transition in the Myosin VI Converter Contributes to the Variable Step Size. Biophysical Journal. 2011;101(10):2436–2444. doi:10.1016/j.bpj.2011.09.044.

38. Lesage A, Lelièvre T, Stoltz G, Hénin J. Smoothed Biasing Forces Yield Unbiased Free Energies with the Extended-System Adaptive Biasing Force Method. The Journal of Physical Chemistry B. 2017;121(15):3676–3685. doi:10.1021/acs.jpcb.6b10055.

39. Kästner J, Thiel W. Bridging the Gap Between Thermodynamic Integration and Umbrella Sampling Provides a Novel Analysis Method: "Umbrella Integration". The Journal of Chemical Physics. 2005;123(14):144104. doi:10.1063/1.2052648.

40. Shirts MR, Chodera JD. Statistically Optimal Analysis of Samples from Multiple Equilibrium States. The Journal of Chemical Physics. 2008;129(12):124105. doi:10.1063/1.2978177.

41. Shirts MR, Ferguson AL. Statistically Optimal Continuous Free Energy Surfaces from Biased Simulations and Multistate Reweighting. Journal of Chemical Theory and Computation. 2020;16(7):4107–4125. doi:10.1021/acs.jctc.0c00077.

42. Hill TL. Free Energy Transduction and Biochemical Cycle Kinetics. Mineola, N.Y: Dover Publications; 2005.

43. Sirigu S, Hartman JJ, Planelles-Herrero VJ, Ropars V, Clancy S, Wang X, et al. Highly Selective Inhibition of Myosin Motors Provides the Basis of Potential Therapeutic Application. Proceedings of the National Academy of Sciences. 2016; p. 201609342. doi:10.1073/pnas.1609342113.

44. Akter F, Ochala J, Fornili A. Binding Pocket Dynamics along the Recovery Stroke of Human β-Cardiac Myosin. PLOS Computational Biology. 2023;19(5):e1011099. doi:10.1371/journal.pcbi.1011099.

45. Yang Y, Yu H, Cui Q. Extensive Conformational Transitions Are Required to Turn On ATP Hydrolysis in Myosin. Journal of Molecular Biology. 2008;381(5):1407–1420. doi:10.1016/j.jmb.2008.06.071.

46. Baldo AP, Tardiff JC, Schwartz SD. Mechanochemical Function of Myosin II: Investigation into the Recovery Stroke and ATP Hydrolysis. The Journal of Physical Chemistry B. 2020;124(45):10014–10023. doi:10.1021/acs.jpcb.0c05762.

47. Chakraborti A, Baldo AP, Tardiff JC, Schwartz SD. Investigation of the Recovery Stroke and ATP Hydrolysis and Changes Caused Due to the Cardiomyopathic Point Mutations in Human Cardiac β Myosin. The Journal of Physical Chemistry B. 2021; p. acs.jpcb.1c03144. doi:10.1021/acs.jpcb.1c03144.

48. Porter JR, Meller A, Zimmerman MI, Greenberg MJ, Bowman GR. Conformational Distributions of Isolated Myosin Motor Domains Encode Their Mechanochemical Properties. eLife. 2020;9:e55132. doi:10.7554/eLife.55132.

49. Meller A, Lotthammer JM, Smith LG, Novak B, Lee LA, Kuhn CC, et al. Drug Specificity and Affinity Are Encoded in the Probability of Cryptic Pocket Opening in Myosin Motor Domains. eLife. 2023;12:e83602. doi:10.7554/eLife.83602.

50. Chakraborti A, Tardiff JC, Schwartz SD. Insights into the Mechanism of the Cardiac Drug Omecamtiv Mecarbil-A Computational Study. The Journal of Physical Chemistry B. 2022;doi:10.1021/acs.jpcb.2c06679.

51. Phillips JC, Braun R, Wang W, Gumbart J, Tajkhorshid E, Villa E, et al. Scalable Molecular Dynamics with NAMD. Journal of Computational Chemistry. 2005;26(16):1781–1802. doi:10.1002/jcc.20289.

52. Abraham MJ, Murtola T, Schulz R, Páll S, Smith JC, Hess B, et al. GROMACS: High Performance Molecular Simulations through Multi-Level Parallelism from Laptops to Supercomputers. SoftwareX. 2015;1–2:19–25. doi:10.1016/j.softx.2015.06.001.

53. Fiorin G, Klein ML, Hénin J. Using Collective Variables to Drive Molecular Dynamics Simulations. Molecular Physics. 2013;111(22-23):3345–3362. doi:10.1080/00268976.2013.813594.

54. Hénin J, Chipot C. Overcoming Free Energy Barriers Using Unconstrained Molecular Dynamics Simulations. The Journal of Chemical Physics. 2004;121(7):2904. doi:10.1063/1.1773132.

55. Blanc FE, Cecchini M. An Asymmetric Mechanism in a Symmetric Molecular Machine. The Journal of Physical Chemistry Letters. 2021;12(13):3260–3265. doi:10.1021/acs.jpclett.1c00404.

56. Johnson ME, Hummer G. Characterization of a Dynamic String Method for the Construction of Transition Pathways in Molecular Reactions. The Journal of Physical Chemistry B. 2012;116(29):8573–8583. doi:10.1021/jp212611k.

57. Maragliano L, Roux B, Vanden-Eijnden E. Comparison between Mean Forces and Swarms-of-Trajectories String Methods. Journal of Chemical Theory and Computation. 2014;10(2):524–533. doi:10.1021/ct400606c.

58. Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et al. Scikit-Learn: Machine Learning in Python. MACHINE LEARNING IN PYTHON. 2011; p. 6.

59. Wereszczynski J, McCammon JA. Nucleotide-Dependent Mechanism of Get3 as Elucidated from Free Energy Calculations. Proceedings of the National Academy of Sciences. 2012;109(20):7759–7764.

60. E W, Ren W, Vanden-Eijnden E. String Method for the Study of Rare Events. Physical Review B. 2002;66(5). doi:10.1103/PhysRevB.66.052301.

61. Jones E, Oliphant T, Peterson P. {SciPy}: Open Source Scientific Tools for {Python}. 2001–///;.

62. Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, et al. SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python. Nature Methods. 2020;17(3):261–272. doi:10.1038/s41592-019-0686-2.