Drivers of thermal tolerance breadth of plants across contrasting biomes

Verónica F. Briceño[1*], Pieter A. Arnold[1*^], Alicia M. Cook[2], Stephanie K. Courtney Jones[1,3],

Rachael V. Gallagher[4], Kris French[5], León A. Bravo[6], Adrienne B. Nicotra[1], Andy Leigh[2^]

[*] Joint lead authors

[1] Division of Ecology and Evolution, Research School of Biology, Australian National

University, Canberra, Australian Capital Territory, Australia.

[2] University of Technology Sydney, School of Life Sciences, Ultimo, New South Wales,

Australia

[3] Ecosystem and Threatened Species, Northwest Branch, NSW Department of Planning and

Environment, Dubbo, New South Wales, Australia

[4] Hawkesbury Institute for the Environment, Western Sydney University, Sydney, New South

Wales, Australia.

[5] Centre for Sustainable Ecosystem Solutions, School of Earth, Atmospheric and Life Sciences,

University of Wollongong, Wollongong, New South Wales, Australia.

[6] Departamento de Ciencias Agronómicas y Recursos Naturales, Facultad de Ciencias

Agropecuarias y Medioambiente, Universidad de la Frontera, Región de la Araucanía, Temuco,

Chile.

* Corresponding authors:

^ Pieter Arnold: pieter.arnold@anu.edu.au , Andy Leigh: andrea.leigh@uts.edu.au

Abstract

1. The climate variability hypothesis (CVH) predicts that species from environments with

more variable temperatures should have wider thermal tolerance breadth. This hypothesis

has not yet been tested thoroughly across diverse plants. Here, we asked how local

climate predictors (including precipitation, mean and extreme temperatures and thermal

variability) are associated with species physiological thermal limits.

2. Measures of lower ($T_{crit-cold}$) and upper ($T_{crit-hot}$) photosystem II thermal tolerance

thresholds were used to determine thermal tolerance breadth (TTB), along with ice

nucleation temperature ($T_{nucleation}$, freezing tolerance) of 69 plant species sampled from

the field across three contrasting biomes: alpine, desert and coastal temperate rainforest.

3. All measured thermal tolerance metrics ($T_{crit-cold}$, $T_{nucleation}$, $T_{crit-hot}$ and TTB) differed

among biomes. Notably, desert species had the most cold and heat tolerant leaves, and

therefore the widest TTB, whereas species in alpine and temperate biomes had similar

TTB. For plants in all biomes, TTB exceeded the thermal range of their local climate.

4. Overall, two Principal Component axes of local climate drivers explained substantial

variation in all tolerance metrics. Extreme hot, dry climates improved freezing and heat

tolerance. High thermal variability and low minimum temperatures also improved

freezing tolerance but were unrelated to heat tolerance or TTB. Species explained a

significant amount of variation among all metrics, but this was not due to phylogenetic

relatedness. We discuss how the remaining variation could be due to microclimate-driven

plasticity, leaf traits or thermoregulatory mechanisms.

5. **Synthesis**. Our results provide partial support for the climate variability hypothesis in

plants: photosystem thermal tolerance breadth was greatest in more thermally variable

biomes. This relationship was largely driven by cold tolerance, with variation in heat

tolerance explained better by mean and extreme temperatures. Therefore, we conclude

that the CVH alone is not sufficient to explain variation in plant thermal tolerance, with

many other aspects of climate, environment and biology being potentially important

drivers.

Introduction

Desert and alpine plant species live close to the thermal limits of biological processes. Unlike

those in temperate regions, plants living in extreme environments are exposed to both very low

and very high temperatures. Given the important influence of temperature in shaping species

distributions (Tattersall *et al.* 2012; Garcia-Pichel *et al.* 2013), the innate thermal tolerance

thresholds (the lowest and highest temperature that a species can withstand) and thermal

tolerance breadth (TTB; the range of temperature between upper and lower thermal tolerance

thresholds) of species might reflect the temperature variation that characterises their

environment. The climate variability hypothesis (CVH; Janzen 1967) predicts that species with

relatively wide thermal tolerance ranges are likely to evolve in environments with variable

temperatures compared to thermal specialists with relatively narrow tolerance ranges that

evolved in thermally stable environments.

The CVH has been applied across many environmental gradients (e.g. elevation, latitude,

climate) and timescales of thermal variability (e.g. Gutiérrez-Pesquera *et al.* 2016; Shah *et al.*

2021; Bovo *et al.* 2023), thus it can be interpreted differently. Here, we follow the interpretation

that the *annual temperature range* has shaped thermal tolerance breadth over evolutionary time,

allowing organisms to tolerate *seasonal shifts* in temperatures (Kefford *et al.* 2022; Dewenter *et*

al. 2024). The CVH is a cornerstone of thermal ecology; it has been tested broadly on animals

(Addo-Bediako, Chown & Gaston 2000; Compton *et al.* 2007; Calosi *et al.* 2010; Baudier *et al.*

2018) but on relatively few plant species. There is also limited consistent evidence supporting

the CVH in plants, and even studies on the same genus show mixed support (e.g. *Mimulus*;

Wooliver, Tittes & Sheth 2020; Querns *et al.* 2022; Chiono & Paul 2023).

71   Cold and heat tolerance thresholds that define the bounds of TTB are rarely studied

concomitantly in plants (Geange *et al.* 2021). In natural systems, cold tolerance studies are

dominant in alpine environments where plants can withstand very low temperatures and tolerate

extracellular ice formation and the resulting cytoplasmic dehydration (Sakai & Larcher 1987;

Larcher 2003). For example, alpine plants are known to tolerate extreme cold, many avoiding

damage below –5 °C or –10 °C (Neuner & Pramsohler 2006; Harris *et al.* 2024) and resisting

freezing below –15 °C (Venn, Morgan & Lord 2013; Neuner *et al.* 2020) or even below –30 °C

(Bannister *et al.* 2005). Heat tolerance studies have historically been primarily focussed on warm

and desert environments, where heat tolerance has often been associated with native microhabitat

conditions (Knight & Ackerly 2002; Curtis *et al.* 2016). Desert plants exhibit tolerance to heat

often in excess of 50 °C (Monson & Williams 1982; Downton, Berry & Seemann 1984;

Seemann, Berry & Downton 1984; Zhu *et al.* 2018), and some species have a remarkable ability

to tolerate over 55 °C during heatwave conditions (Andrew *et al.* 2023). Temperate plants tend to

have less capacity to tolerate heat and cold, many only tolerating cold to –5 °C (Bannister &

Lord 2006) and heat to 45 °C (Bilger, Schreiber & Lange 1984; O'Sullivan *et al.* 2017). Despite

focus on either cold or heat tolerance in either climate extreme, plants in these biomes can be

exposed to both cold and heat extremes, even within growing seasons. As the climate continues

to warm and extreme climatic events increase, understanding the cold and heat tolerance of

plants that grow in a range of biomes will be essential (Geange *et al.* 2021).

Temperature increases due to climate change has motivated an increase in comparative

studies and syntheses of plant thermal tolerance, particularly in relation to climate variability and

the occurrence of increasing extreme climatic events (Rummukainen 2012; Seneviratne *et al.*

2021; Andrew *et al.* 2023; Harris *et al.* 2024). An assessment of plant heat tolerance in four

continents showed a strong pattern of heat tolerance increasing as the mean maximum

temperature of the warmest month increased (O'Sullivan *et al.* 2017). A global meta-analysis

encompassing a wide range of species and biomes suggests that cold tolerance increases towards

the poles and heat tolerance increases towards the equator, where the latitudinal cline is a coarse

proxy for mean annual temperature (Lancaster & Humphreys 2020). However, substantial

variation in these tolerance values is due to methodological differences among studies and

locations (Perez *et al.* 2021). Heat and cold tolerance are rarely measured together within the

same study, let alone with equivalent methods, but both these aspects are important for

accurately and precisely quantifying thermal tolerance breadth (Geange *et al.* 2021; Perez *et al.*

2021).

Differing patterns of climate among contrasting biomes provide an opportunity to test the

CVH by exploring the role of climate variability in upper and lower thermal tolerance thresholds

and on thermal tolerance breadth. While broad analyses indicate that maximum temperatures can

influence heat tolerance (O'Sullivan *et al.* 2017) and mean annual temperature influences both

cold and heat tolerance (Lancaster & Humphreys 2020), the climate predictors of thermal

tolerance breadth, are virtually unknown for plants. Given that water availability affects both

plant heat tolerance (Curtis *et al.* 2016; Cook *et al.* 2021; Marchin *et al.* 2022) and freezing

tolerance (Sierra-Almeida, Reyes-Bahamonde & Cavieres 2016), precipitation is also likely to

influence thermal tolerance breadth. If we can reveal general patterns underlying variation in,

and the climate drivers of, thermal tolerance thresholds and breadth among co-occurring plant

species within and among biomes, we can strengthen predictions of which species and where

will likely be most vulnerable to changing thermal conditions.

Here we sought to determine the climate drivers of leaf-level physiological thermal

tolerance thresholds and thermal tolerance breadth of plant species growing in extreme and

comparatively benign environments in the context of the climate variability hypothesis (CVH).

Our field campaigns covered three biomes found in south-eastern Australia that contrast strongly

in their thermal environments. For simplification, hereafter we refer to these as alpine, desert and

temperate biomes. The alpine biome is characterised by both strong daily temperature

oscillations and strong temperature seasonality, with a cold winter and relatively mild summer;

the desert biome is arid, has strong daily temperature oscillations and strong temperature

seasonality, with particularly hot summers; the temperate biome has relatively mild daily and

seasonal oscillations, with mild winters and high humidity year-round (Loidi, Navarro-Sánchez

& Vynokurov 2022).

We used chlorophyll fluorescence to determine the low and high photosystem II (PSII)

thermal tolerance thresholds and TTB of 22 alpine, 24 desert and 23 temperate species during the

growing season. To examine the relationships between climate variability and thermal tolerance

thresholds, we extracted long-term, locally downscaled climate parameters for each sampling

locality. We addressed three objectives in this study. First, we assessed if and how the thermal

tolerance thresholds and TTB of a broad range of plant species differed among the contrasting

biomes. We hypothesised that thermal tolerance thresholds would reflect the climate of the

respective biomes. That is, colder threshold temperatures in the alpine and hotter threshold

temperatures in the desert, with temperate species having both milder cold and heat tolerance

thresholds. Second, we tested whether species from the three biomes differed in their TTB. We

hypothesised that species from the more thermally variable biomes (alpine and desert) would

have wider thermal tolerance breadth than those from the thermally stable biome (temperate), in

accordance with the CVH. Third, we assessed whether variation in thermal tolerance thresholds

and TTB could be explained by local climate variables, and whether these relationships were

linear or not. We hypothesised that climate extremes and variability would have a stronger

association with thermal tolerance thresholds and TTB than mean annual climate values,

supporting the CVH.

Materials and Methods

Study locations

Sites used in this study were designated to one of three biomes in New South Wales (NSW),

Australia: alpine (Monaro Ngarigo, Ngunnawal, Wiradjuri and Walgalu Country, Kosciuszko

National Park), desert (Ngemba and Barkindji Country, Gundabooka National Park) and

temperate (coastal temperate rainforests of Dharawal, Wadi Wadi and Yuin Country in the

Illawarra Region: Royal National Park, Illawarra State Conservation Area and Bass Point

Conservation Area). This study was conducted under approval from the NSW Department of

Planning, Industry and Environment (Scientific Licence SL102330). Sampling sites incorporated

a range of vegetation types sampled in each biome (mean climatic details from sites are inset in

Fig. 1). Our specific study biomes represent distinct domains, ecozones, biomes, and sub-biomes

of the world by climatic definitions (Loidi, Navarro-Sánchez & Vynokurov 2022). Specifically,

alpine falls within the cryocratic domain (sub-biome 1b: tundras of the temperate mountains in

cryoro belt); desert is in the xerocratic domain (sub-biome 7b: temperate desert and semi-desert);

and temperate is in the mesocratic domain (sub-biome 4a: lauroid evergreen forest of the

lowlands) (Loidi, Navarro-Sánchez & Vynokurov 2022).

Species selection

A total of 69 species (22 alpine, 24 desert and 23 temperate) were selected to encompass a

representative sample of growth forms and plant families and where possible, included con-

generic and/or con-familial species between two or more biomes (Table S1). We included

community dominants and a range of less abundant species to represent a cross section of species

occupying each biome. Species replicates were sampled across days within each sampling period

to incorporate daily variation in thermal tolerance thresholds.

Leaf sample collection

Mature, fully expanded leaves were harvested from five to seven individuals per species. For

shrubs and trees, small branches were collected from the northern-facing side of the plant at a

height of 1-1.5 m. For smaller growth forms, entire individuals or rosettes were collected.

Samples were kept in cool sealed plastic bags with moistened paper towel until they were

processed (approximately 1-2 h after collection). In the laboratory, leaves (size ranging from

0.25 cm^2 to 5 cm^2) were cut to around 1 cm^2 of uniform photosynthetic tissue, avoiding

prominent venation, damaged or discoloured areas or any structures that would reduce thermal

contact of the leaf material during the thermal tolerance assays. For species with smaller leaves,

we placed several leaves directly next to each other to fit an approximate area of 1 cm^2. Leaf

samples were kept on wet florist foam in humid dark containers until thermal tolerance assays

were conducted. We selected when to collect field samples to capture the peak growing season in

each biome; thus, sampling and thermal tolerance measurements occurred in January 2020 for

the desert biome, March 2020 for the temperate biome, and December 2020 for the alpine biome.

Thermal tolerance threshold assays

Photosystem II (PSII) is a protein complex embedded in the thylakoid membrane of plant cells

that captures light energy during photosynthesis. PSII has long been recognised as a thermally

sensitive structure (Schreiber & Berry 1977; Berry & Bjorkman 1980; Seemann, Berry &

Downton 1984). Chlorophyll *a* fluorescence is light that is re-emitted by chlorophyll molecules

in plants after absorbing light energy during photosynthesis, which provides valuable

information about the efficiency and functioning of the photosynthetic process, especially during

stress (Murchie & Lawson 2013). The temperature at which PSII is disrupted causes a rise in

minimal chlorophyll *a* fluorescence (F_0) as a leaf is heated or cooled (Schreiber & Berry 1977;

Neuner & Pramsohler 2006). This critical temperature indicates a threshold beyond which

physiological and photochemical systems become increasingly impaired, and damage can occur

if temperatures are sustained.

Lower and upper thresholds of PSII ($T_{crit-cold}$ and $T_{crit-hot}$)

The critical lower and upper temperature thresholds of PSII ($T_{crit-cold}$ and $T_{crit-hot}$) were measured

using the temperature-dependent (T) change in F_0 (T-F_0 curve) method, that is, the rise in F_0 with

set heating and cooling rates (Neuner & Pramsohler 2006; Arnold *et al.* 2021). One leaf per

individual wherever possible (or multiple small leaves abutting one another for plants with very

small leaves) was used for each of the heating and cooling assays (n = 5-7 replicates per species).

For each set of T-F_0 measurements, we secured 45 samples with double-sided tape to paper that

was then placed on a Peltier plate (CP-121HT; TE-Technology, Inc., Michigan, USA) that was

set to either cool or heat samples. The Peltier plate was controlled using LabView-based control

software (National Instruments, Texas, USA) and adapted code from TE-Technology, Inc.

Double-glazed glass was placed on top of the leaf samples on the plate to avoid water

condensation, particularly at freezing temperatures, and to compress samples to ensure maximum

leaf contact with the Peltier plate for uniform thermal transfer (see Arnold *et al.* 2021 for more

details).

Chlorophyll fluorescence was measured with a Pulse Amplitude Modulated (PAM)

chlorophyll fluorescence imaging system (Maxi-Imaging-PAM; Heinz Walz GmbH, Effeltrich,

Germany) mounted above the Peltier plate. Leaves were first dark-adapted, and the PSII

photochemistry maximum quantum efficiency (F_V/F_M) was measured to check the starting

function of leaves before another dark adaptation period of 15 min prior to the assay. During

temperature ramping, F_0 was measured every 20 s with a weak blue low frequency (1 Hz) pulse

modulated measuring light (0.5 μmol photons m^{-2} s^{-1}). Leaf temperatures were simultaneously

measured with 40-gauge type-T thermocouples (OMEGA Engineering, Singapore) attached to

the undersides of leaves and logged with a multi-channel datalogger (DT85 DataTaker; Lontek,

Australia). For hot T-F_0 measurements, samples were heated from ambient temperature (15-20

°C) to 60 °C at the rate of 30 °C h^{-1} (i.e. 0.5 °C min^{-1}, over ~ 1.5 h). For cold T-F_0 measurements,

the leaves were cooled down from ambient temperature to at least –20 °C, at the rate of 15 °C h^{-1}

(i.e. 0.25 °C min^{-1}, over ~ 2.5-3 h). These heating and cooling rates are moderate rates of

temperature change that are considered appropriate for interspecific comparisons (Arnold *et al.*

2021). The starting ambient temperature of each heating/cooling assay depended on the

temperature of the laboratory space used at each field site, which was generally ~15 °C for alpine

and temperate, and ~20 °C for desert biomes.

$T_{crit-cold}$ and $T_{crit-hot}$ were each determined as the point of transition between slow-rise and

fast-rise phases of F_0 values that occur during steady cooling or heating (Fig. S1; see also Arnold

et al. 2021). These critical temperature points were derived from breakpoint regression models

fitted to each T-F_0 curve, using the *segmented* R package (Muggeo 2017) (R code available from

https://github.com/pieterarnold/Tcrit-extraction). $T_{crit\text{-}cold}$ and $T_{crit\text{-}hot}$ were used to represent the

lower and upper bounds of the PSII thermal tolerance breadth, TTB (TTB = $T_{crit\text{-}hot} - T_{crit\text{-}cold}$).

Ice nucleation temperatures

Freezing tolerance is tied to the temperatures at which ice formation occurs, called the ice

nucleation temperature ($T_{nucleation}$). Measuring $T_{nucleation}$ is required to understand whether

temperature-dependent damage to PSII occurs in association with ice formation. During $T_{crit\text{-}cold}$

assays, the temperature of each leaf sample was measured continuously. $T_{nucleation}$ can be

observed as a small rise in leaf temperature release of heat energy (an exothermic reaction) when

there is a sudden state transition and ice crystals form (i.e. nucleation). By plotting leaf

temperature over time as samples cool, we identified the temperature at which the exothermic

reaction initiated, which is $T_{nucleation}$ (Larcher 2003).

Local climate variables for sampling locations

Climate data were obtained from gridded datasets at 1 km resolution based on spatial

interpolations of long-term (1981–2010) conditions accessed from the CHELSA v2.1 database

(Karger *et al.* 2017). We extracted the following local climate variables at each location that the

leaves of each species were collected: mean annual precipitation (MAP), mean annual

temperature (MAT), mean minimum temperature of the coldest month (MinT), mean maximum

temperature of the warmest month (MaxT). We calculated annual thermal range as: T_{range} =

MaxT – MinT. Table S2 includes local climate variables for the complete list of species and

sampling locations.

Statistical analyses

All analyses and data visualisations were conducted in the R Environment for Statistical

Computing version 4.5.1 (R Core Team 2023). Phylogenetic relatedness can contribute to

explaining variation in physiological traits. Therefore, prior to model fitting, we generated a

plausible phylogenetic tree using the R package *V.phylomaker* (Jin & Qian 2022) and calculated

the phylogenetic signal and statistical significance for the four response variables ($T_{crit-hot}$, $T_{crit-cold}$,

$T_{nucleation}$ and TTB) using the *phylosignal* package (Keck *et al.* 2016). Both these tests of

phylogenetic signal and incorporating phylogenetic relatedness in models described below

allowed us to account for the shared evolutionary history across the 69 species that contributed to

thermal tolerance thresholds.

To assess whether the five key local climate variables (MAP, MAT, MinT, MaxT, T_{range})

could be analysed as independent variables or not, we generated a correlation matrix using the R

package *corrplot* (Wei & Simko 2021). Since almost all local climate variables were moderately

to strongly correlated with each other, we elected to conduct a Principal Components Analysis

(PCA) with all five variables. The PCA approach means that we cannot state explicitly that MAT

or MAP or T_{range} are strong explanatory variables individually *per se*. Rather, the distinct biomes

are defined clearly by climate variables in combination. The first two Principal Component axes

(PC1 and PC2, respectively) explained 99.6% of the variation among the five climate variables;

therefore, we used these two axes as local climate variables in analyses. The PC1 axis represents

a continuum from cool and wet to extremely hot and dry. The PC2 axis represents a continuum

from thermally stable with benign minimum temperatures to thermally variable with cold

minimum temperatures.

Our first objective was to assess if and how the thermal tolerance thresholds and TTB of

a broad range of plant species differed among the contrasting biomes, while our second objective

was to test whether species from the more thermally variable alpine and desert biomes had wider

thermal tolerance breadth than temperate biome species. To address both these objectives, we

first calculated summary statistics (means and standard errors) for each thermal tolerance

threshold metric by biome and species. Next, we fitted individual linear models for the four

response variables and conducted analysis of variance (ANOVA). All response variables showed

highly significant differences among biomes based on ANOVA; therefore, we then computed

and reported pairwise differences using Tukey's Honest Significant Difference method to

compare pairs of biomes.

Our third objective was to assess whether variation in thermal tolerance thresholds and

TTB could be explained by local climate variables, and whether these relationships were linear

or not. To also account for variation in thermal tolerance thresholds and TTB due to phylogenetic

relatedness, we tested the relationships between the thermal tolerance threshold variables and

local climate variables using linear mixed effects regression models using the *Almer* function

from the *evolvability* R package (Bolstad *et al.* 2014). The *Almer* function is built on the *lmer*

function from the *lme4* R package (Bates *et al.* 2015) but incorporates a correlated random

effects structure *A*, which we used to fit a phylogenetic variance–covariance matrix to account

for phylogenetic relatedness among species. For each response variable ($T_{crit-hot}$, $T_{crit-cold}$,

$T_{nucleation}$, TTB), we fit four candidate models that included fixed effects of both climate

variables, PC1 and PC2, as either linear or second-degree (quadratic) polynomial terms in

combination (i.e. fixed effects in model 1 = PC1 + PC2; model 2 = PC1 + PC1^2 + PC2; model 3

= PC1 + PC2 + PC2^2; model 4 = PC1 + PC1^2 + PC2 + PC2^2). Each candidate model included

three random effects: species (69 levels), growth form (10 levels), and a structured phylogenetic

variance–covariance matrix. Variance explained by differences among species could therefore be

separated from variance explained by phylogenetic relatedness. We calculated Akaike's

Information Criterion (AIC) values for each model fit by maximum likelihood to allow for model

comparisons, and then calculated Akaike's weights using the *MuMIn* R package (Bartoń 2023) to

identify the most parsimonious (final) model for each thermal tolerance threshold response

variable. We then refit the final model using restricted estimate maximum likelihood (REML) to

provide unbiased parameter estimates. Variance proportions (as %) were reported for species,

growth form, and phylogeny by squaring the random effects coefficients (standard deviations)

and calculating the proportion of the random effect coefficient (e.g. species) out of the total (all

random effects including the residual) variance. Goodness-of-fit was evaluated by calculating

R^2_{GLMM} values (Johnson 2014) to show variance explained by fixed effects (marginal R^2; R^2m)

and the variance explained by both fixed and random effects (conditional R^2; R^2c). To test for

spatial autocorrelation based on sampling location, we investigated the residuals of final models

for each thermal tolerance threshold and fitted separate sets of models for each biome subset,

following the model 1 structure. We then used Moran's *I* test with inverse distance weights ($1/d^2$)

based on sampling latitude and longitude to test for significant spatial autocorrelation in all

model residuals with the *spdep* package (Bivand 2022).

Results

Objective 1: Differences in thermal tolerance thresholds among the contrasting biomes

There were significant differences among biomes for all thermal tolerance thresholds (Fig. 2A-

C). A full list of species and their individual thermal tolerance thresholds and breadth is

presented in Table S1. Unexpectedly, desert species exhibited the most extreme cold thresholds

$T_{crit-cold}$ (–13.2 ± 0.3 °C), followed by alpine (–10.9 ± 0.4 °C) and then temperate species (–8.8 ±

0.3 °C), all of which differed significantly from one another (Fig. 2A; Tables S1 and S3). Similar

to the results of $T_{crit-cold}$, $T_{nucleation}$ values were lowest in desert species (–16.1 ± 0.4 °C), followed

by alpine (–10.8 ± 0.5 °C) and highest in temperate species (–7.5 ± 0.4 °C). Again, $T_{nucleation}$

values among biomes differed significantly from one another (Fig. 2B; Tables S1 and S3).

Interestingly, $T_{crit-cold}$ values for alpine and temperate species were not significantly different

from their $T_{nucleation}$, whereas in desert species, ice formation occurred at significantly lower

temperatures than $T_{crit-cold}$ (Fig. 2A,B; Fig. S2A,B; Table S4). As predicted, desert species had

significantly higher heat tolerance $T_{crit-hot}$ (49.3 ± 0.5 °C), followed by temperate (46.5 ± 0.4 °C)

and then alpine (43.1 ± 0.4 °C) species (Fig. 2C; Tables S1 and S3).

Objective 2: Test whether thermal tolerance breadth differs among biomes

Thermal tolerance breadth (TTB) was significantly wider in desert species ($T_{crit-cold}$ - $T_{crit-hot}$ = −

13.2 - 49.3 °C, TTB = 62.5 ± 0.6 °C), than temperate ($T_{crit-cold}$ - $T_{crit-hot}$ = –8.8 - 46.5 °C, TTB =

55.3 ± 0.5 °C) or alpine species ($T_{crit-cold}$ - $T_{crit-hot}$ = –10.9 - 43.1 °C, TTB = 53.9 ± 0.6 °C), but

alpine and temperate species had similar TTB (Fig. 2D; Tables S1 and S3). There was

considerable variation among species within biomes (Fig. S3). Within the desert biome, TTB

was more similar across species (lower variability; TTB range = 56.2 - 70.3 °C; Δ = 14.1 °C),

compared to the alpine biome, which had the greatest variation among species (TTB range =

45.9 - 63.1 °C; Δ = 17.2 °C) and the temperate biome with moderate variation (TTB range = 44.8

- 60.4 °C; Δ = 15.6 °C). The rank order of species' TTB from widest to narrowest generally

reflected their biome, where 17 of the top 20 widest TTBs were all desert species (Fig. S3). By

contrast, 10 of the 20 narrowest TTBs were alpine species, and temperate species had generally

moderate to narrow TTB (Fig. S3).

Consistent across all three biomes, TTB was much wider than the extreme temperature

range calculated for the sampling sites; thus, most thermal tolerance thresholds extended beyond

the minimum and maximum temperatures expected for each species in their local environment.

The only notable exception was for the alpine biome species, where their cold threshold was

closer to the mean minimum temperature of our alpine sites (Fig. 3).

Objective 3: Local climate drivers of variation in thermal tolerance thresholds and breadth

Most of the local climate variables were strongly correlated with each other (Fig. 4A); therefore,

by conducting a PCA, 99.6% of the climate space among the three biomes could be described

along two axes (PC1 and PC2; Table S5; Fig. 4B). PC1 explained 77.2% of the variation among

the five climate variables and was strongly loaded negatively by MAP and positively by MAT

and MaxT (Fig. 4B,C). Negative PC1 values can be reasonably interpreted as cooler and wetter

climate, while positive PC1 values indicate a hotter (mean and extreme temperatures) and drier

climate (Fig. 4B). PC2 explained 22.4% of the variation among the five climate variables and

was strongly loaded positively by T_{range} and negatively by MinT (Fig. 4B,D). Positive PC2

values describe a climate with colder extremes and greater range in thermal extremes across the

year, while negative PC2 values indicate a climate with less extreme cold temperatures and

greater thermal stability through the year (Fig. 4B). No evidence for spatial autocorrelation was

detected from residuals in any model (Table S6).

We investigated the relationships between the Principal Components of climate variables

at the local site of plant sampling and each of the thermal tolerance metrics. Model comparisons

(AIC and weights) for model selection are shown in Table S7. We found that $T_{crit-cold}$ had a

significant non-linear relationship with PC1 but no significant relationship with PC2 (Table 1,

Fig. 5A,B). That is, the highest cold tolerance (i.e. most negative) was detected in plants in the

desert, followed by those in alpine, then those in temperate biomes. $T_{nucleation}$ had a significant

(negative) linear relationship with PC1 and significant non-linear relationship with PC2 (Table 1,

Fig. 5C,D). That is, the greatest freezing tolerance (i.e. ice nucleation at colder temperatures) was

detected in plants growing in hot, dry climate (desert). Plants from more thermally stable and

less extreme temperate climate were less tolerant of freezing, but freezing tolerance increased

sharply in climates with colder extremes and greater range in thermal extremes across the year

(both desert and alpine) (Fig. 5D). Species from environments with greater seasonality (i.e.

climate variability) therefore froze at lower temperatures than species in environments where

temperature is more stable (Fig. 5D).

Heat tolerance had a significant linear relationship with PC1 but not PC2 (Table 1). T_{crit-}

$_{hot}$ was highest in plants from hot, dry climate and it declined as the climate became increasingly

cool and wet (Fig. 5E,F). Thermal tolerance breadth had a significant non-linear relationship

with PC1 but was not associated with PC2 (Table 1, Fig. 5G,H). TTB was highest in hot, dry

climates and it declined non-linearly as climates became cooler and wetter, where TTB was not

different between alpine and temperate species (Fig. 5G). For all thermal tolerance thresholds,

among-species differences explained a substantial proportion of variance (variance proportions

due to species from Table 1 as percentage: $T_{crit-cold}$ = 26.3%, $T_{nucleation}$ = 42.8%, $T_{crit-hot}$ = 24.8%,

TTB = 26.8%). Plant growth form explained some variance, which was greatest for TTB

(variance proportions due to growth form from Table 1 as percentage: $T_{crit-cold}$ = 13.4%, $T_{nucleation}$

= 8.9%, $T_{crit-hot}$ = 1.3%, TTB = 11.2%). Among-species differences were not attributable to

phylogenetic relatedness (Table 1). All four phylogenetic signal metrics consistently identified

no evidence of phylogenetic signal in the thermal tolerance thresholds or breadth (Table S8).

Discussion

The climate variability hypothesis (CVH) predicts that species will evolve wider thermal

tolerance breadth (TTB) in environments with more variable temperatures and be selected for

thermal specialisation (narrow TTB) in thermally stable environments. The CVH has been tested

in a few plant systems across different contexts, with limited support (Molina-Montenegro &

Naya 2012; Cuesta *et al.* 2020; Wooliver, Tittes & Sheth 2020; Querns *et al.* 2022; Chiono &

Paul 2023). For example, two studies on *Mimulus guttatus* did not find support for stronger

seasonality being associated with TTB based on thermal performance curves of relative growth

rate (Querns *et al.* 2022; Chiono & Paul 2023). By contrast, another study across the geographic

range of *Mimulus cardinalis* found that TTB of relative growth rate was indeed broader in

populations found in more seasonal climates (Wooliver, Tittes & Sheth 2020). Here we explored

the CVH in diverse plants by applying high-throughput measurements of heat and cold tolerance

thresholds of photosystem II concurrently to determine TTB of species growing in two opposite

extremes and thermally variable biomes (desert and alpine) and one benign, thermally stable

biome (temperate). Overall, we found that mean annual and mean maximum temperatures, and

mean annual precipitation had a stronger relationship with TTB than did thermal variability and

mean minimum temperatures, which provides only partial support for the CVH. Below we

explore our findings in terms of physiology and the ecological implications of these results for

species persistence in a changing climate.

Heat and cold tolerance thresholds vary significantly among biomes (Objective 1)

Based on the CVH framework, we predicted that the thermal tolerance thresholds of alpine and

desert species would be skewed to their respective biome extremes; that is, colder threshold

temperatures in the alpine and hotter threshold temperatures in the desert, with temperate species

having both milder cold and heat tolerance thresholds. Here we found partial support for this

prediction. Temperate species had heat tolerance intermediate between alpine and desert but

were less cold tolerant than either alpine or desert; desert species were more heat tolerant than

alpine and temperate species but were also more tolerant of cold and freezing than alpine species.

The apparently counterintuitive result that desert species were more tolerant of cold and

resistant to freezing (more extreme $T_{crit-cold}$ and $T_{nucleation}$ values; Fig S2) than either alpine or

temperate species may be partially explained by the physiological response to drought

conditions. That is, plants exposed to drought and/or freezing share a similar physiological

response to cell dehydration (Anisko & Lindstrom 1996; Blake & Hill 1996; Lintunen, Hölttä &

Kulmala 2013). Plants can use osmotic adjustments to offset water loss from cells along the

water potential gradient caused by both drying and freezing (Siminovitch & Cloutier 1983;

Larcher 2003). In response to freezing conditions, plants typically either tolerate ice formation

by coping with cellular dehydration (Scholz *et al.* 2012) or avoid ice formation by supercooling,

which refers to preventing ice formation below the equilibrium freezing temperature of the leaf

tissue (Pearce 2001). Here, the desert species that we studied generally had ice nucleation

temperatures ($T_{nucleation}$) that were much lower than their $T_{crit-cold}$, from which we can infer a high

supercooling capacity that delays freezing (Fig. S2). There are several examples that demonstrate

the link between freezing and dehydration. Extreme $T_{nucleation}$ found in Andean alpine species can

be attributed to low soil moisture rather than low temperatures (Sierra-Almeida, Reyes-

Bahamonde & Cavieres 2016). Drought also improves both heat and cold tolerance in an alpine

grass (Sumner *et al.* 2022), and drought can increase heat tolerance by raising intracellular sugar

content, which stabilises thylakoid membranes (Hüve *et al.* 2006; Brestic *et al.* 2012). Therefore,

we postulate that the extreme tolerance in these desert species reflects a secondary effect of their

adaptations to aridity.

Alpine plants generally tolerated cold and freezing very well, as expected. Retaining high

cold tolerance remains necessary for protecting investment in valuable photosynthetic tissue due

to exposure to subzero temperatures without insulative snow during growing seasons (Slatyer,

Umbers & Arnold 2022). The relatively high heat tolerance of some of our alpine species may

seem surprising; however, these heat tolerance ranges are similar to recent measures of heat

tolerance in other alpine species (43-55 °C, Sumner & Venn 2022; Arnold *et al.* 2024; Harris *et*

al. 2024; Morales *et al.* 2025). Alpine plants can be exposed to very high temperatures,

especially those with prostrate growth forms (Körner 2003). Leaf temperature of European

alpine species can reach 50 °C in summer on calm sunny days, and leaf heat tolerance thresholds

in these plants can exceed 50 °C (Buchner & Neuner 2003). Maximum leaf temperature has been

measured at nearly 40 °C in early summer in Australian alpine species (Bird *et al.*, unpublished

data) and may yet be much higher in hot and still conditions in late summer. Therefore, despite

the apparently high heat tolerance thresholds, these could be breached in alpine species if heat

interacts with other environmental stress, such as high light, which can cause heat tolerance to

decrease in some European alpine species (Buchner & Neuner 2003; Buchner *et al.* 2015).

The tolerance of temperate plants was generally less than that of plants from the more

extreme biomes, reflecting the more stable and less extreme climate of the biome. These

temperate species tolerated temperatures well outside their typical range. Nevertheless, under

warmer conditions in the future, vapour pressure deficit, which drives higher evaporative

demand, will increase in wet temperate and tropical rainforests (Binks *et al.* 2023). Plants

growing in these conditions will have greater water loss, and if water becomes limiting,

transpirational cooling capacity will diminish, consequently pushing leaf temperature closer to

heat tolerance thresholds (Zhou *et al.* 2023).

Thermal tolerance breadth is wider in more thermally variable alpine and desert biomes

and exceeds the thermal range of each biome (Objective 2)

The thermal tolerance breadth (TTB) values based on PSII tolerance that we report here are

notably wide—wider than the thermal ranges in each environment and wider than many other

measurements of TTB in other organisms. Across a range of marine and terrestrial ectotherm

taxa, the widest recorded TTB was 60 °C in terrestrial arthropods (Sunday, Bates & Dulvy

2011), while the widest TTB in our dataset was 70.3 °C in the desert species *Rhagodia*

spinescens, with several other desert species also exceeding 60 °C. Within each biome, we found

species with a range of wide and narrow TTB (Fig. S3). For example, several alpine species

across growth forms had comparatively narrow TTB, such as the snow bed species *Psychrophila*

introloba (TTB = 48.9 °C) and the tree line species *Eucalyptus pauciflora* subsp. *niphophila*

(TTB = 47.7 °C). The narrow TTB of some alpine species was mainly driven by their relatively

low heat tolerance thresholds ($T_{crit-hot} \approx 40$ °C), whereas other co-occurring species, *Hovea*

montana and *Epacris paludosa* had TTB values similar to those of many desert species (~63 °C),

which was driven by their extreme cold tolerance ($T_{crit-cold} \approx -18$ °C). The notable within-biome

variation in TTB might be due to variation in microclimatic niche among species, which can

override broad-scale climate drivers of physiological tolerance (Curtis *et al.* 2016; Curtis, Knight

& Leigh 2019; Aparecido *et al.* 2020).

Although thermal seasonality was greater in the alpine than in the temperate regions

sampled here, the average TTB for these two biomes was remarkably similar (Fig. 2D).

Temperate species appeared to tolerate colder temperatures than expected, and it is possible that

their relatively wide thermal tolerance is a legacy of exposure to climate extremes through

geological time (Byrne *et al.* 2008). The climate legacy effect on thermal tolerance is suggested

to be stronger for cold than for heat tolerances across both animal and plant taxa (Bennett *et al.*

2021), which might be an additional explanation as to why some temperate and desert species

have greater freezing tolerance than expected based on their current climate envelopes. Although

climate metrics based on air temperatures for our alpine biome confirm that thermal range and

seasonality are large in this biome, these mountain plants can spend ~4 months under the

insulating cover of snow, which stabilises plant temperature (Briceño *et al.* 2014). The

dampening of climate extremes at a local scale (i.e. microclimate) might also contribute to

similar TTB values between alpine and temperate species.

We find that when considering climate variability from a discrete biome perspective, our

results provide only partial support for the CVH. That is, plants from the climatically variable

desert biome have the widest TTB, but temperate and alpine plants do not have different TTB

despite the greater daily and seasonal climate variability in the alpine environment. Investigating

the climatic conditions occurring in each biome provides further insight into the drivers behind

each thermal tolerance threshold and TTB.

Thermal tolerance thresholds and breadth are driven by different climatic variables

(Objective 3)

Thermal tolerance is difficult to predict based on individual bioclimatic parameters, such as

mean annual temperature and mean annual precipitation (Perez & Feeley 2021). Here, the

Principal Component axis that combined MAT, MAP and MaxT explained considerable

variation in $T_{crit\text{-}cold}$ and $T_{nucleation}$. As discussed above, the much lower $T_{nucleation}$ than $T_{crit\text{-}cold}$ in

desert species suggests that $T_{nucleation}$ is more strongly associated with aridity than is $T_{crit\text{-}cold}$,

supporting the idea of shared mechanisms of tolerance to freezing and dehydration under

drought. Although plants growing in extremely hot and dry environments do have higher heat

tolerance, the high correlation of MAP with both MAT and MaxT means that the role of

temperature versus precipitation cannot be completely separated. For a range of common-grown

Australian *Acacia* species from widespread climatic and geographic origins, precipitation did not

strongly drive variation in $T_{crit-hot}$ (Andrew *et al.* 2023). Even so, water availability and

temperature are intrinsically linked in terms of plant stress responses. Water limitation can

drastically reduce stomatal conductance and therefore cooling capacity during exposure to heat,

increasing leaf temperature and affecting heat tolerance thresholds (Cook *et al.* 2021).

Local climate, particularly MAT and MaxT, seems to play a significant role in driving

variation in $T_{crit-hot}$ in our field study, which contrasts with a common garden study of diverse

species, for which origin climate generally had weak predictive power for heat tolerance (Perez

& Feeley 2021). It is important to reiterate that MinT and MaxT as analysed here are mean

values. Daily or hourly extreme temperatures can far exceed MinT and MaxT for short periods;

therefore, the apparently wide buffer that TTB may provide could occasionally be much

narrower than these patterns suggest. However, it is also important to recognise that many plants

have the capacity to regulate their leaf temperature. A controlled environment study on a subset

of species studied here found that alpine and desert plants cooled their leaves by 1-3 °C below air

temperature during a 38 °C heat event, while temperate plants were frequently 1-2 °C above air

temperature (Arnold *et al.* 2025). Strong thermal decoupling has been found in less heat-tolerant

alpine plant species as an effective heat avoidance strategy (Morales *et al.* 2025). When water is

not limiting, some species can thermoregulate to prevent leaves from reaching damaging

temperatures, but that strategy may be nullified during hotter droughts (Cook *et al.* 2021; Arnold

et al. 2025). Underpinning thermoregulation strategies are combinations of leaf structural traits

(leaf density, thickness, water content) and stomatal behaviour that also vary among species and

across biomes (Arnold *et al.* 2025).

A large proportion of the variation in thermal tolerance thresholds in our study was also

attributable to among-species differences that were unrelated to phylogeny. The contribution of

species differences could reflect their different leaf traits, particularly those contributing to

variation in leaf temperature, the signal of which may be amplified in the field (Leigh *et al.*

2017). Recent studies have shown that a major driver of variation in heat tolerance is leaf

temperature, rather than air temperature (Perez & Feeley 2020; Cook *et al.* 2021). Leaves can

avoid heat stress through different mechanisms that decouple leaf from air temperature, such as

anatomical traits (Leigh *et al.* 2012; Leigh *et al.* 2017; Tserej & Feeley 2021; Arnold *et al.*

2025), leaf inclination (Ball, Cowan & Farquhar 1988) and transpirational cooling (Drake *et al.*

2018; Deva *et al.* 2020), all of which can vary among co-existing species (Urban *et al.* 2017;

Marchin *et al.* 2022).

The capacity for different species to respond to increasingly extreme and variable

temperatures by changing their TTB is only recently emerging. We found that TTB increased

non-linearly (hyper-allometrically) with higher MAT and MaxT and lower MAP (Fig. S4). That

is, plants exposed to the high mean temperatures, coupled with aridity and heat extremes in

deserts had disproportionally wider TTB than temperate or alpine plants. TTB of high elevation

species from temperate and tropical regions appears to differ only at the end of the temperate

growing season, due to an increase in cold acclimation (Sklenář *et al.* 2023). In a common-

garden experiment with seedlings from the same biomes that were tested here, TTB narrowed

when exposed to a combination of hot days and freezing nights in controlled climate chambers

(Harris *et al.* 2024). However, that study found a minimal effect of biome, which contrasts with

the significant biome-based differences in thermal tolerance thresholds observed here for plants

in situ. The distinct responses between these contexts imply that life stage and current growing

conditions could be important modifiers of thermal tolerance.

The effects of immediate growth conditions can outweigh other environmental and

genetic sources of variation in trait changes in plants grown in common gardens (Arnold *et al.*

2024). In alpine species at least, both cold and heat tolerance of PSII can vary widely among

genetically similar populations distributed along an elevational gradient, suggesting intraspecific

variance in thermal tolerance is driven by acclimation to local conditions at a fine scale (Danzey

et al. 2024). Whether PSII thermal tolerance is heritable and under selection in wild plants

remains largely unknown, but evidence so far suggests it is not (Arnold *et al.* 2024). PSII thermal

tolerance can be highly heritable in rice species (Robson *et al.* 2023). Yet, evidence from wheat

shows $T_{crit-hot}$ in field-planted crops is strongly dependent on growth temperature during the days

preceding anthesis, and that $T_{crit-hot}$ is far less heritable in the field than in controlled

environments (Coast *et al.* 2022). Together, these studies and our findings here suggest that PSII

thermal tolerance thresholds and TTB will be dependent both on the physical characteristics that

differentiate species and local climate factors at fine spatial and temporal scales.

Conclusions

While there were substantial differences in thermal tolerance thresholds among biomes, we

found limited support for the climate variability hypothesis. Rather than climate variability,

warmer mean annual temperature, maximum temperature, and lower precipitation were stronger

drivers of wider TTB, particularly in desert plants, which were more tolerant of both extreme

heat and cold than alpine or temperate plants. Thermal tolerance breadth of PSII in leaves was

extremely wide in all biomes—wider than the typical air temperature extremes to which plants

are naturally exposed in these biomes. Sustained extreme temperatures within the bounds of

thermal tolerance breadth could still have far-reaching effects, depending on species life history,

morphological traits and plasticity, as well as microclimatic and temporal heterogeneity. While

local climate partially explains variability in plant thermal tolerance thresholds and breadth,

incorporating additional traits, leaf temperature and microhabitat information through time are

important next steps to better understand the vulnerability of different vegetation communities to

climate change and extreme climatic events.

Acknowledgements

We acknowledge the Traditional Custodians of Country throughout Australia, especially across

the lands on which this study took place, and recognise their continuing connection to these

lands, waters and natural communities. We pay our respects to Aboriginal and Torres Strait

Islander cultures and to Elders past, present and emerging. We thank the National Parks and

Wildlife Service for assistance and support, with special thanks to M. Hams, S. Doak and R.

Bjork at Gundabooka National Park, M. Schroder at Kosciuszko National Park, and J. Erskine, J.

Lemmon and P. Nagle at Royal National Park and Illawarra State Conservation Area. We thank

the following people for their assistance in the field: C. Bellotto, Z. Brown, A. Crimmins, C.

Cunningham, P. Cunningham, L. Danzey, M. Davies, L. Fuller, J.M. Hoskins, T. Salisbury, E.

Rogers and K. Spooner. This study was conducted under approval from the NSW Department of

Planning, Industry and Environment (Scientific Licence SL102330). We thank the anonymous

reviewers and editors that provided constructive feedback that improved this manuscript. The

work was supported by an Australian Research Council Linkage Project grant (LP180100942) in

partnership with the Save our Species program, New South Wales Government Department of

Planning, Industry and Environment.

Conflict of interest statement

The authors have no conflicts of interest to disclose.

Author Contributions

Verónica F. Briceño, Alicia M. Cook, Stephanie K. Courtney Jones, Adrienne B. Nicotra, León

A. Bravo and Andy Leigh designed the study. Pieter A. Arnold, Rachael V. Gallagher, Verónica

F. Briceño and Alicia M. Cook collated and analysed the data. Verónica F. Briceño and Pieter A.

Arnold led the writing and revisions of the manuscript. All authors contributed to intellectual

conceptualisation, data collection, interpretation of results and writing. All authors meet criteria

for authorship.

Data availability statement

Data and code supporting this study are available in the figshare repository at

https://doi.org/10.6084/m9.figshare.29964764 (Briceño *et al.* 2025).

References

Addo-Bediako, A., Chown, S.L. & Gaston, K.J. (2000) Thermal tolerance, climatic variability and latitude. *Proceedings of the Royal Society of London B: Biological Sciences,* **267,** 739-745. https://doi.org/10.1098/rspb.2000.1065.

Andrew, S.C., Arnold, P.A., Simonsen, A.K. & Briceño, V.F. (2023) Consistently high heat tolerance acclimation in response to a simulated heatwave across species from the broadly distributed *Acacia* genus. *Functional Plant Biology,* **50,** 71-83. https://doi.org/10.1071/FP22173.

Anisko, T. & Lindstrom, O.M. (1996) Cold hardiness and water relations parameters in *Rhododendron* cv. Catawbiense Boursault subjected to drought episodes. *Physiologia Plantarum,* **98,** 147-155. https://doi.org/10.1111/j.1399-3054.1996.tb00686.x.

Aparecido, L.M.T., Woo, S., Suazo, C., Hultine, K.R. & Blonder, B. (2020) High water use in desert plants exposed to extreme heat. *Ecology Letters,* **23,** 1189-1200. https://doi.org/10.1111/ele.13516.

Arnold, P.A., Briceño, V.F., Gowland, K.M., Catling, A.A., Bravo, L.A. & Nicotra, A.B. (2021) A high-throughput method for measuring critical thermal limits of leaves by chlorophyll imaging fluorescence. *Functional Plant Biology,* **48,** 634-646. https://doi.org/10.1071/FP20344.

Arnold, P.A., Wang, S., Notarnicola, R.F., Nicotra, A.B. & Kruuk, L.E.B. (2024) Testing the evolutionary potential of an alpine plant: phenotypic plasticity in response to growth temperature outweighs parental environmental effects and other genetic causes of variation. *Journal of Experimental Botany,* **75,** 5971-5988. https://doi.org/10.1093/jxb/erae290.

Arnold, P.A., White, M.J., Cook, A.M., Leigh, A., Briceño, V.F. & Nicotra, A.B. (2025) Plants originating from more extreme biomes have improved leaf thermoregulation. *Annals of Botany***,** mcaf080. https://doi.org/10.1093/aob/mcaf080.

Ball, M., Cowan, I. & Farquhar, G. (1988) Maintenance of leaf temperature and the optimisation of carbon gain in relation to water loss in a tropical mangrove forest. *Functional Plant Biology,* **15,** 263-276. https://doi.org/10.1071/PP9880263.

Bannister, P. & Lord, J.M. (2006) Comparative winter frost resistance of plant species from southern Africa, Australia, New Zealand, and South America grown in a common environment (Dunedin, New Zealand). *New Zealand Journal of Botany,* **44,** 109-119. https://doi.org/10.1080/0028825X.2006.9513011.

Bannister, P., Maegli, T., Dickinson, K.J.M., Halloy, S.R.P., Knight, A., Lord, J.M., Mark, A.F.
& Spencer, K.L. (2005) Will loss of snow cover during climatic warming expose New
Zealand alpine plants to increased frost damage? *Oecologia,* **144,** 245-256.
https://doi.org/10.1007/s00442-005-0087-3.

Bartoń, K.A. (2023) Package 'MuMIn'. Multi-model inference. http://cran.r-
project.org/web/package=MuMIn), R package verision 1.47.5.

Bates, D., Mächler, M., Bolker, B. & Walker, S. (2015) Fitting linear mixed-effects models using
lme4. *Journal of Statistical Software,* **67,** 1-48. https://doi.org/10.18637/jss.v067.i01.

Baudier, K.M., D'Amelio, C.L., Malhotra, R., O'Connor, M.P. & O'Donnell, S. (2018) Extreme
insolation: climatic variation shapes the evolution of thermal tolerance at multiple scales.
The American Naturalist, **192,** 347-359. https://doi.org/10.1086/698656.

Bennett, J.M., Sunday, J., Calosi, P., Villalobos, F., Martinez, B., Molina-Venegas, R., Araujo,
M.B., Algar, A.C., Clusella-Trullas, S., Hawkins, B.A., Keith, S.A., Kuhn, I., Rahbek, C.,
Rodriguez, L., Singer, A., Morales-Castilla, I. & Olalla-Tarraga, M.A. (2021) The
evolution of critical thermal limits of life on Earth. *Nature Communications,* **12,** 1198.
https://doi.org/10.1038/s41467-021-21263-8.

Berry, J. & Bjorkman, O. (1980) Photosynthetic response and adaptation to temperature in
higher plants. *Annual Review of Plant Physiology,* **31,** 491-543.
https://doi.org/10.1146/annurev.pp.31.060180.002423.

Bilger, H.-W., Schreiber, U. & Lange, O.L. (1984) Determination of leaf heat resistance:
comparative investigation of chlorophyll fluorescence changes and tissue necrosis
methods. *Oecologia,* **63,** 256-262. https://doi.org/10.1007/BF00379886.

Binks, O., Cernusak, L.A., Liddell, M., Bradford, M., Coughlin, I., Bryant, C., Palma, A.C.,
Hoffmann, L., Alam, I., Carle, H.J., Rowland, L., Oliveira, R.S., Laurance, S.G.W.,
Mencuccini, M. & Meir, P. (2023) Vapour pressure deficit modulates hydraulic function
and structure of tropical rainforests under nonlimiting soil water supply. *New Phytologist,*
240, 1405-1420. https://doi.org/10.1111/nph.19257.

Bivand, R. (2022) R packages for analyzing spatial data: A comparative case study with areal
data. *Geographical Analysis,* **54,** 488-518. https://doi.org/10.1111/gean.12319.

Blake, J. & Hill, R. (1996) An examination of the drought and frost tolerance of *Banksia*
marginata (Proteaceae) as an explanation of its current widespread occurrence in
Tasmania. *Australian Journal of Botany,* **44,** 265-281.
https://doi.org/10.1071/BT9960265.

Bolstad, G.H., Hansen, T.F., Pélabon, C., Falahati-Anbaran, M., Perez-Barrales, R. &
Armbruster, W.S. (2014) Genetic constraints predict evolutionary divergence in
Dalechampia. *Philosophical Transactions of the Royal Society of London B: Biological
Sciences,* **369,** 20130255. https://doi.org/10.1098/rstb.2013.0255.

Bovo, R.P., Simon N.M., Provete, D.B., Lyra, M., Navas, C.A. & Andrade, D.V. (2023) Beyond
Janzen's hypothesis: How amphibians that climb tropical mountains respond to climate
variation. *Integrative Organismal Biology,* **5,** obad009.
https://doi.org/10.1093/iob/obad009.

Brestic, M., Zivcak, M., Kalaji, H.M., Carpentier, R. & Allakhverdiev, S.I. (2012) Photosystem
II thermostability in situ: Environmentally induced acclimation and genotype-specific
reactions in *Triticum aestivum* L. *Plant Physiology and Biochemistry,* **57,** 93-105.
https://doi.org/10.1016/j.plaphy.2012.05.012.

Briceño, V.F., Arnold, P.A., Cook, A.M., Courtney Jones, S.K., Gallagher, R.V., French, K.,
Bravo, L.A., Nicotra, A.B. & Leigh, A. (2025) Data supporting: Drivers of thermal
tolerance breadth of plants across contrasting biomes: do mean or seasonality in climate
indices matter more? *figshare Repository*. https://doi.org/10.6084/m9.figshare.29964764.

Briceño, V.F., Harris-Pascal, D., Nicotra, A.B., Williams, E. & Ball, M.C. (2014) Variation in
snow cover drives differences in frost resistance in seedlings of the alpine herb *Aciphylla
glacialis. Environmental and Experimental Botany,* **106,** 174-181.
https://doi.org/10.1016/j.envexpbot.2014.02.011.

Buchner, O. & Neuner, G. (2003) Variability of heat tolerance in alpine plant species measured
at different altitudes. *Arctic, Antarctic and Alpine Research,* **35,** 411-420.
https://doi.org/10.1657/1523-0430(2003)035[0411:Vohtia]2.0.Co;2.

Buchner, O., Stoll, M., Karadar, M., Kranner, I. & Neuner, G. (2015) Application of heat stress
in situ demonstrates a protective role of irradiation on photosynthetic performance in
alpine plants. *Plant, Cell & Environment,* **38,** 812-826.
https://doi.org/10.1111/pce.12455.

Byrne, M., Yeates, D.K., Joseph, L., Kearney, M., Bowler, J., Williams, M.A.J., Cooper, S.,
Donnellan, S.C., Keogh, J.S., Leys, R., Melville, J., Murphy, D.J., Porch, N. & Wyrwoll,
K.H. (2008) Birth of a biome: insights into the assembly and maintenance of the
Australian arid zone biota. *Molecular Ecology,* **17,** 4398-4417.
https://doi.org/10.1111/j.1365-294X.2008.03899.x.

Calosi, P., Bilton, D.T., Spicer, J.I., Votier, S.C. & Atfield, A. (2010) What determines a species'
geographical range? Thermal biology and latitudinal range size relationships in European
diving beetles (Coleoptera: Dytiscidae). *Journal of Animal Ecology,* **79,** 194-204.
https://doi.org/10.1111/j.1365-2656.2009.01611.x.

Chiono, A. & Paul, J.R. (2023) The Climatic Variability Hypothesis and trade-offs in thermal
performance in coastal and inland populations of *Mimulus guttatus*. *Evolution,* **77,** 870-
880. https://doi.org/10.1093/evolut/qpad005.

Coast, O., Posch, B.C., Rognoni, B.G., Bramley, H., Gaju, O., Mackenzie, J., Pickles, C., Kelly,
A.M., Lu, M., Ruan, Y.-L., Trethowan, R. & Atkin, O.K. (2022) Wheat photosystem II
heat tolerance: evidence for genotype-by-environment interactions. *The Plant Journal,*
111, 1368-1382. https://doi.org/10.1111/tpj.15894.

Compton, T.J., Rijkenberg, M.J.A., Drent, J. & Piersma, T. (2007) Thermal tolerance ranges and
climate variability: A comparison between bivalves from differing climates. *Journal of*
Experimental Marine Biology and Ecology, **352,** 200-211.
https://doi.org/10.1016/j.jembe.2007.07.010.

Cook, A.M., Berry, N., Milner, K.V. & Leigh, A. (2021) Water availability influences thermal
safety margins for leaves. *Functional Ecology,* **35,** 2179-2189.
https://doi.org/10.1111/1365-2435.13868.

Cuesta, F., Tovar, C., Llambí, L.D., Gosling, W.D., Halloy, S., Carilla, J., Muriel, P., Meneses,
R.I., Beck, S., Ulloa Ulloa, C., Yager, K., Aguirre, N., Viñas, P., Jácome, J., Suárez-
Duque, D., Buytaert, W. & Pauli, H. (2020) Thermal niche traits of high alpine plant
species and communities across the tropical Andes and their vulnerability to global
warming. *Journal of Biogeography,* **47,** 408-420. https://doi.org/10.1111/jbi.13759.

Curtis, E.M., Gollan, J., Murray, B.R. & Leigh, A. (2016) Native microhabitats better predict
tolerance to warming than latitudinal macro-climatic variables in arid-zone plants.
Journal of Biogeography, **43,** 1156-1165. https://doi.org/10.1111/jbi.12713.

Curtis, E.M., Knight, C.A. & Leigh, A. (2019) Intracanopy adjustment of leaf-level thermal
tolerance is associated with microclimatic variation across the canopy of a desert tree
(*Acacia papyrocarpa*). *Oecologia,* **189,** 37-46. https://doi.org/10.1007/s00442-018-4289-
x.

Danzey, L.M., Briceño, V.F., Cook, A.M., Nicotra, A.B., Peyre, G., Rossetto, M., Yap, J.-Y.S. &
Leigh, A. (2024) Environmental and biogeographic drivers behind alpine plant thermal
tolerance and genetic variation. *Plants*.

Deva, C.R., Urban, M.O., Challinor, A.J., Falloon, P. & Svitáková, L. (2020) Enhanced leaf
cooling is a pathway to heat tolerance in common bean. *Frontiers in Plant Science,* **11,**
19. https://doi.org/10.3389/fpls.2020.00019.

Dewenter, B.S., Shah, A.A., Hughes, J., Poff, N.L., Thompson, R. & Kefford, B.J. (2024) The
thermal breadth of temperate and tropical freshwater insects supports the climate
variability hypothesis. *Ecology and Evolution,* **14,** e10937.
https://doi.org/10.1002/ece3.10937.

Downton, W.J.S., Berry, J.A. & Seemann, J.R. (1984) Tolerance of photosynthesis to high
temperature in desert plants. *Plant Physiology,* **74,** 786-790.
https://doi.org/10.1104/pp.74.4.786.

Drake, J.E., Tjoelker, M.G., Vårhammar, A., Medlyn, Belinda E., Reich, P.B., Leigh, A.,
Pfautsch, S., Blackman, C.J., López, R., Aspinwall, M.J., Crous, K.Y., Duursma, R.A.,
Kumarathunge, D., De Kauwe, M.G., Jiang, M., Nicotra, A.B., Tissue, D.T., Choat, B.,
Atkin, O.K. & Barton, C.V.M. (2018) Trees tolerate an extreme heatwave via sustained
transpirational cooling and increased leaf thermal tolerance. *Global Change Biology,* **24,**
2390-2402. https://doi.org/10.1111/gcb.14037.

Garcia-Pichel, F., Loza, V., Marusenko, Y., Mateo, P. & Potrafka, R.M. (2013) Temperature
drives the continental-scale distribution of key microbes in topsoil communities. *Science,*
340, 1574-1577. https://doi.org/10.1126/science.1236404.

Geange, S.R., Arnold, P.A., Catling, A.A., Coast, O., Cook, A.M., Gowland, K.M., Leigh, A.,
Notarnicola, R.F., Posch, B.C., Venn, S.E., Zhu, L. & Nicotra, A.B. (2021) The thermal
tolerance of photosynthetic tissues: a global systematic review and agenda for future
research. *New Phytologist,* **229,** 2497-2513. https://doi.org/10.1111/nph.17052.

Gutiérrez-Pesquera, L.M., Tejedo, M., Olalla-Tárraga, M.Á., Duarte, H., Nicieza, A. & Solé, M.
(2016) Testing the climate variability hypothesis in thermal tolerance limits of tropical
and temperate tadpoles. *Journal of Biogeography,* **43,** 1166-1178.
https://doi.org/10.1111/jbi.12700.

Harris, R.J., Alvarez, P.R., Bryant, C., Briceño, V.F., Cook, A.M., Leigh, A. & Nicotra, A.B.
(2024) Acclimation of thermal tolerance in juvenile plants from three biomes is
suppressed when extremes co-occur. *Conservation Physiology,* **12,** coae027.
https://doi.org/10.1093/conphys/coae027.

Hüve, K., Bichele, I., Tobias, M. & Niinemets, Ü. (2006) Heat sensitivity of photosynthetic
electron transport varies during the day due to changes in sugars and osmotic potential.

Plant, Cell & Environment, **29,** 212-228. https://doi.org/10.1111/j.1365-
3040.2005.01414.x.

Janzen, D.H. (1967) Why mountain passes are higher in the tropics. *The American Naturalist,*
101, 233-249. https://doi.org/10.1086/282487.

Jin, Y. & Qian, H. (2022) V.PhyloMaker2: An updated and enlarged R package that can generate
very large phylogenies for vascular plants. *Plant Diversity,* **44,** 335-339.
https://doi.org/10.1016/j.pld.2022.05.005.

Johnson, P.C.D. (2014) Extension of Nakagawa & Schielzeth's R^2_{GLMM} to random slopes models.
Methods in Ecology and Evolution, **5,** 944-946. https://doi.org/10.1111/2041-
210X.12225.

Karger, D.N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R.W., Zimmermann,
N.E., Linder, H.P. & Kessler, M. (2017) Climatologies at high resolution for the earth's
land surface areas. *Scientific Data,* **4,** 170122. https://doi.org/10.1038/sdata.2017.122.

Keck, F., Rimet, F., Bouchez, A. & Franc, A. (2016) phylosignal: an R package to measure, test,
and explore the phylogenetic signal. *Ecology and Evolution,* **6,** 2774-2780.
https://doi.org/10.1002/ece3.2051.

Kefford, B.J., Ghalambor, C.K., Dewenter, B., Poff, N.L., Hughes, J., Reich, J. & Thompson, R.
(2022) Acute, diel, and annual temperature variability and the thermal biology of
ectotherms. *Global Change Biology,* **28,** 6872-6888. https://doi.org/10.1111/gcb.16453.

Knight, C.A. & Ackerly, D.D. (2002) An ecological and evolutionary analysis of photosynthetic
thermotolerance using the temperature-dependent increase in fluorescence. *Oecologia,*
130, 505-514. https://doi.org/10.1007/s00442-001-0841-0.

Körner, C. (2003) *Alpine Plant Life,* Second edn. Springer, Berlin.

Lancaster, L.T. & Humphreys, A.M. (2020) Global variation in the thermal tolerances of plants.
Proceedings of the National Academy of Sciences of the United States of America, **117,**
13580-13587. https://doi.org/10.1073/pnas.1918162117.

Larcher, W. (2003) *Physiological plant ecology: Ecophysiology and stress physiology of*
functional group. Springer-Verlag

Leigh, A., Sevanto, S., Ball, M.C., Close, J.D., Ellsworth, D.S., Knight, C.A., Nicotra, A.B. &
Vogel, S. (2012) Do thick leaves avoid thermal damage in critically low wind speeds?
New Phytologist, **194,** 477-487. https://doi.org/10.1111/j.1469-8137.2012.04058.x.

Leigh, A., Sevanto, S., Close, J.D. & Nicotra, A.B. (2017) The influence of leaf size and shape
on leaf thermal dynamics: does theory hold up under natural conditions? *Plant, Cell &*
Environment, **40,** 237-248. https://doi.org/10.1111/pce.12857.

Lintunen, A., Hölttä, T. & Kulmala, M. (2013) Anatomical regulation of ice nucleation and
cavitation helps trees to survive freezing and drought stress. *Scientific Reports,* **3,** 2031.
https://doi.org/10.1038/srep02031.

Loidi, J., Navarro-Sánchez, G. & Vynokurov, D. (2022) Climatic definitions of the world's
terrestrial biomes. *Vegetation Classification and Survey,* **3,** 231-271.
https://doi.org/10.3897/VCS.86102.

Marchin, R.M., Backes, D., Ossola, A., Leishman, M.R., Tjoelker, M.G. & Ellsworth, D.S.
(2022) Extreme heat increases stomatal conductance and drought☐induced mortality risk
in vulnerable plant species. *Global Change Biology,* **28,** 1133-1146.
https://doi.org/10.1111/gcb.15976.

Molina-Montenegro, M.A. & Naya, D.E. (2012) Latitudinal patterns in phenotypic plasticity and
fitness-related traits: Assessing the Climatic Variability Hypothesis (CVH) with an
invasive plant species. *PLoS One,* **7,** e47620.
https://doi.org/10.1371/journal.pone.0047620.

Monson, R.K. & Williams, G.J. (1982) A correlation between photosynthetic temperature
adaptation and seasonal phenology patterns in the shortgrass prairie. *Oecologia,* **54,** 58-
62. https://doi.org/10.1007/BF00541108.

Morales, L.V., Sierra-Almeida, A., Sandoval-Urzúa, C. & Arroyo, M.T.K. (2025) Thermal
decoupling may promote cooling and avoid heat stress in alpine plants. *Plants,* **14,** 2023.
https://doi.org/10.3390/plants14132023.

Muggeo, V.M.R. (2017) Interval estimation for the breakpoint in segmented regression: a
smoothed score-based approach. *Australian & New Zealand Journal of Statistics,* **59,**
311-322. https://doi.org/10.1111/anzs.12200.

Murchie, E.H. & Lawson, T. (2013) Chlorophyll fluorescence analysis: a guide to good practice
and understanding some new applications. *Journal of Experimental Botany,* **64,** 3983-
3998. https://doi.org/10.1093/jxb/ert208.

Neuner, G., Huber, B., Plangger, A., Pohlin, J.-M. & Walde, J. (2020) Low temperatures at
higher elevations require plants to exhibit increased freezing resistance throughout the
summer months. *Environmental and Experimental Botany,* **169,** 103882.
https://doi.org/10.1016/j.envexpbot.2019.103882.

Neuner, G. & Pramsohler, M. (2006) Freezing and high temperature thresholds of photosystem 2
compared to ice nucleation, frost and heat damage in evergreen subalpine plants.
Physiologia Plantarum, **126,** 196-204. https://doi.org/10.1111/j.1399-3054.2006.00605.x.

O'Sullivan, O.S., Heskel, M.A., Reich, P.B., Tjoelker, M.G., Weerasinghe, L.K., Penillard, A.,
Zhu, L., Egerton, J.J.G., Bloomfield, K.J., Creek, D., Bahar, N.H.A., Griffin, K.L., Hurry,
V., Meir, P., Turnbull, M.H. & Atkin, O.K. (2017) Thermal limits of leaf metabolism
across biomes. *Global Change Biology,* **23,** 209-223. https://doi.org/10.1111/gcb.13477.

Pearce, R.S. (2001) Plant freezing and damage. *Annals of Botany,* **87,** 417-424.
https://doi.org/10.1006/anbo.2000.1352.

Perez, T.M. & Feeley, K.J. (2020) Photosynthetic heat tolerances and extreme leaf temperatures.
Functional Ecology, **34,** 2236-2245. https://doi.org/10.1111/1365-2435.13658.

Perez, T.M. & Feeley, K.J. (2021) Weak phylogenetic and climatic signals in plant heat
tolerance. *Journal of Biogeography,* **48,** 91-100. https://doi.org/10.1111/jbi.13984.

Perez, T.M., Feeley, K.J., Michaletz, S.T. & Slot, M. (2021) Methods matter for assessing global
variation in plant thermal tolerance. *Proceedings of the National Academy of Sciences of*
the United States of America, **118,** e2024636118.
https://doi.org/10.1073/pnas.2024636118.

Querns, A., Wooliver, R., Vallejo-Marín, M. & Sheth, S.N. (2022) The evolution of thermal
performance in native and invasive populations of *Mimulus guttatus*. *Evolution Letters,* **6,**
136-148. https://doi.org/10.1002/evl3.275.

R Core Team (2023) *R: A language and environment for statistical computing*. R Foundation for
Statistical Computing, Vienna, Austria.

Robson, J.K., Ferguson, J.N., McAusland, L., Atkinson, J.A., Tranchant-Dubreuil, C., Cubry, P.,
Sabot, F., Wells, D.M., Price, A.H., Wilson, Z.A. & Murchie, E.H. (2023) Chlorophyll
fluorescence-based high-throughput phenotyping facilitates the genetic dissection of
photosynthetic heat tolerance in African (*Oryza glaberrima*) and Asian (*Oryza sativa*)
rice. *Journal of Experimental Botany,* **74,** 5181-5197.
https://doi.org/10.1093/jxb/erad239.

Rummukainen, M. (2012) Changes in climate and weather extremes in the 21st century. *Wiley*
Interdisciplinary Reviews: Climate Change, **3,** 115-129. https://doi.org/10.1002/wcc.160.

Sakai, A. & Larcher, W. (1987) *Frost survival of plants: Responses and adaptation to freezing*
stress. Springer-Verlag Berlin.

Scholz, F.G., Bucci, S.J., Arias, N., Meinzer, F.C. & Goldstein, G. (2012) Osmotic and elastic

adjustments in cold desert shrubs differing in rooting depth: coping with drought and

subzero temperatures. *Oecologia,* **170,** 885-897. https://doi.org/10.1007/s00442-012-

2368-y.

Schreiber, U. & Berry, J.A. (1977) Heat-induced changes of chlorophyll fluorescence in intact

leaves correlated with damage of the photosynthetic apparatus. *Planta,* **136,** 233-238.

https://doi.org/10.1007/BF00385990.

Seemann, J.R., Berry, J.A. & Downton, W.J. (1984) Photosynthetic response and adaptation to

high temperature in desert plants: a comparison of gas exchange and fluorescence

methods for studies of thermal tolerance. *Plant Physiology,* **75,** 364-368.

https://doi.org/10.1104/pp.75.2.364.

Seneviratne, S.I., Zhang, X., Adnan, M., Badi, W., Dereczynski, C., Di Luca, A., Ghosh, S.,

Iskandar, I., Kossin, J., Lewis, S., Otto, F., Pinto, I., Satoh, M., Vicente-Serrano, S.M.,

Wehner, M. & Zhou, B. (2021) Weather and Climate Extreme Events in a Changing

Climate. *Climate Change 2021: The Physical Science Basis. Contribution of Working*

Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate

Change (eds V. Masson-Delmotte, P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger,

N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R.

Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu & B. Zhou), pp. 1513–1766.

Cambridge University Press, Cambridge, UK and New York, NY, USA.

Shah, A.A., Woods, H.A., Havird, J.C., Encalada, A.C., Flecker, A.S., Funk, W.C., Guayasamin,

J.M., Kondratieff, B.C., Poff, N.L., Thomas, S.A., Zamudio, K.R. & Ghalambor, C.K.

(2021) Temperature dependence of metabolic rate in tropical and temperate aquatic

insects: Support for the Climate Variability Hypothesis in mayflies but not stoneflies.

Global Change Biology, **27,** 297-311. https://doi.org/10.1111/gcb.15400.

Sierra-Almeida, A., Reyes-Bahamonde, C. & Cavieres, L.A. (2016) Drought increases the

freezing resistance of high-elevation plants of the Central Chilean Andes. *Oecologia,*

181, 1011-1023. https://doi.org/10.1007/s00442-016-3622-5.

Siminovitch, D. & Cloutier, Y. (1983) Drought and freezing tolerance and adaptation in plants:

Some evidence of near equivalences. *Cryobiology,* **20,** 487-503.

https://doi.org/10.1016/0011-2240(83)90037-8.

Sklenář, P., Jaramillo, R., Wojtasiak, S.S., Meneses, R.I., Muriel, P. & Klimeš, A. (2023)

Thermal tolerance of tropical and temperate alpine plants suggests that 'mountain passes

are not higher in the tropics'. *Global Ecology and Biogeography,* **32,** 1073-1086.

https://doi.org/10.1111/geb.13678.

Slatyer, R.A., Umbers, K.D.L. & Arnold, P.A. (2022) Ecological responses to variation in

seasonal snow cover. *Conservation Biology,* **36,** e13727.

https://doi.org/10.1111/cobi.13727.

Sumner, E.E. & Venn, S.E. (2022) Thermal tolerance and growth responses to *in situ* soil water

reductions among alpine plants. *Plant Ecology & Diversity,* **15,** 297-308.

https://doi.org/10.1080/17550874.2022.2160674.

Sumner, E.E., Williamson, V.G., Gleadow, R.M., Wevill, T. & Venn, S.E. (2022) Acclimation to

water stress improves tolerance to heat and freezing in a common alpine grass.

Oecologia, **199,** 831-843. https://doi.org/10.1007/s00442-022-05245-1.

Sunday, J.M., Bates, A.E. & Dulvy, N.K. (2011) Global analysis of thermal tolerance and

latitude in ectotherms. *Proceedings of the Royal Society B: Biological Sciences,* **278,**

1823-1830. https://doi.org/10.1098/rspb.2010.1295.

Tattersall, G.J., Sinclair, B.J., Withers, P.C., Fields, P.A., Seebacher, F., Cooper, C.E. &

Maloney, S.K. (2012) Coping with thermal challenges: physiological adaptations to

environmental temperatures. *Comprehensive Physiology,* **2,** 2151-2202.

https://doi.org/10.1002/cphy.c110055.

Tserej, O. & Feeley, K.J. (2021) Variation in leaf temperatures of tropical and subtropical trees

are related to leaf thermoregulatory traits and not geographic distributions. *Biotropica,*

53, 868-878. https://doi.org/10.1111/btp.12919.

Urban, J., Ingwers, M.W., McGuire, M.A. & Teskey, R.O. (2017) Increase in leaf temperature

opens stomata and decouples net photosynthesis from stomatal conductance in *Pinus*

taeda and *Populus deltoides x nigra. Journal of Experimental Botany,* **68,** 1757-1767.

https://doi.org/10.1093/jxb/erx052.

Venn, S.E., Morgan, J.W. & Lord, J.M. (2013) Foliar freezing resistance of Australian alpine

plants over the growing season. *Austral Ecology,* **38,** 152-161.

https://doi.org/10.1111/j.1442-9993.2012.02387.x.

Wei, T. & Simko, V. (2021) R package 'corrplot': Visualization of a Correlation Matrix.

https://github.com/taiyun/corrplot.

Wooliver, R., Tittes, S.B. & Sheth, S.N. (2020) A resurrection study reveals limited evolution of

thermal performance in response to recent climate change across the geographic range of

the scarlet monkeyflower. *Evolution,* **74,** 1699-1710. https://doi.org/10.1111/evo.14041.

Zhou, Y., Kitudom, N., Fauset, S., Slot, M., Fan, Z., Wang, J., Liu, W. & Lin, H. (2023) Leaf

thermal regulation strategies of canopy species across four vegetation types along a

temperature and precipitation gradient. *Agricultural and Forest Meteorology,* **343,**

109766. https://doi.org/10.1016/j.agrformet.2023.109766.

Zhu, L., Bloomfield, K.J., Hocart, C.H., Egerton, J.J.G., O'Sullivan, O.S., Penillard, A.,

Weerasinghe, L.K. & Atkin, O.K. (2018) Plasticity of photosynthetic heat tolerance in

plants adapted to thermally contrasting biomes. *Plant, Cell & Environment,* **41,** 1251-

1262. https://doi.org/10.1111/pce.13133.

Figure legends

Fig. 1. Locations of study sites across three biomes in New South Wales, Australia. Desert sites

(red) in Gundabooka National Park (GNP), coastal temperate rainforest sites (green) in the

Illawarra Region (temperate), and alpine sites (blue) in Kosciuszko National Park (KNP). GNP

sites spanned open eucalypt woodland, mulga and chenopod shrublands between elevations of

100 to 224 m a.s.l. KNP sites encompassed feldmark, herbfields and sphagnum bog vegetation

formations between 1792 and 1855 m a.s.l. In the temperate environments, collections were

conducted across the Illawarra, including the tall open forest and rainforest communities between

100 to 400 m a.s.l. MAP = mean annual precipitation, MAT = mean annual temperature, MinT =

mean minimum temperature of the coldest month, MaxT = mean maximum temperature of the

warmest month, T_{range} = MaxT - MinT.

Fig. 2. Thermal tolerance thresholds and breadth across the three biomes. (A) Cold tolerance

threshold ($T_{crit-cold}$), (B) ice nucleation temperature ($T_{nucleation}$), (C) heat tolerance threshold (T_{crit-}

$_{hot}$) and (D) thermal tolerance breadth (TTB = $T_{crit-hot}$ - $T_{crit-cold}$), where desert has n = 24 species,

temperate has n = 23 species and alpine has n = 22 species. Small transparent points represent

individual replicates, and large solid points represent biome means ± SE. Lowercase letters inset

in each plot denote significant differences among biomes from post-hoc Tukey's Honest

Significant Differences tests within each thermal tolerance metric (see also Table S2).

Fig. 3. Mean thermal tolerance breadth (TTB = $T_{crit-hot}$ - $T_{crit-cold}$) for the alpine, temperate and

desert biomes, relative to the mean thermal range of the local environments (T_{range} = MaxT -

MinT). Red squares and lines represent desert, green triangles and lines represent temperate and

blue circles and lines represent alpine biomes. Cross (×) symbols represent the mean $T_{nucleation}$

and plus (+) symbols represent the mean annual temperature for each biome. Solid lines connect

the mean $T_{crit-cold}$ to the mean $T_{crit-hot}$ (large symbols) for each biome (length of line = TTB).

Dashed lines connect the mean MinT to the mean MaxT (small symbols) for each biome (length

of line = T_{range}). The light blue vertical dashed line represents 0 °C. Each species has n = 5-6

replicates for each thermal tolerance threshold value.

Fig. 4. Correlations and Principal Component (PC) composition of local climate variables (MAP

= mean annual precipitation, MAT = mean annual temperature, MinT = mean minimum

temperature of the coldest month, MaxT = mean maximum temperature of the warmest month,

T_{range} = annual thermal range (MaxT - MinT)). (A) Correlation matrix of local climate variables,

(B) Principal Components Analysis biplot of PC1 and PC2 showing biome position in climate

PC space, and contribution of individual local climate variables to (C) PC1 and (D) PC2.

Fig. 5. Local climate variables (Principal Components, PC1 and PC2; see Fig. 4) that predict

variation in thermal tolerance thresholds across all biomes. Regressions between (A) $T_{crit-cold}$ and

PC1, (B) $T_{crit-cold}$ and PC2, (C) $T_{nucleation}$ and PC1, (D) $T_{nucleation}$ and PC2, (E) $T_{crit-hot}$ and PC1, (F)

$T_{crit-hot}$ and PC2, (G) TTB and PC1 and (H) TTB and PC2. Arrows and annotations indicate

climate variable loadings on PC1 and PC2 axes: positive PC1 values = hotter and drier climate

(+MAT, –MAP, +MaxT); negative PC1 values = cooler and wetter climate (–MAT, +MAP, -

MaxT). Positive PC2 values = greater range in thermal extremes across the year and colder

extremes (+T_{range}, –MinT) negative PC2 values = greater annual thermal stability and less

extreme cold temperatures (–T_{range}, +MinT), see Fig. 4 and Table S5. Large solid points are

species-level means ± SE and small transparent points are individual replicates. Points are

coloured by biome and regressions are predicted model fits ± SE from the best models reported

in Table 1, where solid regression lines represent significant relationships.

Tables

Table 1. Results of linear mixed-effects regression models that tested the relationships between

local climate factors (summarised as Principal Components, PC1 and PC2) and thermal tolerance

thresholds.

Cold tolerance threshold: $T_{crit\text{-}cold}$ (°C)							
Fixed	**Coefficient ± SE**	*t*-**value**	*p*-**value**	**Random**	**Coefficient**	R^2 **type** [*]	R^2 **value**
Intercept	-10.488 ± 1.167	-8.985	**<0.001**	Growth form	1.308	R^2m	0.195
PC1	-15.179 ± 6.606	-2.299	**0.022**	Species	1.829	R^2c	0.515
PC1^2	-38.459 ± 15.449	-2.487	**0.013**	Phylogeny	0.081	$R^2c - R^2m$	0.320
PC2	0.451 ± 0.774	0.582	0.561	Residual	2.768		
Freezing tolerance: $T_{nucleation}$ (°C)							
Fixed	**Coefficient ± SE**	*t*-**value**	*p*-**value**	**Random**	**Coefficient**	R^2 **type** [*]	R^2 **value**
Intercept	-11.432 ± 0.585	-19.551	**<0.001**	Growth form	1.149	R^2m	0.430
PC1	-0.935 ± 0.205	-4.567	**<0.001**	Species	2.516	R^2c	0.725
PC2	-49.355 ± 7.135	-6.917	**<0.001**	Phylogeny	0.000	$R^2c - R^2m$	0.295
PC2^2	-15.707 ± 6.469	-2.428	**0.016**	Residual	2.670		
Heat tolerance threshold: $T_{crit\text{-}hot}$ (°C)							
Fixed	**Coefficient ± SE**	*t*-**value**	*p*-**value**	**Random**	**Coefficient**	R^2 **type** [*]	R^2 **value**
Intercept	46.360 ± 0.415	111.662	**<0.001**	Growth form	0.518	R^2m	0.250
PC1	1.318 ± 0.184	7.148	**<0.001**	Species	2.236	R^2c	0.446
PC2	-0.166 ± 0.328	-0.504	0.615	Phylogeny	0.000	$R^2c - R^2m$	0.196
				Residual	3.856		
Thermal tolerance breadth: TTB (°C)							
Fixed	**Coefficient ± SE**	*t*-**value**	*p*-**value**	**Random**	**Coefficient**	R^2 **type** [*]	R^2 **value**
Intercept	57.440 ± 0.928	61.920	**<0.001**	Growth form	2.035	R^2m	0.273
PC1	65.441 ± 10.543	6.207	**<0.001**	Species	3.146	R^2c	0.550
PC1^2	49.280 ± 24.352	2.024	**0.044**	Phylogeny	0.000	$R^2c - R^2m$	0.276
PC2	-1.102 ± 1.216	-0.906	0.365	Residual	4.789		

Note: PC1 represents an axis from cooler and wetter climate (negative values) to hotter mean and extreme
temperatures and drier climate (positive values). PC2 represents an axis from a climate with less extreme
cold temperatures and greater thermal stability through the year (negative values) to a climate with colder
extremes and greater range in thermal extremes across the year (positive values). [*]R^2m represents the
variance explained by all fixed effects, R^2c represents the variance explained by all fixed and random
effects, and $R^2c - R^2m$ represents the variance explained by all random effects.

Temperate – Illawarra Region

MAP = 1285mm
MAT = 16.5°C
MinT = 7.4°C
MaxT = 24.6°C
Trange = 17.2°C

N

Desert – Gundabooka NP

0 2.5 5 km

MAP = 332mm
MAT = 20.2°C
MinT = 4.5°C
MaxT = 36.0°C
Trange = 31.5°C

0 2.5 5 km

Alpine – Kosciuszko NP

MAP = 1764mm
MAT = 4.5°C
MinT = -5.2°C
MaxT = 16.5°C
Trange = 22.7°C

0 2.5 5 km

0 25 50 km