Drivers of thermal tolerance breadth of plants across contrasting biomes: do mean or seasonality in climate indices matter more?

Verónica F. Briceño[1], Alicia M. Cook[2], Stephanie K. Courtney Jones[1,3], Pieter A. Arnold[1], Rachael V. Gallagher[4], Kris French[5], León A. Bravo [6], Adrienne B. Nicotra[1] and Andrea Leigh[2]

[1]Division of Ecology and Evolution, Research School of Biology, Australian National University, Canberra, ACT, Australia.

[2]School of Life Sciences, Faculty of Science, University of Technology Sydney, Broadway, NSW, Australia.

[3]Ecosystem and Threatened Species, Northwest Branch, NSW Department of Planning and Environment, Dubbo, NSW, Australia

[4]Hawkesbury Institute for the Environment, Western Sydney University, Sydney, Australia.

[5]Centre for Sustainable Ecosystem Solutions, School of Earth, Atmospheric and Life Sciences, University of Wollongong, Wollongong, Australia.

[6]Departamento de Ciencias Agronómicas y Recursos Naturales, Facultad de Ciencias Agropecuarias y Medioambiente, Universidad de la Frontera, Región de la Araucanía, Temuco, Chile.

Corresponding author: Andy Leigh: andrea.leigh@uts.edu.au ORCiD 0000-0003-3568-2606

ORCiD details:
Verónica F. Briceño
Alicia M. Cook 0000-0003-3594-3220
Stephanie K. Courtney Jones 0000-0002-0905-0624
Pieter A. Arnold 0000-0002-6158-7752
Rachael V. Gallagher 0000-0002-4680-8115
Kris French 0000-0001-6938-2017
León A. Bravo 0000-0003-4705-4842
Adrienne B. Nicotra 0000-0001-6578-369X
Andrea Leigh 0000-0003-3568-2606

Figures x 6
Tables x 2
Supplementary tables x 3

Abstract

1. The Climate Variability Hypothesis (CVH) predicts that species from environments with more variable temperatures should have *wide* thermal tolerance breadth. We addressed this question in plants and asked which climate predictors (e.g., temperature, precipitation, variability/seasonality) best predict species thermal limits.

2. Measures of low ($T_{crit-cold}$) and high ($T_{crit-hot}$) photosystem II thermal thresholds were used to determine thermal tolerance breadth (TTB), along with ice nucleation temperature (NT) of 69 plant species in alpine, desert and temperate biomes.

3. Tolerance metrics $T_{crit-cold}$, NT and $T_{crit-hot}$ all differed among biomes. Notably, desert species had the most cold and heat tolerant leaves with the widest TTB. Alpine and temperate biomes had similar TTB and each biome TTB exceeded their climate thermal ranges. Overall, climate drivers explained the most variation (~50%) in TTB and NT, with species a second strong predictor. The climate variables best explaining tolerance differed for cold and heat: species from more variable (higher temperature seasonality) environments (alpine and desert) were more cold tolerant, whereas mean annual temperature (MAT) was the best predictor of $T_{crit-hot}$ with species in higher MAT environments having higher heat tolerances. TTB and NT were explained by both seasonality and MAT. Unexplained variation could be due to microclimate-driven plasticity, leaf traits or thermoregulatory mechanisms.

4. Our results provide some support for the CVH. Depending on the thermal limits examined, climate means and seasonality remain important predictors of plant thermal tolerance.

Introduction

Desert and alpine plant species live at the thermal limits of biological processes. Unlike those in

temperate regions, plants living in extreme environments are exposed to both very low and very

high temperatures and thus to wide thermal range and highly variable thermal regimes, both

among days and seasons. Given the important influence of temperature in shaping species

distributions (Tattersall *et al.*, 2012; Garcia-Pichel *et al.*, 2013) the innate thermal tolerance

limits (lowest and highest temperature that a species can withstand) and thermal tolerance

breadths (range of temperature between upper and lower thermal tolerance limits) of species are

predicted to correlate with the temperature variation characterising their environment. The

climate variability hypothesis (CVH; Janzen, 1967) predicts that species with *wide* thermal

tolerance ranges are likely to evolve in environments with *variable* temperatures compared to

thermal specialists with *narrow* tolerance ranges that evolved in thermally *stable* environments.

This hypothesis is a cornerstone of thermal ecology, but has to date been primarily explored in

animals (Addo-Bediako *et al.*, 2000; Compton *et al.*, 2007; Calosi *et al.*, 2010; Baudier *et al.*,

2018).

Cold and heat tolerance thresholds are rarely studied concomitantly. Cold tolerance studies

predominate in alpine environments where plants can withstand very low temperatures and

tolerate extracellular ice formation and the resulting cytoplasmic dehydration (Sakai & Larcher,

1987; Larcher, 2003). Heat tolerance studies, historically, are primarily focussed on warm and

desert environments (Geange et al., 2021). Mechanisms for tolerating extreme heat have been

demonstrated in a range of desert species, including heat avoidance through increased

transpiration (Aparecido *et al.*, 2020) with microhabitat conditions setting the scene for heat

tolerance (Curtis *et al.*, 2016), whereas in other cases, these differences disappear under common

garden conditions (Downton *et al.*, 1984; Knight & Ackerly, 2002; Knight & Ackerly, 2003). In

spite of great progress, a recent systematic review revealed that less than 5% of more than 3000

studies examining thermal tolerance of plants considered both heat and cold tolerance thresholds

together (Geange *et al.*, 2021). The current focus on either cold or heat tolerance overlooks the

fact that plants in extreme biomes can experience both cold and heat stress.

Temperature increases associated with climate change have motivated an increase in studies on

plant heat tolerance, particularly in relation to climate variability and the occurrence of aseasonal

extreme events (Rummukainen, 2012; Seneviratne *et al.*, 2012). However, *in situ* field sampling

studies rarely encompass multiple biomes or a broad range of thermal environments across
landscapes. There remains a critical gap in our understanding of how plants in extreme *vs.*
benign environments tolerate both heat and cold stress *in situ*, which limits our capacity to
predict where the most thermally sensitive species occur.

Differing patterns of climate among contrasting biomes provide an opportunity to explore the
role of climate variability in upper and lower thermal thresholds and on thermal tolerance
breadth. Of the few studies that have compared thermal thresholds among biomes, Smillie and
Nott (1979) found that heat tolerance was lower in alpine species, followed by temperate and
tropical species. In a rare study considering both upper and lower thermal thresholds together,
Cunningham and Read (2006) show that, in common garden conditions, temperate origin species
have lower heat and higher cold tolerance than tropical species. Meta-analytical studies across a
wide range of species and biomes suggest that heat tolerance decreases and cold tolerance
increases with latitude; however, these shifts in tolerance do not keep pace with an observed
latitudinal temperature gradient (Lancaster & Humphreys, 2020). Further, predictors of thermal
tolerance variation with latitude may include mean temperature of the warmest month and mean
annual temperature (O'Sullivan *et al.*, 2017; Lancaster & Humphreys, 2020), though threshold-
based predictors (e.g. number of days over 35 degrees per year; Maher *et al.*, 2019) are
increasingly being explored as drivers of heat tolerance (Mora *et al.*, 2015). Given that water
status influences plant heat tolerance (Curtis *et al.*, 2016; Marchin *et al.*, 2022) and freezing
tolerance (Sierra-Almeida *et al.*, 2016), precipitation parameters are also likely to affect thermal
thresholds. In addition, upper and lower thermal thresholds can vary in relation to the methods
used (Lancaster & Humphreys, 2020; Perez *et al.*, 2021), influencing the ability of large-scale
meta-analyses to capture such sources of variation and likely limiting their power to reach
generalisable conclusions about any association between thermal limits and climate variables.
Finally, although we have evidence that climate variables influence heat and cold thermal
thresholds independently, the climate predictors of the difference between these thresholds,
thermal tolerance breadth, are unknown. If we can reveal general patterns underlying variation
and the climate drivers of physiological thermal tolerance thresholds among co-occurring species
within and among biomes, we can strengthen predictions of which and where species are likely
to be most vulnerable to changing thermal conditions.

Here we assess leaf-level thermal tolerance breadth (TTB) of plant species in extreme and benign

environments and the climate drivers of these thresholds to test the CVH. We used chlorophyll

fluorescence to determine the low and high photosystem II (PSII) thermal thresholds and thermal

tolerance breadth of 22 alpine, 24 desert and 23 temperate rainforest species (temperate

hereafter) during the growing season. We also determined ice nucleation temperatures to

examine freezing tolerance in all species. To determine relationships between climate variability

and physiological thresholds, we extracted long term climate parameters for each sampling

locality. First, we hypothesised that the physiological thermal limits will be skewed towards the

extremes of the environments; colder temperatures in the alpine; and warmer temperatures in the

desert respectively, with temperate species having milder cold and heat tolerance thresholds.

Second, we tested if species from more thermally variable biomes have high thermal tolerance to

both heat and cold, and thus wider thermal tolerance breadth, in accordance with the CVH.

Thirdly, we assessed whether the variation in tolerance thresholds and TTB is explained by

climate variability (e.g., seasonality), supporting the CVH, rather than mean climate values.

Materials and Methods

Study locations

Sites used in this study were designated to one of three biomes in New South Wales,

Australia: alpine (Kosciuszko National Park), desert (Gundabooka National Park), and wet

temperate (Illawarra Region: Royal National Park, Illawarra State Conservation Area and Bass

Point Conservation Area) (**Fig. 1**).

Species selection

A total of 69 species (22 alpine, 24 desert and 23 wet temperate species; **Table S1**) were selected

to encompass a representative sample of growth forms and plant families and where possible,

included congeneric and/or confamilial species between two or more biomes. We included

community dominants and a range of less abundant species to represent a cross section of species

occupying each biome. Species replicates were sampled across days within each sampling period

to incorporate daily variation in thermal tolerance thresholds.

Fig. 1: Locations of study sites across three biomes in NSW, Australia. Desert sites (red) in Gundabooka National Park (GNP), temperate sites (green) in the Illawarra Region, and alpine sites (blue) in Kosciuszko National Park (KNP). GNP sites span open eucalypt woodland, mulga and chenopod shrublands between elevations of 100 to 224 m a.s.l. KNP sites encompassed feldmark, herbfields and sphagnum bog vegetation formations between 1792 and 1855 m a.s.l. In the wet temperate environments, collections were conducted across the Illawarra Subtropical Rainforest including the Tall Open Forest and Rainforest communities between 100 and 400 m a.s.l. MAT = mean annual temperature, MAP = mean annual precipitation.

Leaf sample collection

Mature, fully expanded leaves were harvested from five to seven individuals per species. For shrubs and trees, small branches were collected from the northern facing side of the plant at a height of 1-1.5 m. For smaller growth forms, entire individuals or rosettes were collected. Samples were kept in cool sealed plastic bags with moistened paper towel until they were processed (approx. 1-2 hours after collection). In the field laboratory, leaves (size ranging 0.25 cm^2 to 5 cm^2) were cut (or combined) to fit an approximate area of 1 cm^2 and kept on wet florist foam in humid dark containers until thermal tolerance assays were performed. Measurements occurred during January 2020, March 2020 and December 2020 for the desert, wet temperate and

98   alpine biomes, respectively. Sampling periods were selected to capture peak growing season in
99   each region.

Thermal tolerance assays

The photosynthetic machinery, specifically photosystem II (PSII), embedded in the thylakoid
membrane has long been recognised as a thermally sensitive structure (Schreiber & Berry, 1977;
Berry & Bjorkman, 1980; Seemann *et al.*, 1984). The temperature at which the PSII are
disrupted causes a rise in minimal chlorophyll a fluorescence (F_0) as a leaf is heated or cooled
(Schreiber & Berry, 1977; Neuner & Pramsohler, 2006). This critical temperature indicates a
threshold beyond which physiological and photochemical systems become increasingly impaired
and damage can occur, if temperatures are sustained. Freezing tolerance is tied to the
temperatures at which ice formation occurs (ice nucleation, IN), the measuring of which is
required to understand whether temperature dependent damage to photosynthetic machinery
occurs in association with ice formation.

Lower and upper thresholds of PSII ($T_{crit-cold}$ and $T_{crit-hot}$)

The critical low and high temperature thresholds of PSII ($T_{crit-cold}$ and $T_{crit-hot}$) were measured
using T- F_0 curves, i.e., the rise in F_0 (Neuner & Pramsohler, 2006; Arnold et al., 2021) with set
heating or cooling rates. One leaf per individual (or several small leaves abutting one another)
was used for each of the heating and cooling assays with n = 5-7 replicates per species. For each
assay run, 45 samples were secured with double-sided tape to paper on a peltier plate (CP-
121HT; TE-Technology, Inc., Michigan, USA), used to cool or heat samples at a set rate using
LabView-based control software (National Instruments, Texas, USA) and adapted code from
TE-Technology, Inc. A vacuum sealed double-glazed glass was placed on top of the leaf samples
on the plate to avoid water condensation, particularly at freezing temperatures, and compress
samples to ensure maximum leaf contact with the Peltier plate (see Arnold *et al.*, 2021 for more
details).

Chlorophyll fluorescence was measured with a Pulse Amplitude Modulated (PAM) chlorophyll
fluorescence imaging system (Maxi-Imaging-PAM; HeinzWalz GmbH, Effeltrich, Germany)
mounted above the Peltier plate. Leaves were dark-adapted first and the PSII photochemistry
maximum quantum efficiency (F_V/F_M) measured to check starting function of leaves before
another dark adaption for 15min prior to assay. Circular areas of interest were created in the

middle of leaf samples avoiding edges for F_0 measurements using the Maxi-Imaging-PAM

software. During temperature ramping, F_0 was measured every 20 s with a weak blue low

frequency (1 Hz) pulse modulated measuring light (0.5 μmol photons $m^{-2} s^{-1}$). Leaf temperatures

were simultaneously measured with 40-gauge type-T thermocouples (OMEGA Engineering,

Singapore) attached to undersides of leaves and logged with a multi-channel DT85 Datataker

datalogger (Lontek, Australia). For hot T-F_0 measurements, samples were heated to 60°C at a

rate of 30°C h^{-1} (0.5°C/min). For cold T-F_0 measurements, the leaves were cooled down to -20°C

or -25°C when logistically possible, at a rate of 15°C h^{-1} (0.25°C/min). Temperature rates were

selected as moderate temperature changes per Arnold et al., (2021). The starting temperature of

each heating/cooling assay depended on average field lab temperature, generally 15°C for alpine

and temperate, and 20°C for desert measurements.

$T_{crit-cold}$ and $T_{crit-hot}$ were determined as inflection points on the T- F_0 curves, using the *segmented*

R package. For further description of T- F_0 curve see Arnold et al., (2021). $T_{crit-cold}$ and $T_{crit-hot}$

were used to represent the lower and upper bounds of the PSII thermal tolerance breadth, TTB

(TTB = $T_{crit-hot}$ - $T_{crit-cold}$).

Ice nucleation temperatures

The ice nucleation temperature was estimated by analysing the freezing exotherms of leaf tissue,

using the leaf temperature data collected during $T_{crit-cold}$ measurements. Temperature data

collected for each leaf sample was plotted against time to visualize the freezing exotherms (small

peak in temperature). The temperature at which the exotherm began was recorded as the

nucleation temperature (Larcher, 2003).

Climate variables for sampling locations

Climate data were obtained from gridded datasets at 1 km resolution based on spatial

interpolations of long-term (1981-2010) conditions accessed from the CHELSA v2.1 database

(Karger *et al.*, 2017). For each species, a suite of climate variables including means, minimum,

maximum and seasonality variables for temperature and precipitation were extracted for each

species at leaf collections sites (see **Table S2** for complete list and sampling location

coordinates).

Statistical analyses

To test the Climate Variability Hypothesis ability to predict photosystem thermal tolerance, we examined climate by two designations. First, as a-prior categorical variable 'biome' (aims 1 & 2), and secondly, by a comprehensive selection of continuous climate variables (aim 3).

Differences in thermal tolerance metrics among biomes were tested with Linear Mixed Models (LMMs, Bates *et al.*, 2015) from the *lme4* package and post-hoc Tukey's Honest Significant Differences were used to compare biome means. Models included biome (alpine, desert and temperate) as fixed terms and species as a random term. Growth form as a random effect did not significantly improve models thus was excluded. Because the aim of this study was to test the Climate Variability Hypothesis in relation to thermal tolerance limits and TTB, other variables such as local air and leaf temperatures or finer scale environmental characteristics such as aspect were not included in models. Plant evolutionary history was considered in analysis, see below.

LMMs were also used to determine the relative contribution of climate variables to the variation in each thermal tolerance metric. For the contribution of climate variables, LMMs included the climate variables as fixed terms and species as a random term. Several of our climate variables were strongly correlated and to avoid collinearity, we excluded some of those variables. All full models included mean annual temperature (MAT, °C), mean annual precipitation (MAP, mm), temperature seasonality (BIO4, standard deviation of the monthly mean temperatures) and precipitation seasonality (standard deviation of the monthly precipitation estimates expressed as a percentage of the annual mean of those estimates). The best models were selected by backward selection using the function *step* from the *emdi* package (Kreutzmann *et al.*, 2019). The package *lmerTest* (Kuznetsova *et al.*, 2017) was used with *lme4* to calculate LMM degrees of freedom and *P*-values . The *MuMIn* package (Johnson, 2014) was used to calculate the variance explained by fixed effects, (marginal R^2, mR^2) and the variance explained by both fixed and random effects (conditional R^2, cR^2). The Intraclass Correlation Coefficient (ICC) describes the proportion of the total variance in the data that is explained by the random effects alone.

Prior to model fitting we assessed if the shared evolutionary history across our 69 species may also explain differences in thermal tolerance measures. We generated a phylogenetic tree using the R package *V.phylomaker* (Jin & Qian, 2022) and calculated the phylogenetic signal and statistical significance for each $T_{crit-cold}$, $T_{crit-hot}$, TTB and nucleation temperature using Pagel's λ

and Blombergs's K (**Table S3**) with the *phylosignal* package (Keck et al., 2016). No evidence of

phylogenetic signal was present and thus phylogenetic non-independence was not detected

therefore, we did not include the phylogenetic structure in any of our models. All analyses and

visualizations were carried out in R version 3.6.1 (R Core Team, 2018).

Results

Biome differences in thermal tolerance

There were significant differences among biomes in all the individual thermal tolerance metrics

measured on plant species (**Fig. 2a-d**, **Table 1**). Unexpectedly, desert species exhibited the most

extreme cold thresholds (-13.3 ± 0.5°C), followed by alpine (-11.0 ± 0.5°C) and then temperate

species (-8.8 ± 0.5°C), all of which differed significantly from one another (**Fig. 2a**, **Table 1**). As

predicted, desert species had significantly higher heat tolerance (49.3 ± 0.6°C), followed by

temperate (46.6 ± 0.6°C), then alpine (43.1 ± 0.6°C) species (**Fig. 2b**, **Table 1**). Similar to $T_{crit\text{-}cold}$

values, nucleation temperatures were coldest in desert species (-16.0 ± 0.6°C), followed by

alpine (-10.8 ± 0.6°C) and warmest in temperate species (-7.5 ± 0.6°C); again, these differed

significantly from one another (**Fig. 2c**, **Table 1**). Interestingly, $T_{crit\text{-}cold}$ values for alpine and

temperate species were very close to their nucleation temperature, whereas in desert species, ice

formation occurred at markedly lower temperatures than $T_{crit\text{-}cold}$ (**Fig. 2a, c**).

Fig. 2: Heat (T$_{crit-hot}$) and cold (T$_{crit-cold}$) thresholds, ice nucleation temperatures (NT) and thermal tolerance breadth (TTB) for desert (n = 24 spp), temperate (n = 23 spp) and alpine (n = 22 spp) biomes. Circles represent individual replicates, triangles represent biome means across species; letters denote significant differences between biomes tested using linear mixed models and post-hoc Tukey's Honest Significant Differences test for comparing means with significance at 0.05 alpha level.

At a species level, temperate species *Oplismenus hirtellus* was the least cold tolerant (-4.0°C,
T$_{crit-cold}$), whereas two alpine species *Epacris paludosa* and *Hovea montana* were the most cold
tolerant (-18.5°C T$_{crit-cold}$) and the desert species *Atriplex stipitata* had the most extreme
nucleation temperature (-20.5°C). The alpine species *Pimelea ligustrina* was the least heat
tolerant (38.1°C, T$_{crit-hot}$) and the desert species *Rhagodia spinescens* was the most heat tolerant
(T$_{crit-hot}$ = 54.9°C, **Fig. 3**, **Table S1**).

Fig. 3: Thermal tolerance breadth (TTB = T$_{crit-hot}$-T$_{crit-cold}$) for all the studied species. Species are sorted top to bottom from widest to narrowest TTB. Red circles and lines represent desert species (n = 24 spp), green triangles and lines represent wet temperate species (n = 23 spp) and blue squares and lines represent alpine species (n = 22 spp). Black horizontal lines represent the standard error. The vertical dashed line represents 0°C. Each TTB, T$_{crit-hot}$ and T$_{crit-cold}$ value is from n = 5-6 replicates per species.

Table 1: Linear mixed model output for all thermal limits considering biome as a fixed effect and species as random effect. Model biome mean estimates were significant with P values of <0.001.

Biome	$T_{crit-hot}$		$T_{crit-cold}$		NT		TTB	
	Estimate	Std.Error	Estimate	Std.Error	Estimate	Std.Error	Estimate	Std.Error
Alpine	43.1	0.6	-11.0	0.5	-10.8	0.6	54.1	0.9
Desert	49.3	0.6	-13.3	0.5	-16.0	0.6	62.6	0.8
Temperate	46.6	0.6	-8.8	0.5	-7.5	0.6	55.4	0.9
	Variance		Variance		Variance		Variance	
Species	5.0		5.2		7.4		12.3	
Residual	14.9		7.7		7.1		23.0	

The thermal tolerance breadth (TTB) was significantly wider in desert species (62.6 ± 0.8°C)

than temperate (55.4 ± 0.9°C) or alpine species (54.1 ± 0.6°C) (**Fig. 2d**, **Table 1**). However,

there was also considerable variation among species within biomes (**Fig. 3**). Within the desert

biome TTB was more similar across species (lower variability), (56.2 to 70.3°C TTB, Δ14.1°C),

compared to the alpine biome which had the greatest variation among species (45.9 to 63.1°C

TTB, Δ17.2°C) followed by the wet temperate species (44.8 and 60.4°C TTB, Δ15.6°C). For

species from all three biomes, TTB was much wider than the extreme temperature range

calculated for the sampling sites, thus most thermal thresholds lay well beyond the minimum and

maximum temperatures expected for each species in their local environment. The only notable

exception was for the alpine biome species, where the cold threshold was closer to the minimum

temperatures representing our alpine sites (**Fig. 4**).

Fig. 4: Mean thermal tolerance breadth (TTB, solid lines) for alpine, temperate and desert species and their respective mean thermal range (T_{range} = mean MaxT- mean MinT based on the locations of the sample sites, see **Table S2**). Black lines represent standard errors.

Contribution of climate variables to the variation on thermal tolerance metrics

We investigated the relationship between climate variables at the site of plant sampling and each
of the thermal tolerance traits using backward model selection to identify the most parsimonious
linear mixed models. None of the precipitation variables included in our models explained
considerable variation in thermal tolerance thresholds. Whereas the best models included either
mean annual temperature or temperature seasonality or both.

$T_{crit-cold}$ was best explained by temperature seasonality (19% of the variation in $T_{crit-cold}$), although
a much larger proportion of the variation in $T_{crit-cold}$ was explained by species (32%, **Table 2**).
There was a negative relationship between $T_{crit-cold}$ and temperature seasonality, indicating that
species from environments with greater seasonality, our alpine and desert sites, had a more
extreme $T_{crit-cold}$ than species from the less seasonal wet temperate rainforest (**Fig. 5a**). By
contrast, rather than seasonality, $T_{crit-hot}$ was best explained by mean annual temperature (MAT,
23%), with a much smaller proportion of the variation in this variable explained by species
(19%, **Table 2**). There was a positive relationship between $T_{crit-hot}$ and site MAT (**Fig. 5b**), such
that species from, on average, warmer environments, also had higher $T_{crit-hot}$.

Fig. 5: Climatic variables that best predicted the variation in thermal thresholds across all biomes. Environmental variables are temperature seasonality (°C) and mean annual temperature (MAT, °C). Best models included: for $T_{crit-cold}$, seasonality; for $T_{crit-hot}$, MAT; for TTB both seasonality and MAT. See **Table 2** for statistics. Solid circles are species means, open circles are individual replicates which are jittered around mean x axis points.

TTB was best predicted by seasonality and mean annual temperature (**Table 2**), although a
similar proportion of the variance was explained by the species effect compared to climate
variables (29% and 24% respectively). There was a positive relationship between TTB and
seasonality as well as with mean annual temperature (**Fig. 5c,d**). Notably, TTB was wider in
species from desert biomes, which experience considerably greater seasonality and MAT than
those in alpine and temperate environments.

Table 2: Best models predicting climatic drivers of physiological thermal limits across all species. Linear mixed model fit using a restricted maximum likelihood approach and t-tests using Satterthwaite's method. Climatic conditions at the sites where leaf samples were collected were included as fixed effects and species as a random effect. These models were selected using a backward model selection using the function *step* from the *emdi* package in R. Along with species, mean annual temperature (MAT) and temperature seasonality were significant predictors in our models.

$T_{crit-cold}$ ~ Seasonality + (1|Species)								
	Estimate	**SE**	**t-value**	**Df**	**P-value**	**mR²**	**ICC**	**cR²**
Intercept	-4.20	1.21	-3.48	70	<0.001			
Seasonality	-1.41	0.24	-5.82	70	<0.001			
	Variance			**N**		0.19	0.32	0.52
Species	5.25			69				
Residual	7.66	2.76						
$T_{crit-hot}$ ~ MAT + (1|Species)								
	Estimate	**SE**	**t-value**	**Df**	**P-value**	**mR²**	**ICC**	**cR²**
Intercept	41.00	0.81	50.24	61	<0.001			
MAT	0.38	0.05	7.22	63	<0.001			
	Variance			**N**		0.23	0.19	0.43
Species	5.22			69				
Residual	14.87							
TTB ~ Seasonality + MAT + (1|Species)								
	Estimate	**SE**	**t-value**	**DF**	**P-value**	**mR²**	**ICC**	**cR²**
Intercept	42.39	2.11	20.06	67	<0.001			
Seasonality	1.88	0.07	4.91	68	<0.001			
MAT	0.42	0.07	5.51	64	<0.001			
	Variance			**N**		0.29	0.24	0.53
Species	13.14			69				
Residual	22.94							
NT ~ MAT + Seasonality + (1|Species)								
	Estimate	**SE**	**t-value**	**DF**	**P-value**	**mR²**	**ICC**	**cR²**
Intercept	3.15	1.52	2.06	65	0.043			
MAT	-0.14	0.05	-2.68	66	0.009			
Seasonality	-2.60	0.27	-9.38	65	<0.001			
	Variance			**N**		0.45	0.27	0.72
Species	7.22			69				
Residual	7.12							

249   Unlike the cold PSII thermal threshold, both mean annual temperature (MAT) and seasonality
250   explained a considerable proportion of variation (45%) among ice nucleation (NT) temperatures
251   while species had comparatively less of an effect (27%) (**Table 2**). There was a negative
252   relationship between nucleation temperature and temperature seasonality, indicating that species

from environments with higher seasonality froze at lower temperatures than species in environments where temperature is more stable (**Fig. 6a**). There was also a negative relationship between nucleation temperature and MAT, where desert species froze at more extreme temperatures than most species in colder climates (**Fig. 6b**).

Fig. 6: Climatic variables (temperature seasonality and mean annual temperature (MAT)) that best explained the variation in ice nucleation temperature (NT), see **Table 2** for statistics.

Discussion

The climate variability hypothesis (CVH) predicts that species will evolve wider thermal tolerance breadths in environments with more variable temperatures and thermal specialisation (narrow TTB) in thermally stable environments, a concept largely supported in animals (i.e Compton *et al.*, 2007; Sunday *et al.*, 2011). Of the few studies testing this hypothesis on plants, investigations have included phenotypic plasticity increases with latitude (Molina-Montenegro & Naya, 2012), species distribution ranges with seasonality across elevation (Mumladze *et al.*, 2017) and the breadth of photosynthetic thermal response with environmental temperature variability (Perez *et al.*, 2023). Here we explored the CVH by sampling for thermal tolerance metrics including TTB on species growing in three contrasting biomes, two extreme and more variable, one benign and more thermally stable.

Using the CVH framework, we hypothesised that the thermal tolerance limits of alpine and desert species would be skewed to their respective habitat extremes (aim 1). These more skewed

tolerances in the thermally extreme environments would also drive wider TTB than the more
benign temperate thermal environment (aim 2). We found that variation in tolerance breadth in
the more seasonal desert and alpine environments was indeed wider than temperate species, with
desert TTB values skewed towards higher temperatures, alpine TTB values skewed towards
colder temperatures and wet temperate TTB sitting in between. However, the mean tolerance
breadths of alpine and temperate species were not statistically different, while desert species had
both greater heat and cold thresholds. When looking at the climate drivers of tolerance breadth,
we found that not only thermal seasonality, but also mean annual temperature and predicted
TTB, providing some, but not conclusive support for the CVH (aim 3). When we explored the
climate drivers of thermal thresholds, we found that temperature seasonality, our proxy for
climate variability, and MAT both contributed to explaining the variation in TTB: seasonality
best explained the variation in $T_{crit-cold}$, whereas MAT was the strongest predictor of $T_{crit-hot}$.
Below we explore our findings in terms of physiology and the ecological implications of these
results for species persistence in a changing climate.

TTB exceeds thermal ranges in each biome and varies within biomes

Thermal tolerance breadth values reported here are notably wide; wider than the thermal ranges
in each environment and larger than TTB in other organisms. For instance, across a range of
marine and terrestrial ectotherm taxa, the widest recorded TTB was 60°C in terrestrial arthropods
(Sunday *et al.*, 2011), while the widest TTB in our dataset was 70°C in the desert species
Rhagodia spinescens, with several other desert species also exceeding 60°C. Within each biome,
we found species with a range of wide and narrow TTB (**Fig. 3**). For example, several alpine
species across growth forms had relatively narrow TTB, such as the snow bed species
Psychrophila introloba (45°C TTB) and the treeline species *Eucalyptus pauciflora* subsp.
niphophila (47°C TTB). The narrow TTB of some alpine species is mainly driven by their low
heat tolerance ($T_{crit-hot}$ = around 40°C), while other co-occurring species (*Hovea montana* and
Epacris paludosa) had TTB similar to desert species (around 63°C) driven by extreme cold
tolerance (around -18°C). This within-biome variation in TTB might be due to variation in
microclimate 'preferences' of species within biomes, which vary depending on species growth
forms, topography and different transpirational cooling limitations (Curtis *et al.*, 2016; Curtis *et*
al., 2019; Aparecido *et al.*, 2020).

Although thermal seasonality was greater in alpine than temperate regions sampled here, the

average TTB for these two biomes is remarkably similar (**Fig. 2d**). Temperate species can

tolerate colder temperatures than expected, and it is possible that their wide thermal tolerance is a

legacy of exposure to climate extremes through geological time (Byrne *et al.*, 2008). The climate

legacy effect on thermal tolerance seems to be stronger for cold than heat tolerances (Bennett *et*

al., 2021) and might explain why some temperate and desert species have greater freezing

tolerance than expected by their current climate. Although metrics based on air temperatures for

our alpine biome (e.g., MAT, temperature seasonality) confirm that thermal range and

seasonality are high in this biome, mountain plants spend 4-5 months under the insulating cover

of snow, which stabilizes temperatures (Briceño *et al.*, 2014). This dampening of climate

extremes might contribute to the similarity between alpine and temperate species TTB.

Heat and cold thermal limits vary among biomes

Plants exposed to drought and/or freezing share a similar physiological response: cell

dehydration (Anisko & Lindstrom, 1996; Blake & Hill, 1996; Lintunen *et al.*, 2013). The

physiological response to drought conditions may partially explain our somewhat

counterintuitive result that desert species were more freezing tolerant (more extreme $T_{crit-cold}$ and

nucleation temperature) than either alpine or temperate species. Plants can use osmotic

adjustments to offset water loss from cells along the water potential gradient caused by both

drying and freezing (Siminovitch & Cloutier, 1983; Larcher, 2003). Thus, the ability of desert

species to cope with dehydration induced by long periods of drought might also confer the ability

to cope with freezing, despite the fact that these desert species are unlikely to ever be exposed to

such extreme freezing temperatures. In addition, leaf adaptations to aridity such as smaller

leaves, thick cuticles, smaller, more densely packed cells, thicker cell walls with small capillary

structures and small xylem vessel diameter (Dörken *et al.*, 2020) also can prevent ice formation,

which increases supercooling capacity and freezing tolerance (Lintunen *et al.*, 2013). This may

reflect that species living in environments where freezing temperatures are rare use supercooling

as a mechanism to avoid freezing damage, while in alpine environments species tolerate ice

formation. Indeed, the desert species studied here generally had nucleation temperatures much

lower than their $T_{crit-cold}$, from which we can infer a high supercooling capacity that effectively

serves to delay freezing. Indeed, Andean alpine species' low ice nucleation temperatures at the

end of the growing season are attributed to low soil moisture, thus potential water stress, rather

than low temperatures.(Sierra-Almeida *et al.*, 2016). Likewise, drought exposure can influence

heat tolerance through raising intracellular sugar contents which stabilize thylakoid membranes,

increasing heat tolerance (Huve *et al.,* 2006; Brestic *et al.,* 2012). The extreme freezing tolerance

and high heat tolerances found in our desert species might reflect a secondary outcome of plant

adaptation to aridity.

The relatively high heat tolerance of some of our alpine species is at first counterintuitive, as

alpine areas are conventionally seen as primarily cold environments. Yet, alpine plants can be

exposed to very high temperatures, especially those with prostrate growth forms (Körner, 2003).

Studies on leaf temperature in alpine species in Europe have shown that leaf temperature can

reach up to 50°C in summer, especially on calm sunny days, with leaf heat tolerance higher than

50°C (Buchner & Neuner, 2003). Maximum leaf temperature has been measured at 38°C in early

summer in one Australian alpine species (Danzey *et al*., 2021 unpublished data) and may be

higher in suitable hot and still conditions in late summer. Therefore, thermal thresholds could be

breached in alpine species if the combination of heat with other environmental stress, such as

high light, decreased heat tolerances, as found for some European alpine species (Buchner &

Neuner, 2003; Buchner *et al.*, 2015).

Heat and cold thermal limits relate to different macroclimatic temperature variables

Predicting organismal responses to changing temperature regimes is a significant global

challenge and it requires a mechanistic understanding of how temperature affects species

physiology and distributions. Although the physiological mechanisms underpinning freezing and

cold tolerance (see Guy, 2003 for a review) and heat tolerance (Wahid *et al.*, 2007) have been

established in model plants, responses in species exposed to natural conditions remain

understudied. While the thermal limits of plants have recently been shown to broadly vary with

latitude (O'Sullivan *et al.*, 2017; Lancaster & Humphreys, 2020), comparisons employing

standard techniques across multiple biomes remain rare, particularly for both heat and cold

tolerance concomitantly (Geange *et al.*, 2021). Importantly, our capacity to predict thermal

tolerance from key bioclimate parameters such as MAT and MAP remains complex. In our

cross-biome dataset, we found that neither mean annual temperature or precipitation explained

variation in cold tolerance ($T_{crit\text{-}cold}$), which was instead strongly predicted by thermal

seasonality. For heat tolerance, again climate precipitation was not a significant driver, aligning

with the findings for Australian *Acacia* species in an experimental heatwave (Andrew *et al.*,

2023). On the other hand, mean annual temperature explained 23% of the variance in $T_{crit-hot}$,

which contrasts a common garden study of 123 taxonomically diverse species, for which climate

variables generally had very weak predictive power (<5%) for heat tolerance (Perez & Feeley,

2021).

Despite some clear macroclimatic patterns in our study, a large proportion of the variation in

cold and, to a lesser extent, heat tolerance was explained by species (32% and 19%,

respectively). The contribution of species could reflect micro-climate adaptation among species

or differences in species' leaf traits that contribute to within-species variation in leaf temperature,

the signal of which may be amplified when sampled *in situ*. Mahan and Upchurch (1988)

proposed that the lower limit of temperature for a plant is controlled by its environment, while

the upper limit can be controlled by the plant, suggesting that plants are "limited homeotherms".

Recent studies have shown that what drives variation in heat tolerance is leaf temperature, rather

than large scale measures of air temperature (Perez & Feeley, 2020; Cook *et al.*, 2021). Leaves

can avoid heat stress through different mechanisms that decouple leaf from air temperature such

as anatomical traits (Leigh *et al.*, 2012; Leigh *et al.*, 2017; Tserej & Feeley, 2021), patterns of

display such as leaf inclination (Ball *et al.*, 1988) and transpirational cooling through stomatal

conductance and can vary among co-existing species (Urban *et al.*, 2017; Drake *et al.*, 2018;

Deva *et al.*, 2020; Marchin *et al.*, 2022). The thermoregulatory patterns and physical properties

of leaves can differ greatly among species, such that different species exhibit different leaf

temperatures even when they co-exist in close proximity. Leaf to air temperature decoupling

(Blonder *et al.*, 2020) might also explain why relationships between heat tolerance and latitude

are negligible or smaller than expected for the temperature span considered (Curtis *et al.*, 2016;

O'Sullivan *et al.*, 2017; Lancaster & Humphreys, 2020).

Species' TTB is a more holistic measure of species thermal limits given that it encompasses both

lower and upper limits. Unlike for animals, global patterns of variation in the TTB of plants have

received little attention. Despite the existence of large thermal tolerance datasets, the lack of

paired, concurrently measured low and high thermal thresholds measurements for a given species

constrains our ability to adequately assess TTB globally. We found that TTB was best predicted

by the combination of seasonality and mean annual temperature: species exposed to the high

seasonality and mean annual temperatures in our desert biome had wider thermal tolerance

breadth. However, species played an equal role in explaining variation in TTB and the capacity

for species to respond to increasingly extreme high temperatures and greater climate variability

by broadening their TTB is unknown. Previous research generally suggests that plant niche

thermal evolution is constrained by heat tolerance (Aruajo et al 2013) as evident through reduced

variation in heat tolerance across landscapes (Lancaster and Humphreys, 2020). However, we

found the reverse, with heat tolerance having greater variability (~30°C) than cold tolerance

(~20°C) across our diverse biomes. Future studies could examine whether wider TTB or more

extreme thermal thresholds in fact equate to greater plasticity to warming and future climate

extremes.

Narrow thermal physiological limits have been identified as an aspect of species vulnerability to

climate change (Williams *et al.*, 2008; Pacifici *et al.*, 2015). The use of PSII thermal tolerance

metrics determined at the leaf-level, however, requires careful interpretation. When plants reach

their critical thermal limits, damage to PSII can occur, leading to decreased electron transport

rate and photosynthetic failure, thus compromising leaf survival. If this risks whole plant

survival, however is less clear. In 21 species, heat tolerance thresholds have been correlated with

carbon assimilation limits (Perez *et al,* 2021) which influence plant productivity. To further

unpick these patterns, we need to better understand the relationship between physiological

thermal limits with plant survival and fitness and how plastic these metrics are over space, time

and across the plant life cycle – for example, thermal tolerance of seedlings may be more critical

than that of adults.

Conclusion

Thermal tolerance breadth (TTB), a parameter broadly used in animal thermal biology literature,

has not previously been measured using a consistent method in such a diverse range of plant

species and environments. Here we concomitantly measured the lower ($T_{crit-cold}$ and NT) and

upper ($T_{crit-hot}$) thermal thresholds of PSII and quantified the TTB of 69 plant species across three

biomes: two climatically extreme (alpine and desert) and one more stable (wet temperate). Our

results show partial support for the climate variability hypothesis. Looking at climate from a

discreet biome category perspective, TTB only partially supported the CVH with the climatically

variable desert biome having the widest but only significantly different TTB. Using coarse

climate variables at sampling locations, we found support for CVH with temperature seasonality

predicting cold tolerance, TTB and NT but was lacking in heat tolerance (MAT not seasonality

was the significant predictor) with species explaining substantial portions of variation.

Interestingly we found that leaf-level TTB was extremely wide, much wider than the air

temperature extremes to which plants are exposed in these biomes. It is important to remember,

however, that these measured thermal limits represent a point beyond optimal photosynthetic

function and consistent temperatures inside these limits could have far-reaching effects,

depending on species life history, morphological traits and plasticity, as well as microclimatic

and temporal heterogeneity. Incorporating species traits, leaf temperature and microhabitat

information over space and time in future work is important next step to improving predictive

power of thermal tolerance metrics for understanding vulnerability of different vegetation

communities to climate extremes.

Acknowledgements

We acknowledge the Traditional Custodians of Country throughout Australia, especially across

the lands on which this study took place, and recognise the continuing connection to lands,

waters and communities across the landscapes. We pay our respects to Aboriginal and Torres

Strait Islander cultures and to Elders past, present and emerging. We thank the National Parks

and Wildlife Service for assistance and support, with special thanks to M. Hams, S. Doak and R.

Bjork at Gundabooka National Park, M. Schroder at Kosciuszko National Park, and J. Erskine, J.

Lemmon and P. Nagle at Royal National Park and Illawarra State Conservation Area. We thank

the following for assistance in the field: C. Bellotto, Z. Brown, A. Crimmins, C. Cunningham, P.

Cunningham, L. Danzey, M. Davies, L. Fuller, J.M. Hoskins, T. Salisbury, E. Rogers and K.

Spooner.

Author Contribution

445   VFB, AMC, SKCJ, ABN, LABR and AL designed the study, PAA, RVG, VGB and AMC

446   collated and analysed the data, all authors contributed to data collection, interpretation of results

447   and writing.

Conflict of interest

449   The authors have no conflicts of interest to disclose.

Declaration of funding

This work was supported by the Australian Research Council Linkage Project grant

(LP180100942) in partnership with the Save our Species program, DPE, NSW.

Ethics information

This study was conducted under approval from the NSW Department of Planning, Industry and

Environment (Scientific Licence SL102330).

References
458
Addo-Bediako A, Chown SL, Gaston KJ. 2000. Thermal tolerance, climatic variability and
latitude. *Proc Biol Sci* **267**(1445): 739-745.
Andrew SC, Arnold PA, Simonsen AK, Briceno VF. 2023. Consistently high heat tolerance
acclimation in response to a simulated heatwave across species from the broadly
distributed Acacia genus. *Functional Plant Biology* **50**(1): 71-83.
Anisko T, Lindstrom OM. 1996. Cold hardiness and water relations parameters in
Rhododendron cv. Catawbiense Boursault subjected to drought episodes. *Physiologia*
Plantarum **98**(1): 147-155.
Aparecido LMT, Woo S, Suazo C, Hultine KR, Blonder B. 2020. High water use in desert
plants exposed to extreme heat. *Ecology Letters* **23**(8): 1189-1200.
Arnold PA, Briceño VF, Gowland KM, Catling AA, Bravo LA, Nicotra AB. 2021. A high-
throughput method for measuring critical thermal limits of leaves by chlorophyll imaging
fluorescence. *Funct Plant Biol* **48**(6): 634-646.
Ball M, Cowan I, Farquhar G. 1988. Maintenance of Leaf Temperature and the Optimisation
of Carbon Gain in Relation to Water Loss in a Tropical Mangrove Forest. *Functional*
Plant Biology **15**(2): 263-276.
Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting linear mixed-effects models using
lme4. *Journal of Statistical Software* **67**(1): 1-48.
Baudier KM, D'Amelio CL, Malhotra R, O'Connor MP, O'Donnell S. 2018. Extreme
Insolation: Climatic Variation Shapes the Evolution of Thermal Tolerance at Multiple
Scales. *The American Naturalist* **192**(3): 347-359.
Bennett JM, Sunday J, Calosi P, Villalobos F, Martinez B, Molina-Venegas R, Araujo MB,
Algar AC, Clusella-Trullas S, Hawkins BA, et al. 2021. The evolution of critical
thermal limits of life on Earth. *Nature Communications* **12**(1): 9.
Berry J, Bjorkman O. 1980. Photosynthetic Response and Adaptation to Temperature in
Higher Plants. *Annual Review of Plant Physiology* **31**(1): 491-543.
Blake J, Hill R. 1996. An Examination of the Drought and Frost Tolerance of *Banksia*
marginata (Proteaceae) as an Explanation of Its Current Widespread Occurrence in
Tasmania. *Australian Journal of Botany* **44**(3): 265-281.
Blonder B, Escobar S, Kapás RE, Michaletz ST. 2020. Low predictability of energy balance
traits and leaf temperature metrics in desert, montane and alpine plant communities.
Functional Ecology **34**(9): 1882-1897.
Brestic, M., Zivcak, M., Kalaji, H.M., Carpentier, R. & Allakhverdiev, S.I. 2012.
Photosystem II thermostability in situ: Environmentally induced acclimation and
genotype-specific reactions in Triticum aestivum L. *Plant Physiology and Biochemistry*,
57: 93-105.
Briceño VF, Harris-Pascal D, Nicotra AB, Williams E, Ball MC. 2014. Variation in snow
cover drives differences in frost resistance in seedlings of the alpine herb Aciphylla
glacialis. *Environmental and Experimental Botany* **106**(0): 174-181.

Buchner O, Neuner G. 2003. Variability of heat tolerance in alpine plant species measured at
different altitudes. *Arctic Antarctic and Alpine Research* **35**(4): 411-420.
Buchner O, Stoll M, Karadar M, Kranner I, Neuner G. 2015. Application of heat stress in
situ demonstrates a protective role of irradiation on photosynthetic performance in alpine
plants. *Plant Cell and Environment* **38**(4): 812-826.
Byrne M, Yeates DK, Joseph L, Kearney M, Bowler J, Williams MAJ, Cooper S,
Donnellan SC, Keogh JS, Leys R, et al. 2008. Birth of a biome: insights into the
assembly and maintenance of the Australian arid zone biota. *Molecular Ecology* **17**(20):
4398-4417.
Calosi P, Bilton DT, Spicer JI, Votier SC, Atfield A. 2010. What determines a species'
geographical range? Thermal biology and latitudinal range size relationships in European
diving beetles (Coleoptera: Dytiscidae). *J Anim Ecol* **79**(1): 194-204.
Compton TJ, Rijkenberg MJA, Drent J, Piersma T. 2007. Thermal tolerance ranges and
climate variability: A comparison between bivalves from differing climates. *Journal of*
Experimental Marine Biology and Ecology **352**(1): 200-211.
Cook AM, Berry N, Milner KV, Leigh A. 2021. Water availability influences thermal safety
margins for leaves. *Functional Ecology* **35**(10): 2179-2189.
Cunningham SC, Read J. 2006. Foliar temperature tolerance of temperate and tropical
evergreen rain forest trees of Australia. *Tree Physiology* **26**(11): 1435-1443.
Curtis EM, Gollan J, Murray BR, Leigh A. 2016. Native microhabitats better predict tolerance
to warming than latitudinal macro-climatic variables in arid-zone plants. *Journal of*
Biogeography **43**(6): 1156-1165.
Curtis EM, Knight CA, Leigh A. 2019. Intracanopy adjustment of leaf-level thermal tolerance
is associated with microclimatic variation across the canopy of a desert tree (Acacia
papyrocarpa). *Oecologia* **189**(1): 37-46.
Deva CR, Urban MO, Challinor AJ, Falloon P, Svitákova L. 2020. Enhanced Leaf Cooling Is
a Pathway to Heat Tolerance in Common Bean. *Frontiers in Plant Science* **11**.
Dörken VM, Ladd PG, Parsons RF. 2020. Anatomical aspects of xeromorphy in arid-adapted
plants of Australia. *Australian Journal of Botany* **68**(3): 245-266.
Downton WJS, Berry JA, Seemann JR. 1984. Tolerance of photosynthesis to high temperature
in desert plants. *Plant Physiology* **74**(4): 786-790.
Drake JE, Tjoelker MG, Vårhammar A, Medlyn Belinda E, Reich PB, Leigh A, Pfautsch S,
Blackman CJ, López R, Aspinwall MJ, et al. 2018. Trees tolerate an extreme heatwave
via sustained transpirational cooling and increased leaf thermal tolerance. *Global Change*
Biology **24**(6): 2390-2402.
Garcia-Pichel F, Loza V, Marusenko Y, Mateo P, Potrafka RM. 2013. Temperature Drives
the Continental-Scale Distribution of Key Microbes in Topsoil Communities. *Science*
340(6140): 1574-1577.
Geange SR, Arnold PA, Catling AA, Coast O, Cook AM, Gowland KM, Leigh A,
Notarnicola RF, Posch BC, Venn SE, et al. 2021. The thermal tolerance of
photosynthetic tissues: a global systematic review and agenda for future research. *New*
Phytol **229**(5): 2497-2513.
Guy CL. 2003. Freezing tolerance of plants: current understanding and selected emerging
concepts. *Canadian Journal of Botany* **81**(12): 1216-1223.
Hüve, K., Bichele, I., Tobias, M. & Niinemets, Ü. 2006. Heat sensitivity of photosynthetic
electron transport varies during the day due to changes in sugars and osmotic potential.
Plant, Cell & Environment, **29:** 212-228.

Janzen DH. 1967. Why mountain passes are higher in the tropics. *The American Naturalist*
101(919): 233-249.

Jin Y, Qian H. 2022. V.PhyloMaker2: An updated and enlarged R package that can generate
very large phylogenies for vascular plants. *Plant Diversity* **44**(4): 335-339.

Johnson PCD. 2014. Extension of Nakagawa & Schielzeth's $R^{2\text{GLMM}}$ to random slopes models.
Methods in Ecology and Evolution **5**(9): 944-946.

Karger DN, Conrad O, Böhner J, Kawohl T, Kreft H, Soria-Auza RW, Zimmermann NE,
Linder HP, Kessler M. 2017. Climatologies at high resolution for the earth's land
surface areas. *Scientific Data* **4**(1): 170122.

Keck F, Rimet F, Bouchez A, Franc A. 2016. phylosignal: an R package to measure, test, and
explore the phylogenetic signal. *Ecology and Evolution* **6**(9): 2774-2780.

Knight CA, Ackerly DD. 2002. An ecological and evolutionary analysis of photosynthetic
thermotolerance using the temperature-dependent increase in fluorescence. *Oecologia*
130(4): 505-514.

Knight CA, Ackerly DD. 2003. Evolution and plasticity of photosynthetic thermal tolerance,
specific leaf area and leaf size: congeneric species from desert and coastal environments.
New Phytologist **160**(2): 337-347.

Körner C. 2003. *Alpine Plant life*. Berlin: Springer.

Kreutzmann AK, Pannier S, Rojas-Perilla N, Schmid T, Templ M, Tzavidis N. 2019. The R
Package emdi for Estimating and Mapping Regionally Disaggregated Indicators. *Journal*
of Statistical Software **91**(7): 1-33.

Kuznetsova A, Brockhoff PB, Christensen RHB. 2017. lmerTest Package: Tests in Linear
Mixed Effects Models. *Journal of Statistical Software* **82**(13): 1-26.

Lancaster LT, Humphreys AM. 2020. Global variation in the thermal tolerances of plants.
Proceedings of the National Academy of Sciences **117**(24): 13580-13587.

Larcher W. 2003. *Physiological Plant Ecology : Ecophysiology and stress physiology of*
Functional Group: Springer-Verlag

Leigh A, Sevanto S, Ball MC, Close JD, Ellsworth DS, Knight CA, Nicotra AB, Vogel S.
2012. Do thick leaves avoid thermal damage in critically low wind speeds? *New*
Phytologist **194**(2): 477-487.

Leigh A, Sevanto S, Close JD, Nicotra AB. 2017. The influence of leaf size and shape on leaf
thermal dynamics: does theory hold up under natural conditions? *Plant, Cell &*
Environment **40**(2): 237-248.

Lintunen A, Hölttä T, Kulmala M. 2013. Anatomical regulation of ice nucleation and
cavitation helps trees to survive freezing and drought stress. *Scientific Reports* **3**(1): 2031.

Mahan JR, Upchurch DR. 1988. Maintenance of constant leaf temperature by plants—I.
Hypothesis-limited homeothermy. *Environmental and Experimental Botany* **28**(4): 351-
357.

Maher T, Mirzaei M, Pascovici D, Wright IJ, Haynes PA, Gallagher RV. 2019. Evidence
from the proteome for local adaptation to extreme heat in a widespread tree species.
Functional Ecology **33**(3): 436-446.

Marchin RM, Backes D, Ossola A, Leishman MR, Tjoelker MG, Ellsworth DS. 2022.
Extreme heat increases stomatal conductance and drought-induced mortality risk in
vulnerable plant species. *Global Change Biology* **28**(3): 1133-1146.

Molina-Montenegro MA, Naya DE. 2012. Latitudinal Patterns in Phenotypic Plasticity and
Fitness-Related Traits: Assessing the Climatic Variability Hypothesis (CVH) with an
Invasive Plant Species. *PloS one* **7**(10): e47620.

Mora C, Caldwell IR, Caldwell JM, Fisher MR, Genco BM, Running SW. 2015. Suitable
Days for Plant Growth Disappear under Projected Climate Change: Potential Human and
Biotic Vulnerability. *Plos Biology* **13**(6): 15.
Muggeo VMR. 2017. Interval estimation for the breakpoint in segmented regression: a
smoothed score-based approach. *Australian & New Zealand Journal of Statistics* **59**: 311-
322.
Mumladze L, Asanidze Z, Walther F, Hausdorf B. 2017. Beyond elevation: testing the
climatic variability hypothesis vs. Rapoport's rule in vascular plant and snail species in
the Caucasus. *Biological Journal of the Linnean Society* **121**(4): 753-763.
Neuner G, Pramsohler M. 2006. Freezing and high temperature thresholds of photosystem 2
compared to ice nucleation, frost and heat damage in evergreen subalpine plants.
Physiologia Plantarum **126**(2): 196-204.
O'Sullivan OS, Heskel MA, Reich PB, Tjoelker MG, Weerasinghe LK, Penillard A, Zhu L,
Egerton JJG, Bloomfield KJ, Creek D, et al. 2017. Thermal limits of leaf metabolism
across biomes. *Global Change Biology* **23**(1): 209-223.
Pacifici M, Foden WB, Visconti P, Watson JEM, Butchart SHM, Kovacs KM, Scheffers
BR, Hole DG, Martin TG, Akçakaya HR, et al. 2015. Assessing species vulnerability
to climate change. *Nature Climate Change* **5**(3): 215-224.
Perez TM, Andino JEG, Rivas-Torres G, Feeley KJ. 2023. Climate Constrains Photosynthetic
Strategies in Darwin's Daisies: A Test of the Climatic Variability and Jack-of-All-Trades
Hypotheses. *American Naturalist* **201**(1): 78-90.
Perez TM, Feeley KJ. 2020. Photosynthetic heat tolerances and extreme leaf temperatures.
Functional Ecology **34**(11): 2236-2245.
Perez TM, Feeley KJ. 2021. Weak phylogenetic and climatic signals in plant heat tolerance.
Journal of Biogeography **48**(1): 91-100.
Perez TM, Feeley KJ, Michaletz ST, Slot M. 2021. Methods matter for assessing global
variation in plant thermal tolerance. *Proceedings of the National Academy of Sciences of*
the United States of America **118**(30): 2.
Perez, T.M., Socha, A., Tserej, O. & Feeley, K.J. 2021. Photosystem II heat tolerances
characterize thermal generalists and the upper limit of carbon assimilation. *Plant, Cell*
and Environment, **44:** 2321-2330. https://doi.org/10.1111/pce.13990
R Core Team 2018. R: A language and environment for statistical computing. Vienna, Austria:
R Foundation for Statistical Computing.
R Core Team 2020. R: A language and environment for statistical computing. Vienna, Austria:
R Foundation for Statistical Computing.
Rummukainen M. 2012. Changes in climate and weather extremes in the 21st century. *Wiley*
Interdisciplinary Reviews: Climate Change **3**(2): 115-129.
Sakai A, Larcher W. 1987. *Frost survival of Plants: Responses and adaptation to freezing*
stress: Springer-Verlag Berlin.
Schreiber U, Berry JA. 1977. Heat-induced changes of chlorophyll fluorescence in intact leaves
correlated with damage of the photosynthetic apparatus. *Planta* **136**(3): 233-238.
Seemann JR, Berry JA, Downton WJ. 1984. Photosynthetic response and adaptation to high
temperature in desert plants : a comparison of gas exchange and fluorescence methods for
studies of thermal tolerance. *Plant Physiol* **75**(2): 364-368.
Seneviratne S, Nicholls N, Easterling D, Goodess C, Kanae S, Kossin J, Luo Y, Marengo J,
McInnes K, Rahimi M 2012. Managing the Risks of Extreme Events and Disasters to
Advance Climate Change Adaptation. A Special Report of Working Groups I and II of

the Intergovernmental Panel on Climate Change (IPCC): Cambridge University Press
Cambridge.
Sierra-Almeida A, Reyes-Bahamonde C, Cavieres LA. 2016. Drought increases the freezing
resistance of high-elevation plants of the Central Chilean Andes. *Oecologia* **181**(4):
1011-1023.
Siminovitch D, Cloutier Y. 1983. Drought and freezing tolerance and adaptation in plants:
Some evidence of near equivalences. *Cryobiology* **20**(4): 487-503.
Smillie R, Nott R. 1979. Heat Injury in Leaves of Alpine, Temperate and Tropical Plants.
Functional Plant Biology **6**(1): 135-141.
Sunday JM, Bates AE, Dulvy NK. 2011. Global analysis of thermal tolerance and latitude in
ectotherms. *Proceedings of the Royal Society B: Biological Sciences* **278**(1713): 1823-
1830.
Tattersall GJ, Sinclair BJ, Withers P, Fields PA, Seebacher F, Cooper CE, Maloney S 2012.
Coping with Thermal Challenges: Physiological Adaptations to Environmental
Temperatures. *Comprehensive Physiology*.
Tserej O, Feeley KJ. 2021. Variation in leaf temperatures of tropical and subtropical trees are
related to leaf thermoregulatory traits and not geographic distributions. *Biotropica* **53**(3):
868-878.
Urban J, Ingwers MW, McGuire MA, Teskey RO. 2017. Increase in leaf temperature opens
stomata and decouples net photosynthesis from stomatal conductance in Pinus taeda and
Populus deltoides x nigra. *Journal of Experimental Botany* **68**(7): 1757-1767.
Wahid A, Gelani S, Ashraf M, Foolad MR. 2007. Heat tolerance in plants: An overview.
Environmental and Experimental Botany **61**(3): 199-223.
Williams SE, Shoo LP, Isaac JL, Hoffmann AA, Langham G. 2008. Towards an Integrated
Framework for Assessing the Vulnerability of Species to Climate Change. *Plos Biology*
6(12): e325.