Evolution of olfactory sensitivity, preferences and behavioral responses in Mexican cavefish: fish personality matters

Maryline Blin, Louis Valay, Manon Kuratko, Marie Pavie and Sylvie Rétaux

Paris-Saclay Institute of Neuroscience,

CNRS and University Paris-Saclay, 91400, Saclay, France

Author for correspondence: sylvie.retaux@cnrs.fr

Running title: Cavefish olfactory behavior

Key words: evolution, amino acids, preference, sensitivity, personality, genetics

Abstract

Animals are adapted to their natural habitats and lifestyles. Their brains perceive the external world *via* their sensory systems, compute information together with that of internal states and autonomous activity, and generate appropriate behavioral outputs. However, how do these processes evolve across evolution? Here, focusing on the sense of olfaction, we have studied the evolution in olfactory sensitivity, preferences and behavioral responses to six different food-related amino acid odors in the two eco-morphs of the fish *Astyanax mexicanus*. To this end, we have developed a high-throughput behavioral setup and pipeline of quantitative and qualitative behavior analysis, and we have tested 489 six-week-old *Astyanax* larvae. The blind, dark-adapted morphs of the species showed markedly distinct basal swimming patterns and behavioral responses to odors, higher olfactory sensitivity and a strong preference for alanine, as compared to their river-dwelling eyed conspecifics. In addition, we discovered that fish have an individual "swimming personality", and that this personality influences their capability to respond efficiently to odors and find the source. Importantly, the personality traits that favored significant responses to odors were different in surface fish and cavefish. Moreover, the responses displayed by second-generation cave x surface F2 hybrids suggested that olfactory-driven behavior and olfactory sensitivity is a quantitative genetic trait. Our findings show that olfactory processing has rapidly evolved in cavefish at several levels: detection threshold, odor preference, and foraging behavior strategy. Cavefish is therefore an outstanding model to understand the genetic, molecular and neurophysiological basis of sensory specialization in response to environmental change.

Introduction

With more than 26.000 species representing half of vertebrates, bony fishes are extremely diverse and have colonized all possible ecological niches (Helfman et al., 2009), making them outstanding models to study the neurophysiological, genetic and evolutionary underpinnings of adaptive behaviors. To colonize, survive and thrive in various environments, sensory systems are particularly crucial as they serve as windows to the external world, and they display an exceptional diversity in fishes. From the visual system of deep-sea fish shaped to catch the rare photons or the point-like bioluminescent signals to the mobile barbels covered with taste buds to probe the sea bottom and detect buried prey of bottom feeders, examples abound (de Busserolles et al., 2020; Kiyohara et al., 2002).

The anatomical sensory specializations, that is the relative importance taken by specific sensory organs or brain sensory areas, arise during embryonic development through regulatory processes that control the patterning of the neuroepithelium and that define boundaries between presumptive neural regions (Krubitzer et al., 2011). The relative investment in different sensory modules governs anatomical and behavioral specializations. Among fish, sand-dwelling cichlids that rely almost exclusively on vision to execute behaviors develop a large optic tectum, whereas rock-dwelling cichlids, which inhabit complex environments, have small optic tecta but an enlarged telencephalon, and these variations arise early in development (Sylvester et al., 2010). Similarly, in the embryos of dark-adapted blind Mexican cavefish, the presumptive eyefield territory is reduced but the presumptive olfactory epithelium is increased in size, presumably as a compensation for the loss of visual modality in the dark environment (Agnès et al., 2022; Hinaux et al., 2016a; Pottin et al., 2011; Torres-Paz et al., 2019).

While the comparative anatomy of fish brains is reasonably well documented, behavioral studies have mainly focused on a few model species, primarily zebrafish (Kalueff et al., 2013). The behavioral correlates of sensory systems diversity and evolution in fishes are poorly described and understood, which hampers the interpretation of cross-species comparisons. For example, the size and complexity of fish olfactory organs is highly variable and correlates with the richness of odorant receptors repertoire, defining a morpho-genomic space in which olfactory specialists and non-specialists species distribute (Burguera et al., 2023; Policarpo et al., 2022). On the lower end, sunfish or pipefish have a small and flat olfactory epithelium and possess ~30 odorant receptor genes. On the higher end, polypteriforms have large and complex olfactory rosettes with up to 300 lamellae and possess ~1300 odorant receptor genes. Yet, the differences in olfactory behaviors, sensitivity and preferences between these species are completely unknown.

To start addressing the question of the evolution of olfactory sensory-driven behaviors in an amenable laboratory fish model, we used the blind cave and the river dwelling morphs of the Mexican tetra, the characiform *Astyanax mexicanus*. The species has become an established model for evolutionary biology at large, including evolution of behaviors (Duboué and Keene, 2016; Hinaux et al., 2016b; Kowalko, 2020; Yoshizawa, 2016). Cavefish embryos and larvae have larger olfactory pits than surface fish and their neuronal composition is changed (Blin et al., 2018; Hinaux et al., 2016a), while adults of the two forms have a similar olfactory rosette with 20-25 lamellae (Schemmel, 1967). The recent divergence between the two forms probably did not allow for a substantial evolution of their olfactory receptor gene repertoire, which is almost identical in the two eco-morphotypes (245 genes in cavefish, 233 genes in surface fish, from recent genome assemblies) (Policarpo et al., 2022). Yet, cavefish larvae tested in groups display outstanding olfactory detection capacities, as they are attracted to extremely low

concentration of the amino acid alanine (10^{-10}M) while surface fish larvae have a more "classical" threshold (10^{-5}M) (Hinaux et al., 2016a) which approximates levels of free amino acids found in natural waters (Hara, 1994). In wild natural caves too, groups of cavefish adults respond to low concentrations of odorants (Blin et al., 2020). We hypothesized that the evolution of olfactory system development and olfactory skills in blind cavefish is an adaptive trait that, together with other constructive sensory changes in mechano-sensory and gustatory systems (Varatharasan et al., 2009; Yamamoto et al., 2009; Yoshizawa and Jeffery, 2011; Yoshizawa et al., 2014), may have contributed to their adaptation and survival in the extreme environment of their subterranean habitat. Focusing on olfaction, here we developed a sensitive, high-throughput behavioral assay and a pipeline for analysis to compare the types of behavioral responses elicited by food-related, amino acid odors in cavefish and surface fish, at population and individual levels. We determined their olfactory preferences and sensitivity thresholds for six amino acids, and we analyzed their behavioral responses in detail. We also described olfactory-driven behaviors in second-generation (F2) hybrids resulting from crosses between surface and cave morphs, as an attempt to understand the genetic component of the "olfaction trait" in *Astyanax* morphs. We discovered that the behaviors triggered by odorant stimulation has markedly evolved in cavefish.

Results

A high-throughput, sensitive behavioral assay to compare individual's behaviors in *Astyanax* morphotypes

Our goal was to compare 1) olfactory discrimination capacities, 2) odor preferences and 3) behavioral responses to olfactory stimuli in Pachón cavefish (CF), surface fish (SF) and F2 hybrids (F2) individuals. As the three types of fish are markedly different in terms of basal swimming activity and patterns, we first sought to characterize the diversity of these behaviors in order to be able to interpret accurately fish responses to odors.

Six weeks old fish were habituated either 1h or 24h in their individual test box and they were first recorded for 30 minutes in the dark, without any stimulus (Figure 1; see Methods). We observed and categorized several typical and distinctive "baseline" swimming behaviors for individual fish (Figure 2A): random/haphazard swim (R), wall following (WF; defined as the fish continuously following the 4 sides of the box and turning around it, in a clockwise or counterclockwise fashion), large or small circles (C), thigmotactism (T, along the X- or the Y-axis of the box; defined as the fish swimming back and forth along one of the 4 sides of the box), or combinations thereof. The distribution of these different types of swimming patterns was significantly different in Pachón CF, SF and F2 types of fish (Figure 2B; Fisher's exact test; see also Methods and Suppl. Fig8A for another representation using Correspondence Analysis). A majority of SF swam in a random pattern (blue shades), while a majority of CF performed wall following (red/brown shades) and F2 fish showed more diversified swimming patterns. Importantly, the distribution of these baseline-swimming patterns was affected as a function of the habituation time, in all three types of fish (Figure 2B; p=0.044 for SF, p=0.0005

for CF and p=0.0005 for F2, Fisher's exact tests). Moreover, thigmotactism behavior, which

we have previously shown to represent stress behavior (Pierre et al., 2020), was frequent in SF

after 1h habituation (70% of individuals) but reduced after 24h acclimation (37%). In the same

line, the swimming speed was different for all fish types when computed after 1h or after 24h

of habituation (Figure 2C). This suggested that natural, unperturbed behavior is better observed

after long, 24h habituation, which we applied thereafter.

As described above at population level, cavefish, surface fish and F2 displayed overall

markedly different basal swimming patterns (Figure 2B; see also Methods and Suppl. Fig8A

for another representation using Correspondence Analysis). Yet, at individual level, there was

a substantial degree of variability across individuals within a morphotype. That is, even though

most SF tended to swim randomly (75% after 24h habituation; blue shades on graphs) and most

CF performed wall following (50%; red/brown shades), these behaviors were not exclusive as

some SF individuals displayed thigmotactism behavior or some CF individuals swam in circles.

Moreover and importantly, the type of swim pattern of each individual fish was remarkably

stable and reproducible along several days of recordings, and swimming kinetics (position in

the box, swimming speed, number of round trips in X and Y) showed no or very little variation

(Figure 2D and Suppl. Fig1AA'). Together, these observations strengthened the importance of

1) using long habituation times, and 2) studying individual behaviors. They further highlighted

an overlooked aspect of fish behavioral analyses: fish may have a "personality" that we sought

to take into account when comparing behavioral responses to odors below.

Finally, Pachón CF, SF and F2 hybrids possess markedly different sensory apparatuses and

capacities, not only for chemo- but also for mechano-sensation (Lunsford et al., 2022;

Yoshizawa et al., 2010). Therefore, to test behavioral responses specific to olfactory stimuli,

we wanted to find a way to deliver odors that would be vibration-less. We choose to inject

odorant solutions inside tubes attached at the two extremities of the test box, designed to stop

the wave and flow that would otherwise arise at the surface of the water upon injection (Figure

1). The efficacy of the device was demonstrated by the lack of perturbation in swimming

behavior, neither qualitative (i.,e., behavioral response and swimming pattern; Figure 3AB) nor

quantitative (i.,e., swimming kinetics; Figure 3C) for the 3 types of fish when water/control

injections were performed on one side of the test box (or two sides; Suppl. Fig1BC). This

suggested that our way to deliver odorant solutions did not generate vibratory perturbations.

Moreover, when no injection was made, individual fish behaviors were stable over one hour of

test (Suppl. Fig1D), allowing to observe accurately the effects of an olfactory stimulus if applied

after 30 minutes. Finally, all recordings were performed in the dark under infra-red lights to

neutralize the visual modality in sighted fish, a procedure which did not affect SF behavior

(Suppl. Fig1E). Hence, the behavioral responses to odors recorded thereafter in Pachón CF, SF

and F2 are driven by olfaction and can be compared (see also Hinaux et al., 2016a).

Behavioral responses to alanine in *Astyanax* morphotypes

Amino acids are food-related odorant cues for fish. Next, we characterized individual

behavioral responses elicited by different concentrations of alanine, a potent aliphatic amino

acid cue for most fish species including *Astyanax*.

Injection of alanine 10^{-2}M, 10^{-3}M, 10^{-4}M or 10^{-5}M (hence 10^{-4}M, 10^{-5}M, 10^{-6}M or 10^{-7}M final

concentration in the odorant third of the testing box, respectively) induced a strong behavioral

response in cavefish (Figure 4A, 10^{-4}M injection shown; Suppl. Fig2A for other examples).

Upon stimulus, CF decreased the numbers of back-and-forth swims along the X- and the Y-

axes of the box (Figure 4A, top left graph and Figure 4FG), and they shifted and restricted their swimming activity towards the odorant side of the arena, close to the odor delivery tube (Figure 4A, bottom left graph, and Figure 4E). A tendency to decreasing swimming speed was observed but was significant only for 10^{-3}M injections (Figure 4D). Alanine injection also markedly changed CF swimming patterns, as they completely switched from WF and T to R and C swimming modes (Figure 4H and Suppl. Fig2D; Fisher's exact tests p values: 0.014, 0.0005, 0.0005 and 0.099 for 10^{-2}M, 10^{-3}M, 10^{-4}M or 10^{-5}M alanine injections, respectively; see also Methods and Suppl. Fig8B for another representation using Correspondence Analysis). In sum, CF individuals displayed robust behavioral responses and attraction to alanine. Of note, further decreasing the concentration of injected alanine to 10^{-6}M, 10^{-7}M, 10^{-8}, 10^{-9}M and 10^{-10}M had milder to no effects on cavefish. Swimming patterns remained globally unchanged (not shown), but swimming speed was decreased (p= 0.005 for 10^{-7}M; n=16), numbers of round trips in Y were decreased (p= 0.012 for 10^{-7}M; n=16), and numbers of round trips in X were decreased (p= 0.017 for 10^{-7}M; p= 0.029 for 10^{-8}M; p= 0.003 for 10^{-9}M; n=16 each; data not shown). This suggested that in the present experimental setup, cavefish can detect very low concentrations of alanine, in agreement with previously published data (Hinaux et al., 2016a).

Conversely, surface fish responses were more subtle and seemed restricted to some individuals (Figure 4B; Suppl. Fig2B for other examples). Notably, the numbers of back-and-forth swims along the X- and the Y-axes of the box were mostly unchanged and the position in the box did not vary for SF upon alanine injection (Figure 4B, top and bottom left graphs and Figure 4E-G). The swimming speed was unchanged (Figure 4D). Swimming patterns were globally unaffected upon alanine injection (Figure 4H and Suppl. Fig2D; Fisher's exact tests p values: 0.16, 0.54 and 0.80 for 10^{-2}M, 10^{-3}M or 10^{-4}M alanine injections, respectively; Suppl. Fig8B for another representation using Correspondence Analysis). As shown on Suppl. Fig2B, some

rare SF individuals appeared to change their swimming mode upon alanine 10^{-2}M (high

concentration) injection, suggesting that they did perceive and react to the odorant stimulus.

However, these rare individual responses were "diluted" at population level by the pooling of

all fish in the distribution graphs. In summary, SF behavioral responses were modest and

markedly different from CF.

Finally, F2 were tested with alanine 10^{-3}M, a concentration that elicits strong responses in CF

but not in SF. Upon stimulus, F2's change in behaviors were similar to CF (Figure 4C; Suppl.

Fig2C for other examples): they decreased back-and-forth swimming activity, decreased

swimming speed, and swam close to the odor source (Figure 4D-G). They also shifted their

swimming patterns (Suppl. Fig2D; Fisher's exact test p=0.0065), including a loss of wall

following mode that was reminiscent of the trend observed in CF. As an illustration, the

individual shown in Figure 4C displays a striking change from a "large circles" to a "random"

swimming pattern accompanied with variations in swimming kinematics.

Overall, these data suggested that, compared to SF, CF detection threshold and behavioral

responses to the amino acid alanine have significantly evolved.

Behavioral responses to serine and cysteine in *Astyanax* morphotypes

We next systematically tested behavioral responses to injections of 10^{-2}M, 10^{-3}M or 10^{-4}M of

serine (polar amino acid, hydroxyl group) and cysteine (non-polar, sulfur containing), two other

potent amino acid olfactory cues for fish (Figure 5 and Suppl. Fig3).

Serine elicited behavioral responses similar to alanine: CF (as well as F2 hybrids, Suppl. Fig3)

moved towards the odor source, whereas SF did not (Figure 5AB and D). Swimming speed was

unchanged in all fish types (Figure 5C), but a decrease or an increase of back-and-forth swimming activity was observed in CF and SF, respectively (Figure 5EF). At population level, significant changes in swimming patterns were observed for the three fish types upon 10^{-2}M serine (Figure 5G; Fisher's exact p-values 0.0005, 0.001 and 0.0085 for SF, CF and F2, respectively; Suppl. Fig8B for Correspondence Analysis). Of note, and contrarily to alanine, the lower concentrations of serine tested (10^{-4}M and 10^{-3}M in syringe; 10^{-6}M and 10^{-5}M in box, respectively) were unable to trigger a robust behavioral change in cavefish (not shown, see Table1), suggesting that their detection threshold for serine is higher (not as good as) than for alanine. Regarding SF, responses were also observed only for 10^{-2}M serine injections.

Cysteine also produced significant effects (Figure 5H-N and Suppl. Fig3). Swimming speed and numbers of back-and-forth swims along the Y-axis increased only in SF and for the 3 concentrations tested (Figure 5J and 5M for 10^{-2}M) (for 10^{-3}M: p= 0.01 and p=2e-05, n=32; for 10^{-4}M: p=2e-05 and p= 0.00028, n=32). CF were attracted to the odorant side for the 3 concentrations tested (p=0.013 to 0.025; n=16 to 38 each), whereas SF did not change position (Figure 5K) except for cysteine 10^{-3}M to which they seemed to be repelled (p=0.0003, n=32). The three fish types changed significantly their swimming patterns (Figure 5N; Fisher's exact p-values 0.03, 0.017 and 0.04 for SF, CF and F2, respectively; Suppl. Fig8B for Correspondence Analysis). After injection of cysteine at the lower concentrations of 10^{-3}M or 10^{-4}M, CF behavioral responses were similar to those observed for alanine and serine, i.e., a decrease in back-and-forth swimming activity along the X-axis and a significant change in position towards the odor source, without changing swimming speed (not shown, see Table1). In sum, the different fish types show diverse responses to different concentrations of different amino acids.

Behavioral responses to lysine, histidine and leucine in *Astyanax* morphotypes

Finally, we choose to examine responses triggered by high concentrations (10^{-2}M) of three other, less studied amino acids: lysine, histidine (both polar and positively charged) and leucine (aliphatic, like alanine) (Figure 6 and Suppl. Fig4).

For these three odors, CF responses were conspicuous and could include a shift of swim position towards the odor's source, a decrease of back-and-forth swimming activity with decrease in swimming speed, and significant changes in swim patterns at population level (except for leucine) (Figure 6A-G and Suppl. Fig4; Suppl. Fig8B for Correspondence Analysis). Regarding SF, changes were restricted to increases in back and forth swimming activity for histidine, without change in swimming speed or position in the box (Figure 6A-D and Suppl. Fig4). These three amino acids did not elicit changes in swimming pattern in SF at population level (Figure 6E-G). Finally, F2 also showed specific qualitative (swimming patterns) and quantitative responses to each of these three amino acids. Together, these data suggested that CF, SF and F2 detected and responded in their own and specific way to high concentrations of lysine, histidine and leucine.

Table 1A provides a summary of the behavioral responses, averaged at population level, for the three fish types and the six amino acids tested. In the cases when significant behavioral responses were observed at population level, i.e., when responses are strongest and most stereotyped, different and opposed features emerged between CF and SF (and F2). As a "rule", CF responses included a decrease in swimming activity (decrease in speed and back and forth swims along X- and Y-axes) together with an attraction to odor source and a change in swim pattern (Table 1A, red arrows). Conversely, SF responses mostly corresponded to an increase

in swimming activity, with little or no attraction to odor and moderate change in swim pattern (Table 1A, blue arrows). In sum, a given sensory stimulus could trigger markedly different, morph-specific behavioral responses.

Behavioral responses to odors at individual level

Analyses as above performed at population level may mask or blur effects or phenotypes in the case when not all individuals respond in a stereotyped manner. As fish did express individual behavioral features in our experimental paradigm, we sought to perform further analyses at individual level, and to calculate individual response scores to the different odors. To do so, and to take into account the different components of the behavioral response, we summed the absolute values of indexes of speed, back and forth trips in X and Y, position and pattern changes (see Methods; Figure 7A). The threshold of score for which an individual fish was considered to respond significantly to a given stimulus was set at 1.5, from examination of the individual scores of fish in control conditions after perfusion of water (Suppl. Fig5). Visual inspection of the responses to amino acid odors for fish who had an individual score just above or just below this threshold confirmed that it was accurate.

For alanine 10^{-3}M, the representation of individual behavioral responses shows that CF responded in a very stereotyped manner (all lines following the same "curve") whereas SF responses were more diverse (lines cross) (Figure 7A, 1^{st} column). Moreover, 73% of CF but only 19% of SF had an individual score above threshold - a marked difference that was visible on the distribution of scores (Figure 7AB). F2 fish on the other end had a bimodal distribution of their olfactory scores for alanine 10^{-3}M (Figure 7B).

For other odors, the difference in olfactory scores of individual CF and SF was less obvious

(Figure 7B and Suppl. Fig5), suggesting that taking into consideration individual's variations

of swimming behavior can unmask responses that are not visible when averaged at population

level. For example, in response to cysteine 10^{-2}M, SF (51% with score >1.5) and F2 (55% with

score >1.5) had similar or slightly better scores than CF (39% with score >1.5) (Figure 7AB,

3^{rd} column). Of note, these results had not shown up in Table 1A summarizing responses

averaged at population level. Noteworthy, SF had mixed, diverse responses to all odors studied

(lines crossing on all graphs). Conversely, CF showed non-homogenous responses to serine,

histidine, leucine and lysine but highly stereotyped responses to both alanine and histidine

(Figure 7A and Suppl. Fig5). A summary of individual response scores to the different odors is

given in Table 1B.

Individual personality and individual behavioral response

The results presented above led us to test whether the "personality" of each fish, represented by

their specific baseline swimming behavior, could influence their ability to respond to odors. To

this end, we plotted individual olfactory scores as a function of individual baseline swimming

patterns (R, WF, TX, TY, or C). We also examined possible correlations between basal

swimming speed or round trip activity and individual olfactory score.

In SF, individual olfactory scores were similar whatever the basal swimming pattern of the fish

(Figure 8A). However, for several odors, those fish exhibiting lower baseline swimming speed

and less round trips in X had better olfactory scores (Figure 8B and Suppl. Fig6A; negative

correlation). By contrast, in CF, fish exhibiting WF or C as baseline pattern had significantly

better individual scores and those swimming randomly (R) had poor scores (Figure 8A) – but

scores were not correlated whatsoever to baseline swimming speed (Figure 8B). Moreover, for alanine 10^{-3}M and histidine 10^{-2}M, the two odors for which more than 70% of CF showed a significant response (Figure 7A), a high number of round trips in X in baseline behavior was also associated to good olfactory scores (in agreement with WF being associated to significant individual response)(Suppl. Fig6A). Finally, F2 that exhibited a WF baseline pattern had better scores than others (like CF) (Figure 8A), and swimming speed was poorly correlated to their olfactory performances (not shown). Therefore, swimming patterns of CF and swimming speed of SF seem correlated to their behavioral response scores.

To further assess whether fish baseline swimming parameters are predictive of the behavioral output when presented with an odor, we next calculated how much of the odor response score variation can be explained by "personality". We found that the PVE (Percentage of Variation Explained) on the individual olfactory score varies significantly with the baseline swimming pattern in CF (but not in SF) (Figure 8C) and with the swimming speed in SF (but not in CF) (Figure 8D). Altogether, these analyses strongly suggest that (1) the fish swimming personality has an influence on its response to olfactory stimulation and can be predictive of the response, and (2) the personality parameter that is important to predict a good individual response is not the same in the two *Astyanax* morphs: in SF speed matters, while in CF swimming pattern matters.

Discussion

We have developed a high throughput, specific olfaction test, together with a pipeline of analysis allowing assessing qualitatively and quantitatively individual's and population's behaviors, in order to describe and compare responses of blind and sighted *Astyanax* to odorant cues. We discovered that cavefish, surface fish and F2 hybrids display different odor preferences and sensitivities and show individual, distinctive, diverse and specific responses to varying concentrations of the six amino acids tested. Using this novel setup where fish were tested in solo in a rectangular box and during one hour (as opposed to testing in groups in a U-shaped box and during 8 minutes in previous studies (Blin et al., 2020; Blin et al., 2018; Hinaux et al., 2016a), we established that cavefish are *bona fide* "alanine specialists" and we analyzed in depth their behavioral responses.

A setup to probe olfaction in fish with markedly different sensory apparatuses and internal states

There is an inherent difficulty to compare sensory-driven behaviors in cave and surface morphs of *Astyanax*: all their sensory systems have evolved in one way or another. Cavefish have no eyes, but they possess enhanced mechanosensory lateral line and chemosensory gustatory and olfactory organs. To decipher behavioral responses driven by a single of these senses, one needs to control carefully the potential influence of the other sensory modalities. Here, we have recorded unimodal behavioral responses driven exclusively by olfaction by performing experiments in the dark to abrogate vision in sighted fish, and by designing a setup where the delivery of the olfactory stimulus is vibration-free. Moreover, we know from previous studies that gustatory taste buds do not participate in responses to amino acids at the concentrations

used because the lesion of the olfactory epithelium abolishes the attraction to high

concentrations of alanine, in both SF and CF (Hinaux et al., 2016a).

Due to a mutation in their Monoamine Oxidase enzyme that interferes with the metabolism of

brain monoamines (Elipot et al., 2014), Pachón cavefish have lower basal cortisol levels, hence

lower basal anxiety than surface fish, when they are long habituated in their home tank (Pierre

et al., 2020). However, the mutation confers cavefish with a much higher stressability after

environmental change, such as the transfer in a novel tank (Pierre et al., 2020). Consequently,

in order to record relevant behavioral responses to olfactory stimuli in unstressed fish, we used

long acclimation times (72 hours) and we tested long habituation times, either 1 hour or 24

hours – as compared to 10 minutes as usually done in most fish studies including zebrafish or

Astyanax. We concluded that 1 hour is too short for proper habituation, as the swimming speed

and patterns were affected for both SF, CF and F2 as compared to 24 hours. Importantly, such

long, 24 hours habituation periods are susceptible to reveal unperturbed and even novel

behaviors: recently applied in a study of social behaviors in cavefish, 3 days of habituation

allowed the analysis of behaviors in a familiar environment and could unmask social

interactions in the so-called 'asocial' cavefish (Iwashita and Yoshizawa, 2021).

Neurophysiological and molecular considerations

In all vertebrates including fish, odorant molecules are recognized by olfactory receptors

expressed at the surface of olfactory sensory neurons, which project onto the olfactory

glomeruli in the olfactory bulb with the one receptor : one glomerulus rule (Axel, 1995;

Braubach et al., 2012; Buck, 2000; Kermen et al., 2013; Koide et al., 2009; Li et al., 2005;

Yoshihara, 2008). Moreover, parallel neural pathways in the olfactory circuitry process

different types of odorants. In zebrafish, the perception of amino acids is mediated via

OlfC/V2R receptors on microvillous sensory neurons that innervate lateral and ventro-medial glomeruli in the bulb. From the periphery to the brain, *Astyanax* surface and cave morphs display some variations in this amino acid signal processing circuitry. SF have 43 and CF have 41 V2R/OlfC receptor genes in their genomes (Policarpo et al., 2022), a minor difference that is unlikely to underlie their differences in olfactory capacities and preferences. CF larvae have higher proportions of microvillous neurons than SF, which together with the larger size of their olfactory epithelium and olfactory bulbs may influence their olfactory sensibility (Blin et al., 2018). *Astyanax* glomerular organization is unknown.

Interestingly, here we have found that CF are strongly attracted and respond to alanine and histidine, two amino acids which, albeit probably not recognized by the same receptor(s), are processed in the same or very close glomeruli in zebrafish larval olfactory bulbs (Li et al., 2005). SF on the other hand seem to show a preference for cysteine (sulfur), the amino acid that is the most potent to evoke electrical responses in the olfactory bulbs of sea breams (Hubbard et al., 2011) or hammerhead sharks (Tricas et al., 2009), and that elicits a strong aversive behavioral response in larval zebrafish (Vitebsky et al., 2005). Therefore, odor preferences have evolved between cavefish and surface fish, as well as between zebrafish and *Astyanax*. In the same line, previously we had reported that chondroitin is a strong attractant for *Astyanax* (Blin et al., 2020), whereas it induces freezing and fear behavior in zebrafish (Mathuru et al., 2012). Such significant variations in odor preferences or value may be adaptive and relate to the differences in the environmental and ecological conditions in which these different animals live. Of note, we have not found an odor that would be repulsive for *Astyanax* so far, and this may relate to their opportunist, omnivorous and detritivore regime (Espinasa et al., 2017; Marandel et al., 2020). However, the reason why Pachón cavefish have become "alanine specialists" remains a mystery and prompts analysis of the chemical ecology of their natural habitat. Alternatively, specialization for alanine may not need to be specific for an olfactory cue

present only, or frequently, or in high amounts in caves. Bat guano for example, which is probably the main source of food in the Pachón cave, must contain and release many amino acids. Enhanced recognition of only one of them - in the present case alanine but evolution may have randomly acted for enhanced recognition of another amino acid – should suffice to confer cavefish with augmented sensitivity to their main source of nutriment.

Cavefish have also evolved regarding olfactory sensitivity. They are able to detect very low concentrations of alanine (aliphatic), hence they have a lower detection threshold than SF. Injection of low concentration alanine $10^{-4}M/10^{-5}M$ (thus $10^{-6}M/10^{-7}M$ in the odorant third of the arena) elicits strong behavioral responses in CF, whereas even higher concentrations of $10^{-3}M$ and $10^{-2}M$ evoke modest responses in SF (zero response at population level, 19% responders at individual level with alanine $10^{-3}M$). We have performed dose-response experiments for 3 out of the 6 amino acids tested and for both morphs. With these, CF also appear capable of detecting low concentrations of cysteine (sulfur). Moreover, they seem to detect better histidine $10^{-2}M$ (polar) as well (69.2% of CF responders, versus 42.1% of SF), and CF but not SF detected lysine (polar) at the relatively high concentration of $10^{-2}M$. From these observations, we can predict that the difference between SF and CF probably does not lie in the molecular evolution of their olfactory receptor repertoire, because it is unlikely that all the receptors recognizing diverse types of amino acids have evolved all at once. Rather, we can hypothesize that evolution occurred at the level of the general regulation of odorant receptor genes expression, or at the level of olfactory processing and computation in the bulbs (Friedrich et al., 2009), or in higher order brain regions.

A variety of behavioral responses to olfactory stimulation

Our study suggests that behavioral responses to a unimodal olfactory sensory stimulus are

complex and have evolved in cavefish (summary on Table 1). When they detect the odor

stimulus, cavefish globally decrease their scanning activity, which might help them to compute

and swim up towards the higher values of concentration in the odor plume. This is associated

to a change in swimming pattern whereby WF and T are eliminated at the expense of R and C,

presumably to facilitate searching. This hypothesis is further supported by their systematic

change of position in the box, meaning that they locate efficiently the odor source. Strikingly,

surface fish display opposite behavioral responses after odor detection: they rather increase

locomotor activity, which corresponds to intense foraging but does not seem optimal to find the

odor source, which is confirmed by their lack of change in position of the box (i.e., they may

not locate efficiently the odor source). This is also consistent with the fact that at individual

level the best SF responders are slow swimmers. Such poor ability to find the odor source may

result from testing in the dark. Indeed, SF behaviors are mostly visually driven (but see Simon

et al., 2019) and they might need multimodal visual + olfactory integration to find food

efficiently. These interpretations are consistent with early work showing that, when competing

in the dark and with limited amounts of food, SF starve and CF thrive (Hüppop, 1987). In sum,

CF foraging strategy has evolved in response to the serious challenge of finding food in the

dark. Future experiments including functional imaging of brain activity in live animals may

reveal the changes in olfactory-driven motor circuits that allowed the evolution of behavioral

outputs in cavefish.

Finally, at population level, F2 fish shared some behavioral response traits with both parental

morphs, and they were often closer to CF. At individual level their responses were also variable

and non-stereotyped (including for alanine 10^{-3}M and histidine 10^{-2}M, the conditions for which

CF showed highly stereotyped responses). Olfactory scores behave as a quantitative trait. The

tools we have developed here will allow the future determination of the genetic underpinnings

of the evolution of olfactory-driven cavefish behaviors and capabilities.

A "personality" for each fish?

Our recordings of several hundred (n=489 total) well-habituated, individual *Astyanax* larvae

highlight an often overlooked aspect of fish behavioral analyses: fish may have a "personality"

or "temperament", which we characterized at the level of their baseline swimming patterns.

This hypothesis is strongly supported by the consistency over time, one day after the other, of

the favorite swimming pattern or combination of patterns and of the swimming kinetics

expressed by individuals when freely swimming without stimulus (most fish were recorded on

4 days distributed over 2 weeks). To our knowledge, this is the first time individual

temperament is taken into consideration in *Astyanax* behavioral studies.

In the context of ecology and evolution, the personality in non-human animals was proposed to

be organized along five primary axes: sociability, boldness, aggressiveness, exploration and

activity (Réale et al., 2007). In fish, studies applying an animal personality approach have

focused to resolve variations in physiological and molecular parameters, suggesting a link

between phenotype and genotype, or between behavior and transcriptome regulation eg (Rey et

al., 2021). Here, we propose that the different temperaments expressed by individual

larvae/juveniles could correspond to the genetic diversity present in the natural populations of

Astyanax, a diversity that we have maintained intentionally along generations in captivity in

our fish facility.

Further, we have assessed whether swimming temperaments could influence the way and the

extent to which fish respond to an olfactory sensory stimulus. Strikingly, we discovered that

baseline swimming patterns and swimming speed do influence fish olfactory responses. In

addition, important personality traits that confer positive behavioral responses are not the same in SF and CF. For cavefish, WF and round trips in X (but not speed) are the key parameters. Wall following behavior has been reported in *Astyanax* and other blind cavefish species (Chen et al., 2022; Patton et al., 2010; Sharma et al., 2009) and was hypothesized to confer foraging advantages, although this had never been directly tested. Here, it seems to be the case for 6-week old juveniles in a rectangular box - but the link may be more elusive when considering a fish swimming in a natural, complex environment. We propose that WF and round trips in X are together the expression of an intense exploratory behavior, which allows CF to better cover their swimming space and thus have a higher probability to detect odorant cues - even at very low concentrations because their detection capacities for some amino acids are excellent. By contrast, a slow swimming speed (but not swimming pattern) is the most critical personality parameter for SF. As SF olfactory concentration detection thresholds are higher than CF, one could imagine that they need more integration time inside the odorant plume to trigger a neuronal and a behavioral response. In any case and importantly, personality parameters that matter in CF and SF to confer good olfactory response scores are distinct. This suggests that the modulation of neuronal circuits underlying both the control of baseline behaviors and olfactory processing has evolved in cavefish.

Conclusion

Previous studies had shown that, when tested in groups, cavefish larvae as well as adults smell better than their surface fish conspecifics (Blin et al., 2020; Blin et al., 2018; Hinaux et al., 2016a). Here, we have established that it is also true for fish in solo, ruling out the possibility that when in group, one "good smeller" individual could drive others to respond through (unknown) communication mode. Using individual behavioral tests, we have discovered that

536   both olfactory sensitivity, olfactory preferences, olfactory behavioral responses and key

537   personality parameters have evolved in cavefish, conferring them with outstanding skills to

538   forage in the darkness of caves.

Methods

Fish samples

Laboratory stocks of *A. mexicanus* surface fish (SF, origin: San Salomon spring, Texas, USA) and cavefish (CF, Pachón population) were obtained in 2004 from the Jeffery laboratory at the University of Maryland, College Park, MD, USA. Fish were maintained at 22°C with a 12h:12h light:dark cycle, fed twice daily with dry and live food. SF and CF brood stock were induced to spawn once every two weeks by thermal shock at 26°C. A F1 hybrid family was generated by crossing a SF female with a CF Pachón male. Second-generation hybrids (F2) were generated by crossing two F1 individuals. Embryos and larvae were reared at 24°C in water produced by our animal house's water production system. All fish used (CF, SF and F2) were exactly six weeks old on the first day of the experiment. SR's authorization for use of *Astyanax mexicanus* in research is 91-116. Animals were treated according to the French and European regulations for animal testing in research (authorization n°APAFIS#8604).

Experimental set-up

The test consisted of filming individual larvae for 30 minutes before olfactory stimulation and for a further 30 minutes afterwards (see Supplemental movies). The behavior laboratory is a soundproofed room maintained at a constant temperature of 24° C. To ensure that eyed fish (SFs and F2 with eyes) could not use the visual modality during the test (unlike CFs and some F2), all experiments (except one) were carried out in the dark. The experimental device was composed of 2 infrared (IR) tables (60x60cm each) over which 32 test boxes were placed. A

custom-built wooden holder supported 8 IR cameras (DIGITNOW!), located 40 cm above the

boxes. Each camera filmed 4 boxes simultaneously (Figure 1).

The individual test boxes (11.5x8.5x4.3 cm, multiroir #45103BOILAB03) were made of

crystal-clear plastic. Odorant solutions (and/or water) were manually injected at the two

opposite sides of the box using 1ml syringes. In order to stop the dispersion wave of injection,

and to assure that we always injected at the same right place, we added 2 injection guides in

each box: we glued two plastic tubes (4 cm long and 1 cm in diameter) at the center of the two

opposite short sides (Y), 0.2 cm from the bottom. In addition, a net (6.5x6.5 cm) was attached

around each tube and reached the bottom of the box, to prevent fish from passing under the tube

during the experiment, thus escaping from the subsequent tracking (Figure 1).

Behavior tests

Larvae were placed individually in a box containing 150 ml of water (clean rearing water), fed

with micro-worms (three drops of concentrated *Panagrellus redivivus* solution placed in the

center of the box) and subjected to a 12:12h day:night cycle for 72 hours acclimation. Then, the

water in each box was changed, the boxes were placed on IR tables for 24 hours, and the larvae

were fasted (=habituation). The day of testing, for reasons of reproducibility, all experiments

started at 11 a.m. The light was switched off, the cameras were switched on and the

experimenter left the room for a 30-minute baseline behavior recording period. At t=30 min,

the experimenter entered the room and switched on the inactinic red lamps. Solutions were

manually injected bilaterally (500 µl of odor/ 500 µl of water) into the injection guides of each

box. The total time needed to perform injections of 32 boxes was 5 minutes. Then, the

experimenter switched off the inactinic lamps and left the room for a 30-minute response

behavior recording period. At the end of recording, the water was changed and each box was

placed back exactly at the same location on IR table for 24 hours and not moved again until the

following day of testing (the larvae were not fed). For the series of experiments with a

habituation period of 1 hour, the water in the boxes was changed every morning, 1 hour before

the start of the test. The typical timeline was as follows: [72 h acclimation and feeding], Week1

[habituation 24h, test day 1, habituation, test day 2, rest and feeding for three days], Week2

[habituation, test day 3, habituation, test day 4]. As the larvae were tested on several consecutive

days, to ensure that the fish were not learning/associating one side with a scent cue, the odor

injection side was reversed each day. The control experiments (unilateral or bilateral injection

of 500 µl rearing water or no injection) were always carried out the day1 when the larvae were

naive. For the "light vs dark" control experiment, SF were filmed 30 minutes in normal light

and 30 minutes in the dark.

Odorant solutions

The odorant solutions were prepared by dilution of amino acids in clean rearing water (L-

Alanine #A7627; L-Serine #S4500; L-Lysine #L5501; L-Histidine #H8000; L-Leucine #L8000

and L-Cysteine #168149; all from Sigma-Aldrich). A volume of 500µl (either water or odor)

was injected with a 1ml syringe (Sigma-Aldrich #Z683531) fitted with a needle (0.8x50mm,

21G; B Braun #4665503). Depending of the test, concentration of solutions injected were 10^{-2}

M or 10^{-3} M or 10^{-4} M. In a total water volume of 150ml, with a box virtually separated into

three, the volume of water in the odor injection zone is 50 ml. By injecting 500 µl of solution

into this zone, we expected to obtain a dilution factor of 100. For example, for a concentration

of [10^{-3} M] in the syringe, the concentration in the odor injection zone will therefore be ~[10^{-5}

M]. All concentrations given in the article indicate the concentration inside the syringe.

When consecutive injections of an odor or different odors at different concentrations were

tested on the same fish, they were performed from the least concentrated to the most

concentrated.

Video editing and tracking

Videos were saved on SD card in AVI format (1208x720 px, 30 fps), then edited using Adobe

Premiere Pro V14.0 to be calibrated at 1 hour long without the 5 minute-injection time. They

were then exported into MPG2 format. The tracking software used was TheRealFishTracker

V.0.4.0 (https://www.dgp.toronto.edu/~mccrae/projects/FishTracker/), developed and freely

available from the University of Toronto. In our hands, it was the only software able to detect

properly transparent CF larvae. The parameters used were Confidence Threshold = 10; Mean

Filter Size = 1; signed Image = *dark object* and all other parameters were the default ones. The

x/y-scales were drawn and given in cm. The output data file provided 30 X/Y coordinates (in

pixels and in cm) per seconds for a 1-hour movie.

Graphical representations, quantifications and statistics

Statistical analyses were carried out with R-3.4.2 software (R Development Core Team, 2016)

using the stats and rstatix libraries, and all graphical representations were designed using the

ggplot2, ggrepel, gghighlight and RGraphics libraries.

Quantitative parameters. From the output data set, pixel coordinates X_{pi} and Y_{pi} were first

recoded using the box center as 0, $X_{cmax} = 1$ on the odor side and $X_{cmin} = -1$ on the water

side. The same was applied to the Y coordinate with Ycmax = 0.74 and Ycmin = -0.74. Fish

position in the box along the X-axis during test was calculated using Xc and Yc at each time

point (30/s). Speed between two time points (5-second time step) was calculated thanks to

pythagoras's theorem using X/Y in cm. Round trip number is the total number of time the fish

crosses the 0.5 and -0.5 position (along X-axis) divided by two. Idem for round trips in Y-axis.

Quantitative parameter means were all calculated over a 15-minute period, from minute 10 to

minute 25 for the 'before' period, and from minute 37 to minute 52 for the 'after' period. Means

are represented by box plots showing the distribution of individual values of each fish (black

dots), the median, the 25^{th}- 75^{th} percentile and outliers indicated by fish ID number. Paired

Mann-Whitney two-tailed tests were performed to compare means of position, speed, or round

trips before and after odor injection.

Non-paired Mann-Whitney two-tailed tests were performed to compare means of speed

between 1h and 24h habituation periods represented in Figure 2C by box plots.

Swimming patterns. The swimming activity of the fish in the test box was systematically

represented under 3 different forms, which allow grasping the details and different aspects of

the behaviors (see Figure 2A). Xc /t coordinates were used with the geom_line() function for

the 'Position (X-axis)' graph, (Xc,Yc) /t coordinates were used with the geom_path() function

for '2D' top view, and the gg3D package for '2D + time' view.

The determination of baseline swimming patterns and swimming patterns after odor injection

was performed manually based on graphical representations such as in Figure 2A or Figure 3A.

Four distinctive baseline behaviors clearly emerged. (1) Random swim (R; defined as

haphazard swimming with no clear pattern, covering entirely or partly the surface of the arena).

(2) Wall following (WF; defined as the fish continuously following along the 4 sides of the box

and turning around it, in a clockwise or counterclockwise fashion). (3) Large or small circles

(C; self-explanatory). (4) Thigmotactism (T, along the X- or the Y-axis of the box; defined as

the fish swimming back and forth along one of the 4 sides of the box). On graphical

representations of swimming pattern distributions, we used the following color code: R in blue,

WF in red, C in green, T in yellow. Of note, many fish swam according to combination(s) of

these four elementary swimming patterns (see descriptions in the legends of Supplemental

figures showing many examples). To fully represent the diversity and the combinations of

swimming patterns used by individual fish, we used an additional color code derived from the

"basic" color code described above and where, for example R+WF is purple. The complete

combinatorial color code is shown in Suppl. Fig7. Of note, in all figures, the swimming pattern

color code does not relate whatsoever with the time color code used in the 2D plus time

representation of swimming tracks such as in Figure 2A. Swimming patterns (i.e., qualitative

variables; compositional dataset) in different conditions were analyzed using Fisher's exact test

comparing the quantity and distribution of different swimming patterns between SF/CF/F2 at

day1, or over day1 to day4, or between 1h *vs* 24 h habituation period, or before *vs* after odor

injection for each morphotype.

To reinforce our conclusions, swimming patterns in different conditions were also compared

using CA (Correspondence Analysis), an appropriate method to analyze compositional data

(Greenacre, 2021). The results are plotted in Suppl. Fig8. The distribution of swimming patterns

in SF/CF/F2 in control experiments, or before *vs* after odor injection for each morphotype

confirm the changes (or absence of changes) in behavioral patterns suggested by the colored

bar plots in main Figures, with confidence ellipses overlapping or not overlapping, depending

on cases. Of note, CA analysis cannot provide statistical support through calculation of p values.

Index and score.

For quantitative swimming parameters: indexes of position, speed and number of round trips changes were calculated using the equation (mean *after* – mean *before*) / (mean *after* + mean *before*) commonly used in olfactory tests in fish (Choi et al., 2021; Koide et al., 2009; Wakisaka et al., 2017). The index can vary from zero to 1. An index close to zero indicates that the values of the considered parameter before or after odor injection are close. The higher the absolute value of the index, the greater the difference. When the parameter increases after injection, the index is positive; when the parameter decreases after injection, the index is negative. The *absolute* value of each index will be used for score calculation below.

For qualitative swimming parameter: the index of pattern change was calculated as the number of new patterns observed after stimulation, divided by the total number of patterns observed after odor injection. An index equal to zero indicates no pattern change after stimulation. The higher the index is, the greater the difference of pattern is.

For each fish, the individual olfactory score for a given odor and concentration was the sum of the *absolute* values of the four indexes: individual score = |position index| + |speed index| +|round trips in X+Y index| +|pattern index|. On Figure 7A and Suppl. Fig5, the 4 parameter's indexes and the total individual fish olfactory scores are represented by dots connected by a colored line for each individual. Those with a score > 1.5 are highlighted and considered to display a significant behavioral response to the stimulus. The 1.5 threshold value was arbitrarily determined after the examination of individual olfactory scores in control experiments (Suppl. Fig5) and the observation that the vast majority of fish have individual scores below this value in the absence of specific olfactory stimulus.

On Figure 8A and 8C, to assess the relationship between individual olfactory score and baseline swimming pattern, we pooled results of odors for which more than 40% of individuals have an

olfactory score >1.5. So for SF, alanine [10^{-2}M] / cysteine [10^{-2}M] / histidine [10^{-2}M] / serine

[10^{-2}M] have been used; for CF, alanine [10^{-2}M] / alanine [10^{-3}M] / alanine [10^{-4}M]/ cysteine

[10^{-2}M] / cysteine [10^{-3}M] / histidine [10^{-2}M] / lysine [10^{-2}M]; and for F2, alanine [10^{-3}M] /

cysteine [10^{-2}M] / histidine [10^{-2}M] (see Table 1B). Mann-Whitney two-tailed with Bonferroni

post hoc tests were performed on each morphotype.

Slopes in linear regressions show relationships between mean speed or number of round trips

before injection and olfactory score. Linear correlations were calculated with Spearman's rank

correlation test followed by student test for the p.values.

Supplementary information and data availability. The R script used to recode raw data into

corrected coordinates, to calculate means and to draw 2D and 2D + time graphs is available on

GitHub (https://github.com/Maryline-Blin/Astyanax-swimming-pattern).

F1 to **F3 Files** correspond to source data files (.txt) containing the raw data presented and

analyzed in this paper. For each type, the files provide the ID numbers of the fish, condition

(day1 to day4 or odor or control type), means of position, speed, X and Y round-trips and

behavior for periods 10-25 minutes and 37-52 minutes, as well as the calculated individual

olfactory score.

Acknowledgments

We thank Camille Lejeune for her participation in the generation of the SF x CF F1 hybrid line, Maxime Policarpo for preliminary testing of tracking softwares and Rose Tatarsky for interesting discussions. Zootechnical care of our *Astyanax* fish facility was taken in charge by the TEFOR Unit. We are indebted to Joël Attia for invaluable help with statistical analyses during the revision of our manuscript. This work was supported by Equipe FRM (Fondation pour la Recherche Médicale) grant EQU202003010144 to SR.

References

Agnès, F., Torres-Paz, J., Michel, P. and Rétaux, S. (2022). A 3D molecular map of
the cavefish neural plate illuminates eye-field organization and its borders in vertebrates.
Development **149**, dev199966.
Axel, R. (1995). The molecular logic of smell. *Sci Am* **273**, 154-9.
Blin, M., Fumey, J., Lejeune, C., Policarpo, M., Leclercq, J., Père, S., Torres-Paz,
J., Pierre, C., Imarazene, B. and Rétaux, S. (2020). Diversity of Olfactory Responses and
Skills in Astyanax Mexicanus Cavefish Populations Inhabiting different Caves. *Diversity* **12**,
1-21.
Blin, M., Tine, E., Meister, L., Elipot, Y., Bibliowicz, J., Espinasa, L. and Rétaux,
S. (2018). Developmental evolution and developmental plasticity of the olfactory epithelium
and olfactory skills in Mexican cavefish. *Dev Biol* **441**, 242-251.
Braubach, O. R., Fine, A. and Croll, R. P. (2012). Distribution and functional
organization of glomeruli in the olfactory bulbs of zebrafish (Danio rerio). *J Comp Neurol* **520**,
2317-39, Spc1.
Buck, L. B. (2000). The molecular architecture of odor and pheromone sensing in
mammals. *Cell* **100**, 611-8.
Burguera, D., Dionigi, F., Kverkova, K., Winkler, S., Brown, T., Pippel, M., Zhang,
Y., Shafer, M., Nichols, A. L. A., Myers, E. et al. (2023). Expanded olfactory system in ray-
finned fishes capable of terrestrial exploration. *BMC Biol* **21**, 163.
Chen, B., Mao, T., Liu, Y., Dai, W., Li, X., Rajput, A. P., Pie, M. R., Yang, J., Gross,
J. B. and Meegaskumbura, M. (2022). Sensory evolution in a cavefish radiation: patterns of
neuromast distribution and associated behaviour in Sinocyclocheilus (Cypriniformes:
Cyprinidae). *Proc Biol Sci* **289**, 20221641.
Choi, J. H., Duboue, E. R., Macurak, M., Chanchu, J. M. and Halpern, M. E.
(2021). Specialized neurons in the right habenula mediate response to aversive olfactory cues.
Elife **10**.
de Busserolles, F., Fogg, L., Cortesi, F. and Marshall, J. (2020). The exceptional
diversity of visual adaptations in deep-sea teleost fishes. *Semin Cell Dev Biol* **106**, 20-30.
Duboué, E. R. and Keene, A. (2016). Investigating the evolution of sleep in the
Mexican cavefish. *Biology and Evolution of the Mexican Cavefish. Chapter 15.* **Eds AC Keen,**
M Yoshizawa, SE McGaugh. Elsevier, Academic Press., 335-359.
Elipot, Y., Hinaux, H., Callebert, J., Launay, J., Blin, M. and Rétaux, S. (2014). A
Mutation in the Enzyme MonoAmine Oxidase Explains Part of the Astyanax Cavefish
Behavioral Syndrome. *Nature Communications* **Apr 10; 5:3647.**
Espinasa, L., Bonaroti, N., Wong, J., Pottin, K., Queinnec, E. and Rétaux, S. (2017).
Contrasting feeding habits of post-larval and adult Astyanax cavefish. *Subterranean Biology*
21, 1-17.
Friedrich, R. W., Yaksi, E., Judkewitz, B. and Wiechert, M. T. (2009). Processing
of odor representations by neuronal circuits in the olfactory bulb. *Ann N Y Acad Sci* **1170**, 293-
7.
Greenacre, M. (2021). Compositional Data Analysis. *Annual Review of Statistics and*
Its Application **8**, 271-299.
Hara, T. J. (1994). Olfaction and gustation in fish: an overview. *Acta Physiol Scand*
152, 207-17.
Helfman, G. S., Collette, B. B., Facey, D. E. and Bowen, B. W. (2009). The diversity
of fishes. Biology, Evolution, and Ecology. Second Edition. *Wiley-Blackwell*.

Hinaux, H., Devos, L., Bibliowicz, J., Elipot, Y., Alié, A., Blin, M. and Rétaux, S. (2016a). Sensory evolution in blind cavefish is driven by early events during gastrulation and neurulation. *Development* **143**, 4521-4532.

Hinaux, H., Rétaux, S. and Elipot, Y. (2016b). Social behavior and aggressiveness in Astyanax. *Biology and Evolution of the Mexican Cavefish. Chapter 17.* **Eds AC Keen, M Yoshizawa, SE McGaugh. Elsevier, Academic Press.**, 335-359.

Hubbard, P. C., Barata, E. N., Ozorio, R. O., Valente, L. M. and Canario, A. V. (2011). Olfactory sensitivity to amino acids in the blackspot sea bream (Pagellus bogaraveo): a comparison between olfactory receptor recording techniques in seawater. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* **197**, 839-49.

Hüppop, K. (1987). Food-finding ability in cave fish (Astyanax fasciatus). *Int. J. Speleol.* **16**, 59-66.

Iwashita, M. and Yoshizawa, M. (2021). Social-like responses are inducible in asocial Mexican cavefish despite the exhibition of strong repetitive behavior. *Elife* **10**.

Kalueff, A. V., Gebhardt, M., Stewart, A. M., Cachat, J. M., Brimmer, M., Chawla, J. S., Craddock, C., Kyzar, E. J., Roth, A., Landsman, S. et al. (2013). Towards a comprehensive catalog of zebrafish behavior 1.0 and beyond. *Zebrafish* **10**, 70-86.

Kermen, F., Franco, L. M., Wyatt, C. and Yaksi, E. (2013). Neural circuits mediating olfactory-driven behavior in fish. *Front Neural Circuits* **7**, 62.

Kiyohara, S., Sakata, Y., Yoshitomi, T. and Tsukahara, J. (2002). The 'goatee' of goatfish: innervation of taste buds in the barbels and their representation in the brain. *Proc Biol Sci* **269**, 1773-80.

Koide, T., Miyasaka, N., Morimoto, K., Asakawa, K., Urasaki, A., Kawakami, K. and Yoshihara, Y. (2009). Olfactory neural circuitry for attraction to amino acids revealed by transposon-mediated gene trap approach in zebrafish. *Proc Natl Acad Sci U S A* **106**, 9884-9.

Kowalko, J. (2020). Utilizing the blind cavefish Astyanax mexicanus to understand the genetic basis of behavioral evolution. *J Exp Biol* **223**.

Krubitzer, L., Campi, K. L. and Cooke, D. F. (2011). All rodents are not the same: a modern synthesis of cortical organization. *Brain Behav Evol* **78**, 51-93.

Li, J., Mack, J. A., Souren, M., Yaksi, E., Higashijima, S., Mione, M., Fetcho, J. R. and Friedrich, R. W. (2005). Early development of functional spatial maps in the zebrafish olfactory bulb. *J Neurosci* **25**, 5784-95.

Lunsford, E. T., Paz, A., Keene, A. C. and Liao, J. C. (2022). Evolutionary convergence of a neural mechanism in the cavefish lateral line system. *Elife* **11**.

Marandel, L., Plagnes-Juan, E., Marchand, M., Callet, T., Dias, K., Terrier, F., Pere, S., Vernier, L., Panserat, S. and Rétaux, S. (2020). Nutritional regulation of glucose metabolism-related genes in the emerging teleost model Mexican tetra surface fish: a first exploration. *R Soc Open Sci* **7**, 191853.

Mathuru, A. S., Kibat, C., Cheong, W. F., Shui, G., Wenk, M. R., Friedrich, R. W. and Jesuthasan, S. (2012). Chondroitin fragments are odorants that trigger fear behavior in fish. *Curr Biol* **22**, 538-44.

Patton, P., Windsor, S. and Coombs, S. (2010). Active wall following by Mexican blind cavefish (Astyanax mexicanus). *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* **196**, 853-67.

Pierre, C., Pradère, N., Froc, C., Ornelas-Garcia, P., Callebert, J. and Rétaux, S. (2020). A mutation in monoamine oxidase (MAO) affects the evolution of stress behavior in the blind cavefish Astyanax mexicanus. *J Exp Biol* **223**.

Policarpo, M., Bemis, K. E., Laurenti, P., Legendre, L., Sandoz, J. C., Rétaux, S. and Casane, D. (2022). Coevolution of the olfactory organ and its receptor repertoire in ray-finned fishes. *BMC Biol* **20**, 195.

Pottin, K., Hinaux, H. and Rétaux, S. (2011). Restoring eye size in Astyanax mexicanus blind cavefish embryos through modulation of the Shh and Fgf8 forebrain organising centres. *Development* **138**, 2467-76.

R Development Core Team. (2016). R: a language and environment for statistical computing. *R Foundation for Statistical Computing, Vienna, Austria* **http://www.R-project.org**.

Réale, D., Reader, S. M., Sol, D., McDougall, P. T. and Dingemanse, N. J. (2007). Integrating animal temperament within ecology and evolution. *Biol Rev Camb Philos Soc* **82**, 291-318.

Rey, S., Jin, X., Damsgard, B., Begout, M. L. and Mackenzie, S. (2021). Analysis across diverse fish species highlights no conserved transcriptome signature for proactive behaviour. *BMC Genomics* **22**, 33.

Schemmel, C. (1967). Vergleichende Untersuchungen an den Hautsinnesorganen ober- and unter-irdisch lebender Astyanax-Foramen. . *Z Morph Tiere* **61**, 255-316.

Sharma, S., Coombs, S., Patton, P. and Burt de Perera, T. (2009). The function of wall-following behaviors in the Mexican blind cavefish and a sighted relative, the Mexican tetra (Astyanax). *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* **195**, 225-40.

Simon, V., Hyacinthe, C. and Rétaux, S. (2019). Breeding behavior in the blind Mexican cavefish and its river-dwelling conspecific. *PLoS One* **14**, e0212591.

Sylvester, J. B., Rich, C. A., Loh, Y. H., van Staaden, M. J., Fraser, G. J. and Streelman, J. T. (2010). Brain diversity evolves via differences in patterning. *Proc Natl Acad Sci U S A* **107**, 9718-23.

Torres-Paz, J., Leclercq, J. and Rétaux, S. (2019). Maternally regulated gastrulation as a source of variation contributing to cavefish forebrain evolution. *Elife* **8**.

Tricas, T. C., Kajiura, S. M. and Summers, A. P. (2009). Response of the hammerhead shark olfactory epithelium to amino acid stimuli. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* **195**, 947-54.

Varatharasan, N., Croll, R. P. and Franz-Odendaal, T. (2009). Taste bud development and patterning in sighted and blind morphs of Astyanax mexicanus. *Dev Dyn* **238**, 3056-64.

Vitebsky, A., Reyes, R., Sanderson, M. J., Michel, W. C. and Whitlock, K. E. (2005). Isolation and characterization of the laure olfactory behavioral mutant in the zebrafish, Danio rerio. *Dev Dyn* **234**, 229-42.

Wakisaka, N., Miyasaka, N., Koide, T., Masuda, M., Hiraki-Kajiyama, T. and Yoshihara, Y. (2017). An Adenosine Receptor for Olfaction in Fish. *Curr Biol* **27**, 1437-1447 e4.

Yamamoto, Y., Byerly, M. S., Jackman, W. R. and Jeffery, W. R. (2009). Pleiotropic functions of embryonic sonic hedgehog expression link jaw and taste bud amplification with eye loss during cavefish evolution. *Dev Biol* **330**, 200-11.

Yoshihara, Y. (2008). Molecular Genetic Dissection of the Zebrafish Olfactory System. *Results Probl Cell Differ*, 97-119.

Yoshizawa, M. (2016). The evolution of sensory adaptation in Astyanax mexicanus. *Biology and Evolution of the Mexican Cavefish. Chapter 13.* **Eds AC Keen, M Yoshizawa, SE McGaugh. Elsevier, Academic Press.**, 335-359.

Yoshizawa, M., Goricki, S., Soares, D. and Jeffery, W. R. (2010). Evolution of a behavioral shift mediated by superficial neuromasts helps cavefish find food in darkness. *Curr Biol* **20**, 1631-6.

Yoshizawa, M. and Jeffery, W. R. (2011). Evolutionary tuning of an adaptive behavior requires enhancement of the neuromast sensory system. *Communicative and Integrative Biology* **4**, 89-91.

879   **Yoshizawa, M., Jeffery, W. R., van Netten, S. M. and McHenry, M. J.** (2014). The
880   sensitivity of lateral line receptors and their role in the behavior of Mexican blind cavefish
881   (Astyanax mexicanus). *J Exp Biol* **217**, 886-95.

Figures and legends

Figure 1: experimental setup for testing behavioral responses to olfactory stimulation in individual, 6 weeks old, *Astyanax mexicanus* larvae.

A, Each fish is placed individually in 150ml water in a rectangular testbox placed on an infrared light emitting table. Control or odorant solution are delivered at the two extremities of the box, inside tubes covered with a net. Medium to high throughput behavioral testing is achieved by parallel recording of 32 testboxes placed under 8 infrared recording cameras (4 testboxes per camera).

B, After one hour or 24 hours habituation, the test consists of one hour IR video recording. The first 30 minutes provide the control/baseline behavior of individual fish. Odorant stimulation is given at 30 minutes, and the behavioral responses are recorded for 30 more minutes.

Figure 2: basal swimming behaviors of 6 week-old CF, SF and F2.

A, Examples of typical and distinctive basal swimming patterns exhibited by *Astyanax mexicanus* morphotypes. The behavioral patterns (Random, Wall Following, Circles, Thigmotactism) and the individual fish shown are indicated. Each pattern is best described by the combination of 3 types of representations. (1) the top left track shows the displacement of the fish along the x-axis of the box, representing back-and-forth swims along its long dimension. (2) the bottom left track is the top view, 2D representation of the trajectory. (3) the right track is the 2D plus time (color-coded) representation and helps, for example, to discern between wall following and thigmotactism.

B, Distribution of basal swimming pattern displayed by SF, F2, and CF after 1h or after 24h of habituation. Numbers of fish tested are indicated. The elementary color code is indicated (R in blue, WF in red, C in green, T in yellow). Exact swimming patterns and combinations are given on the colored plots (Suppl. Fig7 for the full color code). Fisher's exact tests for statistical comparisons between groups. Of note, in this and subsequent figures, the swimming pattern color code does not relate whatsoever with the time color code used in the 2D plus time representation of swimming tracks such as in panel A.

C, Boxplots showing the swimming speed in SF (blue), F2 (green) and CF (red), after 1h or 24h of habituation. Values are mean speed calculated over a period of 15 minutes. Numbers of fish tested are indicated. Mann-Whitney tests p-values are shown.

D, Examples of the stability of the basal swimming pattern over 4 experimental days in 3 individuals, with 24h of habituation time. One SF displaying Random pattern, one CF displaying Wall Following and one F2 displaying Thigmotactism + Random swim are shown.

Figure 3: Controls on the olfactory setup and solution delivery method.

A, Lack of behavioral response in a representative CF individual after injection of water on one side of the testbox.

B, Lack of change in qualitative swimming pattern displayed by SF, CF and F2, before and after water injection on one side. Fisher's exact tests.

C, Lack of change in quantitative swimming kinetics in SF (blue), CF (red) and F2 (green), before (pale colors) and after (dark colors) water injection on one side. In this and the following graphs, boxplots show the swimming speed, the fish position along the X-axis of the testbox, and the numbers of back-and-forth swims along the X (length) and Y (width) axes of the box. All values are averaged over a 15 minutes period, either before or after the injection. A thin line links the value « before » and the value « after », for each individual. Numbers of fish tested are indicated on the "speed" boxplots. Numbers next to dots indicate the identity of outlier individuals.

Figure 4: Behavioral responses to alanine of 6 week-old CF, SF and F2.

A,B,C: Representative individual responses of CF, SF and F2 after injection of alanine at the indicated concentrations. In the two left graphs of each panel, the blue color indicates the water/control injection side and the yellow color indicates the alanine injection side.

D,E,F,G: Box plots showing swimming speed (D), mean position along the X-axis (E) and the number of back and forth trips in X and Y (F,G) in SF (blue), CF (red) and F2 (green), before (lighter color) or after (darker color) the injection of alanine at the indicated concentration. Values are calculated over a 15 minutes period. Black lines link the « before odor » and « after odor » value for each individual. Numbers close to black dots indicate the identity of outlier individuals. P-values from paired Mann-Whitney tests are shown. The number of fish tested is indicated.

H, Change in swimming patterns exhibited by SF and CF after injection of alanine 10^{-4}M. Fisher's exact tests.

Figure 5: Behavioral responses to serine and cysteine 10^{-2}M.

A,B and **H,I**, Representative individual responses of CF and SF after injection of serine (A,B) or cysteine (H,I) at high concentration (10^{-2}M). In the two left graphs, the blue color indicates the water/control injection side and the green (serine) or red (cysteine) color indicates the injection side.

C-F and **J-M**, Box plots showing swimming speed (C,J), mean position along the X-axis (D,K) and the number of back and forth trips in X and Y (EL and FM) in SF (blue), CF (red) and F2 (green), before (lighter color) or after (darker color) injection of serine 10^{-2}M (C-F) or cysteine

10^{-2}M (J-M). Values are calculated over a 15 minutes period. Black lines link the « before odor » and « after odor » value of each individual fish. Numbers close to black dots indicate the identity of outlier individuals. P-values from paired Mann-Whitney tests are shown. The number of fish tested is indicated.

G and **N**, Change in swimming patterns elicited after injection of 10^{-2}M serine (G) or cysteine (N) in SF and CF (F2 not shown). Fisher's exact tests.

Figure 6: Behavioral responses to lysine, histine and leucine 10^{-2}M.

A-D, Box plots showing swimming speed (A), mean position along the X-axis (B) and the number of back and forth trips in X and Y (CD) in SF (blue), CF (red) and F2 (green), before (lighter color) or after (darker color) the injection of the indicated amino acid. Values are calculated over a 15 minutes period. Black lines link the « before odor » and « after odor » value of each individual fish. Numbers close to black dots indicate the identity of outlier individuals. P-values from paired Mann-Whitney tests are shown. The number of fish tested is indicated. See **Supplemental Figure 4** for examples of representative individual responses.

E-G, Change in swimming patterns elicited after injection of 10^{-2}M lysine (E) or histidine (F) or leucine (G) in SF, CF and F2. Fisher's exact tests.

Table 1. Summary of the results.

A, Population level summary. For each amino acid and each concentration tested, arrows indicate whether the considered parameter has changed (increased or decreased). ns indicates no significant change, - indicates the condition was not tested in the fish type (SF in blue, CF in red, F2 in green).

B, Individual level summary. The percentage of fish displaying an individual olfactory score superior to 1.5 is indicated.

Figure 7: Individual olfactory scores of SF, CF and F2 for different odors.

A, Graphs representing index values (i.e., the variation between the 'before' and the 'after' odor condition; the value 0 corresponds to no change) for the 4 response parameters (speed, thigmotactism in X and Y, position and swim pattern) used to calculate the total individual olfactory score for each fish (last column of points on each graph). Each fish is depicted by a colored line linking its 4 different indexes and its final individual olfactory score. The dotted line at value 1.5 indicates the score threshold above which a fish is considered as a responder. The percentages in black and grey indicate the proportion of responders/non-responders, respectively. The colored lines of responders are bright, those of non-responders are pale. Amino acids and concentrations are indicated. Top row: CF; bottom row: SF.

B, Distributions of individual olfactory scores of SF (blue), CF (red) and F2 (green) for different odors. The threshold score (1.5) is indicated by a dotted line.

Figure 8: The individual olfactory score is related to individual fish swimming personality.

A, Box plots showing individual olfactory scores in SF (blue), F2 (green) and CF (red) as a function of their individual baseline swimming pattern. As stated in Methods, only amino acid conditions for which more than 40% of fish were responders (score >1.5) are pooled and plotted on this graph. Mann-Whitney two-tailed with Bonferroni post hoc tests were performed on each morphotype.

B, Regressions to explore the correlation between baseline swimming speed displayed by individual fish before odor injection and their individual olfactory score. Fish and amino acid type and concentrations are indicated. Linear correlations were calculated with Spearman's rank correlation test followed by Student test for the p values. Conditions for which significant correlation was found are labelled with an asterisk.

C, The PVE (Percentage of Variation Explained) on the individual score is plotted as a function of individual baseline behavior, to probe the predictability of the behavioral output as a function of "swimming pattern personality".

D, The PVE (Percentage of Variation Explained) on the individual score is plotted as a function of individual swimming speed, to probe the predictability of the behavioral output as a function of "locomotor activity personality". Linear correlations were calculated with Spearman's rank correlation test followed by Student test for the p values. Conditions for which significant correlation was found are labelled with an asterisk. SF (blue), F2 (green) and CF (red).

A Setup

x32

Infrared camera

500 µl
odor

500 µl
water

Single larva

Injection tube

Net

V = 150 ml

Y=8.5cm

X=11.5cm

Infra-red
light table

B Experimental timeline

odor/water
injection

0 min

30 min=
baseline behaviour

30 min

30 min=
olfactory response behaviour

60 min

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Table 1

	alanine 10⁻⁵M	alanine 10⁻⁴M	alanine 10⁻³M	alanine 10⁻²M	serine 10⁻⁴M	serine 10⁻³M	serine 10⁻²M	cysteine 10⁻⁴M	cysteine 10⁻³M	cysteine 10⁻²M	lysine 10⁻²M	histidine 10⁻²M	leucine 10⁻²M
A — population													
Speed													
SF	-	ns	ns	ns	-	↗	ns	↗	↗	↗	ns	ns	ns
F2	-	-	↘	-	-	ns	ns	-	-	ns	↗	ns	ns
CF	ns	ns	↘	ns	ns	ns	ns	ns	ns	ns	ns	↘	ns
Position													
SF	-	ns	ns	ns	-	ns	ns	ns	↘	ns	ns	ns	ns
F2	-	-	↗	-	-	↗	↗	-	-	ns	ns	↗	ns
CF	↗	↗	↗	↗	ns	↗	↗	↗	↗	↗	ns	↗	↗
NbrtX													
SF	-	ns	ns	ns	-	ns	↗	↗	↗	ns	ns	↗	↗
F2	-	-	↘	-	-	ns	ns	-	-	ns	ns	ns	ns
CF	ns	↘	↘	↘	ns	↘	↘	ns	↘	ns	↘	↘	↘
NbrtY													
SF	-	↗	ns	ns	-	ns	↗	↗	↗	↗	ns	↗	ns
F2	-	-	↘	-	-	ns	↗	-	-	ns	↗	ns	ns
CF	ns	↘	↘	↘	ns	ns	ns	ns	ns	ns	↘	↘	ns
Patterns													
SF	-	ns	ns	ns	-	ns	change	change	ns	change	ns	ns	ns
F2	-	-	change	-	-	ns	change	-	-	change	ns	change	ns
CF	change	change	change	change	ns	ns	change	ns	change	change	change	change	ns
B — individual													
Score > 1.5													
SF	-	3.2%	19.3%	42%	-	25%	41.1%	18.7%	37.5%	51%	21%	42.1%	21%
F2	-	-	50%	-	-	31.3%	18.8%	-	-	55%	18.7%	55%	36.9%
CF	25%	81.3%	73.2%	56.3%	18.7%	6.2%	35.7%	31%	43.8%	39.5%	48.4%	69.2%	28%

Figure 7

Figure 8

Evolution of olfactory sensitivity, preferences and behavioral responses in Mexican cavefish: fish personality matters

Maryline Blin, Louis Valay, Manon Kuratko, Marie Pavie and Sylvie Rétaux

Suppl Figure 1

Suppl Fig1 (with Figure 3). Additional controls for experimental design.

AA', Distribution of swimming patterns exhibited by SF, F2 and CF (number of fish tested is indicated) along 4 testing days, with 24 hours habituation time (**A**). Fisher's exact tests. The color code for the different swimming patterns and combinations is given in the inset below. The table in **A'** shows statistics summary for paired comparison of individual's mean position, swimming speed and the total number of round trips in X plus Y over 4 testing days, using Friedman tests. The results of additional pairwise Wilcoxon comparisons between different days are as follows. All these pairwise comparisons were NS except for CF roundtrip numbers (d1/d2 p=0.04; d1/d3 p=0.0009; all other comparisons NS) and F2 position (d1/d3 p= 0.022; d3/d4 p= 0.042). These results suggest that fish display individual swimming behaviors that are stable through time and may be considered as individual swimming personality or temperament.

B, Lack of change in swimming pattern displayed by SF, CF and F2, before and after water injection on both sides. Fisher's exact tests.

C, Global lack of change in quantitative swimming kinetics in SF (blue), F2 (green) and CF (red), before and after water injection on both side. Mean speed and mean position are shown.

D, Lack of change in quantitative swimming kinetics in SF, CF and F2 when no injection is performed at all. Mean speed and mean position are shown. The result suggests that an individual's behavior is stable throughout the one-hour test.

E, Lack of change in swimming speed or position on the box when SF are recorded in the dark (dark blue), in the light (light blue), or when the light is switched off after 30 minutes.

A Pachón cavefish, alanine 10⁻³M

B Surface fish, alanine 10⁻²M

C F2 hybrids, alanine 10⁻³M

D

Alanine 10⁻²M:

Alanine 10⁻³M:

Alanine 10⁻⁵M:

WF · C · R · T

Swimming patterns

Suppl Figure 2

Suppl Fig2 (with Figure 4). Additional examples of behavioral responses to alanine

A, Examples of CF responses to alanine 10^{-3}M (intermediate concentration). Note the reproducible switch in position toward the odorant side in the box (yellow shading) and the change in behavioral pattern. Note that CF n°8, which was tested twice with the same concentration of alanine, switched from wall following to small circles and thigmotactism along the Y border of the box in both cases. CF n°4 on the other hand swam in large circles before injection, and changed its pattern to display intense thigmotactism accompanied with a few circles.

B, Examples of SF responses to alanine 10^{-2}M (high concentration). Upon injection, SF show changes in swimming patterns and increase their swimming speed, suggesting that the stimulus was perceived. SF n°22 swam very actively randomly. SF n°4 switched from random with small circles to circles plus thigmotactism. SF n°12 added thigmotactism to its random pattern upon stimulus. Some SF, like SF n°8, did not seem to respond, even at such high alanine concentration.

C, Examples of F2 hybrids responses to alanine 10^{-3}M (intermediate concentration). F2 n°14 did not appear to move its position to the odorant part of the box, but a striking change in swimming pattern from intense thigmotactism to random swim was observed. F2 n°7 explored close to the odor source and decreased swimming speed. F2 n°3 and 1 switched from wall following to circles with thigmotactism associated to a change in position towards the odor side.

D, Change in swimming patterns displayed by SF, F2 and CF after injection of alanine 10^{-2}M, 10^{-3}M and 10^{-5}M. Fisher's exact tests.

A Responses to serine 10⁻²M

CF n°27

SF n°40

F2 n°71

F2 n°79

B Responses to cysteine 10⁻²M and 10⁻³M

CF n°73
Cys 10⁻³M

CF n°71
Cys 10⁻²M

CF n°72
Cys 10⁻²M

SF n°34
Cys 10⁻²M

Suppl Figure 3

Suppl Fig3 (with Figure 5). Additional examples of behavioral responses to serine and cysteine

A, responses to serine 10^{-2}M (high concentration). Note the change in behavioral pattern and position in the box observed in CF and F2 but not in SF. The two F2 shown had different baseline swimming patterns: circles in F2 n°71, thigmotactism in F2 n°79. Yet, both individuals showed a change in swim pattern and an attraction to the serine side of the box.

B, responses to cysteine 10^{-2}M (high concentration) and 10^{-3}M (intermediate concentration). CF n°73 gives an example of an individual that did not respond to 10^{-3}M cysteine: no change in position, no change in pattern, no change in swim kinetics. CF n°71 and CF n°72 on the other hand did respond to cysteine 10^{-2}M, the former by an increase in activity, the later by a transient change in position. SF n°34 is a rare example of a SF that was attracted to cysteine.

A Pachón cavefish, lysine 10^{-2}M

B Surface fish, lysine 10^{-2}M

C F2 hybrids, lysine 10^{-2}M

D Pachón cavefish, histidine 10^{-2}M

E Surface fish, histidine 10^{-2}M

F F2 hybrids, histidine 10^{-2}M

G Pachón cavefish, leucine 10^{-2}M

H Surface fish, leucine 10^{-2}M

I F2 hybrids, leucine 10^{-2}M

Suppl Figure 4

Suppl Fig4 (with Figure 6). Examples of behavioral responses to lysine (ABC), histidine (DEF) and leucine (GHI) 10^{-2}M (high concentration)

In each panel, in the two left graphs, the blue color indicates the water/control injection side The lysine injection side is shown in red, the histidine injection side is yellow, the leucine injection side is pink.

A, The 2 CF individuals shown decrease their swimming activity upon injection. Moreover, n°81 and 87 shift their position to the odor side of the box. A change in swimming pattern from intensive wall following to thigmotactism with circles and some random swimming is also observed.

B, The 2 SF individuals shown do not change their swimming pattern (Random) or kinetics in response to lysine.

C, F2 n°35 increased swimming activity upon lysine injection but did not show attraction to odorant side. F2 n° 37 kept its random pattern of swimming and was not attracted to the odorant side.

D, Whether the cavefish individual was initially displaying thigmotactism (CF n°8) or swimming in large circles (CF n°4), the response to histidine included decreasing the swimming activity and restricting locomotion to the odorant side of the box.

E, The two surface fish individuals shown swam slowly and randomly as baseline pattern. Both increased their activity and spent more time in the odorant side.

F, F2 hybrids n°11 and n°1 were both attracted to histidine odorant side and they changed their swimming pattern.

G, CF n°65 showed no response, neither in activity, in position nor in pattern. CF n°4 drastically changed all swimming parameters.

H, Surface fish showed little reaction to leucine. SF n°43 and n°40 persisted in random swimming patterns, without change of position or swimming kinetics.

I, F2 hybrids n°16 and n°13 had markedly different baseline patterns (large circles *versus* random swim, respectively). The former changed its pattern to less active and small circles, suggesting that it perceived leucine injection. The later showed no behavioral response.

Supp Figure 5

Suppl Fig5 (with Figure 7). Individual olfactory scores of SF, CF and F2 in control and experimental conditions.

The graphs represent index values for the 4 response parameters used to calculate the total individual olfactory score for each fish. Each fish is depicted by a colored line linking its 4 different indexes and its final individual olfactory score. Experimental conditions and fish types are indicated. The threshold value for olfactory score was set at 1.5 (dotted line), because very few individuals of the 3 fish types have scores above this value in control experiments when water is perfused in the box on both sides. The percentages in black and grey indicate the proportion of responders/non-responders, respectively. The colored lines of responders are bright, those of non-responders are pale.

A

B

Suppl Figure 6

Suppl Fig6 (with Figure 8). Relationships between the number of round trip in X and the olfactory score

A, Regressions to explore the correlation between baseline number of round trips in X displayed by individual fish before odor injection and their individual olfactory score. Fish and amino acid type and concentrations are indicated. Linear correlations were calculated with Spearman's rank correlation test followed by Student test for the p values. Conditions for which significant correlation was found are labelled with an asterisk.

B, The PVE (Percentage of Variation Explained) on the individual score is plotted as a function of individual swimming parameter (numbers of round trips in X), to probe the predictability of the behavioral output as a function of "locomotor activity personality". Linear correlations were calculated with Spearman's rank correlation test followed by Student test for the p values. Conditions for which significant correlation was found are labelled with an asterisk. SF (blue), F2 (green) and CF (red).

Supp Figure 7

Suppl Fig7 (with all figures). Complete color code for the description of baseline swimming patterns.

A - controls

B- odors

Supp Figure 8

Suppl Fig8 (with all figures). Correspondence Analyses (CA) for assessment of behavioral pattern change after different stimuli.

Confident ellipses for each condition in the behavioral space are shown in blue for SF, green for F2 hybrids, red for CF. The conditions are indicated at the top of the graphs.

A, Control experiments. CA shows that behavioral patterns in the 3 types of fish are clearly distinctive in the behavioral space: confidence ellipses are strictly non-overlapping. Moreover, CA confirms that for the 3 types of fish, the swimming pattern does not vary along the one hour test when no injection is made (nothing/nothing), when water is injected on one side (water/nothing), or when water is injected on both sides ((water/water). In all cases, confidences ellipses overlap in the "before" and "after" condition. These analyses fully confirm the results shown in Figures 2B and 3B with the colored bar plot representation.

B, Odor stimulation experiments. Changes in behavioral patterns is the behavioral space are illustrated for the 6 different amino acid tested. Overlap of confidence ellipses indicate a lack of change in behavioral pattern after odor stimulus, whereas separated confidence ellipses indicate a change in behavioral pattern after odor stimulus. The CA fully confirms the results shown in Table1 and Figures 4-5-6 with the colored bar plot representation.