Title Page

Title:

Recognition of the Nucleotide-Dependent Feature Facilitates the Microtubule End-Binding of
MCAK

Authors and affiliations:

Wei Chen[1] (chenwill@mail.tsinghua.edu.cn)
Yin-Long Song[1] (y.song4@uu.nl)
Jian-Feng He[1] (jianfenghe2015@mail.tsinghua.edu.cn)
Xin Liang[1,*] (xinliang@tsinghua.edu.cn)

[1] IDG/McGovern Institute for Brain Research, School of Life Sciences, Tsinghua University, Bei-
jing 100084, China

Corresponding author:

Name: Xin Liang
Address: Room E118, Medical School, Tsinghua University, 100084 Beijing, China
Tel: +86-10-62772563
Email: xinliang@tsinghua.edu.cn
ORCID: 0000-0001-7915-8094

Key words

MCAK
Microtubule dynamics
End binding
Depolymerization
GTP cap

Recognition of the Nucleotide-Dependent Feature Facilitates the Microtubule End-Binding of MCAK

Wei Chen[1], Yin-Long Song[1], Jian-Feng He[1], Xin Liang[1,*]

[1] IDG/McGovern Institute for Brain Research, School of Life Sciences, Tsinghua University, Beijing 100084, China

* Correspondence: xinliang@tsinghua.edu.cn

Abstract

The growing plus-end is a key regulatory site for microtubule dynamics. MCAK (mitotic centromere-associated kinesin), a microtubule depolymerizing kinesin, is an end-binding regulator of catastrophe frequency. It is intriguing how MCAK specifically binds to dynamic microtubule ends. Here, we measure the end-binding kinetics of MCAK using single-molecule imaging and reveal the end-binding preference. MCAK binds to the entire GTP cap, including the EB cap and the distalmost cap. Further analysis shows that MCAK strongly binds to GTPγS microtubules, suggesting that it could recognize the nucleotide-dependent feature of microtubules. Moreover, the binding preference is independent on the nucleotide state of MCAK, and this feature facilitates the high-affinity end-binding of MCAK. Finally, we show that despite partially sharing the binding regions, MCAK and XMAP215 function in an additive manner, demonstrating a simple logic of how the end-binding regulators work in coordination. In all, our results provide novel insights into understanding how MCAK regulates the dynamics of microtubule ends.

(156 words)

Introduction

Microtubule is a major type of cytoskeleton in eukaryotic cells and supports many cellular processes, such as cell division, polarization and migration (Alberts, 2015). However, the properties of microtubules vary a lot in different cellular processes and structures (Janke and Magiera, 2020; Kapitein and Hoogenraad, 2015; Wittmann et al., 2001). The plasticity implies that cellular microtubules are under complex and accurate regulations. Because microtubule-associated proteins (MAPs) are important components in the regulatory networks of microtubules, their functions and working mechanisms are key to understand cellular regulations on microtubules.

Not only do kinesin superfamily members move cargoes along microtubules, they also act as regulators of microtubule dynamics, for example, the depolymerizing kinesins (e.g. the kinesin-8 and kinesin-13 family members). Kinesin-13 was initially identified in studying spindle functions and neurons (Aizawa et al., 1992; Noda et al., 1995; Walczak et al., 1996; Wordeman and Mitchison, 1995). Later, the functions of kinesin-13 were revealed in more cellular processes, including spindle assembly (Walczak et al., 2002), chromosome segregation (Kline-Smith et al., 2004), directional migration (Zong et al., 2021), ciliary length control (Piao et al., 2009; Vasudevan et al., 2015), and neuronal polarization (Ghosh-Roy et al., 2012; Puri et al., 2021). The loss of kinesin-13 often alters structural and dynamic properties of cellular microtubules, implying that the kinesin-13 class members play regulatory roles. In addition, the subcellular localizations of kinesin-13, such as growing microtubule ends, centrosome and centromere, are active regulatory sites of microtubule dynamics in cells, further suggesting the functions of the kinesin-13 class members in controlling the length, mass and polarity of microtubules in cells.

As a representative model for the kinesin-13 family, MCAK was shown to be a microtubule depolymerase and a catastrophe factor (Desai et al., 1999; Gardner et al., 2011; Hunter et al., 2003). Biochemical and structural analysis suggests that MCAK could deconstruct the protective end structure of dynamic microtubules, and it does so by bending and then breaking off the underlying substrate (i.e. terminal tubulin dimers or protofilaments) (Moores and Milligan, 2006). It was proposed that MCAK targets microtubule ends by rapid lattice diffusion or binding to the end-binding proteins (EBs) (Gouveia et al., 2010; Helenius et al., 2006; Honnappa et al., 2009). Furthermore, the direct binding of MCAK to growing microtubule ends was also observed (Gouveia et al., 2010), implying the presence of specific end-binding sites. Previous studies on the structures of MCAK, propose that MCAK targets to microtubule ends by recognizing the curved protofilaments (Asenjo et al., 2013; Benoit et al., 2018; Tan et al., 2008; Trofimova et al., 2018; Wang et al., 2017). This was mostly based on the findings on stabilized microtubules, short protofilaments or tubulin dimers. It still remains elusive how MCAK binds to dynamic microtubule ends.

In the present study, we characterize the direct binding of MCAK to growing microtubule ends using single-molecule imaging. The key finding is that MCAK binds

to the entire GTP cap, including both the EB cap, where GDP·Pi-tubulin dimers are
thought to gather, and the distalmost cap, where XMAP215 binds to curved protofil-
ament (mostly in GTP state). The high-affinity binding of MCAK to the EB cap was
intriguing. Further analysis shows that MCAK strongly binds to GTPγS microtubules,
thought to be the analogue of GDP·Pi-tubulins. Therefore, we think that MCAK could
recognize the nucleotide-dependent feature at dynamic microtubule ends. Moreover,
the mutant studies show that the binding-preference of MCAK for GDP·Pi-tubulins
facilitates the end-binding and in turn the function of MCAK. Finally, we find that
despite having partially overlapped binding regions, MCAK and XMAP215 could
precisely specify their regulatory function at dynamic microtubule end, thereby
providing novel insights in understanding how microtubule-associated proteins func-
tionally orchestrate at growing microtubule ends.

Results

MCAK shows a binding preference for growing microtubule ends

We set off by measuring the single-molecule binding kinetics of MCAK at growing microtubule ends using the *in vitro* microtubule dynamics assay. This experiment was performed in BRB80 supplemented with 50 mM KCl and 1 mM ATP, providing a nearly physiological ion strength. In this condition, the individual binding events of MCAK can be observed at both the end and the lattice (**Fig. 1A**). By measuring the fluorescence intensity of these binding events, we confirmed that they reflected individual MCAK dimers (**Fig. s1**). Using an in-house software (Song et al., 2020), we determined the binding position of individual MCAK molecules at subpixel resolution by fitting a Gaussian function to their intensity profile along the long axis of microtubules (**Fig. 1B**). Using the position of the plus-end as the spatial reference, we overlaid the end-binding of 152 MCAK molecules (**Fig. 1C**). The distribution showed that the binding regions of MCAK had a length of 162 nm (FWHM, the Full Width at Half Maximum), about 20 layers of tubulin dimers. We defined the binding events of MCAK within this region as "end binding" and those on the lattice as "lattice binding" (**Fig. 1A**).

To understand the binding kinetics, we first compared the apparent association rates of MCAK on the plus-end (k_{on-P}) and lattice (k_{on-L}) in the presence of ATP. The ratio ($R_{E/L}$) of k_{on-P} ((22. 6 $\pm$ 1.4) $\times 10^{-6}$ s^{-1} nM^{-1}, per tubulin dimer, hereinafter) and k_{on-L} ((2.8 $\pm$ 0.3) $\times 10^{-6}$ s^{-1} nM^{-1}) was 13 $\pm$ 1 (n=66 microtubules from 3 assays) (**Fig. 1 D-E**). We then measured the dwell time, which reflects the dissociation rate (k_{off}) (**Fig. s1**). The mean dwell time of MCAK on the lattice was shorter than that at the plus-end (plus end: 0.75 $\pm$ 0.02 s, n=966 events from 3 assays; 0.62 $\pm$ 0.02 s, n=702 events from 3 assays, p<0.001) (**Fig. 1F**). Using these kinetic parameters, we calculated the dissociation constant (K_d) of MCAK for the plus-end (K_{d-P} =69 μM) and lattice (K_{d-L}=1057 μM), respectively. Therefore, MCAK shows a clear end-binding preference. We also observed single-molecule binding events of MCAK to the minus-end (**Fig. 1A**). Given the higher physiological relevance, we focused on the plus-end binding events (referred to as "end-binding" hereinafter) in the following analysis.

To understand the influence of the nucleotide state, we measured the end-binding kinetics of MCAK·AMPPNP, MCAK·ADP or MCAK·APO (the nucleotide-free state). First, the k_{on-P} and k_{on-L} of MCAK·AMPPNP were both reduced in comparison to that of MCAK·ATP (**Fig. 1 D-E**)*,* resulting in an end-binding preference ($R_{E/L}$=11 $\pm$ 2, n=29 microtubules from 2 assays, p>1) that is similar to that of MCAK·ATP. Meanwhile, the end-binding of MCAK·AMPPNP showed a longer dwell time (3.32 $\pm$ 0.24 s, n=231 events from 2 assays, p<0.001), indicating a lower k_{off} (**Fig. 1F and Fig. s1**). As a result of the proportionally reduced k_{on-P} and k_{off}, the end-binding affinity (K_{d-P}=61 μM) of MCAK·AMPPNP was similar to that of MCAK·ATP. Second, MCAK·ADP showed a nearly unchanged k_{on-P} and an increased k_{on-L}, leading to a smaller $R_{E/L}$ (2 $\pm$ 0.3, n=21 microtubules from 2 assays, p<0.001) and suggest a re-

duced, but not the absence of, end-binding preference (**Fig. 1 D-E**). Meanwhile, it
showed a significantly shorter end-binding dwell time (0.46 ± 0.02 s, n=327 events
from 2 assays, p<0.001) than MCAK·ATP (**Fig. 1 F**). Therefore, MCAK·ADP has a
lower end-binding affinity ($K_{\text{d-P}}$=163 μM). Third, APO·MCAK showed an increase in
$k_{\text{on-P}}$ and an even greater increase in $k_{\text{on-L}}$, thereby having a smaller $R_{\text{E/L}}$ (2 ± 0.2, n=36
microtubules from 3 assays, p<0.001) (**Fig. 1 D-E**). It also showed a significantly
longer end-binding dwell time (1.89 ± 0.23 s, n=71 events from 3 assays, p<0.001)
(**Fig. 1 F**), suggesting a lower k_{off} (**Fig. s1**). Therefore, in comparison to MCAK·ATP,
MCAK·APO possesses a higher end-binding affinity ($K_{\text{d-P}}$=1.8 μM) but partially loses
the end-binding preference. Based on these measurements, we conclude that the
end-binding preference and affinity of MCAK depend on its nucleotide state (see
Discussion).

Having measured the end-binding kinetics, we were able to calculate whether the
binding of MCAK on dynamic microtubule ends is due to direct end-binding (3D dif-
fusion) or 1D lattice-diffusion (Gouveia et al., 2010; Helenius et al., 2006). Note that
these two mechanisms are not mutually exclusive. Using our dataset, we measured
that the lattice-diffusion coefficient of MCAK (D) on the lattice of dynamic microtu-
bules was 0.023 ± 0.001 $\mu m^2 \cdot s^{-1}$ (**Fig. 1G**), close to the previously reported value
(Cooper et al., 2010). Because the dwell time of the lattice-binding events was ~0.6 s,
the average length scanned by a MCAK molecule via 1D lattice-diffusion was ~160
nm. Note that this is close to the length the GTP cap. Using a previously reported
Fick's first equation of diffusion (Helenius et al., 2006), we calculated the flux of
MCAK reaching growing ends via 1D diffusion was 0.004 s^{-1}. This was significantly
lower than the arrival rate of MCAK onto the plus end observed in our experiments
(N_p= $k_{\text{on}} \cdot C$ =0.067 s^{-1}. C: MCAK concentration). Therefore, in our experimental con-
dition, 1D lattice-diffusion makes a minor contribution to the end-binding of MCAK
(~ 6%), and the end-binding of MCAK can be thought of as the direct interaction be-
tween diffusive MCAK and the end structure.

MCAK binds to the entire GTP cap

Having recorded and analyzed the end-binding events of MCAK, we wondered
how MCAK could recognize the end of dynamic microtubules. Previous studies,
mostly based on stabilized microtubules, proposed the idea that MCAK binds to mi-
crotubule ends by recognizing curved protofilaments (Asenjo et al., 2013; Benoit et al.,
2018; Trofimova et al., 2018). This model predicts that the binding site of MCAK is
located to the distalmost tip of growing microtubule ends, where the XMAP215 fam-
ily members are thought to bind (i.e. the distalmost cap) (Ayaz et al., 2012; Brouhard
et al., 2008). We tested this prediction by measuring the end-binding regions of
MCAK, EB1 and XMAP215 (**Fig. 2A**). To our surprise, the binding region of MCAK
was longer than that of XMAP215. It extended proximally towards the lattice, and
covered the binding region of EB1 (i.e. the EB cap) (Maurer et al., 2012). To further
understand if MCAK binds to the distalmost cap and the EB cap with different kinet-
ics, we compared the dwell time of the distally located MCAK to that of the proxi-

mally distributed ones (**Fig. 2B**). Note that we defined the MCAK molecules localiz-
ing to the left of the FWHM of XMAP215 binding region as the proximally distribut-
ed, while those localizing to the right of the FWHM of EB1 binding region was dis-
tally distributed. No significant difference was found (τ_{distal}=0.8 ± 0.5, n=28 binding
events; $\tau_{proximal}$=0.7±0.6, n=45 binding events; p>0.3, the two-tailed Mann-Whitney U
test with Bonferroni correction). Note that in all experiments, the morphologies of
growing microtubule ends were comparable (**Fig. 2C**). Based on these results, we
conclude that MCAK recognizes the entire GTP-cap. This result is intriguing as it
suggests that in addition to the curved protofilaments, there are additional end-binding
sites for MCAK. Moreover, this also provides an independent measurement for the
size of the GTP cap at growing microtubule ends, which was previously determined
using the tubulin dilution experiments or by measuring the binding region of EB1
(Duellberg et al., 2016).

MCAK strongly binds to GTPγS microtubules in a nucleotide-independent
manner

How could MCAK bind to the EB cap? The tubulin dimers in the EB cap are
thought to be different from those in the distalmost cap (Maurer et al., 2011). For
example, the distalmost tubulin dimers (mostly GTP-tubulin) often lack lateral con-
strains (Vitre et al., 2008) and adopt a bent shape (Hoeoeg et al., 2011; McIntosh et al.,
2018), while those in the EB cap (mostly GDP·Pi-tubulin) have formed at least some
lateral contacts and had a more straight conformation (Guesdon et al., 2016; Maurer et
al., 2012). To address this issue, we first examined that to what extent, MCAK may
adopt a similar end-binding mechanism as EB1.

We tested the contribution of GDP·Pi-tubulins, which underlies the end-binding
preference of EBs (Maurer et al., 2011; Maurer et al., 2012). Here, we compared the
binding affinity of MCAK·AMPPNP on GTPγS, GMPCPP and GDP microtubules
(**Fig. 3A**), which mimics tubulin dimers in different GTP hydrolysis states. Because
MCAK in the concentration required for a full-lattice decoration would rapidly disas-
semble microtubules, we cannot directly compare the total binding of MCAK on mi-
crotubules as what was previously done for the EBs (Maurer et al., 2011; Maurer et al.,
2012). Therefore, we took an alternative approach by recording the single-molecule
binding of MCAK·AMPPNP (1 nM) on microtubules and measuring fluorescence in-
tensity summation over a period of time (1000 frames, 0.3 s per frame) (**Fig. s2**).
Strikingly, we found that MCAK·AMPPNP bound strongly to GTPγS microtubules
(**Fig. 3 A-C**). Quantitative analysis revealed a significantly stronger binding on
GTPγS microtubules than on GMPCPP or GDP microtubules (**Fig. 3 B-C**). Because
the tubulin dimers in GTPγS microtubules were found to be the analogue of
GDP·Pi-tubulins (Maurer et al., 2011), this result suggests that MCAK·ATP has a
binding preference for GDP·Pi-tubulins.

Because the end-binding affinity and preference of MCAK depends on the nu-
cleotide state, we wondered if the preference for GTPγS microtubules is similar.

Therefore, we performed the same experiments using MCAK·AMPPNP, MCAK·ATP,
MCAK·ADP and MCAK·APO. Although the MCAK variants bound to microtubules
with different affinities (MCAK·APO > MCAK·AMPPNP > MCAK·ADP >
MCAK·ATP) (**Fig. 3B**), they all showed a clear binding preference for GTPγS micro-
tubules over GDP or GMPCPP microtubules (**Fig. 3 B-C**). These observations suggest
that the binding preference to GTPγS microtubules (i.e. GDP·Pi-tubulin) is, to a large
extent, independent on the nucleotide state of MCAK.

MCAK binds to the end region of stabilized GMPCPP microtubules in a nucleo-
tide state-dependent manner

We noted that MCAK showed a nucleotide state-dependence in the binding pref-
erence for dynamic microtubule ends, but the binding to GTPγS microtubules is nu-
cleotide independent. Therefore, we wondered if the binding to curved protofilaments
at the distalmost cap of dynamic microtubules could account for the nucleotide-state
dependence. To address this issue, we measured the binding preference of MCAK at
the end of stabilized GMPCPP microtubules (**Fig. 4A**), where the length and curvature
of protofilaments are thought to be similar to those at the end of dynamic microtu-
bules but has no GDP·Pi-tubulins (Manka and Moores, 2018; McIntosh et al., 2018).
We found that the MCAK in ATP or AMPCPP state showed a binding preference for
the end of GMPCPP microtubules, while MCAK·ADP and MCAK·APO did not (**Fig.**
4B-C). This is similar to the observation that MCAK·ADP and MCAK·APO had a
lower binding preference for dynamic microtubule ends, thereby providing an expla-
nation for the nucleotide state-dependence of the end-binding of MCAK.

The binding preference for GDP·Pi-tubulins facilitates the end-binding of
MCAK

Intuitively, the binding preference for the EB cap, in addition to the previously
established preference for the curved protofilaments, would provide more
high-affinity sites for MCAK, thereby facilitating the end-binding of MCAK. To test-
ed this idea, we studied two mutants, MCAK[K524A] (K524A in the α4 helix) and
MCAK[V298S] (V298S in the L2 loop). The α4 helix of MCAK interacts with the in-
tra-dimeric interface of tubulin heterodimer (**Fig. 5A**), and the K524A mutation is
expected to disrupt the electrostatic interaction between MCAK and tubulin hetero-
dimer (Patel et al., 2016; Wang et al., 2017). The L2 loop of MCAK has a direct con-
tact with the inter-dimeric interface of tubulin heterodimer (**Fig. 5A**), and the V298S
mutation is expected to disrupt the hydrophobic interaction between the L2 loop and
α-tubulin (Wang et al., 2015). We confirmed that both mutants were depolymeriza-
tion-deficient (**Fig. s3**), as previously reported (Wang et al., 2017; Wang et al., 2015).

Compared to MCAK, MCAK[K524A] showed a significantly reduced binding inten-
sity on microtubules. Meanwhile, it still showed a binding preference for GTPγS mi-
crotubules (**Fig. 5B-C**), but the ratios between the binding intensity on GTPγS micro-
tubules and that on GMPCPP or GDP microtubules were both reduced (**Fig. 5D**). This
suggests that MCAK[K524A] had a lower preference for GTPγS microtubules. On the

contrary, we found that MCAK[K524A] showed a higher binding to the end than to the
lattice of GMPCPP microtubules (**Fig. 5 E-F**), suggesting that the binding preference
for curved protofilament is maintained. Therefore, MCAK[K524A] could be a candidate
variant to test how the preference for GDP·Pi-tubulin contributes to the end-binding
preference of MCAK. We found that MCAK[K524A] could bind to dynamic microtubule
ends, but its k_{on-P} and $R_{E/L}$ were both reduced (**Fig. 5 G-H**). Meanwhile, the
end-binding dwell time of MCAK[K524A] was also reduced (**Fig. 5I**), suggesting a high-
er off-rate. These results are consistent with the idea that the higher binding affinity to
GDP·Pi-tubulin directly contributes to the end-binding preference of MCAK at grow-
ing microtubule ends.

By comparison, the binding preferences of MCAK[V298S] for GTPγS microtubules
and the end of GMPCPP microtubules were both reduced (**Fig. 5 B-F**), implying the
reduced preferences for both the EB cap and the distalmost cap. In the dynamics assay,
the k_{on-P}, $R_{E/L}$ and dwell time of MCAK[V298S] were decreased in comparison to those of
MCAK (**Fig. 5 H-I**). We noted that the k_{on-P}, $R_{E/L}$ and dwell time of MCAK[V298S] were
similar to those of MCAK[K524A] (two-tailed Mann-Whitney U test with Bonferroni
correction, p>0.05), suggesting that the binding to the EB cap makes a major im-
portance for the end-binding preference of MCAK at dynamic microtubule ends.

Functional specification of MCAK and XMAP215

Having revealed the novel binding sites of MCAK in dynamic microtubule ends,
we wondered how MCAK may work in coordination with other microtubule plus
tip-binding proteins, for example EB1 and XMAP215. It has been shown that EB1
interacts with and recruits MCAK to growing microtubule ends (Gouveia et al., 2010;
Honnappa et al., 2009; Lee et al., 2008), so it is straightforward that EB1 would facil-
itate the function of MCAK as a catastrophe factor. On the contrary, MCAK and
XMAP215 were shown to antagonistically regulate the assembly of microtubules, and
it was thought that these two molecules may compete directly for the growing ends
(Tournebize et al., 2000). Here, we showed that the binding regions of MCAK and
XMAP215 in growing microtubule ends are partially overlapping. In addition, previ-
ously resolved structural models suggest that MCAK and XMAP215 likely bind to
similar regions on tubulin dimers (**Fig. s4**). Therefore, these two molecules could in-
deed compete for the binding sites in the distalmost cap. However, it is not yet clear to
what extent, the competition from XMAP215 would antagonize the function of
MCAK, especially given the recent finding that XMAP215 itself could promote the
catastrophe of microtubules (Farmer et al., 2021). Therefore, it is intriguing what
would be the collective effect of MCAK and XMAP215.

To address this issue, we studied how they together regulate microtubule grow
rate and catastrophe frequency. Based on our pilot experiments, we chose 20 nM for
MCAK and 50 nM for XMAP215 as the representative concentrations. In the pres-
ence of 20 nM MCAK, the dynamic microtubules could grow but showed a signifi-
cantly increased catastrophe frequency and shorter lifetime (**Fig. 6 A-B**). Meanwhile,

50 nM XMAP215 increased growth rate by more than four times (control: 0.8 ± 0.1
μm min^{-1}, n=97 microtubules from 6 assays; 50 nM XMAP215: 3.7 ± 0.4 μm min^{-1},
n=50 microtubules from 3 assays). We noted that XMAP215 alone caused a mild
shortening in the lifetime of dynamic microtubules (**Fig. 6 A-B**). This is consistent
with the previous report and suggesting the enhanced fluctuations in the end structures
(Farmer et al., 2021). We then added both 20 nM MCAK and 50 nM XMAP215. In
this case, microtubule grew in a similar rate as if there was no MCAK (3.9 ± 0.4 μm
min^{-1}, n=52 microtubules from 3 assays, p=0.5, the two-tailed Mann-Whitney U test
with Bonferroni correction), suggesting that MCAK has no major effect on the func-
tion of XMAP215 as a polymerase. Meanwhile, the lifetime of dynamic microtubules
became shorter than that with only MCAK or XMAP215 (**Fig. 6B**). To understand if
this reflects the simple summation of the separate contributions of MCAK and
XMAP215 or their synergistic effect, we calculated the probability-based catastrophe
frequency in all conditions (**Fig. 6C**) (Gardner et al., 2011).We found that the effects
of MCAK and XMAP215 on the probability of microtubule catastrophe were only
additive, showing that there was no synergic effect in regulating catastrophe frequen-
cy (**Fig. 6C**).

Discussion

This study provides mechanistic insights into understanding the end-binding mechanism of MCAK. The previously established view on the end-binding mechanism of MCAK is based on specific recognition of curved protofilaments at the distalmost tip of growing microtubules (Asenjo et al., 2013; Benoit et al., 2018; Tan et al., 2008). The findings in the present study add to the model by showing that MCAK strongly binds to the EB cap, and this binding preference could facilitate the end-binding and in turn the depolymerizing activity of MCAK. We now discuss our main findings and their implications.

MCAK preferentially binds to the GDP·Pi-tubulin in a nucleotide state-independent manner

Our main finding is that MCAK preferentially binds to the EB cap where the GDP·Pi-tubulins gather, in addition to the previously known binding preference for the curved protofilaments (Asenjo et al., 2013; Benoit et al., 2018; Tan et al., 2008). This conclusion is based on several lines of evidence. First, MCAK·ADP or MCAK·APO shows a significant reduction but not a complete loss in the binding preference for dynamic microtubule ends. However, their binding preference for stabilized microtubule ends is completely absent. The different binding behavior to dynamic and stabilized microtubule ends suggests that other end features, in addition to the curved protofilament, also contribute to the end-binding of MCAK. Second, the localization analysis of MCAK shows that MCAK binds to the entire GTP cap, including the EB cap and the distalmost cap. Third, MCAK shows a binding preference for GTPγS microtubules over GDP or GMPCPP microtubules. Because the tubulin in GTPγS microtubules is considered to be the analogues to the GDP·Pi-tubulins (Maurer et al., 2011; Maurer et al., 2012), MCAK is expected to have a high affinity to the EB cap. Fourth, the loss of the binding preference of MCAK[K524A] for GTPγS microtubules, but not for the ends of GMPCPP microtubules, suggests that the binding preference for GDP·Pi-tubulins is required for the optimal end-binding affinity and preference.

Implications for the working mechanism of MCAK

We think that our findings add to the current working model of MCAK in four aspects. First, the nucleotide state-independent binding preference for GDP·Pi-tubulins provides a mechanism for MCAK staying bound to the GTP cap after ATP hydrolysis. For example, in a speculative scenario (**Fig. 7A**), when one of the motor domains of MCAK is converted to the ADP or APO state after the breakage of the terminal tubulin dimers, it loses the binding affinity to the curved protofilaments (**Fig. 7A**). However, it still preferentially binds to GDP·Pi-tubulins, so it has a larger chance binding back to the EB cap than to the lattice and thereby remains within the GTP cap (**Fig. 7A**). Then, while the ADP is being replaced by a new ATP, this motor domain could diffuse around to explore the working site for the next round of catalysis. In this way, the nucleotide state-independent binding feature of MCAK would fa-

cilitate the binding to the dynamic microtubule ends. This idea is consistent with the
findings that MCAK[K524A], a mutant showing reduced preference for GDP·Pi-tubulins,
had disrupted end-binding affinity, preference and dwell time.

Second, the previous studies showed that MCAK has a direct interaction with
EB1, and EB1 increases the end-binding rate of MCAK (Gouveia et al., 2010;
Honnappa et al., 2009; Lee et al., 2008). Intuitively, there is a gap here that after the
arrival of MCAK at the binding region of EB1, how it could reach the high-affinity
binding site or working site, previously known as the curved protofilament at the dis-
talmost tip (Asenjo et al., 2013; Benoit et al., 2018; Tan et al., 2008). Our finding fills
this gap by showing that MCAK also binds strongly to the GDP·Pi-tubulins where the
EBs bind (**Fig. 7B**). Moreover, MCAK could explore the entire length of GTP cap
(~160 nm) by 1D lattice diffusion within the binding dwell time. In this way, we argue
that MCAK could find the precisely catalyzing site at dynamic microtubule ends by a
combination of 3D diffusion (to first reach the GTP cap) and 1D lattice diffusion
(short range, within the GTP cap).

Third, a previous study showed that the association on-rate is an important regu-
lated parameter for the functions of MCAK (Cooper et al., 2010). The binding affinity
and preference for the EB cap, in addition to the distalmost cap, would intuitively
provide more binding sites, thereby further increasing the binding on-rate of MCAK.
This idea is also consistent with the finding that MCAK[K524A], a mutant with a reduced
preference for GDP·Pi-tubulins, had a reduced end-binding on-rate.

Fourth, despite having partially overlapped binding regions, MCAK and
XMAP215 provide independent controls for catastrophe frequency and growth rate of
microtubules, respectively (**Fig. 7B**). The free combination of these two regulators
suggests a straightforward strategy in the regulations on the length and mass of mi-
crotubules, i.e., simply by controlling their amount (i.e. expression level) or activity
(e.g. via phosphorylation). In this sense, the working scenario of MCAK and
XMAP215 provides a molecular paradigm to understand how the regulators of mi-
crotubules orchestrate their behaviors at the nanoscopic end structure of dynamic mi-
crotubules and reflects a clear logic in the regulatory network of microtubule dynam-
ics.

Materials and Methods

Protein expression and purification

MCAK purification.

GFP-MCAK was expressed using a modified BAC-to-BAC protocol. The purification was performed as previously described (Helenius et al., 2006). Briefly, the cells were lysed by dounce homogenization in the lysis buffer. The lysate was centrifuged at 40,000 rpm for 60 min at 4°C (XPN-100 ultracentrifuge, Type 45 Ti rotor, Beckman, USA). The supernatant was filtered through a 0.45 μm filter (Pall, 66229) and then purified using a cation-exchange column (HiTrap SP-HP, Cat.17115101, GE, USA). The protein was eluted using the elution buffer supplemented with a continuous salt gradient (0.15~1.0 M NaCl). The peak fractions were pooled together and then loaded onto a Ni-sepharose column (QIAGEN, 30210). The column was then washed with the washing buffer I. After the washing step, 3C Protease (final concentration 10 μg/ml) (Thermo Fisher Scientific, 88946) was used to perform on-column his-tag cleavage overnight at 4°C. The eluate was further purified using a gel-filtration column (Superdex 200 increase 10/300 GL column, Cat. 28990944, GE, USA) and eluted using the washing buffer II. The purified MCAK was frozen in liquid N_2 and stored at -80°C. Lysis buffer: 50 mM HEPES pH 7.5, 150 mM NaCl, 5% glycerol, 0.1% Tween 20, 1.5 mM $MgCl_2$, 3 mM EGTA, 1 mM DTT, 0.5 mM Mg-ATP, 10 units/ml Benzonase. Elution buffer: 6.7 mM HEPES pH 7.5, 6.7 mM MES, 6.7 mM sodium acetate, 1.5 mM $MgCl_2$, 10 μM Mg-ATP. Washing buffer I: 50 mM $NaPO_4$ buffer pH 7.5, 300 mM NaCl, 10 mM imidazole, 10% glycerol, 1 mM $MgCl_2$, 10 μM Mg-ATP. Washing buffer II: BRB80, 300 mM KCl, 1 mM DTT and 10 μM Mg-ATP. BRB80: 80 mM PIPES/KOH pH 6.9, 1 mM $MgCl_2$, 1 mM EGTA.

EB1 purification.

EB1-GFP was expressed and purified as previously described (Song et al., 2020). Briefly, EB1-GFP was expressed in the *BL21 E. coli* strain and lysed using sonication in the lysis buffer. The lysate was centrifuged at 40,000 rpm for 60 min at 4°C. The supernatant was filtered through a 0.45 μm filter and loaded onto a Ni-sepharose column. The column was then washed using the washing buffer. After the washing step, 3C Protease (final concentration 10 μg/ml) was used to perform on-column his-tag cleavage overnight at 4°C. The eluate was desalted into the storage buffer using the PD-10 desalting column (GE, 17085101). The final purified protein was frozen in liquid N_2 and stored at -80 °C. Lysis buffer: 50 mM $NaPO_4$ buffer pH 7.5, 300 mM NaCl, 10% glycerol, 10 mM imidazole, 1 mM DTT, 0.1% Tween 20. Washing buffer: 50 mM $NaPO_4$ buffer pH 7.5, 300 mM NaCl, 10% glycerol, 30 mM imidazole, 1 mM DTT. Storage buffer: BRB80, 100 mM KCl, 10% glycerol and 1 mM DTT.

XMAP215 purification.

XMAP215-GFP was expressed using a modified BAC-to-BAC protocol. The purification was performed as previously described (Brouhard et al., 2008). Briefly, the

cells were lysed by dounce homogenization in the lysis buffer. The lysate was centri-
fuged at 40,000 rpm for 60 min at 4°C. The supernatant was filtered through a 0.45
µm filter and then purified using a cation-exchange column. The protein was eluted
using the elution buffer supplemented with a continuous salt gradient (0.15~1.0 M
NaCl). The peak fractions were pooled together and then loaded onto a Ni-sepharose
column. The column was then washed with the washing buffer I. The protein was
eluted using the washing buffer I supplemented with a continuous imidazole gradient
(20-300 mM). The fractions with best purity were pooled together and further purified
using a gel-filtration column (Superdex 200 increase 10/300 GL column, Cat.
28990944, GE, USA) and eluted using the washing buffer II. The final purified
XMAP215 was frozen in liquid N_2 and stored at -80°C. Lysis buffer: 50 mM HEPES
pH 7.5, 50 mM NaCl, 5% glycerol, 0.1% Tween 20, 1 mM DTT, 10 units/ml Benzo-
nase. Elution buffer: 6.7 mM HEPES pH 7.5, 6.7 mM MES, 6.7 mM sodium acetate.
Washing buffer I: 50 mM NaPO4 buffer pH 7.5, 300 mM NaCl, 10 mM imidazole, 10%
glycerol, 1 mM $MgCl_2$, 10 µM Mg-ATP. Washing buffer II: BRB80, 150 mM KCl, 1
mM DTT.

Tubulin preparation.

Tubulin was purified from porcine brain tissues by two cycles of polymerization
and depolymerization followed by affinity purification using the TOG-based column,
as previously described (Gell et al., 2010; Widlund et al., 2012). Tubulin was labeled
with biotin (Thermo Fisher Scientific, 20217), TAMRA (Thermo Fisher Scientific,
C1171) and Alexa Fluor 647 (Thermo Fisher Scientific, A20106) using the NHS es-
ters according to the standard protocols (Gell et al., 2010).

Microtubule depolymerization assay

Microtubules (5% Alexa Fluor 647 labeled and 20% biotin labeled) were pol-
ymerized in the presence of GMPCPP (JenaBioscience, NU-405L) as previously de-
scribed (Song et al., 2020). The GMPCPP-stabilized microtubules were immobilized
on the surface of a cover glass using the biotin-NeutrAvidin protein links (Thermo
Fisher Scientific, 31000). GFP-MCAK was then added into the flow cell in the imag-
ing buffer. The sample was kept at 35°C using a temperature controller (Tokai Hit,
Japan). Images were recorded using a total internal reflection microscope (TIRFM)
(Olympus, Japan) equipped with a 100× 1.49 N.A. oil TIRF objective (Olympus, Ja-
pan) and an Andor 897 Ultra EMCCD camera (Andor, Belfast, UK). Images were
recorded every 5 s with a 100 ms exposure. Imaging buffer: BRB80 supplemented
with 1 mM ATP, 50 mM KCl, 80 mM D-glucose, 0.4 mg/ml glucose oxidase, 0.2
mg/ml catalase, 0.8 mg/ml casein, 1% β-mercaptoethanol, 0.001% Tween 20.

Microtubule dynamics assay

Microtubule dynamics assay was performed as previously described (Song et al.,
2020). Briefly, the immobilized GMPCPP-stabilized microtubules were used as the
template for microtubule growth. Tubulin dimers (13% TAMRA labeled) and the

protein of interest were then added into the flow cell in the imaging buffer. The sam-
ple was kept at 35°C using a temperature controller (Tokai Hit, Japan). Images were
recorded using a TIRFM (Olympus, Japan). To record microtubule dynamics, images
were recorded every 5 s with a 100 ms exposure. To record single molecule binding
events, images were recorded every 100 ms with a 50 ms exposure. Imaging buffer:
BRB80 supplemented with 1 mM ATP, 2 mM GTP, 50 mM KCl, 0.15% sodium car-
boxymethylcellulose, 80 mM D-glucose, 0.4 mg/ml glucose oxidase, 0.2 mg/ml cata-
lase, 0.8 mg/ml casein, 1% β-mercaptoethanol, 0.001% Tween 20.

The diffusion coefficient of MCAK on dynamic microtubule lattices

The diffusion coefficient of MCAK on dynamic microtubule lattices was calcu-
lated as previously described (Helenius et al., 2006). Firstly, we gained the position
coordinates of MCAK at its molecular trajectory on dynamic microtubule lattices us-
ing the plugin of imageJ (Fiji), TrackMate (Tinevez et al., 2017). Note that the mini-
mum track length was kept at 400 ms (100 ms/frame). Then, the mean-squared dis-
placement (MSD) of MCAK at every molecular trajectory (203 trajectories from 3
assays) was calculated against the interval (100 ms/frame) by an in-house script of
MATLAB. Lastly, the MSDs of MCAK was plotted against the time interval, and the
diffusion coefficient, D was obtained by a linear regression analysis ($<x^2>=2Dt$) in
Origin 8.0 (OriginLab Corporation, USA).

The flux of MCAK reaching growing microtubule ends

The calculation of the flux of MCAK reaching growing microtubule ends by
one-dimensional lattice-diffusion was performed according to Fick's first equation of
diffusion (Helenius et al., 2006):

$$J = -D \cdot \partial c / \partial x \ \ (Eq.1)$$

where J is the flux of MCAK reaching growing microtubule ends, $c(x,t)$ is the con-
centration of MCAK on microtubule lattice at position x from the growing microtu-
bule end and time t.

$$\text{When } x = 0 \quad J_0 = -D \cdot c_\infty / x_0 \ \ (Eq. 2)$$

Where

$x_0 = (D/k_{off})^{1/2}$

$$c(x \to \infty) = c_\infty = k_{on} \cdot C / k_{off}$$

Where C is the concentration of MCAK. Therefore, we can calculate the flux by tak-
ing the measured parameters (D, k_{on}, C and k_{off}) into the equation.

Localization analysis at growing microtubule ends

The analysis on the single-molecule localization at growing microtubule ends
was performed as previously described (Song et al., 2020). Briefly, we determined the
position of growing microtubule end by considering the microtubule lattice as a
Gaussian wall and its end as a half-Gaussian. This model was used to fit the microtu-
bule end and determined the position of microtubule end in every frame. The peak
position of the half-Gaussian was used as the origin of the positional axis, which was
used as a reference to measure the localization of individual single-molecule binding
events. We estimated that the accuracy of microtubule end tracking was ~6 nm by
measuring standard error of the distribution of the estimated error in microtubule end
position (Demchouk et al., 2011). To measure the precise localization of individual
binding events at growing microtubule ends, a Gaussian function was used to fit the
intensity profile of individual molecules. We estimated that the accuracy of the meas-
ured position was ~2 nm by measuring standard error of the fitting peak location
(Maurer et al., 2014). The peak location was recorded as the position of each molecule
in every frame. The average peak location over the entire binding period was recorded
as the position of the molecule. The positional distributions of GFP-MCAK,
EB1-GFP, XMAP215-GFP at growing microtubule ends were plotted along the lon-
gitudinal axis of microtubule. Finally, the plots of GFP-MCAK and XMAP215-GFP
were fitted to a Gaussian function and the plots of EB1-GFP was fitted to an exponen-
tially modified Gaussian function, from which we calculated the FWHM as an esti-
mation for the length of the binding region of each molecule.

Microtubule polymerization

GMPCPP, GDP and GTPγS microtubules were polymerized as previously de-
scribed (Manka and Moores, 2018). Briefly, GMCPP-microtubules were polymerized
using a mixture of 10 μM tubulin, 1 mM GMPCPP (JenaBioscience, NU-405L) and 4
mM $MgCl_2$ in BRB80, which was incubated for 2 hr at 37˚C. The polymerized mi-
crotubules were then collected using an Air-Driven Ultracentrifuge (Beckman,
340401), which were then resuspended in BRB80 and stored at 37˚C.
GDP-microtubules were polymerized using a mixture of 40 μM tubulin, 1 mM GTP
(Roche, 10106399001), 4 mM $MgCl_2$ and 4% DMSO (Sigma, 276855). The mixture
was incubated for 30 min at 37˚C. The polymerized microtubules were collected using
an Air-Driven Ultracentrifuge, resuspended in BRB80 with 20 μM taxol (Cell Sig-
naling Technology, 9807) and stored at 37˚C. GTPγS microtubules were obtained us-
ing two rounds of polymerizations. In the first round, a mixture of 40 μM tubulin, 2
mM GTPγS (Roche, 10220647001) and 4 mM $MgCl_2$ in BRB80 was incubated for 5
hours at 37°C. In the second round, 5 μl of the product from the first polymerization
was mixed with additional 40 μM tubulin, 2 mM GTPγS and 4 mM $MgCl_2$ to make a
final 25 μl reaction mixture. This mixture was incubated overnight at 37°C to obtain
long GTPγS microtubules. The polymerized microtubules were collected using an
Air-Driven Ultracentrifuge, resuspended in BRB80 with 20 μM taxol and stored at
37˚C.

MCAK binding to different microtubules

Different nucleotide microtubules were first immobilized in flow cells.
GFP-MCAK or its mutants were then added into the flow cell in the imaging buffer.
Images were recorded every 300 ms with a 100 ms exposure. Imaging buffer: BRB80
supplemented with 1 mM ATP, ADP or AMPCPP, 20 µM taxol, 50 mM KCl, 80 mM
D-glucose, 0.4 mg/ml glucose oxidase, 0.2 mg/ml catalase, 0.8 mg/ml casein, 1%
β-mercaptoethanol, 0.001% Tween 20.

556   The binding intensity of GFP-MCAK on GMPCPP microtubule ends

To measure the binding intensity of GFP-MCAK on GMPCPP microtubule ends,
an intensity profile of GFP-MCAK or the GMPCPP microtubule was plotted along the
microtubule. The average binding intensity of GFP-MCAK of two pixels represents
the end-binding intensity of GFP-MCAK at the corresponding position of the end, to
the lattice from the midpoint of the microtubule intensity attenuation at the microtu-
bule end.

563   Gamma fitting of microtubule lifetimes and catastrophe frequency analysis

564   The lifetime data of dynamic microtubules were fitted to the gamma density
565   function using the dfittool toolbox in the MATLAB (Mathworks, USA). The probabil-
566   ity-based catastrophe frequency is calculated as the ratio of the number of catastrophe
567   events observed at a certain lifetime to the total number of microtubules that reached
568   this lifetime (Gardner et al., 2011).

Acknowledgement

The authors thank Pengpeng Yu (school of life sciences and technology, TongJi University), Chunguang Wang (school of life sciences and technology, TongJi University), Peng Shi (school of basic medical sciences, Peking University) and Congying Wu (school of basic medical sciences, Peking University) for technical assistance. Special thanks to the light microscopy and protein facility in Tsinghua University. We acknowledge our funding from National Natural Sciences Foundation of China (32070704, 32370730) and IDG/McGovern Institute for Brain Research (Tsinghua University).

References

Aizawa, H., Y. Sekine, R. Takemura, Z.Z. Zhang, M. Nangaku, and N. Hirokawa. 1992. KINESIN FAMILY IN MURINE CENTRAL-NERVOUS-SYSTEM. *Journal of Cell Biology*. 119:1287-1296.

Alberts, B. 2015. Molecular biology of the cell.

Asenjo, A.B., C. Chatterjee, D. Tan, V. DePaoli, W.J. Rice, R. Diaz-Avalos, M. Silvestry, and H. Sosa. 2013. Structural Model for Tubulin Recognition and Deformation by Kinesin-13 Microtubule Depolymerases. *Cell Reports*. 3:759-768.

Ayaz, P., X. Ye, P. Huddleston, C.A. Brautigam, and L.M. Rice. 2012. A TOG:alpha beta-tubulin Complex Structure Reveals Conformation-Based Mechanisms for a Microtubule Polymerase. *Science*. 337:857-860.

Benoit, M.P.M.H., A.B. Asenjo, and H. Sosa. 2018. Cryo-EM reveals the structural basis of microtubule depolymerization by kinesin-13s. *Nature Communications*. 9.

Brouhard, G.J., J.H. Stear, T.L. Noetzel, J. Al-Bassam, K. Kinoshita, S.C. Harrison, J. Howard, and A.A. Hyman. 2008. XMAP215 is a processive microtubule polymerase. *Cell*. 132:79-88.

Cooper, J.R., M. Wagenbach, C.L. Asbury, and L. Wordeman. 2010. Catalysis of the microtubule on-rate is the major parameter regulating the depolymerase activity of MCAK. *Nature Structural & Molecular Biology*. 17:77-U98.

Demchouk, A.O., M.K. Gardner, and D.J. Odde. 2011. Microtubule Tip Tracking and Tip Structures at the Nanometer Scale Using Digital Fluorescence Microscopy. *Cellular and Molecular Bioengineering*. 4:192-204.

Desai, A., S. Verma, T.J. Mitchison, and C.E. Walczak. 1999. Kin I kinesins are microtubule-destabilizing enzymes. *Cell*. 96:69-78.

Duellberg, C., N.I. Cade, D. Holmes, and T. Surrey. 2016. The size of the EB cap determines instantaneous microtubule stability. *eLife*. 5:23.

Farmer, V., G. Arpa, S.L. Hall, and M. Zanic. 2021. XMAP215 promotes microtubule catastrophe by disrupting the growing microtubule end. *Journal of Cell Biology*. 220.

Gardner, M.K., M. Zanic, C. Gell, V. Bormuth, and J. Howard. 2011. Depolymerizing Kinesins Kip3 and MCAK Shape Cellular Microtubule Architecture by Differential Control of Catastrophe. *Cell*. 147:1092-1103.

Gell, C., V. Bormuth, G.J. Brouhard, D.N. Cohen, S. Diez, C.T. Friel, J. Helenius, B. Nitzsche, H. Petzold, J. Ribbe, E. Schaeffer, J.H. Stear, A. Trushko, V. Varga, P.O. Widlund, M. Zanic, and J. Howard. 2010. Microtubule Dynamics Reconstituted In Vitro and Imaged by Single-Molecule Fluorescence Microscopy. *In* Microtubules, in Vitro: Microtubules, in Vitro. Vol. 95. L. Wilson and J.J. Correia, editors. 221-245.

Ghosh-Roy, A., A. Goncharov, Y. Jin, and A.D. Chisholm. 2012. Kinesin-13 and Tubulin Posttranslational Modifications Regulate Microtubule Growth in Axon Regeneration. *Dev. Cell*. 23:716-728.

Gouveia, S.M., K. Leslie, L.C. Kapitein, R.M. Buey, I. Grigoriev, M. Wagenbach, I. Smal, E. Meijering, C.C. Hoogenraad, L. Wordeman, M.O. Steinmetz, and A. Akhmanova. 2010. In Vitro Reconstitution of the Functional Interplay between MCAK and EB3 at Microtubule Plus Ends. *Curr. Biol.* 20:1717-1722.

Guesdon, A., F. Bazilel, R.M. Buey, R. Mohan, S. Monier, R.R. Garcia, M. Angevin, C. Heichette, R.

Wieneke, R. Tampe, L. Duchesne, A. Akhmanova, M.O. Steinmetz, and D. Chretien. 2016. EB1 interacts with outwardly curved and straight regions of the microtubule lattice. *Nature Cell Biology*. 18:1102-1108.

Helenius, J., G. Brouhard, Y. Kalaidzidis, S. Diez, and J. Howard. 2006. The depolymerizing kinesin MCAK uses lattice diffusion to rapidly target microtubule ends. *Nature*. 441:115-119.

Hoeoeg, J.L., S.M. Huisman, Z. Seboe-Lemke, L. Sandblad, J.R. McIntosh, C. Antony, and D. Brunner. 2011. Electron tomography reveals a flared morphology on growing microtubule ends. *Journal of Cell Science*. 124:693-698.

Honnappa, S., S.M. Gouveia, A. Weisbrich, F.F. Damberger, N.S. Bhavesh, H. Jawhari, I. Grigoriev, F.J.A. van Rijssel, R.M. Buey, A. Lawera, I. Jelesarov, F.K. Winkler, K. Wuethrich, A. Akhmanova, and M.O. Steinmetz. 2009. An EB1-Binding Motif Acts as a Microtubule Tip Localization Signal. *Cell*. 138:366-376.

Hunter, A.W., M. Caplow, D.L. Coy, W.O. Hancock, S. Diez, L. Wordeman, and J. Howard. 2003. The kinesin-related protein MCAK is a microtubule depolymerase that forms an ATP-hydrolyzing complex at microtubule ends. *Molecular Cell*. 11:445-457.

Janke, C., and M.M. Magiera. 2020. The tubulin code and its role in controlling microtubule properties and functions. *Nature Reviews Molecular Cell Biology* 21:307-326.

Kapitein, L.C., and C.C. Hoogenraad. 2015. Building the Neuronal Microtubule Cytoskeleton. *Neuron*. 87:492-506.

Kline-Smith, S.L., A. Khodjakov, P. Hergert, and C.E. Walczak. 2004. Depletion of centromeric MCAK leads to chromosome congression and segregation defects due to improper kinetochore attachments. *Molecular Biology of the Cell*. 15:1146-1159.

Lee, T., K.J. Langford, J.M. Askham, A. Bruning-Richardson, and E.E. Morrison. 2008. MCAK associates with EB1. *Oncogene*. 27:2494-2500.

Manka, S.W., and C.A. Moores. 2018. The role of tubulin-tubulin lattice contacts in the mechanism of microtubule dynamic instability. *Nature Structural & Molecular Biology*. 25:607-+.

Maurer, S.P., P. Bieling, J. Cope, A. Hoenger, and T. Surrey. 2011. GTP gamma S microtubules mimic the growing microtubule end structure recognized by end-binding proteins (EBs). *Proceedings of the National Academy of Sciences of the United States of America*. 108:3988-3993.

Maurer, S.P., N.I. Cade, G. Bohner, N. Gustafsson, E. Boutant, and T. Surrey. 2014. EB1 Accelerates Two Conformational Transitions Important for Microtubule Maturation and Dynamics. *Curr. Biol.* 24:372-384.

Maurer, S.P., F.J. Fourniol, G. Bohner, C.A. Moores, and T. Surrey. 2012. EBs Recognize a Nucleotide-Dependent Structural Cap at Growing Microtubule Ends. *Cell*. 149:371-382.

McIntosh, J.R., E. O'Toole, G. Morgan, J. Austin, E. Ulyanov, F. Ataullakhanov, and N. Gudimchuk. 2018. Microtubules grow by the addition of bent guanosine triphosphate tubulin to the tips of curved protofilaments. *Journal of Cell Biology*. 217:2691-2708.

Moores, C.A., and R.A. Milligan. 2006. Lucky 13 - microtubule depolymerisation by kinesin-13 motors. *Journal of Cell Science*. 119:3905-3913.

Noda, Y., R. Satoyoshitake, S. Kondo, M. Nangaku, and N. Hirokawa. 1995. KIF2 IS A NEW MICROTUBULE-BASED ANTEROGRADE MOTOR THAT TRANSPORTS MEMBRANOUS ORGANELLES DISTINCT FROM THOSE CARRIED BY KINESIN HEAVY-CHAIN OR KIF3A/B. *Journal of Cell Biology*. 129:157-167.

Patel, J.T., H.R. Belsham, A.J. Rathbone, B. Wickstead, C. Gell, and C.T. Friel. 2016. The
family-specific alpha 4-helix of the kinesin-13, MCAK, is critical to microtubule end
recognition. *Open Biology*. 6.

Piao, T., M. Luo, L. Wang, Y. Guo, D. Li, P. Li, W.J. Snell, and J. Pan. 2009. A microtubule
depolymerizing kinesin functions during both flagellar disassembly and flagellar assembly in
Chlamydomonas. *Proceedings of the National Academy of Sciences of the United States of*
America. 106:4713-4718.

Puri, D., K. Ponniah, K. Biswas, A. Basu, S. Dey, E.A. Lundquist, and A. Ghosh-Roy. 2021. Wnt
signaling establishes the microtubule polarity in neurons through regulation of Kinesin-13.
Journal of Cell Biology. 220.

Song, Y., Y. Zhang, Y. Pan, J. He, Y. Wang, W. Chen, J. Guo, H. Deng, Y. Xue, X. Fang, and X. Liang.
2020. The microtubule end-binding affinity of EB1 is enhanced by a dimeric organization that
is susceptible to phosphorylation. *Journal of Cell Science*. 133.

Tan, D., W.J. Rice, and H. Sosa. 2008. Structure of the Kinesin13-Microtubule Ring Complex.
Structure. 16:1732-1739.

Tinevez, J.-Y., N. Perry, J. Schindelin, G.M. Hoopes, G.D. Reynolds, E. Laplantine, S.Y. Bednarek, S.L.
Shorte, and K.W. Eliceiri. 2017. TrackMate: An open and extensible platform for
single-particle tracking. Methods. 115:80-90.

Tournebize, R., A. Popov, K. Kinoshita, A.J. Ashford, S. Rybina, A. Pozniakovsky, T.U. Mayer, C.E.
Walczak, E. Karsenti, and A.A. Hyman. 2000. Control of microtubule dynamics by the
antagonistic activities of XMAP215 and XKCM1 in Xenopus egg extracts. *Nature Cell*
Biology. 2:13-19.

Trofimova, D., M. Paydar, A. Zara, L. Talje, B.H. Kwok, and J.S. Allingham. 2018. Ternary complex of
Kif2A-bound tandem tubulin heterodimers represents a kinesin-13-mediated microtubule
depolymerization reaction intermediate. *Nature Communications*. 9.

Vasudevan, K.K., Y.-Y. Jiang, K.F. Lechtreck, Y. Kushida, L.M. Alford, W.S. Sale, T. Hennessey, and J.
Gaertig. 2015. Kinesin-13 regulates the quantity and quality of tubulin inside cilia. *Molecular*
Biology of the Cell. 26:478-494.

Vitre, B., F.M. Coquelle, C. Heichette, C. Garnier, D. Chretien, and I. Arnal. 2008. EB1 regulates
microtubule dynamics and tubulin sheet closure in vitro. *Nature Cell Biology*. 10:415-U481.

Walczak, C.E., E.C. Gan, A. Desai, T.J. Mitchison, and S.L. Kline-Smith. 2002. The
microtubule-destabilizing kinesin XKCM1 is required for chromosome positioning during
spindle assembly. *Curr. Biol.* 12:1885-1889.

Walczak, C.E., T.J. Mitchison, and A. Desai. 1996. XKCM1: a Xenopus kinesin-related protein that
regulates microtubule dynamics during mitotic spindle assembly. *Cell*. 84:37-47.

Wang, W., S. Cantos-Fernandes, Y. Lv, H. Kuerban, S. Ahmad, C. Wang, and B. Gigant. 2017. Insight
into microtubule disassembly by kinesin-13s from the structure of Kif2C bound to tubulin.
Nature Communications. 8.

Wang, W., T. Shen, R. Guerois, F. Zhang, H. Kuerban, Y. Lv, B. Gigant, M. Knossow, and C. Wang.
2015. New Insights into the Coupling between Microtubule Depolymerization and ATP
Hydrolysis by Kinesin-13 Protein Kif2C. *Journal of Biological Chemistry*. 290:18721-18731.

Widlund, P.O., M. Podolski, S. Reber, J. Alper, M. Storch, A.A. Hyman, J. Howard, and D.N. Drechsel.
2012. One-step purification of assembly-competent tubulin from diverse eukaryotic sources.

Molecular Biology of the Cell. 23:4393-4401.

Wittmann, T., A. Hyman, and A. Desai. 2001. The spindle: a dynamic assembly of microtubules and

motors. *Nature Cell Biology*. 3:E28-E34.

Wordeman, L., and T.J. Mitchison. 1995. IDENTIFICATION AND PARTIAL CHARACTERIZATION

OF MITOTIC CENTROMERE-ASSOCIATED KINESIN, A KINESIN-RELATED

PROTEIN THAT ASSOCIATES WITH CENTROMERES DURING MITOSIS. *Journal of*

Cell Biology. 128:95-105.

Zong, H., M. Hazelbaker, C. Moe, S.C. Ems-McClung, K. Hu, and C.E. Walczak. 2021. Spatial

regulation of MCAK promotes cell polarization and focal adhesion turnover to drive robust

cell migration. *Molecular Biology of the Cell*. 32:590-604.

Figure 1. MCAK preferentially binds to growing microtubule ends

(A) The representative kymographs of single-molecule GFP-MCAK (green) binding events on dynamic microtubules (red, tubulin: 16 µM) growing from GMPCPP microtubule seeds (blue) in the presence of 1 mM ATP, AMPPNP, ADP and APO (the nucleotide-free state). The binding events at the plus-end and lattice were indicated by a white arrowhead and a blue arrowhead, respectively. Vertical bar: 5 s; horizontal bar: 2 µm.

(B) The plot showing the representative intensity profiles of a single GFP-MCAK molecule (green) and a growing microtubule end (red). The peak position of a GFP-MCAK molecule was recorded as the binding location (dashed line in green).

(C) The spatial distributions of the binding sites of GFP-MCAK (green, n=152 events) along the long axis of microtubules. The averaged intensity profile of the growing

microtubule ends (red) was shown as the positional reference.

(D) Statistical quantification of the apparent association constant (k_{on}) of GFP-MCAK
on the plus-end and lattice of dynamic microtubules in the presence of 1 mM ATP
(n=66 microtubules from 3 assays), AMPPNP (n=29 microtubules from 2 assays),
ADP (n=21 microtubules from 2 assays) and APO (n=36 microtubules from 3 assays).
The statistical comparison was made versus the k_{on} of MCAK on the corresponding
location (i.e. plus end or lattice) in the ATP condition.

(E) Statistical quantification of $R_{E/L}$ in the presence of ATP, AMPPNP, ADP and APO.
The statistical comparison was made versus the $R_{E/L}$ of the plus-end binding events in
the ATP condition.

(F) Statistical quantification of the dwell time of GFP-MCAK on the growing end
and lattice of dynamic microtubules in the presence of ATP (n=966 events from 3 as-
says for the plus end; n=702 events from 3 assays for lattice), AMPPNP (n=231
events from 2 assays for the plus end; n=142 events from 2 assays for lattice), ADP
(n=327 events from 2 assays for the plus end; n=1184 events from 2 assays for lattice)
and APO (n=71 events from 3 assays for the plus end; n=573 events from 3 assays for
lattice). The statistical comparison was made versus dwell time of the binding events
on the corresponding location (i.e. plus end or lattice) in the ATP condition.

(G) The mean-squared displacement (MSD) of GFP-MCAK was plotted against the
time interval (t, 0.1s per frame). The diffusion coefficient (D) of 0. 023 $\mu m^2 s^{-1}$ was
obtained using a linear fitting ($<x^2>= 2Dt$). The error bar represented SEM (n=203
trajectories).

In panel **D**, **E** and **F**, all the data were presented as mean ± SEM. All the comparisons
were performed using the two-tailed Mann-Whitney U test with Bonferroni correction,
n.s., no significance; ***, p<0.001.

Figure 2. MCAK binds to the entire GTP cap of growing microtubule ends

(**A**) The representative kymograph showing the individual binding events of GFP-MCAK (1 nM, with 1 mM ATP, left), EB1-GFP (10 nM, middle) or XMAP215-GFP (1 nM, right) at growing microtubule ends (red, tubulin: 12 μM). The plots showed the representative intensity profiles of a single molecule (green) and a growing microtubule end (red). The peak position of individual molecules was determined using a Gaussian fit (dashed line in green). Vertical bar: 5 s; horizontal bar: 2 μm.

(**B**) The spatial distributions of the binding sites of GFP-MCAK (green, n=123 events), EB1-GFP (black, n=184 events) and XMAP215-GFP (blue, 142 events) along the long axis of microtubules. The averaged intensity profile of the growing microtubule ends (red) was shown as the positional reference. The region within the FWHM of the blue curve was considered to be the binding region of XMAP215. The MCAK molecules localized to the left of this region was considered to be proximally distributed (grey bar). The region within the FWHM of the black curve was considered to be the binding region of EB1. The MCAK molecules localized to the right of this region was considered to be distally distributed (orange bar).

(**C**) The averaged intensity profiles of the growing microtubule ends were used to quantify the localization of GFP-MCAK (green, 123 microtubules), EB1-GFP (black, 184 microtubules) and XMAP215-GFP (blue, 142 microtubules). Note that the morphologies of microtubule ends in three conditions were nearly identical. In panels **B** and **C**, the red arrows indicated the direction of microtubule growth.

Figure 3. MCAK strongly binds to GTPγS microtubules in a nucleotide-independent
manner

(A) The representative projection images of GFP-MCAK binding to GTPγS (red ar-
rowhead), GDP (purple arrowhead) and GMPCPP microtubules (cyan arrowhead) in
the presence of 1 mM AMPPNP (left), 1 mM ATP (left middle), 1 mM ADP (right
middle) and at the APO state (right). Note that the binding on GDP or GMPCPP mi-
crotubules was compared to that on GTPγS microtubules in the same flow cell. Scale
bar: 5 μm.

(B) Statistical comparison of the normalized fluorescence intensity of GFP-MCAK
on different microtubules in the presence of 1 mM AMPPNP (112 GTPγS microtu-
bules, 56 GMPCPP microtubules, 56 GDP microtubules from 3 assays), 1 mM ATP
(88 GTPγS microtubules, 60 GMPCPP microtubules, 28 GDP microtubules from 3
assays), 1 mM ADP (95 GTPγS microtubules, 51 GMPCPP microtubules, 44 GDP
microtubules from 3 assays) or at the APO state (66 GTPγS microtubules, 30 GMP-
CPP microtubules, 36 GDP microtubules from 3 assays). All data were normalized to
the binding intensity of GFP-MCAK (1 nM) on GTPγS microtubules in the AMPPNP
condition.

(C) The ratios of the binding intensity of GFP-MCAK on GTPγS microtubules to that
on GDP or GMPCPP microtubules in various nucleotide conditions. The dashed line

799 on the plot represented 1. The statistical comparisons were performed between the
800 ratios and 1.

801 In panel **B** and **C**, All the data were presented as mean ± std. All the comparisons
802 were performed using the two-tailed Mann–Whitney U test with Bonferroni correc-
803 tion, ***, $p<0.001$.

Figure 4. MCAK strongly binds to the ends of GMPCPP microtubules in a nucleotide state-dependent manner

(A) The representative projection images and intensity profiles showing GFP-MCAK binding (upper) at the ends of GMPCPP microtubules (lower) in the presence of 1 mM AMPPNP (left), 1 mM ATP (left middle), 1 mM ADP (right middle) or at the APO state (right). Red arrowhead: the end-binding of MCAK. Red bar: ends. Black bar: lattice. Scale bar=2 μm.

(B) Statistical quantification of the binding intensity of GFP-MCAK on the lattice and end of GMPCPP microtubules in the presence of 1 mM AMPPNP (47 microtubules from 3 assays), 1 mM ATP (40 microtubules from 3 assays), 1 mM ADP (45 microtubules from 3 assays) or at the APO state (52 microtubules from 3 assays). All data were normalized to the binding of GFP-MCAK (1 nM) on GTPγS microtubules in the AMPPNP condition. All the data were presented as mean ± std. The statistical analysis was performed using the two-tailed pair *t*-test by Bonferroni correction, ***, p<0.001.

(C) The ratios between the binding intensity of GFP-MCAK at the end and the intensity on the lattice of GMPCPP microtubules in various nucleotide conditions. The ratios were presented as mean ± std. The dashed line on the plot represents 1. The statistical comparisons were performed between the ratios and 1 using the two-tailed Mann-Whitney U test with Bonferroni correction, ***, p<0.001.

Figure 5. The preference for GDP·Pi-tubulins facilitates the binding preference of MCAK for growing microtubule ends

(A) The structural model of the MCAK^{sN+M}-tubulin complex (PDB ID: 5MIO). The cartoon schematic in upper panel showed the domain organization of MCAK. Loop2 and α4 helix, two regions mediating the interaction between MCAK and tubulin, were enlarged and the key sites (K524, V298) were highlighted in red. The corresponding residues of tubulin that may interact with K524 and V298 were highlighted in green.

(B) The representative projection images of GFP-MCAK^{K524A} and GFP-MCAK^{V298S} binding to GTPγS (red arrowhead), GDP (purple arrowhead) and GMPCPP microtubules (cyan arrowhead) in the presence of 1 mM AMPPNP. Scale bar: 5 μm.

(C) Statistical quantification of the binding intensity of GFP-MCAK^{K524A} (144
GTPγS microtubules, 72 GMPCPP microtubules, 62 GDP microtubules from 3 assays)
and GFP-MCAK^{V298S} (124 GTPγS microtubules, 65 GMPCPP microtubules, 59 GDP
microtubules from 3 assays) on different microtubules. Note that all data were nor-
malized to the binding intensity of GFP-MCAK on GTPγS microtubules in the
AMPPNP condition. The data were presented as mean ± std.

(D) The ratios of the binding intensity of GFP-MCAK^{K524A} or GFP-MCAK^{V298S} on
GTPγS microtubules to that on GDP or GMPCPP microtubules in the presence of 1
mM AMPPNP. Purple dashed line: The GTPγS/GDP ratio of GFP-MCAK. The cyan
dashed line: the GTPγS/GMPCPP ratio of GFP-MCAK. The data were presented as
mean ± std. The statistical comparisons were made versus the corresponding value of
GFP-MCAK.

(E) The representative projection images and intensity profiles showing the binding
of GFP-MCAK^{K524A} and GFP-MCAK^{V298S} to the end (red arrowhead in the upper
panels) and lattice of GMPCPP microtubules (lower panels) in the presence of
AMPPNP. Red bar: ends. Black bar: lattice. Scale bar=2 μm.

(F) Statistical quantification of the binding intensity of GFP-MCAK^{K524A} (88 micro-
tubules from 3 assays) and GFP-MCAK^{V298S} (68 microtubules from 3 assays) on the
lattice and end of GMPCPP microtubules in the presence of AMPPNP (lower panel).
All the data were normalized to the binding of GFP-MCAK (1 nM) on GTPγS micro-
tubules in the AMPPNP condition. All the binding intensity data were presented as
mean ± std. The statistical analyses were performed using the two-tailed paired t-test
by Bonferroni correction, n.s., no significance; ***, p<0.001. The inset (upper) show-
ing the ratios of the end-binding intensity of GFP-MCAK^{K524A} or GFP-MCAK^{V298S} to
the lattice-binding intensity on GMPCPP microtubules in the presence of AMPPNP.
The ratios were presented as mean ± std. The statistical comparisons were made ver-
sus the ratio of GFP-MCAK (the dashed line shown in **Fig. 4C)**.

(G) The representative kymographs of the single-molecule binding events of
GFP-MCAK^{K524A} (10 nM, green) and GFP-MCAK^{V298S} (30 nM, green) on growing
microtubules (red, tubulin: 16 μM) in the presence of 1 mM ATP. The end-binding
events were indicated using white arrowheads and enlarged. For the original kymo-
graphs, vertical bar: 2 s; horizontal bar: 2 μm. For the enlarged kymographs, vertical
bar: 1 s; horizontal bar: 0.5 μm.

(H) Statistical quantification of the apparent association constant (k_{on-P}) of
GFP-MCAK^{K524A} (33 microtubules from 3 assays) and GFP-MCAK^{V298S} (23 micro-
tubules from 2 assays) on growing microtubule ends in the presence of ATP (lower).
The statistical comparisons were made versus the k_{on-P} of GFP-MCAK. The inset
(upper) showing the statistical quantification of $R_{E/L}$ of GFP-MCAK^{K524A} and
GFP-MCAK^{V298S}. The $R_{E/L}$ of both mutants were compared to that of GFP-MCAK in
the ATP condition. All the data were presented as mean ± SEM.

(I) Statistical quantification of the dwell time of GFP-MCAK^{K524A} (19 binding events
from 3 assays) and GFP-MCAK^{V298S} (43 binding events from 3 assays) on growing
microtubule ends in the presence of ATP. The dwell time were presented as mean ±
SEM. The dwell time of GFP-MCAK^{K524A} and GFP-MCAK^{V298S} were compared to
that of GFP-MCAK.

In panel **C**, **D**, **F** (the upper inset), **H** and **I**, all the comparisons were performed using
the two-tailed Mann-Whitney U test with Bonferroni correction, n.s., no significance;
*, p<0.05; **, p<0.01; ***, p<0.001.

Figure 6. Functional specification of MCAK and XMAP215 at growing microtubule ends

(**A**) The representative kymographs of dynamic microtubules (tubulin: 10 μM) in the control condition or in the presence of MCAK, XMAP215 or both. Vertical bar: 100 s; horizontal bar: 2 μm.

(**B**) The probability distribution of microtubule lifetime in the experimental conditions indicated in the panel **A**. The lines were gamma fitting curves. Control: 443 microtubules from 7 assays. 20 nM MCAK: 621 microtubules from 7 assays. 50 nM XMAP215: 237 microtubules from 3 assays. 20 nM MCAK+50 nM XMAP215: 283 microtubules from 3 assays.

(**C**) The plots of catastrophe frequency versus the lifetime of microtubules showing how the likelihood of catastrophe depended on the age of microtubules. The data were from the experiments shown in panel **A**.

Figure 7. Cartoon schematics depicting the working model of MCAK at growing microtubule ends

(A) A hypothetic model for the binding cycle of MCAK, given its binding preference on the EB cap. T: MCAK·ATP. D: MCAK·ADP.

(B) The co-operation schematics of MCAK, EB1 and XMAP215 at growing microtubule ends. Pathway 1: the direct binding of MCAK to microtubule ends. Pathway 2: the indirect binding of MCAK to microtubule ends via EB1.

Supplementary Information Appendix

Title:

Recognition of the Nucleotide-Dependent Feature Facilitates the Microtubule End-Binding of
MCAK

Authors and affiliations:

Wei Chen[1] (chenwill@mail.tsinghua.edu.cn)

Yin-Long Song[1] (y.song4@uu.nl)

Jian-Feng He[1] (jianfenghe2015@mail.tsinghua.edu.cn)

Xin Liang[1,*] (xinliang@tsinghua.edu.cn)

[1] IDG/McGovern Institute for Brain Research, School of Life Sciences, Tsinghua University, Bei-
jing 100084, China

Corresponding author:

Name: Xin Liang
Address: Room E118, Medical School, Tsinghua University, 100084 Beijing, China
Tel: +86-10-62772563
Email: xinliang@tsinghua.edu.cn
ORCID: 0000-0001-7915-8094

Supplementary materials:

Figures s1-s4 (with legends)

Figure s1. Single-molecule fluorescence analysis and the apparent off-rates (k_{off}) of
GFP-MCAK

(A) The fluorescence intensity distribution of GFP-MCAK (red, n=509 binding events)
was fitted using a Gaussian function. The background intensity was subtracted. The
intensity range ($\mu \pm 2\sigma$=300 ± 145 A.U.) was used to determine if a fluorescence spot
represents a single dimer.

(B) The apparent off-rate (k_{off}) of GFP-MCAK on growing microtubule ends in the
presence of ATP (minus end, square; plus end, circle), AMPPNP (down triangle), ADP
(up triangle) and APO (diamond). k_{off} was calculated by fitting the dwell time of indi-
vidual GFP-MCAK binding events to a single exponential function.

Figure s2. Projection of single-molecule fluorescence images

The representative images showing how to generate a summation image using 1000
frames of raw images. The GDP and GTPγS microtubule were indicated using green
and yellow arrowheads, respectively. The concentration of GFP-MCAK here was 1
nM. The experiment was performed in the presence of 1 mM AMPPNP. Scale bar: 5
μm.

Figure s3. GFP-MCAK[K525A] and GFP-MCAK[V298S] are depolymerizing-deficient mutants

953
954

(A) The representative kymographs showing the depolymerizing effect of GFP-MCAK, GFP-MCAK[K525A] and GFP-MCAK[V298S] on GMPCPP-stabilized microtubules in the presence of 1 mM ATP. Vertical bar: 100 s, horizontal bar: 2 μm.

955
956
957

(B) Statistical quantification of the depolymerization rates of GFP-MCAK (n=21 microtubules from 3 assays), GFP-MCAK[K525A] (n=11 microtubules from 2 assays) and GFP-MCAK[V298S] (n=52 microtubules from 3 assays). The data were presented as mean ± std. The statistical analysis was performed using the two-tailed Mann-Whitney U test with Bonferroni correction, ***, $p<0.001$.

965 **Figure s4.** Structural models showing the kinesin-13-tubulin contact site and the
966 TOG-tubulin contact site

967 The structural model showing that the TOG domain (PDB ID: 4FFB) and the motor
968 domain of MCAK (PDB ID: 6BBN) shared, to some extent, the binding site on a tu-
969 bulin dimer.