Title Page

Title:

MCAK Recognizes the Nucleotide-Dependent Feature at Growing Microtubule Ends

Authors and affiliations:

Wei Chen[1] (chenwill@mail.tsinghua.edu.cn)

Yin-Long Song[1] (y.song4@uu.nl)

Jian-Feng He[1] (jianfenghe2015@mail.tsinghua.edu.cn)

Xin Liang[1,*] (xinliang@tsinghua.edu.cn)

[1] IDG/McGovern Institute for Brain Research, State Key Laboratory of Complex, Severe, and Rare
Diseases, School of Life Sciences, Tsinghua University, Beijing 100084, China

Corresponding author:

Name: Xin Liang
Address: Room E118, Medical School, Tsinghua University, 100084 Beijing, China
Tel: +86-10-62772563
Email: xinliang@tsinghua.edu.cn
ORCID: 0000-0001-7915-8094

Key words

MCAK
Microtubule dynamics
End binding
Depolymerization
GTP cap

MCAK Recognizes the Nucleotide-Dependent Feature at Growing Microtubule Ends

Wei Chen[1], Yin-Long Song[1], Jian-Feng He[1], Xin Liang[1,*]

[1] IDG/McGovern Institute for Brain Research, School of Life Sciences, Tsinghua University, Beijing 100084, China

* Correspondence: xinliang@tsinghua.edu.cn

Abstract

The growing plus-end is a key regulatory site for microtubule dynamics. MCAK (mitotic centromere-associated kinesin), a microtubule depolymerizing kinesin, is an end-binding regulator of catastrophe frequency. It is intriguing how MCAK specifically binds to growing microtubule ends. Here, we measure the end-binding kinetics of MCAK using single-molecule imaging, and reveal that MCAK binds to the entire GTP cap, including the region where EB1 preferentially binds and the distalmost tip. Further analysis shows that MCAK strongly binds to GTPγS microtubules, and this binding preference is dependent on the nucleotide state of MCAK. This finding suggests that MCAK recognizes the nucleotide-dependent feature of microtubules. Moreover, we show that although MCAK and XMAP215 partly share binding regions at the distalmost ends, their regulatory effects can be additive, highlighting a simple rule of how the end-binding regulators coordinate their activities. Overall, our findings provide new insights into how MCAK regulates microtubule end dynamics.

(148 words)

Introduction

Microtubules are essential components of the eukaryotic cytoskeleton, supporting a wide range of cellular processes, including cell division, polarization, and migration (Alberts, 2015). However, their properties can vary considerably across different cellular contexts and structures (Janke and Magiera, 2020; Kapitein and Hoogenraad, 2015; Wittmann et al., 2001). This variability reflects the complex and tightly regulated mechanisms that govern the dynamics of cellular microtubules, among which the microtubule-associated proteins (MAPs) play crucial roles.

Kinesin superfamily members not only facilitate the transport of cargo along microtubules but also serve as regulators of microtubule dynamics, as exemplified by the kinesin-13 family. Initially identified through studies of spindle functions and neurons (Aizawa et al., 1992; Noda et al., 1995; Walczak et al., 1996; Wordeman and Mitchison, 1995), kinesin-13 has since been implicated in a broad range of cellular processes, including spindle assembly (Walczak et al., 2002), chromosome segregation (Kline-Smith et al., 2004), directional migration (Zong et al., 2021), ciliary length control (Piao et al., 2009; Vasudevan et al., 2015), and neuronal polarization (Ghosh-Roy et al., 2012; Puri et al., 2021).The loss of kinesin-13 often results in altered structural and dynamic properties of cellular microtubules. Moreover, the subcellular localizations of kinesin-13, such as at growing microtubule ends, centrosomes, and centromeres, are key regulatory sites for microtubule dynamics. These observations suggest that kinesin-13 family members are key regulators for the length, mass, and polarity of cellular microtubules.

As a representative member of the kinesin-13 family, MCAK has been characterized as both a microtubule depolymerase and a catastrophe factor (Desai et al., 1999; Gardner et al., 2011; Hunter et al., 2003; Kline-Smith and Walczak, 2002; Walczak et al., 1996). It has been proposed that MCAK targets growing microtubule ends by interacting with end-binding proteins (EBs) or motors, such as Kif18b (Gouveia et al., 2010; Honnappa et al., 2009; McHugh and Welburn, 2023; Tanenbaum et al., 2011). Upon binding, MCAK destabilizes the protective GTP cap at the growing microtubule ends by bending and disassembling the underlying substrate, including terminal tubulin dimers or protofilaments (Benoit et al., 2018; Moores and Milligan, 2006; Wang et al., 2017), thereby increasing the frequency of catastrophe events. MCAK can also function independently (Desai et al., 1999; Gardner et al., 2011; Hunter et al., 2003), potentially targeting stabilized microtubule ends through direct binding or rapid diffusion along the microtubule lattice (Desai et al., 1999; Helenius et al., 2006; Wang et al., 2012). Previous structural studies suggest that MCAK recognizes curved protofilaments at microtubule ends (Asenjo et al., 2013; Benoit et al., 2018; Tan et al., 2008; Trofimova et al., 2018; Wang et al., 2017). However, these studies primarily focus on stabilized microtubules, short protofilaments, or tubulin dimers, leaving the

mechanism by which MCAK binds to growing microtubule ends unresolved.

In this study, we use single-molecule imaging to investigate the direct binding of
MCAK to growing microtubule ends. A key finding is that MCAK binds throughout
the entire GTP cap, encompassing both the EB cap, where EB1 preferentially binds,
and the distalmost cap, where XMAP215 binds to curved protofilaments. Notably, this
is the first discovery that MCAK exhibits a binding preference to the EB cap. Further
analysis reveals that MCAK strongly binds to GTPγS microtubules, which are thought
to mimic GDP·Pi-tubulin conformations enriched in the EB cap. Having confirmed that
MCAK binds to the entire GTP cap structure, we next examine how MCAK co-
regulates microtubule dynamics in collaboration with XMAP215, and reveal their
additive, rather than mutually exclusive, regulatory effects. Overall, this study provides
novel insights into how MCAK binds to and acts at growing microtubule ends.

Results

MCAK shows a binding preference for growing microtubule ends

We began by measuring the single-molecule binding kinetics of MCAK at growing microtubule ends using the in vitro microtubule dynamics assay. This experiment was performed in BRB80 supplemented with 50 mM KCl and 1 mM ATP, providing a near-physiological ion strength of approximately 0.2 M, similar to what was previously reported (Scopes, 2002; Storey, 2004). The size-exclusion chromatography analysis showed that the purified GFP-MCAK exists as dimers under this condition (**Fig. s1**).

We observed MCAK binding events both at growing microtubule ends and along the lattice (**Fig. 1A**). To characterize these events, we measured the fluorescence intensity of each binding event (see Methods) and plot a probability distribution of all binding events (**Fig. s1**). We used the range of fluorescence intensity from the probability distribution ($\mu \pm 2\sigma$: 300 ± 145 A.U.) to identify single MCAK dimers. Using a custom software (Maurer et al., 2014; Song et al., 2020), we determined the end-binding position of each individual MCAK molecule in each frame by fitting a Gaussian function to the intensity profile along the longitudinal axis of the microtubule, with the plus-end serving as the positional reference (**Fig. 1B**). The average position over the dwell time of each MCAK molecule was taken as the position of this binding event (**Fig. s2**). From this, we generated a probability distribution of all binding positions, derived from 152 binding events across three assays (**Fig. 1C**). This analysis revealed that the MCAK binding region was 162 nm in length (FWHM, Full Width at Half Maximum), corresponding to approximately 20 layers of tubulin dimers. We categorized the binding events of MCAK within this region as "end binding" and those located more proximally as "lattice binding".

To investigate the binding kinetics, we first compared the apparent association rates of MCAK on the plus-end (k_{on-P}) and the lattice (k_{on-L}) in the presence of ATP. The ratio ($R_{E/L}$) of k_{on-P} ((22.6 ± 1.4) $\times 10^{-6}$ s^{-1} nM^{-1}, per tubulin dimer, mean$\pm$SEM, hereinafter unless otherwise specified) to k_{on-L} ((2.8 ± 0.3) $\times 10^{-6}$ s^{-1} nM^{-1}) was 13 ± 1 (n=66 microtubules from 3 assays) (**Fig. 1 D-E**). We also measured the dwell time, which reflects the dissociation rate (k_{off}) (**Fig. s1**). MCAK exhibited a longer dwell time at the plus-end than on the lattice (plus end: 0.75 ± 0.02 s, n=966 events from 3 assays; lattice: 0.62 ± 0.02 s, n=702 events from 3 assays, p<0.001) (**Fig. 1F**). Using these kinetic parameters, we calculated the dissociation constant (K_d) of MCAK for individual tubulin dimers at the plus-end as 69 µM. This affinity may appear relatively low, but when considering the microtubule end as a multivalent receptor with multiple MCAK-binding sites (end length=160 nm) - where only a few bound MCAK molecules are required to regulate microtubule dynamics parameters - the effective dissociation constant reaches the nanomolar range (260 nM). The MCAK's lattice binding affinity was even lower (K_{d-L}=1057 µM, per tubulin dimer), demonstrating that MCAK has a

clear end-binding preference. We also observed single-molecule binding events of MCAK at the minus-ends (**Fig. 1A**), with a similar $R_{E/L}$ (13 ± 2, n=28 microtubules from 3 assays) and K_d (40 μM, per tubulin dimer). We focused on the plus-end binding events (referred to as "end-binding" hereinafter), due to their greater physiological relevance.

Having measured the single-molecular binding events, we next investigated whether the binding of MCAK to growing microtubule ends occurs through direct end-binding (3D diffusion) or 1D lattice-diffusion in our experimental conditions (Gouveia et al., 2010; Helenius et al., 2006). These two mechanisms are not mutually exclusive. Our data revealed that the lattice-diffusion coefficient of MCAK (*D*) on the lattice of dynamic microtubules was 0.023 ± 0.001 $\mu m^2 \cdot s^{-1}$ (**Fig. 1G**), comparable to the values reported under high-salt conditions (Cooper et al., 2010; Gouveia et al., 2010) , but lower than that in low-salt conditions (Helenius et al., 2006). Because the dwell time of the lattice-binding events was approximately 0.6 s, the average length scanned by a MCAK molecule via 1D lattice-diffusion was ~160 nm, roughly the length the GTP cap. Using an established method (Helenius et al., 2006), we calculated the flux of MCAK arriving at growing ends via 1D diffusion was 0.004 s^{-1}. This was significantly lower than the observed MCAK arrival rate at the plus-end observed ($N_p = k_{on} \cdot C = 0.067$ s^{-1}. *C*: MCAK concentration). Therefore, in our experimental conditions, 1D lattice-diffusion contributes minimally to the growing end-binding of MCAK (~ 6%), suggesting that the end-binding primarily results from the direct binding pathway.

MCAK binds to the entire GTP cap

We sought to understand how MCAK recognizes growing microtubule ends. Previous studies, mostly based on tubulin dimers, short protofilaments, or stabilized microtubules, suggesting that MCAK binds microtubule ends by recognizing curved protofilaments (Asenjo et al., 2013; Benoit et al., 2018; Trofimova et al., 2018). This model predicts that MCAK binds to the distalmost end of growing microtubules, overlapping with the binding region of XMAP215 family members (i.e. the distalmost cap) (Ayaz et al., 2012; Brouhard et al., 2008). To test this hypothesis, we quantified the end-binding regions of MCAK, EB1, and XMAP215 (**Fig. 2A-B**) by measuring the FWHM of their spatial distribution. To our surprise, the probability distribution of MCAK's positions exhibited a significant difference from that of XMAP215 (p < 10^{-5}, Kolmogorov-Smirnov test). The end-binding region of MCAK (FWHM=185 nm) was longer than that of XMAP215 (FWHM=123 nm) (**Fig. 2B**). It extended proximally towards the lattice, covering the binding region of EB1 (i.e. the EB cap, FWHM=118 nm). To further understand whether MCAK binds to the distalmost cap and the EB cap with different kinetics, we compared the dwell time of MCAK at the distal and proximal locations (**Fig. 2B**). As shown in **Fig.2B**, MCAK molecules located to the left of the XMAP215 binding region were classified as proximally distributed, while those located

to the right of the EB1 binding region were considered distally distributed. No significant difference was found between the two groups (τ_{distal}=0.8 ± 0.1 s, n=28 binding events; $\tau_{proximal}$=0.7±0.1 s, n=45 binding events; p>0.3, the two-tailed Mann-Whitney U test with Bonferroni correction). Additionally, no significant differences were observed in the intensity profiles of the microtubule ends in the presence of MCAK (1 nM), EB1 (10 nM) and XMAP215 (1 nM), respectively (End$_{EB1}$ vs End$_{MCAK}$, p=0.68; End$_{MCAK}$ vs End$_{XMAP215}$, p=0.68; End$_{EB1}$ vs End$_{XMAP215}$, p=1. Kolmogorov-Smirnov test) (**Fig. 2C**).

The results suggest that MCAK's end-binding region partially overlaps with that of XMAP215, with more MCAK positioned behind XMAP215, closer to the lattice. To confirm this, we assessed the colocalization of MCAK and XMAP215. Given the low probability of both molecules simultaneously appearing at microtubule ends at low concentrations, we increased the concentration of XMAP215 to ensure its consistent tracking of growing microtubule ends while simultaneously recording single-molecule binding events of MCAK. In each frame, we measured the distance between the intensity peaks of XMAP215 and MCAK along the microtubule's longitudinal axis (**Fig. 2D**). The average distance over MCAK's binding duration was then calculated to determine its relative position to XMAP215. Subsequently, we plotted a probability distribution of all relative distances (**Fig. 2E**). This distribution peaked at -65 nm, indicating that MCAK predominantly binds behind XMAP215, with 14.5% of binding events occurring within XMAP215's binding region (distance > 0). These findings support the notion that MCAK binds to both the distalmost cap and the EB cap.

The end-binding preference of MCAK is nucleotide state-dependent

It has been reported that MCAK tightly binds to microtubules in both the AMPPNP (a non-hydrolysable ATP analogue, mimicking the ATP-bound state) and nucleotide-free states, but exhibits weaker binding in the ADP-bound state (Friel and Howard, 2011; Helenius et al., 2006; Wang et al., 2012). Moreover, kinesin-13s in the AMPPNP state preferentially form protofilament curls or rings with tubulins (Asenjo et al., 2013; Benoit et al., 2018; Tan et al., 2008). These findings suggest that the interaction between kinesin-13s and microtubules is regulated by their nucleotide-bound state. Given this, we sought to determine whether the end-binding preference of MCAK is also nucleotide state-dependent. To explore this, we measured the end-binding kinetics of MCAK in complex with AMPPNP, ADP, or in the nucleotide-free (APO) state (**Fig. 3A**), under conditions that maintain similar microtubule growth rates (**Fig. s1**).

Comparing the binding kinetics of MCAK in different nucleotide states, we found that k_{on-P} and k_{on-L} of MCAK·AMPPNP were both lower than those of MCAK·ATP (**Fig. 3 B-C**), yet resulting in a similar end-binding preference ($R_{E/L}$=11 ± 2, n=29 microtubules from 2 assays, p>1). However, the binding of MCAK·AMPPNP at the end

and lattice both showed a longer dwell time compared to that of MCAK·ATP ($3.32 \pm$ 0.24 s at the end, n=231 events from 2 assays, $p<0.001$; 1.46 ± 0.21 s at lattice, n=142 events from 2 assays, $p<0.001$), indicating lower k_{off} (**Fig. 3D** and **Fig. s1**). As a result of the proportionally reduced k_{on-P} and k_{off}, the end-binding affinity of MCAK·AMPPNP ($K_{d-P}=61$ μM, per tubulin dimer, hereinafter) was comparable to that of MCAK·ATP. In contrast, MCAK·ADP showed nearly unchanged k_{on-P} but increased k_{on-L}, leading to a lower $R_{E/L}$ (2 ± 0.3, n=21 microtubules from 2 assays, $p<0.001$) and suggesting the reduction, but not a complete loss, of the end-binding preference (**Fig. 3 B-C**). Additionally, MCAK·ADP showed a significantly shorter end-binding and lattice-binding dwell time (0.46 ± 0.02 s at the end, n=327 events from 2 assays, $p<0.001$) compared to MCAK·ATP (**Fig. 3 D**). Consequently, MCAK·ADP has a lower end-binding affinity ($K_{d-P}=163$ μM). MCAK·APO, the nucleotide-free state, showed a large increase in k_{on-P} and k_{on-L}, thereby having a lower $R_{E/L}$ (2 ± 0.2, n=36 microtubules from 3 assays, $p<0.001$) (**Fig. 3 B-C**). It also showed a significantly longer end-binding dwell time (1.89 ± 0.23 s, n=71 events from 3 assays, $p<0.001$) (**Fig. 3D**), indicating a lower k_{off} (**Fig. s1**). As a result, MCAK·APO had a higher end-binding affinity ($K_{d-P}=1.8$ μM), but it partially loses the end-binding preference. Overall, these findings indicate that MCAK's end-binding preference and affinity are dependent on its nucleotide state, with the ATP-bound state exhibiting the highest end-binding preference.

MCAK strongly binds to GTPγS microtubules

The binding region of MCAK at growing microtubule ends encompasses the EB cap. This region consists mostly of GDP·Pi-tubulin dimers, which are thought to form at least some lateral contacts and adopt a straighter conformation (Guesdon et al., 2016; Maurer et al., 2012). This raises the question of how MCAK recognizes the EB cap. To understand this, we examined that to what extent, MCAK may adopt the end-binding mechanism utilized by EB1.

We first investigated the contribution of the nucleotide state-dependent feature of the EB cap, which underlies the end-binding preference of EBs (Maurer et al., 2011; Maurer et al., 2012). Specifically, we compared the binding affinities of MCAK·ATP on GTPγS and GDP microtubules, mimicking the tubulin dimers in the EB cap and the lattice of growing microtubules, respectively (**Fig. 4A**). Because MCAK·ATP at the concentration required for full lattice decoration would rapidly disassemble microtubules, we cannot directly compare the total binding of MCAK on microtubules as what was previously done for the EBs (Maurer et al., 2011; Maurer et al., 2012). Instead, we adopted an alternative approach by recording the binding of MCAK·ATP at a low concentration (10 nM) on microtubules and measuring fluorescence intensity summation over a period of time (1000 frames, 0.3s per frame) (**Fig. s3**). Strikingly, quantitative analysis revealed that MCAK·ATP showed a significantly stronger binding

to GTPγS microtubules than to GDP microtubules (**Fig. 4 B-C**). Because the tubulin
dimers in GTPγS microtubules were considered to be the analogue of GDP·Pi-tubulins
in the EB cap (Manka and Moores, 2018; Maurer et al., 2011; Maurer et al., 2014;
Zhang et al., 2015), this result provides a biochemical basis for the MCAK·ATP's
binding preference for the EB cap. Here, we used EB1 as a control (**Fig. s4**).

Given that the growing end-binding affinity and preference of MCAK depend on
the nucleotide state, we wondered whether its preference for GTPγS microtubules
follows a similar trend. To address this, we performed the experiments for
MCAK·AMPPNP, MCAK·ADP and MCAK·APO. The MCAK variants bound to the
lattice   of   microtubules   with   different   affinities
(MCAK·APO>MCAK·AMPPNP>MCAK·ADP > MCAK·ATP) (**Fig. 4B**), consistent
with the results observed on the lattice of dynamic microtubules. Additionally, MCAK
maintained a binding preference for GTPγS microtubules over GDP microtubules
across all nucleotide states, but the degree of binding preference varied with the
nucleotide state (MCAK·AMPPNP > MCAK·ATP ≈ MCAK·APO >MCAK·ADP) (**Fig.
4C**). This finding indicates that MCAK's binding preference for GTPγS microtubules
(i.e., GDP·Pi-tubulin) is strongest at the ATP state, mirroring its preference for growing
microtubule ends.

**MCAK binds to the end region of stabilized GMPCPP microtubules in a
nucleotide state-dependent manner**

Our findings also support the notion that MCAK binds to the distalmost cap,
consistent with the previous reports that MCAK binds to the curved protofilaments of
microtubule ends (Asenjo et al., 2013; Benoit et al., 2018; Desai et al., 1999; Tan et al.,
2008; Wang et al., 2017). Given the nucleotide state-dependence of the growing end-
binding preference, we questioned whether this binding preference for curved
protofilaments also depends on the nucleotide state of MCAK. To investigate this issue,
we analyzed the binding of MCAK at the ends of stabilized GMPCPP microtubules
(**Fig. 5A**), where curved protofilaments are thought to resemble those at the ends of
dynamic microtubules, but without the presence of GDP·Pi-tubulins (Atherton et al.,
2018; Manka and Moores, 2018; McIntosh et al., 2018). Taxol stabilized GDP-
microtubules were here used as a control (**Fig. s5**). We found that MCAK, in both the
ATP and AMPPNP state, exhibited a binding preference for the ends of GMPCPP-
stabilized microtubules (**Fig. 5A-F**). Similarly, MCAK·ATP showed a binding
preference for the ends of taxol-stabilized GDP microtubules, albeit to a lesser extent
(**Fig. s5**). This observation suggests that the ends of stable microtubules mimic the
structure of growing microtubule ends to varying degrees, with the GMPCPP-stabilized
ends providing the closest resemblance. In contrast, MCAK·APO and MCAK·ADP, did
not show the end-binding preference (**Fig. 5A-F**), consistent with the observation that

MCAK·ADP and MCAK·APO also had a lower binding preference for growing
microtubule ends.

The α4 helix and the L2 loop of MCAK contribute to the binding preference to the
EB cap

Intuitively, the binding preference for the EB cap, in addition to the previously
established preference for the curved protofilaments, provides more binding sites for
MCAK at growing microtubule ends. We sought to identify the key domains
responsible for MCAK's preference for the EB cap. Previous studies have shown that
the α4 helix and the L2 loop are critical for mediating MCAK's interaction with curved
protofilaments (Asenjo et al., 2013; Benoit et al., 2018; Patel et al., 2016; Tan et al.,
2008; Wang et al., 2017). Specifically, the α4 helix of MCAK interacts with the
intradimer interface of the tubulin heterodimer, while the L2 loop contacts the
interdimer interface (**Fig. 6A**). Moreover, the electrostatic interactions between the α4
helix and the L2 loop of MCAK with tubulin heterodimer are important for MCAK's
interactions with microtubule ends (Patel et al., 2016; Tan et al., 2008), as
phosphorylation at T537 disrupts MCAK's microtubule-end binding and
depolymerization activity (Sanhaji et al., 2010). We then wondered whether these
domains also contribute to the binding preference for the EB cap. To test this hypothesis,
we introduced two mutations: the K524A mutation in the α4 helix, which is expected
to disrupt the electrostatic interaction between MCAK and the tubulin heterodimer
(Patel et al., 2016; Wang et al., 2017), and the V298S mutation in the L2 loop, which
is predicted to impair the hydrophobic interaction between the L2 loop and α-tubulin
(Wang et al., 2015). Consistent with previously reports (Wang et al., 2017; Wang et al.,
2015), both mutants exhibited deficiencies in depolymerization and reduced ATPase
activity (**Fig. s6**).

Compared to the wild-type MCAK, both MCAK[K524A] and MCAK[V298S] were
capable of binding to growing microtubule ends (**Fig. 6B**), but their $k_{\text{on-P}}$ and dwell time
at growing microtubule ends were significantly reduced (**Fig. 6C**-**D**). This indicates that
the binding affinity of the mutants to growing microtubule ends is substantially
decreased. Here, due to the minimal to negligible number of observed binding events
of the mutants at growing microtubule ends and the lattice, reliable calculations of $R_{\text{E/L}}$
and k_{off} were not feasible for the mutants. Therefore, we directly assayed the binding
preference for GTPγS microtubules mimicking the EB cap. Both mutants showed
significantly reduced binding intensity and a decreased, but not the loss of, preference
for GTPγS microtubules compared to wild-type MCAK (**Fig. 6E-G** vs **Fig. 4B-C**).
These results suggest that both the α4 helix and the L2 loop contributes the binding
preference for the EB cap.

Functional specification of MCAK and XMAP215

MCAK is a microtubule catastrophe factor, whereas XMAP215 functions as a microtubule polymerase (Brouhard et al., 2008; Desai et al., 1999; Gardner et al., 2011). Both are capable of binding to dynamic microtubule ends and regulating microtubule dynamics. In the previous reports, they were shown to antagonistically regulate microtubule assembly (Barr and Gergely, 2008; Kinoshita et al., 2001; Moriwaki and Goshima, 2016; Tournebize et al., 2000). Additionally, structural models have indicated that both MCAK and XMAP215 bind to curved tubulin dimers (Ayaz et al., 2012; Trofimova et al., 2018), raising the possibility of competition for binding sites in the distalmost cap. However, while XMAP215 was reported to promote microtubule catastrophe (Farmer et al., 2021), our findings showed that MCAK binds more proximally relative to XMAP215 at growing microtubule ends. The extent to which competition from XMAP215 would antagonize the function of MCAK remains to be elucidated. Therefore, the collective effect of MCAK and XMAP215 on microtubule dynamics remains an intriguing question.

To investigate this issue, we studied how MCAK and XMAP215 together regulate microtubule grow rate and catastrophe frequency. Based on our pilot experiments, we chose 20 nM for MCAK and 50 nM for XMAP215 as representative concentrations. In the presence of 20 nM MCAK, dynamic microtubules could grow, but their catastrophe frequency was significantly increased, and their lifetime was shortened (**Fig. 7 A-B**). In contrast, 50 nM XMAP215 increased the growth rate more than fourfold (control: 0.8 ± 0.1 μm min^{-1}, n=97 microtubules from 6 assays; 50 nM XMAP215: 3.7 ± 0.3 μm min^{-1}, n=50 microtubules from 3 assays). Notably, XMAP215 alone also caused a mild shortening of microtubule lifetime (**Fig. 7A-B**) (50 nM XMAP215: 264.6 ± 12.9 seconds, n=237 growing events from 3 assays), as compared to that of the control group (323.4 ± 12.6 seconds of control, n=443 growing events from 6 assays; p>1, the two-tailed Mann-Whitney U test with Bonferroni correction). This is consistent with previous reports suggesting that XMAP215 enhances fluctuations in microtubule end structure (Farmer et al., 2021). When both 20 nM MCAK and 50 nM XMAP215 were added, the microtubule growth rate was comparable to that observed with XMAP215 alone (3.9 ± 0.3 μm min^{-1}, n=52 microtubules from 3 assays; p=0.5, the two-tailed Mann-Whitney U test with Bonferroni correction), suggesting that MCAK does not significantly affect XMAP215's function as a polymerase. However, the lifetime of dynamic microtubules (170.3 ± 7.5 seconds, n=283 growing events from 3 assays) was shorter than that observed with XMAP215 alone ($p=2 \times 10^{-6}$, the two-tailed Mann-Whitney U test with Bonferroni correction), while it was similar to that observed with MCAK alone (196.7 ± 6.6 seconds, n=621 growing events from 6 assays; p>1, the two-tailed Mann-Whitney U test with Bonferroni correction) (**Fig. 7B**). To determine whether this reflects the simple summation of the individual contributions of MCAK and XMAP215 or a synergistic effect, we calculated the probability-based catastrophe

frequency for all conditions (**Fig. 7C**) (Gardner et al., 2011). We found that the effects
of MCAK and XMAP215 on the probability of microtubule catastrophe were additive,
indicating that there was no synergistic effect on catastrophe frequency (**Fig. 7C**). This
suggests that, at least at this concentration combination, MCAK and XMAP215
independently regulate different parameters of microtubule dynamics, namely
catastrophe frequency and growth rate, while simultaneously interacting with
microtubule ends.

Discussion

This study provides mechanistic insights into the end-binding mechanism of
MCAK. The previous established view on the end-binding mechanism of MCAK is
based on its specific recognition of curved protofilaments at the distalmost end of
growing microtubules (Asenjo et al., 2013; Benoit et al., 2018; Tan et al., 2008). The
findings presented here expand on this model by showing that MCAK also binds to the
EB cap, a binding preference that likely facilitates both end-binding and the subsequent
depolymerizing activity of MCAK. Below, we discuss our key findings and their
implications.

MCAK binds to the EB cap at growing microtubule ends

Our primary discovery is that MCAK preferentially binds to the EB cap, where
the GDP·Pi-tubulins accumulate. This conclusion is based on four main lines of
evidence. First, localization analysis demonstrates that MCAK binds to the entire GTP
cap, including both the EB cap and the distalmost cap. Second, MCAK·ADP and
MCAK·APO show the significantly reduced, but not abolished, binding preference for
growing microtubule ends, while their binding preference for curved protofilaments is
absent. These different binding behaviors suggest that additional structural features,
beyond curved protofilaments, contribute to MCAK's growing end-binding preference.
Third, MCAK preferentially binds to GTPγS microtubules over GDP microtubules.
Since the tubulins in GTPγS microtubules are considered to mimic, at least, some of
the key features of the GDP·Pi-tubulins in the EB cap (Maurer et al., 2011; Maurer et
al., 2012), this suggests that MCAK can recognize the molecular features of tubulin
dimers within the EB cap. Fourth, previous cryo-EM studies have shown that GTP
hydrolysis leads to longitudinal compression of tubulin dimers, increasing the contact
area at the interdimer interface (LaFrance et al., 2022; Manka and Moores, 2018; Zhang
et al., 2015). Therefore, the interface in GTPγS microtubules is characteristic, being
larger than that of a straight GMPCPP-tubulin lattice but smaller than that of a straight
GDP-tubulin lattice. We find that the L2 loop contributes to the binding-preference to
the EB cap, and it was previously reported that the L2 loop of MCAK directly contacts
the interdimer interface (**Fig. 6A**) (Trofimova et al., 2018; Wang et al., 2017),
potentially providing a structural basis for MCAK to distinguish the nucleotide-

dependent features of tubulin dimers.

The end-binding affinity of MCAK

In our experiments, we observed that MCAK exhibits a relatively low end-binding
affinity (69 μM), raising the question of its physiological relevance. Several factors
may contribute to this observation. First, the in vitro system used may not fully
recapitulate physiological conditions. For instance, lowering the salt concentration
significantly increases binding affinity (Helenius et al., 2006; Maurer et al., 2011;
McHugh et al., 2019). Additionally, while numerous binding events with short
durations were detected, we excluded transient interactions from our analysis to
facilitate quantification. This likely led to an underestimation of the on-rate and,
consequently, the binding affinity. Furthermore, previous studies have shown that
retaining a His-tag markedly enhances microtubule binding (Maurer et al., 2011; Zhu
et al., 2009). To minimize interference from purification tags, we ensured complete
removal of the His-tag during protein preparation (**Fig. s1**), which may explain the
lower apparent binding affinity in our study compared to others. Finally, a low affinity
is not necessarily unexpected. Considering the microtubule end as a receptor with
multiple binding sites for MCAK, the overall binding affinity is in the nanomolar range
(260 nM). Indeed, only a few MCAK molecules may be sufficient to induce
microtubule catastrophe.

A natural follow-up question is the physiological relevance of MCAK's direct
binding to microtubule ends compared to the well-established EB1-mediated pathway
(Gouveia et al., 2010; Honnappa et al., 2009; Lee et al., 2008; McHugh and Welburn,
2023). The relative contributions of these two pathways depend on MCAK's end-
binding affinity. To estimate their proportional contributions, we employed a simple
model (**Supplementary Appendix-2** and **Fig. s7**). The modeling results suggest that
MCAK's direct end-binding affinity contribute through two distinct mechanisms
(**Supplementary Appendix-2**). First, MCAK establishes an EB1-independent
regulatory pathway. Simultaneously, it enhances the EB1-dependent pathway by
increasing the microtubule-end affinity of the MCAK-EB1 complex. Notably, these two
mechanisms are not mutually exclusive. Particularly intriguing is that the direct
regulatory pathway may account for previously observed experimental phenomena. For
example, prior studies have shown that both MCAK and EB1 accumulate at the
centromere and centrosome during cell division (Bahmanyar et al., 2009; Fodde et al.,
2001; Kline-Smith et al., 2004; Maney et al., 1998; Wordeman and Mitchison, 1995).
However, EB1 neither influences MCAK's localization at these sites nor significantly
modulates its local functions (Domnitz et al., 2012).

Implications for the working mechanism of MCAK

443   We think that our findings add to the current working model of MCAK in several

ways. First, the end-binding preference of MCAK for growing microtubule ends is nucleotide state-dependent. The present study demonstrates that MCAK exhibits the strongest end-binding preference in the ATP-bound state, suggesting that MCAK·ATP could more effectively distinguish between the end and the lattice. This supports the view that MCAK acts at the growing microtubule ends in the ATP-bound state (Asenjo et al., 2013; Benoit et al., 2018; Desai et al., 1999; Tan et al., 2008; Wang et al., 2015).

Second, the binding preference of MCAK to the GDP·Pi-tubulins provides a mechanism for MCAK remaining bound to the GTP cap after ATP hydrolysis. In a speculative scenario (**Fig. 8A**), after the terminal tubulin dimers are removed, the MCAK motor domain transitions to the ADP state and loses its affinity for curved protofilaments. However, it still preferentially binds to GDP·Pi-tubulins, allowing MCAK to stay within the GTP cap. As ADP is replaced by a new ATP, the motor domain could locally diffuse to explore a new working site for the next round of catalysis, which may facilitate MCAK's processivity (Cooper et al., 2010; Friel and Howard, 2011; Helenius et al., 2006; Hunter et al., 2003).

Third, we found that XMAP215 and MCAK independently regulate the growth rate and catastrophe frequency of microtubules, respectively (**Fig. 7** and **8B**). Previous studies have suggested that XMAP215 and MCAK exert antagonistic effects on microtubule assembly (Barr and Gergely, 2008; Kinoshita et al., 2001; Moriwaki and Goshima, 2016; Tournebize et al., 2000), possibly through competition for binding sites. Structural studies have indicated overlap in the binding sites of MCAK and XMAP215 on tubulin dimers (Ayaz et al., 2012; Trofimova et al., 2018), lending support to this hypothesis. However, in the present study, we observed that the binding regions of XMAP215 and MCAK at growing microtubule ends are not entirely overlapping, with MCAK predominantly binding proximally to the binding region of XMAP215. These findings suggest an alternative mechanism for the combined effects of XMAP215 and MCAK. In this model, MCAK and XMAP215 independently regulate catastrophe frequency and growth rate, respectively, modulating microtubule dynamics through additive effects, rather than through direct competition for binding sites.

Materials and Methods

Protein expression and purification

MCAK purification.

GFP-MCAK, its mutants and MCAK-RFP (construct schematics shown in **Fig. s1**) were expressed using a modified BAC-to-BAC protocol. Purification was performed with adaptations based on previously described methods (Helenius et al., 2006). Briefly, the Sf9 cells were lysed by Dounce homogenization with ice-cold lysis buffer. The lysate was centrifuged at 40,000 rpm for 60 min at 4°C (XPN-100 ultracentrifuge, Type 45 Ti rotor, Beckman, USA). Cleared supernatant was filtered through a 0.45 μm filter (Pall, 66229) and then purified using a cation-exchange column (HiTrap SP-HP, Cat.17115101, GE, USA). The protein was eluted using the elution buffer supplemented with a continuous salt gradient (0.15~1.0 M NaCl). Peak fractions were pooled together and then subjected to a Ni-sepharose column (QIAGEN, 30210). The column was then washed with the washing buffer I. After the washing step, 3C Protease (final concentration 10 μg/ml) (Thermo Fisher Scientific, 88946) was employed to perform on-column his-tag cleavage overnight at 4°C. The successful removal of His-tag was subsequently confirmed using Western blot analysis (**Fig. s1**). Cleaved protein was further purified through size-exclusion chromatography (Superdex 200 increase 10/300 GL column, Cat. 28990944, Cytiva, USA) and eluted with the washing buffer II. Purified MCAK aliquots were frozen in liquid N_2 and stored at -80°C.

Lysis buffer: 50 mM HEPES pH 7.5, 150 mM NaCl, 5% glycerol, 0.1% Tween 20, 1.5 mM $MgCl_2$, 3 mM EGTA, 1 mM DTT, 0.5 mM Mg-ATP, 10 units/ml Benzonase. Elution buffer: 6.7 mM HEPES pH 7.5, 6.7 mM MES, 6.7 mM sodium acetate, 1.5 mM $MgCl_2$, 10 μM Mg-ATP. Washing buffer I: 50 mM $NaPO_4$ buffer pH 7.5, 300 mM NaCl, 10 mM imidazole, 10% glycerol, 1 mM $MgCl_2$, 10 μM Mg-ATP. Washing buffer II: BRB80, 300 mM KCl, 1 mM DTT and 10 μM Mg-ATP. BRB80: 80 mM PIPES/KOH pH 6.9, 1 mM $MgCl_2$, 1 mM EGTA.

EB1 purification.

EB1-GFP (construct schematic shown in **Fig. s1**) was expressed and purified as previously described (Song et al., 2020). Briefly, EB1-GFP was expressed in the *BL21 E. coli* strain and lysed by sonication with ice-cold lysis buffer. The lysate was centrifuged at 40,000 rpm for 60 min at 4°C. Cleared supernatant was filtered through a 0.45 μm filter and loaded onto a Ni-sepharose column. The column was then washed using the washing buffer. After the washing step, 3C Protease (final concentration 10 μg/ml) was employed to perform on-column his-tag cleavage overnight at 4°C. Eluates were desalted into storage buffer using the PD-10 desalting column (GE, 17085101). The final purified protein aliquots were frozen in liquid N_2 and stored at -80 °C.

Lysis buffer: 50 mM $NaPO_4$ buffer pH 7.5, 300 mM NaCl, 10% glycerol, 10 mM imidazole, 1 mM DTT, 0.1% Tween 20. Washing buffer: 50 mM $NaPO_4$ buffer pH 7.5, 300 mM NaCl, 10% glycerol, 30 mM imidazole, 1 mM DTT. Storage buffer: BRB80, 100 mM KCl, 10% glycerol and 1 mM DTT.

XMAP215 purification.

XMAP215-GFP-his$_6$ (construct schematic shown in **Fig. s1**) was expressed using a modified BAC-to-BAC protocol. The purification was performed as previously described (Brouhard et al., 2008). Briefly, the cells were lysed by Dounce homogenization with ice-cold lysis buffer. The lysate was centrifuged at 40,000 rpm for 60 min at 4°C. Supernatant was filtered through a 0.45 μm filter and then purified using a cation-exchange column. The protein was eluted using the elution buffer supplemented with a continuous salt gradient (0.15~1.0 M NaCl). Peak fractions were pooled together and then loaded onto a Ni-sepharose column. The column was then washed with the washing buffer I. The protein was eluted using the washing buffer I supplemented with a continuous imidazole gradient (20-300 mM). Eluted fractions with best purity were pooled together and further purified was further purified through size-exclusion chromatography (Superdex 200 increase 10/300 GL column, Cat. 28990944, Cytiva, USA) and eluted using the washing buffer II. The final purified XMAP215 was frozen in liquid N$_2$ and stored at -80°C.

Lysis buffer: 50 mM HEPES pH 7.5, 50 mM NaCl, 5% glycerol, 0.1% Tween 20, 1 mM DTT, 10 units/ml Benzonase. Elution buffer: 6.7 mM HEPES pH 7.5, 6.7 mM MES, 6.7 mM sodium acetate. Washing buffer I: 50 mM NaPO4 buffer pH 7.5, 300 mM NaCl, 10 mM imidazole, 10% glycerol, 1 mM MgCl$_2$, 10 μM Mg-ATP. Washing buffer II: BRB80, 150 mM KCl, 1 mM DTT.

Tubulin preparation.

Tubulin was purified from porcine brain tissues through two cycles of temperature-dependent polymerization and depolymerization followed by affinity purification using the TOG-based affinity column, as previously described (Gell et al., 2010; Widlund et al., 2012). Purified tubulin was labeled with biotin (Thermo Fisher Scientific, 20217), TAMRA (Thermo Fisher Scientific, C1171) and Alexa Fluor 647 (Thermo Fisher Scientific, A20106) using the NHS esters according to the standard protocols (Gell et al., 2010).

Size-exclusion chromatography

To confirm the oligomeric state of the purified proteins, size-exclusion chromatography analysis (Superose™ 6 Increase 5/150 GL column, Cat. 29091597, cytiva, USA) was performed using BRB80 supplemented with 50 mM KCl and 1 mM ATP.

Microtubule depolymerization assay

Microtubules (5% Alexa Fluor 647 labeled and 20% biotin labeled) were polymerized in the presence of GMPCPP (JenaBioscience, NU-405L) as previously described (Song et al., 2020). The GMPCPP-stabilized microtubules were immobilized on the surface of a cover glass using the biotin-NeutrAvidin protein links (Thermo Fisher Scientific, 31000). GFP-MCAK was then added into the flow cell in the imaging buffer. The sample was kept at 35°C using a temperature controller (Tokai Hit, Japan).

Images were recorded using a total internal reflection fluorescence microscope (TIRFM) (Olympus, Japan) equipped with a 100× 1.49 N.A. oil TIRF objective (Olympus, Japan) and an Andor 897 Ultra EMCCD camera (Andor, Belfast, UK). Images were recorded every 5 s with a 100 ms exposure. Imaging buffer: BRB80 supplemented with 1 mM ATP, 50 mM KCl, 80 mM D-glucose, 0.4 mg/ml glucose oxidase, 0.2 mg/ml catalase, 0.8 mg/ml casein, 1% β-mercaptoethanol, 0.001% Tween 20.

Microtubule dynamics assay

Microtubule dynamics assay was performed as previously described (Song et al., 2020). Briefly, the immobilized GMPCPP-stabilized microtubules were used as the template for microtubule growth. Tubulin dimers (13% TAMRA labeled) and the protein of interest were then added into the flow cell in the imaging buffer. The sample was kept at 35°C using a temperature controller (Tokai Hit, Japan). Images were recorded using a TIRFM (Olympus, Japan). To record microtubule dynamics, images were recorded every 5 s with a 100 ms exposure. To record single molecule binding events, images were recorded every 100 ms with a 50 ms exposure. Imaging buffer: BRB80 supplemented with 1 mM ATP, 2 mM GTP, 50 mM KCl, 0.15% sodium carboxymethylcellulose, 80 mM D-glucose, 0.4 mg/ml glucose oxidase, 0.2 mg/ml catalase, 0.8 mg/ml casein, 1% β-mercaptoethanol, 0.001% Tween 20.

Screen for single molecules

The average fluorescence intensity over the dwell time of each binding event was used to represent the fluorescence signal of this binding event. Subsequently, a probability distribution of the fluorescence intensities of all binding events was constructed and fitted by Gaussian fitting. A binding event was deemed to represent a single molecule if its intensity fell within the range ($\mu \pm 2\sigma$) of the probability distribution. In addition, only binding events with a dwell time exceeding the duration corresponding to 2 pixels (~200 ms) were included in the statistical analysis. As a control for the dimeric GFP-MCAK, we used the fluorescence intensity probability distribution of GFP-MCAK^{sN+M}, a monomeric variant of MCAK (Wang et al., 2012) (**Fig. s1**).

The diffusion coefficient of MCAK at the lattice of dynamic microtubules

The diffusion coefficient of MCAK on dynamic microtubule lattices was calculated as previously described (Helenius et al., 2006). First, we tracked MCAK's position along its trajectory on dynamic microtubule lattices using the TrackMate plugin in ImageJ (Fiji) (Tinevez et al., 2017), maintaining a minimum track length of 400 ms (100 ms/frame). Next, the mean-squared displacement (MSD) was computed from 203 trajectories across three assays using MATLAB. Finally, MSD values were plotted against the time interval, and the diffusion coefficient (D) was determined by linear regression analysis ($<x^2> = 2Dt$) using Origin 8.0 (OriginLab Corporation, USA).

The flux of MCAK reaching the growing microtubule ends

594 The of the flux of MCAK reaching growing microtubule ends was calculated by a one-
595 dimensional lattice-diffusion model, based on Fick's first equation of diffusion
596 (Helenius et al., 2006):

$$J = -D \cdot \partial c / \partial x \quad \text{Eq.1}$$

598 where J represents the flux of MCAK reaching growing microtubule ends, $c(x, t)$ is
599 the concentration of MCAK on microtubule lattice at position x from the growing
600 microtubule end and at time t.

$$\text{When } x = 0 \quad J_0 = -D \cdot c_\infty / x_0 \quad \text{Eq. 2}$$

602 where

$$x_0 = (D/k_{\text{off}})^{1/2} \quad \text{Eq. 3}$$

$$c(x \to \infty) = c_\infty = k_{\text{on}} \cdot C / k_{\text{off}} \quad \text{Eq. 4}$$

605 where C is the concentration of MCAK. Therefore, the flux was calculated by taking
606 the measured parameters (D, k_{on}, C and k_{off}) into the equations.

Localization analysis at growing microtubule ends

The single-molecule localization at growing microtubule ends was conducted as
previously described (Maurer et al., 2014; Song et al., 2020). Briefly, Fluorescence
calibration of the captured movie was performed using fluorescent beads (T7279,
Thermo Fisher Scientific). During the dwell time of a single-molecule binding event,
the microtubule lattice was considered as a Gaussian wall and its end as a half-Gaussian
in every frame. The peak position of the half-Gaussian was used as a reference to
measure the intensity profile of single-molecule binding events frame by frame. A
Gaussian function was employed to fit the intensity profile of individual molecules. The
position at the peak of the intensity profile was taken as the single molecule's location
in every frame (**Fig. s2**). Subsequently, the average of these positions was taken to
represent the single-molecule binding event's position during its dwell time (**Fig. s2**).
This procedure was iterated for all single-molecule binding events to determine their
positions during their respective dwell times. Then, the spatial positional distributions
of GFP-MCAK and XMAP215-GFP were fitted by Gaussian fitting and the spatial
positional distribution of EB1-GFP was fitted to an exponentially modified Gaussian
function, from which we calculated the FWHM as an estimation for their length of the
binding regions (**Fig. 2C**).

Location precision analysis

626 The observed image results from the convolution of the fluorescent object with the
627 system's point spread function (PSF). Consequently, the standard deviation of the
628 observed image reflects a combination of measurement error (standard deviation) in
629 determining the object's position and the width of the PSF (Eq. 5) (Bohner et al., 2016;
630 Demchouk et al., 2011). To characterize the PSF of the imaging system, we used

multicolor fluorescent beads (T7279, ThermoFisher Scientific) with a 100 nm diameter (Hirvonen et al., 2009). When measuring the position of a microtubule end in a single image, we modeled the microtubule lattice as a Gaussian wall and its end as a half-Gaussian. Using this fitting approach, we estimated the measurement error (standard deviation) of the convolved microtubule end positions and deconvolved the system's PSF contribution using Eq. 5. The mean standard deviation of end positions over the dwell time of a binding event was then used as the measurement error for the microtubule end position. This procedure was iterated for all microtubule ends with single-molecule binding events to construct their measurement errors. Then, the probability distribution of measurement errors was fitted by Gaussian fitting. The standard deviation of the probability distribution was taken as the precision in estimating microtubule end positions (~1.3 nm). The same calculation method was applied to determine the positional precision of other single molecules (~18 nm).

$$\sigma_I^2 = \sigma_O^2 + \sigma_{PSF}^2 \quad \text{Eq. 5}$$

Microtubule polymerization

GMPCPP, GDP and GTPγS microtubules were polymerized as previously described (Manka and Moores, 2018). Briefly, GMPCPP-microtubules were polymerized using a mixture of 10 μM tubulin, 1 mM GMPCPP (JenaBioscience, NU-405L) and 4 mM MgCl$_2$ in BRB80, which was incubated for 2 hours at 37˚C. The polymerized microtubules were then pelleted using an Air-Driven Ultracentrifuge (Beckman, 340401), resuspended in BRB80 and stored at 37˚C. GDP-microtubules were polymerized using a mixture of 40 μM tubulin, 1 mM GTP (Roche, 10106399001), 4 mM MgCl$_2$ and 4% DMSO (Sigma, 276855). The mixture was incubated for 30 min at 37˚C. The polymerized microtubules were collected using an Air-Driven Ultracentrifuge, resuspended in BRB80 with 20 μM taxol (Cell Signaling Technology, 9807) and stored at 37˚C. GTPγS microtubules were prepared using two rounds of polymerizations. In the first round, a mixture of 40 μM tubulin, 2 mM GTPγS (Roche, 10220647001) and 4 mM MgCl$_2$ in BRB80 was incubated for 5 hours at 37°C. In the second round, 5 μl of the product from the first polymerization was mixed with additional 40 μM tubulin, 2 mM GTPγS and 4 mM MgCl$_2$ to create a final 25 μl reaction mixture. This mixture was incubated overnight at 37°C to obtain long GTPγS microtubules. The polymerized microtubules were collected using an Air-Driven Ultracentrifuge, resuspended in BRB80 with 20 μM taxol and stored at 37˚C.

Microtubule-Stimulated ATPase Assay

The microtubule stimulated ATPase activities of MCAK and its mutants were performed by using the Kinesin ATPase End-Point Biochem Kit (Cytoskeleton, Inc., BK053). The measurements were based on the malachite green phosphate assay to probe inorganic phosphate generated during the reaction. Wild-type MCAK served as the control, and each type of variant was subjected to multiple replicate measurements.

Binding of MCAK to different microtubules

The microtubules in different nucleotide states were first immobilized in flow cells.
GFP-MCAK or its mutants were then added into the flow cell in the imaging buffer.
Images were recorded every 300 ms with a 100 ms exposure. Imaging buffer: BRB80
supplemented with 1 mM ATP, ADP or AMPPNP, 20 μM taxol, 50 mM KCl, 80 mM
D-glucose, 0.4 mg/ml glucose oxidase, 0.2 mg/ml catalase, 0.8 mg/ml casein, 1% β-
mercaptoethanol, 0.001% Tween 20.

678 The binding intensity of GFP-MCAK at GMPCPP microtubule ends

To measure the binding intensity of GFP-MCAK at GMPCPP microtubule ends,
an intensity profile of GFP-MCAK or the GMPCPP microtubule was plotted along the
microtubule. The average binding intensity of GFP-MCAK of three pixels represents
the end-binding intensity of GFP-MCAK at the corresponding position of the end, to
the lattice from the midpoint of the microtubule intensity attenuation at the microtubule
end.

685 Gamma fitting of microtubule lifetimes and catastrophe frequency analysis

686   The lifetime data of dynamic microtubules were fitted to the gamma density
687 function using the dfittool toolbox in MATLAB (Mathworks, USA). The probability-
688 based catastrophe frequency was calculated as the ratio of the number of catastrophe
689 events observed at a certain lifetime to the total number of microtubules that reached
690 this lifetime (Gardner et al., 2011).

Acknowledgement

The authors thank Pengpeng Yu (School of Life Sciences and technology, TongJi University), Chunguang Wang (School of Life Sciences and technology, TongJi University), Peng Shi (School of Basic Medical Sciences, Peking University) and Congying Wu (School of Basic Medical Sciences, Peking University) for technical assistance. Special thanks to the light microscopy and protein facility in Tsinghua University. We also acknowledge our funding from National Natural Sciences Foundation of China (32070704, 32370730), IDG/McGovern Institute for Brain Research (Tsinghua University) and the State Key Laboratory of Complex, Severe, and Rare Diseases.

References

Aizawa, H., Y. Sekine, R. Takemura, Z.Z. Zhang, M. Nangaku, and N. Hirokawa. 1992. KINESIN FAMILY IN MURINE CENTRAL-NERVOUS-SYSTEM. *Journal of Cell Biology*. 119:1287-1296.

Alberts, B. 2015. Molecular biology of the cell.

Asenjo, A.B., C. Chatterjee, D. Tan, V. DePaoli, W.J. Rice, R. Diaz-Avalos, M. Silvestry, and H. Sosa. 2013. Structural Model for Tubulin Recognition and Deformation by Kinesin-13 Microtubule Depolymerases. *Cell Reports*. 3:759-768.

Atherton, J., M. Stouffer, F. Francis, and C.A. Moores. 2018. Microtubule architecture and in cells revealed by cryo-electron tomography. *Acta Crystallogr D*. 74:572-584.

Ayaz, P., X. Ye, P. Huddleston, C.A. Brautigam, and L.M. Rice. 2012. A TOG:alpha beta-tubulin Complex Structure Reveals Conformation-Based Mechanisms for a Microtubule Polymerase. *Science*. 337:857-860.

Bahmanyar, S., W.J. Nelson, and A.I.M. Barth. 2009. Role of APC and Its Binding Partners in Regulating Microtubules in Mitosis. *Adv Exp Med Biol*. 656:65-74.

Barr, A.R., and F. Gergely. 2008. MCAK-Independent Functions of ch-Tog/XMAP215 in Microtubule Plus-End Dynamics. *Molecular and Cellular Biology*. 28:7199-7211.

Benoit, M.P.M.H., A.B. Asenjo, and H. Sosa. 2018. Cryo-EM reveals the structural basis of microtubule depolymerization by kinesin-13s. *Nature Communications*. 9.

Bohner, G., N. Gustafsson, N.I. Cade, S.P. Maurer, L.D. Griffin, and T. Surrey. 2016. Important factors determining the nanoscale tracking precision of dynamic microtubule ends. *Journal of Microscopy*. 261:67-78.

Brouhard, G.J., J.H. Stear, T.L. Noetzel, J. Al-Bassam, K. Kinoshita, S.C. Harrison, J. Howard, and A.A. Hyman. 2008. XMAP215 is a processive microtubule polymerase. *Cell*. 132:79-88.

Cooper, J.R., M. Wagenbach, C.L. Asbury, and L. Wordeman. 2010. Catalysis of the microtubule on-rate is the major parameter regulating the depolymerase activity of MCAK. *Nature Structural & Molecular Biology*. 17:77-U98.

Demchouk, A.O., M.K. Gardner, and D.J. Odde. 2011. Microtubule Tip Tracking and Tip Structures at the Nanometer Scale Using Digital Fluorescence Microscopy. *Cellular and Molecular Bioengineering*. 4:192-204.

Desai, A., S. Verma, T.J. Mitchison, and C.E. Walczak. 1999. Kin I kinesins are microtubule-destabilizing enzymes. *Cell*. 96:69-78.

Domnitz, S.B., M. Wagenbach, J. Decarreau, and L. Wordeman. 2012. MCAK activity at microtubule tips regulates spindle microtubule length to promote robust kinetochore attachment. *Journal of Cell Biology*. 197:231-237.

Farmer, V., G. Arpa, S.L. Hall, and M. Zanic. 2021. XMAP215 promotes microtubule catastrophe by disrupting the growing microtubule end. *Journal of Cell Biology*. 220.

Fodde, R., J. Kuipers, C. Rosenberg, R. Smits, M. Kielman, C. Gaspar, J.H. van Es, C. Bruekel, J. Wiegant, R.H. Giles, and H. Clevers. 2001. Mutations in the tumour suppressor gene cause chromosomal instability. *Nature Cell Biology*. 3:433-438.

Friel, C.T., and J. Howard. 2011. The kinesin-13 MCAK has an unconventional ATPase cycle adapted for microtubule depolymerization. *Embo Journal*. 30:3928-3939.

Gardner, M.K., M. Zanic, C. Gell, V. Bormuth, and J. Howard. 2011. Depolymerizing Kinesins Kip3 and MCAK Shape Cellular Microtubule Architecture by Differential Control of

Catastrophe. *Cell*. 147:1092-1103.

Gell, C., V. Bormuth, G.J. Brouhard, D.N. Cohen, S. Diez, C.T. Friel, J. Helenius, B. Nitzsche, H.
Petzold, J. Ribbe, E. Schaeffer, J.H. Stear, A. Trushko, V. Varga, P.O. Widlund, M. Zanic,
and J. Howard. 2010. Microtubule Dynamics Reconstituted In Vitro and Imaged by Single-
Molecule Fluorescence Microscopy. *In* Microtubules, in Vitro: Microtubules, in Vitro. Vol.
95. L. Wilson and J.J. Correia, editors. 221-245.

Ghosh-Roy, A., A. Goncharov, Y. Jin, and A.D. Chisholm. 2012. Kinesin-13 and Tubulin
Posttranslational Modifications Regulate Microtubule Growth in Axon Regeneration. *Dev.*
Cell. 23:716-728.

Gouveia, S.M., K. Leslie, L.C. Kapitein, R.M. Buey, I. Grigoriev, M. Wagenbach, I. Smal, E.
Meijering, C.C. Hoogenraad, L. Wordeman, M.O. Steinmetz, and A. Akhmanova. 2010. In
Vitro Reconstitution of the Functional Interplay between MCAK and EB3 at Microtubule
Plus Ends. *Curr. Biol.* 20:1717-1722.

Guesdon, A., F. Bazilel, R.M. Buey, R. Mohan, S. Monier, R.R. Garcia, M. Angevin, C. Heichette,
R. Wieneke, R. Tampe, L. Duchesne, A. Akhmanova, M.O. Steinmetz, and D. Chretien.
2016. EB1 interacts with outwardly curved and straight regions of the microtubule lattice.
Nature Cell Biology. 18:1102-1108.

Helenius, J., G. Brouhard, Y. Kalaidzidis, S. Diez, and J. Howard. 2006. The depolymerizing kinesin
MCAK uses lattice diffusion to rapidly target microtubule ends. *Nature*. 441:115-119.

Honnappa, S., S.M. Gouveia, A. Weisbrich, F.F. Damberger, N.S. Bhavesh, H. Jawhari, I. Grigoriev,
F.J.A. van Rijssel, R.M. Buey, A. Lawera, I. Jelesarov, F.K. Winkler, K. Wuethrich, A.
Akhmanova, and M.O. Steinmetz. 2009. An EB1-Binding Motif Acts as a Microtubule Tip
Localization Signal. *Cell*. 138:366-376.

Hunter, A.W., M. Caplow, D.L. Coy, W.O. Hancock, S. Diez, L. Wordeman, and J. Howard. 2003.
The kinesin-related protein MCAK is a microtubule depolymerase that forms an ATP-
hydrolyzing complex at microtubule ends. *Molecular Cell*. 11:445-457.

Janke, C., and M.M. Magiera. 2020. The tubulin code and its role in controlling microtubule
properties and functions. *Nature Reviews Molecular Cell Biology* 21:307-326.

Kapitein, L.C., and C.C. Hoogenraad. 2015. Building the Neuronal Microtubule Cytoskeleton.
Neuron. 87:492-506.

Kinoshita, K., I. Arnal, A. Desai, D.N. Drechsel, and A.A. Hyman. 2001. Reconstitution of
physiological microtubule dynamics using purified components. *Science*. 294:1340-1343.

Kline-Smith, S.L., A. Khodjakov, P. Hergert, and C.E. Walczak. 2004. Depletion of centromeric
MCAK leads to chromosome congression and segregation defects due to improper
kinetochore attachments. *Molecular Biology of the Cell*. 15:1146-1159.

Kline-Smith, S.L., and C.E. Walczak. 2002. The microtubule-destabilizing kinesin XKCM1
regulates microtubule dynamic instability in cells. *Mol Biol Cell*. 13:2718-2731.

LaFrance, B.J., J. Roostalu, G. Henkin, B.J. Greber, R. Zhang, D. Normanno, C.O. McCollum, T.
Surrey, and E. Nogales. 2022. Structural transitions in the GTP cap visualized by cryo-
electron microscopy of catalytically inactive microtubules. *Proceedings of the National*
Academy of Sciences of the United States of America. 119.

Lee, T., K.J. Langford, J.M. Askham, A. Bruning-Richardson, and E.E. Morrison. 2008. MCAK
associates with EB1. *Oncogene*. 27:2494-2500.

Maney, T., A.W. Hunter, M. Wagenbach, and L. Wordeman. 1998. Mitotic centromere-associated

791  kinesin is important for anaphase chromosome segregation. *Journal of Cell Biology*.

792  142:787-801.

793 Manka, S.W., and C.A. Moores. 2018. The role of tubulin-tubulin lattice contacts in the mechanism

794  of microtubule dynamic instability. *Nature Structural & Molecular Biology*. 25:607-+.

795 Maurer, S.P., P. Bieling, J. Cope, A. Hoenger, and T. Surrey. 2011. GTP gamma S microtubules

796  mimic the growing microtubule end structure recognized by end-binding proteins (EBs).

797  *Proceedings of the National Academy of Sciences of the United States of America*.

798  108:3988-3993.

799 Maurer, S.P., N.I. Cade, G. Bohner, N. Gustafsson, E. Boutant, and T. Surrey. 2014. EB1 Accelerates

800  Two Conformational Transitions Important for Microtubule Maturation and Dynamics.

801  *Curr. Biol.* 24:372-384.

802 Maurer, S.P., F.J. Fourniol, G. Bohner, C.A. Moores, and T. Surrey. 2012. EBs Recognize a

803  Nucleotide-Dependent Structural Cap at Growing Microtubule Ends. *Cell*. 149:371-382.

804 McHugh, T., and J.P.I. Welburn. 2023. Potent microtubule-depolymerizing activity of a mitotic

805  Kif18b-MCAK-EB network. *Journal of Cell Science*. 136.

806 McHugh, T., J. Zou, V.A. Volkov, A. Bertin, S.K. Talapatra, J. Rappsilber, M. Dogterom, and J.P.I.

807  Welburn. 2019. The depolymerase activity of MCAK shows a graded response to Aurora

808  B kinase phosphorylation through allosteric regulation. *Journal of Cell Science*. 132.

809 McIntosh, J.R., E. O'Toole, G. Morgan, J. Austin, E. Ulyanov, F. Ataullakhanov, and N. Gudimchuk.

810  2018. Microtubules grow by the addition of bent guanosine triphosphate tubulin to the tips

811  of curved protofilaments. *Journal of Cell Biology*. 217:2691-2708.

812 Moores, C.A., and R.A. Milligan. 2006. Lucky 13 - microtubule depolymerisation by kinesin-13

813  motors. *Journal of Cell Science*. 119:3905-3913.

814 Moriwaki, T., and G. Goshima. 2016. Five factors can reconstitute all three phases of microtubule

815  polymerization dynamics. *Journal of Cell Biology*. 215:357-368.

816 Noda, Y., R. Satoyoshitake, S. Kondo, M. Nangaku, and N. Hirokawa. 1995. KIF2 IS A NEW

817  MICROTUBULE-BASED ANTEROGRADE MOTOR THAT TRANSPORTS

818  MEMBRANOUS ORGANELLES DISTINCT FROM THOSE CARRIED BY KINESIN

819  HEAVY-CHAIN OR KIF3A/B. *Journal of Cell Biology*. 129:157-167.

820 Patel, J.T., H.R. Belsham, A.J. Rathbone, B. Wickstead, C. Gell, and C.T. Friel. 2016. The family-

821  specific alpha 4-helix of the kinesin-13, MCAK, is critical to microtubule end recognition.

822  *Open Biology*. 6.

823 Piao, T., M. Luo, L. Wang, Y. Guo, D. Li, P. Li, W.J. Snell, and J. Pan. 2009. A microtubule

824  depolymerizing kinesin functions during both flagellar disassembly and flagellar assembly

825  in Chlamydomonas. *Proceedings of the National Academy of Sciences of the United States

826  of America*. 106:4713-4718.

827 Puri, D., K. Ponniah, K. Biswas, A. Basu, S. Dey, E.A. Lundquist, and A. Ghosh-Roy. 2021. Wnt

828  signaling establishes the microtubule polarity in neurons through regulation of Kinesin-13.

829  *Journal of Cell Biology*. 220.

830 Sanhaji, M., C.T. Friel, N.-N. Kreis, A. Kraemer, C. Martin, J. Howard, K. Strebhardt, and J. Yuan.

831  2010. Functional and Spatial Regulation of Mitotic Centromere-Associated Kinesin by

832  Cyclin-Dependent Kinase 1. *Molecular and Cellular Biology*. 30:2594-2607.

833 Scopes, R.K. 2002. Enzyme activity and assay. John Wiley & Sons, Ltd, West Sussex, UK.

834 Song, Y., Y. Zhang, Y. Pan, J. He, Y. Wang, W. Chen, J. Guo, H. Deng, Y. Xue, X. Fang, and X.

Liang. 2020. The microtubule end-binding affinity of EB1 is enhanced by a dimeric
organization that is susceptible to phosphorylation. *Journal of Cell Science*. 133.

Storey, K.B. 2004. Functional Metabolism: Regulation and Adaptation. John Wiley & Sons. Inc.,
Hoboken, New Jersey.

Tan, D., W.J. Rice, and H. Sosa. 2008. Structure of the Kinesin13-Microtubule Ring Complex.
Structure. 16:1732-1739.

Tanenbaum, M.E., L. Macurek, B. van der Vaart, M. Galli, A. Akhmanova, and R.H. Medema. 2011.
A Complex of Kit18b and MCAK Promotes Microtubule Depolymerization and Is
Negatively Regulated by Aurora Kinases. *Curr. Biol.* 21:1356-1365.

Tinevez, J.-Y., N. Perry, J. Schindelin, G.M. Hoopes, G.D. Reynolds, E. Laplantine, S.Y. Bednarek,
S.L. Shorte, and K.W. Eliceiri. 2017. TrackMate: An open and extensible platform for
single-particle tracking. *Methods*. 115:80-90.

Tournebize, R., A. Popov, K. Kinoshita, A.J. Ashford, S. Rybina, A. Pozniakovsky, T.U. Mayer, C.E.
Walczak, E. Karsenti, and A.A. Hyman. 2000. Control of microtubule dynamics by the
antagonistic activities of XMAP215 and XKCM1 in Xenopus egg extracts. *Nature Cell
Biology*. 2:13-19.

Trofimova, D., M. Paydar, A. Zara, L. Talje, B.H. Kwok, and J.S. Allingham. 2018. Ternary complex
of Kif2A-bound tandem tubulin heterodimers represents a kinesin-13-mediated
microtubule depolymerization reaction intermediate. *Nature Communications*. 9.

Vasudevan, K.K., Y.-Y. Jiang, K.F. Lechtreck, Y. Kushida, L.M. Alford, W.S. Sale, T. Hennessey,
and J. Gaertig. 2015. Kinesin-13 regulates the quantity and quality of tubulin inside cilia.
Molecular Biology of the Cell. 26:478-494.

Walczak, C.E., E.C. Gan, A. Desai, T.J. Mitchison, and S.L. Kline-Smith. 2002. The microtubule-
destabilizing kinesin XKCM1 is required for chromosome positioning during spindle
assembly. *Curr. Biol.* 12:1885-1889.

Walczak, C.E., T.J. Mitchison, and A. Desai. 1996. XKCM1: A Xenopus kinesin-related protein that
regulates microtubule dynamics during mitotic spindle assembly. *Cell*. 84:37-47.

Wang, W., S. Cantos-Fernandes, Y. Lv, H. Kuerban, S. Ahmad, C. Wang, and B. Gigant. 2017.
Insight into microtubule disassembly by kinesin-13s from the structure of Kif2C bound to
tubulin. *Nature Communications*. 8.

Wang, W., Q. Jiang, M. Argentini, D. Cornu, B. Gigant, M. Knossow, and C. Wang. 2012. Kif2C
Minimal Functional Domain Has Unusual Nucleotide Binding Properties That Are Adapted
to Microtubule Depolymerization. *Journal of Biological Chemistry*. 287:15143-15153.

Wang, W., T. Shen, R. Guerois, F. Zhang, H. Kuerban, Y. Lv, B. Gigant, M. Knossow, and C. Wang.
2015. New Insights into the Coupling between Microtubule Depolymerization and ATP
Hydrolysis by Kinesin-13 Protein Kif2C. *Journal of Biological Chemistry*. 290:18721-
18731.

Widlund, P.O., M. Podolski, S. Reber, J. Alper, M. Storch, A.A. Hyman, J. Howard, and D.N.
Drechsel. 2012. One-step purification of assembly-competent tubulin from diverse
eukaryotic sources. *Molecular Biology of the Cell*. 23:4393-4401.

Wittmann, T., A. Hyman, and A. Desai. 2001. The spindle: a dynamic assembly of microtubules and
motors. *Nature Cell Biology*. 3:E28-E34.

Wordeman, L., and T.J. Mitchison. 1995. IDENTIFICATION AND PARTIAL
CHARACTERIZATION OF MITOTIC CENTROMERE-ASSOCIATED KINESIN, A

KINESIN-RELATED PROTEIN THAT ASSOCIATES WITH CENTROMERES
DURING MITOSIS. *Journal of Cell Biology*. 128:95-105.

Zhang, R., G.M. Alushin, A. Brown, and E. Nogales. 2015. Mechanistic Origin of Microtubule
Dynamic Instability and Its Modulation by EB Proteins. *Cell*. 162:849-859.

Zhu, Z.Q.C., K.K. Gupta, A.R. Slabbekoorn, B.A. Paulson, E.S. Folker, and H.V. Goodson. 2009.
Interactions between EB1 and microtubules: dramatic effect of affinity tags and evidence
for cooperative behavior. *Journal of Biological Chemistry*. 284:32651-32661.

Zong, H., M. Hazelbaker, C. Moe, S.C. Ems-McClung, K. Hu, and C.E. Walczak. 2021. Spatial
regulation of MCAK promotes cell polarization and focal adhesion turnover to drive robust
cell migration. *Molecular Biology of the Cell*. 32:590-604.

Figure 1. MCAK preferentially binds to growing microtubule ends

(A) Representative kymographs of single-molecule GFP-MCAK (green, 10 nM,
measured in terms of monomer concentration unless otherwise stated) binding events
on dynamic microtubules (red, tubulin: 16 mM) growing from GMPCPP microtubule
seeds (blue) in the presence of 1 mM ATP. A plus-end binding event and a lattice binding
event were marked by the white and blue arrowheads, respectively. Scale bars: vertical,
5 s; horizontal, 2 μm.

(B) Representative intensity profiles of a single GFP-MCAK molecule (green, 10 nM)
and a growing microtubule end (red, tubulin: 16 mM) at one frame during the dwell
time of a binding event (corresponding to the white arrowhead in panel **A**). The peak
position of a GFP-MCAK molecule was determined by Gaussian fitting (green dashed
line).

(C) Position probability distributions of GFP-MCAK binding sites (green, n=152
events) along the microtubule longitudinal axis. The red curve represents the fitting
curve of the growing microtubule end as the positional reference. The MCAK binding
region was defined as the area within the FWHM (double black arrow, 162 nm) of the
fitted green curve.

(D) Statistical quantification of on-rate (k_{on}) of GFP-MCAK's binding to the plus-end
versus lattice of dynamic microtubules in the presence of 1 mM ATP (n=66
microtubules from 3 assays).

(E) Statistical quantification of $R_{E/L}$ in the presence of 1 mM ATP (n=66 microtubules
from 3 assays).

(F) Statistical quantification of the dwell time of GFP-MCAK's binding to the growing
end versus lattice of dynamic microtubules in the presence of 1 mM ATP (n=966 events
from 3 assays for the plus end; n=702 events from 3 assays for lattice).

(G) Mean-squared displacement (MSD) of GFP-MCAK plotted against the time

interval (t, 0.1s per frame). The diffusion coefficient (D) was calculated via linear
regression of the equation ($<x^2>= 2Dt$), yielding 0. 023 $\mu m^2 \, s^{-1}$ (error bars represent
SEM, n=203 trajectories).
In panel **D**, **E** and **F**, all the data were presented as mean ± SEM. All comparisons
were performed using two-tailed Mann-Whitney U test with Bonferroni correction, n.s.,
no significance; ***, p<0.001.

Figure 2. MCAK binds to the entire GTP cap of growing microtubule ends

(A) Representative kymographs showing the individual binding events of GFP-MCAK
(1 nM, with 1 mM ATP, left), EB1-GFP (10 nM, middle) or XMAP215-GFP (1 nM,
right) at growing microtubule ends (red, tubulin: 12 μM). The plots showed the
representative intensity profiles of single molecules (green) and growing microtubule
ends (red) at one frame during the dwell time of the individual binding event (white
arrowheads). Peak positions were determined by Gaussian fitting (green dashed line).
Scale bars: vertical, 5 s; horizontal, 2 μm.

(B) Position probability distributions of the binding sites of GFP-MCAK, EB1-GFP
(black, n=184 events) and XMAP215-GFP (blue, 142 events) along the microtubule
longitudinal axis. The fitting curve of the growing microtubule end (red) was used as
the positional reference. The binding region was defined as the area within the FWHM
of the fitted curves. The GFP-MCAK molecules localized to the left of XMAP215-
GFP's binding region was defined as proximally distributed (grey bar), and those
localized to the right of EB1-GFP's binding region was defined as distally distributed
(orange bar).

(C) Averaged intensity profiles of the growing microtubule ends decorated with GFP-MCAK (green, 123 microtubules), EB1-GFP (black, 184 microtubules) and XMAP215-GFP (blue, 142 microtubules). Data were presented as mean ± std, and no significant difference was observed between the three conditions (Kolmogorov-Smirnov test, End_{EB1} vs End_{MCAK}, p=0.68; End_{MCAK} vs $End_{XMAP215}$, p=0.68; End_{EB1} vs $End_{XMAP215}$, p=1). In panels **B** and **C**, the red arrows indicated microtubule growth direction.

(D) Representative kymographs showing concurrent binding events of XMAP215-GFP (25 nM) and MCAK-RFP (20 nM) at growing microtubule ends (tubulin: 12 μM) in the presence of 1 mM ATP. The plots showed the representative intensity profiles of XMAP215-GFP (blue), MCAK-RFP (green) and growing microtubule ends (red) at one frame during the dwell time of the binding event (cyan arrowhead). Peak positions were determined by Gaussian fitting (dashed line). The spacing between the two peaks represents the instantaneous distance between MCAK and XMAP215 (double black arrow). Scale bars: vertical, 5 s; horizontal, 2 μm.

(E) Probability distribution of the distance between XMAP215-GFP and MCAK-RFP (62 events from 3 assays) at growing microtubule ends. In this plot, the position of XMAP215-GFP was taken as the origin. A negative distance indicated a more proximal localization.

Figure 3. The growing end-binding preference of MCAK is dependent on its
nucleotide state

(A) Representative kymographs of single-molecule GFP-MCAK (green) binding
events on dynamic microtubules (red, tubulin: 16 μM) growing from the GMPCPP
microtubule seeds (blue) in the presence of 1 mM AMPPNP (left), 1 mM ADP (middle)
and APO (right) (the nucleotide-free state). The plus-end binding events were indicated
by white arrowheads. Scale bar: vertical, 5 s; horizontal, 2 μm.

(B) Statistical quantification of on-rate (k_{on}) of GFP-MCAK binding to the plus end (E)
versus the lattice (L) of dynamic microtubules in the presence of 1 mM AMPPNP (n=29
microtubules from 2 assays), 1 mM ADP (n=21 microtubules from 2 assays) and APO
(n=36 microtubules from 3 assays). The cyan and purple dashed lines denoted the on-
rate value of MCAK binding at the plus end and lattice in the ATP condition (from **Fig.
1**), respectively.

(C) Statistical quantification of $R_{E/L}$ in the presence of 1 mM AMPPNP (n=29
microtubules from 2 assays), 1 mM ADP (n=21 microtubules from 2 assays) and APO
(n=36 microtubules from 3 assays). Dashed line: the mean value of $R_{E/L}$ in the ATP
condition from **Fig. 1**.

(D) Statistical quantification of the dwell time for GFP-MCAK binding at the growing
ends and lattice of dynamic microtubules in the presence of 1 mM AMPPNP (n=231
events from 2 assays for the plus end; n=142 events from 2 assays for lattice), 1 mM
ADP (n=327 events from 2 assays for the plus end; n=1184 events from 2 assays for
lattice) and APO (n=71 events from 3 assays for the plus end; n=573 events from 3
assays for lattice). The cyan and purple dashed lines denoted the dwell time value of

MCAK binding at the plus end and lattice in the ATP condition (from **Fig. 1**),
respectively.

In the panel **B**, **C** and **D**, all the data were presented as mean $\pm$ SEM. All the
comparisons were made against the corresponding value in the ATP condition from **Fig.**
1 using two-tailed Mann-Whitney U test with Bonferroni correction, n.s., no
significance; ***, $p < 0.001$.

Figure 4. MCAK strongly binds to GTPγS microtubules in a nucleotide-dependent manner

(A) Representative projection images of GFP-MCAK binding to GTPγS microtubules (red) versus GDP microtubules (cyan) in the presence of 1 mM ATP (left, 10 nM GFP-MCAK), 1 mM AMPPNP (left middle, 1nM GFP-MCAK), 1 mM ADP (right middle, 10 nM GFP-MCAK) and at the APO state (right, 0.1 nM GFP-MCAK). Note that the binding on the GDP microtubules was compared to that on the GTPγS microtubules in the same flow cell. The GFP-MCAK binding to the GTPγS and GDP microtubules was indicated by the red and cyan arrowhead, respectively. Scale bar: 5 μm.

(B) Statistical quantification of the normalized fluorescence intensity of GFP-MCAK (1 nM) on different microtubules in the presence of 1 mM ATP (95 GTPγS microtubules, 28 GDP microtubules from 2 assays), 1 mM AMPPNP (133 GTPγS microtubules, 56 GDP microtubules from 3 assays), 1 mM ADP (111 GTPγS microtubules, 44 GDP microtubules from 2 assays) or at the APO state (52 GTPγS microtubules, 36 GDP microtubules from 3 assays). All data were normalized to the binding to GTPγS microtubules binding in the AMPPNP condition.

(C) The binding intensity ratios of GFP-MCAK at GTPγS microtubules to GDP microtubules under various nucleotide conditions. ATP: n=28, from 2 assays. AMPPNP: n=56 GTPγS/GDP from 3 assays. ADP: n=44 from 2 assays. APO: n=36 from 3 assays in APO state. The dashed line represented 1. Statistical comparisons were performed between the ratios and 1.

1014   In panels **B** and **C**, all the data were presented as mean ± SEM. All comparisons
1015   were performed using two-tailed Mann–Whitney U test with Bonferroni correction, ***,
1016 $p<0.001$.

1017

1018

Figure 5. MCAK binds to the ends of GMPCPP microtubules in a nucleotide state-dependent manner

(A, B, C, D) Representative fluorescence projection images and intensity profiles showing GFP-MCAK binding to the ends and lattice of GMPCPP microtubules in the presence of 1 mM ATP (10 nM GFP-MCAK), 1 mM AMPPNP (1 nM GFP-MCAK), 1 mM ADP (10 nM GFP-MCAK) or at the APO state (0.1nM GFP-MCAK). Red arrowhead: the signal of MCAK enriching at the ends. Intensity profiles (right panels) showed spatial intensity signal distributions of the microtubule (black curve) and GFP-MCAK (green curve). Red and black bars indicated the end and the lattice, respectively. Scale bar=2 μm.

(E) Statistical quantification of the normalized binding intensity of GFP-MCAK (1 nM) at the end versus lattice regions of GMPCPP microtubules in the presence of 1 mM ATP

(40 microtubules from 3 assays), 1 mM AMPPNP (47 microtubules from 3 assays), 1
mM ADP (45 microtubules from 3 assays) or at the APO state (52 microtubules from 3
assays). All the data were normalized to the binding intensity of GFP-MCAK at
GMPCPP microtubule lattices in the AMPPNP condition. All the data were presented
as mean ± SEM. Statistical analysis was performed using two-tailed paired *t*-test by
Bonferroni correction, ***, p<0.001.

(F) The end-to-lattice binding intensity ratios of GFP-MCAK at GMPCPP microtubules
under various nucleotide conditions. The ratios were presented as mean ± SEM. The
dashed line represents 1. Statistical comparisons were performed between the ratios and
1 using two-tailed Mann-Whitney U test with Bonferroni correction, ***, p<0.001.

1041

1042

Figure 6. The α4 helix and the L2 loop of MCAK contribute to the preferential binding to the EB cap

(A) Structural model of the MCAK^sN+M-tubulin complex (PDB ID: 5MIO; X-ray diffraction). Top: Domain organization of MCAK. Bottom: Enlarged view of α4 helix and L2 loop, regions mediating the interaction between MCAK and tubulin, with the key sites (K524 and V298) were highlighted in red. The residues in tubulin that potentially interact with K524 and V298 were highlighted in green.

(B) Representative kymographs of single-molecule binding events of GFP-MCAK^K524A (10 nM) and GFP-MCAK^V298S (30 nM) at growing microtubules (red, tubulin: 16 μM) in the presence of 1 mM ATP. The end-binding events were indicated by red arrowheads. Scale bars: vertical, 5 s; horizontal, 2 μm.

(C) Statistical quantification of on-rate (k_{on-P}) of GFP-MCAK^K524A (31 microtubules from 3 assays) and GFP-MCAK^V298S (23 microtubules from 2 assays) at growing microtubule ends in the presence of 1 mM ATP. The data were presented as mean ± SEM and were compared to that of wild-type GFP-MCAK (from Fig. 1).

(D) Statistical quantification of dwell time of GFP-MCAKK524A (18 binding events from 3 assays) and GFP-MCAK^V298S (39 binding events from 3 assays) at growing microtubule ends in the presence of 1 mM ATP. The data were presented as mean ± SEM and were compared to that of wild-type GFP-MCAK in the presence of ATP (from **Fig. 1**). The dashed line: the mean dwell time of wild-type GFP-MCAK.

(E) Representative fluorescence projection images of 5 nM GFP-MCAK^K524A and 10 nM GFP-MCAK^V298S binding to GTPgS (red arrowhead) versus GDP (cyan arrowhead) microtubules in the presence of 1 mM AMPPNP. Scale bar: 5 mm.

(F) Statistical quantification of normalized binding intensity of 1 nM GFP-MCAK^{K524A} (144 GTPgS microtubules and 62 GDP microtubules from 3 assays) and GFP-MCAK^{V298S} (124 GTPgS microtubules, and 59 GDP microtubules from 3 assays) on different microtubules. All the data were normalized to the binding intensity of wild-type GFP-MCAK at GTPgS microtubules in the AMPPNP condition (from **Fig. 4**). Data were presented as mean ± SEM.

(G) The binding intensity ratios of GFP-MCAK^{K524A} or GFP-MCAK^{V298S} on GTPγS microtubules to that on GDP microtubules in the presence of 1 mM AMPPNP. GFP-MCAK^{K524A}: n=62 from 3 assays in AMPPNP state. GFP-MCAK^{V298S}: n=59 GTPγS/GDP binding ratios from 3 assays in AMPPNP state. Dashed line: the GTPγS/GDP ratio of wild-type GFP-MCAK. The data were presented as mean ± SEM and were compared to that for wild-type GFP-MCAK (from **Fig. 4**).

In panel **C**, **D**, **F** and **G**, all comparisons were performed using two-tailed Mann-Whitney U test with Bonferroni correction, n.s., no significance; *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$.

1082

Figure 7. Functional specification of MCAK and XMAP215 at growing microtubule ends

(A) Representative kymographs of dynamic microtubules (tubulin: 10 μM) under four conditions: control (no MAPs), 20 nM MCAK (1 mM ATP), 50 nM XMAP215 or both (1 mM ATP). Scale bars: vertical, 100 s; horizontal, 2 μm.

(B) Probability distribution of the microtubule lifetime under the conditions indicated in the panel **A**. Lines represent gamma function fits. Control: n=443 microtubules from 7 assays. 20 nM MCAK: n=621 microtubules from 7 assays. 50 nM XMAP215: n=237 microtubules from 3 assays. 20 nM MCAK+50 nM XMAP215: n=283 microtubules from 3 assays.

(C) The probability-based catastrophe frequency versus microtubule lifetime showing how the likelihood of catastrophe depended on the age of microtubules. Data derived from panel **A**.

1096

1097

Figure 8. Cartoon schematics depicting the working model of MCAK at growing microtubule ends

(A) A hypothetical model for the binding cycle of MCAK, given its binding preference on the EB cap. Arrows indicate the transitions in nucleotide-dependent MCAK functional states. T: MCAK·ATP. D: MCAK·ADP.

(B) Cooperation schematics of MCAK, EB1 and XMAP215 at growing microtubule ends. Pathway 1: the direct binding of MCAK to microtubule ends. Pathway 2: the indirect binding of MCAK to microtubule ends via EB1.

1106

Supplementary Information Appendix

Title:

MCAK Recognizes the Nucleotide-Dependent Feature at Growing Microtubule Ends

Authors and affiliations:

Wei Chen[1] (chenwill@mail.tsinghua.edu.cn)

Yin-Long Song[1] (y.song4@uu.nl)

Jian-Feng He[1] (jianfenghe2015@mail.tsinghua.edu.cn)

Xin Liang[1,*] (xinliang@tsinghua.edu.cn)

[1] IDG/McGovern Institute for Brain Research, State Key Laboratory of Complex, Severe, and Rare
Diseases, School of Life Sciences, Tsinghua University, Beijing 100084, China

Corresponding author:

Name: Xin Liang
Address: Room E118, Medical School, Tsinghua University, 100084 Beijing, China
Tel: +86-10-62772563
Email: xinliang@tsinghua.edu.cn
ORCID: 0000-0001-7915-8094

Supplementary materials:

Supplementary appendix-1 (Figures s1-s6)

Supplementary appendix-2 (A simple model for the functional contribution of
MCAK's direct end-binding mechanism, including Figure s7-s8)

1127

Supplementary appendix-1

1129

1130 **Figure s1.** Biochemical and functional characterization of purified MCAK, its variants,
1131 EB1 and XMAP215

1132 **(A)** Domain architecture schematics of GFP-MCAK and its mutants, MCAK-RFP,
1133 EB1-GFP, and XMAP215-GFP used in this study.

(B) The western blot result showing the removal of His-tag in purified GFP-MCAK. In the figure, the molecular weight marker lane and the protein sample lane were merged. The molecular weight of GFP-MCAK monomer is 110 kDa. Anti-his antibody was used for detection.

(C-I) Size-exclusion chromatography (Superose™ 6 Increase 5/150 GL) profiles of GFP-MCAK, GFP-MCAK^{K524A}, GFP-MCAK^{V298S}, MCAK-RFP, GFP-MCAK^{sN+M}, EB1-GFP and XMAP215-GFP-his$_6$. The molecular weight of GFP-MCAK, GFP-MCAK^{K524A}, GFP-MCAK^{V298S} and MCAK-RFP monomer were 110 kDa. The molecular weight of GFP-MCAK^{sN+M} was 73 kDa. The molecular weight of EB1-GFP monomer was 60 kDa. The molecular weight of XMAP215-GFP-his$_6$ was 260 kDa. The elution volumes of the peak for GFP-MCAK, GFP-MCAK^{K524A}, GFP-MCAK^{V298S}, MCAK-RFP, GFP-MCAK^{sN+M}, EB1-GFP and XMAP215-GFP-his$_6$ were 1.85 ml, 1.84 ml, 1.91 ml, 1.90 ml, 2.25 ml, 2.11 ml and 1.78 ml, respectively. In panel **C**, the elution volume of the peak corresponded to the molecular weight range between 120 kDa (EB1-GFP dimer) and 260 kDa (XMAP215-GFP-his$_6$), suggesting that GFP-MCAK exists as a dimer under this experimental condition.

(J) SDS-PAGE gels of elution fractions corresponding to peaks from size-exclusion chromatography profiles of GFP-MCAK, GFP-MCAK^{K524A}, GFP-MCAK^{V298S}, MCAK-RFP, GFP-MCAK^{sN+M}, EB1-GFP and XMAP215-GFP-his$_6$ in panels **C-I**.

(K) Fluorescence intensity distributions of single GFP-MCAK^{sN+M} (blue, n=327 binding events from 3 assays) and GFP-MCAK (red, n=509 binding events from 3 assays) molecules were fitted by Gaussian fitting. The background intensity was subtracted. The concentrations of the two proteins were both 10 nM. Dynamic microtubules (tubulin: 16 μM) were polymerized in the presence of 1 mM ATP. The intensity range-1 (blue, $\mu \pm 2\sigma = 196 \pm 79$ A.U.) and range-2 (red, $\mu \pm 2\sigma = 300 \pm 145$ A.U.) were used to determine if a fluorescence spot represents a single monomer and dimer, respectively.

(L) The apparent off-rate (k_{off}) of GFP-MCAK at growing microtubule ends in the presence of ATP (circle), AMPPNP (down triangle), ADP (up triangle) and APO (diamond). Data of dwell time were from **Fig. 1F**. k_{off} was calculated by fitting the dwell time of individual GFP-MCAK binding events to a single exponential function.

(M) Statistical quantification of the growth rates of dynamic microtubules with GFP-MCAK in the presence of 1 mM ATP (57 microtubules from 3 assays), AMPPNP (38 microtubules from 2 assays), ADP (24 microtubules from 2 assays) and APO (34 microtubules from 3 assays). Data were from the experiments shown in panel **Fig. 1A**, which were presented as mean ± SEM. Statistical analysis was performed using two-tailed Mann-Whitney U test with Bonferroni correction, n.s., no significance, ***, p<0.001.

1173

Figure s2. Single-molecule localization analysis at growing microtubule ends

(A) Representative kymograph showing a binding event of GFP-MCAK (1 nM, with 1 mM ATP, left) at the growing microtubule end (red, tubulin: 12 μM). Schematic illustration of MCAK binding to the microtubule end (right). The red cross represented the reference location point. The green dots represented the position of MCAK at every frame during its dwell time. The orange cross indicated the average position of MCAK during the dwell time, which represented the location of MCAK over this period. Scale bars: vertical, 2 s; horizontal, 2 μm.

(B) Snapshot of a single GFP-MCAK molecule (green), and the microtubule (red) at the timepoint marked by a black arrow in panel **A** (left). Intensity profiles of the microtubule (red) and the single molecule (green) at this moment. The peak position of the single molecule was determined using Gaussian fitting (green dashed line), which represented the location of the molecule at this moment. Scale bar: 2 μm.

1187

1188

Figure s3. Projection of single-molecule fluorescence images

1189

1190 Representative images showing how to generate a summation image using 1000 frames
1191 of raw images. The binding intensity of GFP-MCAK at GDP versus GTPγS
1192 microtubule was indicated using green and yellow arrowheads, respectively. The
1193 concentration of GFP-MCAK here was 1 nM. The experiment was performed in the
1194 presence of 1 mM AMPPNP. Scale bar: 5 μm.

1195

1196

Figure s4. EB1-GFP strongly binds to GTPγS microtubules

(A) Representative projection images of EB1-GFP binding to GTPγS microtubules. The
binding at GDP microtubules was compared to that at GTPγS microtubules in the same
flow cell. In GFP channel, the fluorescence intensity of EB1-GFP binding to GTPγS
and GDP microtubules were indicated by red and cyan arrowhead, respectively. Scale
bar: 5 μm.

(B) The binding intensity ratios of EB1-GFP on GTPγS microtubules to that on GDP
microtubules under various nucleotide conditions. 36 GTPγS/GDP binding ratios from
3 assays. Dashed line on the plot represented 1. All the data were presented as mean ±
SEM. Statistical comparison was performed between the ratios and 1. The comparison
was performed using two-tailed Mann–Whitney U test with Bonferroni correction, ***,
p<0.001.

1209

1210

Figure s5. MCAK preferentially binds to the ends of taxol stabilized GDP microtubules

(A) Representative projection images and intensity profiles showing GFP-MCAK (10
mM) binding at the ends of taxol stabilized GDP microtubules in the presence of 1 mM
ATP. Red arrowhead: the end-binding of MCAK events. Intensity profiles (right panels)
showed spatial intensity signal distribution of microtubule (black curve) and GFP-
MCAK (green curve) along microtubule longitudinal axes from left to right. Red and
black bars above the intensity profiles indicated end and lattice regions, respectively.
Scale bar=2 μm.

(B) Statistical quantification of the binding intensity of GFP-MCAK at the end and
lattice of taxol stabilized GDP microtubules in the presence of 1 mM ATP (35
microtubules from 2 assays). Data were scaled relative to the binding intensity of GFP-
MCAK at the lattice of taxol stabilized GDP microtubules. Data were presented as mean
± SEM. Statistical analysis was performed using two-tailed pair *t*-test by Bonferroni
correction, ***, p<0.001.

(C) The ration of the binding intensity of GFP-MCAK at the end to that on the lattice
of GDP microtubules in ATP state. The data were presented as mean ± SEM. Dashed
line on the plot represents 1. Statistical comparisons were performed between the ratios
and 1 using two-tailed Mann-Whitney U test with Bonferroni correction, ***, p<0.001.
1229

1230

Figure s6. GFP-MCAK[K525A] and GFP-MCAK[V298S] are depolymerizing-deficient and ATPase activity-reduced mutants

(A) Representative kymographs showing depolymerization of GMPCPP-stabilized microtubules by GFP-MCAK, GFP-MCAK[K525A] and GFP-MCAK[V298S] in the presence of 1 mM ATP. Scale bars: vertical, 100 s; horizontal, 2 μm.

(B) Statistical quantification of microtubule depolymerization rates of GFP-MCAK (n=21 microtubules from 3 assays), GFP-MCAK[K525A] (n=11 microtubules from 2 assays) and GFP-MCAK[V298S] (n=52 microtubules from 3 assays). Data were presented as mean ± SEM. Statistical analysis was performed using two-tailed Mann-Whitney U test with Bonferroni correction, ***, p<0.001.

(C) Microtubule-stimulated ATPase activities of wild-type GFP-MCAK (6 assays), GFP-MCAK[K525A] (4 assays) and GFP-MCAK[V298S] (4 assays). Data were normalized by the microtubule-stimulated ATPase activity of the wild-type. Statistical analysis was performed using two-tailed Student's t test, ***, p<0.001.

1245

 Supplementary appendix-2

1247   Here, we presented a simple model to calculate the relative contribution of the
1248 direct and EB-dependent end-binding of MCAK (**Fig. s7**).

$$
\begin{array}{ccccc}
\text{MCAK} & + & \text{EB1} & \overset{K_0}{\leftrightharpoons} & \text{MCAK·EB1} \\
+ & & + & & + \\
\text{MTE} & & \text{MTE} & & \text{MTE} \\
K_1 \Downarrow\Uparrow & & \Downarrow\Uparrow K_2 & & \Downarrow\Uparrow K_3 \\
\text{MCAK-MTE} & & \text{EB1-MTE} & & \text{MCAK·EB1-MTE}
\end{array}
$$

Figure s7. A simple model for the end-binding of MCAK and EB1. MCAK can bind to
microtubule growing ends through both the direct (left) and EB-dependent (right)
pathways. The dissociation constants were K_0, K_1, K_2 and K_3, respectively. MTE:
growing microtubule end.

1254   Based on the model, we had the dissociation constants:

$$K_0 = \frac{[MCAK]\cdot[EB1]}{[MCAK\cdot EB1]} \quad \text{Eq. 6}$$

$$K_1 = \frac{[MCAK]\cdot[MTE]}{[MCAK-MTE]} \quad \text{Eq. 7}$$

$$K_2 = \frac{[EB1]\cdot[MTE]}{[EB1-MTE]} \quad \text{Eq. 8}$$

$$K_3 = \frac{[MCAK\cdot EB1]\cdot[MTE]}{[MCAK\cdot EB1-MTE]} \quad \text{Eq. 9}$$

1259   Then, the relative contribution of the direct and EB-dependent end-binding of
1260 MCAK can be expressed as α:

$$\alpha = \frac{[MCAK-MTE]}{[MCAK\cdot EB1-MTE]} = \frac{K_3\cdot[MCAK]}{K_1\cdot[MCAK\cdot EB1]} = \frac{K_3\cdot K_0}{K_1\cdot[EB1]} \quad \text{Eq. 10}$$

Here, we considered two scenarios. In the cytoplasm, both EB1 and MCAK
undergo free diffusion and can associate with each other without restrictions. The
relative concentrations of MCAK and EB1 are critical parameters, but they may vary
across different cell types and remain unknown. We also considered the second scenario
in which MCAK is locally enriched at specific cellular localizations through an EB-
independent mechanism. For example, EB1 does not affect the localization of MCAK
at centromere and centrosome, nor does EB1 significantly affect the function of MCAK
there (Domnitz et al., 2012). Here, we assumed that that the local concentration of
anchored-state MCAK is relatively high; and EB1 remains diffusive and its
concentration is nearly constant, as it is continuously replenished in the local space from
the vast cytoplasmic pool. In both cases, the ratio of K_3 to K_1 emerges as a key
determinant.

K_3 is the end-binding affinity of MCAK·EB1 complex. Intuitively, it depends on the respective microtubule-binding affinities of MCAK and EB1, as well as the cooperativity, if any, of their microtubule-binding behaviors. K_3 can be expressed as:

$$\frac{1}{K_3} = a\frac{1}{K_1} + b\frac{1}{K_2} \quad \text{Eq. 11}$$

$$K_3 = \frac{K_2 \cdot K_1}{aK_2 + bK_1} \quad \text{Eq. 12}$$

where a and b represent the weighting factors of binding sites or cooperativity factors of the binding behaviors. Therefore, K_3 shows a positively correlated, monotonically increasing dependence on K_1, indicating that the increase in the end-binding affinity of MCAK contributes to that of the MCAK·EB1complex. Therefore, we think that MCAK's functional impact at microtubule ends derives not only from its intrinsic end-binding capacity, but also its ability to strengthen the EB1-mediated end association pathway.

In the simplest case, the formation of the MCAK·EB1-MTE complex arises from the binding of either MCAK or EB1 to microtubule ends; and the binding behaviors for MCAK and EB1 are independent ($a=1$; $b=1$). Consequently, K_3 can be expressed as:

$$K_3 = \frac{K_2 \cdot K_1}{K_2 + K_1} \quad \text{Eq. 13}$$

if $K_1 \ll K_2$, then

$$K_3 \approx K_1 \quad \text{Eq. 14}$$

if $K_1 \gg K_2$, then

$$K_3 \approx K_2 \quad \text{Eq. 15}$$

If $K_1 \approx K_2$, then

$$K_3 \approx \frac{K_1}{2} \quad \text{Eq. 16}$$

In our experiments, we measured the dissociation constants of MCAK to growing microtubule ends is 69 μM (K_1). We also performed similar experiments with EB1 and found that EB1 showed the dissociation constant of 722 μM (K_2) for growing microtubule ends (**Fig. s8**), similar to the value reported in our previous report (Song et al., 2020). Therefore, substituting Eq. 14 into Eq. 10, we obtained

$$\alpha \approx \frac{K_0}{[EB1]} \quad \text{Eq. 17}$$

Here, if we assume that the cytoplasmic concentration of EB1 is twice the value of K_0, then $\alpha=0.5$, indicating that 50% of MCAK binds to microtubule ends via the direct binding pathway; even if the EB1 concentration reaches ten times the value of K_0, 10% of MCAK still utilizes the direct binding pathway. Overall, as the EB1

1306   concentration increases relative to K_0, α decreases, reflecting a decline in the proportion

1307   of MCAK that associates with microtubule ends through the direct binding mechanism.

1308

Figure s8. The binding kinetics of single-molecule EB1-GFP binding to growing

microtubule end

(A) Statistical quantification of on-rate (k_{on}) of EB1-GFP's binding to the plus end of

dynamic microtubules (data calculated from Fig. 2, n=73 microtubules from 3 assays).

(B) The apparent off-rate (k_{off}) of EB1-GFP at growing microtubule ends (data

calculated from **Fig. 2**, n=153 bonding events from 3 assays). k_{off} was calculated by

fitting the dwell time of individual GFP-MCAK binding events to a single exponential

function.

1317