1

Full title: Dynamic metabolic and molecular changes during seasonal shrinking in *Sorex araneus*

Short title: Seasonal metabolic and molecular changes in *Sorex araneus*

William R. Thomas[1]*, Cecilia Baldoni[2,3], Yuanyuan Zeng[4], David Carlson[1], Julie Holm-

Jacobsen[4], Marion Muturi[2], Dominik von Elverfeldt[5], John Nieland[4], Dina K. N. Dechmann[2,3],

Angelique Corthals[6¶], Liliana M. Dávalos[1,7¶]*

[1]Department of Ecology and Evolution, Stony Brook University; Stony Brook, NY 11794, United
States.
15
16   [2]Max-Planck Institute of Animal Behavior; Radolfzell, 78315, Germany.
17
18   [3]University of Konstanz; Konstanz, 78464, Germany.
19
20   [4]Health Science and Technology, Aalborg University; Aalborg, 9100, Denmark.
21
22   [5]Division of Medical Physics, Department of Diagnostic and Interventional Radiology, University
23   Medical Center Freiburg, Faculty of Medicine, University of Freiburg, Freiburg, Germany.
24
25   [6]John Jay College of Criminal Justice; New York, NY 10019, United States.
26
27   [7]Consortium for Inter-Disciplinary Environmental Research, Stony Brook University; Stony
28   Brook, NY 11794, United States.
29

31   * Corresponding authors: William R. Thomas and Liliana M. Dávalos

32   **Emails:** William.thomas@stonybrook.edu, Liliana.Davalos@stonybrook.edu

35   ¶ These authors contributed equally to this work.

Abstract

To meet the challenge of wintering in place many high-latitude small mammals reduce energy

demands through hibernation. In contrast, short-lived Eurasian common shrews, *Sorex araneus*,

remain active and shrink, including energy-intensive organs in winter, regrowing in spring in an

evolved strategy called Dehnel's phenomenon. How this size change is linked to metabolic and

regulatory changes to sustain their high metabolism is unknown. We analyzed metabolic,

proteomic, and gene expression profiles spanning the entirety of Dehnel's seasonal cycle in wild

shrews. We show regulatory changes to oxidative phosphorylation and increased fatty acid

metabolism during autumn-to-winter shrinkage, as previously found in hibernating species. But in

shrews we also found upregulated winter expression of genes involved in gluconeogenesis: the

biosynthesis of glucose from non-carbohydrate substrates. Co-expression models revealed

changes in size and metabolic gene expression interconnect via FOXO signaling, whose

overexpression reduces size and extends lifespan in many model organisms. We propose that

while shifts in gluconeogenesis meet the challenge posed by high metabolic rate and active winter

lifestyle, FOXO signaling is central to Dehnel's phenomenon, with spring downregulation

limiting lifespan in these shrews.

Introduction

Wintering adaptations vary widely across animals and the most common strategies are conserving

energy by lowering metabolic rates and activity (e.g. hibernation or torpor) or migration. By

contrast, a handful of species that neither migrate nor slow their metabolism have evolved a

different wintering size plasticity known as Dehnel's phenomenon (1–4). This pattern of growth,

best studied in the Eurasian common shrew, *Sorex araneus*, is unlike that of almost every other

mammal. Instead of growing continuously to adulthood, juvenile *S. araneus* grow to an initial

maximum size in the first summer of their disproportionately brief life (~1 year). In anticipation

of harsh winter conditions, shrews then shrink, reaching a nadir in winter, followed by rapid

regrowth to their adult, breeding size in spring. This plasticity is not just a retooling of overall

body size, but also occurs in many vital organs, including the liver, brain, and spleen (2,5).

Although Dehnel's phenomenon has garnered much interest because of its potential for

regenerating organs (6), metabolic and regulatory changes during this wintering adaptation

remain largely unknown.

Dehnel's phenomenon corresponds to the unique physiological constraints shrews face. *S.*

araneus have one of the highest mammalian metabolic rates measured to date (7), requiring high

and constant food intake throughout the year (8). But supplying energy to maintain this high

metabolic rate is especially difficult in autumn and winter, when temperatures decrease and

resources become scarce (9,10). Therefore, Dehnel's phenomenon is construed as an evolutionary

adaptation that compensates for continuously high energetic demands by reducing body and

organ size, particularly of energy-expensive tissue (11). In support of this hypothesis, absolute

resting metabolic rate decreases with shrew size in winter, despite harsh extrinsic conditions

(10,12,13)

Wintering strategies encompass a spectrum of related traits and physiological processes (14).

Changes in other wintering strategies, such as metabolic shifts observed in hibernation (15,16),

may thus share common mechanisms with those involved in Dehnel's phenomenon. For example,

mechanistic analyses of mammalian hibernation have identified a toolkit of genes that regulates

vast metabolic changes (15–17), including those associated with increased fatty acid oxidation

simultaneous (18), as hibernators rely on gradual fat utilization over winter. Research on shrew

physiology indicates they also rely extensively on lipid metabolism in winter (8) and may share at

least some molecular mechanisms with hibernation.

But *S. araneus* remain active throughout winter, undergoing drastic shrinkage while navigating a

metabolically demanding environment. This active wintering strategy contrasts with hibernation,

suggesting key differences in metabolic and regulatory processes. For example, wintering shrews

show rapid fat turnover, with 50% fat turnover rates decreasing from 4.5 hours in summer

juveniles to 2.5 hours in winter (8). Consequently, metabolic profiles may differ significantly

between these strategies. In hibernators there is also a downregulation of genes involved in

glucose breakdown (18). In general, when energy stores are depleted, glucose is produced from

the liver through gluconeogenesis, which is then released into the bloodstream and metabolized in

the mitochondria via oxidative phosphorylation (19). Unlike hibernators that suppress this

response, wintering shrews seem to increase this starvation response as environments become

harsher (10). Thus, we hypothesized that shrews shrink by upregulating fatty acid metabolism,

with potential similarities to hibernators, but unlike them supplement their energy budget with

glucose to maintain their active winter lifestyle.

To examine metabolic dynamics and regulatory changes throughout Dehnel's phenomenon, and

its potential parallels to other wintering strategies, we collected phenotypic, metabolomic,

transcriptomic, and proteomic data from a shrew population across seasons matching the size

plasticity. We focused on blood, which circulates metabolites across the organisms, and the liver,

where many functions that regulate metabolism take place, including lipid and sugar processing.

Our results reveal seasonal changes in lipid metabolism and gluconeogenesis during shrew size

plasticity, which appears to relate to mitochondrial function. Metabolic and size changes are

regulated by pathways that include FOXO and insulin signaling, yet incur a cost to lifespan. By

 exploring these mechanisms, we offer novel insights into the interplay between metabolism, size,

110 and longevity, shedding light on both the wintering strategy of the common shrew and broader

111 mammalian physiology.

113 Results

114 Dehnel's phenomenon in German shrew population

Comparisons of body mass throughout the cycle confirmed patterns of size changes (Fig 1A).

Analyzing the 24 shrews collected for all stages of Dehnel's phenomenon (large summer

juvenile=5; shrinking autumn juvenile n=4; small winter juvenile n=5; regrowing spring adult

n=5; regrown Summer Adult n=5), we found body mass shrank as expected between summer

juveniles and winter juveniles ($BM_{WiJ-SuJ}$=-1.00 g, $p_{WiJ-SuJ}$<0.028), with regrowth as they

matured to spring adults ($BM_{SpA-WiJ}$=3.724 g, $p_{SpA-WiJ}$<0.0001). Shrew liver mass reached a

minimum as autumn juveniles ($LiM_{AuJ-SuJ}$=-904.1 mg, $p_{AuJ-SuJ}$=0.0949), with the significant

portion of regrowth occurring between the winter juvenile and spring adult phases

($LiM_{SpA-AuJ}$=536.7 mg, $p_{SpA-WiJ}$<0.001). These size changes both validated previous research

and were used to guide metabolomic, transcriptomic, and proteomics characterization of Dehnel's

phenomenon.

Fig 1. Confirming Dehnel's phenomenon and metabolic profiling of the blood metabolome.

(A) Mass change of body and liver through a year of Dehnel's phenomenon. Both the body and

the liver begin to decrease in mass as autumn juveniles, with body mass at a minimum as winter

juveniles, proceeded by regrowth as spring adults. Asterisks represent significant size changes

(adjusted p<0.05). **(B)** Heatmap of 28 statistically significant differentially concentrated

metabolites between stages of Dehnel's phenomenon. Hierarchical clustering using these

significant metabolites groups each profile into each season. **(C)** Three of these metabolites were

lipid metabolites (ethyl palmitoleate, docosahexanoic acid, arachidonic acid), with decreases in

autumn, winter, and spring individuals.

Metabolomic shifts in lipid metabolite concentrations

We characterized the blood plasma metabolome to quantify metabolites throughout the cycle and

evaluated the potential shifts in metabolism during shrinkage and regrowth. We validated 250

metabolites found in all seasons with multiple detection methods. Of these metabolites, we found

28 with significant differences in mean concentration across the stages of Dehnel's phenomenon

($p<0.05$, Fig 1B). Among differentially concentrated metabolites, three were lipid metabolites;

arachidonic acid (AA, $p=0.04$, $F=5.78$), ethyl palmitoleate (EP, $p=0.02$, $F=6.79$), and

docosahexanoic acid (DHA, $p=0.04$, $F=5.86$). These three lipid metabolites showed significant

decreases in peak area, a proxy for concentration, between summer juveniles and other stages of

size change (autumn juveniles, winter juveniles, spring adults), especially in winter juveniles

($AA_{WiJ-SuJ}$=-8.7 x 10^7 peak area, $DHA_{WiJ-SuJ}$=-1.2 x 10^8 peak area, $EP_{WiJ-SuJ}$=-3.1 x 10^6 peak

area), for which the concentrations of DHA and AA were at their minimum.

Regulatory changes associated with metabolic shifts

Results showed many differentially expressed genes in the liver between summer juveniles and

winter juveniles (shrinking stage), with an upregulation of genes related to mitochondrial, lipid,

and glucose metabolism. Comparisons revealed 964 differentially expressed genes ($p_{adj}<0.05$),

with 497 significantly upregulated and 467 significantly downregulated in winter juveniles (Fig

2A). We then used a ranked gene set enrichment test to identify functional effects of RNA

regulatory change. This analysis identified 25 pathways enriched with differentially expressed

genes (21 upregulated and 4 downregulated) (Fig 2D). Oxidative phosphorylation was the most

enriched pathway (p_{adj}<0.001, Normalized Enrichment Score=2.34, [NES]) consisting of 46

genes. This gene set included 21 upregulated genes that encode mitochondrial complex I subunit

proteins (NDUFs) and 19 upregulated genes encoding ATPases and synthases. Other pathways

enriched with upregulated genes included glycolysis and gluconeogenesis (p_{adj}<0.001,

NES=2.09), which contained the gene encoding for glucose-6 phosphatase catalytic subunit 1,

G6PC1 (p_{adj}=0.41, Log Fold Change=0.30, [LFC]), and fatty acid metabolism (p_{adj}<0.01,

NES=1.81).

Fig 2. Seasonal regulatory (transcriptomics and proteomics) change in the shrew liver. (A)

Volcano plots of significant (adjusted p<0.05) differentially expressed genes and proteins

(colored) between (from left to right); summer juveniles vs. winter juveniles (shrunk/RNA),

summer juveniles vs winter juveniles (shrunk/protein), winter juveniles vs spring adults

(regrowth/RNA), and winter juveniles vs spring adults (regrowth/protein) plotted by log fold-

change. Triangles represent lower p-values (-log(p)>10) or log fold changes (abs(LFC)>4)

beyond the limits of the graph**. (B)** Correlations of log fold change and significant gene overlap

between RNA and protein indicate high association between the summer vs. winter juveniles

analyses, and **(C)** winter juveniles vs spring adults. **(D)** Pathway enrichment of summer vs winter

juveniles between RNA (shown) and protein analyses, both with the largest enrichment of

oxidative phosphorylation pathways. **(E)** Normalized gene counts in pathways of interest.

We validated changes in metabolism-related gene expression between summer juveniles and

winter juveniles (shrinking stage) with liver proteomics. Despite the liver proteomics (n=1,377

proteins) data set being smaller than the gene expression set (n=24,205 loci), there was

substantial overlap in direction, significance, and pathway enrichment of changes between the

two. Using a linear model to estimate the correlation between differential change (summer

juveniles vs winter juveniles) in gene expression and protein concentration, we found that LFC in

gene expression was significantly associated with LFC in protein concentration (F=197.5,

df=1373, p<0.001), but the model only explains a small fraction of the variation between them

(R^2=0.13) (Fig 2B). Analyzing differential concentration of proteins, we found 124 proteins

significantly changed (p_{adj}<0.05) between summer juveniles and winter juveniles (78 increased

and 46 decreased in winter) (Fig 2A). Thirty-four of these proteins overlapped with differentially

expressed genes, all but one with matching mRNA and protein direction.

Ranked set enrichment for proteomics identified five enriched pathways (p_{adj}<0.05), four of

which overlapped the RNA pathway enrichment (oxidative phosphorylation, Parkinson's disease,

Alzheimer's disease, Huntington's disease). Oxidative phosphorylation was the most enriched

pathway in both mRNA and protein expression and included 17 overlapping genes/proteins, 11 of

which were found to be significant in at least one of the data sets (*ATP5MG*, *NDUFB4*, *PPA1*,

SDHB, NDUFV3, NDUFA6, NDUFB8, NDUFS3, ATP5PO, NDUFC2, NDUFS6; all in the

same direction). Proteomic analysis did not reveal upregulation in glycolysis/gluconeogenesis

(p_{adj}=1.00, NES=0.66) pathway, though all but one overlapping gene also increased in

concentration in proteomics, including three aldehyde dehydrogenase genes (*ALDH3A2*,

p_{adj}<0.01, LFC=0.64; *ALDH1B1*, p_{adj}<0.01, LFC=0.81; *ALDH9A1*, p_{adj}=0.47, LFC=0.35) (Fig

2E). Similarly, proteomic set enrichment analysis did not validate fatty acid metabolism

enrichment (p_{adj}=0.91, NES=1.00), but there was overlap (11 of 18 genes), including genes that

encode proteins that transport (*ACADM*, p_{adj}=0.17, LFC=0.40; *ACADVL*, p_{adj}=0.26, LFC=0.35)

and breakdown of long fatty acid chains (*ACOX1*, p_{adj}<0.001, LFC=0.81).

Comparisons between winter juveniles and spring adults (growth) showed extensive change in

gene expression coincident with metabolic change, but correlation with proteomics was weak. Of

2,462 significantly differentially expressed genes, 1,130 were upregulated and the rest

downregulated in spring adults (Fig 2A). Yet, differential mRNA expression only significantly

enriched two pathways, including the downregulation of cell cycle (p_{adj}<0.05, NES=-1.77) and

proteosome (p_{adj}<0.05, NES=-1.91) pathways. Comparing protein concentration between winter

juveniles and spring adults (growth), 183 proteins showed significant change (42 increased and

141 decreased) (Fig 2A). While 41 of these overlapped with significant differentially expressed

genes, nearly a quarter of them (10) were in the opposite direction. This is unsurprising, as the

model between protein LFC and mRNA LFC does not explain a large portion of the variation

(R^2=0.02, F=22.73, df=1374, p<0.001) (Fig 2C). Ranked protein enrichment corroborated

minimal correlation, as none of the 6 significant pathways: starch and sucrose metabolism

(p_{adj}<0.01, NES=2.33), TCA cycle (p_{adj}<0.01, NES=2.28), cysteine and methionine metabolism

(p_{adj}<0.05, NES=2.13), valine, leucine, and isoleucine degradation (p_{adj}<0.05, NES=1.92),

spliceosome (p_{adj}<0.05, NES=-1.71), and cell adhesion molecules (p_{adj}<0.05, NES=-1.75),

overlapped with mRNA gene set analyses. There were, however, individual genes validated in

change of direction with proteomic data found in upregulated processes in winter juveniles that

decreased in spring adults (LFC<0): oxidative phosphorylation (*NDUFB4*, *PPA1*),

gluconeogenesis (*ALDH3A2*, *PGK1*, *ALDH1B1*), and fatty acid metabolism (*ACOX1*).

Weighted Gene Correlation Network

Differential expression analyses focused on genes of high effect between two size extremes, but

large modules comprising genes of *small* effect could also play pivotal roles in regulating

metabolic changes across all seasons. We characterized interactions in expression among genes

and identified modules of coexpressed liver genes correlated with liver size using weighted gene

co-expression network analysis (WGCNA). We identified 37 modules (Fig 3), 10 of which were

significantly associated with liver size ($p < 0.05$). Among these, the "grey60" module is a small

gene network most positively correlated with larger liver size (n=352 genes, $p < 0.001$, r=0.81,

df=21) and the "red" module is a large network positively correlated with small liver size

(n=3,268 gene, $p = 0.001$, r=-0.63, df=21). Although no KEGG pathways were significant after

multiple test correction when testing for gene set enrichment from the "grey60" module

significant genes (n=342 genes, 10 genes were not significantly associated to network), metabolic

pathways was highly significant ($p < 0.01$). Thirty-four KEGG pathways from the "red" module

significant genes (n=2,706 genes) were significant after multiple test correction ($p_{adj} < 0.05$). These

pathways included FOXO signaling ($p_{adj} < 0.01$), as well as pathways upstream (MAPK signaling,

$p_{adj} < 0.05$) and downstream (insulin resistance, $p_{adj} < 0.01$; longevity regulating pathways,

$p_{adj} < 0.05$) of this pathway.

Fig 3. Gene networks highly correlated with seasonal liver size plasticity. Pathway

enrichment of the large (n=3268 genes) "red" module highlighted functional associations to

FOXO signaling and insulin resistance (plotted genes), while the small (n=342 genes) "grey60"

module related to metabolic pathways. Central to FOXO signaling and insulin resistance

pathways is the FOXO1 gene (n=19 connections)

Discussion

Eurasian common shrews, *Sorex araneus*, require constant food intake to sustain their

extraordinarily metabolic rates throughout the year (7). Maintaining their energetic budget is

particularly difficult in winter, as temperatures drop, as does available food. Within a German

shrew population, we found clear signs of a seasonal size plasticity known as Dehnel's

phenomenon (1–3). These shrews radically decrease their size in autumn through winter, with

recovery in the ensuing spring. Using this population with confirmed Dehnel's phenomenon, we

could then identify seasonal changes to metabolism and underlying regulatory processes.

We hypothesized that, as in hibernators, wintering relies on fat utilization to maintain metabolic

homeostasis in winter, but differs as shrews rapidly turnover fat (8) during shrinkage.

Transcriptomic analyses of winter juvenile shrews revealed upregulation of key lipid metabolism

genes, including those involved in fatty acid oxidation: acyl-CoA dehydrogenase medium chain

(*ACADM*), acyl-CoA dehydrogenase very long chain (*ACADVL*), and acyl-CoA oxidase 1

(*ACOX1*). *ACOX1*, which encodes the rate-limiting enzyme for oxidizing very long chain fatty

acids, is critical for inter-organ metabolic communication, such that liver-specific knockouts of

this gene in mice increase adipose tissue browning and circulating levels of polyunsaturated

omega-3 fatty acids (20). In shrunken winter juveniles we observe the same correlation, an

increase in *ACOX1* expression paired with a decrease in circulating levels of polyunsaturated

fatty acids (DHA and AA). This contrasts with hibernators: circulating DHA and AA rise

significantly in wintering bears and hibernating arctic ground squirrels (16,21–23). Regulatory

similarities and metabolomic differences indicate both wintering strategies rely upon lipid

metabolism in the liver. But while hibernators conserve fat reserves over winter, shrews exhibit

rapid fat turnover (8), decreasing levels of polyunsaturated fatty acids as they maintain activity in

winter.

Regulatory changes related to mitochondrial function further highlight differences in metabolic

processes between Dehnel's phenomenon and hibernation. The strongest signal in both RNA and

protein expression was the upregulation of oxidative phosphorylation pathways in winter

juveniles, including 21 genes encoding mitochondrial complex I subunit proteins (NDUFs) and

the succinate dehydrogenase gene, *SDHB*. Hibernators conserve energy in part through reduced

liver mitochondrial oxidation in response to cold temperature (24,25), decreasing SDH activity

(26,27) that in turn impairs cellular respiration as they lower body temperature and metabolic

rate. In contrast, winter juvenile shrews show upregulated oxidative phosphorylation pathways

compared to summer juveniles. While Dehnel's phenomenon reduces resting metabolic rate in

winter (12), shrews remain active and do not reduce their body temperature as hibernators do.

Mitochondrial regulation in shrews is opposite to that of hibernators and may use known

temperature-sensitive oxidative phosphorylation pathways to improve mitochondrial efficiency,

enhance metabolic plasticity, or maintain internal temperatures despite the reduction in metabolic

rate and body size. This hypothesis can be further tested by examining mitochondrial oxygen

consumption in the liver throughout Dehnel's phenomenon.

During autumn and winter, shrews appeared to meet their energetic needs by activating responses

similar to those triggered by starvation. When blood sugar is low, mammals promote

gluconeogenesis, a process in which the liver produces free glucose from non-carbohydrate

substrates to avoid cellular starvation (19). This process has previously been linked to seasonal

size change in shrews (10). Glucose-6-phosphatase (G6P) enzyme levels peak during autumn and

decline through spring, suggesting gluconeogenesis is highest in autumn and early winter when

snow is too scarce to offer insulation, increasing the need for glucose to prevent starvation. We

discovered an upregulation of gluconeogenesis pathways in winter juveniles, including the

overexpression of aldehyde dehydrogenases (*ALDH1B1* and *ALDH3A2*) and *G6PC1*. *G6PC1*

encodes one of three subunits of G6PC, a key gluconeogenesis enzyme in the final step of

gluconeogenesis, which catalyzes the formation of D-glucose that can then be released into the

blood. We propose the upregulation of this gene, along with others in the pathway, regulates an

increase in gluconeogenesis (10). Alongside elevated lipid metabolism and oxidative

phosphorylation, these findings indicate shrews maximize all available energy resources during

winter (Fig 4).

Fig 4. Summary of dynamic changes in shrew metabolism associated with seasonal size

change, with proposed regulation of Dehnel's cycle. Depleted lipid metabolites in the blood

and increased RNA expression of fatty acid metabolism genes highlights the importance of rapid

fat turnover in winter juveniles. FOXO signaling regulates gluconeogenesis, which is also

upregulated in winter juveniles, providing another energy source. Both FOXO signaling and fatty

acid metabolism promote changes in oxidative phosphorylation, found in both transcriptomic and

proteomic analyses, and have large effects on mitochondrial respiration in the liver.

Large changes in mRNA and proteins only weakly supported one another when comparing winter

juveniles to spring adults, but changes in gene networks across all seasons parallel regrowth. We

identified a large network negatively correlated with liver size and enriched with genes involved

in FOXO and insulin signaling. Central to this network is the forkhead box transcription factor 1

(*FOXO1*), upregulated during shrinkage and downregulated during growth. In mice, FOXO1

stimulates gluconeogenesis that promotes hepatic glucose production and release (28–30), which

can subsequently impede processes like lipid metabolism in response to fasting glucose levels

(29–32). In shrinking and regrowing shrews, FOXO signaling may thus regulate organismal

energy homeostasis and size by simultaneously regulating both glucose and lipid use to meet

shifting energy demands.

Increased FOXO signaling may, however, incur costs to shrew longevity. In model organisms

including *C. elegans*, *Drosophila*, mice and humans, FOXO1 overexpression can reduce size and

extend lifespan, while inhibition leads to aging and senescence (33–36). Compared to other

mammals, *S. araneus* evolved Dehnel's phenomenon, a rare size plasticity (1–3), as well as

extreme and disproportionately high metabolic rate (37) and short lifespan (38). FOXO1 cycling

could explain this unique combination of traits. Should findings from model organisms apply in

the shrew, FOXO1-dependent glucose utilization regulates shrinking reducing mortality during

harsh winters but undermines shrew longevity when reversed in spring as a terminal investment

for growth and reproduction. Genes within this network are therefore key candidates for

exploring both the regulatory mechanisms of size change in Dehnel's phenomenon and metabolic

change correlated with lifespan in mammals.

Despite valuable insights, there are limitations and questions outside the scope of this study.

Metabolic and regulatory change was extensively characterized throughout seasonal size changes,

but functional validation of current hypotheses is needed. For example, future studies could

analyze mitochondrial respiration to detect changes in seasonal efficiency or probe blood glucose

levels. Seasonal size change occurs in other shrew organs, and several mechanisms proposed here

involve interorgan communication, including communication between the hypothalamus and liver

through the endocrine system. Thus, it is important to investigate other tissues, such as adipocytes

in response to changes in lipid turnover, or the brain, which is rich in fats and experiences

extraordinary size plasticity, particularly in the hippocampus and cortex. Additionally, we could

investigate proposed regulatory mechanisms of longevity by examining if shrews that do not

undergo Dehnel's phenomenon, and if, lacking FOXO signaling, they exhibit extended lifespans.

Such future tests could reveal insights into natural neurodegenerative processes, their links to

metabolic dysfunction and longevity, and provide targets for potential treatments for aging and

neurodegeneration.

Conclusion

Dehnel's phenomenon was known to decrease energy spent on large and energy expensive tissue

(8,13,37,39), but associated metabolic and regulatory shifts were unknown. Analyzing changes to

the shrew liver and blood plasma, we discovered a decrease in polyunsaturated fatty acid

metabolites and altered oxidative phosphorylation pathways, potentially related to mitochondrial

respiration, alongside upregulated lipid and glucose metabolism during winter shrinkage. These

changes distinguish Dehnel's phenomenon from other wintering strategies. As in hibernation,

Dehnel's phenomenon alters mitochondrial function and relies upon increased lipid metabolism

to meet winter energetic demands, but rapidly depletes fats, supplementing energetic demands

with gluconeogenesis. These metabolic fluctuations and subsequent size change seem to be

regulated by FOXO and insulin signaling pathways that regulate both size and lifespan in several

model organisms (33–36). Along with key genes (*ACOX1*, *SDHB*, NDUFs), these pathways are

strong candidate regulators for Dehnel's phenomenon and link metabolism, size, and lifespan in

mammals.

Material and Methods

Sample Collection and RNA Sequencing

Eurasian common shrews were trapped using insulated wooden traps in Radolfzell, Germany, at

five different stages of development: large-brained summer juveniles (n=5), shrinking autumn

juveniles (n=4), small winter juveniles (n=5), regrowing spring adults (n=5), and regrown

summer adults (n=5). Shrews were aged and sexed based on tooth wear, fur appearance, and

gonad development prior to euthanasia. Fewer females than males were sampled in juvenile

stages (n=6 females). At that age, shrews are prepubescent, not sexually dimorphic, and only

minor sex dimorphism was detected in the size change of representative brain regions in a

previous study (Lázaro et al. 2018). Female under sampling is unlikely to influence gene

expression, but sex was used as a covariate in subsequent models, accounting for this skew.

Blood was collected prior to perfusion through the heart, let stand for 15 minutes, spun down at

1200 rpm, the serum pipetted off and then stored at -80°C. Shrews were perfused under

anesthesia via the vascular system with PAXgene Tissue Fixative. Livers were removed and

weighed to confirm the occurrence of DP and then placed immediately into PAXgene Tissue

Stabilizer. All samples were placed in stabilizer (2-24 hours after extraction) and then stored in

liquid nitrogen. RNA was extracted with a modified Qiagen Micro RNAeasy protocol used for

small amounts of mammalian tissue (40). RNA was sent to Genewiz for quality control, library

preparation, and sequencing. Quantity of RNA was measured with a Nanodrop spectrophotometer

and quality of RNA assessed with RNA ScreenTape. Libraries were prepared with standard

polyA selection and sequenced as 150bp pair-end reads for a target of 15-25 million reads per

sample. Body and liver weight were tested for significant change in mass using a t-test between

seasons.

Metabolomics

Metabolite analysis was carried out by MS-Omics using a Thermo Scientific Vanquish LC

(UPLC) coupled to Thermo Q Exactive HF MS. Identification of compounds were performed at

four levels; Level 1: identification by retention times (compared against in-house authentic

standards), accurate mass (with an accepted deviation of 3ppm), and MS/MS spectra, Level 2a:

identification by retention times (compared against in-house authentic standards), accurate mass

(with an accepted deviation of 3 ppm). Level 2b: identification by accurate mass (with an

accepted deviation of 3ppm), and MS/MS spectra. MetaboAnalyst was used to test for changes in

concentration between seasons (41). Concentrations for each sample were normalized by the

average concentration of summer adult samples, log transformed and auto scaled. A one-way

ANOVA was used to test for any significant seasonal difference in metabolite concentration, with

p-values corrected with a false discovery rate of 0.05. Within each significant metabolite, t-tests

were used to identify pairwise differences in mean between each season. Samples were

hierarchically clustered using significant genes and visualized using a heatmap.

Differential Gene Expression Analysis

Tests for differential gene expression were conducted to identify genes whose expression changes

during Dehnel's phenomenon. We trimmed adapters and filtered reads using fastp (42) and

removed samples with extreme read counts (~10-fold). Gene counts were quantified by pseudo

aligning to the 19,296 protein coding genes of the *S. araneus* genome (sorAra2;

GCF_000181275.1) using Kallisto (43). Counts were normalized for each tissue using DESeq2

median of ratios (44). In the liver, summer juveniles and winter juveniles were compared to infer

regulatory processes underpinning shrinkage, while winter juveniles were tested against spring

adults to explore processes underlying regrowth. Normalized counts for each gene were fitted

with a negative binomial generalized linear model and then we used a Wald test to examine the

statistical association between the log-fold change and the season parameter, with p-values

corrected using the Benjamini and Hochberg procedure (45). We then identified significantly

enriched pathways with differentially expressed genes using a ranked gene set enrichment of

KEGG pathways using fgsea (46).

Weighted Gene Co-expression Network Analysis

We used the WGCNA package (47) package to create co-expression networks for the liver and

hippocampus and determined if these correlations between genes are significantly correlated with

our shrinkage and regrowth phenotypes. We calculated the Pearson correlation for expression

between each gene pair, which was then transformed to an adjacency matrix using a soft-

thresholded power adjacency function described in Langfelder and Horvath (47). The power

function, which reduced the spurious connectivity between genes from noise by increasing

sensitivity, was determined by maximizing the scale-free topology criterion (maximizing model

fit while saturating mean connectivity). The resulting adjacency matrix was then used to calculate

a topological overlap matrix (dissimilarity matrix), in which dissimilarity distances grouped genes

into modules using fuzzy cmeans hierarchical clustering. We removed any module with fewer

than 25 genes, and merged gene modules that were found to have similar eigengene values. Then,

eigengene significance between the correlation of liver size to each module's eigengene value

was calculated. Cytoscape (48) was used to visualize significant networks and calculate network

statistics such as node (gene) connectivity. Gene set enrichment of each module was analyzed

using DAVID Gene Functional Classification Tool to test for GO enrichment (49).

Proteomics

Liver tissue from *Sorex araneus* was isolated and protein homogenates were prepared by

transferring the liver tissue to a dissection buffer containing 10% sucrose (VWR), imidazole

(Merck Millipore), EDTA (Sigma-Aldrich), Pefabloc (Sigma-Aldrich) and leupeptin (VWR). The

mixture was homogenized (T10 basic ULTRA-TURAX homogenizer IKA®) for 20 sec; then,

samples were centrifuged at 1000 g for 15 min. Supernatants were stored at -20°C. Proteomic

sample preparation was performed using the PreOmics, Cat. No.: 00027 iST 96x kit. The samples

were vacuum centrifuged overnight, and the dry peptide product was stored at -80°C until

analysis.

Peptides were resuspended in a solution containing 2% acetonitrile (ACN), 0.1% formic acid

(FA) and 0.1% TFA, and then, peptides were briefly sonicated. Five micrograms of total peptide

material were analyzed per liquid chromatography–mass spectrometry analysis. Samples were

analyzed using a UPLC-nanoESI MS/MS setup with a NanoRSLC system (Dionex, Sunnyvale,

CA, USA). The system was coupled online with an emitter for nanospray ionization (New

Objective PicoTip 360-20-10) to a Q Exactive HF mass spectrometer (Thermo Scientific,

Waltham, USA). The peptide material was loaded onto a 2-cm trapping reversed-phase Acclaim

PepMap RSLC C18 column (Dionex) and separated using an analytical 75-cm reversed-phase

Acclaim PepMap RSLC C18 column (Dionex). Both columns were kept at 60°C. The sample was

eluted with a gradient of 90% solvent A (0.1% FA, 0.1% TFA) and 10% solvent B (0.1% FA,

0.1% TFA in ACN), which was increased to 7% solvent B on a 1-min ramp gradient at a constant

flow rate of 300 nL/min. Subsequently, the gradient was raised to 30% solvent B on a 45-min

ramp gradient. The mass spectrometer was operated in positive mode, selecting up to 20

precursor ions with a mass window of m/z 1.6 based on highest intensity for higher-energy

collisional dissociation (HCD) fragmenting at a normalized collision energy of 27. Selected

precursors were dynamically excluded for fragmentation for 30 s.

A label-free relative quantitation analysis was performed using MaxQuant 1.5.7.4 software (50).

Raw files were searched against the *S. araneus* genome

(GCF_027595985.1_mSorAra2.pri_genomic.gtf). All standard settings were employed with

carbamidomethylation (C) as a static peptide modification and deamidation (NQ), oxidation (M),

formylation (N-terminal and K), and protein acetylation (N-terminal) as variable modifications.

The output contained a list of proteins identified <1% false discovery rate, and their abundances

were further filtered and processed using the Perseus v1.5.6.0 platform (51). All reverse hits that

identified proteins were removed from further analysis, and the data were log2-transformed to

approximate a normal distribution. Two or more unique peptides were required for protein

quantitation. Additionally, a non-zero quantitation value in at least 70% of the samples in one of

the groups was required for the quantifiable proteins.

We inferred potential functional changes associated with differences in protein concentrations

and analyzed correlations with transcriptomic data. We identified significantly enriched pathways

of protein using a ranked gene set enrichment of KEGG pathways using fgsea (46). Additionally,

to investigate the relationship between seasonal changes in protein concentration and RNA

abundance, we performed a linear regression analysis. Differences in protein concentration were

modeled as a function of the corresponding gene expression change using the lm function in R.

Acknowledgments

We thank Joshua Rest, Krishna Veeramah, and Tanya Lama who have provided helpful feedback

on initial results and previous versions of the manuscript. We thank Michal Oklinski for help with

initial dissections.

References

1.   Dehnel A. Studies of the genus Sorex L. Ann Univ M Curie-Sklod. 1949;4(2):17–104.
2.   Pucek Z. Seasonal and age changes in the weight of internal organs of shrews. Acta
Theriol (Warsz). 1965;10:369–438.

3. Lázaro J, Hertel M, Sherwood CC, Muturi M, Dechmann DKN. Profound seasonal
changes in brain size and architecture in the common shrew. Brain Struct Funct [Internet].
2018;223(6):2823–40. Available from: http://dx.doi.org/10.1007/s00429-018-1666-5
4. Lázaro J, Dechmann DKN. Dehnel's phenomenon. Curr Biol. 2021;31(10):R463–5.
5. Pucek M. Water contents and seasonal changes of the brain-weight in shrews. Acta
Theriol (Warsz). 1965;10(1955):353–67.
6. Ray S, Li M, Koch SP, Mueller S, Boehm-Sturm P, Wang H, et al. Seasonal plasticity in
the adult somatosensory cortex. Proc Natl Acad Sci U S A. 2020;117(50):32136–44.
7. Genoud M, Isler K, Martin RD. Comparative analyses of basal rate of metabolism in
mammals : data selection does matter. Biol Rev. 2018;93:404–38.
8. Keicher L, O'Mara MT, Voigt CC, Dechmann DKN. Stable carbon isotopes in breath
reveal fast metabolic incorporation rates and seasonally variable but rapid fat turnover in
the common shrew (Sorex araneus). J Exp Biol. 2017;220(15):2834–41.
9. Churchfield S, Rychlik L, Taylor JRE. Food resources and foraging habits of the common
shrew, Sorex araneus: Does winter food shortage explain Dehnel's phenomenon? Oikos.
2012;121(10):1593–602.
10. Hyvärinen H. Wintering strategy of voles and shrews in Findland. Winter Ecol Small
Mamm. 1984;10:139–48.
11. Lázaro J, Hertel M, Muturi M, Dechmann DKN. Seasonal reversible size changes in the
braincase and mass of common shrews are flexibly modified by environmental conditions.
Sci Rep. 2019;9(1):1–10.
12. Taylor JRE, Rychlik L, Churchfield S. Winter reduction in body mass in a very small,
nonhibernating mammal: Consequences for heat loss and metabolic rates. Physiol
Biochem Zool. 2013;86(1):9–18.
13. Schaeffer PJ, O'Mara MT, Breiholz J, Keicher L, Lázaro J, Muturi M, et al. Metabolic
rate in common shrews is unaffected by temperature, leading to lower energetic costs
through seasonal size reduction. R Soc Open Sci. 2020;7(4).
14. Auteri GG. A conceptual framework to integrate cold-survival strategies: Torpor,
resistance and seasonal migration. Biol Lett. 2022;18(5):1–6.
15. Boyer BB, Barnes BM. Molecular and metabolic aspects of mammalian hibernation.
Bioscience. 1999;49(9):713–24.
16. Chazarin B, Storey KB, Ziemianin A, Chanon S, Plumel M, Chery I, et al. Metabolic
reprogramming involving glycolysis in the hibernating brown bear skeletal muscle. Front
Zool. 2019;16(1):1–21.
17. Faherty SL, Luis Villanueva-Cañas J, Klopfer PH, Albà MM, Yoder AD. Gene expression
profiling in the hibernating primate, Cheirogaleus medius. Genome Biol Evol.
2016;8(8):2413–26.
18. Villanueva-Cañas JL, Faherty SL, Yoder AD, Albà MM. Comparative genomics of
mammalian hibernators using gene networks. Integr Comp Biol. 2014;54(3):452–62.
19. Rui L. Energy Metabolism in the Liver Liangyou Rui. Physiol Behav. 2014;176(5):139–
48.
20. Lu D, He A, Tan M, Mrad M, El Daibani A, Hu D, et al. Liver ACOX1 regulates levels of
circulating lipids that promote metabolic health through adipose remodeling. Nat
Commun. 2024;15(1).
21. Elaine Epperson L, Karimpour-Fard A, Hunter LE, Martin SL. Metabolic cycles in a
circannual hibernator. Physiol Genomics. 2011;43(13):799–807.
22. Drew KL, Rice ME, Kuhn TB, Smith MA. Neuroprotective adaptations in hibernation:
Therapeutic implications for ischemia-reperfusion, traumatic brain injury and
neurodegenerative diseases. Free Radic Biol Med. 2001;31(5):563–73.

23.   Rice SA, Mikes M, Bibus D, Berdyshev E, Reisz JA, Gehrke S, et al. Omega 3 fatty acids
stimulate thermogenesis during torpor in the Arctic Ground Squirrel. Sci Rep [Internet].
2021;11(1):1–14. Available from: https://doi.org/10.1038/s41598-020-78763-8

24.   Chaffee RR, Hoch FL, Lyman CP. Mitochondrial oxidative enzymes and
phosphorylations in cold exposure and hibernation. Am J Physiol. 1961;201(1):29–32.

25.   Staples JF, Brown JCL. Mitochondrial metabolism in hibernation and daily torpor: A
review. J Comp Physiol B Biochem Syst Environ Physiol. 2008;178(7):811–27.

26.   Armstrong C, Staples JF. The role of succinate dehydrogenase and oxaloacetate in
metabolic suppression during hibernation and arousal. J Comp Physiol B Biochem Syst
Environ Physiol. 2010;180(5):775–83.

27.   Gehnrich SC, Aprille JR. Hepatic gluconeogenesis and mitochondrial function during
hibernation. Comp Biochem Physiol -- Part B Biochem. 1988;91(1):11–6.

28.   Matsumoto M, Pocai A, Rossetti L, DePinho RA, Accili D. Impaired Regulation of
Hepatic Glucose Production in Mice Lacking the Forkhead Transcription Factor Foxo1 in
Liver. Cell Metab. 2007;6(3):208–16.

29.   Xiong X, Tao R, DePinho RA, Dong XC. Deletion of Hepatic FoxO1/3/4 Genes in Mice
Significantly Impacts on Glucose Metabolism through Downregulation of
Gluconeogenesis and Upregulation of Glycolysis. PLoS One. 2013;8(8):1–11.

30.   Zhang W, Patil S, Chauhan B, Guo S, Powell DR, Le J, et al. FoxO1 regulates multiple
metabolic pathways in the liver effects on gluconeogenic, glycolytic, and lipogenic gene
expression. J Biol Chem [Internet]. 2006;281(15):10105–17. Available from:
http://dx.doi.org/10.1074/jbc.M600272200

31.   Tikhanovich I, Cox J, Weinman SA. Forkhead box class O transcription factors in liver
function and disease. J Gastroenterol Hepatol. 2013;28(1):125–31.

32.   Yang Z, Roth K, Agarwal M, Liu W, Petriello MC. The transcription factors CREBH,
PPARa, and FOXO1 as critical hepatic mediators of diet-induced metabolic dysregulation.
J Nutr Biochem [Internet]. 2021;95:108633. Available from:
https://doi.org/10.1016/j.jnutbio.2021.108633

33.   Gami MS, Wolkow CA. Studies of Caenorhabditis elegans DAF-2/insulin signaling reveal
targets for pharmacological manipulation of lifespan. Aging Cell. 2006;5(1):31–7.

34.   Hwangbo DS, Garsham B, Tu MP, Palmer M, Tatar M. Drosophila dFOXO controls
lifespan and regulates insulin signalling in brain and fat body. Nature.
2004;429(6991):562–6.

35.   Katic M, Kahn CR. The role of insulin and IGF-1 signaling in longevity. Cell Mol Life
Sci. 2005;62(3):320–43.

36.   Satoh A, Brace CS, Rensing N, Cliften P, Wozniak DF, Herzog ED, et al. Sirt1 extends
life span and delays aging in mice through the regulation of Nk2 Homeobox 1 in the DMH
and LH. Cell Metab [Internet]. 2013;18(3):416–30. Available from:
http://dx.doi.org/10.1016/j.cmet.2013.07.013

37.   Taylor JRE. Evolution of Energetic Strategies in Shrews. In: Evolution of Shrews. 1998.
p. 309–46.

38.   Healy K, Guillerme T, Finlay S, Kane A, Kelly SBA, McClean D, et al. Ecology and
mode-of-life explain lifespan variation in birds and mammals. Proc R Soc B. 2014;281.

39.   Pucek Z. Seasonal and age change in shrews as an adaptive process. Symp zool Soc Lond.
1970;26:189–207.

40.   Yohe LR, Davies KTJ, Simmons NB, Sears KE, Dumont ER, Rossiter SJ, et al.
Evaluating the performance of targeted sequence capture, RNA-Seq, and degenerate-
primer PCR cloning for sequencing the largest mammalian multigene family. Mol Ecol
Resour. 2020;20(1):140–53.

41. Pang Z, Chong J, Zhou G, De Lima Morais DA, Chang L, Barrette M, et al.
MetaboAnalyst 5.0: Narrowing the gap between raw spectra and functional insights.
Nucleic Acids Res. 2021;49:388–96.
42. Chen S, Zhou Y, Chen Y, Gu J. Fastp: An ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics. 2018;34(17):i884–90.
43. Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq
quantification. Nat Biotechnol. 2016;34(5):525–7.
44. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol. 2014;15(550):1–21.
45. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate : A Practical and
Powerful Approach to Multiple Testing. J R Stat Soc. 1995;57(1):289–300.
46. Korotkevich G, Sukhov V, Budin N, Atryomov MN, Sergushichev A. Fast gene set
enrichment analysis. bioRxiv. bioRxiv. 2021;
47. Langfelder P, Horvath S. WGCNA: An R package for weighted correlation network
analysis. BMC Bioinformatics. 2008;9.
48. Shannon P, Markiel A, Ozier O, Nitin SB, Wang JT, Ramage D, et al. Cytoscape: A
Software Environment for Integrated Models. Genome Res [Internet]. 2003;13(22):2498–
504. Available from: http://ci.nii.ac.jp/naid/110001910481/
49. Huang DW, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene
lists using DAVID bioinformatics resources. Nat Protoc. 2009;4(1):44–57.
50. Tyanova S, Temu T, Cox J. The MaxQuant computational platform for mass
spectrometry-based shotgun proteomics. Nat Protoc [Internet]. 2016;11(12):2301–19.
Available from: http://dx.doi.org/10.1038/nprot.2016.136
51. Tyanova S, Temu T, Sinitcyn P, Carlson A, Hein MY, Geiger T, et al. The Perseus
computational platform for comprehensive analysis of (prote)omics data. Nat Methods.
2016;13(9):731–40.
620

Supporting Information

Data Availability

Further details of experimental protocols are described in the Expanded View. Data openly

available on Dryad https://doi.org/10.5061/dryad.pc866t1w3 with reproducible code found on

GitHub: https://github.com/wrthomas315/Dehnels_Seasonal_RNAseq2/tree/main/data. Primary

sequencing was deposited on the NCBI SRA, Bioproject PRJNA941271.

Author Contributions and Funding Disclosure

LD, DD, AC, and JN conceived and funded project. CB, MM, and DD collected, measured, and

sampled shrews. WRT designed and conducted genomic analyses. WRT, JHJ, JN, and AC

analyzed metabolite concentrations. WRT visualized data and wrote initial draft. LD, DD, JN,

DC, and AC contributed to review and editing of draft. All authors contributed to data

interpretation. WRT and research were supported by the Human Frontiers in Science Program

Award (RGP0013/2019) to DD, LMD, and JN, and by the Stony Brook University Presidential

Innovation and Excellence Fund to LMD.

Competing Interests

The authors declare that they have no conflict of interest.