Molecular mechanisms of seasonal brain shrinkage and regrowth in *Sorex araneus*

William R. Thomas[1]*†, Dina K. N. Dechmann[2,3], John Nieland[4], Cecilia Baldoni[2,3], David Carlson[1], Dominik von Elverfeldt[5], Julie Holm-Jacobsen[4], Marion Muturi[2], Angelique Corthals[6]‡, Liliana M. Dávalos[1,7]‡*

[1]Department of Ecology and Evolution, Stony Brook University; Stony Brook, NY 11794, United States.

[2]Max-Planck Institute for Animal Behavior; Radolfzell, 78315, Germany.

[3]University of Konstanz; Konstanz, 78464, Germany.

[4]Health Science and Technology, Aalborg University; Aalborg, 9100, Denmark.

[5]Division of Medical Physics, Department of Diagnostic and Interventional Radiology, University Medical Center Freiburg, Faculty of Medicine, University of Freiburg, Freiburg, Germany.

[6]John Jay College of Criminal Justice; New York, NY 10019, United States.

[7]Consortium for Inter-Disciplinary Environmental Research, Stony Brook University; Stony Brook, NY 11794, United States.

† Lead author

‡ These authors contributed equally to this work

* Corresponding authors: William R. Thomas and Liliana M. Dávalos

Emails: William.thomas@stonybrook.edu, Liliana.Davalos@stonybrook.edu

Author Contributions: LD, DD, AC, and JN conceived and funded project. CB, MM, and DD collected, measured, and sampled shrews. WRT designed and conducted genomic analyses. WRT, JHJ, JN, and AC analyzed metabolite concentrations. WRT visualized data and wrote initial draft. LD, DD, JN, DC, and AC contributed to review and editing of draft. All authors contributed to data interpretation.

Competing Interest Statement: Authors disclose no competing interests.

Keywords: Dehnel's phenomenon, neurodegeneration, winter, metabolism, shrew

Abstract
Mammalian brains typically grow through development. In contrast, the Eurasian common shrew *Sorex araneus* has evolved a wintering strategy that shrinks many organs in autumn, including their brain, then rapidly regrow in spring. To explore the molecular underpinnings of this unique brain size change, we analyzed multi-organ, season-specific transcriptomics and metabolomic data. Simultaneous with brain shrinkage, we discovered system-wide metabolic shifts from lipid to glucose metabolism, similar to hypoxia responses, as well as neuroprotection of brain metabolic homeostasis through reduced cholesterol efflux. Analyses of co-expression models uncovered a finely tuned brain-liver crosstalk underlying these shifts, with the brain shrinkage module comprising of markers associated with longevity-regulating pathways and pathways involved in neurodegeneration, including Parkinson's disease and Huntington's disease. We propose metabolic shifts are central to seasonal brain size changes in *S. araneus*, with potential implications for mammalian neurodegeneration and neuroregeneration.

Main Text
Introduction
Animals vary widely in their wintering adaptations. The most common ones are hibernation or migration, but a handful of species that cannot migrate or slow their metabolism have evolved wintering size plasticity: Dehnel's phenomenon (DP) (*1*, *2*). This pattern of growth, best studied in the Eurasian common shrew, *Sorex araneus*, is unlike that of almost every other mammal. Instead of growing continuously to adulthood, juvenile *S. araneus* grow to an initial maximum size in the first summer of their disproportionately brief life (~1 year). Shrews then shrink, reaching a nadir in winter, followed by rapid regrowth to their adult, breeding size in spring. This size plasticity also occurs in the brain, with a 26% volume decrease through winter and in contrast with typical unidirectional brain growth in mammals (*2*). In spring, the brain then partially regrows. Although DP has garnered much interest because of its potential for regenerating organs, especially the brain, the molecular underpinnings of this wintering adaptation remain largely unknown (*3*).

Dehnel's phenomenon corresponds to the unique physiological constraints shrews face. *S. araneus* have one of the highest mammalian metabolic rates measured to date (*4*), requiring high and constant food intake throughout the year. Research on shrew physiology has shown fat turnover in *S. araneus* is constantly high, with 50% turnover rates increasing from 4.5 hour in summer juveniles to 2.5 hours in winter (*5*). Yet, the amount of stored lipids in the liver is also highest during the winter months (*6*). Meanwhile, energy consumption per unit mass remains constant year-round (*7*). Given these high energy demands and high lipid turnover, DP is construed as an adaptation that compensates for the scarcity of resources in winter by reducing body size, particularly of energy-expensive tissue such as the brain (*5*, *8*). In humans, the brain accounts for 20% of glucose metabolized (*9*), and contains 25% of the unesterified cholesterol found in the body (*10*). But this high metabolic demand also makes the human brain vulnerable: metabolic changes have been implicated in the onset of human neurodegeneration (*11*–*13*). Therefore, we hypothesized that molecular changes underlying seasonal size change in shrew size may parallel those previously identified implicated in neurodegenerative diseases.

Changes in glucose and lipid metabolism have also been identified in hibernation (*14*, *15*), and Dehnel's phenomenon could share some of those regulatory mechanisms, which include reduced feeding, thermoregulation and hypoxia. Mechanistic analyses of mammalian hibernation have identified a toolkit of genes that regulates vast metabolic changes (*14*–*16*) associated with metabolic pathways, circadian rhythm, thermogenesis, and feeding behavior (*17*). For example, research across different species of hibernators has identified increased fatty acid oxidation simultaneous with downregulation of genes associated with the breakdown of glucose as a molecular metabolic switch underlying winter energy savings (*16*). Notably, these metabolic changes can have profound negative effects on the brain, specifically hypoxia (*18*). Thus, evolved neuroprotective responses to mitigate the harmful effects of metabolic fluctuations have been discovered in hibernating mammals (*19*, *20*). In the Arctic ground squirrel, the negative consequences of oxidative stress and glucose deprivation through hibernation are diminished by modifying neural ion channels (*21*). But *S. araneus* remain awake throughout the winter and thus experience a changing and metabolically demanding environment. To enable winter survival while maintaining activity, shrews are therefore likely to exhibit different regulatory patterns for metabolic functions suppressed in hibernators.

Metabolic changes associated with hibernation also show similarities to the metabolic dysregulation found in human neurodegenerative diseases. Both high fat and high sucrose diets have negative consequences on memory, mood, and other cognitive processes, akin to changes in lipid and glucose utilization under hibernation (*22*). Rats fed diets high in saturated fats and sugars perform worse in place recognition tests and exhibit higher oxidative stress and inflammation markers in the hippocampus (*23*, *24*). This occurs despite the protective functions of the blood-brain barrier (BBB), a semipermeable system of blood vessels that regulate the

entrance of molecules into the brain (*25*), insulating the brain from metabolic changes. In
humans, BBB permeability increases with age, leading to diminished protection from metabolic
shifts (*26*) and neurodegenerative diseases (*27*, *28*). Metabolic molecules that can cross the
BBB, such as insulin and glucose (*29*), can thus have large effects on the brain. Circulating
insulin and glucose increase in diseases such as type-2 diabetes and obesity, is associated with
reduced brain size, motor activity, functional connectivity, and increased risk of developing
Alzheimer's and Parkinson's diseases (*11–13*, *30*). As in both hibernation and human
neurodegenerative disease, brain shrinkage and regrowth phenotypes may therefore be
associated with metabolic shifts and transport of circulating molecules to the brain through the
BBB.

To explore the metabolic hypothesis and its potential parallels, we collected phenotypic,
metabolomic, and transcriptomic data from a shrew population across seasons, examining gene
expression dynamics change throughout the entirety of DP size plasticity. We focused on: 1)
blood, which circulates metabolites across the organisms, 2) liver, where many functions that
regulate metabolism take place, including glucose processing, and 3) brain, where the
anticipatory and predictable changes in volume occur. Our results show that shrews survive the
winter through processes that in model organisms are pathological, while avoiding detrimental
health effects. Our analyses reveal an unequivocal metabolic basis for DP, with implications for
metabolic- and age-related human neurodegeneration.
110
Results
Confirming Dehnel's phenomenon in the German shrew population
Comparisons of body and organ mass throughout the cycle confirmed patterns of size changes in
the brain and spleen, and similar but slightly earlier growth in the liver. (Figure 1B/2A, full results
shown in Supplemental Data). Analyzing the 24 shrews collected for all stages of DP (large
brained Summer Juvenile=5; shrinking Autumn Juvenile n=4; small brained Winter Juvenile n=5;
regrowing Spring Adult n=5; regrown Summer Adult n=5), we found body mass shrank as
expected between Summer Juveniles and Winter Juveniles ($BM_{WiJ-SuJ}$=-1.00g, $p_{WiJ-SuJ}$<0.028),
with regrowth as they mature to Spring Adults ($BM_{SpA-WiJ}$=3.724, $p_{SpA-WiJ}$<0.0001). Shrew liver
mass (Figure 1) reached a minimum as Autumn Juveniles ($LiM_{AuJ-SuJ}$=-94.1mg/g of BM,
$p_{AuJ-SuJ}$=0.0949). Although regrowth began slightly early as Winter Juveniles ($LiM_{WiJ-AuJ}$=70.4/g
of BM, $p_{WiJ-AuJ}$=1.0), the significant portion of regrowth still occurred between the Winter Juvenile
and Spring Adult phases ($LiM_{SpA-AuJ}$=607.0mg/g of BM, $p_{SpA-WiJ}$=0.001). Brain (Figure 2) and
spleen mass followed the same pattern, as Summer Juveniles reached a minimum by the Winter
Juvenile phase ($BrM_{WiJ-SuJ}$=-54.4mg, $p_{WiJ-SuJ}$=0.048; $SpM_{WiJ-SuJ}$=-53.1mg/g of BM,
$p_{WiJ-SuJ}$<0.001), with regrowth into Spring Adults ($BrM_{SpA-WiJ}$=71.2mg, $p_{SpA-WiJ}$=.297,
$SpM_{SpA-WiJ}$=63.7mg/g of BM, $p_{SpA-WiJ}$<0.01), although not significantly in the brain. Overall, these
size changes validated previous research and are further used to guide transcriptomic
characterization of DP.

Reversible metabolic shifts from lipid metabolism to gluconeogenesis during shrinkage
We characterized the plasma metabolome to quantify metabolites throughout the cycle and
evaluated the potential shifts in metabolism during shrinkage and regrowth. Of 250 metabolites
validated by multiple detection methods in all seasons, we found 28 with significantly different
concentrations varying by stage of DP (p<0.05, Figure 1A). Of these differentially concentrated
metabolites, three were lipid metabolites; arachidonic acid (p=0.04), ethyl palmitoleate (p=0.02),
and docosahexanoic acid (p=0.04). Compared to both Summer Juveniles and Adults, these three
lipid metabolites were generally lowest in Autumn Juveniles, Winter Juveniles, and Spring Adults.
This could indicate either limited lipid resource availability as size change occurs and, potentially,
suppressed lipid metabolism, or lipids metabolized to generate energy are consumed faster than
being generated.
141

To determine if changes in the metabolome were the result of the regulatory effects of gene
expression on size change and suppressed lipid metabolism, we analyzed variation across five
seasons and quantified differential expression of the metabolically active liver. Instead of resulting
from ontogenetic or gradual changes of expression during the shrew lifespan, most of the liver
differentially expressed genes (Figure 1C) fell into two temporal clusters, 8 and 12 (Figure 1D),
corresponding to genes transiently up- and downregulated during the wintering cycle. Several key
transcription factors (*31, 32*) were differentially expressed mirroring liver size changes, including
PPARs, CREBPs, RXRs, and FOXOs. While FOXO1 and FOXO3 promote gluconeogenesis and
are reversibly upregulated in Autumn Juveniles, PPARA, PPARD, RXRA, and RXRG are both
reversibly downregulated and associated with lipid metabolism (Supplemental Table 1, Figure
1E). All the mentioned liver nutrient sensing transcription factors and the genes they control are
differentially expressed mirroring size changes (Supplemental Table 1), indicating large seasonal
shifts in metabolism associated with brain size plasticity. We also identified a single clock gene,
PER1, reversibly upregulated in Autumn Juveniles, and the upregulation of genes associated with
endocrine signaling from the liver to the blood (POMC, VDR, NRG1, FGFR4, FGF21).
Differentially expressed liver nutrient sensing, clock transcription factors, and endocrine genes
are known to control gene expression and regulate metabolism, indicating a switch from lipid to
glucose fuel during autumnal shrinkage and initial decline of circulating lipid availability. This
switch parallels mechanisms implicated in evolutionary adaptations to wintering and hypoxia, as
well as hypothesized metabolic dysregulation associated with neurodegeneration (*12, 33*).
162
Differential expression analyses focused on genes of high effect, but large modules comprising
genes with small effect could also play a pivotal role in regulating seasonal metabolic changes.
We characterized interactions in expression among genes and identified modules of co-regulated
liver-expressed genes related to seasonal changes in body and liver size using weighted gene
co-expression network analysis (WGCNA). We identified 37 modules (Supplemental Figure 2A),
of which ten are significantly associated with growth or size changes (Figure 3A). Among these,
one module, cyan, encompassed 2,816 genes and correlated with all reversible body and liver
size changes. Unlike other modules also significantly associated with summer controls, co-
expression in the cyan module is exclusively related to size changes. Therefore, genes in the
cyan module are associated with the cyclic shifts defining DP through transient activation and
deactivation. In this module, we identified 76 significantly enriched pathways that together form
16 pathway clusters (Figure 3B, Supplemental Data). The most enriched of these pathway
clusters comprises four pathways (Figure 3C): insulin resistance, longevity regulating pathway,
AMPK signaling pathway, and longevity regulating pathway (across multiple species); and
included transcription factors (PPARA, FOXO1, CREB3L2, CREB3L3), their downstream targets
(G6PC), PPARG coactivators (PPARGC1A, PPARGC1B), and regulators of inflammation
(HNF4A, NFKB1). Therefore, metabolic and longevity-related co-expression in the shrew is
associated with these transcription factors, which are also involved in metabolic and age-related
progression of neurodegeneration in model organisms.
182
Decreased brain cholesterol efflux, signatures of neurodegeneration, and a brain-liver axis
We evaluated the effect of metabolic shifts on the shrew brain, which may drive observed size
plasticity, while also coordinating seasonal changes in metabolism. First, to disentangle markers
of shrinkage from those associated with ontogeny, we compared gene expression across stages
in the hippocampus, which shrinks then regrows, and the cortex, which shrinks massively, but
minimally regrows. In the hippocampus, we found 266 differentially expressed genes, only 25 of
which were of high effect (Figure 2C). In contrast, the cortex showed 540 differentially expressed
genes, with 96 downregulated and 65 upregulated genes of high effect at the Winter Juvenile
phase (Figure 2C). Temporal gene clustering analyses revealed fewer distinct clusters in the
hippocampus than the liver and cortex, but, as with the liver, most differentially expressed genes
are in Clusters 3 and 11, showing transient up and down regulation consistent with DP
(Supplemental Figure 4). For the cortex, most of the high effect differentially expressed genes
grouped in Clusters 1 and 11, with genes in Cluster 1 showing a gradual increase in expression

through the life cycle and genes in Cluster 11 being permanently downregulated between the Autumn Juvenile and Winter Juvenile periods. Despite less variation in gene expression between stages in the brain than in the liver, we identified unique brain-region-specific responses to DP and gene expression patterns consistent with both ontogenetic brain development and DP, especially in the hippocampus.

Testing for differential expression during size change in the hippocampus and cortex, we found few overlapping differentially expressed genes, but both tissue-specific gene sets were enriched for the cholesterol metabolism pathway (Figure 2D), including the apolipoproteins APOA1 and APOH (Figure 2B). In each tissue, the cholesterol metabolism pathway shows a similar trend: cholesterol metabolism upregulated during initial growth and regrowth in the hippocampus, and downregulated during cortex shrinkage, indicating this pathway's crucial role in both shrinkage and regrowth. Gene co-expression network analyses of the hippocampus also revealed 34 modules (Supplemental Figure 3), three of which correlated with brain shrinkage (Figure 4A). The most significant of these modules (grey) comprised 346 genes, and is associated with both metabolic processes and glycolysis, as well as enriched with genes —such as CASP3, PPARG, CREB3L2, CALM3— in pathways known to underlie neurodegenerative pathogenesis including pathways of neurodegeneration, Parkinson's disease, and Huntington's disease (Figure 4C). Although centered on the key glycolytic enzyme ALDOA (Figure 4B), the regulator of lipid metabolism and transcription factor PPARG is also central to this module. While gene expression patterns in the brain and the liver are distinct, both related through signals of reversible fluctuations in metabolic energy sources. Evidence of both a switch to glucose metabolism in the liver and downregulation of cholesterol metabolism in the brain during shrinkage is consistent with a nadir in circulating lipid metabolites in winter. We therefore propose that shifts in the regulation of lipids in the liver and brain, mirrored by shifting metabolite concentrations in the blood, indicate a pivotal role for regulatory coordination via a brain-liver crosstalk in DP.

Discussion
For the first time we characterized a range of molecular shifts associated with brain shrinkage and regrowth in *Sorex araneus*. Previous research suggested winter size reduction reduced energy expenditure of large and energy expensive tissue (*5*, *7*, *34*, *35*), but at a cost for the brain, which experiences architectural change and cognitive decline during shrinkage (*2*, *36*). Exploring this metabolic hypothesis, stage-by-stage analysis of blood metabolites indicates similarities to hypoxic strategies instead of hibernation, while analysis of RNA brain expression during shrinkage revealed both markers of decreased cholesterol efflux and correlations with pathways of neurodegenerative disease. These changes in brain expression occurred after the body and liver showed drastic yet reversible metabolic shifts from lipid to glucose metabolism during shrinkage and regrowth. Circadian clock genes appeared to regulate shifts in liver metabolism, along with changes in hepatokine and endocrine signals that can permeate the BBB. We therefore propose a crosstalk between the brain and liver underlies metabolic and seasonal size changes in DP, despite the protective functions of the BBB that isolate brain from body metabolism. This crosstalk between brain and liver metabolism in the healthy shrew going through its annual cycle shows parallels to mechanisms of human neurodegeneration.

If shrew metabolic shifts paralleled those in hibernation, an increase in lipid metabolites would be observed during shrinkage. Instead, we found autumn, winter, and spring reductions in arachidonic acid, ethyl palmitoleate, and docosahexanoic acid (Figure 1A). Increased concentration of these lipid metabolites has been found in hibernators that shift fuel source from carbohydrates to lipids in winter including the arctic ground squirrel (*37*), hedgehog (*38*), echidna (*39*), brown bear (*15*), and hamster (*40*, *41*). By contrast, shrew metabolite variation during winter more closely resembles conserved hypoxia responses previously found in naked mole rats (*42*), yaks (*43*), and even goldfish (*44*). In hypoxic naked mole rats, stored liver glucose is released into the blood to fuel carbohydrate metabolism (*45*), as the mole rats remain active with only slight reductions in metabolic rate (*46*). Like mole rats in their fossorial environment, semi-fossorial

250   shrews stay active throughout the winter. Instead of mimicking hibernation, shrew metabolites
251   indicate a hypoxia-like shift to glucose as fuel as a lipid conservation mechanism for periods of
252   energy deficit beginning with autumn shrinkage. We argue this metabolic shift is the foundation of
253   the observed seasonal brain shrinkage.
254
While metabolomic data implicated hypoxia-like constraints on circulating lipids, how gene
regulation mediated these decreases was unresolved. By analyzing seasonal differential
expression of metabolically important liver transcription factors, we discovered a switch in autumn
shrinkage to a new primary fuel: glucose. While lipid metabolism genes (e.g., PPARA, PPARD,
RXRA, and RXRG) are downregulated, increased PER1 expression signals metabolic demands
met by glucose. Indicating this switch, two forkhead box transcription factors (FOXOs) associated
with glucose metabolism are significantly upregulated during autumn shrinkage (Figure 2C). In
mice, FOXO1 and FOXO3 stimulate gluconeogenesis that promotes hepatic glucose production,
with glucose released into the blood for energy utilization throughout the body (*47–49*). FOXO1
and FOXO3 also impede lipid metabolism in response to fasting glucose levels (*32, 47, 48, 50*).
Upregulation of these transcription factors in autumn shrews, later reversed in spring, may
regulate organismal energy homeostasis by promoting glucose utilization as mobilization of
stored lipids decreases during shrinkage, as in hypoxia.
268
FOXO upregulation maintains shrew metabolic homeostasis during challenging environmental
conditions and lipid scarcity, but this incurs costs to both shrew brain size and longevity. The liver
co-expression network (cyan) most associated with size change was enriched for genes involved
in glucose utilization and longevity associated pathways, which encompassed both *FOXO1* and
SIRT1 (Figure 3). In model organisms including *C. elegans*, *Drosophila*, mice, and humans,
FOXO1 overexpression can decrease size and extend lifespan, while inhibition leads to aging
and senescence (*51–54*). In the shrew, extreme brain size plasticity in response to FOXO1
dependent glucose utilization reduces mortality during harsh winters but may undermine shrew
longevity when reversed in spring, as a terminal investment for growth and reproduction. FOXO1
cycling could explain why compared to other mammals of similar size, *S. araneus* have high
metabolic rates (*35*) and short lifespan (*55*), both extreme and disproportionate to their body size.
Neurodegeneration is also considered an age-related neurological disorder. Enrichment of
metabolically related- and longevity regulating genes, including *FOXO1* and *SIRT1*, in this cyan
gene co-expression network further highlights interactions among metabolic shifts, size, and age-
related effects. Genes in this network are therefore candidates for potential therapeutic reversals
of brain size decline after the onset of symptoms.
285
Although metabolite and liver transcriptomics results implicated overall shrew metabolism as a
regulator of size plasticity, we sought to identify both the impacts of and effects on the brain.
Within the hippocampal gene networks (Figure 4), there was an enrichment of genes centered on
the key glycolytic enzyme ALDOA. *ALDOA* encodes an important enzyme in glycolysis that
converts dietary glucose into energy by catalyzing the conversion of fructose biphosphate into
glyceraldehyde 3 phosphate (*56, 57*). Genes in the network surrounding this hub, CASP3,
PPARG, CREB3L2, and CALM3, enrich pathways of neurodegeneration, including Huntington's
disease and Parkinson's disease. *ALDOA* is central in this network, indicating that glycolysis
regulation and shifts in glucose metabolism during shrew brain shrinkage directly regulate
previously identified human neurodegenerative markers. This result further highlights similarities
in the mechanisms of both shrew brain shrinkage induced via metabolic change and human
neurodegeneration.
298
Despite these glucose-related changes, we found the shrew's shrinking brain attempts to
maintain internal lipid homeostasis by reducing cholesterol efflux, likely as neuroprotection.
Changes in lipid metabolism can occur in the brain to compensate for energy deficits by self-
catabolizing fats (*58*), but maintenance of metabolic homeostasis is important for neural health
(*59, 60*). Therefore, self-catabolism of the shrew brain would undermine function during the winter

months. Except for RXRG, lipid metabolism transcription factors differentially expressed in the
liver during seasonal size change (PPARA, PPARB, RXRA) were *not* differentially expressed in
the shrew brain. Instead, differential expression analyses indicate limits to brain cholesterol efflux
during shrinkage via apolipoproteins (APOs; Figure 2). These signals suggest the shrew brain
protects itself from bodily lipid constraints by limiting cholesterol efflux during lipid scarcity and
high turnover in winter.
310
Lastly, we propose a brain-liver crosstalk regulates shifts in glucose and lipid metabolism, organ
size-changes, and neuroprotective changes in brain cholesterol efflux. Although the brain
communicates extrinsic environmental signals to the rest of the body, the blood brain barrier limits
the relay of environmental information back to the brain. We identified several gene expression
patterns associated with molecules that both cross this semi-permeable barrier and are likely
involved in brain size change. As circadian clock gene PER1 is downregulated, there is
upregulation of genes involved in endocrine signaling in the liver. Endocrine signaling between
the brain and liver has been shown to regulate glucose homeostasis (*61*) and is promoted by
circadian clock genes (*62*). Additionally, the large liver co-expression module of seasonal size
change (cyan) is significantly enriched with AMPK signaling genes in (cyan, Figure 3C). Across
mammals, AMPK signaling restores energy balance by transitioning between anabolic and
catabolic pathways in response to circulating glucose levels through tissue-specific cross-
modulation (*63, 64*). In shrews, liver-specific change in AMPK signaling, in tandem with circadian
and endocrine signaling, modulates both brain and liver expression changes in response to the
transition from lipids to glucose as main fuel source.
326
Coordinated changes in gene expression of the liver and brain, the proposed brain-liver axis, are
ideal candidate mechanisms of brain shrinkage. Functional relationships, or axes, as the brain
receives stimuli from peripheral tissues can induce vast changes in brain function, expression,
and phenotype. These stimuli range from inflammatory responses sent from the spleen (*65*) to
regulation of nutrient metabolism through signals from the gut microbiota (*66*). In mammals,
alterations to the brain-liver axis have been implicated in several models of neurodegenerative
disease (*67–70*). For example, in mice models of Parkinson's disease, nigrostriatal degeneration
results in hepatic mitochondrial dysfunction, causing metabolic dysregulation that self-
perpetuates initial neurodegenerative symptoms (*68*). Reversals of candidate genes and
pathways within this brain-liver axis, however, indicate these processes in the shrew are self-
perpetuating. Instead, the reversals suggest effective self-regulation that not only drives
shrinkage, but also allows brain regrowth. Expression reversals via the brain-liver axis are thus
important targets for functional tests and future therapeutic treatments.
340
Dehnel's phenomenon was known to be an energy saving adaptation that incurred some
cognitive compromise (*2*), but its molecular underpinnings were largely unknown. In shrew liver
and blood plasma, we discovered shifts from lipid to sugar metabolism during shrinkage, in
tandem with upregulated indicators of neurodegeneration in the brain. But in contrast with
mammalian models for neurodegenerative diseases, the shrew brain also enters a state of fat
conservation by seemingly limiting cholesterol efflux in the winter, the most metabolically
demanding phase of the shrew's brief life. Based on our results, we propose this combination of
sugar-dependent whole-body metabolism and limited brain cholesterol efflux is an evolved form
of neuroprotection within the constraints of the shrew's environment and life history. Our study
thus sheds light on relationships between metabolic regulation and neurodegenerative disease,
with implications for therapeutic medical research.

Materials and Methods
Further details of experimental protocols are described in SI Appendix. Code and data can be
found on GitHub: https://github.com/wrthomas315/Dehnels_Seasonal_RNAseq/tree/main/data
355
356

Sample Collection and RNA Sequencing
Eurasian common shrews were trapped with insulated wooden traps in Radolfzell, Germany, at
five different stages of development: large-brained Summer Juveniles (n=5), shrinking Autumn
Juveniles (n=4), small brained Winter Juveniles (n=5), regrowing Spring Adults (n=5), and
regrown Summer Adults (n=5) (Supplemental Table 1). Shrews were aged and sexed based on
tooth wear, fur appearance, and gonad development prior to euthanasia. Fewer females than
males were sampled in juvenile stages (n=6 females), this has minimal effect on gene expression
as shrews are prepubescent and not sexually dimorphic at that age. Only minor sex dimorphism
was detected in the size change of representative brain regions in a previous study (*2*). Blood
was collected prior to perfusion through the heart, let stand for 15 minutes, spun down at 1200
rpm, the serum pipetted off and then stored at -80°C. Shrews were perfused under anesthesia via
the vascular system with PAXgene Tissue Fixative. All organs were removed and weighed to
confirm the occurrence of DP. Thoracic organs were then placed immediately into PAXgene
Tissue Stabilizer, while the brain was submerged in cold fixative, dissected into each region, and
then placed into PAXgene Tissue Stabilizer. All samples were placed in stabilizer (2-24 hours
after extraction) and then stored in liquid nitrogen. RNA was extracted with a modified Qiagen
Micro RNAeasy protocol used for small amounts of mammalian sensory tissue. Tissues extracted
included the liver, cortex, and hippocampus. The hippocampus and cortex were chosen, as they
both undergo substantial non sex-biased shrinkage (*2*), while also the regions most affected by
neurovegetative diseases such as Alzheimer's disease (*71*). Therefore, these may be the most
informative for future therapeutic purposes. RNA was sent to Genewiz for quality control, library
preparation, and sequencing. Quantity of RNA was measured with a Nanodrop
spectrophotometer and quality of RNA assessed with RNA ScreenTape. Libraries were prepared
with standard polyA selection and sequenced as 150bp pair-end reads for a target of 15-25
million reads per sample. To confirm the occurrence of DP, thoracic organs were normalized by
body weight and tested for significant change in mean normalized mass using a t-test between
stages. For the brain, mass was not normalized as change in size itself is of interest, regardless
of parallel changes in body mass.

Temporal Clustering Differential Gene Expression Analysis
We trimmed adapters and filtered reads using fastp (*72*) and removed samples with extreme read
counts (~10-fold). Gene counts were quantified by pseudo aligning to the 19,296 protein coding
genes of the *S. araneus* genome (sorAra2; GCF_000181275.1) using Kallisto (*73*). Counts were
normalized for each tissue using DESeq2 (*74*) median of ratios. We used these normalized gene
expressions to also analyze gene expression through time using the package TCseq (*75*). To
remove genes with relatively little change in expression, we filtered each organ set for genes with
an absolute fold change less than 0.5 between any two stages. Next, we filtered any gene with
extremely low expression, such that two samples had to have greater than ten normalized reads.
We then transformed each gene expression profile across stages to z-scores, and then used
fuzzy cmeans clustering to identify clusters of similar gene expression profiles. The number of
resultant gene clusters was found by bootstrapping (n=20) a gap statistic that minimized the
within-cluster distance (standard error from center). Seasonal cluster membership was compared
to high-effect significant results from differential gene expression analyses below.

Differential Gene Expression Analysis
Tests for differential gene expression were conducted between the phenotypically extreme stages
of the same tissue. For the liver, Autumn Juveniles and Spring Adults were compared, which also
overlap with the spleen, hippocampus, and cortex shrinking and regrowth phases, such that we
could make inferences about potential extrahepatic signals. Winter Juveniles (hippocampus
minimum) and Spring Adults (hippocampus regrowth) were compared. For the cortex, Summer
Juveniles (before shrinkage) and Winter Juveniles (cortex minimum) were compared, as the
cortex does not enter a period of substantial regrowth (*2*). Normalized counts for each gene was
fitted with a negative binomial generalized linear model and then we used a Wald test to examine
if differences in gene expression was the result of the season parameter, with p-values corrected

using the Benjamini and Hochberg procedure (*76*). We filtered our significant results for significant genes higher and lower than 1.58 log-fold change (absolute 3-fold change), to identify how many significant differentially expressed genes were of high effect, largely for gene set enrichments, thus reducing the number of resultant genes to improve interpretability. The hippocampus data had very few differentially expressed genes when filtered (n=25), thus, log-fold change filtering was omitted for this tissue. We then identified enriched KEGG Gene Ontology (GO) pathways in the gene data sets using DAVID functional annotation (*77*).

Weighted Gene Co-expression Network Analysis
We used the WGCNA package (*78*) package to create co-expression networks for the liver and hippocampus and determined if these correlations between genes are significantly correlated with our shrinkage and regrowth phenotypes. Network construction parameters and code can be found in the SI Appendix and on GitHub. Cytoscape (*79*) was used to visualize significant networks and calculate network statistics such as node (gene) connectivity. Identification of potential function of each module were analyzed using DAVID Gene Functional Classification Tool to test for GO enrichment (*77*).

Metabolomics
Metabolite analysis was carried out by MS-Omics using a Thermo Scientific Vanquish LC (UPLC) coupled to Thermo Q Exactive HF MS. Identification of compounds were performed at four levels; Level 1: identification by retention times (compared against in-house authentic standards), accurate mass (with an accepted deviation of 3ppm), and MS/MS spectra, Level 2a: identification by retention times (compared against in-house authentic standards), accurate mass (with an accepted deviation of 3ppm). Level 2b: identification by accurate mass (with an accepted deviation of 3ppm), and MS/MS spectra. MetaboAnalyst was used to test for changes in concentration between seasons (*80*). Concentrations for each sample were normalized by the average concentration of our Summer Adults samples and log transformed and auto scaled. A one-way ANOVA was used to test for any significant seasonal difference in metabolite concentration, with p-values corrected with a false discovery rate of 0.05. Within each significant metabolite, t-tests were used to identify pairwise differences in mean between each season. Samples were hierarchically clustered using significant genes and visualized using a heatmap.

Acknowledgments
We thank Joshua Rest, Krishna Veeramah, and Tanya Lama who have provided helpful feedback on initial results and previous versions of the manuscript. We thank Michal Oklinski for help with initial dissections.

Funding
WRT and research were supported by the Human Frontiers in Science Program Award (RGP0013/2019) to DD, LMD, and JN, and by the Stony Brook University Presidential Innovation and Excellence Fund to LMD.

References
1.   Z. Pucek, Seasonal and age changes in the weight of internal organs of shrews. *Acta Theriol. (Warsz).* **10**, 369–438 (1965). doi:10.4098/at.arch.65-31.
2.   J. Lázaro, M. Hertel, C. C. Sherwood, M. Muturi, D. K. N. Dechmann, Profound seasonal changes in brain size and architecture in the common shrew. *Brain Struct. Funct.* **223**, 2823–2840 (2018). doi:10.1007/s00429-018-1666-5. http://dx.doi.org/10.1007/s00429-018-1666-5.
3.   S. Ray, M. Li, S. P. Koch, S. Mueller, P. Boehm-Sturm, H. Wang, M. Brecht, R. K. Naumann, Seasonal plasticity in the adult somatosensory cortex. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 32136–32144 (2020). doi:10.1073/pnas.1922888117.
4.   M. Genoud, K. Isler, R. D. Martin, Comparative analyses of basal rate of metabolism in mammals : data selection does matter. *Biol. Rev.* **93**, 404–438 (2018).

doi:10.1111/brv.12350.

5.   L. Keicher, M. T. O'Mara, C. C. Voigt, D. K. N. Dechmann, Stable carbon isotopes in
breath reveal fast metabolic incorporation rates and seasonally variable but rapid fat
turnover in the common shrew (Sorex araneus). *J. Exp. Biol.* **220**, 2834–2841 (2017).
doi:10.1242/jeb.159947.

6.   H. Hyvärinen, Wintering strategy of voles and shrews in Findland. *Winter Ecol. Small*
Mamm. **10**, 139–148 (1984).

7.   P. J. Schaeffer, M. T. O'Mara, J. Breiholz, L. Keicher, J. Lázaro, M. Muturi, D. K. N.
Dechmann, Metabolic rate in common shrews is unaffected by temperature, leading to
lower energetic costs through seasonal size reduction. *R. Soc. Open Sci.* **7** (2020).
doi:10.1098/rsos.191989.

8.   J. Lázaro, M. Hertel, M. Muturi, D. K. N. Dechmann, Seasonal reversible size changes in
the braincase and mass of common shrews are flexibly modified by environmental
conditions. *Sci. Rep.* **9**, 1–10 (2019). doi:10.1038/s41598-019-38884-1.

9.   P. Mergenthaler, U. Lindauer, G. A. Dienel, A. Meisel, Sugar for the brain: The role of
glucose in physiological and pathological brain function. *Trends Neurosci.* **36**, 587–597
(2013). doi:10.1016/j.tins.2013.07.001.

10.   D. A. Elliott, C. S. Weickert, B. Garner, Apolipoproteins in the brain: Implications for
neurological and psychiatric disorders. *Clin. Lipidol.* **5**, 555–573 (2010).
doi:10.2217/clp.10.37.

11.   R. Agrawal, F. Gomez-Pinilla, "Metabolic syndrome" in the brain: Deficiency in omega-3
fatty acid exacerbates dysfunctions in insulin receptor signalling and cognition. *J. Physiol.*
590, 2485–2499 (2012). doi:10.1113/jphysiol.2012.230078.

12.   C. Sims-Robinson, B. Kim, A. Rosko, E. L. Feldman, How does diabetes accelerate
Alzheimer disease pathology? *Nat. Rev. Neurol.* **6**, 551–559 (2010).
doi:10.1038/nrneurol.2010.130.

13.   H. Macpherson, M. Formica, E. Harris, R. M. Daly, Brain functional alterations in Type 2
Diabetes – A systematic review of fMRI studies. *Front. Neuroendocrinol.* **47**, 34–46
(2017). doi:10.1016/j.yfrne.2017.07.001. http://dx.doi.org/10.1016/j.yfrne.2017.07.001.

14.   B. B. Boyer, B. M. Barnes, Molecular and metabolic aspects of mammalian hibernation.
Bioscience **49**, 713–724 (1999). doi:10.2307/1313595.

15.   B. Chazarin, K. B. Storey, A. Ziemianin, S. Chanon, M. Plumel, I. Chery, C. Durand, A. L.
Evans, J. M. Arnemo, A. Zedrosser, J. E. Swenson, G. Gauquelin-Koch, C. Simon, S.
Blanc, E. Lefai, F. Bertile, Metabolic reprogramming involving glycolysis in the hibernating
brown bear skeletal muscle. *Front. Zool.* **16**, 1–21 (2019). doi:10.1186/s12983-019-0312-
2.

16.   J. L. Villanueva-Cañas, S. L. Faherty, A. D. Yoder, M. M. Albà, Comparative genomics of
mammalian hibernators using gene networks. *Integr. Comp. Biol.* **54**, 452–462 (2014).
doi:10.1093/icb/icu048.

17.   S. L. Faherty, J. Luis Villanueva-Cañas, P. H. Klopfer, M. M. Albà, A. D. Yoder, Gene
expression profiling in the hibernating primate, Cheirogaleus medius. *Genome Biol. Evol.*
8, 2413–2426 (2016). doi:10.1093/gbe/evw163.

18.   P. W. Hochachka, Defense Strategies against Hypoxia and Hypothermia. *Science (80-.).*
231, 234–241 (1986).

19.   K. L. Drew, M. E. Rice, T. B. Kuhn, M. A. Smith, Neuroprotective adaptations in
hibernation: Therapeutic implications for ischemia-reperfusion, traumatic brain injury and
neurodegenerative diseases. *Free Radic. Biol. Med.* **31**, 563–573 (2001).
doi:10.1016/S0891-5849(01)00628-1.

20.   K. R. Dave, S. L. Christian, M. A. Perez-Pinzon, K. L. Drew, Neuroprotection: Lessons
from hibernators. *Comp. Biochem. Physiol. - B Biochem. Mol. Biol.* **162**, 1–9 (2012).
doi:10.1016/j.cbpb.2012.01.008.

21.   A. P. Ross, S. L. Christian, H. W. Zhao, K. L. Drew, Persistent tolerance to oxygen and
nutrient deprivation and N-methyl-D-aspartate in cultured hippocampal slices from
hibernating Arctic ground squirrel. *J. Cereb. Blood Flow Metab.* **26**, 1148–1156 (2006).

doi:10.1038/sj.jcbfm.9600271.

22.   J. A. Davis, J. R. Paul, L. J. McMeekin, S. R. Nason, J. P. Antipenko, S. D. Yates, R. M.
Cowell, K. M. Habegger, K. L. Gamble, High-Fat and High-Sucrose Diets Impair Time-of-
Day Differences in Spatial Working Memory of Male Mice. *Obesity* **28**, 2347–2356 (2020).
doi:10.1002/oby.22983.

23.   A. Jayaraman, D. Lent-Schochet, C. J. Pike, Diet-induced obesity and low testosterone
increase neuroinflammation and impair neural function. *J. Neuroinflammation* **11**, 1–14
(2014). doi:10.1186/s12974-014-0162-y.

24.   M. P. Mattson, T. V. Arumugam, Hallmarks of Brain Aging: Adaptive and Pathological
Modification by Metabolic States. *Cell Metab.* **27**, 1176–1199 (2018).
doi:10.1016/j.cmet.2018.05.011. https://doi.org/10.1016/j.cmet.2018.05.011.

25.   T. S. Reese, M. J. Karnovsky, Fine structural localization of a blood-brain barrier to
exogenous peroxidase. *J. Cell Biol.* **34**, 207–217 (1967). doi:10.1083/jcb.34.1.207.

26.   G. A. Rosenberg, Neurological diseases in relation to the blood-brain barrier. *J. Cereb.*
Blood Flow Metab. **32**, 1139–1151 (2012). doi:10.1038/jcbfm.2011.197.
http://dx.doi.org/10.1038/jcbfm.2011.197.

27.   B. D. Zipser, C. E. Johanson, L. Gonzalez, T. M. Berzin, R. Tavares, C. M. Hulette, M. P.
Vitek, V. Hovanesian, E. G. Stopa, Microvascular injury and blood-brain barrier leakage in
Alzheimer's disease. *Neurobiol. Aging* **28**, 977–986 (2007).
doi:10.1016/j.neurobiolaging.2006.05.016.

28.   B. V. Zlokovic, Neurovascular pathways to neurodegeneration in Alzheimer's disease and
other disorders. *Nat. Rev. Neurosci.* **12**, 723–738 (2011). doi:10.1038/nrn3114.

29.   A. D. Wong, M. Ye, A. F. Levy, J. D. Rothstein, D. E. Bergles, P. C. Searson, The blood-
brain barrier: An engineering perspective. *Front. Neuroeng.* **6**, 1–22 (2013).
doi:10.3389/fneng.2013.00007.

30.   D. Wang, J. M. Pascual, H. Yang, K. Engelstad, X. Mao, J. Cheng, J. Yoo, J. L. Noebels,
D. C. De Vivo, A mouse model for Glut-1 haploinsufficiency. *Hum. Mol. Genet.* **15**, 1169–
1179 (2006). doi:10.1093/hmg/ddl032.

31.   P. Nguyen, V. Leray, M. Diez, S. Serisier, J. Le Bloc'H, B. Siliart, H. Dumon, Liver lipid
metabolism. *J. Anim. Physiol. Anim. Nutr. (Berl).* **92**, 272–283 (2008). doi:10.1111/j.1439-
0396.2007.00752.x.

32.   Z. Yang, K. Roth, M. Agarwal, W. Liu, M. C. Petriello, The transcription factors CREBH,
PPARa, and FOXO1 as critical hepatic mediators of diet-induced metabolic dysregulation.
J. Nutr. Biochem. **95**, 108633 (2021). doi:10.1016/j.jnutbio.2021.108633.
https://doi.org/10.1016/j.jnutbio.2021.108633.

33.   A. P. Corthals, Multiple sclerosis is not a disease of the immune system. *Q. Rev. Biol.* **86**,
287–321 (2011). doi:10.1086/662453.

34.   Z. Pucek, Seasonal and age change in shrews as an adaptive process. *Symp. zool. Soc.*
Lond. **26**, 189–207 (1970).

35.   J. R. E. Taylor, Evolution of Energetic Strategies in Shrews. *Evol. Shrews*, 309–346
(1998).

36.   J. Lázaro, M. Hertel, S. LaPoint, M. Wikelski, M. Stiehler, D. K. N. Dechmann, Cognitive
skills of common shrews (Sorex araneus) vary with seasonal changes in skull size and
brain mass. *J. Exp. Biol.* **221**, 1–6 (2018). doi:10.1242/jeb.166595.

37.   S. A. Rice, M. Mikes, D. Bibus, E. Berdyshev, J. A. Reisz, S. Gehrke, I. Bronova, A.
D'Alessandro, K. L. Drew, Omega 3 fatty acids stimulate thermogenesis during torpor in
the Arctic Ground Squirrel. *Sci. Rep.* **11**, 1–14 (2021). doi:10.1038/s41598-020-78763-8.
https://doi.org/10.1038/s41598-020-78763-8.

38.   S. Laukola, Seasonal changes in the fatty acid spectrum of the hedgehog's white and
brown adipose tissue. *Ann. Zool. Fennici* **17**, 191–201 (1980).

39.   F. Falkenstein, G. Körtner, K. Watson, F. Geiser, Dietary fats and body lipid composition
in relation to hibernation in free-ranging echidnas. *J. Comp. Physiol. - B Biochem. Syst.*
Environ. Physiol. **171**, 189–194 (2001). doi:10.1007/s003600000157.

40.   S. Giroud, C. Frare, A. Strijkstra, A. Boerema, W. Arnold, T. Ruf, Membrane Phospholipid

Fatty Acid Composition Regulates Cardiac SERCA Activity in a Hibernator, the Syrian
Hamster (Mesocricetus auratus). *PLoS One* **8** (2013). doi:10.1371/journal.pone.0063111.

41.   F. Geiser, M. Klingenspor, B. M. McAllan, A Functional Nexus between Photoperiod
Acclimation, Torpor Expression and Somatic Fatty Acid Composition in a Heterothermic
Mammal. *PLoS One* **8** (2013). doi:10.1371/journal.pone.0063803.

42.   M. R. Pergande, V. G. Amoroso, T. T. A. Nguyen, W. Li, E. Vice, T. J. Park, S. M.
Cologna, PPARα and PPARγSignaling Is Enhanced in the Brain of the Naked Mole-Rat, a
Mammal that Shows Intrinsic Neuroprotection from Oxygen Deprivation. *J. Proteome Res.*
20, 4258–4271 (2021). doi:10.1021/acs.jproteome.1c00131.

43.   X. Meng, J. Zhou, J. Li, R. Wan, H. Yu, Q. Wei, UPLC-QTOF/MS-based metabolomic
analysis of plasma reveals altitude effects on yaks (Bos grunniens). *Thai J. Vet. Med.* **51**,
551–559 (2021). doi:10.14456/tjvm.2021.66.

44.   E. Farhat, E. D. Turenne, K. Choi, J. M. Weber, Hypoxia-induced remodelling of goldfish
membranes. *Comp. Biochem. Physiol. Part B* **237**, 1–8 (2019).
doi:10.1016/j.cbpb.2019.110326. https://doi.org/10.1016/j.cbpb.2019.110326.

45.   M. E. Pamenter, Adaptations to a hypoxic lifestyle in naked mole-rats. *J. Exp. Biol.* **225**
(2022). doi:10.1242/jeb.196725.

46.   A. M. Kirby, G. D. Fairman, M. E. Pamenter, Atypical behavioural, metabolic and
thermoregulatory responses to hypoxia in the naked mole rat (Heterocephalus glaber). *J.*
Zool. **305**, 106–115 (2018). doi:10.1111/jzo.12542.

47.   W. Zhang, S. Patil, B. Chauhan, S. Guo, D. R. Powell, J. Le, A. Klotsas, R. Matika, X.
Xiao, R. Franks, K. A. Heidenreich, M. P. Sajan, R. V. Farese, D. B. Stolz, P. Tso, S. H.
Koo, M. Montminy, T. G. Unterman, FoxO1 regulates multiple metabolic pathways in the
liver effects on gluconeogenic, glycolytic, and lipogenic gene expression. *J. Biol. Chem.*
281, 10105–10117 (2006). doi:10.1074/jbc.M600272200.
http://dx.doi.org/10.1074/jbc.M600272200.

48.   X. Xiong, R. Tao, R. A. DePinho, X. C. Dong, Deletion of Hepatic FoxO1/3/4 Genes in
Mice Significantly Impacts on Glucose Metabolism through Downregulation of
Gluconeogenesis and Upregulation of Glycolysis. *PLoS One* **8**, 1–11 (2013).
doi:10.1371/journal.pone.0074340.

49.   M. Matsumoto, A. Pocai, L. Rossetti, R. A. DePinho, D. Accili, Impaired Regulation of
Hepatic Glucose Production in Mice Lacking the Forkhead Transcription Factor Foxo1 in
Liver. *Cell Metab.* **6**, 208–216 (2007). doi:10.1016/j.cmet.2007.08.006.

50.   I. Tikhanovich, J. Cox, S. A. Weinman, Forkhead box class O transcription factors in liver
function and disease. *J. Gastroenterol. Hepatol.* **28**, 125–131 (2013).
doi:10.1111/jgh.12021.

51.   M. S. Gami, C. A. Wolkow, Studies of Caenorhabditis elegans DAF-2/insulin signaling
reveal targets for pharmacological manipulation of lifespan. *Aging Cell* **5**, 31–37 (2006).
doi:10.1111/j.1474-9726.2006.00188.x.

52.   D. S. Hwangbo, B. Garsham, M. P. Tu, M. Palmer, M. Tatar, Drosophila dFOXO controls
lifespan and regulates insulin signalling in brain and fat body. *Nature* **429**, 562–566
(2004). doi:10.1038/nature02549.

53.   A. Satoh, C. S. Brace, N. Rensing, P. Cliften, D. F. Wozniak, E. D. Herzog, K. A. Yamada,
S. I. Imai, Sirt1 extends life span and delays aging in mice through the regulation of Nk2
Homeobox 1 in the DMH and LH. *Cell Metab.* **18**, 416–430 (2013).
doi:10.1016/j.cmet.2013.07.013. http://dx.doi.org/10.1016/j.cmet.2013.07.013.

54.   M. Katic, C. R. Kahn, The role of insulin and IGF-1 signaling in longevity. *Cell. Mol. Life*
Sci. **62**, 320–343 (2005). doi:10.1007/s00018-004-4297-y.

55.   K. Healy, T. Guillerme, S. Finlay, A. Kane, S. B. A. Kelly, D. McClean, D. J. Kelly, I.
Donohue, A. L. Jackson, N. Cooper, Ecology and mode-of-life explain lifespan variation in
birds and mammals. *Proc. R. Soc. B Biol. Sci.* **281** (2014). doi:10.1098/rspb.2014.0298.

56.   S. Du, Z. Guan, L. Hao, Y. Song, L. Wang, L. Gong, L. Liu, X. Qi, Z. Hou, S. Shao,
Fructose-bisphosphate aldolase a is a potential metastasis-associated marker of lung
squamous cell carcinoma and promotes lung cell tumorigenesis and migration. *PLoS One*

627   **9** (2014). doi:10.1371/journal.pone.0085804.

628   57.   J. H. T. Potter, R. Drinkwater, K. T. J. Davies, N. Nesi, M. C. W. Lim, L. R. Yohe, H. Chi,
629   X. Zhang, I. Levantis, B. K. Lim, C. C. Witt, G. Tsagkogeorga, M. dos Reis, Y. Liu, W.
630   Furey, M. J. Whitley, D. Aksentijevic, L. M. Dávalos, S. J. Rossiter, Nectar-feeding bats
631   and birds show parallel molecular adaptations in sugar metabolism enzymes. *Curr. Biol.*
632   **31**, 1–8 (2021). doi:10.1016/j.cub.2021.08.018.

633   58.   C. N. Brocker, D. P. Patel, T. J. Velenosi, D. Kim, T. Yan, J. Yue, G. Li, K. W. Krausz, F. J.
634   Gonzalez, Extrahepatic PPAR modulates fatty acid oxidation and attenuates fasting-
635   induced hepatosteatosis in mice. *J. Lipid Res.* **59**, 2140–2152 (2018).
636   doi:10.1194/jlr.M088419. http://dx.doi.org/10.1194/jlr.M088419.

637   59.   D. Val-Laillet, S. Layec, S. Guérin, P. Meurice, C. H. Malbert, Changes in brain activity
638   after a diet-induced obesity. *Obesity* **19**, 749–756 (2011). doi:10.1038/oby.2010.292.
639   http://dx.doi.org/10.1038/oby.2010.292/nature06264.

640   60.   R. Agrawal, Y. Zhuang, B. P. Cummings, K. L. Stanhope, J. L. Graham, P. J. Havel, F.
641   Gomez-Pinilla, Deterioration of plasticity and metabolic homeostasis in the brain of the
642   UCD-T2DM rat model of naturally occurring type-2 diabetes. *Biochim. Biophys. Acta - Mol.*
643   *Basis Dis.* **1842**, 1313–1323 (2014). doi:10.1016/j.bbadis.2014.05.007.
644   http://dx.doi.org/10.1016/j.bbadis.2014.05.007.

645   61.   A. Kalsbeek, E. Foppen, I. Schalij, C. Van Heijningen, J. van der Vliet, E. Fliers, R. M.
646   Buijs, Circadian control of the daily plasma glucose rhythm: An interplay of GABA and
647   glutamate. *PLoS One* **3** (2008). doi:10.1371/journal.pone.0003194.

648   62.   F. Gachon, U. Loizides-Mangold, V. Petrenko, C. Dibner, Glucose homeostasis:
649   Regulation by peripheral circadian clocks in rodents and humans. *Endocrinology* **158**,
650   1074–1084 (2017). doi:10.1210/en.2017-00218.

651   63.   S.-C. Lin, D. G. Hardie, AMPK: Sensing Glucose as well as Cellular Energy Status. *Cell*
652   *Metab.* **27**, 299–313 (2018). doi:10.1016/j.cmet.2017.10.009.
653   https://doi.org/10.1016/j.cmet.2017.10.009.

654   64.   Y. Minokoshi, T. Alquier, H. Furukawa, Y. B. Kim, A. Lee, B. Xue, J. Mu, F. Foufelle, P.
655   Ferré, M. J. Birnbaum, B. J. Stuck, B. B. Kahn, AMP-kinase regulates food intake by
656   responding to hormonal and nutrient signals in the hypothalamus. *Nature* **428**, 569–574
657   (2004). doi:10.1038/nature02440.

658   65.   B. Zhang, J. Zhong, Z. Gao, A Brain-Spleen Axis Regulates Humoral Immunity. *Neurosci.*
659   *Bull.* **37**, 427–429 (2021). doi:10.1007/s12264-020-00610-7.
660   https://doi.org/10.1007/s12264-020-00610-7.

661   66.   S. Z. Wang, Y. J. Yu, K. Adeli, Role of gut microbiota in neuroendocrine regulation of
662   carbohydrate and lipid metabolism via the microbiota-gut-brain-liver axis. *Microorganisms*
663   **8**, 8–10 (2020). doi:10.3390/microorganisms8040527.

664   67.   N. Zeltser, I. Meyer, G. V. Hernandez, M. J. Trahan, R. K. Fanter, M. Abo-Ismail, H.
665   Glanz, C. R. Strand, D. G. Burrin, M. R. la Frano, R. Manjarín, M. Maj, Neurodegeneration
666   in juvenile Iberian pigs with diet-induced nonalcoholic fatty liver disease. *Am. J. Physiol. -*
667   *Endocrinol. Metab.* **319**, E592–E606 (2020). doi:10.1152/AJPENDO.00120.2020.

668   68.   M. Vairetti, A. Ferrigno, V. Rizzo, G. Ambrosi, A. Bianchi, P. Richelmi, F. Blandini, M. T.
669   Armentero, Impaired hepatic function and central dopaminergic denervation in a rodent
670   model of Parkinson's disease: A self-perpetuating crosstalk? *Biochim. Biophys. Acta -*
671   *Mol. Basis Dis.* **1822**, 176–184 (2012). doi:10.1016/j.bbadis.2011.11.008.
672   http://dx.doi.org/10.1016/j.bbadis.2011.11.008.

673   69.   Z. Huang, H. W. Lin, Q. Zhang, X. Zong, Targeting Alzheimer's Disease: The Critical
674   Crosstalk between the Liver and Brain. *Nutrients* **14** (2022). doi:10.3390/nu14204298.

675   70.   M. Trapecar, E. Wogram, D. Svoboda, C. Communal, A. Omer, T. Lungjangwa, P.
676   Sphabmixay, J. Velazquez, K. Schneider, C. W. Wright, S. Mildrum, A. Hendricks, S.
677   Levine, J. Muffat, M. J. Lee, D. A. Lauffenburger, D. Trumper, R. Jaenisch, L. G. Griffith,
678   Human physiomimetic model integrating microphysiological systems of the gut, liver, and
679   brain for studies of neurodegenerative diseases. *Sci. Adv.* **7** (2021).
680   doi:10.1126/SCIADV.ABD1707.

71. X. Wang, M. L. Michaelis, E. K. Michaelis, Functional Genomics of Brain Aging and
Alzheimers Disease: Focus on Selective Neuronal Vulnerability. *Curr. Genomics* **11**, 618–
633 (2010). doi:10.2174/138920210793360943.
72. S. Chen, Y. Zhou, Y. Chen, J. Gu, Fastp: An ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics **34**, i884–i890 (2018). doi:10.1093/bioinformatics/bty560.
73. N. L. Bray, H. Pimentel, P. Melsted, L. Pachter, Near-optimal probabilistic RNA-seq
quantification. *Nat. Biotechnol.* **34**, 525–527 (2016). doi:10.1038/nbt.3519.
74. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol.* **15**, 1–21 (2014). doi:10.1186/s13059-014-
0550-8.
75. L. G. Mengjun, TCseq: Time course sequencing data analysis. R package version 1.8.0.
1–8 (2019).
76. Y. Benjamini, Y. Hochberg, Controlling the False Discovery Rate : A Practical and
Powerful Approach to Multiple Testing. *J. R. Stat. Soc.* **57**, 289–300 (1995).
77. D. W. Huang, B. T. Sherman, R. A. Lempicki, Systematic and integrative analysis of large
gene lists using DAVID bioinformatics resources. *Nat. Protoc.* **4**, 44–57 (2009).
doi:10.1038/nprot.2008.211.
78. P. Langfelder, S. Horvath, WGCNA: An R package for weighted correlation network
analysis. *BMC Bioinformatics* **9** (2008). doi:10.1186/1471-2105-9-559.
79. P. Shannon, A. Markiel, O. Ozier, S. B. Nitin, J. T. Wang, D. Ramage, N. Amin, B.
Schwikowski, T. Ideker, Cytoscape: A Software Environment for Integrated Models.
Genome Res. **13**, 2498–2504 (2003). doi:10.1101/gr.1239303.metabolite.
http://ci.nii.ac.jp/naid/110001910481/.
80. Z. Pang, J. Chong, G. Zhou, D. A. De Lima Morais, L. Chang, M. Barrette, C. Gauthier, P.
É. Jacques, S. Li, J. Xia, MetaboAnalyst 5.0: Narrowing the gap between raw spectra and
functional insights. *Nucleic Acids Res.* **49**, 388–396 (2021). doi:10.1093/nar/gkab382.
707
708

Figures

710
Figure 1. Metabolic profiling of the blood metabolome and liver transcriptomes. (**A**) Heatmap of
28 statistically significant differentially concentrated metabolites between stages of Dehnel's
phenomenon. Hierarchical clustering using these significant metabolites groups each profile into
each season. Three of these metabolites were lipid metabolites (ethyl palmitoleate,
docosahexanoic acid, arachidonic acid), with decreases in autumn, winter, and spring individuals.
(B) Mass change in liver through five stages of Dehnel's phenomenon. Unlike other organs, the
liver reaches a minimum as an Autumn Juvenile, with regrowth beginning as a Winter Juvenile.
Asterisks represent significant size changes (adjusted p<0.05). **(C)** Volcano plot of significant
(adjusted p<0.05) differentially expressed genes (colored) between phenotypic extremes of liver
size change (Spring Adult vs Autumn Juvenile) plotted by log fold-change. Vertical thresholds
represent a 1.58 log fold-change in differential expression of high effect between stages (dark
colors). **(D)** Temporal gene expression clusters 8 (n=95 significant genes) and 12 (n=79
significant genes), with the most significant differentially expressed genes. **(E)** Patterning of gene
expression across stages of Dehnel's phenomenon for transcription factors pivotal roles in lipid
metabolism and gluconeogenesis. Lipid metabolism and gluconeogenesis appear to have inverse
cycles, with lipid metabolism decreasing in the fall and increasing in the spring, with
gluconeogenesis following the opposite trend.
728

Figure 2. Seasonal plasticity in brain phenotype and expression patterns. **(A)** Brain mass change through five stages of Dehnel's phenomenon. As does body mass, the brain begins to shrink as Autumn Juveniles, reaching a minimum as Winter Juveniles, and regrows as Spring Adults. Asterisks represent significant size changes (adjusted p<0.05). **(B)** 11 genes were differentially expressed in both brain regions, including the apolipoproteins APOA1 and APOH, which both show winter decreases followed by slight spring upregulation. **(C)** Volcano plot of significant (adjusted p<0.05) differentially expressed genes (colored) in *hippocampal regrowth* (Spring Adults vs Winter Juveniles) and *cortex shrinkage* (Winter Juveniles vs Summer Juveniles) plotted by log fold-change. Vertical thresholds represent a 1.58 log fold-change which show differential expression of high effect between stages. **(D)** APOA1 and APOH are found in the cholesterol metabolism pathway, which was enriched for differentially expressed genes in both shrinkage and regrowth phenotypes.

Figure 3. Gene network analyses across liver and body phenotypes of Dehnel's phenomenon. (**A**) Modules (n=37) of high gene co-expression in the liver, including 21 modules with significant correlation with at least one change phenotype (Controls, Liver Shrinkage, Liver Growth, Body Shrinkage, and Body Growth). The 2816 gene cyan module was significant in all size changes, tests for pathway enrichment found 77 significantly enriched pathways, with 59 shown after filtering (removed pathways without 2 neighbors for simplicity). (**B**) Gene pathways clustered in 16 clusters GO-pathways based on gene similarity (**C**) Cluster 1 was the most enriched cluster (Enrichment Score 3.56; higher ES occurs with lower geometric mean of encompassed pathway p-values) containing four pathways; Insulin resistance, Longevity regulating pathway, AMPK signaling pathway, and Longevity regulating pathway-multiple species.

Figure 4. Gene network analyses across hippocampal size change. (**A**) Modules (34) of high gene co-expression in the hippocampus, including ten modules significantly correlated with target phenotypes (Regrowth, Shrinkage, and Control). The 346 gene grey module was the most significant module associated within the shrinkage phenotype (p = 0.01, r = 0.55). (**B**) Visualization of gene network (filtered and removed edges with weighted correlations < 0.02 and nodes with only a single neighbor). ALDOA was the most central to the network, with 17 connections to other nodes. (**C**) Pathway enrichment analysis also identified 6 pathways to be enriched for genes in the grey module: Pathways of Neurodegeneration, Parkinson's disease, Huntington's disease, NF-kappa beta Signaling, Dilated Cardiomyopathy, and Biosynthesis.