Multiple independent acquisitions of ACE2 usage in MERS-related coronaviruses

Cheng-Bao Ma[1,7], Chen Liu[1,7], Young-Jun Park[2,3,7], Jingjing Tang[4,7], Jing Chen[4,7], Qing Xiong[1], Jimin Lee[2], Cameron Stewart[2], Daniel Asarnow[2], Jack Brown[2], M. Alejandra Tortorici[2], Xiao Yang[1], Ye-Hui Sun[1], Yuan-Mei Chen[1], Xiao Yu[1], Jun-Yu Si[1], Peng Liu[1], Fei Tong[1], Mei-Ling Huang[1], Jing Li[1], Zheng-Li Shi[4,5]*, Zengqin Deng[4,6]*, David Veesler[2,3]*, Huan Yan[1,8,] *

Affiliations:
[1]State Key Laboratory of Virology, College of Life Sciences, TaiKang Center for Life and Medical Sciences, Wuhan University; Wuhan, Hubei, 430072, China.
[2]Department of Biochemistry, University of Washington; Seattle, WA 98195, USA.
[3]Howard Hughes Medical Institute, University of Washington; Seattle, WA 98195, USA.
[4]Key Laboratory of Virology and Biosafety, Wuhan Institute of Virology, Chinese Academy of Sciences; Wuhan, China.
[5]Guangzhou Laboratory, Guangzhou International Bio Island; Guangzhou, China
[6]Hubei Jiangxia Laboratory; Wuhan, China.
[7]These authors contributed equally to this work.
[8]Lead contact
*Correspondence: Zheng-Li Shi: zlshi@wh.iov.cn (Z.L.S), dengzengqin@wh.iov.cn (Z.D.), dveesler@uw.edu (D.V.), huanyan@whu.edu.cn (H.Y.)

SUMMARY

The angiotensin-converting enzyme 2 (ACE2) receptor is shared by various coronaviruses with distinct receptor-binding domain (RBD) architectures, yet our understanding of these convergent acquisition events remains elusive. Here, we report that two European bat MERS-related coronaviruses (MERSr-CoVs) infecting *Pipistrellus nathusii* (P.nat), MOW15-22 and PnNL2018B, use ACE2 as their receptor, with narrow ortholog specificity. Cryo-electron microscopy structures of the MOW15-22 RBD-ACE2 complex unveil an unexpected and entirely distinct binding mode, mapping 50Å away from that of any other known ACE2-using coronaviruses. Functional profiling of ACE2 orthologs from 105 mammalian species led to the identification of host tropism determinants, including an ACE2 N432-glycosylation restricting viral recognition, and the design of a soluble P.nat ACE2 mutant with potent viral neutralizing activity. Our findings reveal convergent acquisition of ACE2 usage for merbecoviruses found in European bats, underscoring the extraordinary diversity of ACE2 recognition modes among coronaviruses and the promiscuity of this receptor.

Graphic abstract

INTRODUCTION

Human MERS-CoV belongs to the *Merbecovirus* subgenus (also known as lineage C β-coronavirus or group 2c coronavirus) and is the causative agent of Middle-East respiratory syndrome (MERS) with a case-fatality rate of 36%.[1] The evolutionary trajectory and host-switching history of MERS-CoV remain unclear. While dromedary camels are established intermediate hosts of MERS-CoV, increasing evidence from newly identified viral sequences indicates that Old World vesper bats (Vespertilionidae) are natural merbecovirus reservoirs.[2–7] However, no currently known bat MERS-related coronaviruses (MERSr-CoVs) exhibit close similarity to human or camel MERS-CoV at the whole-genome level.[8,9] NeoCoV, identified in African Laephotis bats (Cape serotine), represents the closest known relative to MERS-CoV, sharing 85.5% whole genome nucleotide sequence identity.[10]

Receptor usage determines host tropism and transmission of coronaviruses[11]. Dipeptidyl peptidase-4 (DPP4) has been documented as the entry receptor for several merbecoviruses, including MERS-CoV, HKU4, MjHKU4r, HKU25, and BtCoV-422.[12–18] NeoCoV and the closely related PDF-2180 are two bat MERSr-CoVs that harbor markedly different spike (S) glycoproteins relative to that of MERS-CoV, and their cognate receptor remained elusive for a decade. For a long time, ACE2 was solely considered the host receptor recognized by members of sarbecoviruses (e.g., SARS-CoV-1 and SARS-CoV-2) and setracoviruses (e.g., NL63) through two different ACE2-binding modes with largely overlapping footprints in spite of their distinct receptor-binding domain (RBD) architectures.[19–23] We recently revealed that NeoCoV and PDF-2180 also use the ACE2 receptor with a broad tropism across mammals but low efficiency in using human ACE2 (hACE2).[9,24] Cryo-electron microscopy (cryo-EM) analysis of the NeoCoV and the PDF-2180 RBDs in complex with *Pipistrellus pipistrellus* (P.pip) ACE2 unveiled a distinct binding mode from that of SARS-CoV-1/SARS-CoV-2 and of NL63 involving extensive interactions with ACE2 N linked-glycans. The footprint of their RBDs shared few residues with that of SARS-CoV-1/2 or NL63, highlighting a convergent evolutionary history of ACE2 acquisition in these viruses.[9] This discovery suggested that a receptor switch might have occurred during MERS-CoV emergence in animals, such as bats and camels, potentially through recombination between a NeoCoV-like MERSr-CoV and a DPP4-using merbecovirus (e.g., HKU4 or BtCoV-422).[8,13] Furthermore, these findings underscored the complexity of receptor binding modes used by merbecoviruses with diverse RBD sequences, many of which remain enigmatic.

Here, we discovered that two MERSr-CoVs circulating in European bats, designated MOW15-22 and PnNL2018B, utilize a subset of mammalian ACE2 orthologs as entry receptors. Cryo-EM analyses reveal that these viruses recognize ACE2 through an unprecedented binding mode, mapping 50Å away from the footprint of any other coronaviruses, emphasizing multiple independent acquisitions of ACE2 usage in bat MERSr-CoVs. This study sheds light on the global diversity and distribution of ACE2-using merbecoviruses and unexpected convergent evolution events, underscoring the zoonotic potential associated with these viruses.

RESULTS

Prediction of a distinct receptor recognition mode utilized by two MERSr-CoVs

In light of the unexpected ACE2 usage of NeoCoV and PDF-2180, we investigated whether other merbecoviruses utilize ACE2 as a receptor. We conducted phylogenetic analyses of representative merbecovirus sequences, with a special emphasis on viruses belonging to MERSr-CoVs, as defined by amino acid sequence identity of the five concatenated replicase domains (3CLpro, NiRAN, RdRp, ZBD, and HEL1) greater than 92.4% compared with MERS-CoV (Figure 1A).[25] At the whole-genome level, MOW15-22 and PnNL2018B (formerly PN-βCoV) are two bat MERSr-CoVs with the highest genetic similarity to MERS-CoV after NeoCoV and PDF-2180 (Figure 1A). At the S glycoprotein level, the MOW15-22 and PnNL2018B S glycoproteins cluster together and are distantly related to all other MERSr-CoVs. By contrast, hedgehog merbecovirus (Erinaceus coronavirus, EriCoV) HKU31 exhibits the highest sequence identity with NeoCoV and PDF-2180, although HKU31 is not known to use ACE2 (Figure 1B).[9] Simplot analysis comparing several viral genome sequences with MOW15-22 reveals both MERS-CoV and NeoCoV display an extensive divergence within the S_1 subunit region (Figure 1C). Consistent with the phylogenetic tree based on S glycoproteins, pairwise analysis shows MOW15-22 and PnNL2018B share only 31-36% and 57-60% amino acid sequence identities with RBDs and S glycoproteins of other merbecoviruses, respectively (Figure 1D). Alignment of the region corresponding to the NeoCoV receptor-binding motif (RBM) reveals the presence of two insertions and a disulfide bond specific of MOW15-22 and PnNL2018B (Figure 1E). Only two (positions $N502_{MOW15-22}$ and $G566_{MOW15-22}$) out of nine NeoCoV residues critical for interactions with P.pip ACE2 are conserved with MOW15-22 and PnNL2018B (Figure 1E).[9] Furthermore, the two insertions and elongated α-helix in the RBM present in AlphaFold2 predicted structures[26] potentially affect receptor recognition (Figure 1F). Overall, these distinct sequence and structural features indicate that MOW15-22 and PnNL2018B may utilize a previously uncharacterized receptor binding mode.

Figure 1. Prediction of a distinct receptor recognition mode utilized by two MERSr-CoVs.
(**A-B**) Phylogenetic trees of representative merbecoviruses were generated using complete genome nucleotide sequences (A) or S glycoprotein amino acid sequences (B) with the IQ-tree method. NL63 was set as an outgroup. Amino acid sequence identities of five replicase domains for coronavirus classification, host species, and receptor-related information (receptor usage, binding mode, and availability of RBD/receptor complex structure) are indicated. (**C**) Simplot analysis of the complete genome sequence similarity of several MERSr-CoVs analyzed based on the MOW15-22 genome. The right panel magnifies the RBD and adjacent regions. (**D**) Pairwise RBD and S amino acid sequence identities of indicated merbecoviruses. (**E**) RBM sequence alignment of the indicated merbecoviruses. MOW15-22 and PnNL2018B-specific insertions and disulfides are indicated in red and yellow, respectively. Red/black asterisks: residues crucial for NeoCoV interactions with P.pip ACE2 that are conserved/not conserved with MOW15-22 and PnNL2018B. The MOW15-22 residue numbering is shown. (**F**) Cryo-EM structures or AlphaFold2-predicted RBD structures of representative merbecoviruses. Magenta indicates putative RBMs, blue represents the RBM helix, and the MOW15-22 and PnNL2018B-specific insertions (ins) are indicated. The yellow sticks represent specific disulfide bonds in RBM insertion 1.

MOW15-22 and PnNL2018B use ACE2 as receptor

MOW15-22 and PnNL2018B were recently sampled in *Pipistrellus nathusii* (P.nat, the common host) in Russia (Moscow region) and the Netherlands, respectively (Figure 2A).[27,28] P.nat inhabits a wide range of Europe and undertakes seasonal long-distance migrations, typically from northeast to southwest Europe.[29–31] Previous reports have proposed that DPP4 may serve as the entry receptor for MOW15-22 and PnNL2018B based on molecular docking analyses.[27,28]

However, in view of the unique sequence and structural features of MOW15-22 and PnNL2018B, we set out to investigate receptor usage of MOW15-22 and PnNL2018B by assessing the ability of human and P.nat ACE2 and DPP4 orthologs to support S-mediated pseudovirus entry and viral antigen bindings. We found that P.nat ACE2, but not human ACE2 (hACE2), human DPP4, or P.nat DPP4, promoted MOW15-22 and PnNL2018B pseudovirus entry and RBD or S_1 binding (Figures 2B-2E). Biolayer interferometry (BLI) analysis showed that the soluble dimeric P.nat ectodomain, but not human ACE2, bound to immobilized MOW15-22 RBD with an apparent affinity (K_D, app) of 149.9 nM (Figures 2F-2G). Further investigations of the contributions of the four domains present in the S_1 subunit revealed that the MOW15-22 RBD is sufficient for binding to P.nat ACE2 whereas the SD1 and SD2 domains seem required for better PnNL2018B engagement (Figures 2H-2J). Therefore, we recommend using the S_1 subunits of PnNL2018B to assess ACE2 binding efficiency.

Collectively, these data along with a recent report showing that HKU5 uses *Pipistrellus abramus* (P.abr) ACE2 as an entry receptor[32] expand the geographic distribution of ACE2-using merbecoviruses to all three continents of the Old World, with Pipistrellus bats as important reservoir hosts that should be closely monitored (Figure 2K).

Figure 2. MOW15-22 and PnNL2018B use ACE2 as receptor. (**A**) Geographical distribution of *Pipistrellus nathusii* in Europe. The purple regions represent its habitat, data retrieved from the IUCN (International Union for Conservation of Nature) Red List of Threatened Species, and the distribution chart was generated using Geoscene Pro. Red asterisks indicate viral discovery locations. (B) Expression levels of human or P.nat ACE2 and DPP4 orthologs in HEK293T cells. (**C-D**) P.nat ACE2 but not DPP4 supports MOW15-22 and PnNL2018B RBD-hFc binding (C) and pseudovirus (PSV) entry (D) in HEK293T cells. Mean ± SD and unpaired two-tailed t-tests for D. n=4 biological replicates. RLU: Relative light unit. (**E**) Flow cytometry analysis of MOW15-22 and PnNL2018B S1 binding to P.nat ACE2, hACE2 or P.nat DPP4 transiently expressed in HEK293T cells. Gray: cells transfected with vector control. Mean ± SD for three technical repeats. (**F-G**) BLI analyses of binding kinetics of the soluble dimeric P.nat ACE2 (F) or hACE2 (G) ectodomain to immobilized MOW15-22 RBD-hFc. Analysis was conducted with curve-fitting kinetic with global fitting (1:1 binding model). NA: not applicable. (**H**) Binding of recombinant MOW15-22 and PnNL2018B S1 subunit constructs to P.nat ACE2 expressed at the surface of HEK293T cells detected by fluorescence microscopy. (**I**) Expression levels of recombinant proteins comprising different domains of the PnNL2018B S_1 subunit. Equal volumes of protein-containing supernatants used for purification were loaded for Western blot analysis. (**J**) Binding of PnNL2018B S_1 subunit constructs at various concentrations to HEK293T cells transiently expressing P.nat ACE2 analyzed by fluorescence microscopy. (**K**) Confirmed DPP4 or ACE2 usage of representative merbecoviruses is displayed with blue and red backgrounds, respectively. Discovery locations and natural hosts are indicated. Scale bars: 100 μm for C, H, and J.

Multi-species ACE2 tropism of MOW15-22 and PnNL2018B

To explore the potential host range of MOW15-22 and PnNL2018B, we assessed the ability of various ACE2 orthologs transiently transfected in HEK293T cells to promote RBD/S_1 subunit binding and pseudovirus entry. We used a receptor library comprising 105 ACE2 orthologs from 52 bats and 53 non-bat mammalian species with validated expression (Figure S1).[24] MOW15-22 efficiently used bat ACE2 from P.nat, P.par, P.dav, and a few others to a lesser extent (Figure 3A). PnNL2018B could only use P.nat and L.bor ACE2 efficiently for entry. In contrast to NeoCoV and PDF-2180, MOW15-22 and even more so PnNL2018B had limited tropism for non-bat mammalian ACE2s (Figure 3B), with only a few orthologs from Carnivora and Primates species, such as dog (C.fam) and common marmoset (C.jac) supporting MOW15-22 entry (Figure 3B).

Concurring with the entry data, P.nat, P.dav, P.par, and less prominent for several other mammalian ACE2s, supported binding of the MOW15-22 S_1 subunit whereas only P.nat ACE2 supported efficient PnNL2018B RBD/S_1 binding, with higher sensitivity observed in S_1 binding assays (Figures 3A-3B and Figure S2). Conversely, ACE2 from Pipistrellus bats, P.pip and P.kuh, did not support MOW15-22 or PnNL2018B RBD/S_1 binding, indicating host ACE2 specialization during virus evolution. As a result, two Pteronotus bat ACE2s but not the additional two Pipstrellus bat orthologs tested were functional receptors, underscoring the importance of specific sequence determinants over phylogenetic relationships of the bat hosts. BLI analysis showed that the soluble P.dav ACE2s bound to immobilized MOW15-22 RBD with an apparent affinity (K_D, app) of 8.6 nM, outperforming the binding of P.nat ACE2 (Figure 3C). We also tested several DPP4 orthologs from bats, humans, and hedgehogs, and none of them showed any detectable receptor function (Figure 3D-F).

We observed pseudovirus entry for several ACE2 orthologs for which only weak or no MOW15-22 S_1 binding could be detected (Figure 3B, Figure S2C, and S3A-B), as previously

described for other coronaviruses.[9,33] This apparent discrepancy may be attributed to the use of TPCK-treated trypsin to enhance pseudovirus entry with weak receptors and to the multivalent presentation of S trimers on pseudoviruses markedly increasing binding avidity, as compared to dimeric RBD or S_1-Fc.[34] Accordingly, L.bor ACE2 promoted pseudovirus entry and membrane fusion in the presence of exogenous trypsin, although no MOW15-22 S_1 binding to this ACE2 was detected (Figure S3C-F). The addition of exogenous trypsin can thus improve pseudovirus entry assay sensitivity to identify weakly functional ACE2 orthologs.[34]

Figure 3. Multi-species ACE2 tropism of MOW15-22 and PnNL2018B. (**A-B**) Heat map representing MOW15-22 PSV entry into and RBD/S_1 subunit binding to HEK293T cells transiently expressing various ACE2 orthologs from bats (A) or other mammalian species (B). Different mammalian orders are denoted with distinctly colored backgrounds, from left to right: Carnivora, Primates, Artiodactyla, Rodentia, Cetacea, Perissodactyla, Diprotodontia, Pholidota, Erinaceomorpha, and Lagomorpha. Data are normalized relative to P.nat ACE2 and shown as mean values. n=3 biological repeats for PSV entry. Data representative of 2-3 independent experiments for antigen binding assays. (**C**) BLI analysis of binding kinetics of the soluble recombinant dimeric P.dav ACE2 ectodomain to immobilized biotinylated MOW15-22 RBD. (**D**) Expression levels of several mammalian DPP4 orthologs in HEK293T cells. (**E**) PSV entry efficiency of MOW15-22 and PnNL2018B in HEK293T expressing the indicated receptors. Dashed lines: threshold of the background entry. MEAN ± SEM and unpaired two-tailed t-tests. n=3 biological replicates. (**F**) MOW15-22 and PnNL2018B RBD-hFc binding to HEK293T cells transiently expressing the indicated receptors assessed by fluorescence microscopy. Scale bars: 100 μm.

See also Figure S1-S3.

Host ACE2 tropism determinants for MOW15-22 and PnNL2018B

We previously identified four host range molecular determinants (designated A-D) on the ACE2 receptor for NeoCoV/PDF-2180.[24] To assess their relevance for species-specific recognition by MOW15-22 and PnNL2018B, we generated P.dav, P.par, P.pip, and L.bor ACE2 mutants harboring unfavorable changes for NeoCoV S-mediated entry. Unexpectedly, none of these alterations markedly affected MOW15-22 S receptor utilization (Figure S4). Moreover, exchanging residues 1-400 between M.bla ACE2 and P.dav ACE2 had no effect on their ability to promote MOW15-22 RBD binding and pseudovirus entry in spite of comprising all of the corresponding residues recognized by NeoCoV (Figure 4A). Additional sequence swaps led to the identification of residues 400-450 and subsequently the $N428_{M.blaACE2}$ glycosylation site ($Y430_{P.davACE2}$) as critical host range determinants for both MOW15-22 and PnNL2018B, with the presence of this oligosaccharide abolishing RBD or S_1 binding with P.dav ACE2 (Figure 4A and Figure S5A-D). However, the restrictive effect of this glycan is less pronounced on MOW15-22/PnNL2018B S_1 than MOW15-22/PnNL2018B RBD binding to P.nat ACE2, underscoring possible differences of interactions with different ACE2s (Figures S5E-5F). Furthermore, the N432C mutation, which abrogates the glycosylation site, was insufficient to enable strong binding of the MOW15-22 RBD or S_1 to P.pip ACE2 without the additional substitutions of residues 500-600 or of residues $E589_{P.pipACE2}$ (K in P.nat ACE2) and $K597_{P.pipACE2}$ (E in P.nat ACE2) (Figure 4B). Consistently, P.nat ACE2 chimera with residues 500-600 replaced by P.pip ACE2 equivalents lost MOW15-22 RBD/S_1 binding. Furthermore, substitutions at position $K287_{P.natACE2}$ (K in P.nat ACE2) were identified to modulate MOW15-22 and PnNL2018B RBD and S_1 binding in ACE2 swap experiments testing P.nat ACE2 and P.pip ACE2-N432C (Figure 4C). Given that the residues identified as governing receptor utilization are spatially close to each other but far from the NeoCoV/PDF-2180 footprint on ACE2 along with their different phenotypes of sensitivity to specific ACE2 glycans, we hypothesized that MOW15-22 and PnNL2018B might rely on a distinct mode of receptor engagement (Figure 4D-E).

Viruses	Determinant positions (ACE2)	N-glycan contribution	Favorable residues	Unfavorable residues
NeoCoV	54-56	(+)	NIT	NII
	305	NA	E	K/Q
	329-330	(+)	NNS	EKS
	337-340	NA	SDGR	GNVQ
MOW15-22	287	NA	K	Q
	432-434	(-)	YET/CET	NET
	589	NA	K/R	E
	597	NA	E/D	K

Figure 4. Host ACE2 tropism determinants for MOW15-22 and PnNL2018B. (**A**) MOW15-22 RBD and S_1 subunit binding to HEK293T cells transiently expressing ACE2 chimeras with sequence swaps between M.bla and P.dav ACE2s analyzed by fluorescence microscopy. (**B**) Sequence swaps between P.pip ACE2 N432C mutant and P.nat ACE2 and their impact on indicated viral antigen bindings. (**C**) Sequence swaps between P.pip ACE2 N432C/E589K/K598E mutant and P.nat ACE2 and their impact on indicated viral antigen bindings. (**D**) Summary of the critical determinants for MOW15-22 receptor recognition and host range determination identified through genetic swaps. Key residues discussed in the text are highlighted in blue on the P.pip ACE2 surface (PDB 7WPO). The N432-glycan is indicated in magenta. (**E**) Summary of ACE2 residues and glycans governing receptor utilization for MOW15-22 or PnNL2018B, respectively. The contribution of N-glycans in receptor binding and representative favorable/unfavorable residues in ACE2 orthologs are indicated. (+): support binding; (-): restrict binding; NA: not applicable (not glycosylation site). The P.pip ACE2 residue numbering is shown. Residue positions, $N432_{P.pip\ ACE2}$ glycans (in magenta), and swap strategies were indicated with the schematic below each panel in A, B, and C. Scale bars:100 μm. See also Figure S4-S5.

Structural basis for MOW15-22 binding with bat ACE2

To understand the molecular basis of ACE2 utilization by these recently identified merbecoviruses, we determined a cryo-EM structure of the MOW15-22 RBD bound to the dimeric P.dav ACE2 ectodomain at 2.8Å resolution (Figure 5A, Figure S6, and Table S1). Two MOW15-22 RBD extensions spanning residues 505-535 and 563-571 protrude at the distal end of the receptor-binding motif (RBM) to interact with the ACE2 peptidase domain (Figure 5A). Specifically, the MOW15-22 RBM interacts with a glycan-free P.dav ACE2 surface comprising residues P282, Y283, E285, K286, P287, P427, E428, D429, Y430, E431, E433, I434, L437, T534, P536, H538, R587, P588, N591, W592, E595, Q596 and S598, encompassing the identified host range-associated P.dav ACE2 residues (E285, Y430, R587, and E595) (Figure 4D). An average surface of ~850Å^2 is buried at the interface between the RBD and ACE2 with key interactions including (Figure 5B-C): (**i**) $K509_{MOW15-22}$ forming a salt bridge with $E595_{ACE2}$; (**ii**) a salt bridge triad involving $K286_{ACE2}$ with $D563_{MOW15-22}$ and $D567_{MOW15-22}$; (**iii**) a hydrogen bond network between $Y430_{ACE2}$, $T526_{MOW15-22}$, and $R571_{MOW15-22}$; (**iv**) hydrogen bonding between $N591_{ACE2}$ and $E523_{MOW15-22}$; (**v**) $Y283_{ACE2}$ that is hydrogen-bonded to the $D508_{MOW15-22}$ backbone carbonyl and tucked in between $V507_{MOW15-22}$ and $P511_{MOW15-22}$; and (**vi**) extensive hydrophobic interactions mediated by $V507_{MOW15-22}$, $I521_{MOW15-22}$, and $F524_{MOW15-22}$ with ACE2.

The MOW15-22 RBD largely adopts a canonical merbecovirus architecture comprising a core folded as a five-stranded antiparallel β-sheet with two α-helices and a four-stranded antiparallel β-sheet RBM with one α-helix (Figure 5D). However, the MOW15-22 RBD is set apart from other merbecoviruses due to its elongated and twisted distal RBM extensions, one of them being stabilized by the C506-C520 disulfide, which form the ACE2-binding surface, and to the elongated α-helix flanking the RBM on the opposite side. Furthermore, the MOW15-22 RBD harbors two N-linked glycans protruding from the base of the RBM, which are conserved in PnNL2018B S but not among other merbecoviruses (Figure 5D). The reduced P.dav ACE2 binding and utilization observed for the PnNL2018B RBD, relative to MOW15-22, is explained by the conservation of 14 out of 21 ACE2-interacting residues (Figure 5D and Table S2).

Strikingly, the MOW15-22 RBD recognizes ACE2 at a site located more than 50Å away from the sites recognized by SARS-CoV-2, NL63, or NeoCoV (Figures 5E-5F). This suggests that ACE2-utilization was acquired multiple times independently among these coronaviruses, including at least twice for MERSr-CoVs. Our data reveal that P.dav ACE2 lacks a glycan at

position Y430, which is otherwise conserved in several other ACE2 orthologs (e.g., hACE2/P.pip N432 and M.bla N428 glycans), and would impede MOW15-22 RBD binding sterically, concurring with the lack of binding observed to a P.dav Y430N glycan knockin mutant (Figure 5C). Accordingly, ACE2 orthologs lacked this glycan (e.g. P. par ACE2, C.jac, and L.bor ACE2) supporting efficient MOW15-22 or PnNL2018B S pseudovirus entry into cells (Figures 3A-3B), and these viruses are therefore not expected to acquire efficient hACE2 utilization without major adaptations.

To further assess the contribution of the identified receptor binding determinants, we analyzed the impact of residue changes at key positions of P.nat, P.pip, human, P.kuh, and C.jac orthologs (Figure 5G). P.nat ACE2 mutants harboring F285G and W594G mutations with reduced hydrophobicity dampened binding to MOW15-22 or PnNL2018B, while three hACE2 residue substitutions (Q287K, N432C, and E589K) and two P.kuh ACE2 residue substitutions (K436I and E589K) promoted detectable binding of MOW15-22 and PnNL2018B (Figures 5H 5I and Figure S7A-7B). The C.jac ACE2 S432N glycan knock-in mutation abolished receptor function of the sole primate ortholog supporting entry, further underscoring the key role of this oligosaccharide as a tropism barrier. Conversely, the C.jac ACE2 Q287K and Q598L point mutations with additional interactions significantly improved binding affinity to both MOW15-22 and PnNL2018B (Figure S7C-7D). Concurring with the structural data, introducing unfavorable changes to the MOW15-22 RBM, either by reducing interaction or abolishing the C506-C520 disulfide, significantly reduced RBD binding and pseudovirus entry in P.nat or P.dav ACE2-expressing HEK293T cells (Figures 5J-5K and Figure S7E).

Figure 5. Structural basis for MOW15-22 binding with bat ACE2. (**A**) CryoEM structure of the MOW15-22 RBD bound to the P. dav ACE2 ectodomain. (**B-C**) Zoomed-in views of key interactions mediating MOW15-22 RBD(yellow) binding to P.dav ACE2 (dark green). Selected salt bridges and hydrogen bonds are shown as black dotted lines. (**D**) Comparison of merbecovirus RBD architectures. MOW15-22, MERS-CoV (PDB 6Q04), HKU4 (PDB 4QZV), NeoCoV (PDB 7WPO). RBD core subdomains are indicated in gray. (**E**) Comparison of the binding modes of the MOW15-22, NeoCoV (PDB 7WPO), SARS-CoV-2/SARS-CoV-1 (PDB 7TN0), and NL63 (PDB 3KBH) RBDs on bat (not shown for clarity) or hACE2 (PDB 6M1D, B0AT1 not shown for clarity). (**F**) RBD footprints of the four ACE2-using coronaviruses. N-terminus labeled in green. (**G**) Schematic of critical residues responsible for species-specific receptor recognition in selected ACE2 orthologs. (**H**) P. nat ACE2 mutants with reduced MOW15-22 RBD or S_1 subunit binding due to unfavorable substitutions to hydrophobic interactions. (**I**) human or P.kuh ACE2 mutants with improved MOW15-22 RBD or S_1 subunit binding due to favorable substitutions mapping to the N432-glycosylation site, hydrophobic interactions, or polar interactions. (**J-K**) MOW15-22 RBM mutants with reduced ability to utilize P.dav and P. nat ACE2 orthologs for RBD binding (J) and pseudovirus entry (K). Scale bars:100 μm. Binding in panels H, I, and J was analyzed by fluorescence microscopy. MEAN ± SD and unpaired two-tailed t-tests for K. n=3 biological replicates.
See also Figure S6-S7.

Characterization and inhibition of MOW15-22 and PnNL2018B ACE2-mediated entry

MERS-CoV efficiently utilizes endogenous host proteases, such as furin and TMPRSS2, for cellular entry.[35] MERS-CoV S harbors a polybasic (furin) cleavage site at the junction between the S_1/S_2 subunits, leading to proteolytic processing during biogenesis.[36] Conversely, MOW15-22 and PnNL2018B S glycoproteins lack an S_1/S_2 furin cleavage site and remain uncleaved upon incorporation into VSV pseudovirus particles, suggesting distinct protease requirements for cellular entry compared to MERS-CoV (Figures 6A-B). Accordingly, MOW15-22 and PnNL2018B S glycoproteins are highly dependent on the addition of exogenous trypsin to promote cell-cell fusion, indicating a potential reliance on endocytic pathways for cellular entry in Caco2 cells, which endogenously express TMPRSS2 at their surface (Figures 6C-6D).[37]

To compete viral entry using soluble ACE2, we designed P.nat ACE2 recombinant ectodomain chimeras carrying P. dav ACE2 residues 404-432 without (P. nat ACE2.M1) or with the Q598L substitution (P.nat ACE2.M2) which exhibited enhanced binding affinity for both MOW15-22 and PnNL2018B RBD-Fc constructs relative to wildtype P. nat ACE2, especially the off-rate that is most critical for neutralization (Figures 6E-6F, Figure S8A-C).[9] Cryo-EM analysis of the complex between the MOW15-22 RBD and P. nat ACE2.M2 confirmed the retention of a native binding mode and rationalized the enhanced binding (Figure 6G, Figure S9, and Table S1). We observed concentration-dependent soluble P.nat ACE2.M2-mediated inhibition of MOW15-22 and PnNL2018B S pseudoviruses with a half-maximal inhibitory concentration (IC_{50}) of 19.22 ng/ml, outperforming wild-type P.nat and P.dav ACE2s (Figure 6H), further confirming the potential of these ACE2 mutants for developing entry inhibitors.

To identify other countermeasures against these divergent merbecoviruses, we assessed the ability of entry inhibitors, monoclonal antibodies, nanobodies, and peptide fusion inhibitors to block S-mediated entry in HEK293T cells expressing P.dav ACE2 (Figures 6I-6R and Figure S10). Endosomal entry inhibitors, such as E64D, bafilomycin A, and Apilimod, inhibited pseudovirus entry in a dose-dependent manner (Figures 6I-6K) whereas the TMPRSS2 inhibitor camostat did not, likely due to the lack of TMPRSS2 expression in this cell line or their low

efficiency in using this protease (Figure 6L).[37–39] These results confirmed that MOW15-22 S- and PnNL2018B S-mediated entry can occur through an endosomal pathway in these target cell types. Whereas MERS-CoV RBD-directed nanobodies were ineffective against MOW15-22 and PnNL2018B S pseudoviruses (Figure S10A),[40,41] broadly neutralizing S_2 subunit-directed monoclonal antibodies targeting the stem helix (S2P6) or the fusion peptide/S_2' cleavage site (76E1) retained activity, consistent with the conservation of the targeted S_2 epitopes and the marked divergence of their RBDs (Figures 6N-6R and Figures S10B-D).[42,43] Furthermore, the HR1-directed EK1C4 peptide and the hACE2-targeting antibody H11B11 efficiently neutralized the entry of MOW15-22 and PnNL2018B pseudoviruses (Figures 6M, Q, R and Figure S10E).[39,44] Collectively, these data demonstrate that neutralizing antibodies targeting the fusion machinery (S_2 subunit) or the host receptor prevent ACE2-mediated entry and could be potentially used as countermeasures against these viruses.

Figure 6. Characterization and inhibition of MOW15-22 and PnNL2018B ACE2-mediated entry. (**A**) Quantification of S glycoprotein incorporation in VSV pseudotypes analyzed by Western blot detecting the C-terminal-fused HA tags**.** VSV-M was used as a loading control. (**B**) Sequence analysis of S glycoprotein S_1/S_2 junction (dashed box) from the indicated viruses, with arginine highlighted in red. (**C-D**) MOW15-22 S mediated cell-cell fusion using Caco2 cells overexpressing P.nat ACE2 upon addition of TPCK-treated trypsin. The signal resulting from the reconstitution of split sfGFP (C) or RLuc (D) is indicated. (**E-F**) BLI analyses of binding kinetics of soluble dimeric ACE2 ectodomains from P.nat ACE2.M1 (E) and P.nat ACE2.M2 (F) carrying P.dav ACE2 equivalent residues (red) to immobilized MOW15-22 RBD-hFc. Global fitting to the data using a 1:1 binding model is shown in black. (**G**) Superimposition of the cryoEM structures of the MOW15-22 RBD (gold) in complex with P.dav ACE2 (not shown for clarity) and of the MOW15-22 RBD (blue) bound to P.nat ACE2.M2 (dark green) superimposed based on the ACE2 peptidase domain. (**H**) Dose-dependent inhibition of MOW15-22 pseudovirus entry by soluble P.dav, P.nat, P.nat.M2 and human ACE2s in Caco2 cells stably expressing P.nat ACE2. IC_{50} values are indicated. N.D.: Not detectable. (**I-M**) Inhibitory activities of small compounds or peptide inhibitors against MOW15-22 and PnNL2018B pseudoviruses using HEK293T cells stably expressing P.dav ACE2 (for MOW15-22) or L.bor ACE2 (for PnNL2018B). Endosomal acidification inhibitor bafilomycin A1 (Baf-A1) (I), and the PIKfyve inhibitor Apilimod (J), cathepsin L inhibitor E64d (K), TMPRSS2 inhibitor camostat (L), S_2-targeting HR1 peptide fusion inhibitor EK1C4 (M). (**N**) Schematic representation of the epitopes targeted by the antibodies used in the pseudovirus neutralization assays. (**O-R**) Dose-dependent inhibition of MOW15-22 and PnNL2018B pseudoviruses by S2P6 and 76E1 (O-P), or H11B11 (Q-R) in the indicated cell lines.n=2 biological replicates for H-R. Scale bars:200 μm. See also Figure S8-S10.

MOW15-22 and PnNL2018B S-mediated propagation supported by ACE2

Finally, we utilized a reverse genetic system to create propagation-competent VSV (pcVSV) to characterize the MOW15-22 and PnNL2018B S glycoprotein-mediated propagation supported by different ACE2. The VSV-G genes in these recombinant viruses were replaced with the MOW15-22 or the PnNL2018B S glycoprotein gene, with an additional GFP gene for visualization (Figure 7A-7B). We successfully rescued the viruses and observed a dose-dependent trypsin enhancement of amplification in Caco2 cells expressing P.nat ACE2, as evidenced by syncytia formation (Figure 7C). Although viral propagation was not observed in Caco2 cells overexpressing hACE2, abrogation of the N432 glycan (via the N432C mutation) resulted in detectable amplification of pcVSV-MOW15-22. Furthermore, the E589K or Q287K/E589K mutations in the hACE2 N432C background further improved the amplification of pcVSV-MOW15-22 (Figure 7D) and enabled the detection of pcVSV-PnNL2018B amplification (Figure 7E), concurring with our binding data (Figure 5H). Despite the restriction on viral recognition imposed by the hACE2 N432 glycan, the two viruses could nevertheless utilize P.nat ACE2 harboring a C432N (glycan knockin) mutation, as demonstrated by the observed viral propagation and S_1 subunit binding (Figures 7F-7G and Figure S5E-F). The observed phenotype indicates that these viruses hold the potential to partially overcome the steric tropism barrier imposed by the N432 glycan, possibly through compensatory interactions in other ACE2 regions.

Figure 7. MOW15-22 and PnNL2018B S-mediated propagation supported by ACE2. (**A**) Genetic organization and amplification of pcVSV-MOW15-22 and pcVSV-PnNL2018B. (**B**) Immunofluorescence analyzing the expression of P.nat and human ACE2 mutants stably expressing in Caco2 cells. (**C**) GFP signal from TPCK-trypsin enhanced amplification of pcVSV-MOW15-22 and pcVSV-PnNL2018B in Caco2 cells with or without the expression of P.nat ACE2 at 24hpi. (**D-E**) Propagation of pcVSV-MOW15-22(D) and pcVSV-PnNL2018B(E) in Caco2 cells stably expressing hACE2 carrying N432C(1M), Q287K/N432C(2M), Q287K/N432C/E589K(3M) mutants, with P.nat ACE2 as a positive control. (**F**) Representative fluorescence images of pcVSV-MOW15-22 and pcVSV-PnNL2018B amplification in Caco2 cells expressing P.nat ACE2 or P.nat ACE2 C432N mutants at 12, 24 and 36-hour post-infection (hpi). (**G**) pcVSV-MOW15-22 propagation kinetics in Caco2 cells expressing P.nat ACE2 or P.nat ACE2 C432N mutants at the indicated time points. MEAN ± SD and n=3 biological replicates. RFU: relative fluorescence unit. TPCK-treated trypsin was present in the cells at 20 µg/mL in D, E, F, and G. Scale bars:100 µm for B and 200 µm for C-E.

DISCUSSION

Coronaviruses exhibit remarkable variations in RBD sequences, resulting in diverse receptor usage across different viruses.[45] While coronaviruses within the same genus or subgenus typically share similar RBD core structures, RBM variability can result in entirely different receptor usage.[9,46,47] Conversely, phylogenetically distant coronaviruses can convergently evolve to engage the same receptor during evolution. For example, APN is a receptor shared by many α-coronaviruses and δ-coronaviruses,[47–50] whereas ACE2 serves as a functional receptor for three viral subgenera that belong to α-coronaviruses (*Setracoviruses*) and β-coronaviruses (*Sarbecoviruses* and *Merbecoviruses*) .[21,51–53]

The discovery of ACE2 usage in European bat MERSr-CoVs expands the diversity of ACE2-using merbecoviruses beyond NeoCoV and PDF-2180 and reveals that coronaviruses belonging to the same subgenus or even species (e.g. MOW15-22 and NeoCoV) can engage entirely distinct surfaces of a same receptor (e.g. P.dav ACE2). This study emphasizes the need to characterize receptor usage and binding modes experimentally, instead of solely relying on *in silico* predictions.

Our data further support the hypothesis that ACE2 usage was convergently opted by different coronaviruses with remarkable differences in their RBD or RBM sequences and structures. Furthermore, a recent report showed that HKU5 exhibited a specialized adaptation to its host ACE2 (P.abr), suggesting a potential fifth ACE2 binding mode that remains to be elucidated.[32] Notably, these ACE2-using merbecoviruses with distinct binding modes evolved in three different Old World continents, suggesting the convergent evolution of ACE2 adaptation and the importance of investigating the global prevalence and distribution of ACE2-using merbecoviruses. This ACE2 preference likely confers certain evolutionary advantages in transmission, as exemplified by the highly transmissible SARS-CoV-2 omicron variants.[54,55] However, a recent study reported that PnNL2180B (PN-βCoV) primarily exhibits intestinal tropism in its natural bat host, suggesting a potential fecal-oral route used by these viruses in bats .[28] Given that airborne transmission is the major route of all known ACE2-using human coronaviruses, it is important to investigate whether tissue tropism and transmission route changes when ACE2-using viruses jump from bats to other mammals, especially humans.

Interestingly, ACE2 glycans can play contrasting roles in host range determination of ACE2-using merbecoviruses. For example, unlike the positive role of ACE2 glycans (e.g. N54 and N329 glycans) in NeoCoV and PDF-2180 recognition[9], the N432 glycan in interaction

interface restricted efficient MOW15-22 and PnNL2018B binding, contributing to narrower ACE2 tropism, particularly for PnNL2018B for which we only detected binding to its host P. nat ACE2[9]. Common marmoset (C.jac) ACE2, which is the only primate ortholog supporting MOW15-22 S-mediated entry, lacks an N432 glycan which is otherwise conserved in other primate ACE2s, explaining its distinct phenotypes. This specialized adaptation to host ACE2s lacking an N432-glycan is expected to prevent ACE2 utilization from many other species, including humans. However, the efficient amplification of pcVSV-MOW15-22 and pcVSV-PnNL2018B supported by a P.nat ACE2 mutant (C432N) harboring an N432 glycan suggests putative evolutionary pathways by which these viruses could overcome this obstacle and the usefulness of the countermeasures identified here. Notably, the VM314 viral sequence, identified in samples collected in the Netherlands in 2008 based on an RNA-dependent RNA polymerase (RdRp) gene fragment that is phylogenetically close to MOW15-22 and PnNL2018B, may already be adapted to its host (P.pip) ACE2 carrying an N432 glycan. Future studies should be conducted to obtain the corresponding S glycoprotein sequences and to verify whether these viruses recognize ACE2 in a similar way to MOW15-22.[56]

Previous studies proposed that the emergence of the DPP4-using MERS-CoV may be associated with a receptor switch from ACE2 to DPP4 through recombination.[9] This could have occurred between an ACE2-using bat MERSr-CoV and a yet-to-be-identified DPP4-using merbecovirus, such as HKU4 or CoV-422.[15,57] We note that CoV-422 was also suggested to have originated from recombinations leading to receptor switch[15]. Additionally, it has been suggested that ACE2-using merbecoviruses might have arisen through recombination between ancestral viruses of bats and hedgehogs.[58] However, thus far, we and others have not detected solid evidence of ACE2 usage in testing bat or hedgehog ACE2 for coronaviruses HKU31 or other EriCoVs.[9,59,60] These observations raise intriguing and important questions regarding the evolution trajectory of receptor usages of merbecoviruses and whether ACE2 receptor usage is the more ancestral trait for merbecoviruses than DPP4.

Our study profoundly changes our understanding of ACE2-using merbecoviruses by identifying and characterizing two bat merbecoviruses with previously unknown ACE2 binding modes, involving binding sites located 50Å away from that recognized by any other coronaviruses. This discovery also underscores the likelihood of the existence of other yet-to-be-discovered ACE2-using merbecoviruses, further expanding the diversity and geographic distribution of these viruses with spillover potential. Although the pathogenicity and transmission abilities of these viruses remain unclear, enhanced surveillance along with identification of viral inhibitors are warranted to proactively detect and prepare for potential zoonosis caused by ACE2-using merbecoviruses.

Acknowledgments:

This study was supported by the National Key R&D Program of China (2023YFC2605500 to H.Y. and Z.L.S.), the National Natural Science Foundation of China (NSFC) projects (82322041, 32270164, 32070160, 32188101 to H.Y., 323B2006 to C.B.M.), the Fundamental Research Funds for the Central Universities (2042023kf0191, 2042022kf1188 to H.Y.), Natural Science Foundation of Hubei Province (2023AFA015 to H.Y.), the National Institute of Allergy and Infectious Diseases (P01AI167966, DP1AI158186 and 75N93022C00036 to D.V.), an Investigators in the Pathogenesis of Infectious Disease Awards from the Burroughs Wellcome Fund (D.V.), the University of Washington Arnold and Mabel Beckman cryoEM center and the National Institute of Health grant S10OD032290 (to D.V.). D.V. is an Investigator of the Howard Hughes Medical Institute and the Hans Neurath Endowed Chair in Biochemistry at the University of Washington. The CAS Pioneer Hundred Talents Program to Z.D.

We are grateful to Michael Hiller (LOEWE Centre for Translational Biodiversity Genomics), Camila Mazzoni (Leibniz Institute for Zoo and Wildlife), and Huabin Zhao (Wuhan University) for sharing the ACE2 and DPP4 coding sequences of *Pipistrellus nathusii*. We thank Qiang Ding (Tsinghua University) and Qihui Wang (CAS Key Laboratory of Pathogenic Microbiology & Immunology, China) for sharing some mammalian ACE2 plasmids. We thank Mingzhou Chen (Hubei University) for providing the recombinant vaccinia virus expressing T7 RNA polymerase (vvT7). We thank the Center for Instrumental Analysis and Metrology of Wuhan Institute of Virology for providing technical assistance in cryoEM sample preparation and data collection, as well as the core facilities of the Key Laboratory of Virology, Wuhan University for providing instrumental and technical support.

Author contributions:
H.Y., Y.J.P., M.A.T., D.V., Z.D., Z.L.S, C.B.M., C.L., and Q.X., conceived the project; J.C., C.B.M., and C.L. conducted bioinformatic analysis; C.B.M. and C.L., designed and carried out experiments to generate clones, purify proteins, dissect host range determinants and test entry inhibitors. C.B.M., C.L., Y.J.P, Q.X., X.Yu., P.L., J.Y.S., M.L.H, F.T., Y.H.S., and Y.M.C. established assays and methods used in this study; X.Yang rescued the pcVSV-MOW15-22 and pcVSV-PnNL2018B viruses. J.L., J.B., and C.S. produced recombinant glycoproteins and carried out binding and entry assays. Y.J.P. carried out cryo-EM sample preparation, data collection, and processing of the P.dav ACE2-MOW15-22 RBD complex. D.A. helped with data collection. Y.J.P. and D.V. built and refined the P.dav ACE2-bound and MOW15-22 RBD structure; J.T. carried out cryo-EM sample preparation, data collection, and processing of the P.nat ACE2.M2-MOW15-22 RBD complex; Z.D. and J.T. built and refined the P.nat.ACE2.M2 ACE2-MOW15-22 RBD complex structure. H.Y., and D.V., wrote an initial draft of the manuscript with input from all authors.

Competing interests: H.Y. has submitted a patent application to the China National Intellectual Property Administration for the utilization of propagation-competent VSV to evaluate amplification of ACE2-using merbecoviruses in bat ACE2-expressing cells.

1 **STAR ★ METHODS**
2 **KEY RESOURCES TABLE**
3

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Alexa Fluor 488-conjugated goat anti-human IgG	Thermo Fisher Scientific	Cat#A11013
Mouse anti-FLAG antibody M2	Sigma-Aldrich	Cat#F1804
Anti-VSV-M(23H12)Antibody	kerafast	Cat# EB0011
Anti-HA-tag mAb	MBL	Cat#MBL-M180-3
Alexa Fluor 594-conjugated goat anti-mouse IgG	Thermo Fisher Scientific	Cat#A32742
Goat anti-human IgG(H+L) Cross Absorbed Secondary Antibody	Thermo Fisher Scientific	Cat#A11013
β-Tubulin mAb	Immmuno Way	Cat#YM3030
H11B11	Du et al. [44]	N/A
76E1	Sun et al. [43]	N/A
S2P6	Pinto et al. [42]	N/A
Bacterial and virus strains		
VVT7	Mingzhou Chen's lab, Wuhan University	N/A
pcVSV-MOW15-22	This study	N/A
pcVSV-PnNL2018B	This study	N/A
VSV-dG-GFP-fLuc	Ma et al. [24]	N/A
Chemicals, peptides, and recombinant proteins		
TPCK-treated trypsin	Sigma-Aldrich	Cat#T8802
Polybrene	Beyotime	Cat#C0351
Dulbecco's Modified Eagle Medium	Monad, China	Cat#CC00101S

Penicillin/Streptomycin	Biosharp	Cat#BL505A
Fetal Bovine Serum	Excell	Cat#FSP500-11I323
GeneTwin reagent	Biomed	Cat#TG101
Trypsin-EDTA (0.25%)	Gibco	Cat#25200072
SMM 293-TII Expression Medium	Sino Biological	Cat#M293TII
Pierce Protein A/G Plus Agarose	Thermo Scientific	Cat#20424
1M Tris-HCI, pH 8.0	Thermo Scientific	Cat#15568025
Strep-Tactin XT 4Flow high-capacity resin	IBA	Cat#2-5030-002
Hanks' Balanced Salt Solution	ServiceBio	Cat#G4204
Poly-L-lysine hydrobromide	Sigma	Cat#P1399
PMSF	Beyotime	Cat#ST506
Triton X-100	Solarbio	Cat#T8200
GelGreen	Biosharp	Cat#BS355A
D-Biotin	Shanghai yuanye Bio-Technology Co., Ltd	Cat#S13004
EK1-C4 peptide	LuLu's lab, Fudan University	N/A
E64d	Aladdin	Cat#E123225
Camostat	MedChemExpress	Cat#HY-13512
Bafilomycin A1	MedChemExpress	Cat#HY-100558
Critical commercial assays		
Bright-Glo Luciferase Assay Kit	Promega	Cat#E2650
EnduRen live-cell substrate	Promega	Cat#E6481
Omni-ECL Femto Light Chemiluminescence Kit	EpiZyme	Cat#SQ201
Omni-Easy Instant BCA Protein Assay Kit	Epizyme	Cat#SZJ102

Genejet gel extraction kit	Thermo	Cat#K0692
Deposited data		
P.nat ACE2.M2-bound MOW15-22 RBD cryo-EM structure	**This study**	**PDB:8ZUF**
P.dav ACE2-bound MOW15-22 RBD cryo-EM structure	**This study**	**PDB: 9C6O**
P.nat ACE2 and P.nat DPP4 coding sequences	**This study**	**https://data.mendeley.com/preview/mw3gyj7ncr?a=6a5b0b4f-ad01-4f5c-b573-f73a89404b1a**
Original western blot images	**This study**	**https://data.mendeley.com/preview/wjns8j3z6y?a=3d8afefb-cd8b-4bba-aeab-3948f7d94236**
CryoEM data collection, processing, and model refinement statistics.	**This study**	**https://data.mendeley.com/preview/xxksdm3dhn?a=cbbba022-4359-4572-9795-f0e4c7c8c7ce**
Experimental models: Cell lines		
HEK293T	**ATCC**	**CRL-3216**
HEK293T	**ATCC**	**CRL-11268**
Caco2	**ATCC**	**HTB-37**
BHK21	**ATCC**	**CCL-10**
I1-Hybridoma cell lines	**ATCC**	**CRL-2700**
Expi293F	**ATCC**	**A14527**
Recombinant DNA		
pVSV-dG-GFP-MOW15-22-S	**This paper**	**N/A**
pVSV-dG-GFP-PnNL2018B-S	**This paper**	**N/A**
rLucN(1-155)-sfGFP1-7(1-157)	**This paper**	**N/A**
sfGFP8-11(158-231)-rLuc(156-311)	**This paper**	**N/A**
Software and algorithms		

MEGA-X version 10.1.8	Kumar et al. [61]	https://www.megasoftware.net/
IQ-TREE	Minh et al. [62]	http://igtree.cibiv.univie.ac.at/
iTOL (v6)	Letunic and Bork [63]	https://itol.embl.de/
UCSF ChimeraX (V.1.7.1)	Goddard et al. [64]	https://www.rbvi.ucsf.edu/chimerax
ColabFold v1.5.5: AlphaFold2 using MMseqs2	Mirdita et al. [65]	https://colab.research.google.com/github/sokrypton/ColabFold/blob/main/AlphaFold2.ipynb
PDBePISA	Krissinel and Henrick [66]	https://www.ebi.ac.uk/pdbe/pisa/pistart.html
ClustalW	Thompson et al. [67]	https://www.genome.jp/tools-bin/clustalw
Other		
MERS-CoV RBD structure	Lu et al. [17]	PDB:4KQZ
HKU4 RBD structure	Wang et al. [13]	PDB:4QZV
NeoCoV RBD structure	Xiong et al. [9]	PDB:7WPO
HKU5 RBD structure	Han et al. [68]	PDB:5XGR

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Huan Yan (严欢) (huanyan@whu.edu.cn)

Materials availability

All plasmids generated in this study are available with a completed Materials Transfer Agreement. Correspondence and requests for materials can be addressed to the lead contact.

Data and code availability

- The cryoEM structures have been deposited to the electron microscopy data bank and protein data bank with accession numbers EMD-45253, PDB-9C6O (P.dav ACE2-bound MOW15-22 RBD) and EMD-60483, PDB-8ZUF (P.nat ACE2.M2-bound MOW15-22 RBD). Original western blot images, cryo-EM data statistics, P.nat ACE2 and P.nat DPP4 coding sequences have been deposited at Mendeley and are publicly available as of the date of publication. The DOI is listed in the key resources table. Accession numbers are listed in the key resources table. Microscopy data reported in this paper will be shared by the lead contact upon request.
- This paper does not report original code.
- Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

Cell lines and culture conditions

HEK293T (CRL-3216), HEK293T (ATCC, CRL-11268) Caco2 (HTB-37), BHK21 (CCL-10), and I1-Hybridoma (CRL-2700) cell lines were obtained from the American Type Culture Collection (ATCC). Expi293F (A14527) cells were obtained from Thermo Fisher Scientific. These cells were maintained in Dulbecco's Modified Eagle Medium (DMEM, Monad, China) supplemented with 1% PS (Penicillin/Streptomycin) and 10% Fetal Bovine Serum (FBS). The I1-Hybridoma cell line, which produces a neutralizing antibody targeting the VSV glycoprotein (VSV-G), was cultured in Minimum Essential Medium (MEM) with Earles's balances salts, 2.0 mM of L-glutamine (Gibico), and 10% FBS. All cell lines were cultured at 37℃ with 5% CO_2 and routinely passaged every 2-3 days. HEK293T or Caco2 stable cell lines overexpressing various receptors were generated using lentivirus transduction and antibiotic selection. The stable cells based on HEK293T or Caco2 were selected and maintained in the growth medium with puromycin (1 μg /ml).

METHOD DETAILS

Plasmids and vectors

Plasmids expressing wild-type (WT) or mutated bat and non-bat mammalian ACE2 orthologs [24,33] were constructed by inserting human codon-optimized sequences with/without specific mutations into a lentiviral transfer vector (pLVX-EF1a-Puro, Genewiz) with C-terminus 3×FLAG tags (DYKDHD-G-DYKDHD-I-DYKDDDDK) and single FLAG tags (DYKDDDDK) for non-bat mammalian ACE2 orthologs[24]. P.nat ACE2 and DPP4 coding sequences are retrieved from the genome of *Pipistrellus nathusii* (GCA_963693515.1), P.nat ACE2 and P.nat DPP4 coding sequences have been deposited at Mendeley with DOI listed in the key resources table. [31]. For pseudovirus production, human codon-optimized spike sequences of MOW15-22 (USL83011.1), PnNL2018B (WDE20340.1), SARS-CoV-2 (YP_009724390.1) carrying D614G mutation, MERS-CoV (YP_009047204.1), HKU4 (AWH65899), NeoCoV (AGY29650.2) and HKU31 (QGA70692.1) were cloned into the pCAGGS vector with C-terminal deletions (residues 13-15) for improving the pseudovirus assembly efficiency. The DNA fragments for cloning ACE2 chimera or mutants were generated by overlap extension PCR or gene synthesis and verified by commercial DNA sequencing. For the expression of recombinant CoVs RBD-hFc fusion proteins, plasmids were constructed by inserting NeoCoV RBD (residues 380-585),

MOW15-22 RBD (residues 360-610aa), PnNL2018B RBD (residues 361-606) coding sequences into the pCAGGS vector containing an N-terminal CD5 secretion signal peptide (MPMGSLQPLATLYLLGMLVASVL) and C-terminal hFc-twin-strep tandem tags for purification and detection. Plasmids expressing soluble ACE2 ectodomain proteins were generated by inserting sequences from hACE2 (residues 18-740)，P.dav ACE2 (residues 18-738)，and P.nat ACE2 (residues 18-739) into pCAGGS vector, with an N-terminal CD5 secretion signal peptide and a C-terminal twin-strep-3×FLAG tag (WSHPQFEKGGGSGGGSGGGSAWSHPQFEK-GGGRSDYKDHDGDYKDHDIDYKDDDDK)[9]. The construct for expressing soluble P.nat ACE2.M1 was generated by overlapping PCR to replace residues of 404-430 with P.dav ACE2 corresponding sequences. The P.nat ACE2.M2 was generated by further introducing a Q598L point mutation to P.nat ACE2.M1. For CryoEM analysis, the P.dav ACE2 ectodomain (residues 17-723) construct harbors an N-terminal signal peptide (MPMGSLQPLATLYLLGMLVASVL) with avi tag followed by seven residues flexible linker and C-terminal octa-histidine tag was codon-optimized, synthesized, and inserted the pcDNA3.1(+) vector by Genscript. The MOW15-22 RBD construct encoding S residues 351-599 containing N-terminal signal peptide (MGILPSPGMPALLSLVSLLSVLLMGCVA), with an avi tag followed by eight residues flexible linker and C-terminal octa-histidine tag were codon optimized, synthesized, and inserted into the pcDNA3.1(+) vector by Genscript.

Protein expression and purification

HEK293T cells were transfected with corresponding plasmids using GeneTwin reagent (Biomed, TG101-01). Subsequently, the culture medium of the transfected cells was replenished with the SMM 293-TII Expression Medium (Sino Biological, M293TII) 4-6 hours post-transfection, and the protein-containing supernatant was collected every three days for 2-3 batches. Antibodies and recombinant RBD-hFc proteins were purified using Pierce Protein A/G Plus Agarose (Thermo Scientific, 20424). In general, Fc-containing proteins were enriched by the agarose, washed with wash buffer (100 mM Tris/HCl, pH 8.0, 150 mM NaCl, 1 mM EDTA), eluted using the Glycine buffer (100 mM in H_2O, pH 3.0), and immediately neutralized with 1/10 volume of 1M Tris-HCl, pH 8.0 (15568025, Thermo Scientific). Proteins with twin-strep tag were purified using Strep-Tactin XT 4Flow high-capacity resin (IBA, 2-5030-002), washed by wash buffer (100 mM Tris/HCl, pH 8.0, 150 mM NaCl, 1 mM EDTA), and then eluted with buffer BXT (100 mM Tris/HCl, pH 8.0, 150 mM NaCl, 1 mM EDTA, 50 mM biotin). All eluted proteins were concentrated using Ultrafiltration tubes, buffer-changed to PBS, and stored at -80℃. Protein concentrations were determined by the Omni-Easy Instant BCA Protein Assay Kit (Epizyme, ZJ102).

Antigen-hFc live-cell binding assay

RBD- or S_1-hFc live-cell binding assays were conducted following a previously described protocol[33]. The coronavirus RBD- or S_1-hFc recombinant proteins were diluted in DMEM at indicated concentrations and incubated with HEK293T cells transiently expressing different ACE2 for 30 minutes at 37℃ at 36 hours post-transfection. Subsequently, cells were washed once with Hanks' Balanced Salt Solution (HBSS) and incubated with 1 μg/mL of Alexa Fluor 488-conjugated goat anti-human IgG (Thermo Fisher Scientific; A11013) diluted in HBSS/1% BSA for 1 hour at 37℃. After another round of washing with HBSS, the cell nuclei were stained with Hoechst 33342 (1:10,000 dilution in HBSS) for 30 minutes at 37℃. The images were captured using a fluorescence microscope (MI52-N). The relative fluorescence intensities (RFUs)

of the stained cells were determined by a Varioskan LUX Multi-well Luminometer (Thermo Scientific). The heatmap presentation in Figure 2A, B, and S3C, D, was set based on RFUs subjected with background RLUs in cells without ACE2 expression. Color ranges and thresholds were adjusted to show binding efficiencies consistent with the fluorescence images captured by the microscope.

Flow cytometry

To analyze S_1-hFc binding through flow cytometry, HEK293T cells transiently expressing the indicated ACE2 orthologs were detached with 5 mM EDTA/PBS at 36 hours post-transfection. The cells were washed twice with cold PBS and incubated with MOW15-22 or PnNL2018B S_1-hFc proteins at 2 μg/mL concentrations at 4℃ for 30 minutes. Subsequently, cells were incubated with Alexa Fluor 488-conjugated goat anti-human IgG to stain the RBD (Thermo Fisher Scientific; A11013) at 4℃ for 1 hour. Afterward, cells were fixed with 4% PFA, permeabilized with 0.25% Triton X-100, blocked with 1% BSA/PBS at 4℃, and then incubated with mouse antibody M2 (Sigma-Aldrich, F1804) diluted in PBS/1% BSA for 1 hour at 4℃, followed by incubation with Alexa Fluor 594-conjugated goat anti-mouse IgG (Thermo Fisher Scientific; A32728) diluted in 1% BSA/PBS for 1 hour at 4℃. For all samples, 10,000 ACE2-expressing live cells (gated based on SSC/FSC and FLAG-fluorescence intensity and SSC/FSC) were analyzed using a CytoFLEX Flow Cytometer (Beckman).

Biolayer interferometry (BLI) binding assay

Protein binding kinetics was assessed using BLI assays with the Octet RED96 instrument (Molecular Devices) at 25℃, and shaking at 1000 rpm. Specifically, RBD-hFc or S1-hFc recombinant proteins were diluted to 20 μg/mL and immobilized on Protein A (ProA) biosensors (ForteBio, 18-5010), which were then incubated with soluble dimeric bat ACE2-ectodomain proteins, which were two-fold serial-diluted in kinetic buffer (PBST), starting from 1,000 nM. A well incubated with kinetic buffer (PBST) only serves as a background control. The kinetic parameters and the apparent binding affinities (due to ACE2 dimerization) between the RBD-hFc and ACE2 were analyzed using Octet Data Analysis software 12.2.0.20 with global curve fitting using a 1:1 binding model.

P.dav ACE2 binding to the MOW15-22 RBD (Figure 2C) was performed at 30℃ and shaking at 1,000 rpm. The biotinylated MOW-15-22 RBD was diluted to 10μg/mL in 10× Octet kinetics buffer (Sartorius) and loaded onto hydrated Streptavidin biosensors to 1nm shift, equilibrated in 10x Octet kinetics buffer for 60 seconds, and dipped into P. davyi ACE2 at 600nM, 200nM, 66.6nM and 22nM for 300s to observe association. Dissociation was followed by dipping biosensors in a 10× Octet kinetics buffer for 300s. Baseline subtraction was done by subtracting the response from unloaded SA tips dipping into 600nM P. davyi ACE2. In addition, association phases were aligned to 0 seconds and 0 response in Octet Data Analysis HT software and the processed results were exported. Global fitting using a 1:1 binding model was carried out using the Octet Data Analysis HT software Sensorgrams and plotted in GraphPad Prism10.

Cell-cell fusion assays

A cell-cell fusion assay based on dual-split proteins (DSPs) was conducted in Caco2 cells stably expressing ACE2 receptors. To assess the S glycoprotein-receptor interaction-mediated membrane fusion between cells, group A cells were transfected with S glycoprotein and rLucN(1-155)-sfGFP1-7(1-157) expressing plasmids, while group B cells were transfected with S glycoprotein and sfGFP8-11(158-231)-rLuc(156-311) expressing plasmids. After 12 hours of

transfection, both groups of cells were trypsinized, mixed, and seeded into a 96-well plate at 8×10^4 cells per well. Subsequently, the cells were washed once with DMEM and then incubated with DMEM with or without indicated concentrations of TPCK-treated trypsin (Sigma-Aldrich, T8802) for 10 min at room temperature at 24 h post-transfection. Six hours later, nuclei were stained with Hoechst 33342 (1:5,000 dilution in HBSS) for 30 min at 37 °C, and images of syncytia formation with green fluorescence were subsequently captured using a fluorescence microscope (MI52-N; Mshot). To measure live-cell luciferase activity after cell-cell fusion, 20 μM of EnduRen live-cell substrate (Promega, E6481) was added to the cells in DMEM and incubated for at least 1 hour before detection using the Varioskan LUX Multi-well Luminometer (Thermo Fisher Scientific).

Pseudovirus production

VSV-dG-based pseudovirus (PSV) carrying trans-complementary S glycoproteins from various coronaviruses were produced following a modified protocol as previously described[69]. Briefly, HEK293T cells were transfected with plasmids expressing coronaviruses S glycoproteins. At 24 hours post-transfection, cells were transduced with 1.5×10^6 TCID$_{50}$ VSV-G glycoprotein-deficient VSV expressing GFP and firefly luciferase (VSV-dG-GFP-fLuc, constructed and produced in-house) diluted in DMEM with 8 μg/mL polybrene for 4-6 hours at 37 °C. After three PBS washes, the culture medium was replenished with either DMEM+10% FBS or SMM 293-TII Expression Medium (Sino Biological, M293TII), along with the presence of the neutralizing antibody (from I1-mouse hybridoma) targeting the VSV-G to eliminate the background due to any remaining VSV-dG-GFP-fLuc. Twenty-four hours later, the pseudovirus containing supernatant was clarified through centrifugation at 12,000 rpm for 5 minutes at 4°C, aliquoted, and stored at -80°C. The TCID$_{50}$ of the PSV was calculated using the Reed-Muench method[70,71]. Heatmap presentation in Figure 3A, 3B, and S3A was set based on RLU with thresholds set based on entry background in cells without ACE2 expression. The value of P.nat ACE2 was set as a baseline of 100%, and the color of the largest value was set based on the ortholog showing the highest RLU, unless otherwise specified.

Single-round pseudovirus entry assay

Single-round pseudovirus entry assays were conducted using cells transiently or stably expressing different ACE2 orthologs. Approximately 3×10^4 trypsinized cells were incubated with pseudovirus (2×10^5 TCID$_{50}$/100 μL) in a 96-well plate to facilitate attachment and viral entry simultaneously. Before inoculation, pseudoviruses were typically treated with 100 μg/mL TPCK-trypsin (Sigma-Aldrich, T8802). Specifically, pseudoviruses produced in serum-free SMM 293-TII Expression Medium were incubated with TPCK-treated trypsin for 10 minutes at room temperature, and the proteolytic activity was neutralized by FBS in the culture medium. Intracellular luciferase activity (Relative light units, RLU) was measured using the Bright-Glo Luciferase Assay Kit (Promega, E2620) and detected with a GloMax 20/20 Luminometer (Promega) or Varioskan LUX Multi-well Luminometer (Thermo Fisher Scientific) at 18 hours post-infection.

pcVSV-CoV amplification assay

Plasmids for rescuing propagation-competent (pc) VSV-CoV expressing MOW15-22 and PnNL2018B S glycoproteins (pVSV-dG-GFP-MOW15-22-S and pVSV-dG-GFP-PnNL2018B-S) were generated by replacing the fLuc coding sequences with coronavirus spike coding sequences based on the vector pVSV-dG-GFP-fLuc. The propagation-competent recombinant VSVs were created by replacing the VSV-G gene with genomically encoded S glycoprotein genes and

additionally incorporating a GFP-expressing cassette for visualization. Reverse genetics was applied to rescue pcVSV pseudotypes expressing MOW15-22 and PnNL2018B S glycoproteins along with a GFP reporter, following a modified protocol from previous descriptions[69]. Specifically, BHK21 cells, seeded in a 6-well plate at 80% confluence, were inoculated with 5 MOI of recombinant vaccinia virus expressing T7 RNA polymerase (vvT7, a kind gift from Mingzhou Chen's lab, Wuhan University) for 45 minutes at 37°C. After removing vvT7, cells were subsequently transfected with pVSV-dG-GFP-S vector plasmids and helper plasmids (pVSV-dG-GFP-S: pBS-N: pBS-P: pBS-G: pBS-L=5:3:5:8:1). pcVSV-CoV containing supernatant (P0) was collected 48 hours post-transfection and 0.22-μm filtered to remove vvT7. For enhanced virus amplification, the passage 1 (P1) virus carrying VSV-G proteins was produced by inoculating P0 supernatant with Caco2 cells 24 hours post-transfection of plasmids expressing VSV-G protein. Subsequently, P1 viruses were further amplified in Caco2 cells stably expressing P.dav ACE2 or L.bor ACE2, without the ectopic expression of VSV-G and in the presence of anti-VSVG (I1-Hybridoma supernatant), producing passage 2 (P2) viruses carrying MOW15-22 and PnNL2018B S glycoproteins, respectively. In a typical virus amplification assay, 3×10^4 trypsinized cells stably expressing the indicated ACE2 were incubated with pcVSV-CoV (1×10^4 $TCID_{50}$/100 μL) in a 96-well plate with or without TPCK-treated trypsin treatment. GFP images or RFU were collected at indicated time points post-infection by a fluorescence microscope (MI52-N) or a Varioskan LUX Multi-well Luminometer (Thermo Fisher Scientific).

Pseudotyped virus neutralization assays
For soluble ACE2 neutralization assays, serial dilutions of recombinant proteins were prepared in DMEM. Pseudotyped viruses (1×10^5 $TCID_{50}$ per well) were mixed with 25 μl of each dilution for 1 hour at 37 °C, and then incubated P.nat ACE2-expressing Caco2 cells seeded at 2×10^4 cells per well in a 96-well plate. After 16–20 hours post-infection, the luciferase activity was measured as described in the single-round pseudotype virus entry assays. For S2P6, 76E1 antibody neutralization assays, pseudoviruses were incubated with 3-fold serial dilutions of antibodies in DMEM for 1 hour at 37 °C, and then applied to HEK293T stably expressing the P.dav or L.bor ACE2. For the H11B11 neutralization assay, Caco2 cells stably expressing the hACE2.3M were seeded on the 96-well plate one day before infection, 3-fold serial dilutions of H11B11 were diluted in the culture medium and then incubated with the cells for 1 hour at 37 °C. Pseudoviruses were then added to the cells with the previous H11B11-containing medium. Neutralizing efficiencies were assessed at 18 hours post-infection by detecting the intracellular luciferase activity (RLU).

Western blot
For detecting the cellular expression of ACE2 or DPP4 receptors with C-terminal FLAG tags, cells at 24 hours post-transfection were lysed in 1% TritonX/PBS+1 mM PMSF (Beyotime, ST506) for 10 minutes at 4°C. The lysate was clarified after centrifugation at 12,000 rpm for 5 minutes at 4°C and then incubated at 98°C for 10 minutes after mixing with the 1/5 volume of 5×SDS loading buffer. Following gel electrophoresis and membrane transfer, the membranes were blocked with 5% skimmed milk in PBST for 2 hours at room temperature. Subsequently, the membrane was incubated with 1 μg/mL anti-FLAG mAb (Sigma, F1804) or anti-β-tubulin (Immmuno Way, YM3030) mAb diluted in PBST containing 1% milk overnight at 4°C. After four washes with PBST, the blots were incubated with Horseradish peroxidase (HRP)-conjugated secondary antibody AffiniPure Goat Anti-Mouse in 1% skim milk diluted in PBST and incubated for one hour at room temperature. Finally, the blots were washed four times again by PBST and visualized using an Omni-ECL Femto Light Chemiluminescence Kit (EpiZyme,

SQ201) through a ChemiDoc MP Imaging System (Bio-Rad). To examine the S glycoprotein packaging efficiency, the PSV-containing supernatant was concentrated using a 30% sucrose cushion (30% sucrose, 15 mM Tris-HCl, 100 mM NaCl, 0.5 mM EDTA) at 20,000×g for 30 minutes at 4°C. The concentrated virus pellet was re-suspended in 1×SDS loading buffer and incubated at 95°C for 10 minutes, followed by western blot detecting the S glycoproteins by C-terminal HA tags and with the VSV-M serving as a loading control. Original western blot images have been deposited at Mendeley with DOI listed in the key resources table.

Immunofluorescence assay

Immunofluorescence assays were conducted to determine the expression levels of ACE2 orthologs with C-terminal fused FLAG tags. Specifically, the transfected cells were fixed and permeabilized by incubation with 100% methanol for 10 minutes at room temperature. Subsequently, the cells were incubated with a mouse antibody M2 (Sigma-Aldrich, F1804) diluted in PBS/1% BSA for one hour at 37°C. After a PBS wash, the cells were incubated with Alexa Fluor 594-conjugated goat anti-mouse IgG (Thermo Fisher Scientific, A32742) secondary antibody diluted in 1% BSA/PBS for one hour at 37°C. The images were captured and merged with a fluorescence microscope (Mshot, MI52-N) after the nucleus was stained blue with Hoechst 33342 reagent (1:5,000 dilution in PBS).

Recombinant glycoprotein production for cryo-EM analysis

The P.dav ACE2 and P.nat ACE2.M2 ectodomain or MOW15-22 RBD were produced and purified using Expi293F or ExpiCHO-S cells. Expi293F cells were grown to a density of 3×10^6 cells/mL and transfected using the ExpiFectamine 293 Transfection Kit (ThermoFisher Scientific) and expression was carried out for 4 days post-transfection at 37°C with 8% CO_2. The P.dav ACE2 ectodomain and MOW15-22 RBD were purified from clarified supernatants using HisTrap HP affinity columns (Cytiva) and washed with ten column volumes of 10 mM imidazole, 25 mM sodium phosphate pH 8.0, and 300 mM NaCl before elution with two column volumes of 300 mM imidazole, 25 mM sodium phosphate pH 8.0, and 300 mM NaCl. Purified P.dav ACE2 ectodomain or MOW15-22 RBD were then further purified by size exclusion chromatography using a Superdex 200 Increase 10/300 GL column (Cytiva) and concentrated using centrifugal filters (Amicon Ultra) before being flash frozen. P.nat ACE2.M2 with twin-strep tag was purified with Strep-Tactin XT 4Flow high-capacity resin (IBA, 2-5030-002), washed with ten column volumes of wash buffer (100 mM Tris/HCl, pH 8.0, 150 mM NaCl), and then eluted with a buffer containing 100 mM Tris/HCl, pH 8.0, 150 mM NaCl, 50 mM biotin. P.nat ACE2.M2 and MOW15-22 RBD were initially mixed at a molar ratio of 1:1.5 and incubated for 1h on ice, and then were further purified in a Superose 6 Increase 10/300 GL column (Cytiva) with a buffer containing 20 Tris/HCl, pH 8.0, 150 mM NaCl. The fractions containing the complex were collected and concentrated to 1 mg/ml for further use.

CryoEM sample preparation, data collection, and data processing

Complex formation was performed by mixing a 1:2.5 molar ratio of the P.dav ACE2 ectodomain (residues 17-723 with C-terminal octa-histidine tag) with the MOW15-22 RBD (residues 351-599 with C-terminal octa-histidine tag) before incubation for 1 hour at room temperature. CryoEM grids of the complex were prepared using two separate methods and data were combined during data processing. For the first dataset, 3 µL of 4 mg/ml complex with 6 mM 3-[(3-Cholamidopropyl)dimethylammonio]-2-hydroxy-1-propanesulfonate (CHAPSO) were applied onto freshly glow discharged R 2/2 UltrAuFoil grids[72] prior to plunge freezing using a

vitrobot MarkIV (ThermoFisher Scientific) with a blot force of 0 and 5.5 sec blot time at 100% humidity and 22°C. 14,561 movies were collected from UltrAuFoil grids with CHAPSO detergent with a defocus range comprised between -0.2 and -3.5 μm, and stage tilt angle of 0, 20° and 30°[73]. For the second dataset, 3 μL of 0.2 mg/mL complex was added to the glow discharged side of R 2/2 UltrAuFoil grids and 1μL was added to the back side before plunging into liquid ethane using a GP2 (Leica) with 6 sec blot time. 8,238 movies were collected with a defocus range comprised between -0.2 and -3.5 μm. The data were acquired using an FEI Titan Krios transmission electron microscope operated at 300kV and equipped with a Gatan K3 direct detector and Gatan Quantum GIF energy filter, operated in zero-loss mode with a slit width of 20 eV. Automated data collection was carried out using SerialEM[74] or Leginon[75] at a nominal magnification of 105,000× with a pixel size of 0.843 Å. The dose rate was adjusted to 9 counts/pixel/s, and each movie was acquired in counting mode fractionated in 100 frames of 40 ms. Movie frame alignment, estimation of the microscope contrast-transfer function parameters, particle picking, and extraction were carried out using Warp[76]. Particles were extracted with a box size of 192 pixels with a pixel size of 1.686Å. Two rounds of reference-free 2D classification were performed using CryoSPARC[77] to select well-defined particle images from each dataset. Particles belonging to classes with the best resolved RBD and ACE2 density were selected. To improve particle picking further, we trained the Topaz[78] picker on Warp-picked particle sets belonging to the selected classes after 2D classification. The particles picked using Topaz were extracted and subjected to 2D classification using cryoSPARC, which improved the number of unique 2D views. The two different particle sets picked from Warp and Topaz were merged and duplicate particle picks were removed using a minimum distance cutoff of 160Å. Initial model generation was done using ab-initio reconstruction in cryoSPARC and used as references for a heterogenous 3D refinement in cryoSPARC. After two rounds of ab-initio reconstructions and heterogeneous refinements to remove junk particles, 3D refinement was carried out using non-uniform refinement with per-particle defocus refinement in cryoSPARC[79] and the particles were transferred from cryoSPARC to Relion using pyem (https://github.com/asarnow/pyem) to be subjected to the Bayesian polishing procedure implemented in Relion[80] during which particles were re-extracted with a box size of 320 pixels and a pixel size of 1.0 Å. After ab-initio reconstructions and heterogeneous refinements to remove RBD unbound class, subsequent 3D refinement used non-uniform refinement along with per-particle defocus refinement in cryoSPARC. To further improve the density of the RBD:ACE2 interface, local refinement was performed using cryoSPARC with a soft mask comprising one ACE2 peptidase domain and the bound RBD. Local resolution estimation, filtering, and sharpening were carried out using cryoSPARC to yield the final reconstruction at 2.8 Å resolution comprising 705,956 particles. Reported resolutions are based on the 0.143 gold-standard Fourier shell correlation (FSC) criterion and Fourier shell correlation curves were corrected for the effects of soft masking by high-resolution noise substitution[81,82].

For P.nat ACE2.M2 and MOW15-22 RBD complex, 3.5 μL of the purified complex at 0.8 mg/mL was applied to glow-discharged Cu R1.2/1.3 holey carbon grid (200 mesh, Quantifoil) with a final concentration of 1% glycerol. After incubation for 20 s, the grids were blotted with force 0 for 2 s at 4°C and 100% humidity, and plunge-frozen into liquid ethane using Vitrobot Mark IV (FEI Thermo Fisher). Grids were then transferred to a CRYO ARM 300 electron microscope (JEOL, Japan) operating at 300 kV equipped with a K3 direct electron detector (Gatan, USA). Cryo-EM images were recorded automatically using Serial-EM software with a super-resolution pixel size of 0.475 Å /pixel at defocus values ranging from -0.5 to -2.5 μm at a calibrated magnification of 50,000. Data were collected at a frame rate of 40 frames per second. The total electron dose was 40 e-/Å2. 4,214 movies were collected and subjected to patch motion

correction and patch CTF estimation in cryoSPARC. Particles were automatically picked with a diameter of 120 Å from 200 micrographs. ~50,000 particles representing good classes were used to train a topaz model for auto-picking. A subset of 785,038 particles was selected after two rounds of 2D classification, followed by an ab initio reconstitution and a heterogeneous refinement. One class representing intact P.nat ACE2.M2-MOW15-22 RBD complex was further selected for heterogeneous refinement requesting three classes. One class containing 244,733 particles with good features was subjected to non-uniform refinement, yielding a 3.3 Å reconstruction.

Model building and refinement

UCSF Chimera[83] was used to rigid-body dock models into the sharpened cryoEM map and adjustments and refinement were carried out with Coot[84] and Rosetta[85,86] using sharpened and unsharpened maps. Validation used Molprobity[87], Phenix[88] and Privateer[89].

Bioinformatic and structural analysis

Sequence alignments of different bats ACE2 were performed using either the MUSCLE algorithm by MEGA-X (version 10.1.8) or ClustalW software (https://www.genome.jp/tools-bin/clustalw) with slight position adjustment to align indels. Phylogenetic trees were generated using the maximal likelihood method in IQ-TREE (http://igtree.cibiv.univie.ac.at/) (1000 Bootstraps) and refined with iTOL (v6) (https://itol.embl.de/). The structures of PnNL2018B, and HKU31 RBDs were predicted using alphaFold2.ipynb-Colaboratory[26]. The structure of NeoCoV RBD & P.pip ACE2 complex (7WPO), MERS-CoV RBD (4KQZ), HKU4 RBD (4QZV), HKU5 RBD (5XGR), MOW15-22 RBD, HKU31 RBD, and PnNL2018B RBD were visualized and analyzed using the ChimeraX(V.1.7.1). Analysis of buried surface area and identification of interface residues were assisted by PISA[66].

QUANTIFICATION AND STATISTICAL ANALYSIS

Most experiments related to pseudovirus infection were conducted 2-3 times with 2-4 biological repeats, technical repeats are indicated in the legends. Representative results were shown. All data were presented by MEAN ± SD. Unpaired two-tailed t-tests were conducted for all statistical analyses for two independent groups using GraphPad Prism 8. $P<0.05$ was considered significant. * $p<0.05$, ** $p<0.01$, *** $p<0.005$, and **** $p<0.001$. No data were excluded for data analysis.

References

1. CSR MERS outbreaks. World Health Organization - Regional Office for the Eastern Mediterranean. https://www.emro.who.int/health-topics/mers-cov/mers-outbreaks.html.

2. Alagaili, A.N., Briese, T., Mishra, N., Kapoor, V., Sameroff, S.C., Burbelo, P.D., de Wit, E., Munster, V.J., Hensley, L.E., Zalmout, I.S., et al. (2014). Middle East respiratory syndrome coronavirus infection in dromedary camels in Saudi Arabia. MBio *5*, e00884–14.

3. Meyer, B., Müller, M.A., Corman, V.M., Reusken, C.B.E.M., Ritz, D., Godeke, G.-J., Lattwein, E., Kallies, S., Siemens, A., van Beek, J., et al. (2014). Antibodies against MERS coronavirus in dromedary camels, United Arab Emirates, 2003 and 2013. Emerg. Infect. Dis. *20*, 552–559.

4. Memish, Z.A., Cotten, M., Meyer, B., Watson, S.J., Alsahafi, A.J., Al Rabeeah, A.A., Corman, V.M., Sieberg, A., Makhdoom, H.Q., Assiri, A., et al. (2014). Human infection with MERS coronavirus after exposure to infected camels, Saudi Arabia, 2013. Emerg. Infect. Dis. *20*, 1012–1015.

5. Azhar, E.I., El-Kafrawy, S.A., Farraj, S.A., Hassan, A.M., Al-Saeed, M.S., Hashem, A.M., and Madani, T.A. (2014). Evidence for camel-to-human transmission of MERS coronavirus. N. Engl. J. Med. *370*, 2499–2505.

6. Mohd, H.A., Al-Tawfiq, J.A., and Memish, Z.A. (2016). Middle East Respiratory Syndrome Coronavirus (MERS-CoV) origin and animal reservoir. Virol. J. *13*, 87.

7. Anthony, S.J., Gilardi, K., Menachery, V.D., Goldstein, T., Ssebide, B., Mbabazi, R., Navarrete-Macias, I., Liang, E., Wells, H., Hicks, A., et al. (2017). Further Evidence for Bats as the Evolutionary Source of Middle East Respiratory Syndrome Coronavirus. MBio *8*. 10.1128/mBio.00373-17.

8. Tolentino, J.E., Lytras, S., Ito, J., and Sato, K. (2024). Recombination analysis on the receptor switching event of MERS-CoV and its close relatives: implications for the emergence of MERS-CoV. Virol. J. *21*, 84.

9. Xiong, Q., Cao, L., Ma, C., Tortorici, M.A., Liu, C., Si, J., Liu, P., Gu, M., Walls, A.C., Wang, C., et al. (2022). Close relatives of MERS-CoV in bats use ACE2 as their functional receptors. Nature *612*, 748–757.

10. Ithete, N.L., Stoffberg, S., Corman, V.M., Cottontail, V.M., Richards, L.R., Schoeman, M.C., Drosten, C., Drexler, J.F., and Preiser, W. (2013). Close relative of human Middle East respiratory syndrome coronavirus in bat, South Africa. Emerg. Infect. Dis. *19*, 1697–1699.

11. Lu, G., Wang, Q., and Gao, G.F. (2015). Bat-to-human: spike features determining "host jump" of coronaviruses SARS-CoV, MERS-CoV, and beyond. Trends Microbiol. *23*, 468–478.

12. Wang, N., Shi, X., Jiang, L., Zhang, S., Wang, D., Tong, P., Guo, D., Fu, L., Cui, Y., Liu,

X., et al. (2013). Structure of MERS-CoV spike receptor-binding domain complexed with human receptor DPP4. Cell Res. *23*, 986–993.

13. Wang, Q., Qi, J., Yuan, Y., Xuan, Y., Han, P., Wan, Y., Ji, W., Li, Y., Wu, Y., Wang, J., et al. (2014). Bat origins of MERS-CoV supported by bat coronavirus HKU4 usage of human receptor CD26. Cell Host Microbe *16*, 328–337.

14. Lau, S.K.P., Zhang, L., Luk, H.K.H., Xiong, L., Peng, X., Li, K.S.M., He, X., Zhao, P.S.-H., Fan, R.Y.Y., Wong, A.C.P., et al. (2018). Receptor Usage of a Novel Bat Lineage C Betacoronavirus Reveals Evolution of Middle East Respiratory Syndrome-Related Coronavirus Spike Proteins for Human Dipeptidyl Peptidase 4 Binding. J. Infect. Dis. *218*, 197–207.

15. Tse, L.V., Hou, Y.J., McFadden, E., Lee, R.E., Scobey, T.D., Leist, S.R., Martinez, D.R., Meganck, R.M., Schäfer, A., Yount, B.L., et al. (2023). A MERS-CoV antibody neutralizes a pre-emerging group 2c bat coronavirus. Sci. Transl. Med. *15*, eadg5567.

16. Raj, V.S., Mou, H., Smits, S.L., Dekkers, D.H.W., Müller, M.A., Dijkman, R., Muth, D., Demmers, J.A.A., Zaki, A., Fouchier, R.A.M., et al. (2013). Dipeptidyl peptidase 4 is a functional receptor for the emerging human coronavirus-EMC. Nature *495*, 251–254.

17. Lu, G., Hu, Y., Wang, Q., Qi, J., Gao, F., Li, Y., Zhang, Y., Zhang, W., Yuan, Y., Bao, J., et al. (2013). Molecular basis of binding between novel human coronavirus MERS-CoV and its receptor CD26. Nature *500*, 227–231.

18. Chen, J., Yang, X., Si, H., Gong, Q., Que, T., Li, J., Li, Y., Wu, C., Zhang, W., Chen, Y., et al. (2023). A bat MERS-like coronavirus circulates in pangolins and utilizes human DPP4 and host proteases for cell entry. Cell *186*, 850–863.e16.

19. Wu, K., Li, W., Peng, G., and Li, F. (2009). Crystal structure of NL63 respiratory coronavirus receptor-binding domain complexed with its human receptor. Proc. Natl. Acad. Sci. U. S. A. *106*, 19970–19974.

20. Li, F., Li, W., Farzan, M., and Harrison, S.C. (2005). Structure of SARS coronavirus spike receptor-binding domain complexed with receptor. Science *309*, 1864–1868.

21. Hofmann, H., Pyrc, K., van der Hoek, L., Geier, M., Berkhout, B., and Pöhlmann, S. (2005). Human coronavirus NL63 employs the severe acute respiratory syndrome coronavirus receptor for cellular entry. Proc. Natl. Acad. Sci. U. S. A. *102*, 7988–7993.

22. Li, W., Moore, M.J., Vasilieva, N., Sui, J., Wong, S.K., Berne, M.A., Somasundaran, M., Sullivan, J.L., Luzuriaga, K., Greenough, T.C., et al. (2003). Angiotensin-converting enzyme 2 is a functional receptor for the SARS coronavirus. Nature *426*, 450–454.

23. Walls, A.C., Park, Y.-J., Tortorici, M.A., Wall, A., McGuire, A.T., and Veesler, D. (2020). Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. Cell *181*, 281–292.e6.

24. Ma, C., Liu, C., Xiong, Q., Gu, M., Shi, L., Wang, C., Si, J., Tong, F., Liu, P., Huang, M.,

et al. (2023). Broad host tropism of ACE2-using MERS-related coronaviruses and
determinants restricting viral recognition. Cell Discov *9*, 57.

25. International Committee on Taxonomy of Viruses Executive Committee (2020). The new
scope of virus taxonomy: partitioning the virosphere into 15 hierarchical ranks. Nat
Microbiol *5*, 668–674.

26. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., et al. (2021). Highly accurate
protein structure prediction with AlphaFold. Nature *596*, 583–589.

27. Speranskaya, A.S., Artiushin, I.V., Samoilov, A.E., Korneenko, E.V., Khabudaev, K.V.,
Ilina, E.N., Yusefovich, A.P., Safonova, M.V., Dolgova, A.S., Gladkikh, A.S., et al. (2023).
Identification and Genetic Characterization of MERS-Related Coronavirus Isolated from
Nathusius' Pipistrelle (Pipistrellus nathusii) near Zvenigorod (Moscow Region, Russia). Int.
J. Environ. Res. Public Health *20*. 10.3390/ijerph20043702.

28. Mols, V.C., Lamers, M.M., Leijten, L.M., Breugem, T.I., van de Bildt, M.W., van den Doel,
P.B., Lina, P.H., Koopmans, M.P., Haagmans, B.L., Kuiken, T., et al. (2023). Intestinal
Tropism of a Betacoronavirus (Merbecovirus) in Nathusius's Pipistrelle Bat (Pipistrellus
nathusii), Its Natural Host. J. Virol. *97*, e0009923.

29. Juste, J., and Paunović, M. (2016). IUCN Red List of Threatened Species: Pipistrellus
nathusii. IUCN Red List of Threatened Species.

30. Schmieder,   P.D.   Pipistrellus   nathusii.
https://www.eurobats.org/About_eurobats/Protected_bat_species/Pipistrellus_nathusii.

31. Driller, M., Brown, T., Currie, S.E., Hiller, M., Winkler, S., Pippel, M., Voigt, C.C., Fickel,
J., and Mazzoni, C.J. (2024). A Haplotype-resolved reference genome of a long-distance
migratory bat, Pipistrellus nathusii (Keyserling & Blasius, 1839). DNA Res.
10.1093/dnares/dsae018.

32. Letko, M. (2024). Functional assessment of cell entry and receptor use for merbecoviruses.
bioRxiv. 10.1101/2024.03.13.584892.

33. Yan, H., Jiao, H., Liu, Q., Zhang, Z., Xiong, Q., Wang, B.-J., Wang, X., Guo, M., Wang,
L.-F., Lan, K., et al. (2021). ACE2 receptor usage reveals variation in susceptibility to
SARS-CoV and SARS-CoV-2 infection among bat species. Nat Ecol Evol *5*, 600–608.

34. Menachery, V.D., Dinnon, K.H., 3rd, Yount, B.L., Jr, McAnarney, E.T., Gralinski, L.E.,
Hale, A., Graham, R.L., Scobey, T., Anthony, S.J., Wang, L., et al. (2020). Trypsin
Treatment Unlocks Barrier for Zoonotic Bat Coronavirus Infection. J. Virol. *94*.
10.1128/JVI.01774-19.

35. Kleine-Weber, H., Elzayat, M.T., Hoffmann, M., and Pöhlmann, S. (2018). Functional
analysis of potential cleavage sites in the MERS-coronavirus spike protein. Sci. Rep. *8*,
16597.

36. Millet, J.K., and Whittaker, G.R. (2014). Host cell entry of Middle East respiratory syndrome coronavirus after two-step, furin-mediated activation of the spike protein. Proc. Natl. Acad. Sci. U. S. A. *111*, 15214–15219.

37. Hoffmann, M., Kleine-Weber, H., Schroeder, S., Krüger, N., Herrler, T., Erichsen, S., Schiergens, T.S., Herrler, G., Wu, N.-H., Nitsche, A., et al. (2020). SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. Cell *181*, 271–280.e8.

38. Riva, L., Yuan, S., Yin, X., Martin-Sancho, L., Matsunaga, N., Pache, L., Burgstaller-Muehlbacher, S., De Jesus, P.D., Teriete, P., Hull, M.V., et al. (2020). Discovery of SARS-CoV-2 antiviral drugs through large-scale compound repurposing. Nature *586*, 113–119.

39. Xia, S., Liu, M., Wang, C., Xu, W., Lan, Q., Feng, S., Qi, F., Bao, L., Du, L., Liu, S., et al. (2020). Inhibition of SARS-CoV-2 (previously 2019-nCoV) infection by a highly potent pan-coronavirus fusion inhibitor targeting its spike protein that harbors a high capacity to mediate membrane fusion. Cell Res. *30*, 343–355.

40. Stalin Raj, V., Okba, N.M.A., Gutierrez-Alvarez, J., Drabek, D., van Dieren, B., Widagdo, W., Lamers, M.M., Widjaja, I., Fernandez-Delgado, R., Sola, I., et al. (2018). Chimeric camel/human heavy-chain antibodies protect against MERS-CoV infection. Sci Adv *4*, eaas9667.

41. Wrapp, D., De Vlieger, D., Corbett, K.S., Torres, G.M., Wang, N., Van Breedam, W., Roose, K., van Schie, L., VIB-CMB COVID-19 Response Team, Hoffmann, M., et al. (2020). Structural Basis for Potent Neutralization of Betacoronaviruses by Single-Domain Camelid Antibodies. Cell *181*, 1004–1015.e15.

42. Pinto, D., Sauer, M.M., Czudnochowski, N., Low, J.S., Tortorici, M.A., Housley, M.P., Noack, J., Walls, A.C., Bowen, J.E., Guarino, B., et al. (2021). Broad betacoronavirus neutralization by a stem helix-specific human antibody. Science *373*, 1109–1116.

43. Sun, X., Yi, C., Zhu, Y., Ding, L., Xia, S., Chen, X., Liu, M., Gu, C., Lu, X., Fu, Y., et al. (2022). Neutralization mechanism of a human antibody with pan-coronavirus reactivity including SARS-CoV-2. Nat Microbiol *7*, 1063–1074.

44. Du, Y., Shi, R., Zhang, Y., Duan, X., Li, L., Zhang, J., Wang, F., Zhang, R., Shen, H., Wang, Y., et al. (2021). A broadly neutralizing humanized ACE2-targeting antibody against SARS-CoV-2 variants. Nat. Commun. *12*, 5000.

45. Millet, J.K., Jaimes, J.A., and Whittaker, G.R. (2021). Molecular diversity of coronavirus host cell entry receptors. FEMS Microbiol. Rev. *45*. 10.1093/femsre/fuaa057.

46. Li, F. (2015). Receptor recognition mechanisms of coronaviruses: a decade of structural studies. J. Virol. *89*, 1954–1964.

47. Tortorici, M.A., Walls, A.C., Joshi, A., Park, Y.-J., Eguia, R.T., Miranda, M.C., Kepl, E., Dosey, A., Stevens-Ayers, T., Boeckh, M.J., et al. (2022). Structure, receptor recognition, and antigenicity of the human coronavirus CCoV-HuPn-2018 spike glycoprotein. Cell *185*,

2279–2291.e17.

48. Yeager, C.L., Ashmun, R.A., Williams, R.K., Cardellichio, C.B., Shapiro, L.H., Look, A.T.,
and Holmes, K.V. (1992). Human aminopeptidase N is a receptor for human coronavirus
229E. Nature *357*, 420–422.

49. Tresnan, D.B., Levis, R., and Holmes, K.V. (1996). Feline aminopeptidase N serves as a
receptor for feline, canine, porcine, and human coronaviruses in serogroup I. J. Virol. *70*,
8669–8674.

50. Li, W., Hulswit, R.J.G., Kenney, S.P., Widjaja, I., Jung, K., Alhamo, M.A., van Dieren, B.,
van Kuppeveld, F.J.M., Saif, L.J., and Bosch, B.-J. (2018). Broad receptor engagement of
an emerging global coronavirus may potentiate its diverse cross-species transmissibility.
Proc. Natl. Acad. Sci. U. S. A. *115*, E5135–E5143.

51. Zhou, P., Yang, X.-L., Wang, X.-G., Hu, B., Zhang, L., Zhang, W., Si, H.-R., Zhu, Y., Li,
B., Huang, C.-L., et al. (2020). A pneumonia outbreak associated with a new coronavirus of
probable bat origin. Nature *579*, 270–273.

52. Kuhn, J.H., Li, W., Choe, H., and Farzan, M. (2004). Angiotensin-converting enzyme 2: a
functional receptor for SARS coronavirus. Cell. Mol. Life Sci. *61*, 2738–2743.

53. Liu, K., Pan, X., Li, L., Yu, F., Zheng, A., Du, P., Han, P., Meng, Y., Zhang, Y., Wu, L., et
al. (2021). Binding and molecular basis of the bat coronavirus RaTG13 virus to ACE2 in
humans and other species. Cell *184*, 3438–3451.e10.

54. Cameroni, E., Bowen, J.E., Rosen, L.E., Saliba, C., Zepeda, S.K., Culap, K., Pinto, D.,
VanBlargan, L.A., De Marco, A., di Iulio, J., et al. (2022). Broadly neutralizing antibodies
overcome SARS-CoV-2 Omicron antigenic shift. Nature *602*, 664–670.

55. McCallum, M., Czudnochowski, N., Rosen, L.E., Zepeda, S.K., Bowen, J.E., Walls, A.C.,
Hauser, K., Joshi, A., Stewart, C., Dillen, J.R., et al. (2022). Structural basis of SARS-CoV-
2 Omicron immune evasion and receptor engagement. Science *375*, 864–868.

56. Reusken, C.B.E.M., Lina, P.H.C., Pielaat, A., de Vries, A., Dam-Deisz, C., Adema, J.,
Drexler, J.F., Drosten, C., and Kooi, E.A. (2010). Circulation of group 2 coronaviruses in a
bat species common to urban areas in Western Europe. Vector Borne Zoonotic Dis. *10*,
785–791.

57. Zaki, A.M., van Boheemen, S., Bestebroer, T.M., Osterhaus, A.D.M.E., and Fouchier,
R.A.M. (2012). Isolation of a novel coronavirus from a man with pneumonia in Saudi
Arabia. N. Engl. J. Med. *367*, 1814–1820.

58. He, W.-T., Hou, X., Zhao, J., Sun, J., He, H., Si, W., Wang, J., Jiang, Z., Yan, Z., Xing, G.,
et al. (2022). Virome characterization of game animals in China reveals a spectrum of
emerging pathogens. Cell *185*, 1117–1129.e8.

59. Lau, S.K.P., Luk, H.K.H., Wong, A.C.P., Fan, R.Y.Y., Lam, C.S.F., Li, K.S.M., Ahmed,
S.S., Chow, F.W.N., Cai, J.-P., Zhu, X., et al. (2019). Identification of a Novel

Betacoronavirus (Merbecovirus) in Amur Hedgehogs from China. Viruses *11*. 10.3390/v11110980.

60. Corman, V.M., Kallies, R., Philipps, H., Göpner, G., Müller, M.A., Eckerle, I., Brünink, S., Drosten, C., and Drexler, J.F. (2014). Characterization of a novel betacoronavirus related to middle East respiratory syndrome coronavirus in European hedgehogs. J. Virol. *88*, 717–724.

61. Kumar, S., Stecher, G., Li, M., Knyaz, C., and Tamura, K. (2018). MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. Mol. Biol. Evol. *35*, 1547–1549.

62. Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von Haeseler, A., and Lanfear, R. (2020). IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. Mol. Biol. Evol. *37*, 1530–1534.

63. Letunic, I., and Bork, P. (2024). Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic tree display and annotation tool. Nucleic Acids Res. 10.1093/nar/gkae268.

64. Goddard, T.D., Huang, C.C., Meng, E.C., Pettersen, E.F., Couch, G.S., Morris, J.H., and Ferrin, T.E. (2018). UCSF ChimeraX: Meeting modern challenges in visualization and analysis. Protein Sci. *27*, 14–25.

65. Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., and Steinegger, M. (2022). ColabFold: making protein folding accessible to all. Nat. Methods *19*, 679–682.

66. Krissinel, E., and Henrick, K. (2007). Inference of macromolecular assemblies from crystalline state. J. Mol. Biol. *372*, 774–797.

67. Thompson, J.D., Gibson, T.J., and Higgins, D.G. (2002). Multiple sequence alignment using ClustalW and ClustalX. Curr. Protoc. Bioinformatics *Chapter 2*, Unit 2.3.

68. Han, X., Qi, J., Song, H., Wang, Q., Zhang, Y., Wu, Y., Lu, G., Yuen, K.-Y., Shi, Y., and Gao, G.F. (2017). Structure of the S1 subunit C-terminal domain from bat-derived coronavirus HKU5 spike protein. Virology *507*, 101–109.

69. Whitt, M.A. (2010). Generation of VSV pseudotypes using recombinant ΔG-VSV for studies on virus entry, identification of entry inhibitors, and immune responses to vaccines. J. Virol. Methods *169*, 365–374.

70. Reed, L.J., and Muench, H. (1938). A Simple Method for Estimating Fifty Per Cent Endpoints.

71. Nie, J., Li, Q., Wu, J., Zhao, C., Hao, H., Liu, H., Zhang, L., Nie, L., Qin, H., Wang, M., et al. (2020). Quantification of SARS-CoV-2 neutralizing antibody by a pseudotyped virus-based assay. Nat. Protoc. *15*, 3699–3715.

72. Russo, C.J., and Passmore, L.A. (2014). Electron microscopy: Ultrastable gold substrates for electron cryomicroscopy. Science *346*, 1377–1380.

73. Tan, Y.Z., Baldwin, P.R., Davis, J.H., Williamson, J.R., Potter, C.S., Carragher, B., and Lyumkis, D. (2017). Addressing preferred specimen orientation in single-particle cryo-EM through tilting. Nat. Methods *14*, 793–796.

74. Mastronarde, D.N. (2005). Automated electron microscope tomography using robust prediction of specimen movements. J. Struct. Biol. *152*, 36–51.

75. Suloway, C., Pulokas, J., Fellmann, D., Cheng, A., Guerra, F., Quispe, J., Stagg, S., Potter, C.S., and Carragher, B. (2005). Automated molecular microscopy: the new Leginon system. J. Struct. Biol. *151*, 41–60.

76. Tegunov, D., and Cramer, P. (2019). Real-time cryo-electron microscopy data preprocessing with Warp. Nat. Methods *16*, 1146–1152.

77. Punjani, A., Rubinstein, J.L., Fleet, D.J., and Brubaker, M.A. (2017). cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. Nat. Methods *14*, 290–296.

78. Bepler, T., Morin, A., Rapp, M., Brasch, J., Shapiro, L., Noble, A.J., and Berger, B. (2019). Positive-unlabeled convolutional neural networks for particle picking in cryo-electron micrographs. Nat. Methods *16*, 1153–1160.

79. Punjani, A., Zhang, H., and Fleet, D.J. (2020). Non-uniform refinement: adaptive regularization improves single-particle cryo-EM reconstruction. Nat. Methods *17*, 1214–1221.

80. Zivanov, J., Nakane, T., Forsberg, B.O., Kimanius, D., Hagen, W.J., Lindahl, E., and Scheres, S.H. (2018). New tools for automated high-resolution cryo-EM structure determination in RELION-3. Elife *7*. 10.7554/eLife.42166.

81. Rosenthal, P.B., and Henderson, R. (2003). Optimal determination of particle orientation, absolute hand, and contrast loss in single-particle electron cryomicroscopy. J. Mol. Biol. *333*, 721–745.

82. Chen, S., McMullan, G., Faruqi, A.R., Murshudov, G.N., Short, J.M., Scheres, S.H.W., and Henderson, R. (2013). High-resolution noise substitution to measure overfitting and validate resolution in 3D structure determination by single particle electron cryomicroscopy. Ultramicroscopy *135*, 24–35.

83. Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng, E.C., and Ferrin, T.E. (2004). UCSF Chimera--a visualization system for exploratory research and analysis. J. Comput. Chem. *25*, 1605–1612.

84. Emsley, P., Lohkamp, B., Scott, W.G., and Cowtan, K. (2010). Features and development of Coot. Acta Crystallogr. D Biol. Crystallogr. *66*, 486–501.

85. Frenz, B., Rämisch, S., Borst, A.J., Walls, A.C., Adolf-Bryfogle, J., Schief, W.R., Veesler, D., and DiMaio, F. (2019). Automatically Fixing Errors in Glycoprotein Structures with Rosetta. Structure *27*, 134–139.e3.

86. Wang, R.Y.-R., Song, Y., Barad, B.A., Cheng, Y., Fraser, J.S., and DiMaio, F. (2016).
Automated structure refinement of macromolecular assemblies from cryo-EM maps using
Rosetta. Elife *5*. 10.7554/eLife.17219.

87. Chen, V.B., Arendall, W.B., 3rd, Headd, J.J., Keedy, D.A., Immormino, R.M., Kapral, G.J.,
Murray, L.W., Richardson, J.S., and Richardson, D.C. (2010). MolProbity: all-atom
structure validation for macromolecular crystallography. Acta Crystallogr. D Biol.
Crystallogr. *66*, 12–21.

88. Liebschner, D., Afonine, P.V., Baker, M.L., Bunkóczi, G., Chen, V.B., Croll, T.I., Hintze,
B., Hung, L.W., Jain, S., McCoy, A.J., et al. (2019). Macromolecular structure
determination using X-rays, neutrons and electrons: recent developments in Phenix. Acta
Crystallogr D Struct Biol *75*, 861–877.

89. Agirre, J., Iglesias-Fernández, J., Rovira, C., Davies, G.J., Wilson, K.S., and Cowtan, K.D.
(2015). Privateer: software for the conformational validation of carbohydrate structures. Nat.
Struct. Mol. Biol. *22*, 833–834.

Supplementary information for

Multiple independent acquisitions of ACE2 usage in MERS-related coronaviruses

Cheng-Bao Ma[1]#, Chen Liu[1]#, Young-Jun Park[2,3]#, Jingjing Tang[4]#, Jing Chen[4]#, Qing Xiong[1],
Jimin Lee[2], Cameron Stewart[2], Daniel Asarnow[2], Jack Brown[2], M. Alejandra Tortorici[2], Xiao
Yang[1], Ye-Hui Sun[1], Yuan-Mei Chen[1], Xiao Yu[1], Jun-Yu Si[1], Peng Liu[1], Fei Tong[1], Mei-Ling
Huang[1], Jing Li[1], Zheng-Li Shi[4,5]*, Zengqin Deng[4,6]*, David Veesler[2,3]*, Huan Yan[1,7,]*

The PDF document file includes:

Figure S1 to S10 and Tables S1-S2

A

Bat ACE2 orthologs (FLAG)

R.eag_	E.spe_	P.ale_	P.gig_	E.hel_	M.sob_	C.sph_	C.bra_	R.fer_	R.sin
R.cor_	R.mal_	R.aff_	R.sha_	R.alc_	R.pea_	H.arm_	H.pra_	H.gal_	M.lyr
T.mel_	N.lep_	A.cau_	C.per_	A.jam_	S.hon_	T.cir_	T.sau_	P.dis_	V.spe
D.rot_	M.hir_	P.par_	P.dav_	M.bla_	T.bra_	M.mol_	M.sch_	M.nat_	E.fus
N.hum_	P.pip_	P.kuh_	L.bor_	A.cin_	A.pal_	M.fea_	M.myo_	M.dav_	M.bra
M.luc_	P.nat_	hACE2_	Vector_						

B

Non-bat mammalian ACE2 orthologs (FLAG)

P.lar_	L.can_	F.cat_	P.con_	P.pard_	V.vul_	C.fam_	U.arc_	A.mel_	M.erm
M.put_	N.sch_	E.jub_	Z.cal_	P.anu_	T.gel_	M.fas_	R.rox_	P.tep_	P.tro
G.gor_	P.abe_	N.leu_	C.jac_	S.ape_	S.bol_	Cfer_	S.scr_	B.tau_	B.mut
B.bub_	C.hir_	O.ari_	O.vir_	P.leu_	C.gri_	Mmus_	Rnor_	J.jac_	I.tri
O.orc_	T.tru_	G.mel_	L.vex_	N.asi_	P.cat_	C.sim_	E.cab_	P.cin_	M.jav
E.eur_	O.cun_	Vector_							

Bats

P.dav_	L.bor_	P.nat_

Figure S1. Validation of the expression of ACE2 orthologs from bats or other mammalian species, related to Figure 3 (A-B) Immunofluorescence analyzing the expression of bat (A) or mammalian (B) ACE2 orthologs in HEK293T cells by detecting 3×FLAG tags fused to the C-terminal of the receptors. Scale bars:100 μm.

A

MOW15-22 S1-hFc binding

P.lar _	L.can _	F.cat _	P.con_	P.pard_	V.vul_	C.fam _	U.arc_	A.mel _	M.erm_
M.put _	N.sch _	E.jub _	Z.cal_	P.anu _	T.gel_	M.fas _	R.rox_	P.tep _	P.tro _
G.gor _	P.abe _	N.leu _	C.jac _	S.ape _	S.bol _	Cfer _	S.scr_	B.tau _	B.mut _
B.bub _	C.hir _	O.ari _	O.vir_	P.leu _	C.gri _	Mmus _	Rnor_	J.jac _	I.tri _
O.orc _	T.tru _	G.mel_	L.vex_	N.asi _	P.cat_	C.sim _	E.cab_	P.cin _	M.jav_
E.eur _	O.cun_	P.dav_	L.bor _	P.nat_	hACE2_	Vector_			

B

PnNL2018B S1-hFc binding

P.lar _	L.can _	F.cat _	P.con_	P.pard_	V.vul_	C.fam _	U.arc_	A.mel _	M.erm_
M.put_	N.sch _	E.jub _	Z.cal_	P.anu _	T.gel_	M.fas _	R.rox_	P.tep _	P.tro _
G.gor _	P.abe _	N.leu _	C.jac_	S.ape _	S.bol _	Cfer _	S.scr_	B.tau _	B.mut _
B.bub _	C.hir _	O.ari _	O.vir_	P.leu _	C.gri _	Mmus _	Rnor_	J.jac _	I.tri _
O.orc _	T.tru _	G.mel_	L.vex_	N.asi _	P.cat_	C.sim _	E.cab_	P.cin _	M.jav_
E.eur_	O.cun_	P.dav_	L.bor _	P.nat_	hACE2_	Vector_			

C

Selected MOW15-22/PnNL2018B RBD/S_1-hFc binding(10 µg/mL,high concentration)

	Vector	P.nat	L.can	P.con	P.pard	V.vul	C.fam	M.erm	C.jac	O.ari	M.jav
ACE2 FLAG											
MOW15-22 RBD											
MOW15-22 S1											
PnNL2018B RBD											
PnNL2018B S1											

Figure S2. MOW15-22 and PnNL2018B S glycoproteins exhibit narrow ACE2 tropism, related to Figure 3. (**A-B**) Images showing MOW15-22 (A) and PnNL2018B (B) S_1 subunit binding to HEK293T cells transiently expressing the indicated mammalian ACE2 orthologs analyzed by fluorescence microscopy. (**C**) Fluorescence images showing MOW15-22 and PnNL2018B RBD/S_1 subunit binding at a higher protein concentration (10 µg/mL) in selected mammalian ACE2 orthologs supporting MOW15-22 entry in Figure. 3B. Scale bars:100 µm.

Figure S3. Trypsin-dependence of bat ACE2 mediated MOW15-22 membrane fusion and
PSV entry, related to Figure 3. (**A-B**) Heat map (A) and bar graph (B) of MOW15-22 PSV
entry efficiency mediated by several ACE2 orthologs in the presence or absence of 100 μg/mL
TPCK-treated trypsin. Orthologs showing the highest entry-supporting efficiency (P.dav ACE2
for MOW15-22 and L.bor for PnNL2018B) were set as 100%. Red highlights the two bat ACE2s
supporting efficient MOW15-22 RBD binding. The dashed line indicates the background signal.
MEAN ± SD and unpaired two-tailed t-tests. n=3 biological replicates. (**C-D**) MOW15-22 spike-
mediated cell-cell membrane fusion in HEK293T cells stably expressing the indicated ACE2
orthologs in the presence of various concentrations of TPCK-treated trypsin. Fusion efficiency is
indicated by GFP intensity (C) and live-cell Renilla luciferase activity (D) through the
reconstitution of dual-split reporter proteins (DSPs). MEAN ± SD and unpaired two-tailed t-tests.
n=3 biological replicates. (**E-F**) MOW15-22 PSV entry efficiency in HEK293T cells stably
expressing ACE2 orthologs with the indicated concentration of TPCK-treated trypsin, as
indicated by GFP intensity (E) and luciferase (F). n=3 biological replicates for F. Scale bars in C
and E:200 μm.
16

Figure S4. ACE2 determinants critical for NeoCoV/PDF-2180 do not impact MOW15-
22/PnNL2018B ACE2 recognition, related to Figure 4. (**A**) Structural presentation of the four
host range determinants A-D critical for P.Pip ACE2 recognition by NeoCoV (PDB 7WPO). (**B**)
Schematic illustration of P.dav and P.par ACE2 mutants with sequences of indicated
determinants replaced by residues unfavorable for NeoCoV recognition. Glycosylation sites in
determinants A and C are indicated with ¥. (**C**) MOW15-22 and NeoCoV RBD binding to
HEK293T cells transiently expressing the indicated wild-type (WT) or mutants ACE2 orthologs.
The expression level of indicated ACE2 orthologs was verified by immunofluorescence (C, top
panel). Scale bars: 100 μm. (**D**) Western blot analysis of the expression of indicated WT and
mutated ACE2 in HEK293T cells. MW: molecular weight. (**E-F**) PSV entry efficiency of
PnNL2018B in HEK293T cells transiently expressing the L.bor (E) and P.pip (F) ACE2 mutants
affecting NeoCoV receptor recognition. n=3 biological replicates. RLU: relative light unit.

Figure S5. Mapping of ACE2 host range determinants for MOW15-22 and PnNL2018B by genetic swap analysis, related to Figure 4. (A-B) Virus entry-promoting efficiencies (A) and a sequence alignment of residues 400-450 (B) of the selected ACE2 orthologs. Residue numbering is based on P.dav ACE2, and the red dashed box indicates a region participating in host range determination (residues 420-432). ¥ : glycosylation, +:supportive, -: not supportive, +/-:weakly supportive. (C-D) Determinant mapping based on sequence swaps between residues 400-500 of M.bla and P.dav ACE2. Schematic illustration of chimeric ACE2 mutants enhancing M.bla ACE2 binding (C) abolishing P.dav ACE2 binding (D). RBD binding to (middle) and pseudovirus entry into (right) HEK293T cells transiently expressing the indicated ACE2 swap mutants are shown. MEAN ± SD and unpaired two-tailed t-tests. n=3 biological replicates. (E-F) Binding efficiencies of recombinant proteins comprising different domains of the MOW15-22 (C) or PnNL2018B (D) S_1 subunit to Caco2 cells stably expressing P.nat ACE2 or a P.nat ACE2 C432N mutant. Scale bars: 100 μm.

Figure S6. CryoEM data processing workflow for the P.dav ACE2-bound MOW15-22 RBD complex, related to Figure 5. (A-B) Representative electron micrograph (A) and 2D class averages of the P.dav ACE2-bound MOW15-22 RBD complex (B) embedded in vitreous ice. The scale bars represent 100 nm and 200Å, respectively. **(C)** Gold-standard Fourier shell correlation curve. The 0.143 cutoff is indicated by a horizontal dashed line. **(D)** Local resolution estimation was calculated using cryoSPARC and plotted on the sharpened map. **(E)** Data processing flowchart. CTF: contrast transfer function; NUR: non-uniform refinement.

Figure S7. Validation of critical residues for interactions between MOW15-22/PnNL2018B
and different ACE2 orthologs, related to Figure 5. (A) P. nat ACE2 mutants with reduced
PnNL2018B RBD or S_1 binding due to unfavorable substitutions of key interacting residues. (**B**)
P.pip, human, and P.kuh ACE2 mutants with enhanced ability to promote PnNL2018B RBD or
S_1 binding. (**C-D**) C.jac ACE2 mutants promoting enhanced MOW15-22 and PnNL2018B RBD
or S_1 binding and PSV entry or abolishing receptor function via introducing the N432 glycan
knock-in mutation. MEAN ± SD and unpaired two-tailed t-tests. n=3 biological replicates for D.
(**E**) VSV packaging efficiencies of MOW15-22 S harboring the indicated mutations.VSV-M
serves as a loading control. Scale bars in A, B, and C:100 μm.

11
12

Figure S8. Binding of different soluble P.nat ACE2 mutants to PnNL2018B S₁-hFc, related
2 **to Figure 6.** BLI analyses of binding kinetics of soluble dimeric ACE2 ectodomains from
3 wildtype (WT) P.nat ACE2 (**A**), P.nat ACE2.M1 (**B**), or P.nat ACE2.M2 (**C**) to the immobilized
4 PnNL2018B S₁-hFc. Analysis was conducted with global fitting (1:1 binding model) and the fit
5 to the data is shown in black.
6
7

1
2 **Figure S9. CryoEM data processing workflow for the P.nat ACE2.M2-bound MOW15-22**
3 **RBD complex, related to Figure 6.** (**A-B**) Representative electron micrograph and 2D class
4 averages of the P.nat ACE2.M2 bound MOW15-22 RBD complex embedded in vitreous ice. (**C**)
5 Flowchart of cryo-EM data processing. (**D**) Fourier shell correlation (FSC) curve was calculated
6 using two independent half maps, and resolution was estimated using the FSC=0.143 cutoff.

8
9

Figure S10. The neutralizing activity of monoclonal antibodies against PSV entry of
indicated merbecoviruses, related to Figure 6. (**A**) Neutralization efficiency of MERS-CoV
RBD-directed nanobodies against MERS-CoV and MOW15-22 pseudotyped viruses. MEAN ±
SD and unpaired two-tailed t-tests. n=3 biological replicates. (**B-C**) Sequence alignment
displaying corresponding sequences of S2P6 (B) and 76E1 (C) epitopes of indicated
coronaviruses. Red dashed box: S2P6 epitope. The MOW15-22 residue numbering is shown in B
and C. (**D**) Dose-dependent inhibition of NeoCoV pseudovirus entry by antibodies targeting the
stem helix (S2P6) or the S_2'/fusion peptide (76E1) in HEK293T cells expressing P.pip ACE2.
(**E**) The hACE2-specific antibody H11B11 does not inhibit MERS-CoV pseudovirus entry in
Caco2-P.nat ACE2.3M cells. n=2 biological replicates for D and E.

Table S1. CryoEM data collection, processing, and model refinement statistics, related to Figure 5.

	MOW15-22 RBD - P.dav ACE2 PDB 9C6O EMD 45253	MOW15-22 RBD - P.nat ACE2.M2 PDB 8ZUF EMD 60483
Data collection and processing		
Magnification	105,000	50,000
Voltage (kV)	300	300
Electron exposure (e–/Å^2)	60	40
Defocus range (μm)	-0.2 to -3.5	-0.5 to -2.5
Pixel size (Å)	0.843	0.95
Symmetry imposed	C1	C1
Initial particle images (no.)	5,091,367	1,844,603
Final particle images (no.)	705,956	244,733
Map resolution (Å) FSC threshold	2.8 0.143	3.3 0.143
Refinement		
Model resolution (Å) FSC threshold	3.0 0.5	3.6 0.5
Map sharpening B factor (Å^2)	-101.3	-152.3
Model composition Non-hydrogen atoms Protein residues Ligands	6,164 792 12	6,603 805 6
B factors (Å^2) Protein Ligand	17.25 27.55	21.42 39.89
R.m.s. deviations Bond lengths (Å) Bond angles (°)	0.010 1.041	0.004 0.777
Validation MolProbity score Clashscore Poor rotamers (%)	1.09 1.62 0.18	1.7 7.5 0.14
Ramachandran plot Favored (%) Allowed (%) Disallowed (%)	96.93 2.17 0.90	95.87 3.88 0.25

Table S2. Conservation of receptor-interacting residues between MOW15-22 and PnNL2018B, , related to Figure 5.

MOW15-22 residue number	MOW15-22 residue identity	PnNL2018B residue identity
505	G	G
506	C	C
507	V	V
508	D	G
509	K	Q
511	P	P
519	I	T
520	C	C
521	I	L
522	P	E
523	E	E
524	F	F
527	E	F
535	K	K
563	D	D
564	S	S
565	T	V
567	D	D
569	I	I
570	W	W
571	R	R

56